

# Network Structural Dependency in the Human Connectome Across the Life-Span

Markus D. Schirmer<sup>a,b,c,1\*</sup>, Ai Wern Chung<sup>d\*</sup>, P. Ellen Grant<sup>d</sup> and Natalia S. Rost<sup>a</sup>

<sup>a</sup>Stroke Division & Massachusetts General Hospital, J. Philip Kistler Stroke Research Center, Harvard Medical School, Boston, USA

<sup>b</sup>Computer Science and Artificial Intelligence Lab, Massachusetts Institute of Technology, Cambridge, USA

<sup>c</sup>Department of Population Health Sciences, German Centre for Neurodegenerative Diseases (DZNE), Germany

<sup>d</sup>Fetal-Neonatal Neuroimaging & Developmental Science Center, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

\*Authors contributed equally.

## Abstract

Principles of network theoretical topology have been widely studied in the human connectome. Of particular interest is the modular nature of the human brain, where the connectome is divided into subnetworks and subsequently studied to investigate changes with development, aging or disease. Here, we present a new, weighted network measure called the Network Dependency Index (NDI), which can be used to identify an individual region's importance to the global functioning of the network. Importantly, we utilize NDI to differentiate four different subnetworks (Tiers) in the human connectome following Gaussian Mixture Model fitting. After identification of these subnetworks, we analyze their individual topological aspects with respect to age and compare it to the rich-club topology, which we use to define three subnetworks (rich-club, feeder and seeder). Our results first demonstrate the efficacy of NDI to identify more consistent, central nodes of the connectome across age-groups, when compared to the rich-club framework. Stratifying the connectome accordingly by NDI led to consistent subnetworks across the life-span revealing distinct patterns associated with age where, e.g., the key relay nuclei and cortical regions are contained in a subnetwork with highest NDI. The divisions of the human connectome derived from our data-driven NDI framework have the potential to reveal topological alterations due to disease and/or development, which are not detectable by utilizing the connectome as a whole.

---

<sup>1</sup> Corresponding author: [mschirmer1@mgh.harvard.edu](mailto:mschirmer1@mgh.harvard.edu)

# Introduction

Network theoretical principles have been readily applied to the human connectome to investigate its structural and functional organization. Typically, a brain network comprises of nodes representing brain regions that are connected by edges representing either reconstructed white matter pathways from diffusion-weighted magnetic resonance imaging (dMRI) or functional correlations from functional MRI (fMRI). Furthermore, edges between two nodes may be weighted to reflect the strength of the connection (Bullmore and Sporns, 2009; Fornito et al., 2013). There has been an ongoing quest to identify the regions in brain networks that are critical for efficient network functioning (see e.g. (Hagmann et al., 2008)), as well as defining modules, or sets of nodes, that distinguish themselves from other nodes in the connectome. One network organizational principle that has been established in the human connectome is the rich club (RC) - a core subnetwork of brain regions that are strongly connected to form a high-cost, high-capacity backbone. The RC has been shown to be critical for effective communication in the connectome (van den Heuvel et al., 2012; Van Den Heuvel and Sporns, 2011) and has been studied in healthy subjects (Grayson et al., 2014; Schirmer and Chung, 2018; Van Den Heuvel and Sporns, 2011; Zhao et al., 2015), with development (Ball et al., 2014) and in disease (Collin et al., 2013; Daianu et al., 2015; Ray et al., 2014). While an important topological aspect in brain networks, defining RC membership is not straight-forward. Typically, an RC-regime is identified over a range of degrees where the RC coefficient is significantly greater than a distribution of equivalent RC coefficients computed from random networks. From this range, a single degree,  $k$ , is chosen and nodes with a degree greater or equal to  $k$  are deemed to form the RC subnetwork. The lack of consensus in choosing this  $k$ -th threshold is in part due to the variations in network construction and weightings employed in the field, leading to great variability in RC coefficients (Van Den Heuvel and Sporns, 2011). As such, the neuroscience community has yet to establish an optimal way of defining RC nodes.

In this work, we present an alternative for defining regions of the brain that are integral for network efficiency, and which can be consistently identified across the life-span, based on the network dependency index (NDI; (Woldeyohannes and Jiang, 2018)). First introduced in network communication science for critical node detection, NDI quantifies a node's importance as defined by the impact on the network's performance given the node's failure (or removal) from the system - simply put, NDI measures the dependency of the network on any given node. It also has the advantage of evaluating nodal importance by incorporating the network's efficiency for information

transport as measured by topological distance. Ordering brain regions by NDI then allows us to stratify the connectome into subnetworks according to nodal importance. We present a comparative investigation between RC and NDI frameworks for detecting a core subnetwork and their corresponding ‘peripheral’ subnetworks. We demonstrate the efficacy of NDI in the human connectome by applying our framework on a large, open-source, normal population from the NKI-Rockland study with an age range of 4 to 85 years. By doing so, we derive normative values and evaluate NDI with age to contextualize our findings with other literature in the field. In the following sections, we describe both RC and our NDI framework to define a core subnetwork from which to further stratify the entire connectome. In Results, we compare the core regions detected by both frameworks, and their corresponding network characteristics in relation to age. We end with a Discussion of our primary NDI findings in comparison to the RC formalism.

## Materials and Methods

### Study design and patient population

In this work we utilize data from the NKI-Rockland life-span study (Nooner et al., 2012). Preprocessed connectome data were obtained from the USC Multimodal Connectivity database<sup>2</sup>. MRI acquisition details are available elsewhere (Brown et al., 2012). In brief, a total of 196 connectomes of healthy participants are computed, from 3T dMRI acquisitions (64 gradient directions; TR=10000ms; TE=91ms; voxel size=2mm<sup>3</sup>; *b*-value=1000 s/mm<sup>2</sup>). Following eddy current and motion correction, diffusion tensors are modelled and deterministic tractography performed using fiber assignment by continuous tracking (angular threshold 45°; (Mori et al., 1999)). Regions of interest (ROI) are based on the Craddock atlas (Craddock et al., 2012), resulting in 188 ROIs and connections are weighted by the number of streamlines connecting pairs of ROIs. Here, we normalize each connectome by the maximum streamline count for each subject so that the connection weights  $w_{ij}$  within each subject are  $w_{ij} \in [0, 1]$ .

For part of our analysis, we divide the 196 participants into four age groups: U20 ≤ 20 years, 20 years < U40 ≤ 40 years, 40 years < U60 ≤ 60 years, 60 years < O60 ≤ 80 years. Four subjects were above 80 years old (81, 82, 83 and 85 years). As there were only four subjects, we included them in the O60 group. Table 1 characterizes the study cohort and groups.

---

<sup>2</sup><http://umcd.humanconnectomeproject.org>

Table 1: NKI-Rockland life-span study cohort characterization and their stratification by age (in years).

	Overall	U20	U40	U60	O60
<b>N</b>	196	53	67	47	29
<b>Age (Mean (SD))</b>	35.0 (20.0)	13.8 (4.1)	27.4 (5.9)	47.4 (5.4)	71.0 (6.8)
<b>Sex (Male; %)</b>	58.1	54.7	56.7	72.3	44.8

The available connectivity matrices include both cortical and subcortical structures. Before analysis, regions in the brainstem and cerebellum were removed from the network, resulting in 170 regions.

## Group and cohort connectomes

A group-averaged connectome of weighted matrices can be computed in two steps (Van Den Heuvel and Sporns, 2011). First, we calculate a binarized, group-average adjacency matrix by retaining edges that are present in at least 90% of the subjects in each group. Weights are subsequently added to the group-averaged adjacency matrix by taking the average weight of each connection across the group, generating a weighted group-averaged connectome  $W_{\text{group}}$ . In addition to calculating a group connectome for each age group, we also compute a cohort-based connectome,  $W_{\text{cohort}}$ , across all 196 subjects.

## Rich-club framework

We utilize  $W_{\text{group}}$  to subsequently calculate the weighted RC parameter  $\Phi_{\text{group}}(k)$  (Opsahl et al., 2008), where  $k$  denotes the degree of nodes. The RC parameter  $\Phi_{\text{group}}(k)$  is normalized relative to a set of comparable random networks of equal size and with similar connectivity distribution. Here, we generate 1000 random networks while preserving weight, degree and strength distributions of  $W_{\text{group}}$  (Rubinov and Sporns, 2010). For each of these random realizations of the graph, we calculate the weighted RC parameter  $\Phi_{\text{rand}}(k)$ . Finally, the normalized weighted RC parameter is calculated as

$$\Phi_{group}^{norm}(k) = \frac{\Phi_{norm}(k)}{\Phi_{rand}(k)}$$

For this metric,  $\Phi_{group}^{norm}(k) > 1$  denotes the presence of a rich-club. In our analysis, we select

$$k_{max}^{group}: \max(k), \text{ for which } \Phi_{group}^{norm}(k) > 1,$$

as the degree of the RC nodes of a given group, which allows us to determine the RC members with a degree of at least  $k$ .

### Rich-club subnetworks

For each group connectome, we first determine  $k_{max}^{group}$ , as implemented in the Brain Connectivity Toolbox (Rubinov and Sporns, 2010). Statistical significance of  $\Phi_{group}^{norm}(k) > 1$  is assessed by performing a right-sided t-test for each  $k$  ( $p < 0.05$  after Bonferroni correction, where the number of tests equals the maximum degree within each group connectome). After establishing the RC nodes, remaining nodes in the group connectome were categorized according to their relationship to the RC subnetwork. All non-RC nodes were identified as feeders ( $F$ ) if they were connected to an RC node, or seeders ( $S$ ) if they shared no connection with an RC node (Schirmer and Chung, 2018).

### Network dependency index framework

The NDI score has recently been described in unweighted networks (Woldeyohannes and Jiang, 2018). Here, we extend their formalism to allow for use in weighted networks. Given a connectivity matrix  $W_{group} = \{w_{ij}\}$  in a network  $G$  with  $n$  nodes, we first calculate the topological distance matrix  $D$  between all node pairs using the inverse of the connection strength  $w_{ij}$  between nodes  $i$  and  $j$  as an initial topological distance. The information measure  $(I^n)_{ij}$  between nodes  $i$  and  $j$  can then be calculated as  $1/D_{ij}$ , resulting in an information measure matrix  $I^n$ . As a next step, we normalize  $I^n$  by the maximum information measure  $I_{max} = \max(w_{ij})$ .  $I_{max}$  can be defined for each connectivity matrix  $W$  individually, however, as we compare NDIs across groups, we define it as the maximum of  $I_{max}$  across all 196 connectivity matrices.

To determine the NDI of node  $m$ , we first create a subnetwork  $G^{-m}$ , by removing node  $m$ . We subsequently calculate  $(I_{-m}^n)_{ij}$  for all remaining node pairs  $(i,j)$  in  $G^{-m}$ . If removal of node  $m$  results in disconnected components, some path-lengths may become infinite. For such paths, the loss of information measure is set to 1. Finally, each node  $i \in G^{-m}$  is assigned an accumulated loss in information measure

$$\Delta I_i = \sum_j (I^n)_{ij} - (I_{-m}^n)_{ij}.$$

The NDI of node  $m$  is then given as the mean of  $\Delta I_i$  over all nodes  $i$  in  $G^{-m}$ , where the closer the value is to 1, the higher the information loss and subsequently the more important node  $m$  is to the network. This analysis is then repeated for all nodes in the network, resulting in a  $n \times 1$  dimensional feature vector of NDI scores for the network.

## NDI subnetworks

For each group connectome, we calculate its NDI scores. We determine regions of importance by ranking their NDI score from highest to lowest for comparison with RC nodes. Additionally, we model the natural-log-transformed NDI of each group connectome, using a Gaussian Mixture Model with three Gaussian distributions (GMM<sub>3</sub>). Using the halfway point between the Gaussian centers  $\mu_i$ , we define four subnetworks, Tiers 1 to 4, of decreasing NDI importance (Table 2).

*Table 2: Conditions for subnetwork definition into Tiers using NDI.*

Tier	1	2	3	4
NDI condition	$NDI > \frac{\mu_2 - \mu_3}{2}$	$\frac{\mu_1 - \mu_2}{2} < NDI \leq \frac{\mu_2 - \mu_3}{2}$	$0 < NDI \leq \frac{\mu_1 - \mu_2}{2}$	$NDI = 0$

## Statistical analysis

### Network organization by age group

Our first analysis is to investigate the consistency and distributions of NDI scores for each subnetwork in relation to rich-club subnetworks. Consistency of NDI across age groups is subsequently tested using Spearman's rank correlation coefficient. To characterize different

topological aspects of the subnetworks derived from each framework, we calculate three network measures (Rubinov and Sporns, 2010), namely transitivity (T), global efficiency (E) and assortativity (a).

## Age-dependent subnetwork trends within the cohort

We also investigate the associations of each network measure with age in both RC- and NDI-defined subnetworks. In order to make the subnetworks uniform across the age groups, we repeat the RC and NDI analyses without stratifying by age group and instead use the population averaged connectome  $W_{\text{cohort}}$ . Furthermore, we apply the  $W_{\text{cohort}}$  defined subnetworks from both frameworks to each subject's individual connectome for analysis with age. Associations are estimated by fitting a linear model to the subject-level data, given by

$$\text{measure} = a * \text{age} + b,$$

with slope  $a$  and offset  $b$ . All statistical analyses were performed using MATLAB. The code used to calculate the statistics and generate the figures, as well as an implementation for computing NDI is available at <https://github.com/mdschirmer/NDI>.

# Results

## Network organization by age group

Table 3 details the  $k_{\text{max}}^{\text{group}}$  computed from each group-averaged connectome and the number of corresponding regions determined to form the rich-club subnetwork.

*Table 3: Normalized RC analysis of the four age groups, identifying the degree  $k_{\text{max}}^{\text{group}}$ , and the corresponding number of nodes.*

	U20	U40	U60	O60
$k_{\text{max}}^{\text{group}}$	49	48	49	53
# nodes	11	10	14	5

In some cases, the atlas assigns multiple nodes to the same anatomical label. Figure 1 shows the rich-club regions identified for each of our four age groups using  $k_{max}^{group}$  shown in Table 3. It highlights 14 regions as belonging to the rich-club across the life-span cohort, with eight regions appearing in at least three of the four age groups. These regions consist bilaterally of the caudate, thalamus, pallidum, and parahippocampal posterior regions, as well as the right putamen.




Figure 1: Rich-club and the top ten brain regions with highest NDI scores defined from the connectome ( $W_{group}$ ) for each age group.

For NDI, the ten regions with highest score are also indicated in Figure 1. Comparing NDI across the four age groups demonstrates high consistency with an average Spearman's correlation coefficient of  $0.98 \pm 0.01$ . Using the subnetworks defined by the rich-club, we investigate the NDI scores of RC, F and S. Figure 2 summarizes the NDI scores stratified by RC subnetworks and age groups. RC nodes, on average, have the highest NDI scores of the group connectome, with lower scores for F, followed by S regions.






Figure 2: A comparison of NDI scores stratified by RC subnetworks for each age group.

## Subnetwork definition using NDI

We observe that each connectome's NDI follows a mixture of normal distributions after a natural log-transform. Based on the NDI scores within each group connectome, we fit a GMM<sub>3</sub> to each of the distributions (see Figure 3). The average centers of the Gaussians across all four  $W_{\text{group}}$  are determined to be  $-14.88 \pm 0.67$ ,  $-11.21 \pm 0.06$ , and  $-8.24 \pm 0.13$ . The halfway point between consecutive Gaussians ( $-13.05$  and  $-9.73$ ) divide the NDI into three sections. Subsequently, with the inclusion of the  $\text{NDI} = 0$  Tier, each node is assigned to one of the four Tiers according to their NDI value for our analysis.




Figure 3: Natural log-transformed NDI distribution for each age group. Each distribution has a GMM with three Gaussians fitted to it. The centers of these Gaussians are indicated by black diamonds, while the probability density function is shown in orange.

Network theoretical measures computed from the subnetworks derived via both the RC and NDI frameworks are plotted in Figure 4.




Figure 4: Network measures stratified by age group for each of the investigated subnetworks. Top: NDI-based subnetworks following their differentiation in Table 2, with increasing NDI from left to right. Bottom: RC-based subnetworks, stratified by their connectivity profile with respect to the rich-club.

## Age-dependent subnetwork associations within the cohort

Figure 5 shows the regions stratified by subnetworks from  $W_{\text{cohort}}$ , as defined from NDI and RC frameworks. NDI nodes were classified according to Gaussian means -14.76, -10.75 and -7.93 for Tiers 1 to 3, respectively, and RC node were defined with  $k_{\text{max}} = 47$ .




Figure 5: NDI- (left) and RC-derived (right) subnetworks from cohort-averaged connectome. a) Assignments of cortical regions by subnetwork for each framework, where corresponding left and right hemispheric regions were combined into a single region label. b) Brain regions plotted by subnetwork label for each framework.

Figure 6 shows the association of network measures calculated from RC- and NDI-based subnetworks with age, including the linear model. For both frameworks, we see, on average, good separation of each subnetwork investigated in at least one of the three network measures consistently across age. Global efficiency demonstrates the highest separation between subnetworks in both frameworks, with RC and Tier 1 regions having the highest levels of efficiency. We see clear differentiation between regions in the three remaining subnetworks (Tiers 2, 3 and 4) in the NDI-framework, as well as between the two remaining subnetworks in the RC-framework (F and S). This separation between subnetworks can also be observed in terms of transitivity, where Tiers 3 and 4, as well as F and S overlap in part. For assortativity, there is less of an overlap between subnetworks derived from NDI compared to the RC-framework.




Figure 6: Network measures plotted against age for NDI- (top) and RC-based (bottom) subnetworks defined from the full cohort connectome. Linear regressions lines for each network measure and subnetwork are plotted.

Fitted parameters of the linear regressions are summarized in Table 4. All subnetworks show increasing efficiency with age, while assortativity decreases. In the RC-based subnetworks, S decreases in transitivity, as does the Tier 4 subnetwork (NDI=0). All other subnetworks exhibit increasing transitivity with age.

*Table 4: Results of linear regression with slope  $a$  and offset  $b$  for each network measure with age and for RC- and NDI-based subnetworks. Statistically significant regression parameters are in bold ( $p < 0.05$ ).*

	T				E				a			
NDI	1	2	3	4	1	2	3	4	1	2	3	4
a	1.42E-4	2.64E-5	2.56E-5	-1.89E-6	1.63E-4	1.63E-4	1.08E-4	4.70E-5	-3.18E-4	-1.00E-3	-4.09E-4	-1.55E-3
b	8.07E-2	2.88E-2	1.29E-2	1.06E-2	2.19E-1	8.08E-2	2.82E-2	1.09E-2	-5.42E-2	1.40E-1	6.75E-2	2.34E-1
RC	RC	F	S		RC	F	S		RC	F	S	
a	1.40E-4	4.52E-6	-8.39E-5		1.59E-5	1.22E-4	1.40E-4		-4.90E-4	-8.71E-4	-5.08E-4	
b	1.11E-2	1.77E-2	2.79E-2		2.94E-1	6.12E-2	2.57E-2		-4.35E-2	9.77E-2	2.79E-2	

## Discussion

In this work we extended the NDI framework to the analysis of weighted connectomes and utilized the nodal indices to identify four distinct subnetworks in the human connectome. Importantly, our subnetwork definition is data driven, without any manually chosen parameters, and shows higher consistency across age groups, compared to RC-based subnetworks. Investigating NDI, stratified by RC-based subnetworks, revealed generally higher NDI for nodes that also belong to the rich-club. This is to be expected, as the rich-club has been shown to be an integral part for effective information transport in networks. Feeders and seeders have lower NDI, where seeders almost uniformly exhibit an NDI of zero. This is consistent with the notion that these nodes are ‘local’ or ‘peripheral’ in the network. However, we observe that some nodes belonging to feeder regions exhibit higher importance for network functioning, with NDI values that are comparable to rich-club members. The rich-club is often coined as a backbone for information transport in brain

networks. While this aspect has been previously shown, due to the limited assessment of rich-club regions based simply on nodal degree (a localized measure), their importance as a backbone structure is only partially interrogated. NDI, however, investigates the relevance of each region for efficient information transport within the entire brain network by incorporating topological distance information, a more globally informed measure than nodal degree. In doing so, nodal NDI may be more reflective of pathologies or structural changes that implicate entire pathways, making it advantageous oversimply counting the number of edges immediately appended to a node.

We observe that NDI follows a mixture of normal distributions after a natural log-transform. Here, we modeled the natural log-transformed NDI of each group connectome and the cohort connectome using a Gaussian Mixture Model with three distributions. By using the halfway point between Gaussian centers (where the probability of a node belonging to either one of the distributions is equal) we distinguish four different subnetworks. Importantly, the Gaussian center estimations are consistent, whether computed between age groups or from the cohort connectome. Furthermore, NDI demonstrated greater stability in identifying more recurring regions with high NDI values in all four age groups than the RC framework (Figure 1). Stability in identifying regions integral to brain architecture and functioning is desirable for comparative purposes particularly given that they have been similarly detected across species, age, and disease (Ball et al., 2014; Daianu et al., 2015; Grayson et al., 2014; Heuvel and Sporns, 2011; Schirmer and Chung, 2018; van den Heuvel et al., 2013). This has led to a recent uptake in employing *a priori* RC nodes for network analysis (Collin et al., 2016, 2014; van den Heuvel et al., 2013; Wierenga et al., 2018). Utilizing these NDI subnetwork definitions and investigating their network topology with commonly used network measures, we observe distinct patterns for each of the four subnetworks.

Our first analysis investigates age-associated changes in network measures by stratifying the cohort according to age. In both frameworks we observe clear, mostly non-linear patterns in each subnetwork. Similar patterns have been observed previously, e.g. with strength, following an inverse U-shape for RC members (Zhao et al., 2015). However, the membership of each region to a subnetwork (at the RC and Tier 1 level, for example) is not constant with age group in this analysis. While refinement and reassignment of region membership to any subnetwork is possible, comparisons between age groups becomes difficult. In our second experiment, we homogenized the regions that define the subnetworks in both frameworks by utilizing the cohort

connectome. Analyzing the trends of the investigated network measures also showed significant patterns associated with age in each subnetwork for both framework. Specifically, we observe a general increase in efficiency with age, most likely reflecting a refinement of the information transport within each subnetwork. Assortativity shows a general decrease with age and for all subnetworks, indicating that the connections between nodes of different degrees are strengthened. Transitivity demonstrates mixed patterns for individual subnetworks. It generally increases in RC and F subnetworks, as well as in Tiers 1, 2 and 3, indicative of a within-subnetwork strengthening over age. However, for S and Tier 4, transitivity decreases, reflective of less tightly integrated subnetworks. This may be the result of a distribution of the limited resources in the brain, which favors pre-existing, highly integrated regions. These results suggest that our NDI framework defines organizations of age-associated network principles of T, E and a with good separation between subnetworks (Figure 6), indicative of more highly modularized groupings of nodes, when compared with the RC framework, across the ages.

Regarding the functional significance of the NDI-derived subnetworks, it can be clearly noted that the gray matter regions (both subcortical nuclei and cortical regions) follow a unique, largely non-redundant pattern (Figure 5). This is particularly noticeable in separation of the Tier 1 regions from the Tiers 2-4, given that the major components of the Tier 1 subnetwork represent the key relay nuclei in the brain (such as thalamus or the basal ganglia represented by caudate, pallidum, putamen) or cortical regions (such as insular, posterior cingulate gyrus) responsible for processing and redistribution of essential information flow from the primary (motor, sensory, language) cortex to the related association cortices and beyond. Greater overlap appears to exist between the cortical regions represented in Tiers 2-4, which potentially reflects the redundancy necessary for operations involving larger and more complex clusters of gray matter structures. This redundancy may serve to ensure the dynamic connectivity that engages multiple brain functions necessary to sustain defined tasks. Future studies on the functional topography of NDI-derived subnetworks should aim to elucidate the significance and versatility of the tier-specific region interactions, as well as the potential influences that may disrupt or enhance them.

There are some limitations to our study. For our framework, we decided to use three Gaussians to model the distributions of NDI in our groups. While we observe good separation and consistent Gaussian centers across the cohort and four age group connectomes, other choices may be valid. Future studies may investigate different choices of modeling the NDI distribution, potentially leading to a higher number of separable subnetworks. Additionally, other studies have employed

alternative means to investigate versions of the rich club, such as the participation coefficient, community and/or the distribution index (Grayson et al., 2014; Van Den Heuvel and Sporns, 2011). While these network measures are closely related to the NDI, the purpose of our work was not to use NDI as a direct measure of network topology, but to differentiate and identify meaningful subnetworks in the connectome. However, NDI can be utilized as a nodal measure to inform the impact of localized damage, e.g. in conditions such as stroke or brain tumors and to study differences in network topology, which will be the aim of future studies. Limitations in any life-span analysis include defining the age groups in the cohort, and also how to standardize network estimates across subjects and/or age groups. We utilized age groups to investigate potential changes in subnetwork definition based on differences in the group connectome in the first part of the presented analysis. In particular, subjects in the youngest age group (<20 years; U20) include biological changes such as myelination which can modify the structural connectome globally, as part of normal development. The level in which to subdivide by age during periods of neurological development (or aging) is not trivial. Therefore, in this work we did not focus on these developmental stages, warranting dedicated, fine-grained studies in the U20 group.

We have demonstrated our NDI framework to successfully identify highly consistent subnetworks across the life-span which can be used to uncover topological aspects of the human connectome. By investigating topology according to subnetworks, we found distinct patterns of increasing efficiency with age and differential changes in transitivity in relation to nodal importance. The combined stability and sensitivity in detecting distinct subnetworks shows NDI to have great promise for assessing the global implications of a disease in a network across pathologies and developmental time-frames.

## Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 753896.

## References

Ball, G., Aljabar, P., Zebari, S., Tusor, N., Arichi, T., Merchant, N., Robinson, E.C., Ogundipe, E., Rueckert, D., Edwards, A.D., 2014a. Rich-club organization of the newborn human brain. *Proceedings of the National Academy of Sciences* 111, 7456–7461.



- Ball, G., Aljabar, P., Zebari, S., Tusor, N., Arichi, T., Merchant, N., Robinson, E.C., Ogundipe, E., Rueckert, D., Edwards, A.D., Counsell, S.J., 2014b. Rich-club organization of the newborn human brain. *PNAS* 111, 7456–7461. <https://doi.org/10.1073/pnas.1324118111>
- Brown, J.A., Rudie, J.D., Bandrowski, A., Van Horn, J.D., Bookheimer, S.Y., 2012. The UCLA multimodal connectivity database: a web-based platform for brain connectivity matrix sharing and analysis. *Frontiers in neuroinformatics* 6, 28.
- Bullmore, E., Sporns, O., 2009. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience* 10, 186.
- Collin, G., de Nijs, J., Hulshoff Pol, H.E., Cahn, W., van den Heuvel, M.P., 2016. Connectome organization is related to longitudinal changes in general functioning, symptoms and IQ in chronic schizophrenia. *Schizophr. Res.* 173, 166–173. <https://doi.org/10.1016/j.schres.2015.03.012>
- Collin, G., Kahn, R.S., de Reus, M.A., Cahn, W., van den Heuvel, M.P., 2013. Impaired rich club connectivity in unaffected siblings of schizophrenia patients. *Schizophrenia Bulletin* 40, 438–448.
- Collin, G., Kahn, R.S., Reus, D., A, M., Cahn, W., Heuvel, V.D., P, M., 2014. Impaired Rich Club Connectivity in Unaffected Siblings of Schizophrenia Patients. *Schizophr Bull* 40, 438–448. <https://doi.org/10.1093/schbul/sbt162>
- Craddock, R.C., James, G.A., Holtzheimer III, P.E., Hu, X.P., Mayberg, H.S., 2012. A whole brain fMRI atlas generated via spatially constrained spectral clustering. *Human brain mapping* 33, 1914–1928.
- Daianu, M., Jahanshad, N., Nir, T.M., Jack, C.R., Weiner, M.W., Bernstein, M., Thompson, P.M., 2015a. Rich club analysis in the Alzheimer's disease connectome reveals a relatively undisturbed structural core network. *Hum Brain Mapp* 36, 3087–3103. <https://doi.org/10.1002/hbm.22830>
- Daianu, M., Jahanshad, N., Nir, T.M., Jack Jr, C.R., Weiner, M.W., Bernstein, M.A., Thompson, P.M., Initiative, A.D.N., 2015b. Rich club analysis in the Alzheimer's disease connectome reveals a relatively undisturbed structural core network. *Human brain mapping* 36, 3087–3103.
- Fornito, A., Zalesky, A., Breakspear, M., 2013. Graph analysis of the human connectome: promise, progress, and pitfalls. *Neuroimage* 80, 426–444.
- Grayson, D.S., Ray, S., Carpenter, S., Iyer, S., Dias, T.G.C., Stevens, C., Nigg, J.T., Fair, D.A., 2014a. Structural and functional rich club organization of the brain in children and adults. *PLoS one* 9, e88297.

- Grayson, D.S., Ray, S., Carpenter, S., Iyer, S., Dias, T.G.C., Stevens, C., Nigg, J.T., Fair, D.A., 2014b. Structural and functional rich club organization of the brain in children and adults. *PLoS ONE* 9, e88297. <https://doi.org/10.1371/journal.pone.0088297>
- Hagmann, P., Cammoun, L., Gigandet, X., Meuli, R., Honey, C.J., Wedeen, V.J., Sporns, O., 2008. Mapping the structural core of human cerebral cortex. *PLoS biology* 6, e159.
- Heuvel, M.P. van den, Sporns, O., 2011. Rich-Club Organization of the Human Connectome. *J. Neurosci.* 31, 15775–15786. <https://doi.org/10.1523/JNEUROSCI.3539-11.2011>
- Mori, S., Crain, B.J., Chacko, V.P., Van Zijl, P.C., 1999. Three-dimensional tracking of axonal projections in the brain by magnetic resonance imaging. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society* 45, 265–269.
- Nooner, K.B., Colcombe, S., Tobe, R., Mennes, M., Benedict, M., Moreno, A., Panek, L., Brown, S., Zavitz, S., Li, Q., 2012. The NKI-Rockland sample: a model for accelerating the pace of discovery science in psychiatry. *Frontiers in neuroscience* 6, 152.
- Opsahl, T., Colizza, V., Panzarasa, P., Ramasco, J.J., 2008. Prominence and control: the weighted rich-club effect. *Physical review letters* 101, 168702.
- Ray, S., Miller, M., Karalunas, S., Robertson, C., Grayson, D.S., Cary, R.P., Hawkey, E., Painter, J.G., Kriz, D., Fombonne, E., 2014. Structural and functional connectivity of the human brain in autism spectrum disorders and attention-deficit/hyperactivity disorder: a rich club-organization study. *Human brain mapping* 35, 6032–6048.
- Rubinov, M., Sporns, O., 2010. Complex network measures of brain connectivity: uses and interpretations. *Neuroimage* 52, 1059–1069.
- Schirmer, M.D., Chung, A.W., 2018a. Structural subnetwork evolution across the life-span: rich-club, feeder, seeder, in: *International Workshop on Connectomics in Neuroimaging, LNCS*. Springer, pp. 1–10. [https://doi.org/10.1007/978-3-030-00755-3\\_15](https://doi.org/10.1007/978-3-030-00755-3_15)
- Schirmer, M.D., Chung, A.W., 2018b. Structural subnetwork evolution across the life-span: rich-club, feeder, seeder, in: *Connectomics in Neuroimaging Workshop*. Presented at the MICCAI, LNCS, Springer, pp. 134–143. [https://doi.org/10.1007/978-3-030-00755-3\\_15](https://doi.org/10.1007/978-3-030-00755-3_15)
- van den Heuvel, M.P., Kahn, R.S., Goñi, J., Sporns, O., 2012. High-cost, high-capacity backbone for global brain communication. *Proceedings of the National Academy of Sciences* 109, 11372–11377.
- Van Den Heuvel, M.P., Sporns, O., 2011. Rich-club organization of the human connectome. *Journal of Neuroscience* 31, 15775–15786.

- van den Heuvel, M.P., Sporns, O., Collin, G., Scheewe, T., Mandl, R.C.W., Cahn, W., Goñi, J., Hulshoff Pol, H.E., Kahn, R.S., 2013. Abnormal rich club organization and functional brain dynamics in schizophrenia. *JAMA Psychiatry* 70, 783–792.  
<https://doi.org/10.1001/jamapsychiatry.2013.1328>
- Wierenga, L.M., van den Heuvel, M.P., Oranje, B., Giedd, J.N., Durston, S., Peper, J.S., Brown, T.T., Crone, E.A., The Pediatric Longitudinal Imaging, Neurocognition, and Genetics Study, 2018. A multisample study of longitudinal changes in brain network architecture in 4-13-year-old children. *Hum Brain Mapp* 39, 157–170.  
<https://doi.org/10.1002/hbm.23833>
- Woldeyohannes, Y.T., Jiang, Y., 2018. Measures for Network Structural Dependency Analysis. *IEEE Communications Letters*.
- Zhao, T., Cao, M., Niu, H., Zuo, X.-N., Evans, A., He, Y., Dong, Q., Shu, N., 2015. Age-related changes in the topological organization of the white matter structural connectome across the human lifespan. *Human brain mapping* 36, 3777–3792.