

Sleep-like slow oscillations induce hierarchical memory association and synaptic homeostasis in thalamo-cortical simulations

Cristiano Capone¹, Elena Pastorelli^{1,2}, Bruno Golosio^{3,4}, and Pier Stanislaw Paolucci^{1,*}

¹INFN Sezione di Roma

²PhD Program in Behavioural Neuroscience, “Sapienza” University of Rome

³Dipartimento di Fisica, Università di Cagliari

⁴INFN Sezione di Cagliari

*Corresponding author: Pier Stanislaw Paolucci, pier.paolucci@roma1.infn.it

Abstract

The occurrence of sleep is widespread over the large majority of animal species, suggesting a specific evolutionary strategy. The activity displayed in such a state is known to be beneficial for cognitive functions, stabilizing memories and improving the performances in several tasks. Despite this, a theoretical and computational approach to achieve the understanding of such mechanism is still lacking. In this paper we show the effect of sleep-like activity on a simplified thalamo-cortical model which is trained to encode and retrieve images of handwritten digits. We show that, if spike-timing-dependent-plasticity (STDP) is active during sleep, the connections among groups of neurons associated to instances of the same class (digit) are enhanced, and that the internal representation is hierarchical and orthogonalized. Such effect might be beneficial to the network to obtain better performances in retrieval and classification tasks and to create hierarchies of categories in integrated representations. The model leverages on the coincidence of top-down contextual information with bottom-up sensory flow during the training phase and on the integration of top-down predictions and bottom-up pathways during deep-sleep-like slow oscillations.

Authors’ summary

A simplified thalamo-cortical model is trained with examples drawn from the MNIST set of handwritten digits, while receiving simultaneous bottom-up sensory stimulation and selective lateral sub-threshold activation. During a training phase, spike-timing-dependent-plasticity (STDP) sculpts a pre-sleep synaptic matrix which associates training examples with well separated groups of cortical neurons and creates top-down synapses toward the thalamus. Then, the network is induced to produce deep-sleep-like slow oscillations (SO), disconnected from sensory and lateral stimuli, and driven by its internal activity. During sleep up-states, thalamic cells are activated by top-down predictive stimuli produced by cortical neural groups and respond with a forward feedback, recruiting other cortical neurons. This induces two simultaneous and beneficial effects: a reduction in the strength of synapses inside the groups created by the training and an increase of synapses that associate examples of characters belonging to the same digit category. The fine structure of the synaptic matrix reflects the categorization of examples and a hierarchical structure within each category. During post-sleep retrieval activity the correlation between neurons reflects the richer hierarchical structure of the underlying synaptic matrix.

1 Introduction

Human brains spend about one-third of their life-time sleeping. Sleep is present in every animal species that has been studied (G. Tononi, C. Cirelli [1]), notwithstanding two negative facts: the danger caused by sleep, that diminishes the capability to defend from predators and other threats, and the reduction of time available for activities targeting immediate rewards (e.g hunting or gathering food). Having survived the evolutionary selection in all species, sleep must therefore provide strong advantages. A notable fact is that newborns’ human brains occupy the majority of their time asleep, nevertheless they learn at a very fast rate. For this and other motivations for studying sleep, also in relation to consciousness, see e.g. [2].

Even if occasional awakening does not seriously impair brain biology and cognitive functions, in the long term chronic deprivation of sleep produces measurable effects on cognition, mood and health [3].

Here we present a simple thalamo-cortical simulation model that incurs interesting modifications induced by deep-sleep-like activity: simultaneous hierarchical association of memories and differential synaptic homeostasis. In the retrieval and training phases, the model is stimulated by bottom-up sensory features and top-down selection stimuli. In the sleep phase, it runs according to top-down and bottom-up mechanisms that are internal to the model.

1.1 Experimental and theoretical cues

The proposed model and the observed effects are compatible with proposals coming from a few experimental facts and theoretical cues. We mention them hereafter.

A first guideline is the "organizing principle" proposed by M. Larkum in [4]. It is about the general organization of the cerebral cortex architecture and about specific cellular mechanisms supporting this architectural scheme, implemented on pyramidal neurons populating Layer 5. The architecture is grounded on the separation of intra-areal and inter-areal contextual information (reaching the apical dendrites of pyramidal neurons) from the feed-forward flow of area specific information targeting its basal synapses. Cellular mechanisms, like Ca^{++} spikes, promote the detection of coincidence between contextual and specific activity. High-frequency bursts of spikes are emitted when the coincidence is detected.

Relying on these features we introduced external lateral stimuli in our thalamo-cortical model. This way, the model changes the perceptual threshold of specific subsets of neurons during the presentation of examples in the training phase. For each example, a vector of features is computed and is projected toward a cortical network by a thalamic neural network. Due to the change in the perceptual threshold, spike-timing-dependent-plasticity (STDP) creates stronger bottom-up (thalamo-cortical) and top-down (cortico-thalamic) connections between a subset of cortical and the thalamic neurons. The subset is specific for the current training example. We used examples extracted from the MNIST set of handwritten digits, with features computed using a histogram of oriented gradients (HOG) descriptor. For an easier control, the sub-threshold activation induced by predictive connections is delivered to disjoint groups of cortical neurons (one group per training example) and it is set to an identical value for all neurons and examples.

A second class of leading ideas derive from the proposed role of sleep in the predictive processing framework (see [5]). Bucci and Grasso make hypotheses about the peculiarity of prediction in sleep and discuss the role of sleep in relation with the Integrated Information Theory of consciousness [6]. They notice that during sleep the external perceptions are, at least, strongly attenuated, and the majority of the motor system is blocked. This suggests the need for a different mechanism enabling the brain to perform prediction during sleep, compared to those used when awake.

Along this line, during the sleep stage we switched off in our model both lateral inputs and sensory stimulation and used a non specific stimulation. We follow the hypotheses that predictive backward activation, projected from the cortex to the thalamus, might have a beneficial role in the computational performances. Indeed, they supported hierarchical associations between memories of training examples during sleep-like oscillations.

We considered highly plausible the search of a role of sleep as an off-line mechanism of multi-objective synaptic optimization, in addition to biological restoration (see [1, 7]). A first motivation for the brain disconnecting from the external world would be to set a better energy consumption regime during next wakefulness cycle. This might be obtained by reducing firing rates and the amplitude of evoked post-synaptic potentials. It could be reached by pruning non necessary synapses and by reducing synaptic weights within the limits imposed by the conservation of adequate coding for memories. An ideal off-line optimization could also associate existing memories, create novel conceptual hierarchies and enable the computation of additional logical predicates. In contrast with the energy-related optimization objective, this would generally require the strengthening of specific synapses.

In synthesis, we designed a minimal biologically inspired computational model in order to quantitatively explore the dynamics of memories during sleep and of network homeostasis mechanisms. This study also paves the way to include plasticity in large scale simulation of Slow Wave Activity (SWA) on cortical areas with biologically plausible long-range lateral interconnects (see Pastorelli et al, [8] based on a columnar description), to extend our observations from SO to SWA. The long term goal is to compare simulation results with experimental measures of the effects of different stages of sleep on neural circuits.

Also would be interesting to test and compare different hypotheses of sleep functions which are not yet unified in a coherent framework [1, 7]. We hope to contribute to refine these hypotheses: they could find a conciliation path according to our preliminary simulation results, that point to the possibility of having SO promoting both a hierarchical association of memories and differential synaptic homeostasis mechanisms.

1.2 Focus on the effect of sleep-like activity on a trained network

In this work, we consider the modeling of the effects of deep-sleep-like SO induced after a training performed with the network on the MNIST dataset of handwritten digits.

SO are considered the default emergent activity of the cortical network (see Sanchez-Vives, Massimini, Mattia [9]) and are observed during the deepest of physiological non-REM sleep stages as an alternation between *down-states* (characterized by nearly silent neurons) and *up-states* (in which a subset of neurons goes in a high firing rate regime) occurring at a frequency in the range [0.5, 4] Hz (*delta* band). We set the model SO at a comparable frequency.

SO activity is expected to play two complementary roles, which are separately mediated by up-states and down-states. Down-states would play a purely biological function, with a lower immediate impact on cognitive performance. The role of down-states, with a majority of neurons put in a silent state for a long fraction of deep sleep time, would serve a restoration purpose, enabling periodic biological maintenance and recovery, as it happens in the whole body when at rest [10].

Our modeling and investigation is focused only on the effects mediated by the up-states dynamics. By recruiting population of neurons firing at high rates, up-states can substantially modify synaptic weights. Measurable after-sleep effects can be observed on the structure of the synaptic matrix, on the correlation between the activity of neurons, on the computational properties of the network and on the energy consumption, that depends on the firing rates and on the post-synaptic evoked potentials.

Cognitive effects of chronic deprivation are probably due to the lack of positive effects of both down-states (biological restoration) and up-states (optimization of the working point of the networks and of its interconnect).

2 Methods

Network architecture The network, which is designed to be a minimal thalamo-cortical model [11] (see Fig.1), is composed of two populations of cortical neurons (one excitatory CX and one inhibitory IN) and two thalamic populations (one excitatory TH and one inhibitory RE). The connection probability is $p = 1.0$ for the populations connected by the arrows in Fig.1. We define *lateral prediction* as a signal which selectively facilitates neurons to learn new stimuli and which is assumed to come from same (or higher) order cortices, while *top-down prediction* is the signal flowing through $CX \rightarrow TH$ connections.

AdEx The neurons in the network are adaptive exponential (adEx) point like single compartment neurons.

$$\begin{cases} \frac{dV(t)}{dt} = -\frac{V(t)-E_l}{\tau_m} + \frac{\Delta V}{\tau_m} e^{\left(\frac{V(t)-\theta}{\Delta V}\right)} + \frac{I(t,V(t))}{C} - \frac{W(t)}{C} \\ \frac{dW(t)}{dt} = -\frac{W(t)}{\tau_w} + b \sum_k \delta(t-t_k) + a(V(t) - E_l) \end{cases} \quad (1)$$

where the synaptic input I is defined as

$$I(t, V(t)) = \sum_{\alpha} g_{\alpha}^{syn}(t)(V(t) - E_{\alpha}^{syn}) \quad (2)$$

where $V(t)$ is the membrane potential of the neuron and $\alpha = e, i$ defines the excitatory (e) and the inhibitory (i) input. The population dependent parameters are: τ_m the membrane time constant, C the membrane capacitance, E_l the reversal potential, θ the threshold, ΔV the exponential slope parameter, W the adaptation variable, a and b the adaptation parameters, and g_{α}^{syn} the synaptic conductance, defined as




Figure 1: **Thalamo-cortical model description.** Sketch of the structure of the simplified thalamo-cortical model considered, which is composed of an excitatory and an inhibitory population both for the cortex (CX, IN) and for the thalamus (TH,RE). Connectivity structure is represented by solid lines. The visual input is fed into the model through the thalamic population, mimicking the biological visual pathways. In the training phase a lateral stimulus enhances a specific subset of CX neurons to preferentially represent the stimulus.

$$g_{\alpha}^{syn}(t) = \sum_k \Theta(t - t_k) Q_{\alpha} \exp(-(t - t_k)/\tau_{\alpha}^{syn}) \delta(t - t_k) \quad (3)$$

We define the spiking time of the neuron when the membrane potential reaches the threshold $V_{spike} = \theta + 5\Delta V$. t_k^{α} indicates the times of pre-synaptic spikes received by the neuron from synapse type α with characteristic time τ_{α}^{syn} and its synaptic efficacy Q_{α} .

STDP plasticity We used spike-timing-dependent-plasticity (STDP) which potentiates w_{ij} when spike occurs earlier in neuron j (spike time t_j) than in neuron i (spike time t_i) and viceversa. We considered STDP in its multiplicative form [12–14] which is described by the following equation

$$\begin{cases} \Delta w_{ij} = \lambda(w_{max} - w_{ij})e^{-|t_i - t_j|/\tau} & \text{potentiation} \\ \Delta w_{ij} = -\alpha\lambda w_{ij}e^{-|t_i - t_j|/\tau} & \text{depression} \end{cases} \quad (4)$$

where w_{max} is the maximum weights value, α is the asymmetry parameter between potentiation and depression, λ is the learning rate and τ is the STDP timescale.

Network size and training set The number of thalamic (TH) neurons are the same as the dimensionality of the visual input ($N_{TH} = 324$), while the number of cortical neurons is 20 times the number of images used to train the network ($N_{CX} = 180$ in this paper).

Indeed, the training set is composed of 9 elements, that is, three examples of images for three different classes of handwritten digits.

The number of inhibitory neurons are $N_{IN} = 200$ and $N_{RE} = 200$.

Training algorithm Every time a new training image is presented to the network through the thalamic pathway, the facilitation signal coming from lateral prediction, induces selectively a new set of 20 neurons to encode for that specific stimulus. The learning mechanism is allowed by symmetric ($\alpha = 1.0$) STDP plasticity present in the $CX \rightarrow CX$, $CX \rightarrow TH$ and $TH \rightarrow CX$ connections which shapes the weights structure.

Slow oscillations After the training stage, the sleep-like slow oscillations activity is induced by providing a non-specific stimulus inside the cortex and increasing the strength of SFA parameter ($b = 60$, in eq.(1)). In this stage asymmetric STDP plasticity ($\alpha = 3.0$) is active in the recurrent CX connectivity, inducing sleep-induced modification in the synaptic weights structure.

Incremental training The architecture described in previous section is an evolution of the spiking network model [15]. In that model, lateral prediction projected toward cortical neurons selected groups of target neurons during an incremental training of MNIST handwritten characters. The training presented 25 novel characters per training step (see Fig. 2). After each training step, we measured the growth in the recognition rate. The network reached a recognition rate of 92% after the presentation of 350 example characters (i.e. 35 examples per digit). The coincidence of bottom-up thalamo-cortical input and lateral predictions drove STDP in sculpturing the connectivity matrix that supported the recognition. We stress again that in the sleep phase the network dynamics is driven by internal activity, spontaneously generated by the cortex. The thalamic populations participates to the SO activity thanks to the top-down cortico-thalamic connections that we included in the architecture.




Figure 2: **Incremental training without slow oscillation.** In this simplified model, thalamo-cortical synapses grow when cortical neurons also receive a simultaneous lateral/top-down facilitation. Cortico-thalamic top-down predictions that revealed to be essential during sleep were still missing in the previous model. Notwithstanding this missing feature, 92% recognition rate was achieved after the exposure to 35 examples per digit.

Simulation engine We used the NEST simulation engine, the high-performance general purpose simulator developed by the NEST Initiative, release 2.12 [16].

Model revision The release associated to the software discussed in this paper is stored in the APE Parallel/Distributed Computing lab svn repository, version n. 1835, date 23-10-2018.

3 Results

In Fig. 3 is reported an example of activity time-course in the CX and TH populations during the training phase, retrieval phase, and the early stage of sleep phase.

Looking at the activity (Fig. 3) and at the structure of the pre-sleep matrix (Fig. 4A) we observe that, during the initial stages of simulated SO, up-states are sustained by those neural groups associated with higher internal interconnections during the training phase. The consequent dynamics can be understood along the following lines. Each cortical group preferentially recruits the group of thalamic neurons from which it received the stronger stimulation during training. This happens through the set of top-down cortico-thalamic connections learned during the training. In turn, thalamic neurons preferentially activate other cortical neurons sharing similarities in the set of training features. We notice that, during the simulated sleep activity, this happens through a bottom-up activation that is not related to the incoming of sensory stimulation. The activation of neural groups associated to different training examples induces, through STDP, not only a synaptic pruning but also the creation of stronger synapses between groups of neurons that share enough commonality in the features they received during the training.

The introduction of lateral stimulation raises the perceptive thresholds of selected groups of neurons during the pre-sleep training phase. The induction of sustained activity in neurons that received both thalamic stimuli and selective lateral prediction enabled the sculpture of well defined groups of synapses associated to individual training examples.




Figure 3: **Sketch of cortico-thalamic activity.** Activity produced during training phase, recognition and SO in the CX (top) and TH (bottom) populations.

The structure of the synaptic matrix before and after sleep is reported in Fig. 4A-B, where is showed the morphing from the initial flat structure (which reflects the individual training examples) towards a hierarchical structure (embedding the categories of learned digits).




Figure 4: **SO effects on connectivity structure.** **A.** Synaptic weights matrix of the recurrent connectivity of CX population, before (left) and after (right) the occurrence of sleep-like activity. The yellow squares represent high weights emerged between neurons encoding the visual input related to the same object (single instance of 0,1,2 ... image). Red solid lines separate the neurons encoding visual inputs related to different classes of objects (0,1,2 ...). **B** Scatter-plot of the same synaptic weights before and after sleep. **C** Synaptic weights after sleep, separated in three groups, synapses between neurons encoding the same object (yellow), the same class (but not the same object, red) and different classes (blue).

The birth of novel synapses among examples in the same digit category and the simultaneous homeostatic reduction of synapses linking neurons grouped by the single training example is clearly visible in

Fig. 4A-left (potentiation) and Fig. 4A-right (depression). The differential effect on synapses can also be observed in Fig. 4C where the histogram of synaptic weights after-sleep is reported.

The changes in the structure of the synaptic weights matrix induces changes in the activity expressed by the network in the retrieval phase, which can be appreciated analyzing the change in the correlation matrix between neural populations before and after sleep (see Fig. 5A).

Fig. 5B reports the difference of correlation evaluated after and before sleep, showing the decorrelation (blue regions) of the populations encoding different classes, and the correlation (red regions) of the ones coding the same class. Such information is reported also in Fig. 5C, where are shown the correlation changes for populations in the same class (blue) and in different classes (green). Such effect might provide benefits in retrieval and classification tasks.




Figure 5: **SO effects on internal representation.** **A.** Activity correlation between all pairs of populations representing the single images before (left) and after (right) sleep. **B.** Correlation difference between after and before sleep. **C** Histogram of differences for populations encoding the same class (blue) and different classes (green).

4 Discussion

In the study described by this first report, we use the model to simulate the effect of deep sleep-like slow oscillations (SO) on the structure of the synaptic matrix and of the neuron groups created by the training.

In our simulations, SO induce two beneficial and apparently contradictory effects. The first is a reduction in the strength of synapses inside the groups created by the training, the second a simultaneous increase of synapses that associate examples of characters belonging to the same category of digits.

As a result, the final structure of the synaptic matrix is more complex and hierarchically organized. Moreover, the correlation of neural groups during post-sleep retrieval activity reflects the hierarchical structure supported by the underlying synaptic matrix, with cooperation between neurons trained by different examples. Also, during the sleep phase, the network displays a rich internal dynamics that evolves across sleep time. Indeed, the composition of neuronal groups recruited during the up states changes sensibly between the initial and the final stages of sleep.

5 Acknowledgments

This work has been supported by the European Union Horizon 2020 Research and Innovation program under the FET Flagship Human Brain Project (SGA2 grant agreement SGA2 n. 785907), System and Cognitive Neuroscience subproject, WaveScaleS experiment. This work is part of the activities performed by the INFN APE Parallel/Distributed Computing laboratory. We would like to thank the members of the APE lab for their strenuous support, and critical discussions: Roberto Ammendola, Andrea Biagioni, Fabrizio Capuani, Paolo Cretaro, Giulia De Bonis, Ottorino Frezza, Alessandro Lonardo, Francesca Lo Cicero, Michele Martinelli, Luca Pontisso, Francesco Simula and Piero Vicini. We also thank for their comments Marco Celotto, Chiara De Luca and Paolo Muratore.

References

- [1] G. Tononi and C. Cirelli, “Sleep and the price of plasticity: From synaptic and cellular homeostasis to memory consolidation and integration,” *Neuron*, vol. 81, no. 1, pp. 12 – 34, 2014.
- [2] G. Tononi and al., “Center for sleep and consciousness - research,” October 2018.
- [3] W. D. Killgore, “Effects of sleep deprivation on cognition,” in “*Human sleep and cognition*” (G. A. Kerkhof and H. P. van Dongen, eds.), vol. 185 of *Progress in Brain Research*, pp. 105 – 129, Elsevier, 2010.
- [4] M. Larkum, “A cellular mechanism for cortical associations: an organizing principle for the cerebral cortex,” *Trends in Neurosciences*, vol. 36, no. 3, pp. 141 – 151, 2013.
- [5] A. Bucci and M. Grasso, “Sleep and dreaming in the predictive processing framework,” in *Philosophy and Predictive Processing* (T. K. Metzinger and W. Wiese, eds.), ch. 6, Frankfurt am Main: MIND Group, 2017.
- [6] G. Tononi and C. Koch, “Consciousness: here, there and everywhere?,” *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 370, no. 1668, 2015.
- [7] B. Watson, D. Levenstein, J. Greene, J. Gelinias, and G. Buzsáki, “Network homeostasis and state dynamics of neocortical sleep,” *Neuron*, vol. 90, no. 4, pp. 839 – 852, 2016.
- [8] E. Pastorelli, P. S. Paolucci, F. Simula, A. Biagioni, F. Capuani, P. Cretaro, G. D. Bonis, F. L. Cicero, A. Lonardo, M. Martinelli, L. Pontisso, P. Vicini, and R. Ammendola, “Gaussian and exponential lateral connectivity on distributed spiking neural network simulation,” in *2018 26th Euromicro International Conference on Parallel, Distributed and Network-based Processing (PDP)*, vol. 00, pp. 658–665, Mar 2018.
- [9] M. V. Sanchez-Vives, M. Massimini, and M. Mattia, “Shaping the default activity pattern of the cortical network,” *Neuron*, vol. 94, no. 5, pp. 993 – 1001, 2017.
- [10] V. V. Vyazovskiy and K. D. Harris, “Sleep and the single neuron: the role of global slow oscillations in individual cell rest,” *Nature Reviews Neuroscience*, vol. 14, pp. 443 – 451, 2013.
- [11] A. Destexhe, “Self-sustained asynchronous irregular states and up-down states in thalamic, cortical and thalamocortical networks of nonlinear integrate-and-fire neurons,” *Journal of computational neuroscience*, vol. 27, no. 3, p. 493, 2009.
- [12] R. Güttig, R. Aharonov, S. Rotter, and H. Sompolinsky, “Learning input correlations through non-linear temporally asymmetric hebbian plasticity,” *Journal of Neuroscience*, vol. 23, no. 9, pp. 3697–3714, 2003.
- [13] W. Gerstner, R. Ritz, and J. L. Van Hemmen, “Why spikes? hebbian learning and retrieval of time-resolved excitation patterns,” *Biological cybernetics*, vol. 69, no. 5-6, pp. 503–515, 1993.
- [14] L. Abbott and S. Song, “Temporally asymmetric hebbian learning, spike limiting and neural response variability,” in *Advances in neural information processing systems*, pp. 69–75, 1999.
- [15] E. Pastorelli, C. Capone, G. Stegel, B. Golosio, and P. S. Paolucci, “Incremental training of spiking neural network with contextual prediction - in preparation,” *ArXiv*, vol. , p. , 2018.
- [16] S. e. a. Kunkel, “Nest 2.12.0,” Mar. 2017.