

# A stochastic modeling framework for single cell migration: coupling contractility and focal adhesions

Aydar Uatay

Technische Universität Kaiserslautern, Felix-Klein-Zentrum für Mathematik  
Paul-Ehrlich-Str.31, 67663 Kaiserslautern, Germany  
(uatay at mathematik.uni-kl.de)

September 3, 2022

## Abstract

The interaction of the actin cytoskeleton with cell-substrate adhesions is necessary for cell migration. While the trajectories of motile cells have a stochastic character, investigations of cell motility mechanisms rarely elaborate on the origins of the observed randomness. Here, guided by a few fundamental attributes of cell motility, we construct a minimal stochastic cell migration model from ground-up. The resulting model couples a deterministic actomyosin contractility mechanism with stochastic cell-substrate adhesion kinetics, and yields a well-defined piecewise deterministic process. Numerical simulations reproduce several experimentally observed results, including anomalous diffusion, tactic migration, and contact guidance. This work provides a basis for the development of cell-cell collision and population migration models.

**Keywords:** cell motility, focal adhesions, piecewise deterministic process, superdiffusion, tactic migration, cell contractility.

**AMS Classification:** 92B05, 92C05, 92C10, 92C17, 60J25.

## 1 Introduction

Cell migration is essential for embryogenesis, wound healing, immune surveillance, and progression of diseases, such as cancer metastasis. During embryogenesis, coordinated collective migration to particular sites is essential to the development of an organism. Tissue repair and immune response rely on directed movement of cells following external cues, which are produced after the tissue layer is damaged and infected with pathogens. Likewise, cancer cells migrate away from the tumor into surrounding tissue and distant organs to form metastases, which is the leading cause of death among cancer patients.

Cell motility is a cyclical process, involving morphological changes of the cell body and adhesive contacts with the underlying substrate [33]. The cycle can be divided into three

steps: protrusion of the leading edge, assembly of adhesions to the substrate at the front and disassembly in the rear, and contraction of the cell body, thereby producing locomotion [1]. This type of crawling movement requires transmission of contractile forces, generated within the cell cytoskeleton, through substrate adhesions. Such mechanical interactions between various cellular structures is attained by integrating numerous signaling molecules, the most prominent of which are Rho GTPases [51], [54]. While there are other modes of motility relying on, for example, flagellar activity (e.g. spermatazoa or *E.coli*) or rolling in the bloodstream (e.g. leukocytes), here the focus is on the crawling type of movement.

Due to cell motility being a highly complex and not fully elucidated process, mathematical modeling of the corresponding phenomena is a challenging task. Numerous approaches have been developed, each being able to capture a certain aspect of the process (see [28], [70] for extensive reviews on whole cell motility models and [41] for a review on modeling of its critical components). For example, free boundary and phase-field models of steadily migrating cells in [9], [46], [56], [58] are able to reproduce cell morphology as a result of mechanical and biochemical interactions. In their hybrid motility model, Marée et al. [38], [39] explored the mechanochemical interaction in detail by considering the Rho GTPase signaling circuit. A common feature of these continuum models is that they are not able to reproduce experimentally observed random paths of migrating cells. Stochastic motility models of eukaryotic cells have been proposed, for example, in [16], [26], where cell-substrate interactions have also been considered. There stochasticity is driven by a Gaussian process, although there is evidence that the paths of the migrating cells follow a non-Gaussian process [17]. Another way to include randomness is based on velocity jump process, as proposed by [47] in the context of cell motility. Extending the model and including cell-substrate interactions, population migration models have been proposed in [18], [31]. Here, however, our focus is solely on single cell migration.

In this paper and the follow-up works, we strike a middle ground. By proposing a minimal cell representation including a few cellular structures essential for cell motility, we aim at reproducing stochastic migratory paths in various experimental settings. In our model, stochasticity arises as a result of mechanochemical coupling between the cell cytoskeleton and the substrate through adhesive contacts. Specifically, the random events under consideration are the (de)adhesion events of the cell migration cycle, whereas deterministic locomotion and contraction occur between arrival of the events. Here, we do not make any prior assumptions about the distributions that the events and their arrival times follow. Rather, we only consider major determinants of an arbitrary adhesion site dynamics. Thereby, the construction of our cell motility model will result in a piecewise deterministic process [13], [14]. Our simulations reproduce experimental observations, such as superdiffusive scaling of the squared displacement [17], [36], [37], biased migration in the presence of an external cue, contact guidance [52], and directed movement due to asymmetric contractility (and in the absence of guidance cues) [65], [68]. Moreover, due to the minimal character of the model and our bottom-up approach, this work serves as a basis for modeling cell-cell collisions and population migration, which will be presented in follow-up papers.

This article is organized as follows. In Section 2 we briefly describe the cell motility

cycle and the relevant agents. We then introduce the minimal cell representation and describe the deterministic cell motion between the cycle steps. In Section 3 we construct a stochastic model of adhesion events, which signify transitions of cycle stages. In Section 4 we combine the deterministic and stochastic components of the migration cycle to obtain a well-defined piecewise deterministic Markov process. In Section 5 we specify the kinetics of adhesion events. Numerical simulations are performed in Section 6. A discussion of the results and future outlook are presented in Section 7.

## 2 The cell motility model




Figure 1: (a) Schematic diagram of the cell migration cycle and the implicated cellular structures. Actin polymerization at the front pushes the membrane allowing protrusions to form. Then, adhesions assemble at the front and disassemble at the rear. Finally, deadhesion and cell contraction produce locomotion, pulling the body forward. The black arrows overlaying the stress fibers show the inwardly directed contractile forces. (b) Schematic representation of a cell with  $M = 8$  focal adhesions. Solid black lines represent stress fibers while red circles represent focal adhesions. Red arrows indicate the direction and magnitude of applied traction force  $\mathbf{F}_i$ ,  $i = 1, \dots, 8$ . Dashed line and corresponding red circle represent an absent stress fiber and unbound focal adhesion, respectively. The central red arrow indicates the net force  $\mathbf{F}$  on  $\mathbf{x}_n$ .

The cell migration cycle begins with protrusion of the leading edge as a result of actin polymerization (Figure 1a). The polymerization process in lamellipodia is mediated by the Arp2/3 complex, which acts downstream of signaling pathways responsible for cell polarization [54]. Next, the protrusions are stabilized due to formation of focal adhesions

(FAs) in the lamellae (regions behind the lamellipodia), which link the actin cytoskeleton to the extracellular matrix (ECM). A FA is a multiprotein integrin-based adhesion cluster, which mature in a Rho GTPase dependent manner [51]. Furthermore, FA maturation depends on the applied tension and occurs concomitantly with actomyosin bundle formation [22]. These bundles, called stress fibers (SFs), generate contractile forces due to non-muscle myosin II motors. Due to increased tension in cell rear, FAs rupture. Finally, deadhesion leads to cell body translocation due to cytoskeletal contraction.

In order to construct the mathematical model, we make the following observations. First, FA unbinding leads to remodeling of the SF configuration (and of the entire cytoskeleton) and cell movement, whereas assembly of new FAs leads to restructuring only. Second, FA events need not occur in the order described above. Several adhesions might be assembled (disassembled) before deadhesion (adhesion) occurs. Note also that while the contractile machinery is important, the dynamic instability of adhesions is what drives the migratory process, for stable FAs prevent retraction. Thus, we consider only interactions of SFs and FAs. Moreover, we do not consider the actin polymerization process and simplify the migration cycle to two steps: after FA assembly occurs, a cell does not move, but reconfigures SFs; after disassembly, a cell does both. Neglecting the polymerization process and the reduction to binding/unbinding events can be justified by the fact that one of the major consequences of the leading edge protrusions is promotion of FA assembly. Because repolarization of migrating cells occurs frequently as an outcome of intricate biochemical activity, then, in order to keep the model tractable, we do not explicitly model cell polarity. Instead, (de)adhesion frequency is indicative of (rear)front.

## 2.1 The cell representation

Consider the situation in the Figure 1b. The disk represents a cell. Let the radius be  $R_{cell}$  and let the position of the center at time  $t$  be  $\mathbf{x}(t) \in \mathbb{R}^2$ . Suppose there are  $M$  equally spaced adhesion sites  $\mathbf{x}_i(t) \in R_{cell}\mathbb{S}^1$ ,  $i = 1, \dots, M$  on a cell circumference with constant relative distance. Let  $\mathbf{Y}(t) \in \{0, 1\}^M$  be a vector of focal adhesion states at time  $t$ , i.e.  $Y_i(t) = 0, 1$  correspond to unbound and bound FA at node  $i$ , respectively.

Since the traction stresses are oriented inward, transmitted to ECM by FAs, and generated by contractile SFs, then the FAs on the circumference must be one of the ends of SFs. Suppose the other end of all SFs at time  $t$  is at the position  $\mathbf{x}_n(t) \in \Omega_{cell} := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 \leq R_{cell}^2\}$  (in a cell's reference frame with origin at  $\mathbf{x}$ ), i.e. all SFs are connected at  $\mathbf{x}_n$ . Since stress fibers behave like Hookean springs on extension, but readily buckle under compression [44], then, inspired by Guthardt Torres et al. [27], the force  $\mathbf{F}_i$  at focal adhesion  $i$  is given by:

$$\mathbf{F}_i = \begin{cases} \left(T_i + EA \frac{L_i - L_0}{L_0}\right) \mathbf{e}_i, & L_0 < L_i \\ T_i \mathbf{e}_i, & L_c \leq L_i \leq L_0 \\ \frac{L_i - L_c + \delta}{\delta} T_i \mathbf{e}_i, & L_c - \delta \leq L_i < L_c \end{cases} \quad (2.1)$$




Figure 2: (a) Magnitude of  $\mathbf{F}_i$  in red. The blue dashed line correspond to the profile of  $F_i$  if we were to treat the fiber as a Hookean spring with constant  $EA/L_0$ . (b) Schematic representation of stress fiber contraction. As the fiber contracts below the rest length  $L_0$ , buckling occurs. As myosin mediated contraction causes the fiber to contract below the rest length  $L_0$ , buckling occurs due to lack of resistance to compression. Below the critical length  $L_c$ , the fiber ceases to contract due to vanishing interfilament distance. Modified from [44].

where  $T_i$  is the magnitude of contractile force due to myosin motors,  $EA$  is the one-dimensional Young's modulus,  $L_0$  and  $L_c$  are, respectively, rest and critical lengths,  $L_i = \|\mathbf{x}_n - \mathbf{x}_i\|$  and  $\mathbf{e}_i = \frac{\mathbf{x}_n - \mathbf{x}_i}{L_i}$  is the unit vector along the  $i^{\text{th}}$  SF, and  $\delta$  is a small positive constant.

Note that since  $\mathbf{x}_i(t) \in R_{\text{cell}}\mathbb{S}^1$  and FA sites are equally spaced, then in polar coordinates we have:

$$\mathbf{x}_i(t) = R_{\text{cell}}(\cos(\theta_i), \sin(\theta_i))^T, \quad \theta_i(t) := \theta_1(t) + (i-1)\frac{2\pi}{M},$$

and so  $\mathbf{F}_i = \mathbf{F}_i(\mathbf{x}_n, \theta_1)$ .

Since the total force at  $\mathbf{x}_n$  is

$$\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) := - \sum_{i=1}^M Y_i \mathbf{F}_i(\mathbf{x}_n, \theta_1), \quad (2.2)$$

then, assuming negligible inertial effects (due to the viscous nature of cytoplasm) and constant  $\mathbf{Y}$ :

$$\beta_{\text{cell}} \dot{\mathbf{x}}_n = \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}), \quad (2.3)$$

where  $\beta_{\text{cell}}$  is the drag coefficient in the cytoplasm.

The representation of a cell in such a way is justified for the following reasons:

- The first case in (2.1) is due to the Hookean behavior of SFs upon extension and myosin tension generation. Furthermore, stress fiber laser ablation experiments [30], [32], [57] revealed that the initial instantaneous response (elastic behavior due to the SF length dependence in the first case) is followed by slower contraction due to myosin activity (force dependence on  $T_i$ ) in the remaining portion of the fiber. Combined with stress fiber buckling, we obtain the second case in (2.1). Deguchit et al. [15] also found that SF contraction ceased after reaching a certain critical length. This implies that  $F_i = 0$  when  $L_i < L_c$ . For technical reasons we assume  $F_i$  is piecewise continuous - hence the remaining case in (2.1). We also assumed for simplicity that myosin generated force  $T_i$  in each SF is constant.
- The traction stresses are largely applied on the cell periphery and their magnitude decays rapidly towards the center [42], [63]. Thus, the cell body SF ends are at or near mechanical equilibrium. Since contractile forces are generated by SFs, then a cell body SF end must be balanced by all other SFs (due to the equilibrium). Hence, it is reasonable to have a single connecting node of radial SFs which is either at mechanical equilibrium (for stationary cells) or tends to it.
- Paul et al. [49] demonstrated in their active cable network model, combined with force application originating from nuclear region on FAs by star-like SF arrangement, results in cells acquiring morphologies typical for motile cells. Since the distribution of forces applied on FAs determines which one ruptures, then it also influences the motion of a cell (due to retraction). Since we are primarily interested in cell migration it is justified to assume that this architecture represents a realistic situation. Furthermore, Oakes et al. [45] found that modeling SFs embedded in contractile networks, where only SFs actively contract, yields a behavior mimicking the experimental results - the cytoskeletal flow was directed along the stress fibers. In the same study, the authors concluded that it is appropriate to treat an SF as a 1D viscoelastic contractile element, which also justifies neglecting inertia in Equation (2.2).
- Since motile cells assume a wide variety of cell shapes and continuously remodel their actin cytoskeleton, one can view this representation as a cell shape normalization (it is implicitly assumed that a cell volume remains constant). That is, Figure 1b depicts a cell and forces applied on FAs normalized to a circle. Möhl et al. [42] applied shape normalization technique to a timelapse series data of migrating keratinocytes and demonstrated that this allows consistent analysis of FA dynamics, actin flow and traction forces. In view of their results, a particular cell traction force map and FA configuration normalized to a circle can be effectively captured by our representation.

Note that  $\mathbf{x}_n(t) \in \Omega_{cell}$  for  $t > 0$ , proved below.

**Proposition 2.1.** *Let  $\mathbf{x}_n \in \partial\Omega_{cell}$ ,  $\theta_1 \in [0, 2\pi)$ , and  $\mathbf{Y} \in \{0, 1\}^M$  be arbitrary. Let  $\mathbf{n}$  be outward unit normal at  $\mathbf{x}_n$ . Then  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \mathbf{n} \leq 0$  with equality sign if and only if  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) = 0$ .*

*Proof.* Suppose  $\exists \mathbf{x}_n \in \partial\Omega_{cell}$ ,  $\exists \theta_1 \in [0, 2\pi)$ , and  $\exists \mathbf{Y} \in \{0, 1\}^M$  such that

$$\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \mathbf{n} = \sum_{i=1}^M Y_i(-\mathbf{F}_i(\mathbf{x}_n, \theta_1)) \cdot \mathbf{n} > 0. \quad (2.4)$$

Since  $\mathbf{x}_n \in \partial\Omega_{cell}$ , then in polar coordinates we have  $\mathbf{x}_n = R_{cell}(\cos(\theta), \sin(\theta))^T$ . Note that  $\forall i$  such that  $\theta_i \notin [\theta - \pi/2, \theta + \pi/2]$  we have:

$$Y_i(-\mathbf{F}_i(\mathbf{x}_n, \theta_1)) \cdot \mathbf{n} = Y_i \|\mathbf{F}_i(\mathbf{x}_n, \theta_1)\| |(-\mathbf{e}_i \cdot \mathbf{n})| = \frac{Y_i \|\mathbf{F}_i(\mathbf{x}_n, \theta_1)\|}{L_i} (\mathbf{x}_i - \mathbf{x}_n) \cdot \mathbf{n} \leq 0,$$

since  $\mathbf{x}_i \cdot \mathbf{n} < 0$  and  $\mathbf{x}_n \cdot \mathbf{n} > 0$ . Thus  $\exists i \in \{1, \dots, M\}$  such that  $\theta_i \in [\theta - \pi/2, \theta + \pi/2]$ ,  $Y_i = 1$  and  $(-\mathbf{F}_i(\mathbf{x}_n, \theta_1)) \cdot \mathbf{n} > 0$ . Otherwise Inequality (2.4) does not hold and we are done. But then,

$$(\mathbf{x}_i - \mathbf{x}_n) \cdot \mathbf{n} > 0 \Rightarrow (\cos(\theta_i), \sin(\theta_i))^T \cdot \mathbf{n} > 1,$$

which is false, since both are unit vectors. Therefore,  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \mathbf{n} \leq 0$  for arbitrary  $\mathbf{x}_n \in \partial\Omega_{cell}$  and  $\mathbf{Y} \in \{0, 1\}^M$ .

Suppose  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \mathbf{n} = 0$ . Then, we have (by the above considerations):

$$\sum_{i'} Y_{i'} \mathbf{F}_{i'}(\mathbf{x}_n, \theta_1) \cdot \mathbf{n} = \sum_i Y_i (-\mathbf{F}_i(\mathbf{x}_n, \theta_1)) \cdot \mathbf{n} \geq 0,$$

where  $i'$  is such that  $\theta_{i'} \notin [\theta - \pi/2, \theta + \pi/2]$  and  $i$  is such that  $\theta_i \in [\theta - \pi/2, \theta + \pi/2]$ . If we have equality, then obviously  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) = 0$ . If there is inequality, then  $\exists i$  such that  $\theta_i \in [\theta - \pi/2, \theta + \pi/2]$ ,  $Y_i = 1$  and  $(-\mathbf{F}_i(\mathbf{x}_n, \theta_1)) \cdot \mathbf{n} > 0$ , which is also false (see above).

Therefore,  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) = 0$ . ■

**Corollary 2.2.** *Let  $\mathbf{x}_n(0) \in \Omega_{cell}$  be arbitrary and let  $\theta_1(t)$  be given. Suppose  $\mathbf{x}_n \in C^1([0, \infty))$  is a solution of (2.3). Then,  $\mathbf{x}_n(t) \in \Omega_{cell} \forall t > 0$  and  $\forall \mathbf{Y}(t) \in \{0, 1\}^M$ .*

*Proof.* Due to (2.3) it suffices to show that  $\forall \mathbf{x}_n \in \partial\Omega_{cell}$  we have  $\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \mathbf{n} \leq 0$ , which follows from Proposition 2.1. ■

## 2.2 The cell migration cycle

Recall that during the migration cycle, deadhesion leads to cell body translocation, while adhesion binding does not. In both cases actomyosin contractility leads to reconfiguration of the cytoskeleton occurs. Here we show how our cell representation can describe the reconfiguration and cell body motion following binding and unbinding events.

Without loss of generality assume that an event occurred at  $t = 0$ . Let  $\tau > 0$  be the time of the next adhesion event, be it binding or unbinding. Let  $\mathbf{Y}(0) \in \{0, 1\}^M$ ,  $\mathbf{x}(0) \in \mathbb{R}^2$ , and  $\mathbf{x}_n(0) \in \Omega_{cell}$  be arbitrary. Then,  $\mathbf{Y}(t) = \text{const.}$  for  $t \in [0, \tau)$ .

We assume  $\theta_1(t = 0) = 0$ . Since the FA sites are equally spaced, it is sufficient to consider  $\theta_1(t)$  only.

### 2.2.1 Focal adhesion binding

Following an FA binding, we suppose that a cell becomes stationary (i.e. cell centroid remains constant). However, a newly formed FA and the associated SF lead to cytoskeletal reshaping. Thus, we have the following system of ODEs for  $t \in [0, \tau)$ :

$$\begin{aligned}\dot{\mathbf{x}} &= 0 \\ \dot{\mathbf{x}}_n &= \beta_{cell}^{-1} \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \\ \dot{\theta}_1 &= 0.\end{aligned}\tag{2.5}$$

### 2.2.2 Focal adhesion unbinding

Following unbinding event, cytoskeletal contraction leads to cell body movement. Due to the circular geometry, the contractile forces induce both rotational and translational motion.

Note that the bound focal adhesions are able to slide for short distances [42]. Oakes et al. [45] found that the cytoskeleton behaves like an elastic solid on timescales up to one hour. Provided the time  $\tau$  between adhesion events is small enough, the following is justified.

The force  $\mathbf{F}$  along the radial vector  $\hat{\mathbf{r}}(\mathbf{x}_n)$  is acting on the cell center, thereby inducing translational motion (see Figure 3). On the other hand, the rotational motion is produced due to  $\mathbf{F}$  acting along the tangential vector  $\hat{\boldsymbol{\varphi}}(\mathbf{x}_n)$ . The radial and tangential components of the force  $\mathbf{F}$  are given by:

$$\begin{aligned}F_r &:= \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\mathbf{r}}(\mathbf{x}_n) \\ F_\varphi &:= \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\boldsymbol{\varphi}}(\mathbf{x}_n),\end{aligned}$$

where  $\mathbf{x}_n = (x_{n,1}, x_{n,2})$  and

$$\hat{\mathbf{r}}(\mathbf{x}_n) = \frac{\mathbf{x}_n}{\|\mathbf{x}_n\|}, \quad \hat{\boldsymbol{\varphi}}(\mathbf{x}_n) = \left( -\frac{x_{n,2}}{\|\mathbf{x}_n\|}, \frac{x_{n,1}}{\|\mathbf{x}_n\|} \right)^T.$$

Note that the characteristic Reynolds number  $Re$  is given by

$$Re = \frac{\rho \cdot s \cdot L}{\nu} \sim 10^{-6} - 10^{-4},$$

where we assumed the surrounding fluid is water (with corresponding values for density  $\rho$  and viscosity  $\nu$ ), and that characteristic cell speed  $s$  and size




Figure 3: Force diagram showing transmission of internally generated contractile forces into translational and rotational motion.  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\varphi}}$  are radial and angular unit vectors, respectively.  $\dot{\theta}_1$  is angular velocity,  $\mathbf{F}$  is a net contractile force,  $F_r$  and  $F_\varphi$  are radial and tangential components of  $\mathbf{F}$ ,  $\mathbf{x}$  and  $R_{cell}$  are cell center and radius, respectively.



$L$  are  $0.1 - 1\mu m/s$ ,  $L = 10 - 50\mu m$ , respectively. Thus, neglecting inertia, we have:

$$\begin{aligned}\dot{\mathbf{x}} &= \beta_{ECM}^{-1} \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\mathbf{r}}(\mathbf{x}_n) \hat{\mathbf{r}}(\mathbf{x}_n) \\ \dot{\mathbf{x}}_n &= \beta_{cell}^{-1} \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \\ \dot{\theta}_1 &= \beta_{rot}^{-1} \|\mathbf{x}_n\| \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\varphi}(\mathbf{x}_n)\end{aligned}\quad (2.6)$$

where  $\beta_{ECM}$  and  $\beta_{rot}$  are, respectively, translational and rotational drag coefficients in the ECM.

### 2.3 Specification of kinematics

It is convenient to transform the system above into nondimensional form. In order to do so, we define the following scales:

- Spatial and cell length scales are defined by cell radii  $R_{cell}$ .
- Time scale is defined by some constant  $t_e$ .
- Force scale is defined by characteristic force  $F_b$ .

The constants are to be specified later. Whence we define the new variables:

$$\tilde{\mathbf{x}} := \frac{\mathbf{x}}{L}, \quad \tilde{\mathbf{x}}_n := \frac{\mathbf{x}_n}{R_{cell}}, \quad \tilde{t} := \frac{t}{t_e}$$

and transform  $\mathbf{F}_i$  from (2.1):

$$\tilde{\mathbf{F}}_i := \frac{\mathbf{F}_i}{F_b} = \begin{cases} \left( \tilde{T}_i + \widetilde{EA} \frac{\tilde{L}_i - \tilde{L}_0}{\tilde{L}_0} \right) \tilde{\mathbf{e}}_i, & \tilde{L}_0 < \tilde{L}_i \\ \tilde{T}_i \tilde{\mathbf{e}}_i, & \tilde{L}_c \leq \tilde{L}_i \leq \tilde{L}_0 \\ \frac{\tilde{L}_i - \tilde{L}_c + \tilde{\delta}}{\tilde{\delta}} \tilde{T}_i \tilde{\mathbf{e}}_i, & \tilde{L}_c - \tilde{\delta} \leq \tilde{L}_i < \tilde{L}_c \end{cases}, \quad (2.7)$$

where

$$\begin{aligned}\tilde{L}_i &= \frac{L_i}{R_{cell}}, & \tilde{L}_0 &= \frac{L_0}{R_{cell}}, & \tilde{L}_c &= \frac{L_0}{R_{cell}}, & \tilde{\delta} &= \frac{\delta}{R_{cell}}, \\ \tilde{T}_i &= \frac{T_i}{F_b}, & \widetilde{EA} &= \frac{EA}{F_b}, & \tilde{\mathbf{e}}_i &= \frac{\tilde{\mathbf{x}}_n - \tilde{\mathbf{x}}_i}{\tilde{L}_i}, & \tilde{\mathbf{x}}_i &= \frac{\mathbf{x}_i}{R_{cell}}.\end{aligned}$$

Note that we have  $\tilde{\mathbf{x}}_n \in \tilde{\Omega}_{cell} := \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 \leq 1\}$  and  $\tilde{\mathbf{x}}_i \in \mathbb{S}^1$ .

Let

$$\tilde{\mathbf{F}} := \mathbf{F}/F_b, \quad \tilde{\beta}_{cell} := \frac{R_{cell}}{t_e F_b} \beta_{cell}, \quad \tilde{\beta}_{ECM} := \frac{R_{cell}}{t_e F_b} \beta_{ECM}, \quad \tilde{\beta}_{rot} := \frac{1}{t_e R_{cell} F_b} \beta_{rot}.$$

To complete the specification of cell kinematics between adhesion events, we introduce another discrete variable  $\mu \in \{0, 1\}$ :

$$\mu = \begin{cases} 1, & \text{if the last event was unbinding} \\ 0, & \text{if the last event was binding.} \end{cases}$$

Then, plugging in the rescaled quantities and dropping tildes, it follows from equations (2.5), (2.6) that the system evolves according to the following ODE system between the FA events:

$$\begin{aligned} \dot{\mathbf{x}} &= \mu \beta_{ECM}^{-1} \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\mathbf{r}}(\mathbf{x}_n) \hat{\mathbf{r}}(\mathbf{x}_n) \\ \dot{\mathbf{x}}_n &= \beta_{cell}^{-1} \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \\ \dot{\theta}_1 &= \mu \beta_{rot}^{-1} \|\mathbf{x}_n\| \mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\boldsymbol{\varphi}}(\mathbf{x}_n) \end{aligned} \quad (2.8)$$




Figure 4: Schematic representation of the migration cycle between adhesion events. Suppose just before an event occurs at time  $t = 0$ , the cell is in state  $I$ . If at time  $t = 0$  (de)adhesion occurs, the cell jumps into state  $(II')II$  and the system evolves according to (2.8) until the next event occurs at time  $t = \tau$ , after which the cycle begins anew. The scenarios can be characterized as “run” and “tumble” phases in the bottom and top panels, respectively.

### 3 FA event model

In the previous section we constructed a model of cell motion between FA events. Following [25], here we construct a stochastic model describing the random adhesion/deadhesion events and their arrival times.

#### 3.1 Focal adhesion events

Since there are  $M$  FAs and since each FA can participate in only two reactions (binding and unbinding), then there are  $2M$  total possible reactions. We adopt the following convention for enumerating reactions: reaction  $j$  corresponds to a *binding* reaction of the FA site  $i = (j + 1)/2$  if  $j$  is odd; otherwise reaction  $j$  corresponds to an *unbinding* reaction of the FA site  $i = j/2$ . Let  $\mathbf{Y}(t)$  be defined as before.

Let  $a_j(\mathbf{y}, t)dt$  be the probability, given  $\mathbf{Y}(t) = \mathbf{y} \in \{0, 1\}^M$ ,  $(\mathbf{x}(t), \mathbf{x}_n(t), \theta_1(t))$  and time  $t$ , that reaction  $j$  will occur in the time interval  $[t, t + dt)$ . For clarity, we suppress here the dependence of the rate  $a_j(\mathbf{y}, \cdot)$  on  $(\mathbf{x}(\cdot), \mathbf{x}_n(\cdot), \theta_1(\cdot))$ . We assume that the rate  $a_j$  satisfies the following:

$$a_j(\mathbf{y}, t) = \begin{cases} 0, & \text{if } j \text{ is odd and } y_{(j+1)/2} = 1 \\ 0, & \text{if } j \text{ is even and } y_{j/2} = 0 \\ \neq 0, & \text{else.} \end{cases} \quad (3.1)$$

That is, if the FA is (un)bound, the probability of the (un)binding reaction is zero; if the FA is (un)bound, the probability of (binding) unbinding is nonzero. This implies that  $a_j(\mathbf{y}, t) \neq 0$  for at least one  $j \in \{1, \dots, 2M\}$ .<sup>1</sup>

**Lemma 3.1.** *Let  $\mathbf{Y}(t) = \mathbf{y}$ . Then the probability that no FA event occurs in the time interval  $[t, t + dt)$  is  $1 - \sum_{j=1}^{2M} a_j(\mathbf{y}, t)dt + o(dt)$ .*

*Proof.* Using the definition of  $a_j$ , the probability that reaction  $j$  does not happen is  $1 - a_j(\mathbf{y}, t)dt$ . Then, the probability that no FA reaction occurs is:

$$\prod_{j=1}^{2M} (1 - a_j(\mathbf{y}, t)dt) = 1 - \sum_{j=1}^{2M} a_j(\mathbf{y}, t)dt + o(dt).$$

■

Let  $K(\tau, j|t, \mathbf{y})d\tau$  be the probability, given  $\mathbf{Y}(t) = \mathbf{y}$  and  $(\mathbf{x}(t), \mathbf{x}_n(t), \theta_1(t))$  at time  $t$ , that the *next* reaction will occur in the time interval  $[t + \tau, t + \tau + d\tau)$  and will be reaction  $j$ . Here, again, we suppress for clarity the dependence on  $\mathbf{x}, \mathbf{x}_n, \theta_1$ .

**Proposition 3.2.** *Let  $\tau > 0$  and  $\mathbf{Y}(t) = \mathbf{y}$ . Then,*

$$K(\tau, j|t, \mathbf{y}) = a_j(\mathbf{y}, t + \tau) \exp \left( - \int_t^{t+\tau} \sum_{j'=1}^{2M} a_{j'}(\mathbf{y}, \tau') d\tau' \right).$$

<sup>1</sup>For each FA site  $i \in \{1, \dots, M\}$ , either  $a_{2i-1}$  or  $a_{2i}$  is nonzero.

*Proof.* Let  $P(\tau|t, \mathbf{y})$  denote the probability that no reaction occurs in the time interval  $[t, t + \tau)$ , given  $\mathbf{y}$  (and  $\mathbf{x}, \mathbf{x}_n, \theta_1$ ) at time  $t$ . Then, by Lemma 3.1:

$$P(\tau + d\tau|t, \mathbf{y}) = P(\tau|t, \mathbf{y}) \left( 1 - \sum_{j=1}^{2M} a_j(\mathbf{y}, t) d\tau + o(d\tau) \right) \Rightarrow$$

$$\frac{P(\tau + d\tau|t, \mathbf{y}) - P(\tau|t, \mathbf{y})}{d\tau} = -P(\tau|t, \mathbf{y}) \sum_{j=1}^{2M} a_j(\mathbf{y}, t) + P(\tau|t, \mathbf{y}) \frac{o(d\tau)}{d\tau}.$$

Letting  $d\tau \rightarrow 0$  we obtain the following ODE:

$$\frac{d}{d\tau} P(\tau|t, \mathbf{y}) = -P(\tau|t, \mathbf{y}) \sum_{j=1}^{2M} a_j(\mathbf{y}, t).$$

Since  $P(0|t, \mathbf{y}) = 1$ , the solution  $P(\tau|t, \mathbf{y})$  is given by:

$$P(\tau|t, \mathbf{y}) = \exp \left( - \int_t^{t+\tau} \sum_{j=1}^{2M} a_j(\mathbf{y}, \tau') d\tau' \right).$$

We have then:

$$K(\tau, j|t, \mathbf{y}) = P(\tau|t, \mathbf{y}) a_j(\mathbf{y}, t + \tau) = a_j(\mathbf{y}, t + \tau) \exp \left( - \int_t^{t+\tau} \sum_{j'=1}^{2M} a_{j'}(\mathbf{y}, \tau') d\tau' \right). \quad (3.2)$$

■

Let  $K_{time}(\tau|t, \mathbf{y}) d\tau$  be the probability that the next reaction will occur in the time interval  $[t + \tau, t + \tau + d\tau)$ , given  $\mathbf{Y}(t) = \mathbf{y}$  and  $(\mathbf{x}(t), \mathbf{x}_n(t), \theta_1(t))$  at time  $t$ .

Let  $K_{index}(j|\tau, t, \mathbf{y})$  be the probability that the index of the next reaction is  $j$  given  $\mathbf{Y}(t) = \mathbf{y}$ ,  $(\mathbf{x}(t), \mathbf{x}_n(t), \theta_1(t))$  at time  $t$  and given that the reaction will occur at time  $t + \tau$ .

By elementary probability theory (using the definition of conditional probability), we know that

$$K(\tau, j|t, \mathbf{y}) d\tau = K_{index}(j|\tau, t, \mathbf{y}) K_{time}(\tau|t, \mathbf{y}) d\tau.$$

Due to equation (3.2), we see that:

$$K_{index}(j|\tau, t, \mathbf{y}) = \frac{a_j(\mathbf{y}, t + \tau)}{a_0(\mathbf{y}, t + \tau)}$$

$$K_{time}(\tau|t, \mathbf{y}) = a_0(\mathbf{y}, t + \tau) \exp \left( - \int_t^{t+\tau} a_0(\mathbf{y}, \tau') d\tau' \right), \quad (3.3)$$

where

$$a_0(\mathbf{y}, t) = \sum_{j=1}^{2M} a_j(\mathbf{y}, t),$$

and  $a_0 \neq 0$  due to equation (3.1).

Obviously,

$$\sum_{j=1}^{2M} K_{index}(j|\tau, t, \mathbf{y}) = 1$$

$$\int_0^\infty K_{time}(\tau|t, \mathbf{y}) = 1.$$

Thus, if  $T$  is (random) time until the next reaction, then its probability density function given by  $K_{time}$ , its survival function  $S(s)$  is given by (without loss of generality, suppose that  $t = 0$ ):

$$\mathbb{P}(T > s) = S(s) = \exp\left(-\int_0^s a_0(\mathbf{y}, \tau') d\tau'\right), \quad (3.4)$$

and its (cumulative) distribution function is given by  $1 - S(s)$ .<sup>1</sup> Note that the distribution of a random variable is uniquely determined by its distribution function.

Using the proof of Proposition 3.2 one has the following:

**Proposition 3.3.** *Let  $\tau > 0$  and let  $\hat{K}(\tau|t, \mathbf{y})d\tau$  be the probability of more than one FA event occurring in the time interval  $[t + \tau, t + \tau + d\tau)$ , given the state of the system at time  $t$ . Then  $\hat{K}(\tau|t, \mathbf{y}) = 0$ .*

*Proof.* By the proof of Proposition 3.2:

$$\hat{K}(\tau|t, \mathbf{y})d\tau = P(\tau|t, \mathbf{y})o(d\tau),$$

since, following the definition of  $a_j$ , the probability of more than one reaction occurring in time interval  $[t, t + d\tau)$  is  $o(d\tau)$ .

The result follows from dividing the equation above by  $d\tau$ , letting  $d\tau \rightarrow 0$  and seeing that  $\frac{o(d\tau)}{d\tau} \rightarrow 0$ . ■

## 3.2 Combining the cell motility and the FA event model

With the results of the previous section we can now formally state the cyclical mesenchymal cell motility model as a stochastic process (see Figure 4).

Let  $t = 0$ ,  $\mathbf{x}(0)$ ,  $\mathbf{x}_n(0)$ ,  $\theta_1(0)$ ,  $\mu$  be given and  $\mathbf{Y}(0) = \mathbf{y}^0$ .

- The time  $T_1$  of the FA event is chosen such that  $\mathbb{P}(T_1 > s) = S(s)$ .
- The system evolves according to (2.8) for  $t \in [0, T_1)$ .

---

<sup>1</sup>One can check this by differentiating the distribution function, given by  $1 - S(s)$ , with respect to  $s$ .

- At time  $t = \tau$ , the index  $j$  of the FA event is chosen with probability  $K_{index}(j|T_1, 0, \mathbf{y}^0)$ .  $\mathbf{Y}$  and  $\mu$  jump to new values:

$$\mathbf{Y}(t = \tau) = \begin{cases} \mathbf{y}^0 + \mathbf{e}_i, & i = (j + 1)/2, \text{ if } j \text{ is odd} \\ \mathbf{y}^0 - \mathbf{e}_i, & i = j/2, \text{ else} \end{cases},$$

$$\mu(t = \tau) = \begin{cases} 0, & \text{if } j \text{ is odd} \\ 1, & \text{else} \end{cases},$$

where  $\mathbf{e}_i \in \mathbb{R}^M$  is the standard basis vector. Note that due to (3.1), we always have  $\mathbf{Y}(t = \tau) \in \{0, 1\}^M$ , since the probability of (un)binding of (un)bound FA is zero.

- The cycle starts anew with initial time  $t = T_1$  and initial values of other variables at this time: starting at  $t = T_1$  we choose the time  $T_2$  of the FA event such that

$$\mathbb{P}(T_2 > s|T_1) = \exp\left(-\int_{T_1}^{T_1+s} a_0(\mathbf{y}, \tau')d\tau'\right),$$

- The system evolves according to (2.8) for  $t \in [T_1, T_1 + T_2)$  and so on.

One sees that the cyclical process described above is a Markov process, since the evolution of the system depends only on the current state. This completes formal specification of the model. In the following we will show that this process is well-defined.

## 4 Piecewise deterministic process

In this section we first briefly overview a class of piecewise deterministic processes, first introduced by Davis [13]. We then show how the deterministic equations, describing motion between stochastic focal adhesion events, can be combined to yield a well-defined piecewise deterministic Markov process (PDMP).

### 4.1 PDMP overview

Let  $A$  be countable and let  $\Gamma \subset \mathbb{R}^d$  be open. Let  $\mathbf{X}_t \in \Gamma$  and let  $\mathbf{H}_\nu : \Gamma \rightarrow \mathbb{R}^d$  for  $\nu \in A$ .

Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  be a filtered probability space, where  $\Omega$  is a sample space,  $\mathcal{F}$  is a  $\sigma$ -algebra on  $\Omega$ ,  $(\mathcal{F}_t)_{t \geq 0}$  is a (natural) filtration, and  $\mathbb{P}$  is a probability measure. Let  $E := \{(\nu, \boldsymbol{\xi}) : \nu \in A, \boldsymbol{\xi} \in \Gamma\}$  and let  $(E, \mathcal{E})$  be a Borel space. For details see [Chapter 2 in [14]].

We can define the piecewise deterministic process on the state space  $(E, \mathcal{E})$  (for a more detailed general treatment see Davis [14]) by the following objects<sup>1</sup>:

---

<sup>1</sup>Here we first provide a constructive definition. Verification of the conditions and their explicit representation corresponding to our case of cell motility is postponed for the sake of clearer exposition

I Vector fields  $(\mathbf{H}_\nu, \nu \in A)$  such that for all  $\nu \in A$  there exists a unique global solution  $\mathbf{X}_t \in \Gamma$  to the following equation:

$$\begin{aligned} \frac{d}{dt} \mathbf{X}_t &= \mathbf{H}_\nu(\mathbf{X}_t) \\ \mathbf{X}_0 &\in \Gamma. \end{aligned} \tag{4.1}$$

Let  $\phi_\nu : [0, \infty) \times \Gamma \rightarrow \Gamma$  denote the flow corresponding to Equation (4.1), i.e.

$$\phi_\nu(t, \mathbf{X}_0) = \mathbf{X}_t.$$

II A measurable function  $a_0 : E \rightarrow \mathbb{R}_+$  such that the function  $s \mapsto a_0(\nu, \phi_\nu(s, \mathbf{X}_0))$  is integrable.

III A transition measure  $Q : \mathcal{E} \times E \rightarrow [0, 1]$ , such that for fixed  $C \in \mathcal{E}$ ,  $(\nu, \boldsymbol{\xi}) \mapsto Q(C; (\nu, \boldsymbol{\xi}))$  is measurable for  $(\nu, \boldsymbol{\xi}) \in E$ , and  $Q(\cdot; (\nu, \boldsymbol{\xi}))$  is a probability measure for all  $(\nu, \boldsymbol{\xi})$  on  $(E, \mathcal{E})$ .

Let  $(\nu^0, \mathbf{X}^0) \in E$  at time  $t = 0$  be given. Let a survival function  $S$  be defined similarly as in equation (3.4):

$$S(t, (\nu, \mathbf{X})) := \exp\left(-\int_0^t a_0(\nu, \phi_\nu(s, \mathbf{X})) ds\right). \tag{4.2}$$

Let  $T_1$  be the first jump time such that

$$\mathbb{P}(T_1 > t | (\nu^0, \mathbf{X}^0)) = S(t, (\nu^0, \mathbf{X}^0)),$$

and let  $(\nu^1, \mathbf{X}^1)$  be distributed according to the probability law  $Q(\cdot, \phi_{\nu^0}(T_1, \mathbf{X}^0))$ . Then, the motion of  $(\nu_t, \mathbf{X}_t)$  for  $t \leq T_1$  is given by:

$$(\nu_t, \mathbf{X}_t) = \begin{cases} (\nu^0, \phi_{\nu^0}(t, \mathbf{X}^0)), & t < T_1, \\ (\nu^1, \mathbf{X}^1), & t = T_1. \end{cases}$$

At time  $t = T_1$  the next jump time  $T_2$  is distributed such that

$$\mathbb{P}(T_2 - T_1 > s | (\nu_{T_1}, \mathbf{X}_{T_1})) = S(s, (\nu_{T_1}, \mathbf{X}_{T_1})).$$

The value of the process at the jump time  $T_2$  is determined by the measure  $Q(\cdot, \phi_{\nu_{T_1}}(T_2, \mathbf{X}_{T_1}))$  and the process continues in a similar way. Thus, we have a well-defined piecewise deterministic process [13].

**Theorem 4.1** ([13]). *The process  $(\nu_t, \mathbf{X}_t)_{t \geq 0}$  is a homogeneous Markov process.*

## 4.2 Cell motility and PDMP

In this section we show that the cyclical cell motility model described in Section 3.2 is a well-defined PDMP.

One can show that  $\mathbf{F}_i(\mathbf{x}_n, \theta_1)$  satisfies Lipschitz condition for  $(\mathbf{x}_n, \theta_1) \in \Omega_{cell} \cup [0, 2\pi) := D_{cell}^1$ . Furthermore, one can show that  $\beta_{cell}^{-1}\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y})$  and  $\mu\beta_{rot}^{-1}\|\mathbf{x}_n\|\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\varphi}(\mathbf{x}_n)$ , given by

$$\begin{aligned}\beta_{cell}^{-1}\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) &= -\beta_{cell}^{-1} \sum_{i=1}^M Y_i \mathbf{F}_i(\mathbf{x}_n, \theta_1) \\ \mu\beta_{rot}^{-1}\|\mathbf{x}_n\|\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\varphi}(\mathbf{x}_n) &= -\mu\beta_{rot}^{-1}\|\mathbf{x}_n\| \sum_{i=1}^M Y_i \mathbf{F}_i(\mathbf{x}_n, \theta_1) \cdot \hat{\varphi}(\mathbf{x}_n)\end{aligned}$$

also satisfy the Lipschitz condition for  $(\mathbf{x}_n, \theta_1) \in D_{cell}$  and arbitrary  $\mu \in \{0, 1\}$ ,  $\mathbf{Y} \in \{0, 1\}^M$ .

**Proposition 4.2.** *Let  $\mathbf{x}(0) = \mathbf{x}_0$ ,  $(\mathbf{x}_n(0), \theta_1(0)) \in D_{cell}$ . Let  $\mu \in \{0, 1\}$ ,  $\mathbf{Y} \in \{0, 1\}^M$ . Then there exists a unique solution of the system*

$$\begin{aligned}\dot{\mathbf{x}} &= \mu\beta_{ECM}^{-1}\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\mathbf{r}}(\mathbf{x}_n)\hat{\mathbf{r}}(\mathbf{x}_n) \\ \dot{\mathbf{x}}_n &= \beta_{cell}^{-1}\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \\ \dot{\theta}_1 &= \mu\beta_{rot}^{-1}\|\mathbf{x}_n\|\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\varphi}(\mathbf{x}_n),\end{aligned}\tag{4.3}$$

for  $t > 0$ .

*Proof.* Note that since the evolution of  $\mathbf{x}$  is decoupled from the other two equations, it is sufficient to prove the claim for the following subsystem:

$$\begin{aligned}\dot{\mathbf{x}}_n &= \beta_{cell}^{-1}\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \\ \dot{\theta}_1 &= \mu\beta_{rot}^{-1}\|\mathbf{x}_n\|\mathbf{F}(\mathbf{x}_n, \theta_1, \mathbf{Y}) \cdot \hat{\varphi}(\mathbf{x}_n)\end{aligned}\tag{4.4}$$

Since the right hand side of this system is Lipschitz on  $D_{cell}$  and  $(\mathbf{x}_n(0), \theta_1(0)) \in D_{cell}$ , then there exists unique solution of the subsystem (4.4) for time  $t \leq t_{D_{cell}}$ , where  $t_{D_{cell}} = \inf \{t^* > 0 | \mathbf{x}_n(t^*) \notin \Omega_{cell}\}$  is the exit time from  $D_{cell}$ . By Corollary 2.2, we see that  $t_{D_{cell}} = \infty$ . ■

Let  $A := \{1, 2, \dots, 2^{M+1}\}$  and let  $\alpha : A \rightarrow \{0, 1\} \times \{0, 1\}^M$  be a bijection. This is simply a mapping such that  $\alpha(\nu) = (\mu, \mathbf{Y}) \in \{0, 1\} \times \{0, 1\}^M$  corresponds to a particular FA and cell motion state (recall that the latter can either be moving or stationary).

---

<sup>1</sup>The restriction to the interval  $[0, 2\pi)$  is due to periodic dependence on  $\theta_1$  in the definition of  $\mathbf{x}_i$ . See Section 2



Let  $(\mathbf{x}(0), \mathbf{x}_n(0), \theta_1(0)) \in \Gamma := \mathbb{R}^2 \times \Omega_{cell} \times [0, 2\pi)$  and denote  $\mathbf{X}_t = (\mathbf{x}(t), \mathbf{x}_n(t), \theta_1(t))$ . Moreover, let  $\mathbf{H}_\nu : \Gamma \rightarrow \mathbb{R}^5$  be such that

$$\mathbf{H}_\nu(\mathbf{X}) := \begin{pmatrix} \alpha_1(\nu) \beta_{ECM}^{-1} \mathbf{F}(\mathbf{x}_n, \boldsymbol{\alpha}_2(\nu), \theta_1) \cdot \hat{\mathbf{r}}(\mathbf{x}_n) \hat{\mathbf{r}}(\mathbf{x}_n) \\ \beta_{cell}^{-1} \mathbf{F}(\mathbf{x}_n, \boldsymbol{\alpha}_2(\nu), \theta_1) \\ \alpha_1(\nu) \beta_{rot}^{-1} \|\mathbf{x}_n\| \mathbf{F}(\mathbf{x}_n, \boldsymbol{\alpha}_2(\nu), \theta_1) \cdot \hat{\boldsymbol{\varphi}}(\mathbf{x}_n) \end{pmatrix}, \quad (4.5)$$

where  $\alpha_1$  and  $\boldsymbol{\alpha}_2$  are, respectively, the first and the second components of  $\boldsymbol{\alpha}^1$ .

Let the probability  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  and state space  $(E, \mathcal{E})$  be defined as in the previous section.

We now specify objects (I,II,III) described in Section 4.1.

I By Proposition 4.2 we see that for all  $\nu \in A$ , there exists a unique global solution to (4.1).

II Note that in our case the rate function  $a_0$  is given by (recalling Section 3.1)<sup>2</sup>:

$$a_0(\nu, \mathbf{X}_t) = a_0(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_t) = \sum_{j=1}^{2M} a_j(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_t).$$

Thus, for the integrability condition to be satisfied, we assume that each probability rate function  $a_j$  is integrable along the solution of equation (4.1). An exact form of the rates  $a_j$  satisfying this condition will be given in the subsequent section. Note that  $a_0$  is nonzero, which follows from (3.1).

III In our case, the measure  $Q(\cdot; (\nu, \boldsymbol{\xi}))$  is given by (recalling Section 3):

$$\begin{aligned} Q(\{\eta\} \times d\boldsymbol{\xi}'; (\nu, \boldsymbol{\xi})) &= \delta_{\boldsymbol{\xi}}(d\boldsymbol{\xi}') \sum_{j=1}^M \delta_{\alpha_1(\eta), 0} \frac{a_j^+(\boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi})}{a_0(\boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi})} \delta_{\alpha_2(\eta)_j, 1} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i} \\ &\quad + \delta_{\alpha_1(\eta), 1} \frac{a_j^-(\boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi})}{a_0(\boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi})} \delta_{\alpha_2(\eta)_j, 0} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i}, \end{aligned} \quad (4.6)$$

where  $\delta$  is the Kronecker delta function,  $\delta_{\boldsymbol{\xi}}(\cdot)$  is the Dirac measure at  $\boldsymbol{\xi}$ ,  $a_j^+ = a_{2j-1}$  and  $a_j^- = a_{2j}$  correspond to, respectively, the binding and unbinding probability rates at FA site  $j$ , and  $\boldsymbol{\alpha}_2(\cdot)_i$  is the  $i^{\text{th}}$  component of the vector  $\boldsymbol{\alpha}_2(\cdot)$ .

The justification for choosing the functions above stems from our deductions in Section 3.1. In particular, the rate function  $a_0$  in (II) is due to (3.3): the probability density function

<sup>1</sup>In this work, the subscript index in  $\boldsymbol{\alpha}_2$  always denotes the second component of  $\boldsymbol{\alpha}$ .

<sup>2</sup>We abuse the notation introduced in Section 3.1:  $a_j(\boldsymbol{\alpha}_2(\nu), t) = a_j(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_t) = a_j(\nu, \mathbf{X}_t)$  for  $j = 0, \dots, 2M$ .

of the jump time  $T_{k+1}$ , given that  $T_k \leq t < T_{k+1}$ , is given by:

$$\begin{aligned} K_{time}(\cdot|t, \boldsymbol{\alpha}_2(\nu_t)) &= a_0(\boldsymbol{\alpha}_2(\nu_t), \mathbf{X}_t) \exp\left(-\int_0^\cdot a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) ds\right) \\ &= a_0(\boldsymbol{\alpha}_2(\nu_{T_k}), \mathbf{X}_t) \exp\left(-\int_0^\cdot a_0(\nu_{T_k}, \phi_{\nu_{T_k}}(s, \mathbf{X}_t)) ds\right), \end{aligned}$$

which corresponds to the survival function given by (4.2).

We now turn our attention to the measure  $Q$  in (4.6). The components of  $\mathbf{X}_t$  do not jump, and vary continuously in time, i.e. if  $T_k$  is the jump time, then  $\mathbf{X}_{T_k^-} = \mathbf{X}_{T_k}$  (see Section 3.2), hence the Dirac measure  $\delta_{\boldsymbol{\xi}}(\cdot)$  at  $\boldsymbol{\xi}$  in (4.6). Clearly, such transition of the continuous component  $\mathbf{X}_t$  of the PDMP at a jump time is probabilistically independent of the transition of the discrete component  $\nu$ . Hence we have the product of the Dirac measure with the sum, which is a discrete measure for the transition of the discrete component.

By Proposition 3.3, there is only one FA event at a jump time. Hence, for the probability of transition  $\nu \rightarrow \eta$  to be nonzero, the vectors of FA states  $\boldsymbol{\alpha}_2(\nu)$  and  $\boldsymbol{\alpha}_2(\eta)$  must differ only by one component. Consider the following example to illustrate the jump mechanism.

*Example.* Let  $M = 4$ ,  $T_k \leq t < T_{k+1}$  and suppose  $T_{k+1}$ ,  $\mathbf{X}_t$  are given. Let  $\nu, \eta \in A$  be such that  $\boldsymbol{\alpha}(\nu) = (\mu_\nu, \mathbf{Y}_\nu)$  and  $\boldsymbol{\alpha}(\eta) = (\mu_\eta, \mathbf{Y}_\eta)$ , where

$$\mathbf{Y}_\nu = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \mathbf{Y}_\eta = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}.$$

Then, by equation (3.3), the probability of the transition  $\nu \rightarrow \eta$  is given by:

$$\begin{aligned} \delta_{\mu_\eta, 0} K_{index}(3|T_{k+1} - t, t, \mathbf{Y}_\nu) &= \delta_{\mu_\eta, 0} \frac{a_3(\boldsymbol{\alpha}_2(\nu), \phi_\nu(T_{k+1} - t, \mathbf{X}_t))}{a_0(\boldsymbol{\alpha}_2(\nu), \phi_\nu(T_{k+1} - t, \mathbf{X}_t))} \\ &= \delta_{\alpha_1(\eta), 0} \frac{a_3(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})}{a_0(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})} \\ &= \delta_{\alpha_1(\eta), 0} \frac{a_2^+(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})}{a_0(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})}. \end{aligned}$$

Clearly, the transition  $\mathbf{Y}_\nu \rightarrow \mathbf{Y}_\eta$  corresponds to the binding event at FA site 2, explaining the Kronecker delta term (see Sections 2.3, 3.2). Now, consider the sum in (4.6) for this example. We see that

$$\prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i} \neq 0, \text{ for } j = 2 \text{ only.}$$

We therefore obtain

$$Q(\{\eta\} \times d\boldsymbol{\xi}'; (\nu, \mathbf{X}_{T_{k+1}})) = \delta_{\mathbf{X}_{T_{k+1}}} (d\boldsymbol{\xi}') \delta_{\alpha_1(\eta), 0} \frac{a_2^+(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})}{a_0(\boldsymbol{\alpha}_2(\nu), \mathbf{X}_{T_{k+1}})} \delta_{\alpha_2(\eta)_{2,1}}.$$

Note that if at time  $t$  the vector of FA states is given by  $\mathbf{Y}_\nu$ , then there are  $M$  possible FA state vectors into which a transition can occur with nonzero probability:

$$\left\{ \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

Similarly as with the rate function  $a_0$ , we can derive equation (4.6) from the principles we established before.

**Proposition 4.3.** *The transition probability measure  $Q(\cdot, (\nu, \boldsymbol{\xi}))$  is given by equation (4.6) for each  $(\nu, \boldsymbol{\xi}) \in E$ .*

*Proof.* Let  $(\nu, \boldsymbol{\xi}) \in E$ ,  $\{\eta\} \times d\boldsymbol{\xi}' \in \mathcal{E}$ . Let  $(N, \boldsymbol{\Xi})$  and  $(N_-, \boldsymbol{\Xi}_-)$  be  $E$ -valued random variables before and after the jumps. Then,

$$\begin{aligned} Q(\{\eta\} \times d\boldsymbol{\xi}'; (\nu, \boldsymbol{\xi})) &= \mathbb{P}((N, \boldsymbol{\Xi}) \in \{\eta\} \times d\boldsymbol{\xi}' | (N_-, \boldsymbol{\Xi}_-) = (\nu, \boldsymbol{\xi})) \\ &= \mathbb{P}(\{\eta\} \times d\boldsymbol{\xi}' | (\nu, \boldsymbol{\xi})), \end{aligned}$$

where we omitted the random variables for notational convenience. Then we have:

$$\mathbb{P}(\{\eta\} \times d\boldsymbol{\xi}' | (\nu, \boldsymbol{\xi})) = \mathbb{P}(d\boldsymbol{\xi}' | \{\eta\}, (\nu, \boldsymbol{\xi})) \mathbb{P}(\{\eta\} | (\nu, \boldsymbol{\xi})) = \delta_{\boldsymbol{\xi}}(d\boldsymbol{\xi}') \mathbb{P}(\eta | (\nu, \boldsymbol{\xi})),$$

since  $\boldsymbol{\Xi} = \boldsymbol{\Xi}_-$  a.s., by construction of the process. Since  $\boldsymbol{\alpha}$  is a bijection, we have<sup>1</sup>

$$\begin{aligned} \mathbb{P}(\eta | (\nu, \boldsymbol{\xi})) &= \mathbb{P}((\alpha_1(\eta), \boldsymbol{\alpha}_2(\eta)) | (\alpha_1(\nu), \boldsymbol{\alpha}_2(\nu)), \boldsymbol{\xi}) \\ &= \mathbb{P}(\alpha_1(\eta), \boldsymbol{\alpha}_2(\eta) | \boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi}), \end{aligned}$$

since, by construction of the cell motility process, the new FA state  $\boldsymbol{\alpha}_2(\eta)$  is determined independently of whether a cell was moving or not (represented by  $\alpha_1(\nu) \in \{0, 1\}$ ) and the new motility state  $\alpha_1(\eta)$  is determined only by which FA event took place (binding or unbinding), regardless of whether a cell was previously moving or not.

Note that when a jump occurs, then, by Proposition 3.3, only one of  $j = 1, \dots, 2M$  possible (binding and unbinding) FA events occurs. Thus, for  $j, j' \in \{1, \dots, 2M\}$  and  $j \neq j'$  the events “reaction  $j$  occurs” and “reaction  $j'$  occurs” are mutually exclusive. We then have, by the definition of conditional probability:

$$\mathbb{P}(\alpha_1(\eta), \boldsymbol{\alpha}_2(\eta) | \boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi}) = \sum_{j=1}^{2M} \mathbb{P}(\alpha_1(\eta), \boldsymbol{\alpha}_2(\eta) | j, \boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi}) \mathbb{P}(j | \boldsymbol{\alpha}_2(\nu), \boldsymbol{\xi}),$$

where

---

<sup>1</sup>If  $(H, \mathbf{Z})$  is a random variable, then, due to  $\boldsymbol{\alpha}$  being a bijection:

$$\{H = \eta\} = \{\boldsymbol{\alpha}^{-1}(H) = \boldsymbol{\alpha}^{-1}(\eta)\} = \{\boldsymbol{\alpha}(H) = \boldsymbol{\alpha}(\eta)\}.$$

- $\mathbb{P}(j|\alpha_2(\nu), \boldsymbol{\xi})$  is the probability that the FA event  $j$  occurs, given  $\alpha_2(\nu)$  and  $\boldsymbol{\xi}$ .
- $\mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j, \alpha_2(\nu), \boldsymbol{\xi})$  is the probability of a jump into cell state  $(\alpha_1(\eta), \alpha_2(\eta))$ , given  $\alpha_2(\nu)$  and  $\boldsymbol{\xi}$ , and that the FA event  $j$  occurred.

Let  $j \in \{1, \dots, M\}$  and  $j^+ = 2j - 1$  and  $j^- = 2j$ . Due to (3.3) we have:

$$\mathbb{P}(j^\pm|\alpha_2(\nu), \boldsymbol{\xi}) = \frac{a_j^\pm(\alpha_2(\nu), \boldsymbol{\xi})}{a_0(\alpha_2(\nu), \boldsymbol{\xi})}. \quad (4.7)$$

Furthermore,

$$\begin{aligned} \mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j^+, \alpha_2(\nu), \boldsymbol{\xi}) &= \delta_{\alpha_1(\eta), 0} \delta_{\alpha_2(\eta), 1} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i} \\ \mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j^-, \alpha_2(\nu), \boldsymbol{\xi}) &= \delta_{\alpha_1(\eta), 1} \delta_{\alpha_2(\eta), 0} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i}. \end{aligned}$$

Therefore,

$$\begin{aligned} &\sum_{j=1}^{2M} \mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j, \alpha_2(\nu), \boldsymbol{\xi}) \mathbb{P}(j|\alpha_2(\nu), \boldsymbol{\xi}) \\ &= \sum_{j=1}^M \mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j^+, \alpha_2(\nu), \boldsymbol{\xi}) \mathbb{P}(j^+|\alpha_2(\nu), \boldsymbol{\xi}) \\ &\quad + \mathbb{P}(\alpha_1(\eta), \alpha_2(\eta)|j^-, \alpha_2(\nu), \boldsymbol{\xi}) \mathbb{P}(j^-|\alpha_2(\nu), \boldsymbol{\xi}) \\ &= \sum_{j=1}^M \delta_{\alpha_1(\eta), 0} \frac{a_j^+(\alpha_2(\nu), \boldsymbol{\xi})}{a_0(\alpha_2(\nu), \boldsymbol{\xi})} \delta_{\alpha_2(\eta), 1} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i} \\ &\quad + \delta_{\alpha_1(\eta), 1} \frac{a_j^-(\alpha_2(\nu), \boldsymbol{\xi})}{a_0(\alpha_2(\nu), \boldsymbol{\xi})} \delta_{\alpha_2(\eta), 0} \prod_{i \neq j}^M \delta_{\alpha_2(\eta)_i, \alpha_2(\nu)_i}. \end{aligned}$$

■

## 5 Adhesion kinetics

While we introduced the probability rates of binding and unbinding events, we have not yet fully specified them. Here we elaborate on how such quantities unambiguously correspond to the relevant subcell dynamics by providing the precise functional forms of the propensity functions.

## 5.1 Unbinding rate

Consider the unbinding rate  $a_j^-$  of FA adhesion site  $j \in \{1, \dots, M\}$  and let  $\mathbf{y} \in \{0, 1\}^M$ ,  $\boldsymbol{\xi} = (\mathbf{x}, \mathbf{x}_n, \theta_1) \in \Gamma$ . Following Bell [6], the bond disassociation rate under applied force is given by:

$$a_j^-(\mathbf{y}, \boldsymbol{\xi}) = k_{off}^0 e^{\|\mathbf{F}_j(\mathbf{x}_n, \theta_1)\|/F_b} y_j, \quad (5.1)$$

where  $k_{off}^0$  is the FA disassociation rate under no load,  $\mathbf{F}_i$  is the force applied at the FA, given by equation (2.1), and  $F_b$  is a characteristic force scale. The last term  $y_j$  simply indicates that only bound FAs can unbind (thus satisfying equation (3.1)). Clearly, the function in (5.1) is integrable. Here we neglect the fact that the force may be applied at the FA (and consequently at the transmembrane receptors) at an angle and assume for tractability of the model that it is applied normal to the FA (hence consider only magnitude).

**Remark.** In the context of cell migration and within the framework of our model, we only consider FA disassembly on cell periphery (including the lamellae). The primary cause of such FA unbinding has mechanical, rather than biochemical nature due to the cell contractile mechanism applying load to the adhesions. Although it is known that Rho family of GTPases (in particular its member RhoA) mediate disassembly of FAs, their effect is indirect: the activity of myosin motors, which generate contractile forces in SFs, is regulated by RhoA [53]. Hence the force dependence of the unbinding rate  $a_j^-$ . Recalling the definition of  $\mathbf{F}_i$  in equation (2.1), we can include such indirect biochemical mediation by considering mediators of the force  $T_i$ . In order to keep the model tractable, we omit the interaction between RhoA and myosin motors.

## 5.2 Binding dynamics

Consider the binding probability rate  $a_j^+$  of FA adhesion site  $j \in \{1, \dots, M\}$  and let  $\mathbf{y} \in \{0, 1\}^M$ ,  $\boldsymbol{\xi} = (\mathbf{x}, \mathbf{x}_n, \theta_1) \in \Gamma$ . The probability rate  $a_j^+$  is given by:

$$a_j^+(\mathbf{y}, \boldsymbol{\xi}) = k_{on,j}(\boldsymbol{\xi})(1 - y_j),$$

where  $k_{on,j} : \Gamma \rightarrow \mathbb{R}^+$  is the effective binding rate at FA site  $j$ . The last term  $(1 - y_j)$  simply indicates that only unbound FAs can bind. Whereas unbinding can be viewed effectively as a bond rupturing under applied tension, a binding reaction, or focal adhesion assembly and maturation, is a highly regulated process. Due to complexity of FA assembly process, we focus on two major mediators of FA formation: Rac activity and contractile forces.

### 5.2.1 Rac dependence

For simplicity, we assume that the probability of FA formation is directly proportional to local Rac concentration. Consider the case of chemotactic cell migration. Leading edge protrusions preferentially form in the direction of a chemoattractant. Since Rac is required

for formation of lamellipodium and FA formation [53], then local Rac activity correlates with the concentration of the chemical cues. Conversely, local Rac activity negatively correlates with chemorepellent.

Let  $Q_{cue} : \mathbb{R}^2 \rightarrow \mathbb{R}^+$  denote the concentration of a cue in the spatial domain and let  $q : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  denote  $Q_{cue}$  dependent concentration of Rac. Clearly,  $q$  is an increasing function for the case of an attractant and a decreasing function for a repellent. Then,

$$k_{on,j}(\boldsymbol{\xi}) \propto q(Q_{cue}(\mathbf{x} + \mathbf{x}_j)),$$

where we recall that  $\mathbf{x}_j$  is the position of  $j^{\text{th}}$  FA site.

For example, we can take  $Q_{cue}(\mathbf{x})$  to be the density of the ECM (or chemoattractant) at  $\mathbf{x} \in \mathbb{R}^2$  and take  $q(x) = x$ . Then, the probability of binding is simply proportional to the ECM (or chemoattractant) density <sup>1</sup>.

### 5.2.2 Force dependence

Note that enlargement of nascent adhesions is concurrent with their maturation into focal adhesions. Thus, enlargement and maturation are synonymous. While the initial stage of adhesion growth is force-independent [11], maturation occurs in a force-dependent manner [22], [61], [69].

However, during such a force-dependent maturation, the positive correlation between the adhesion size and the applied tension exists only in the initial stages of maturation. As FAs increase in size, the force dependence plateaus [61].

That is, the study by Stricker et. al. [61] showed that for mature FAs there is no correlation between applied force and FA size. One can consider an adhesion site as mature when its size reaches  $\sim 1\mu m^2$  (see e.g. [5], [60]).

Choi et. al. [11] showed that nascent adhesions assemble at a rate of  $\sim 1.3\text{min}^{-1} = 0.021\text{s}^{-1}$  reaching a size of  $\sim 0.2\mu m^2$ . Furthermore, it was shown that the formation of these adhesions is independent of fibronectin density [11], stiffness [11] and myosin II activity [2],[11],[66].

Let  $k_{on}^0$  be force-independent FA maturation rate, which we take  $k_{on}^0 = 0.01\text{s}^{-1}$ . We now want to find a function that could represent force dependence of FA maturation rate. Denote this function  $k_{force} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ .

It satisfies the following:

- $k_{force}(0) = k_{on}^0$ , i.e. if there is no force applied, the rate is  $k_{on}^0$ .




Figure 5: Force dependence of unbinding and binding rates.

<sup>1</sup>This corresponds to a bi-molecular reaction rate, which depends on the *number* of one of the reactants (FA) and on the *concentration* (ECM or a chemoattractant) of the other.

- If the applied force is below the characteristic force  $F_b$ , then  $k_{force}$  is greater than the unbinding rate, i.e. it is more likely that an FA increases in size than it ruptures.
- If the characteristic force  $F_b$  is applied, the rate  $k_{force}$  should equal the unbinding rate, i.e. we assume that there is a dynamic equilibrium in some sense.
- If the applied force is larger than  $F_b$ , then the unbinding rate dominates the binding rate. Note that as FA increases in size, the force dependence plateaus [61]. Thus,  $k_{force}$  should plateau around  $F_b$ . We also assume that for large applied forces  $k_{force}$  plateaus at  $k_{on}^0$ , since exceeding loads rupture integrin bonds frequently and impede stable maturation.

The following form of  $k_{force}$  satisfies the conditions above:

$$k_{force}(F) = \begin{cases} \frac{k_{off}^0 e + k_{on}^0}{1 + \exp(-\gamma_1(F - F_1^*)/F_b)} + k_{on}^0 - \epsilon, & F \leq F_b \\ \frac{k_{off}^0 e + k_{on}^0}{1 + \exp(\gamma_2(F - F_2^*)/F_b)} + k_{on}^0, & \text{else} \end{cases}, \quad (5.2)$$

where  $F_1^* = F_b/4$  and  $F_2^* = 5F_b/4$  are the midpoints of the sigmoid functions (see Figure 5). The values of  $\gamma_1, \gamma_2$  and  $\epsilon$  can be found as follows:

Suppose  $F \leq F_b$ . Then,

$$\gamma_1 = -\frac{F_b}{F - F_1^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{k_{force}(F) - k_{on}^0 + \epsilon} - 1 \right).$$

Since  $k_{force}(0) = k_{on}^0$  and  $k_{force}(F_b) = k_{off}^0 e$ , then:

$$\begin{aligned} \gamma_1 &= \frac{F_b}{F_1^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{\epsilon} - 1 \right) \\ \gamma_1 &= -\frac{F_b}{F_b - F_1^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{k_{off}^0 e - k_{on}^0 + \epsilon} - 1 \right). \end{aligned}$$

It follows that  $\epsilon$  is given as the solution of the following equation:

$$\frac{F_b}{F_1^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{\epsilon} - 1 \right) + \frac{F_b}{F_b - F_1^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{k_{off}^0 e - k_{on}^0 + \epsilon} - 1 \right) = 0.$$

Similarly, since  $k_{force}(F_b) = k_{off}^0 e$ , we find:

$$\gamma_2 = \frac{F_b}{F_b - F_2^*} \log \left( \frac{k_{off}^0 e + k_{on}^0}{k_{off}^0 e - k_{on}^0} - 1 \right).$$

The values of  $\gamma_1, \gamma_2$ , and  $\epsilon$  are fixed for a value of  $F_b$ , given in Appendix A.

**Remark.** Chan and Odde [10] showed that adhesion site dynamics depends on substrate stiffness. In particular, they showed that for a stiff substrate the transmembrane bonds rupture more frequently, compared to the case with softer substrate under the same load, since the critical load is reached faster. This mechanism provides means for a cell to assess the surrounding rheology. Within the context of our model, this means that the force  $F_b$  is smaller for the stiffer substrate, thus increasing the disassociation rate for the same load (see (5.1)). Consequently, the force dependent binding rate  $k_{force}$  also changes for the stiffer ECM. In this case, the curves in Figure 5 will shift to the left. Therefore, our model provides an opportunity to include mechanosensitivity of migrating cells by considering the relevant dynamics for individual FAs.

Therefore, the binding propensity rate  $a_j^+$  of an adhesion  $j \in \{1, \dots, M\}$  is given by:

$$a_j^+(\mathbf{y}, \boldsymbol{\xi}) = q(Q_{cue}(\mathbf{x} + \mathbf{x}_j))k_{force}(\|\mathbf{F}_j(\mathbf{x}_n, \theta_1)\|)(1 - y_j). \quad (5.3)$$

## 6 Numerical Simulations

Here we show the results of simulating cell trajectories under different scenarios, which represent various experimental settings, namely:

1. Uniform environment with no cues.
2. Non-uniform environment with external cue gradient and uneven myosin motor activity within a cell.
3. Striped ECM architecture.

Note that the total number of adhesion sites  $M$  is a free parameter, which differs from cell to cell. Nevertheless, it is a crucial parameter, determining whether the motility type is amoeboid or mesenchymal. Amoeboid motility is characterized by a large number of weak adhesions, high turnover, and more contractile cell body. On the other hand, mesenchymal migration relies on fewer, but stronger focal adhesions with slower turnover and lower overall contractility. The cells with former motility type are faster and have more diffusive motion [37], [48]. Note that if  $a_j^\pm \sim O(1)$ , then the rate function is  $a_0 \sim O(M)$ . Therefore, adhesion events occur more frequently for increasing  $M$ , implying higher adhesion turnover. Thus, by varying  $M$ , we can also simulate different types of migration. It is important to stress out that the mode of migration is a spectrum, rather than binary, and a cell may transition from one motility type to the other. For simplicity we do not explicitly simulate such transition. However, it is, of course, possible to do so by simply changing the parameter  $M$  at a transition time and restarting the simulation with initial conditions given by the values of the variables at the switch times.

For all scenarios we employ the same initial conditions for all cells. Namely, at  $t = 0$  the conditions are:

- $\mathbf{x}$  is at the origin,  $\mathbf{x}_n$  is uniformly distributed on a circle with radius  $R_{cell}$ , and  $\theta = 0$ .



- Each FA is in (un)bound state and each cell is in moving state with probability 1/2.

We simulate trajectories of  $n_{cell} := 56$  cells for time  $t_{end} := 600min$ . We divided the time interval into  $n_{time} := 5000$  intervals, at the end of which we recorded the cell centroid positions  $\mathbf{x}$ . For details on the parameters and numerical methods used for simulations, see Appendix A and C, respectively. For details on the data analysis, see Appendix B.

## 6.1 Uniform environment

The results of the simulation with uniform spatial cue  $Q_{cue}$  are presented in Figure 6. Due to the absence of spatial variation of  $Q_{cue}$ , we take  $q = 1$  in equation (5.3).

The cell trajectories with varying number of adhesion sites, depicted in Figure 6 (a-c), show no clear trend and resemble those of a Brownian motion. Indeed, fitting the mean-squared displacement  $msd(t)$  to the curve  $\widehat{msd}(t) = \beta_0 t^{\bar{\beta}}$  (see Appendix B for details), we see that the exponent is  $\bar{\beta} \approx 1$  for the three cases (see Figure 6 (d-f)). This suggests that the cell motion has diffusive characteristics in this scenario. In Figure 6 (g-i) we see the simulated distribution of speeds. The average speeds  $s_{av}$  and the parameters  $\beta_0, \bar{\beta}$  are shown in Table 1. We see that as  $M$  increases, the cell motion becomes progressively faster and more diffusive<sup>1</sup>, which is expected for a dominantly amoeboid type of motility. Because  $\bar{\beta} \sim 1$ , we can estimate the diffusion coefficient  $D = \beta_0/4$ . The obtained values are lower, but within an order of magnitude estimated by Liu et. al. [37], who found that  $D \approx 2.7\mu min^2/min$ . Interestingly, gamma distribution gives a very good fit to the simulated data for various values of  $M$ , suggesting that cell speeds are gamma distributed. Moreover, the obtained values of the average speed  $s_{av}$  fall in the range reported by Liu et. al. [37], who found the speeds to be in the range from  $0.2\mu m/min$  to  $7\mu m/min$ . Although there are very high speeds observed in Figure 6 (i), which seem to be outliers, speeds as high as  $25\mu m/min$  have been observed [21]. As expected, rose plots of normalized velocities show no bias in any particular direction in Figure 6 (j-l).

M	8	16	32
$s_{av}, \mu m/min$	1.7595	2.4845	3.6047
$\bar{\beta}, 1$	1.0683	1.0035	1.0552
$\beta_0, \mu m^2/min^{\bar{\beta}}$	2.1493	3.3971	4.6308

Table 1: Parameters obtained from the simulations.

Although our results show in Figure 6 (d-f) the mean-squared displacement scales diffusively (i.e. linearly) with time, this is not consistent with the reported results. For example, Dietrich et. al. [17], Liang et. al. [36], and Liu et. al. [37] showed that the displacement scales superdiffusively. In these studies it was experimentally found that

<sup>1</sup>Since  $\bar{\beta} \sim 1$ , the slope  $\beta_0$  is a measure of diffusivity. See Appendix B.




Figure 6: Simulation results with  $M = 8, 16, 32$  adhesions in the first, second, and third columns respectively. (a-c) Trajectories of 13 cell centroids  $\mathbf{x}(t)$ . (d-f) Mean-squared displacements  $msd(t)$  (red) and fitted  $\widehat{msd}(t)$  (black) with parameters  $\beta_0$  and  $\bar{\beta}$  (see text for details). The unit of  $\beta_0$  is  $\mu\text{m}^2/\text{min}^{\bar{\beta}}$ . (g-i) Histograms of speed probability density functions and fitted density function of gamma distribution (red) with parameters  $k$  and  $\theta$ . (j-l) Relative frequency histogram of normalized velocities.

the time scaling went as  $\sim t^{\bar{\beta}}$ , where  $\bar{\beta} \approx 1.2 - 1.5$ . The primary reason why, in our case, we have diffusive behavior is that, given a certain state of adhesion sites, there is a complete circular symmetry of the probability rates  $a_j^\pm$  with respect to  $\mathbf{x}_n$  variable. Due to this symmetry, then, the probability of moving in one direction is exactly the same as the probability of moving in the opposite direction if we rotate  $\mathbf{x}_n$  by  $\pi$  radians. Hence, somewhat reminiscent of a random walk, we obtain a diffusive time scaling of the squared displacement. Moreover, this symmetry of the probability rates is somewhat idealistic, since it implies that the signaling activity relevant for adhesion dynamics is homogeneous within a cell. One of the ways to break this symmetry, is to multiply each binding probability rate  $a_j^+$  by  $1 + u$ , where  $u \sim U(-\delta, \delta)$  is uniformly distributed on the interval  $(-\delta, \delta)$  with  $\delta \in (0, 1)$ . Then, on average, the rates are unmodified<sup>1</sup>. This way, we not only simulate a non-homogeneous binding rate (and hence, for example, non-homogeneous Rac activity) within a cell, but also simulate otherwise completely identical copies of cells. Such a modification, where we do not explicitly apply a directed, predefined bias can be referred to as chemokinesis [50].

M	8			16			32		
$\delta$	0.05	0.1	0.15	0.05	0.1	0.15	0.05	0.1	0.15
$\bar{\beta}, 1$	0.9859	1.3184	1.4084	1.0086	1.3505	1.5581	1.1014	1.4299	1.5639
$s_{av},$ $\mu m/min$	1.7656	1.7768	1.7557	2.4918	2.5009	2.5021	3.5818	3.5735	3.6104
$\beta_0,$ $\mu m^2/min^{\bar{\beta}}$	3.0846	0.6934	0.5534	3.4517	0.7103	0.3543	4.1257	0.9536	0.5716

Table 2: Parameters obtained from the simulations with varying  $\delta$ .

The effect of modifying the rates  $a_j^\pm$  with  $\delta = 0.05, 0.1, 0.15$  can be seen in Figure 7. The cell trajectories, depicted in Figure 7 (a-c), show that the motion consists of periods with relatively regular path intermingled with highly irregular and random movement. In Figure 7 (d-f) we see that the rate modification leads to a superdiffusive time scaling of the mean-squared displacement, as the exponent  $\bar{\beta}$  becomes larger than one and falls within the experimentally observed range of values [17], [36], [37]. Moreover, we see that as  $\delta$  increases, so does  $\bar{\beta}$ , and the increase of the latter is more pronounced for larger number of adhesion sites  $M$  (see also Table 2). This is due to the fact that as each adhesion site is modified independently, the variance of the modified rates of a cell grows with the number of FAs, which corresponds to increased cell polarization, and hence more prominent persistent motion resulting in higher values of  $\bar{\beta}$ . However, the distribution of speeds for the

<sup>1</sup>The multiplication factor  $1 + u$  for each  $j = 1, \dots, M$  of every cell is computed at the beginning of simulations and is held fixed thereafter.




Figure 7: Simulation results with  $M = 8, 16, 32$  adhesions in the first, second, and third columns respectively, and with various values of  $\delta$ . (a-c) Trajectories of 27 cell centroids  $\mathbf{x}(t)$  with  $\delta = 0.1$ . (d-f) Mean-squared displacements  $msd(t)$  (solid) and fitted  $\widehat{msd}(t)$  (dash). (g-i) Superimposed histograms of speed probability density functions and fitted density function of gamma distribution (solid red) with average parameters  $k$  and  $\theta$  (see text for details). (j-l) Superimposed histogram of relative frequency of normalized velocities.

corresponding values of  $M$  is virtually identical to the case with the unmodified probability rates (Figure 7 (g-i) and Table 2). The uniform distribution of normalized velocities also remained unchanged (Figure 7 (j-l)). These results suggest that in the absence of spatial cues, the distribution of speeds for a given adhesiveness (represented by the total number of adhesions  $M$ ) remain invariant under symmetry breaking of adhesion binding, while the diffusion type (normal vs. anomalous) does not. Thus, the adhesion number and its turnover is a major determinant of the cell speed, which is consistent with [48].

As cell polarization is required for migration even in the absence of external signals, it is not surprising that our results show that an imbalance of adhesion formation within a cell leads to experimentally observed superdiffusive scaling of the squared displacement [17],[36], [37]. Nevertheless, this highlights a potential mechanism of anomalous diffusion. In the following, we examine whether our model gives biologically consistent results in the case of externally induced polarization.

## 6.2 External cue gradient

We first investigate how cell trajectories are varied in the presence of an external cue gradient. If a cue, for example, is a chemoattractant, then it is well known that adhesion formation in a cell is biased in the direction of the attractant. Thus, to simulate such biased migration, we let the functions  $Q_{cue}$  and  $q$  to have the following form (recall equation (5.3)):

$$Q_{cue}(\mathbf{x}) = \begin{cases} 1 + \delta_E x_2, & \text{if } x_2 \geq 0 \\ 1, & \text{else} \end{cases}$$

$$q(Q_{cue}(\mathbf{x})) = Q_{cue}(\mathbf{x}),$$

where  $\delta_E$  represents the gradient magnitude and  $x_2$  is the second component of  $\mathbf{x}$ . Here, for simplicity we took the identity function for  $q$  and linear a cue gradient in the  $y$  coordinate. This cue can represent, for example, density of ECM or concentration of a chemoattractant. Thus, we simulate, respectively, hapto- or chemotactic migration.

M	8			16			32		
$\delta_E$	0.05	0.1	0.15	0.05	0.1	0.15	0.05	0.1	0.15
$\bar{\beta}, 1$	1.2551	1.5051	1.52	1.3405	1.6963	1.7545	1.5427	1.7722	1.8569
$s_{av},$ $\mu m/min$	1.8136	1.9133	2.0365	2.5235	2.5972	2.6089	3.5819	3.4218	3.3074
$\beta_0,$ $\mu m^2/min^{\bar{\beta}}$	1.0697	0.4625	0.6845	1.0312	0.3496	0.3654	0.8319	0.3412	0.2242

Table 3: Parameters obtained from the simulations with varying  $\delta_E$ .




Figure 8: Simulation results with  $M = 8, 16, 32$  adhesions in the first, second, and third columns respectively, and with various values of  $\delta_E$ . (a-c) Trajectories of 27 cell centroids  $\mathbf{x}(t)$  with  $\delta_E = 0.1$ . (d-f) Mean-squared displacements  $msd(t)$  (solid) and fitted  $\widehat{msd}(t)$  (dash). (g-i) Time-averaged exponents  $\beta_{av}(t)$

In the presence of a cue gradient, we see that the cell trajectories, shown in Figure 8 (a-c), exhibit a clear trend in the direction of an increasing concentration. The corresponding plots of the mean-squared displacements show the superdiffusive time scaling in Figure 8 (d-f), with the exponent  $\bar{\beta} > 1$  for all cases. Notice that as the number of adhesion sites  $M$  increases, so does  $\bar{\beta}$  for the same  $\delta_E$  (see Table 3). Together with the trajectory plots in Figure 8, our results suggest that in the presence of an external gradient, the taxis becomes more prominent and a cell more sensitive to a cue for increasing number of FAs. Moreover, comparing with the case of a uniform environment, we see that although the amoeboid motility is more diffusive in the absence of external cues, it is also more regular




Figure 9: Superimposed histograms of speeds, velocities and adhesion events with  $M = 8, 16, 32$  adhesions in the first, second, and third columns respectively, and with various values of  $\delta_E$ . (a-c) Speed probability density functions and fitted density function of gamma distribution (solid red) with average parameters  $k$  and  $\theta$  (see text for details). (d-f) Relative frequency of normalized velocities. (g-i) Ratio of the number of binding to unbinding events in each sector, such that any given time, only one adhesion site is in each sector

and directed when a cue gradient is present (see Tables 1, 2 vs. Table 3 and Figures 6, 7(a-c) vs. 8(a-c)). In Figure 8 (g-i) we see that the evolution of time-averaged exponents  $\beta_{av}(t)$  (see Appendix B) have three phases. Following the rapid increase in the first phase, there is a gradual decrease in the rate of change in the second phase, followed by stabilization of  $\beta_{av}(t)$  at  $\bar{\beta}$ . Curiously, a similar behavior has also been observed by Dieterich et al. [17].

The distribution of speeds again remained invariant and the average speeds are very close to the cases with no external cues (see Table 3). However, the frequency of normalized

velocities (see Figure 9 (d-f)) show, as expected, that the cell velocities are aligned with the cue gradient. Recall that in the presence of, for example, a chemotactic cue, a cell polarizes so that its adhesion dynamics is aligned with the gradient. In particular, adhesions are preferentially formed at the front (where the chemoattractant concentration is larger), and preferentially ruptured at the back. We can see in Figure 9 (g-i), that our simulation results reproduce such polarized dynamics: the ratio of binding to unbinding events is larger (smaller) than unity in the northern (southern) part of the cells, where the cue is stronger (weaker) relative to cell centroid. Also, for a smaller number of adhesion sites, the effects of increasing the cue gradient are more noticeable effect on the ratios of events (see 9 (g-i)). This is simply due to the increased density of adhesion sites, which leads to smaller relative difference of the cue between them.

### 6.3 Fibrillar architecture of ECM

ECM topography is another important determinant of directed cell migration. In particular, spatial distribution of the ECM fibers guides the motility by inducing cell shape alignment along the adhesive cues, resulting in characteristic directed movement along the fiber tracts [50]. Such guided migration is called contact guidance [50], [52]. Ramirez-San Juan et. al. [52] showed that contact guidance can be modulated by micrometer scale variations of interfiber spacing. Inspired by this study, we simulate how subcellular scale fiber spacing influences cell motility, and whether such ECM architecture yields migration patterns characteristic of contact guidance.




Figure 10: Stripe pattern with  $\delta_G = 0.15, 0.25, 0.35$  on, respectively, left, middle, and right plots. A cell is illustrated such that each FA on a stripe is bound and cell center  $\mathbf{x}$  coincides with  $\mathbf{x}_n$  at the origin.

Similar to the case with an external cue gradient, the functions  $Q_{cue}$  and  $q$  have the following form:

$$Q_{cue}(\mathbf{x}) = \begin{cases} 1, & \text{if } \mathbf{x} \in \Omega_{\delta_G} \\ 0.01, & \text{else} \end{cases}$$

$$q(Q_{cue}(\mathbf{x})) = Q_{cue}(\mathbf{x}),$$



where  $\Omega_{\delta_G}$  represents the stripe pattern,  $\delta_G = 0.15, 0.25, 0.35$  represents the spacing between stripes such that the distances between them is  $\delta_G R_{cell}$  (Figure 10). The stripe width is taken to be  $0.25 R_{cell}$ . Similarly as in [52], these dimensions are chosen so that a cell is spread on multiple stripes.




Figure 11: Simulation results with  $M = 16$ , and  $\delta_G = 0.15, 0.25, 0.35$  in first, second, and third columns, respectively. (a-c) Trajectories of 7 cells and the striped ECM pattern. (d-f) Relative frequencies of normalized velocities. (g-i) Relative frequencies of binding events in each of the 16 cell sectors.

The simulation results, shown in Figure 11, indicate that the cell motility has characteristics of contact guidance. Namely, the trajectories show preferential horizontal cell movement (Figure 11 (a-c)), and the displacements are aligned with the fiber pattern (Figure 11 (d-f)). However, increasing the spacing does not simply lead to a greater adhesion alignment along the horizontal direction, as can be observed in Figure 11 (g-i). Rather,

it is the combination of the ECM pattern and the radial position of FAs that gives rise to, for example, definite x-shaped adhesion binding patterns (Figure 11 (h)). Such binding (and unbinding) pattern leads to fluctuating movement along northwest-southeast and northeast-southwest axis, with the resulting net migration pattern shown in Figure 11 (b). Similarly, the binding pattern shown in Figure 11 (i) with more frequent events along the equator corresponds to a mixture of diagonal and horizontal movements (Figure 11 (c)), as larger interfiber spacing precludes FA binding at the poles and facilitates adhesion along, as well as across the stripes in x-shaped pattern (see also Figure 10 (right) for illustration of a characteristic FA configuration). On the other hand, smaller spacing also leads to horizontal movement, but with more frequent vertical displacement across the stripes (Figure 11). We also found that the average speeds were lower than in previous scenarios (Tables 2, 3):  $1.52\mu\text{m}/\text{min}$ ,  $0.94\mu\text{m}/\text{min}$ , and  $0.87\mu\text{m}/\text{min}$  corresponding to, respectively,  $\delta_G = 0.15, 0.25, 0.35$ . Interestingly, the average speeds reported in [52] were  $\sim 0.6\mu\text{m}/\text{min}$ , although in that study the speeds were nearly constant for varying fiber pattern.

Altogether, our simulations of contact guidance are, for the most part, consistent with the observations reported in the literature. In particular, we obtain the expected guidance of cell movement (Figure 11 (a-f)) and the geometric constraint of adhesion sites (Figure 11 (g-i)) by the fibrillar ECM pattern, in agreement with [50], [52]. Nevertheless, since our model does not explicitly take into account morphological changes in cell shape (recall that in our model cell shape is normalized to a circle; see Section 2) and since cell shape control is essential to contact guidance [50], [52], increasing the interfiber distance does not necessarily lead to greater alignment of cell migration along the ECM fibers in our simulations<sup>1</sup>.

## 6.4 Asymmetric contractility

We now investigate how cell motility is influenced by asymmetrical contractile forces in a cell. Along with preferential adhesion formation, due to, for example, chemotactic gradient, formation of cell rear by increased actomyosin contractile activity serves as an alternative mechanism by which a directed migration can be induced in the absence of such gradient [12]. In particular, local stimulation of contractility leads to directed motility in the direction opposite to the stimulated area, even in the absence of response to chemotactic stimuli [68]. Here we show that our model is also capable of capturing such directed movement, triggered by breaking myosin mediated contractile symmetry.

Recall that  $T_i$  in equation (2.1) denotes the force generated by myosin motors at an adhesion site  $i$ . Instead of taking it constant, we let it vary with the radial position of an FA. Namely, let  $T_i : [0, 2\pi) \rightarrow \mathbb{R}_+$  be defined as:

$$T_i(\theta) = \begin{cases} (1 + \delta_{myo})T_{i_0}, & \text{if } \pi < \theta + (i - 1)\frac{2\pi}{M} < 2\pi \\ T_{i_0}, & \text{else} \end{cases},$$

---

<sup>1</sup>The values of the guidance parameter  $G$ , defined as in [52] (See Appendix B for details), were found to be 0.64, 0.70, 0.64, corresponding to, respectively,  $\delta_G = 0.15, 0.25, 0.35$ .

where  $\delta_{myo} = 0.35, 0.40, 0.45$ ,  $T_{i_0}$  is the constant value used in previous simulations (see Appendix A), and  $\theta + (i - 1)\frac{2\pi}{M}$  is the radial position of the  $i^{\text{th}}$  FA. Thus, contractile forces south of cell equator are larger by 35%, 40%, 45% for corresponding values of  $\delta_{myo}$ . We should, therefore, expect in our simulations that the northern part of a cell becomes the front due to the imposed contractile symmetry breaking, and that cells will move accordingly (see Figure 12 for illustration).




Figure 12: A schematic representation of asymmetric contractility. (Bottom row) Increased contractility causes  $\mathbf{x}_n$  (blue circle) to shift south of otherwise equilibrium point in the center. (Top row) Preferential unbinding of FAs south of equator leads to directed movement indicated by blue arrows. See also Figure 4 (II' and III') for schematic representation of cell motility in case of unbinding event.

Indeed, Figure 13 (a-c) shows, as expected, the trajectories of cells maintaining north-south polarity corresponding to, respectively, front and rear. Since the asymmetry of myosin forces remained during the simulations, the cell's north-south polarity also persisted, resulting in the cell movement that was highly directed along this axis, consistent with [68]. Consequently, we obtain higher values of  $\beta_{av}(t)$ , as shown in Figure 13 (d-f). In particular, for  $\delta_{myo} = 0.45$ , we see that the time scaling of the mean squared displacement is close to ballistic (see Table 4 for values of  $\bar{\beta}$ ). Moreover, increasing the number of FAs leads to more polarized, directed migration. As in the previous cases, neither speed averages (Table 4) nor their distribution (data not shown) changed significantly for a given number of focal adhesions. Interestingly, for  $\delta_{myo} = 0.35$ , the binding is relatively more frequent in the rear (i.e. south of equator) than in the front, and unbinding is relatively more frequent in the front (i.e. north of equator) than in the back (Figure 13) (g-i). This suggests, then, that cells were preferentially moving in the southern direction. However, as can be seen in Figure 14, this is not the case. Although movements southwards is more frequent (due to the above-mentioned event frequencies), the speeds are lower than north-




Figure 13: Simulation results with  $M = 8, 16, 32$  adhesions in the first, second, and third columns respectively, and with various values of  $\delta_{myo}$ . (a-c) Trajectories of 13 cells with  $\delta_{myo} = 0.4$ . (d-f) Time-average exponents  $\beta_{av}$ . (g-i) Ratio of the number of binding to unbinding events in each sector.

ward movements: the ratios of the average speeds directed north to the average speeds directed south were found to be 1.0165, 1.0181, and 1.0858 corresponding to, respectively,  $M = 8, 16, 32$ . The net effect is northward movement. For higher values of  $\delta_{myo}$ , we see that the unbinding is, expectedly, more frequent in the rear, while binding is preferentially in the front. This suggests that there is some threshold value of  $T_i$ , above which we obtain the simulation results that are consistent with the experimental observations. Indeed, the values of  $\delta_{myo}$  correspond to  $T_i = 0.98F_b, 1.0182F_b, 1.0545F_b$ , where  $F_b$  is the characteristic force scale (see Appendix A for the details on the parameter values). Recalling Section 5.2.2, we can see that for forces above  $F_b$ , the unbinding probability rate dominates the

M	8			16			32		
$\delta_{myo}$	0.35	0.40	0.45	0.35	0.40	0.45	0.35	0.4	0.45
$\bar{\beta}, 1$	1.3072	1.6325	1.7759	1.1311	1.7905	1.8524	1.4650	1.8892	1.9353
$s_{av},$ $\mu m/min$	1.6230	1.5639	1.5103	2.3742	2.3798	2.3516	3.5861	3.7414	3.7701
$\beta_0,$ $\mu m^2/min\bar{\beta}$	0.7767	0.3659	0.2493	2.3088	0.2893	0.3037	1.0713	0.9894	1.2787

Table 4: Parameters obtained from the simulations with varying  $\delta_{myo}$ .

binding rate, leading to the observed simulation results.




Figure 14: Simulated trajectories with  $M = 8, 16, 32$  adhesions with  $\delta_{myo} = 0.35$  on left, middle, and right plots, respectively.

**Remark.** Another way to induce contractile asymmetry is, for example, to decrease the myosin force  $T_i$  north of the cell's equator. Then, again, the south of the cell equator is more contractile. However, the simulated trajectories show southward directed movement (data not shown), contrary to what we should expect. Therefore, merely inducing contractile asymmetry is not sufficient. For the expected directed migration to occur, there must be a local increase of contractile forces above some critical level in the prospective cell rear, rather than a local decrease of contractility in the prospective front. Interestingly, Yam et al. [68] were able to initiate directed movement by increasing local actomyosin contraction, while locally decreasing the contractile activity did not lead to migration initiation. More recently, Shellard et al. [59] showed that directed collective cell migration of neural crest cells requires greater contractility at the rear of the clump.

## 7 Discussion and outlook

In this paper we constructed a stochastic model of cell migration using a minimal representation of cellular structures, essential for crawling, such as stress fibers and focal adhesions. Using this representation, and observing that FA assembly and disassembly events of the migration cycle lead to different migratory outcomes, we obtained the equations describing deterministic cell motion between the random occurrence of FA events. After introducing the rates of FA binding and unbinding, we obtained the remaining necessary objects to define a piecewise deterministic Markov process: the distribution of interarrival times and of the next FA event. Note that the forms of these distributions have been derived, rather than simply postulated. Having specified the coupling between SFs and FAs, as well as between the cellular environment and FAs, we performed numerical simulations. We showed that our model is able to reproduce experimental observations, such as: superdiffusive scaling of the mean-squared displacement [17], [36], [37] (Figure 7); biased motility in the presence of external cue (Figure 8); contact guidance [52] (Figure 11). In these cases, the obtained results followed solely due to asymmetric, dynamic instability of FAs in direct response to environmental stimuli. Specifically, it is only the biased FA assembly rate that drives biased cell motility along the cue gradient or the fiber tracts (Figures 9 and 11 (d-i)). That is, preferred velocities were not imposed or chosen in any way, but simply followed as a consequence of front-rear polarity, as the cell front is characterized by preferential FA binding and the rear by unbinding.

Another characteristic of directed migration is the asymmetric contraction of actomyosin bundles. By increasing the force generation of myosin motors in the prospective rear, we obtained directed movement in the opposite direction (Figure 14). Here asymmetric FA dynamics (and so front-rear polarity) was also obtained, but as a consequence of locally induced contractile activity, consistent with [68].

Our simulation results in various settings suggest that the cell speeds follow a gamma distribution (Figures 6 (g-i), 7 (g-i), 9 (a-c)). Furthermore, the number of adhesion sites seems to be a determinant of the gamma distribution, as its parameters are similar under different settings and given number of FAs. These results suggests that cell speeds are independent of biased FA formation, i.e. the bias only alters the directionality and not the speed. It is also interesting to see a correlation between the number of adhesion sites and diffusivity (Table 1), as well as average speed (Tables 1-4). Note that faster and diffusive amoeboid movement is characterized by an increased number of weaker adhesions with high turnover and contractility [48]. Thus, the aforementioned correlation is also consistent with experimental observations. We note that our model is not fit to take into account motility, strongly relying on cell shape control, which is required, for example, in highly mobile cells. However, the simulations reproduce migration along fiber tracts, where cell reshaping takes place [52]. Our results suggest, then, that adhesion along the tracts is sufficient to produce such migration patterns.

Although the model of the internal contractile machinery driving cell locomotion and cytoskeletal remodeling is simple, the resulting numerical simulations explain several experimental observations. Moreover, the cyclical nature of cell motility is captured with our

piecewise deterministic model. Note also that the FA dynamics in our model is independent of a particular cellwise contractility mechanism: only local mechanochemical interactions are relevant. Thus, one possible avenue of improvement is the inclusion of a more complex cell contractility apparatus, which could, for example, regulate shape changes. This can be done, for example, with the help of an active cable network model [27]. However, given the relative simplicity of the stochastic model and its ability to explain a handful of the experimental observations, it is possible to extend the model to include cell-cell collisions in the context of contact inhibition of locomotion (CIL). Here, the collisions lead to cessation of locomotion and to repolarization, such that the formation of new adhesions at the site of contact is inhibited, while contractility is stimulated [55]. Within the framework of our model this can be implemented in a straightforward manner: collisions cause a switch to a non-moving state and the FA probability rates are modified according to contact location. Yet another extension is obtained by treating cells as particles and using kinetic theory, yielding equations governing population migration. Thus, we can achieve a multiscale description of cell motility. Due to limitations in size and scope, cell-cell collisions and population migration will be treated in forthcoming works.

## Acknowledgement

The author acknowledges support of the German Academic Exchange Service (DAAD).

## Appendix A Parameter assessment

Note that the length of myosin mini-filaments is  $\sim 0.3\mu m$  [62] and the interfilament distance is  $\sim 1\mu m$  in an uncontracted fiber [3]. Assuming vanishing interfilament distance (see Figure 2 for illustration and [44] for a review on actomyosin contraction mechanism), then the proportion of the minifilaments to the initial, uncontracted SF length is  $\frac{0.3}{1+0.3} = 0.23$ . Kassianidou et. al. [30] showed that the retraction length scales linearly with the initial length. If the interfilament distance vanishes, the front myosin motor cannot perform a power stroke and step forward, which renders the single motor and the entire minifilament unable to apply contractile stress. Therefore, taking the initial length to be  $\sim R_{cell}$ , we estimate the critical length  $L_c = 0.2R_{cell}$ . Interestingly, Deguchi et. al. [15] found that stress fibers contract, on average, to 20% of their original length. As stress fibers generally span more than half of a cell, and since it was found that there is a preexisting strain [15] in them, we estimate  $L_0 = 1.1R_{cell}$ .

If we take  $R_{cell} = 25\mu m$ , and assuming a SF has at rest the length of  $R_{cell}$ , we can estimate the number of myosin minifilaments in a fiber to be  $25\mu m/1.3\mu m \approx 20$ . It was estimated in [64] that there are 10-30 myosin motors in each minifilament. As each motor produces a force of  $2 - 10pN$  [20], [43], [44], we then estimate  $T_i = 4nN$ . Balaban et. al. [5] found that focal adhesions apply a constant stress of  $\sim 5.5nN/\mu m^2$  over an area of  $\sim 1\mu m^2$  on an elastic substrate. Thus, we take  $F_b = 5.5nN$ . Assuming a preexisting strain

was 0.1 when these measurements were taken, and since  $T_i = 4nN$ , we then estimate the one dimensional Young's modulus  $EA = 15nN$ .

Using Stokes' Law for drag in low Reynolds number regime, the drag coefficient  $\beta_{ECM}$  can be estimated as:

$$\beta_{ECM} = 6\pi\eta_{ECM}R_{cell},$$

where  $\eta_{ECM}$  is the dynamic viscosity of the environment. Assuming that the viscosity  $\eta_{ECM}$  is higher than that of water, and taking into account that the contact between cell surface and the substrate increase the effective friction, we estimate  $\beta_{ECM} \approx 10 - 10^3 \frac{N \cdot s}{m^2} \times R_{cell}$ . Similarly, due to low Reynold number, the rotational drag coefficient  $\beta_{rot}$  is given by:

$$\beta_{rot} = 8\pi\eta_{ECM}R_{cell}^3 \approx 10 - 10^3 \frac{N \cdot s}{m^2} \times R_{cell}^3.$$

In order to obtain estimates for the drag coefficient  $\beta_{cell}$  one needs to have an estimate of cytoplasm viscosity. Assessing the effective cytoplasmic viscosity of migrating cells is a challenging task, since the viscoelastic properties of cytoskeleton (which, among other things, consists of polymer networks) are highly dynamic due to constant remodeling and spatiotemporal mediation of the rheology by various signaling pathways. Particularly, the actin network bundle size and cross-linkers influence viscoelastic properties [23]. Furthermore, the effective viscosity experienced by an experimental probe (or a protein) in polymer solutions depends not only on the type of material properties of the fluid, but also on the size of a probe <sup>1</sup> [29]. Inferring that the body being dragged in the cell is the nucleus with radius  $R_{nucleus}$ , we estimate:

$$\beta_{cell} = 6\pi\eta_{cell}R_{nucleus} \approx 10 - 10^2 \frac{N \cdot s}{m^2} \times R_{nucleus},$$

where  $\eta_{cell}$  is the cytoplasm viscosity.

Note that a focal adhesion is a cluster of transmembrane receptors (integrins) linking the substrate with the cytoskeleton, which is always under tension. The cluster also includes adapter proteins, which interlinks these receptors with the cytoskeleton (see e.g. reviews [8], [24]). Thus, if there is no load on a focal adhesion, then, since the cytoskeleton is always under tension, such a focal adhesion can be treated as a complex of independent bonds to the substrate without a link to SF. In the absence of a load, the average cluster lifetime  $T_{life}$  is given by [19]:

$$T_{life} = \frac{1}{k_0^1 + k_{on}^1} \left[ H_N + \sum_{n=1}^N \left( \frac{k_{on}^1}{k_0^1} \right)^n \frac{1}{n} \binom{N}{n} \right],$$

where  $N$  is the number of bonds in a cluster,  $H_N$  is the  $N$ th harmonic number,  $k_0^1$  and  $k_{on}^1$  are, respectively, unbinding and binding rates of integrins under no load. Note that in

---

<sup>1</sup>Using a small molecule as a probe, the cytoplasm viscosity was found to be  $\simeq 2 - 3 \times 10^{-4} Pa \cdot s$  [40]. With larger probes, the viscosity was found to be  $\simeq 2 - 4 \times 10^{-2} Pa \cdot s$  [29] and  $\simeq 5 \times 10^{-2} Pa \cdot s$  [4]



Parameter	Value	Parameter	Value
$T_i$	$4nN$	$R_{cell}$	$25\mu m$
$EA$	$15nN$	$L_0$	$27.5\mu m$
$F_b$	$5.5nN$	$L_c$	$5\mu m$
$k_{off}^0$	$0.05s^{-1}$	$\beta_{rot}$	$3.25 \times 10^{-15} N \cdot s \cdot m$
$k_{on}^0$	$0.01s^{-1}$	$\beta_{ECM}$	$5 \times 10^{-4} \frac{N \cdot s}{m}$
$R_{nucleus}$	$5\mu m$	$\beta_{cell}$	$5 \times 10^{-3} \frac{N \cdot s}{m}$

Table 5: Parameters used for simulation

the absence of a load, (re-)binding of individual integrins is an independent event, which bears no relation to the FA, since tension is required for an FA to form and sustain itself. We then estimate:

$$k_{off}^0 = 1/T_{life}|_{k_{on}^1=0} = k_0^1/H_N.$$

Li et al.[35] found that  $k_0^1 = 0.012s^{-1}$  for  $\alpha_5\beta_1$ -integrin binding to fibronectin. For  $N = 10^3 - 10^4$  we estimate that  $k_{off}^0 = 0.05s^{-1} - 0.1s^{-1}$ .

## Appendix B Data analysis

Given that the time interval  $[0, t_{end}]$  is divided into  $n_{time}$  subintervals of equal length  $\Delta t$  and given the positions  $\mathbf{x}_i(t_j)$  of cell  $i$  at the time points  $t_j := j\Delta t$ ,  $j = 0, \dots, n_{time}$ , the mean-squared displacement  $msd_i(t_j)$  of cell  $i \in \{1, \dots, n_{cells}\}$  over a time interval of length  $t_j$  is given by:

$$msd_i(t_j) := \frac{1}{n_{time} - j} \sum_{k=1}^{n_{time}-j} \|\mathbf{x}_i(t_{j+k}) - \mathbf{x}_i(t_k)\|^2, \quad (B.1)$$

where  $j = 1, \dots, n_{time} - 1$  and  $n_{cells}$  is the total number of cells. Then, the mean-squared displacement  $msd(t)$  of an ensemble of cells over time interval of length  $t_j$  is defined by:

$$msd(t) := \frac{1}{n_{cells}} \sum_{i=1}^{n_{cells}} msd_i(t_j). \quad (B.2)$$

**Remark.** In general, the (time-averaged) mean-squared displacement  $\langle d^2(t, T) \rangle$  of a particle trajectory  $\mathbf{x}(t)$  at the time  $t$ , time endpoint  $T$  is formally defined as:

$$\langle d^2(t, T) \rangle = \frac{1}{T-t} \int_0^{T-t} \|\mathbf{x}(s+t) - \mathbf{x}(t)\|^2 ds. \quad (B.3)$$

For an ergodic process, we have:

$$\lim_{T \rightarrow \infty} \langle d^2(t, T) \rangle = \langle \mathbf{x}^2(t) \rangle,$$

where  $\langle \mathbf{x}^2(t) \rangle$  is formally defined as:

$$\langle \mathbf{x}^2(t) \rangle = \int \mathbf{x}^2 P_t(d\mathbf{x}),$$

and  $P_t(d\mathbf{x})$  is the probability measure of the underlying stochastic process at time  $t$ . That is, for an ergodic process, and for sufficiently long times, the time average equals the phase space average. However, our cell motility process need not be ergodic and hence, using a quadrature to evaluate the integral in equation (B.3), we obtain time average displacement in equation (B.1). To smooth out trajectory-to-trajectory fluctuations, we then average the displacements over all trajectories in equation (B.2).

For a diffusive motion we expect that  $msd(t) \sim t^{\beta(t)}$  with  $\beta(t) \approx 1$ , while for a ballistic motion  $\beta(t) \approx 2$ . Since  $msd(0) = 0$ , we can estimate the exponent  $\beta(t)$  for  $t \in [\Delta t, t_{end} - \Delta t]$  from the simulated data as:

$$\beta(t) = t \frac{d}{dt} \ln msd(t).$$

Although averaging reduces fluctuations, it does not eliminate them completely. Thus, computing the derivative above will yield a result that may oscillate wildly. Then, in order to investigate how  $\beta$  varies with time, we define the time average  $\beta_{av}(t)$  over the interval  $[\Delta t, t]$  as:

$$\beta_{av}(t) := \frac{1}{t - \Delta t} \int_{\Delta t}^t \beta(s) ds = \frac{1}{t - \Delta t} \left( s \ln msd(s) \Big|_{\Delta t}^t - \int_{\Delta t}^t \ln msd(s) ds \right),$$

where  $t \in [2\Delta t, t_{end} - \Delta t]$ . Then,  $\bar{\beta} := \beta_{av}(t_{end} - \Delta t)$  estimates the time scaling of  $msd$  over the whole time interval. To assess how well  $\bar{\beta}$  reflects the scaling of  $msd$ , we define the following function  $\widehat{msd}(t) := \beta_0 t^{\bar{\beta}}$ , where  $\beta_0$  is found by minimizing the square error:

$$\begin{aligned} \min_{\beta_0} \frac{1}{2} \sum_{j=1}^{n_{time}-1} \left( \beta_0 t_j^{\bar{\beta}} - msd(t_j) \right)^2 &\Rightarrow \\ \beta_0 &= \frac{\sum_{j=1}^{n_{time}-1} msd(t_j) t_j^{\bar{\beta}}}{\sum_{j=1}^{n_{time}-1} t_j^{2\bar{\beta}}}. \end{aligned}$$

We refer to Section 6 for comparisons between  $msd$  and  $\widehat{msd}$ .

Note that because binding events can occur, a cell need not to have moved between the two time points  $t_j$  and  $t_{j+1}$ . Thus, the speed between the consecutive time points may be zero for many time points, which would give an inaccurate statistical assessment of cell speeds. In order to estimate the speeds of a cell  $i$  we use the following procedure:

First, we find  $l_i$ , given by:

$$l_i := \min \{l \in \mathbb{N} : \mathbf{x}_i(t_{l+k}) \neq \mathbf{x}_i(t_k), 0 \leq k < n_{time}, l+k \leq n_{time}\}.$$

Then we find the set of speeds  $S_i$  as:

$$S_i := \left\{ s \in \mathbb{R}^+ : s = \frac{\|\mathbf{x}_i(t_{(k+1)l_i}) - \mathbf{x}_i(t_{kl_i})\|}{l_i \Delta t}, k \in \mathbb{N}, (k+1)l_i \leq n_{time} \right\}.$$

This simply means that to compute speeds we only use a (minimal) time interval, such that a cell  $i$  is guaranteed to change its position. The total set of speeds for  $n_{cells}$  is  $S := \cup_{i=1}^{n_{cells}} S_i$ . The average speed  $s_{av}$  is then defined as arithmetic average:

$$s_{av} := \frac{1}{|S|} \sum_{s \in S} s.$$

The set of normalized velocities  $V_i$  (or, alternatively, displacements) of cell  $i$  is given by:

$$V_i := \left\{ \mathbf{v} \in \mathbb{R}^2 : \mathbf{v} = \frac{\mathbf{x}_i(t_{(k+1)l_i}) - \mathbf{x}_i(t_{kl_i})}{\|\mathbf{x}_i(t_{(k+1)l_i}) - \mathbf{x}_i(t_{kl_i})\|}, k \in \mathbb{N}, (k+1)l_i \leq n_{time} \right\},$$

and the total set of normalized velocities  $V$  is given by  $V := \cup_{i=1}^{n_{cells}} V_i$ .

We define the guidance parameter  $G \in [0, 1]$  similarly as in [52]:

$$G := \frac{1}{|\Theta|} \sum_{\theta \in \Theta} g(\theta),$$

where  $\Theta := \cup_{i=1}^{n_{cells}} \Theta_i$ . The set  $\Theta_i$  of angles between a displacement vector of cell  $i$  and the ECM fibers is defined by

$$\Theta_i := \left\{ \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] : \theta = \arcsin \left( \frac{x_{i,2}(t_{(k+1)l_i}) - x_{i,2}(t_{kl_i})}{\|\mathbf{x}_i(t_{(k+1)l_i}) - \mathbf{x}_i(t_{kl_i})\|} \right), k \in \mathbb{N}, (k+1)l_i \leq n_{time} \right\},$$

where  $x_{i,2}$  is the  $y$ -component of  $\mathbf{x}_i$ . The function  $g : \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow \{0, 1\}$  is given by

$$g(\theta) = \begin{cases} 1, & \text{if } |\theta| \leq \pi/4 \\ 0, & \text{else} \end{cases}.$$

Thus,  $G$  increases when displacements are aligned with the horizontal axis.

## Appendix C Simulation of the PDMP

In order to compute the trajectories of the cell motility process, the following algorithm is used:

---

**Algorithm 1** Simulation of the PDMP
 

---

1. Set  $(\nu_0, \mathbf{X}_0) \in A \times \Gamma$  and  $t = T_0 = 0$
2. For  $k = 0, 1, \dots$   
 Generate interarrival time  $\Delta_k = T_{k+1} - T_k$ , whose distribution function is given by:

$$\mathbb{P}(\Delta_k \leq \tau) = 1 - \exp\left(-\int_t^{t+\tau} a_0(\nu_t, \phi_\nu(s, \mathbf{X}_t)) ds\right) \quad (\text{C.1})$$

Compute  $\mathbf{X}_{t+s^-} := \phi_\nu(s, \mathbf{X}_t)$   
 Set  $t = T_{k+1} = T_k + \Delta_k$   
 Generate  $(\nu_t, \mathbf{X}_t) \sim Q(\cdot; (\nu_{t^-}, \mathbf{X}_{t^-}))$

---

To generate the interarrival time  $\Delta_k$ , we need to solve for  $\tau$  in the following equation:

$$f(\tau) := \int_t^{t+\tau} a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) ds + \ln(1 - u) = 0, \quad (\text{C.2})$$

where  $u$  is uniformly distributed on the interval  $(0, 1)$ . Notice that the evaluation of the integral by a quadrature rule requires computing the solution  $\mathbf{X}_{t+s} = \phi_{\nu_t}(s, \mathbf{X}_t)$  up to time  $s$ , where  $s$  is a quadrature point. Moreover, using an iterative method to solve (C.2) requires computing the integral at each iteration. Therefore, it is imperative to devise an efficient method to sample from distribution (C.1). In the following, we propose a general method to generate the next event time.

### C.1 Simulation of the next event time

Let  $T_k \leq t < T_{k+1}$  and let  $\mathbf{G}(\cdot; h) : \Gamma \rightarrow \Gamma$  represent a numerical method to solve the ODE system

$$\frac{d}{dt} \mathbf{X}_t = \mathbf{H}_{\nu_t}(\mathbf{X}_t)$$

for one time step  $h$ . That is,  $\mathbf{X}_{t+h} = \mathbf{G}(\mathbf{X}_t; h)$  is the numerical solution of the above ODE system at time  $t + h$ .

Let  $[T_k, T_{k+1})^2 \ni (s', t') \mapsto A_0(t', s') := \int_{s'}^{t'} a_0(\nu_t, \phi_{\nu_t}(u, \mathbf{X}_t)) du$  denote the integrated rate function.

The method to find the root  $\tau$  of equation (C.2) is given in Algorithm 2. First, in steps (1-22), we find the upper bound  $\tau_{max}$  by solving the ODE system for  $n$  steps with step size  $h$  and store the solution, the computed rate  $a_0$ , and the integrated rate  $A_0$  at these times steps. Then, for any  $\tau \leq \tau_{max}$  we can compute  $A_0(t + \tau, t)$  upon using the stored value of

$A_0$  at time  $t + \tau_i$ , where  $i = \lfloor \frac{\tau}{h} \rfloor$  and  $\tau_i = ih$ :

$$\begin{aligned} A_0(t + \tau, t) &= \int_t^{t+\tau} a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) ds \\ &= \int_t^{t+\tau_i} a_0(\nu_t, \mathbf{X}_{t+s}) ds + \int_{t+\tau_i}^{t+\tau_i+\tau-\tau_i} a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) ds \\ &= A_0(t + \tau_i, t) + \int_{t+\tau_i}^{t+\tau_i+h_i} a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) ds. \end{aligned}$$

To compute the last integral in the expression above, we interpolate the integrand using the stored values  $a_0(\nu_t, \mathbf{X}_{t+\tau_i})$  and  $a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}})$ . Let

$$I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) := \int_{t+\tau_i}^{t+\tau_i+h_i} a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t))$$

denote the approximation of the integral using interpolated integrand. We can use the following interpolation methods for  $t + \tau_i < s < t + \tau_{i+1}$ :

1. Piecewise constant

$$\text{Forward: } a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) = a_0(\nu_t, \phi_{\nu_t}(t + \tau_i, \mathbf{X}_t)) = a_0(\nu_t, \mathbf{X}_{t+\tau_i}).$$

$$I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) = h_i a_0(\nu_t, \mathbf{X}_{t+\tau_i}). \quad (\text{C.3})$$

$$\text{Backward: } a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) = a_0(\nu_t, \phi_{\nu_t}(t + \tau_{i+1}, \mathbf{X}_t)) = a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}})$$

$$I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) = h_i a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}). \quad (\text{C.4})$$

2. Average:  $a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) = \frac{1}{2} (a_0(\nu_t, \mathbf{X}_{t+\tau_i}) + a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}))$ .

$$I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) = \frac{1}{2} h_i (a_0(\nu_t, \mathbf{X}_{t+\tau_i}) + a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}})). \quad (\text{C.5})$$

3. Piecewise linear:

$$a_0(\nu_t, \phi_{\nu_t}(s, \mathbf{X}_t)) = (s - t - \tau_i) \frac{a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}) - a_0(\nu_t, \mathbf{X}_{t+\tau_i})}{h} + a_0(\nu_t, \mathbf{X}_{t+\tau_i}).$$

$$\begin{aligned} I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) \\ = h_i a_0(\nu_t, \mathbf{X}_{t+\tau_i}) \left(1 - \frac{h_i}{h}\right) + \frac{h_i^2}{2h} a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}). \end{aligned} \quad (\text{C.6})$$

Thus,  $f(\tau)$  can be evaluated using equations (C.3)-(C.6):

$$f(\tau) = A_0(t + \tau_i, t) + I(a_0(\nu_t, \mathbf{X}_{t+\tau_i}), a_0(\nu_t, \mathbf{X}_{t+\tau_{i+1}}); h_i, h) + \ln(1 - u).$$

---

**Algorithm 2** Event time computation
 

---

```

1: procedure INITIALIZATION
2:   Input: Time  $t = T_k$ ;  $(\nu_t, \mathbf{X}_t)$ ; time step  $h$  and ODE method  $\mathbf{G}$ ;  $n, m \in \mathbb{N}$ .
3:   Set  $s_0 := 0, n_0 := 0$ , create  $List_{a_0}, List_{A_0}$ , and  $List_{\mathbf{X}}$ .
4:   Append  $List_{a_0} \leftarrow a_0(\nu_t, \mathbf{X}_t)$ ,  $List_{A_0} \leftarrow 0$ ,  $List_{\mathbf{X}} \leftarrow \mathbf{X}_t$ .
5:   Set  $\tau_{max} := nh$ ,  $s_0 := n_0h$ .
6:   Set initial condition  $\mathbf{X}_s \leftarrow List_{\mathbf{X}}[last]$ .
7:   Set  $A_0^0 := List_{A_0}[last]$ .
8:   for  $i = 1$  to  $n$  do
9:      $s_i := t + s_0 + ih$ .
10:    Compute  $\mathbf{X}_{s_i} := \mathbf{G}(\mathbf{X}_s; h)$  and  $a_0(\nu_t, \mathbf{X}_{s_i})$ .
11:    Compute  $A_0(s_i; t + s_0)$  with quadrature points  $s_j, j = 0, \dots, i$ .
12:     $A_0(s_i; t) := A_0^0 + A_0(s_i; t + s_0)$ .
13:    Append  $List_{a_0} \leftarrow a_0(\nu_t, \mathbf{X}_{s_i})$ ,  $List_{A_0} \leftarrow A_0(s_i; t)$ ,  $List_{\mathbf{X}} \leftarrow \mathbf{X}_{s_i}$ .
14:  end for
15:  Generate  $u \sim U(0, 1)$ .
16:  if  $List_{A_0}[last] < -\ln(1 - u)$  then
17:     $n := n + m$ .
18:     $n_0 = n_0 + m$ .
19:    go to 5.
20:  end if
21:  Output:  $\tau_{max}, List_{a_0}, List_{A_0}$ , and  $List_{\mathbf{X}}$ .
22: end procedure
23: procedure EVALUATION OF  $f$ 
24:   Input:  $\tau$ ; time step  $h$ ;  $List_{a_0}, List_{A_0}$ ; Integrated interpolation method  $I$ .
25:   Set  $i := \lfloor \frac{\tau}{h} \rfloor$  and  $h_i = \tau - ih$ .
26:   Output:  $f(\tau) = List_{A_0}[i] + I(List_{a_0}[i], List_{a_0}[i + 1]; h_i, h) + \ln(1 - u)$ 
27: end procedure
28: procedure EVENT TIME
29:   Find the root  $\tau$  of  $f(\tau) = 0$  using 23 and a root finding method.
30:    $i := \lfloor \frac{\tau}{h} \rfloor$ .
31:   Compute  $\mathbf{X}_{t+\tau} := \mathbf{G}(List_{\mathbf{X}}[i]; \tau - ih)$ .
32:   Output:  $\tau, \mathbf{X}_{t+\tau}$ 
33: end procedure

```

---

Using the interpolations above, we can now employ, for example, Newton's method to find the root of equation (C.2):

$$\tau_{l+1} = \tau_l - \frac{f(\tau_l)}{a_0(\nu_t, \phi_{\nu_t}(\tau_l, \mathbf{X}_t))},$$

or any other root-finding method (note that  $a_0 > 0$ ). Once the root is found, we simply advance the ODE system for one time step as described in Steps (28-33) of the Algorithm 2.

Note that we solve the ODE system for  $n + 1 = \tau_{max}/h + 1$  steps and we also solve for the interarrival time  $\tau$  primarily by using a look-up table. Moreover, obtaining a relatively sharp upper bound  $\tau_{max}$  does not yield a large computational overhead, since one simply can start the Algorithm 2 with a small  $n, m$ . Consequently, choosing an initial guess close to the sharp upper bound for Newton’s method results in a faster convergence. In case of thinning methods (see [34] or [7] for adaptive method), after each rejection one needs to solve the ODE system for time period that is, on average, approximately the same as  $\tau_{max}$  (in the best case scenarios for both methods, i.e. when the bound  $\tau_{max}$  and the bound for the rate function in thinning methods are sharp). Of course, these arguments hold when the computational cost of solving the ODE system is relatively large.

## C.2 Sampling from the transition measure

Given the time  $t$  of the next event and  $\mathbf{X}_t$  we need to sample from the transition distribution  $Q(\cdot, (\nu_{t-}, \mathbf{X}_{t-}))$ . Recalling Section 3.2 and the proof of Proposition 4.3, in order to sample from the transition measure it is sufficient to simulate the index  $j \in \{1, \dots, 2M\}$  of the next reaction, since the continuous component of the process does not jump. The discrete distribution of the next reaction index is given by equation (4.7):

$$\mathbb{P}(j|\alpha_2(\nu_{t-}), \mathbf{X}_{t-}) = \frac{a_j(\alpha_2(\nu_{t-}), \mathbf{X}_{t-})}{a_0(\alpha_2(\nu_{t-}), \mathbf{X}_{t-})}.$$

To simulate from the discrete distribution one can use fairly efficient Vose’s Alias Method [67].

## References

- [1] M. Abercrombie. The croonian lecture, 1978 - the crawling movement of metazoan cells. *Proceedings of the Royal Society of London B: Biological Sciences*, 207(1167):129–147, 1980.
- [2] A. Y. Alexandrova, K. Arnold, S. Schaub, J. M. Vasiliev, J.-J. Meister, A. D. Bershadsky, and A. B. Verkhovsky. Comparative dynamics of retrograde actin flow and focal adhesions: formation of nascent adhesions triggers transition from fast to slow flow. *PLoS one*, 3(9):e3234, 2008.
- [3] Y. Aratyn-Schaus, P. W. Oakes, and M. L. Gardel. Dynamic and structural signatures of lamellar actomyosin force generation. *Molecular Biology of the Cell*, 22(8):1330–1339, 2011.
- [4] D. Arcizet, B. Meier, E. Sackmann, J. O. Rädler, and D. Heinrich. Temporal analysis of active and passive transport in living cells. *Phys. Rev. Lett.*, 101:248103, Dec 2008.

- [5] N. Q. Balaban, U. S. Schwarz, D. Riveline, P. Goichberg, G. Tzur, I. Sabanay, D. Mahalu, S. Safran, A. Bershadsky, L. Addadi, et al. Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates. *Nature Cell Biology*, 3(5):466–472, 2001.
- [6] G. I. Bell. Models for the specific adhesion of cells to cells. *Science*, 200(4342):618–627, 1978.
- [7] A. Bouchard-Côté, S. J. Vollmer, and A. Doucet. The bouncy particle sampler: A non-reversible rejection-free Markov chain Monte Carlo method. *ArXiv e-prints*, Oct. 2015.
- [8] K. Burridge and M. Chrzanowska-Wodnicka. Focal adhesions, contractility, and signaling. *Annual Review of Cell and Developmental Biology*, 12(1):463–519, 1996.
- [9] B. A. Camley, Y. Zhao, B. Li, H. Levine, and W.-J. Rappel. Crawling and turning in a minimal reaction-diffusion cell motility model: coupling cell shape and biochemistry. *Phys. Rev. E*, 95:012401, Jan 2017.
- [10] C. E. Chan and D. J. Odde. Traction dynamics of filopodia on compliant substrates. *Science*, 322(5908):1687–1691, 2008.
- [11] C. K. Choi, M. Vicente-Manzanares, J. Zareno, L. A. Whitmore, A. Mogilner, and A. R. Horwitz. Actin and  $\alpha$ -actinin orchestrate the assembly and maturation of nascent adhesions in a myosin II motor-independent manner. *Nature Cell Biology*, 10(9):1039–1050, 2008.
- [12] L. P. Cramer. Forming the cell rear first: breaking cell symmetry to trigger directed cell migration. *Nature Cell Biology*, 12:628 – 632, 2010.
- [13] M. H. Davis. Piecewise-deterministic Markov processes: A general class of non-diffusion stochastic models. *Journal of the Royal Statistical Society. Series B (Methodological)*, 46(3):353–388, 1984.
- [14] M. H. A. Davis. *Markov models and optimization*. Chapman and Hall, 1993.
- [15] S. Deguchi, T. Ohashi, and M. Sato. Tensile properties of single stress fiber isolated from cultured vascular smooth muscle cells. *Journal of Biomechanics*, 39(14):2603–2610, 2006.
- [16] R. B. Dickinson and R. T. Tranquillo. A stochastic model for adhesion-mediated cell random motility and haptotaxis. *Journal of mathematical biology*, 31(6):563–600, 1993.
- [17] P. Dieterich, R. Klages, R. Preuss, and A. Schwab. Anomalous dynamics of cell migration. *Proceedings of the National Academy of Sciences*, 105(2):459–463, 2008.



- [18] C. Engwer, T. Hillen, M. Knappitsch, and C. Surulescu. Glioma follow white matter tracts: a multiscale dti-based model. *Journal of mathematical biology*, 71(3):551–582, 2015.
- [19] T. Erdmann and U. S. Schwarz. Stability of adhesion clusters under constant force. *Physical Review Letters*, 92(4):108102, 2004.
- [20] J. T. Finer, R. M. Simmons, and J. A. Spudich. Single myosin molecule mechanics: piconewton forces and nanometre steps. *Nature*, 368(6467):113–119, 1994.
- [21] P. Friedl, P. Noble, E. Shields, and K. Zänker. Locomotor phenotypes of unstimulated CD45RA<sup>high</sup> and CD45RO<sup>high</sup> CD4<sup>+</sup> and CD8<sup>+</sup> lymphocytes in three-dimensional collagen lattices. *Immunology*, 82(4):617, 1994.
- [22] M. L. Gardel, I. C. Schneider, Y. Aratyn-Schaus, and C. M. Waterman. Mechanical integration of actin and adhesion dynamics in cell migration. *Annual review of cell and developmental biology*, 26:315–333, 2010.
- [23] M. L. Gardel, M. T. Valentine, J. C. Crocker, A. R. Bausch, and D. A. Weitz. Microrheology of entangled F-actin solutions. *Phys. Rev. Lett.*, 91:158302, Oct 2003.
- [24] B. Geiger, J. P. Spatz, and A. D. Bershadsky. Environmental sensing through focal adhesions. *Nature reviews Molecular cell biology*, 10(1):21, 2009.
- [25] D. T. Gillespie. A rigorous derivation of the chemical master equation. *Physica A: Statistical Mechanics and its Applications*, 188(1-3):404–425, 1992.
- [26] A. Groh and A. K. Louis. Stochastic modelling of biased cell migration and collagen matrix modification. *Journal of mathematical biology*, 61(5):617–647, 2010.
- [27] P. Guthardt Torres, I. B. Bischofs, and U. S. Schwarz. Contractile network models for adherent cells. *Physical Review Letters E*, 85:011913, Jan 2012.
- [28] W. R. Holmes and L. Edelstein-Keshet. A comparison of computational models for eukaryotic cell shape and motility. *PLoS computational biology*, 8(12):e1002793, 2012.
- [29] T. Kalwarczyk, N. Zibacz, A. Bielejewska, E. Zaboklicka, K. Koynov, J. Szymaski, A. Wilk, A. Patkowski, J. Gapiski, H.-J. Butt, and R. Hoyst. Comparative analysis of viscosity of complex liquids and cytoplasm of mammalian cells at the nanoscale. *Nano Letters*, 11(5):2157–2163, 2011.
- [30] E. Kassianidou, A. C. Brand, U. S. Schwarz, and S. Kumar. Geometry and network connectivity govern the mechanics of stress fibers. *Proceedings of the National Academy of Sciences*, 114(10):2622–2627, 2017.
- [31] J. Kelkel and C. Surulescu. A multiscale approach to cell migration in tissue networks. *Mathematical Models and Methods in Applied Sciences*, 22(03):1150017, 2012.

- [32] S. Kumar, I. Z. Maxwell, A. Heisterkamp, T. R. Polte, T. P. Lele, M. Salanga, E. Mazur, and D. E. Ingber. Viscoelastic retraction of single living stress fibers and its impact on cell shape, cytoskeletal organization, and extracellular matrix mechanics. *Biophysics Journal*, 90(10):3762–3773, 2006.
- [33] D. A. Lauffenburger and A. F. Horwitz. Cell migration: a physically integrated molecular process. *Cell*, 84(3):359–369, 1996.
- [34] P. A. Lewis and G. S. Shedler. Simulation of nonhomogeneous Poisson processes by thinning. *Naval Research Logistics (NRL)*, 26(3):403–413, 1979.
- [35] F. Li, S. D. Redick, H. P. Erickson, and V. T. Moy. Force measurements of the  $\alpha5\beta1$  integrin–fibronectin interaction. *Biophysical Journal*, 84(2):1252–1262, 2003.
- [36] L. Li, S. F. Nørrelykke, and E. C. Cox. Persistent cell motion in the absence of external signals: A search strategy for eukaryotic cells. *PLOS ONE*, 3(5):1–11, 05 2008.
- [37] Y.-J. Liu, M. Le Berre, F. Lautenschlaeger, P. Maiuri, A. Callan-Jones, M. Heuzé, T. Takaki, R. Voituriez, and M. Piel. Confinement and low adhesion induce fast amoeboid migration of slow mesenchymal cells. *Cell*, 160(4):659–672, 2015.
- [38] A. F. Marée, A. Jilkine, A. Dawes, V. A. Grieneisen, and L. Edelstein-Keshet. Polarization and movement of keratocytes: a multiscale modelling approach. *Bulletin of mathematical biology*, 68(5):1169–1211, 2006.
- [39] A. F. M. Marée, V. A. Grieneisen, and L. Edelstein-Keshet. How cells integrate complex stimuli: The effect of feedback from phosphoinositides and cell shape on cell polarization and motility. *PLOS Computational Biology*, 8(3):1–20, 03 2012.
- [40] A. M. Mastro, M. A. Babich, W. D. Taylor, and A. D. Keith. Diffusion of a small molecule in the cytoplasm of mammalian cells. *Proceedings of the National Academy of Sciences*, 81(11):3414–3418, 1984.
- [41] A. Mogilner. Mathematics of cell motility: have we got its number? *Journal of mathematical biology*, 58(1-2):105, 2009.
- [42] C. Möhl, N. Kirchgessner, C. Schäfer, B. Hoffmann, and R. Merkel. Quantitative mapping of averaged focal adhesion dynamics in migrating cells by shape normalization. *Journal of Cell Science*, 125(1):155–165, 2012.
- [43] J. E. Molloy, J. E. Burns, J. Kendrick-Jones, R. T. Tregear, and D. C. White. Movement and force produced by a single myosin head. *Nature*, 378(6553):209–212, 1995.
- [44] M. Murrell, P. W. Oakes, M. Lenz, and M. L. Gardel. Forcing cells into shape: the mechanics of actomyosin contractility. *Nature Reviews Molecular Cell Biology*, 16(8):486–498, 2015.

- [45] P. W. Oakes, E. Wagner, C. A. Brand, D. Probst, M. Linke, U. S. Schwarz, M. Glotzer, and M. L. Gardel. Optogenetic control of RhoA reveals zyxin-mediated elasticity of stress fibers. *Nature Communications*, 8:15817, 2017.
- [46] D. Oelz and C. Schmeiser. *How do cells move? Mathematical modeling of cytoskeleton dynamics and cell migration. Cell mechanics: from single scale-based models to multiscale modeling*. Chapman and Hall, 2010.
- [47] H. G. Othmer, S. R. Dunbar, and W. Alt. Models of dispersal in biological systems. *Journal of mathematical biology*, 26(3):263–298, 1988.
- [48] K. Paňková, D. Rösel, M. Novotný, and J. Brábek. The molecular mechanisms of transition between mesenchymal and amoeboid invasiveness in tumor cells. *Cellular and molecular life sciences*, 67(1):63–71, 2010.
- [49] R. Paul, P. Heil, J. P. Spatz, and U. S. Schwarz. Propagation of mechanical stress through the actin cytoskeleton toward focal adhesions: model and experiment. *Biophysical Journal*, 94(4):1470–1482, 2008.
- [50] R. J. Petrie, A. D. Doyle, and K. M. Yamada. Random versus directionally persistent cell migration. *Nature Reviews. Molecular Cell Biology*, 10(8):538–549, 2009.
- [51] M. Raftopoulou and A. Hall. Cell migration: Rho GTPases lead the way. *Developmental Biology*, 265(1):23–32, 2004.
- [52] G. R. Ramirez-San Juan, P. W. Oakes, and M. L. Gardel. Contact guidance requires spatial control of leading-edge protrusion. *Molecular Biology of the Cell*, 28(8):1043–1053, 2017.
- [53] A. J. Ridley. Rho GTPases and cell migration. *Journal of Cell Science*, 114(15):2713–2722, 2001.
- [54] A. J. Ridley, M. A. Schwartz, K. Burridge, R. A. Firtel, M. H. Ginsberg, G. Borisy, J. T. Parsons, and A. R. Horwitz. Cell migration: integrating signals from front to back. *Science*, 302(5651):1704–1709, 2003.
- [55] A. Roycroft and R. Mayor. Molecular basis of contact inhibition of locomotion. *Cellular and Molecular Life Sciences*, 73(6):1119–1130, 2016.
- [56] B. Rubinstein, K. Jacobson, and A. Mogilner. Multiscale two-dimensional modeling of a motile simple-shaped cell. *Multiscale Modeling & Simulation*, 3(2):413–439, 2005.
- [57] R. J. Russell, S.-L. Xia, R. B. Dickinson, and T. P. Lele. Sarcomere mechanics in capillary endothelial cells. *Biophysical journal*, 97(6):1578–1585, 2009.
- [58] D. Shao, W.-J. Rappel, and H. Levine. Computational model for cell morphodynamics. *Physical review letters*, 105(10):108104, 2010.

- [59] A. Shellard, A. Szabó, X. Trepát, and R. Mayor. Supracellular contraction at the rear of neural crest cell groups drives collective chemotaxis. *Science*, 362(6412):339–343, 2018.
- [60] J. R. D. Soiné, C. A. Brand, J. Stricker, P. W. Oakes, M. L. Gardel, and U. S. Schwarz. Model-based traction force microscopy reveals differential tension in cellular actin bundles. *PLoS Computational Biology*, 11(3):e1004076, 2015.
- [61] J. Stricker, Y. Aratyn-Schaus, P. W. Oakes, and M. L. Gardel. Spatiotemporal constraints on the force-dependent growth of focal adhesions. *Biophysical journal*, 100(12):2883–2893, 2011.
- [62] T. M. Svitkina, I. G. Surguchova, A. B. Verkhovsky, V. I. Gelfand, M. Moeremans, and J. De Mey. Direct visualization of bipolar myosin filaments in stress fibers of cultured fibroblasts. *Cell Motility and the Cytoskeleton*, 12(3):150–156, 1989.
- [63] U.S.Schwarz and M.L.Gardel. United we stand integrating the actin cytoskeleton and cellmatrix adhesions in cellular mechanotransduction. *Journal of Cell Science*, 125(13):3051–3060, 2012.
- [64] A. B. Verkhovsky and G. G. Borisy. Non-sarcomeric mode of myosin ii organization in the fibroblast lamellum. *The Journal of Cell Biology*, 123(3):637–652, 1993.
- [65] A. B. Verkhovsky, T. M. Svitkina, and G. G. Borisy. Self-polarization and directional motility of cytoplasm. *Current Biology*, 9(1):11–S1, 1999.
- [66] M. Vicente-Manzanares, J. Zareno, L. Whitmore, C. K. Choi, and A. F. Horwitz. Regulation of protrusion, adhesion dynamics, and polarity by myosins IIA and IIB in migrating cells. *The Journal Of Cell Biology*, 176(5):573–580, 2007.
- [67] M. D. Vose. A linear algorithm for generating random numbers with a given distribution. *IEEE Transactions on Software Engineering*, 17(9):972–975, 1991.
- [68] P. T. Yam, C. A. Wilson, L. Ji, B. Hebert, E. L. Barnhart, N. A. Dye, P. W. Wiseman, G. Danuser, and J. A. Theriot. Actin–myosin network reorganization breaks symmetry at the cell rear to spontaneously initiate polarized cell motility. *J Cell Biol*, 178(7):1207–1221, 2007.
- [69] R. Zaidel-Bar, M. Cohen, L. Addadi, and B. Geiger. Hierarchical assembly of cell–matrix adhesion complexes. *Biochemical Society Transactions*, 32(3):416–420, 2004.
- [70] F. Ziebert and I. S. Aranson. Computational approaches to substrate-based cell motility. *npj Computational Materials*, 2:16019, 2016.