

Co-Toeplitz Quantization: A Simple Case

Stephen Bruce Sontz

Abstract. The author has introduced in a recent paper a new class of operators, called co-Toeplitz operators, with symbols in a co-algebra. This is the categorical dual to Toeplitz operators which have symbols in an algebra. The mapping from a symbol to its co-Toeplitz operator gives a quantization scheme, called co-Toeplitz quantization. A new, quite simple particular case of co-Toeplitz quantization is introduced in this note. Examples are given in order to show some of its properties.

Mathematics Subject Classification (2010). Primary 81S99; Secondary 47B99.

Keywords. Quantization, co-Toeplitz operators, co-algebras.

Contents

1. Introduction	1
2. Manin Quantum Plane	2
3. Divided Power Co-algebra	3
4. Negative Degrees	4
5. Matrix Coalgebra	5
6. Concluding Remarks	6
References	6

1. Introduction

In [2] I have defined co-Toeplitz operators in a dual way in terms of category theory to Toeplitz operators. The structures needed for this definition are a co-algebra \mathcal{C} (see [1]) together with a sesqui-linear form $\langle \cdot, \cdot \rangle$ defined on it. We let $\Delta : \mathcal{C} \rightarrow \mathcal{C} \otimes \mathcal{C}$ denote the co-multiplication of \mathcal{C} . Also, we suppose there is another co-algebra \mathcal{P} which injects into \mathcal{C} by a map $j : \mathcal{P} \rightarrow \mathcal{C}$ and that there is a projection $Q : \mathcal{C} \rightarrow \mathcal{P}$, that is, $Qj = id_{\mathcal{P}}$, the identity on \mathcal{P} .

Then we define the *co-Toeplitz operator* $C_g : \mathcal{P} \rightarrow \mathcal{P}$ to be the linear operator defined by the composition

$$\mathcal{P} \xrightarrow{j} \mathcal{C} \xrightarrow{\Delta} \mathcal{C} \otimes \mathcal{C} \xrightarrow{Q \otimes id} \mathcal{P} \otimes \mathcal{C} \xrightarrow{\pi_g} \mathcal{P}. \quad (1.1)$$

The linear map $\pi_g : \mathcal{P} \otimes \mathcal{C} \rightarrow \mathcal{P}$, where $g \in \mathcal{C}$ is called the *symbol* of the co-Toeplitz operator C_g , is defined in [2] by

$$\pi_g(\phi \otimes f) := \langle g, f \rangle \phi$$

for $\phi \in \mathcal{P}$ and $f \in \mathcal{C}$. (The map π_g is dual to α_g to be defined below.) The *anti-linear* map $g \mapsto C_g$ is called the *co-Toeplitz quantization* of \mathcal{C} . This in general does not involve measure theory. For more details see [2].

The definition (1.1) is the dual diagram to that for a *Toeplitz operator* which is defined for a symbol $g \in \mathcal{A}$, an algebra, as the composition (right to left)

$$\mathcal{P} \xleftarrow{P} \mathcal{A} \xleftarrow{\mu} \mathcal{A} \otimes \mathcal{A} \xleftarrow{\iota \otimes id} \mathcal{P} \otimes \mathcal{A} \xleftarrow{\alpha_g} \mathcal{P}. \quad (1.2)$$

Here \mathcal{P} is a sub-algebra of \mathcal{A} with $\iota : \mathcal{P} \rightarrow \mathcal{A}$ being the inclusion map and $P : \mathcal{A} \rightarrow \mathcal{P}$ being a projection. Also, μ is the multiplication map of \mathcal{A} and $\alpha_g(\phi) := \phi \otimes g$ for $\phi \in \mathcal{P}$.

The definition (1.1) has a particular case: $\mathcal{P} = \mathcal{C}$ and $j = Q = id_{\mathcal{C}}$. Then

$$C_g = \pi_g \Delta : \mathcal{C} \rightarrow \mathcal{C}.$$

In this particular simple case, which is the new idea in this note, the only structures needed are a co-algebra and a sesqui-linear form on it. All the examples in this note fall within this case. This simple case does not have an interesting analogue for Toeplitz operators, since diagram (1.2) reduces to the right regular representation of g acting on \mathcal{A} if we put $\mathcal{P} = \mathcal{A}$.

If the sesqui-linear form is positive definite, then C_g acts in a pre-Hilbert space and so may be considered as a densely defined operator acting in the Hilbert space completion of \mathcal{C} . Thus we can construct models for quantum physics, including creation and annihilation co-Toeplitz operators. (See [2].)

All objects in this paper are vector spaces over the complex numbers, and all arrows are linear maps, except as noted.

2. Manin Quantum Plane

We define \mathcal{C} to be the algebra generated by two elements a, c with the relation $ac = qca$ for some non-zero $q \in \mathbb{C}$, the complex numbers. This is called the *Manin quantum plane*. The notation follows that used in [2]. We define the co-multiplication Δ to be the algebra morphism determined by

$$\Delta(a) = a \otimes a \quad \text{and} \quad \Delta(c) = c \otimes a.$$

This is well defined on \mathcal{C} , since $\Delta(ac - qca) = 0$ as the reader can verify. We note that \mathcal{C} does not have a co-unit, though this has no great importance for our purposes. Clearly, $\mathcal{B} := \{a^i c^j \mid i, j \in \mathbb{N}\}$, is a Hamel basis of \mathcal{C} , where \mathbb{N} denotes the non-negative integers. Since C_g is anti-linear in the symbol $g \in \mathcal{C}$, it suffices to calculate C_g for the basis elements $a^i c^j$. And since C_g

is linear, it suffices to evaluate it on these basis elements. We proceed to do this. First, we see that

$$\Delta(a^k c^l) = (\Delta(a))^k (\Delta(c))^l = (a \otimes a)^k (c \otimes a)^l = a^k c^l \otimes a^{k+l}.$$

Then

$$C_{a^i c^j}(a^k c^l) = \pi_{a^i c^j} \Delta(a^k c^l) = \pi_{a^i c^j}(a^k c^l \otimes a^{k+l}) = \langle a^i c^j, a^{k+l} \rangle a^k c^l.$$

So $C_{a^i c^j}$ is diagonalized by the basis \mathcal{B} with its eigenvalues determined by the sesqui-linear form, and therefore it is neither a creation nor an annihilation operator. Rather $C_{a^i c^j}$ is what is known as a *preservation operator*.

At this point in the calculation it becomes clear that the definition of the sesqui-linear form enters in a fundamental way, namely, different choices for it give different co-Toeplitz quantizations. One simple choice is to choose it so that the basis \mathcal{B} is orthogonal. So we put

$$\langle a^i c^j, a^k c^l \rangle := \delta_{i,k} \delta_{j,l} w(i, j).$$

Here $\delta_{r,s}$ denotes the Kronecker delta. We also take $w(i, j) > 0$ for all $i, j \in \mathbb{N}$ so that this is a positive definite inner product and \mathcal{C} is a pre-Hilbert space. With this inner product we see that

$$C_{a^i c^j}(a^k c^l) = \langle a^i c^j, a^{k+l} \rangle a^k c^l = \delta_{i,k+l} \delta_{j,0} w(i, j) a^k c^l.$$

So $C_{a^i c^j} = 0$ if $j > 0$. Continuing with the case $j = 0$ we see that

$$C_{a^i}(a^k c^l) = \delta_{i,k+l} w(i, 0) a^k c^l.$$

So C_{a^i} is diagonalized by the basis \mathcal{B} with 0 and $w(i, 0)$ being its eigenvalues. We define the degree of the monomial $a^k c^l$ by $\deg a^k c^l := k + l$. Then C_{a^i} is zero on monomials with degree $\neq i$ and is a non-zero multiple of the identity on the finite dimensional vector space spanned by the monomials of degree i .

Another choice for the sesqui-linear form is

$$\langle a^i c^j, a^k c^l \rangle := \delta_{i-j,k-l} \mu(i, j, k, l),$$

where $\mu : \mathbb{N}^4 \rightarrow (0, \infty)$ is a positive weight function. With this choice of sesqui-linear form we have

$$C_{a^i c^j}(a^k c^l) = \langle a^i c^j, a^{k+l} \rangle a^k c^l = \delta_{i-j,k+l} \mu(i, j, k+l, 0) a^k c^l.$$

Thus the eigenvalues of $C_{a^i c^j}$ are 0 and $\mu(i, j, i-j, 0)$. Moreover, $C_{a^i c^j}$ is zero except on the set of monomials of degree $i-j$. So $i < j$ implies that $C_{a^i c^j} = 0$.

3. Divided Power Co-algebra

This is based on Example 2.4.8 in [1]. We let \mathcal{C} be the vector space with basis $\{x_i \mid i \in \mathbb{N}\}$. The co-multiplication Δ is the linear map determined by

$$\Delta(x_n) := \sum_{i+j=n} x_i \otimes x_j. \quad (3.1)$$

The degree of each basis element is defined by $\deg x_n := n$. We also define a sesqui-linear form by

$$\langle x_i, x_j \rangle := w(i) \delta_{i,j}, \quad (3.2)$$

where $w : \mathbb{N} \rightarrow (0, \infty)$ is a strictly positive weight function. So this is an inner product making \mathcal{C} into a pre-Hilbert space. Again, it suffices to compute the co-Toeplitz operators T_g for g in the basis. So we compute as follows:

$$\begin{aligned} C_{x_k}(x_n) &= \pi_{x_k} \Delta(x_n) = \pi_{x_k} \left(\sum_{i+j=n} x_i \otimes x_j \right) \\ &= \sum_{i+j=n} \langle x_k, x_j \rangle x_i = \sum_{i+j=n} \delta_{k,j} w(k) x_i. \end{aligned}$$

Now if $k > n$ we have $\delta_{k,j} = 0$ for all the terms in the last sum, since $j \leq n$. So $C_{x_k}(x_n) = 0$ if $k > n$. For the opposite case $0 \leq k \leq n$ we have

$$C_{x_k}(x_n) = \sum_{i+j=n} \delta_{k,j} w(k) x_i = w(k) x_{n-k}.$$

If we define $x_i := 0$ and $w(i) := 0$ for all integers $i < 0$, then we can write this result as one formula for all $k, n \in \mathbb{N}$:

$$C_{x_k}(x_n) = w(k) x_{n-k}.$$

So for $k > 0$ we have that C_{x_k} decreases degree by k and so is an annihilation operator. On the other hand C_{x_0} is a preservation operator.

4. Negative Degrees

Here we give a modification of the previous example that includes negative degrees. We let $M \geq 1$ be an integer and define \mathcal{C} to be the complex vector space with basis $\{x_i\}$ for integers $i \in [-M, M]$. So $\dim \mathcal{C} = 2M + 1$.

We define $\deg x_i := i$ for $i \in [-M, M]$. For convenience we also define $x_i := 0$ for all integers i with $|i| > M$. We use the same formulas as in the previous example, but with new interpretations. So, the co-multiplication Δ is defined by (3.1), but now for integers $|n| \leq M$. With our definitions only finitely many terms in the (now) infinite sum (3.1) are non-zero. We also define a sesqui-linear form by (3.2) but now for integers $i, j \in [-M, M]$. Again, for convenience we put $w(i) := 0$ for $|i| > M$. The same calculation as in the previous example gives

$$C_{x_k}(x_n) = w(k) x_{n-k}$$

but now for all integers $k, n \in [-M, M]$. There are three cases:

1. C_{x_k} increases degree by $|k|$ if $k < 0$ and is a creation operator.
2. C_{x_k} decreases degree by k if $k > 0$ and is an annihilation operator.
3. C_{x_k} preserves degree if $k = 0$ and is a preservation operator.

So we get the three types of operators relevant to physics by using the basis elements with positive, negative and zero degrees. We also can define a **-operation* (\equiv *conjugation*) on \mathcal{C} by putting $x_i^* := x_{-i}$. This definition is

motivated by the theory of complex variables. Using this as motivation, for each $i > 0$ we then define the elements x_i to be *holomorphic* and the elements x_i^* to be *anti-holomorphic*. (The element x_0 could be defined as being both holomorphic and anti-holomorphic, if one wished. But I opt not to do that.) Then the anti-holomorphic elements are symbols of creation operators while the holomorphic elements are symbols of annihilation operators.

5. Matrix Coalgebra

This example comes from Example 2.4.1 in [1]. Let \mathcal{C} be the vector space with basis $\{E_{i,j} \mid 1 \leq i, j \leq n\}$, where $n \geq 1$ is an integer. So, $\dim \mathcal{C} = n^2$. Of course, the motivation is that $E_{i,j}$ is analogous to the $n \times n$ matrix with all entries 0, except for row i and column j which has the entry 1. Let the co-multiplication be determined by

$$\Delta(E_{i,j}) := \sum_{k=1}^n E_{i,k} \otimes E_{k,j}.$$

(As a curious parenthetical remark, let us note that this vector space has a natural algebra structure motivated by matrix multiplication. However, this does not combine with this co-multiplication to give us a bi-algebra. See [1].) We define an inner product on \mathcal{C} by making the basis $\{E_{i,j}\}$ orthonormal. Then we calculate

$$\begin{aligned} C_{E_{r,s}}(E_{i,j}) &= \pi_{E_{r,s}} \Delta(E_{i,j}) = \pi_{E_{r,s}} \left(\sum_{k=1}^n E_{i,k} \otimes E_{k,j} \right) \\ &= \sum_{k=1}^n \langle E_{r,s}, E_{k,j} \rangle E_{i,k} = \sum_{k=1}^n \delta_{r,k} \delta_{s,j} E_{i,k} = \delta_{s,j} E_{i,r}. \end{aligned}$$

So, $C_{E_{r,s}}(E_{i,j})$ is either zero or another basis element.

Another sesqui-linear form is given by $\langle E_{i,j}, E_{r,s} \rangle := w(i+s) \delta_{i-j, r-s}$ with a weight function $w : \mathbb{N} \rightarrow (0, \infty)$. Then

$$\begin{aligned} C_{E_{r,s}}(E_{i,j}) &= \pi_{E_{r,s}} \Delta(E_{i,j}) = \pi_{E_{r,s}} \left(\sum_{k=1}^n E_{i,k} \otimes E_{k,j} \right) \\ &= \sum_{k=1}^n \langle E_{r,s}, E_{k,j} \rangle E_{i,k} = \sum_{k=1}^n w(r+j) \delta_{r-s, k-j} E_{i,k} = w(r+j) E_{i, j+r-s}, \end{aligned}$$

where we put $E_{i,j} = 0$ if $j \leq 0$ or $j > n$. If we define $\deg E_{i,j} := i+j$, then we see that $C_{E_{r,s}}$ changes degree by $r-s$. So, $C_{E_{r,s}}$ is a creation operator if $r > s$, it is an annihilation operator if $r < s$ and finally it is a preservation operator if $r = s$.

Using the adjoint operation of matrices as motivation, we also define a $*$ -operation by $E_{i,j}^* := E_{j,i}$. We also say that $E_{i,j}$ is *holomorphic* if $i < j$ ('upper triangular') and is *anti-holomorphic* if $i > j$ ('lower triangular'). As previously, the anti-holomorphic $E_{i,j}$ are the symbols of (degree increasing)

creation operators and, on the other hand, the holomorphic $E_{i,j}$ are the symbols of (degree decreasing) annihilation operators. Also the ‘diagonal’ elements $E_{i,i}$, which are self-adjoint (or real) with respect to the $*$ -operation, are symbols of (degree preserving) preservation operators.

6. Concluding Remarks

The quantization of co-algebras is a new field of research with co-Toeplitz quantization being the first theory that achieves this. It is remarkable that any sesqui-linear form defined on a co-algebra \mathcal{C} is sufficient extra structure to give us a co-Toeplitz quantization of \mathcal{C} . It is noteworthy that in some of these examples a $*$ -operation can be defined thereby giving holomorphic and anti-holomorphic elements, which are symbols whose co-Toeplitz operators are annihilation and creation operators, respectively.

Acknowledgement

I thank the organizers of the XXXVII Workshop on Geometric Methods in Physics held in July, 2018 in Białowieża, Poland for the opportunity to present a talk based on my paper [2] and for inviting me to write this note which furthers that continuing research program.

References

- [1] M. Hazewinkel et al., *Algebras, Rings and Modules*, Math. Surv. Mono., Vol. 168, Am. Math. Soc., 2010.
- [2] S.B. Sontz, Co-Toeplitz Operators and their Associated Quantization, eprint: arXiv:1708.01979v3

Stephen Bruce Sontz
Centro de Investigación en Matemáticas, A.C. (CIMAT)
Jalisco s/n
Valenciana
Guanajuato 36023
Mexico
e-mail: `sontz@cimat.mx`