

Massive Scattering Amplitudes in Six Dimensions

Rishabh JHA^{*}, Chethan KRISHNAN^{a†}, K. V. Pavan KUMAR^{a‡}

^a Center for High Energy Physics,
Indian Institute of Science, Bangalore 560012, India

Abstract

We show that a natural spinor-helicity formalism that can describe massive scattering amplitudes exists in $D = 6$ dimensions. This is arranged by having helicity spinors carry an index in the Dirac spinor $\mathbf{4}$ of the massive little group, $SO(5) \sim Sp(4)$. In the high energy limit, two separate kinds of massless helicity spinors emerge as required for consistency with arXiv:0902.0981, with indices in the two $SU(2)$'s of the massless little group $SO(4)$. The tensors of $\mathbf{4}$ lead to particles with arbitrary spin, and using these and demanding consistent factorization, we can fix 3- and 4-point tree amplitudes of arbitrary masses and spins: we provide examples. We discuss the high energy limit of scattering amplitudes and the Higgs mechanism in this language, and make some preliminary comments about massive BCFW recursion.

^{*}rishabh.jha77@gmail.com

[†]chethan.krishnan@gmail.com

[‡]kumar.pavan56@gmail.com

Contents

1	Introduction	3
2	Spinor Helicity in Six Dimensions	4
2.1	Spinor Helicity for Massless Particles in $6D$	4
2.2	Spinor Helicity for Massive Particles in $6D$	5
2.3	The High-Energy Limit	6
3	Three Point Scattering Amplitudes	7
3.1	2–Massless, 1–Massive Particles	7
3.2	1–Massless, 2–Massive Particles	9
3.2.1	Unequal Mass	9
3.2.2	Equal Mass	10
3.3	3–Massive Particles	12
4	Four Point Scattering Amplitudes	12
5	Higgs Mechanism	16
6	Massive Recursion Relations	20
A	Four Dimensions	23
A.1	Results in the Convention $x^\mu = (t, -\vec{x})$ as in arXiv:1709.04891	24
A.1.1	Massless Case	24
A.1.2	Massive Case	25
A.2	Results in the Standard Convention $x^\mu = (t, +\vec{x})$	26
A.2.1	Massless Case	27
A.2.2	Massive Case	27
A.3	Obtaining Massive Spinors via Lorentz Boost	28
B	Six Dimensions	30
B.1	Conventions	30
B.2	Angle/Square Brackets and Lorentz Invariant Objects in $6D$	31
B.3	$p^{AB} = \lambda^{AI}\lambda_I^B$	32

B.4	$p_{AB} = \tilde{\lambda}_{AI}\tilde{\lambda}_B^I$	34
B.5	Massive Dirac Equation	35
C	Summary of Important Conventions/Definitions in $6D$	35

1 Introduction

The philosophy that one can fix a theory based largely on consistency conditions (like unitarity, locality, Lorentz invariance) is an old one and its incarnation in S-matrix bootstrap is sometimes considered to be the roots from which string theory arose. A closely related idea that has lead to substantial progress in the last decade is to use on-shell methods (without direct reliance on local quantum fields) to compute scattering amplitudes. See [1] for a review. The foundational observation here is that the basic objects in a quantum field theory can be viewed as particles and not fields, and the former transform in representations of the little group. Quantum fields (which are introduced as a tool for ensuring manifest locality) on the other hand transform as tensors of the Lorentz group, and it has become increasingly plausible in recent years that working with little group covariant objects (spinor helicity variables) might be a more natural and simpler way to construct scattering amplitudes, since they deal directly with particles.

Typically, most work on scattering amplitudes is in the context of massless particles. This is partly because quantum field theories are typically more complicated in the context of massless particles (they often lead to gauge theories), and therefore the simplifications of the spinor helicity formalism are most apparent for massless particles. However, if our goals are truly ambitious, and we are trying to *derive* a UV completion like string theory from our general expectations about scattering amplitudes, then it is evident that we are likely to require the ability to deal with massive particle as well. In such a set up, scattering amplitudes of massive particles should be understandable in terms of IR deformations of massless scattering amplitudes. This type of a formalism for dealing with massive scattering amplitudes has indeed been developed recently in four dimensions by [2] via an extension of the usual massless helicity spinors. In this paper, we will generalize this approach to six dimensions.

Even though the general philosophy of using little group covariant quantities to describe scattering applies across dimensions, it has turned out that its usefulness is immediate only in certain dimensions: $D = 3, 4, 6$, see [3]. This is because to take full advantage of helicity covariance one needs to work with variables that are manifestly on-shell, and explicitly solving on-shell constraints in a useful way has only been possible in these dimensions¹.

¹These specific dimensions have connections to the possible division algebras.

Among these, the massive case for four has already been done as we noted, three is somewhat trivial (though possibly still quite interesting for various purposes), and therefore we turn to six to see whether we can construct a massive spinor helicity formalism. We find that we can.

2 Spinor Helicity in Six Dimensions

Before proceeding further, we note that any particle can be described by specifying its transformation properties under the little group. In six dimensions, the Lorentz group is $SO(5, 1) \sim SU(4)$ and the little group is $SO(4) \sim SU(2) \times SU(2)$ for massless particles and $SO(5) \sim Sp(4)$ for massive particles. We denote the $SU(4)$ indices using A, B, \dots and the $Sp(4)$ indices are denoted by I, J, \dots whereas the undotted (a, b, \dots) and dotted indices (\dot{a}, \dot{b}, \dots) correspond to the two $SU(2)$ groups.

2.1 Spinor Helicity for Massless Particles in 6D

We begin by reviewing the spinor helicity formalism for massless particles following [4]. We start by decomposing the momentum as follows:

$$p_{AB} \equiv p_\mu \Sigma_{AB}^\mu = \tilde{\lambda}_{A\dot{a}} \tilde{\lambda}_{B\dot{b}} \epsilon^{\dot{a}\dot{b}} \quad (2.1)$$

$$p^{AB} \equiv p_\mu \bar{\Sigma}^{\mu AB} = \lambda^{Aa} \lambda^{Bb} \epsilon_{ab} \quad (2.2)$$

The subscript and superscript Lorentz indices $\{A, B\}$ correspond to fundamental and anti-fundamental representations of $SU(4)$. Σ and $\bar{\Sigma}$ correspond to gamma matrices in six dimensions and their explicit forms are given in the appendix. ϵ_{ab} are used to raise and lower the little group ($SU(2) \times SU(2)$ for massless particles) indices and we work with a convention where $\epsilon_{21} = \epsilon^{12} = 1$ for both the $SU(2)$ groups. Note that p_{AB} and p^{AB} are 4×4 matrices of rank two. Also, p_{AB} and p^{AB} are related as follows:

$$p^{AB} = \frac{1}{2} \epsilon^{ABCD} p_{CD} \quad (2.3)$$

Further, we demand that λ^{Aa} and $\tilde{\lambda}_{A\dot{a}}$ satisfy the Dirac equation i.e., we need that λ^{Aa} and $\tilde{\lambda}_{A\dot{a}}$ solve the following equations:

$$p_{AB} \lambda^{Ba} = 0; \quad p^{AB} \tilde{\lambda}_{A\dot{a}} = 0 \quad (2.4)$$

From the explicit forms of p_{AB} and p^{AB} , we can see that λ^{Aa} and $\tilde{\lambda}_{A\dot{a}}$ are solutions the above equations only if the following relation holds:

$$\lambda^{Aa} \tilde{\lambda}_{A\dot{b}} = 0 \quad (2.5)$$

The explicit forms of λ and $\tilde{\lambda}$ are given in the appendix.

Similar to the case of $4D$, we can also introduce the notation of angle and square brackets here. We use the following notation:

$$\lambda^{Aa} = |p^a\rangle^A; \quad \tilde{\lambda}_{A\dot{a}} = |p_{\dot{a}}]_A \quad (2.6)$$

Equations (2.1) and (2.2) can be written in this notation as follows:

$$p_{AB} = |p_{\dot{a}}]_A |p_{\dot{b}}]_B \epsilon^{\dot{a}\dot{b}}; \quad p^{AB} = |p^a\rangle^A |p^b\rangle^B \epsilon_{ab} \quad (2.7)$$

Before going on to the massive case, let us see how the degrees of freedom counting works in (2.1). The momentum has five independent² parameters. $\tilde{\lambda}_{A\dot{a}}$ has eight independent components initially. But note that the RHS is invariant under one of the $SU(2)$'s of the little group. Since $SU(2)$ has three free complex parameters, we can use them to fix three of the components of $\tilde{\lambda}_{A\dot{a}}$ leaving us with a total of five independent components. A similar counting works for the λ^{Aa} in (2.2).

2.2 Spinor Helicity for Massive Particles in $6D$

Now, we turn to the massive case i.e., we consider particles whose momenta satisfy³ $p_\mu p^\mu = m^2$ where m is the mass of the particle. Here, we decompose the momentum as follows:

$$p_{AB} = \tilde{\lambda}_{AI} \tilde{\lambda}_{BJ} J^{IJ} \equiv |p_I]_A |p_J]_B J^{IJ} \quad (2.8)$$

$$p^{AB} = \lambda^{AI} \lambda^{BJ} J_{IJ} \equiv |p^I\rangle^A |p^J\rangle^B J_{IJ} \quad (2.9)$$

where $\{I, J\}$ are $Sp(4)$ fundamental/ anti-fundamental representation indices and J is an invariant matrix of $Sp(4)$ and its explicit form is given in the appendix. Using the matrix J , we can raise and lower the $Sp(4)$ indices. Also, starting from the following property of symplectic matrices:

$$M^T J M = J \quad (2.10)$$

where $M \in Sp(4)$, we can show that p_{AB} and p^{AB} are invariant under $Sp(4)$ transformations of λ and $\tilde{\lambda}$. Further, demanding that λ and $\tilde{\lambda}$ satisfy the Dirac equation⁴ gives us the following condition:

$$\lambda^{AI} \tilde{\lambda}_A^J = -m J^{IJ} \quad (2.11)$$

²Our counting is for complex momenta.

³Note that we are working with mostly negative signature.

⁴The Dirac equation in the massive case is given as follows: $p_{AB} \lambda^{AI} = m \tilde{\lambda}_B^I$, $p^{AB} \tilde{\lambda}_A^I = m \lambda^{BI}$

Let us now explain how the counting works in the massive case. Consider equation (2.8). The momentum has six independent parameters where mass is one of them. $\tilde{\lambda}$ is a 4×4 matrix and hence there are 16 independent parameters. But, as explained above, p_{AB} is invariant under $Sp(4)$. Hence, we can use ten free parameters of $Sp(4)$ to fix ten components of $\tilde{\lambda}$ leaving us with six degrees of freedom as desired. A similar counting works for λ as well.

2.3 The High-Energy Limit

We conclude this section by commenting on how to connect our formalism for massive particles with that of the spinor helicity formalism of massless particles. Note that this will also be useful later in the paper where we take the high-energy limit of various massive scattering amplitudes and match it with the results of massless amplitudes presented in [4].

We start by writing down⁵ the massive helicity spinors in $6D$ in terms of their massless counterparts as follows:

$$\lambda^{AI} = \sqrt{\frac{E+p}{2p}} \lambda_a^A e^{+Ia} + \sqrt{\frac{E-p}{2p}} \eta^{A\dot{a}} e_{\dot{a}}^{-I} \quad (2.12)$$

$$\tilde{\lambda}_A^I = \sqrt{\frac{E+p}{2p}} \tilde{\lambda}_A^{\dot{a}} e_{\dot{a}}^{-I} + \sqrt{\frac{E-p}{2p}} \tilde{\eta}_{Aa} e^{+Ia} \quad (2.13)$$

where e 's are the basis vectors and their explicit forms are given in the appendix. p is the magnitude of the spatial part of the momentum and E is the energy of the particle i.e., $p^0 = E$. $\{\lambda_a, \tilde{\lambda}^{\dot{a}}\}$ and $\{\eta^{\dot{a}}, \tilde{\eta}_a\}$ are massless helicity spinors that satisfy the massless Dirac equation (2.4) where we compute p_{AB}/p^{AB} with $E = +p$ and $E = -p$ respectively. These spinors satisfy the relations $p^{AB} = \lambda^A \lambda_B^B$ and $p_{AB} = \tilde{\lambda}_{A\dot{a}} \tilde{\lambda}_{\dot{b}}^B$. Furthermore the above expansions are useful in obtaining explicit forms of λ^I and $\tilde{\lambda}^J$ because it is easier to solve for the massless helicity spinors using (2.4) as they are decoupled equations as opposed to massive Dirac equation (B.20) which couples λ^I and $\tilde{\lambda}^J$. We use this strategy to find the explicit forms of λ^I and $\tilde{\lambda}^J$ in the appendix.

The high-energy limit is taken in the later sections as discussed henceforth. Starting from the massive scattering amplitudes, we use the above expansions for λ and $\tilde{\lambda}$ to express the massive amplitudes in terms of massless amplitudes as follows:

$$\mathcal{M}^{I_1 \dots I_N} = \sum_i (e^{+i} e^{-N-i})_{\dot{a}_{i+1} \dots \dot{a}_N}^{I_1 \dots I_N; a_1 \dots a_i} \mathcal{M}_{a_1 \dots a_i}^{\dot{a}_{i+1} \dots \dot{a}_N} \quad (2.14)$$

⁵Algebraically, we are expanding a 4×4 matrix in the basis of 4×2 matrices.

Then the high-energy limit is taken for each term separately i.e.,

$$\text{HE limit of } e^{+i}e^{-N-i} \text{ component} = \lim_{m \rightarrow 0} \mathcal{M}_{a_1 \dots a_i}^{\dot{a}_i+1 \dots \dot{a}_N} \quad (2.15)$$

An example on how to take the high-energy limit is given in the section after the discussion of three point and four point functions.

3 Three Point Scattering Amplitudes

In this section, we present a strategy to construct all the possible three point amplitudes in six dimensions where at least one of the particles is massive.

To construct these amplitudes, we use the fact that the scattering amplitude is Lorentz invariant and little group covariant with respect to each of the particles. That is,

$$\mathcal{M}(\{|1\rangle, |1\rangle\}, \dots \{S|i\rangle, S|i\rangle\} \dots) = S\mathcal{M}(\{|1\rangle, |1\rangle\}, \dots \{|i\rangle, |i\rangle\} \dots) \quad (3.1)$$

where S is an arbitrary little group transformation i.e., $S \in Sp(4)$ for massive particles and $S \in SU(2) \times SU(2)$ for massless particles.

3.1 2–Massless, 1–Massive Particles

Let us start the construction of three point amplitudes where only one of the particles is massive. We take the particles 1 and 2 to be massless and the third particle to be massive with mass m_3 . Here we consider particles⁶ 1 and 2 to have one dotted and one undotted index each. We denote them as $1^{a\dot{a}}$ and $2^{b\dot{b}}$ respectively. The massive particle has N $Sp(4)$ indices and transforms in some representation R of the little group $Sp(4)$ and we denote it as $3^{\{I_1, I_2, \dots, I_N\}R}$. Since the scattering amplitude is covariant under the little group and invariant under the Lorentz group, we can write it as follows:

$$\mathcal{M}_3^{ab\dot{a}\dot{b}\{I_1 I_2 \dots I_N\}R} = \sum_{\alpha} g_{\alpha} \lambda_1^{A_1 a} \lambda_2^{B_1 b} \tilde{\lambda}_{1A_2}^{\dot{a}} \tilde{\lambda}_{2B_2}^{\dot{b}} \tilde{\lambda}_{3C_1}^{I_1} \tilde{\lambda}_{3C_2}^{I_2} \dots \tilde{\lambda}_{3C_N}^{I_N} M_{A_1 B_1}^{A_2 B_2 \{C_1 C_2 \dots C_N\}R} \quad (3.2)$$

where we have used the massive Dirac equation to write the above amplitude only in terms of $\tilde{\lambda}_3$'s. By writing this equation, we have reformulated our problem of constructing scattering amplitudes to constructing the tensor M of the Lorentz group $SU(4)$. Note that the available⁷ $SU(4)$ tensors to construct M are δ_B^A , ϵ_{ABCD} , ϵ^{ABCD} and $\frac{1}{m_3} p_{1,2,3}^{AB}$. Also, we need

⁶Note that the rest of the discussion goes through even when these two particles have more than two little group indices each but to avoid clutter of notation, we work with this choice.

⁷Here, we did not include p_{iAB} in the list of available tensors because of the following relation: $\frac{1}{2}\epsilon^{ABCD}p_{iCD} = p_i^{AB}$.

to construct M such that the resultant scattering amplitude transforms in the representation R of $Sp(4)$ with respect to I_1, \dots, I_N . This can be achieved if we demand that the Lorentz tensor M should transform in the representation R with respect to the indices C_1, \dots, C_N and we have already incorporated this in the above equation. Summarizing, M is sum of all the possible terms that can be constructed using the available tensors of $SU(4)$ that have the right transformation properties. Each of these terms in M can have a different coupling constant. We denote it by g_α where α corresponds to different terms in the sum. Sum over α is taken in (3.2) to indicate sum over various terms in M .

Before giving an example where we write down the scattering amplitude explicitly, let us give a simple counting argument that rules out certain interactions. Consider a general case where there are n number of superscript indices and m number of subscript indices in M . To construct M , suppose that we need to use α number of δ_D^C , β number of ϵ_{ABCD} , γ number of ϵ^{ABCD} and η number of $\frac{1}{m_3} p_i^{AB}$. Then the following relation should hold:

$$n - m = 4\gamma + 2\eta - 4\beta \quad (3.3)$$

Since all the parameters here are non-negative integers, if $(n - m)$ is not even, then we can not construct M . Also, as m and n are related to the number of little group indices, this relation can also be used to see whether a scattering is possible among the given particles. For example, consider the equation (3.2). The values of n and m are $(N + 2)$ and 2 respectively. From the above relation (3.3), it is clear that we can not write down any scattering amplitude if N is odd. For instance, this statement implies that we can not have an interaction with two massless vector particles and a massive half-spin particle.

Next, we give a simple example where we write down the scattering amplitude of two massless fermions and one massive vector particle explicitly. That is, the third particle has two massive little group indices I_1 and I_2 that are in antisymmetric representation⁸ of $Sp(4)$. So, the particles we consider in this example are 1^a , 2^b and $3^{[I_1 I_2]}$. Remembering that we can always convert between λ and $\tilde{\lambda}$ using the massive $6D$ Dirac equation (B.20), we get the following expression:

$$\mathcal{M}_3^{ab[I_1 I_2]} = \sum_\alpha g_\alpha \lambda_1^{Aa} \tilde{\lambda}_{2B}^b \tilde{\lambda}_{3C}^{I_1} \tilde{\lambda}_{3D}^{I_2} M_A^{B[CD]} \quad (3.4)$$

where the possibilities of M are:

$$M_A^{B[CD]} = \frac{1}{m_3} \left(\delta_{AP_3}^C p_3^{DB} - \delta_{AP_3}^D p_3^{CB} - \frac{1}{2} \delta_{AP_3}^B p_3^{CD} \right); \quad \frac{1}{m_3} \delta_A^B \left(p_1^{CD} - \frac{1}{2} p_3^{CD} \right) \quad (3.5)$$

The last term in each of the above expressions is required to remove the trace term in the resulting amplitude. So, the three point scattering amplitude for these particles can be

⁸Note that from the usual antisymmetric combination, the trace part has to be subtracted to obtain the antisymmetric representation of $Sp(4)$. See [5] for example.

written as:

$$\mathcal{M}_3^{ab[I_1 I_2]} = g_1 \left(\langle 1^a 3^{I_1} | \langle 3^{I_2} 2^b | - (I_1 \leftrightarrow I_2) - \frac{m_3}{2} J^{I_1 I_2} \langle 1^a 2^b | \right) + \frac{g_2}{m_3} \langle 1^a 2^b | \left([3^{I_1} | 1 | 3^{I_2}] - \frac{m_3^2}{2} J^{I_1 I_2} \right) \quad (3.6)$$

From this expression, it is easy to see that $J_{I_1 I_2} \mathcal{M}_3^{ab[I_1 I_2]} = 0$ as desired.

3.2 1-Massless, 2-Massive Particles

Next, we consider the case where the particles 1 and 2 are massive and particle 3 is massless. Here, there are two cases. One where $m_1 \neq m_2$ and $m_1 = m_2$. The latter case is subtle and is somewhat qualitatively different from the former. We start with the unequal masses case.

3.2.1 Unequal Mass

Let particle 3 be massless and has one dotted and one undotted index i.e., it is denoted⁹ as $3^{a\dot{a}}$. We assume that the particles 1 and 2 transform in representations R_1 and R_2 respectively of the massive little group $Sp(4)$ and we denote these particles as $1^{\{I_1, I_2, \dots, I_{N_1}\}_{R_1}}$ and $2^{\{J_1, J_2, \dots, J_{N_2}\}_{R_2}}$. Following similar steps as in the previous subsection, the scattering amplitude can be written as follows:

$$\mathcal{M}_3^{\{I_1, \dots, I_{N_1}\}_{R_1} \{J_1, \dots, J_{N_2}\}_{R_2} a\dot{a}} = \sum_{\alpha} g_{\alpha} \tilde{\lambda}_{1A_1}^{I_1} \dots \tilde{\lambda}_{1A_{N_1}}^{I_{N_1}} \lambda_2^{B_1 J_1} \dots \lambda_2^{B_{N_2} J_{N_2}} \lambda_3^{C_1 a} \tilde{\lambda}_{3C_2}^{\dot{a}} M_{C_1 \{B_1 \dots B_{N_2}\}_{R_2}}^{C_2 \{A_1 \dots A_{N_1}\}_{R_1}} \quad (3.7)$$

The list of available $SU(4)$ tensors to construct M in this case are δ_B^A , ϵ_{ABCD} , ϵ^{ABCD} , $\frac{1}{g(m)} p_i^{AB}$ where $g(m)$ is an order one polynomial in $m_{1,2}$ and is required for dimensional reasons.

Consider the following example where massive particles 1 and 2 are both spin $\frac{1}{2}$ and particle 3 is spin 1, then the scattering amplitude is given by:

$$\mathcal{M}_3^{I;J;a\dot{a}} = \sum_{\alpha} g_{\alpha} \tilde{\lambda}_{1A}^I \lambda_2^{BJ} \lambda_3^{C_1 a} \tilde{\lambda}_{3C_2}^{\dot{a}} M_{BC_1}^{AC_2} \quad (3.8)$$

where there are two possibilities for M , namely $M = \delta_{C_1}^A \delta_B^{C_2}$ and $M = \frac{1}{r_1(m)r_2(m)} p_1^{AC_2} p_2^{BC_1}$ where both $r_1(m)$ and $r_2(m)$ have mass dimension 1. Hence, the amplitude can be written as:

$$\mathcal{M}_3^{I;J;a\dot{a}} = g_1 \langle 3^a 1^I | \langle 2^J 3^{\dot{a}} | + \frac{g_2}{f_2(m)} \langle 3^a 2^J | \langle 1^I 3^{\dot{a}} | \quad (3.9)$$

where $f_2(m)$ is a function of $m_{1,2}$ with zero mass dimension.

⁹As before, our method works for any number of indices and to avoid clutter of notation, we restrict to just two indices.

3.2.2 Equal Mass

In the equal mass case where $m_1 = m_2 = m$, from the momentum conservation, we see that $p_3 \cdot p_{1,2} = 0$. From this, we can show¹⁰ that $p_{2AB} \lambda_3^{Aa} \lambda_{3a}^B = \langle 3|p_2|3 \rangle = 0$ and that $p_2^{AB} \tilde{\lambda}_{3A\dot{a}} \tilde{\lambda}_{3B}^{\dot{a}} = [3|p_2|3] = 0$. Since particle 3 is massless, we also have $\langle 3^a|3^b \rangle = 0$. These two equations imply that $p_2|3 \rangle \propto |3 \rangle$. The proportionality constant contains an additional information that is used to modify the vertex factors in amplitude calculations similar in line with the massive 4D case as in [2]. This constant of proportionality is defined through either of the following equations:

$$p_{2AB} \lambda_3^{Aa} = m x^{a\dot{a}} \tilde{\lambda}_{3B\dot{a}} \Rightarrow x^{a\dot{a}} = \frac{\langle \zeta^b | 2 | 3^a \rangle}{m \langle \zeta^b 3_{\dot{a}} \rangle} \quad (3.10)$$

$$p_2^{AB} \tilde{\lambda}_{3A\dot{a}} = m y_{\dot{a}a} \lambda_3^{Ba} \Rightarrow y_{\dot{a}a} = \frac{[\zeta^{\dot{b}} | 2 | 3_{\dot{a}}]}{[\zeta^{\dot{b}} 3^a]} \quad (3.11)$$

where ζ is a reference spinor. Note that x, y are dimensionless but transform nontrivially under the massless little group $SU(2) \times SU(2)$. From the above two equations, we can show that $x^{a\dot{a}} y_{\dot{a}b} = \delta_b^a$. Also, we can show that $x^{a\dot{a}} = -y^{a\dot{a}}$ using the following identities¹¹:

$$\begin{aligned} p_{AB} q^{BC} &= -\frac{1}{2} (\delta_{AP}^C q^{GH} - \delta_{AP}^G q^{CH} + \delta_A^H q^{CG}) q_{GH} \\ \langle p_a q^{\dot{b}} \rangle^{-1} &= -\frac{\langle p^a q_{\dot{b}} \rangle}{2p \cdot q}; \quad \langle p^a | q | p^b \rangle = -2\epsilon^{ab}(p \cdot q) \end{aligned} \quad (3.12)$$

Hence, we use only $x^{a\dot{a}}$ in the further calculations.

With the introduction of this x -factor, the three point amplitudes are different as compared to previous cases in the sense that the scattering amplitudes are now written as polynomials in $\lambda, \tilde{\lambda}$ and x . Let us consider the first example where particles 1 and 2 are massive scalars with equal mass m and particle 3 is a massless vector with one dotted and one undotted index. Then the most general polynomial with proper transformation properties constructed out of $x^{a\dot{a}}, x_{\dot{a}a}, \lambda_3^{Aa}$ and $\tilde{\lambda}_{3B\dot{b}}$ is as follows:

$$\begin{aligned} \mathcal{M}_3^{a\dot{a}} &= g_0 m x^{a\dot{a}} + g_1 \lambda_3^{Aa} \tilde{\lambda}_{3B\dot{a}} M_A^B + g_2 x^{a\dot{b}} \tilde{\lambda}_{3Ab} \tilde{\lambda}_{3B}^{\dot{a}} M^{AB} + g_3 x^{b\dot{a}} \lambda_{3b}^A \lambda_3^{Ba} M_{AB} \\ &\quad + g_4 x^{a\dot{b}} x_{\dot{b}c} \lambda_3^{Ac} \tilde{\lambda}_{3B}^{\dot{a}} M_A^B + g_5 x^{a\dot{b}} x^{b\dot{a}} \tilde{\lambda}_{3Ab} \lambda_{3b}^B M_B^A \end{aligned} \quad (3.13)$$

where we have included an extra factor m in the first term for convenience later on. From the available $SU(4)$ tensors ($\epsilon_{ABCD}, \epsilon^{ABCD}, \frac{1}{m} p_2^{AB}, \frac{1}{m} p_{1AB}, \delta_B^A$), we have $M_A^B = \delta_B^A$, $M^{AB} = \frac{1}{m} p_2^{AB}$ and $M_{AB} = \frac{1}{m} p_{1AB}$. Using these and the fact that $x^{a\dot{a}} x_{\dot{a}b} \sim \delta_b^a$, only the first term survives

¹⁰Without loss of generality we could have assumed momentum of particle 1 instead of particle 2.

¹¹The explicit form of the inverse of $\langle p^a q_{\dot{b}} \rangle$ is for example given in [6].

and hence the amplitude of two scalars equal masses and a massless vector is given by:

$$\mathcal{M}_3^{a\dot{a}} = gm x^{a\dot{a}} \quad (3.14)$$

For the next example, let the particle 1 be massive spin-half particle while particles 2 and 3 are massive scalar and massless vector respectively. Then the most general polynomial constructed out of λ_1^{AI} , $x^{a\dot{a}}$, $x_{\dot{a}a}$, λ_3^{Aa} and $\tilde{\lambda}_{3B\dot{b}}$ is as follows:

$$\begin{aligned} \mathcal{M}_3^{Ia\dot{a}} = & g_0 \sqrt{m} x^{a\dot{a}} \lambda_1^{AI} M_A + \frac{1}{\sqrt{m}} (g_1 \lambda_3^{Aa} \tilde{\lambda}_{3B\dot{a}} \lambda_1^{CI} M_{AC}^B + g_2 x^{ab} \tilde{\lambda}_{3A\dot{b}} \tilde{\lambda}_{3B\dot{a}} \lambda_1^{CI} M_C^{AB} \\ & + g_3 x^{b\dot{a}} \lambda_{3b}^A \lambda_3^{Ba} \lambda_1^{CI} M_{ABC} + g_4 x^{ab} x_{\dot{b}c} \lambda_3^{Ac} \tilde{\lambda}_{3B\dot{a}} \lambda_1^{CI} M_{AC}^B + g_5 x^{ab} x^{b\dot{a}} \tilde{\lambda}_{3A\dot{b}} \lambda_{3b}^B \lambda_1^{CI} M_{BC}^A) \end{aligned} \quad (3.15)$$

From the available tensors of $SU(4)$, we can see that none of M_A , m_C^{AB} , M_{ABC} , M_{AB}^C and M_{BC}^A can be constructed. Hence the amplitude corresponding to this process vanishes.

As a third example, let's consider the process involving two massive spin half particles (particle 1 and 2) with equal mass m and a massless vector particle containing one dotted and one undotted index. Then the scattering amplitude constructed using λ_1^{AI} , λ_2^{BJ} , λ_3^{Ca} , $\tilde{\lambda}_{3D}^{\dot{a}}$ and $x^{a\dot{a}}$ is given by:

$$\begin{aligned} \mathcal{M}_3^{IJa\dot{a}} = & \lambda_1^{AI} \lambda_2^{BJ} \{ g_0 m x^{a\dot{a}} M_{AB} + g_1 \lambda_3^{Ca} \tilde{\lambda}_{3D}^{\dot{a}} M_{ABC}^D + g_2 x^{ab} \tilde{\lambda}_{3C\dot{b}} \tilde{\lambda}_{3D}^{\dot{a}} M_{AB}^{CD} \\ & + g_3 x^{b\dot{a}} \lambda_{3b}^C \lambda_3^{Da} M_{ABCD} + g_4 x^{ab} x^{b\dot{a}} \tilde{\lambda}_{3C\dot{b}} \lambda_{3b}^D M_{ABD}^C \} \end{aligned} \quad (3.16)$$

Now by constructing the various M above using the available $SU(4)$ tensors, we get the following result:

$$\begin{aligned} \mathcal{M}_3^{IJa\dot{a}} = & g_0 m^2 x^{a\dot{a}} (\langle 1^I 2^J \rangle + [1^I 2^J]) \\ & + g_1 \left[(\langle 2^J 3^{\dot{a}} \rangle [1^I 3^a] + \langle 1^I 3^{\dot{a}} \rangle [2^J 3^a]) + x^{ab} (\langle 1^I 3^{\dot{a}} \rangle [3_b 2^J] + \langle 1^I 3_b \rangle [3^{\dot{a}} 2^J]) \right. \\ & \left. + y^{\dot{a}b} (\langle 3^a 1^I \rangle [2^J 3_b] + \langle 3_b 1^I \rangle [2^J 3^a]) \right] \\ & + g_2 x^{ab} [1^I 2^J 3_b 3^{\dot{a}}] + g_3 x^{b\dot{a}} \langle 1^I 2^J 3^a 3_b \rangle + g_4 x^{ab} x^{b\dot{a}} (\langle 2^J 3_b \rangle [1^I 3_b] + \langle 1^I 3_b \rangle [2^J 3_b]) \end{aligned} \quad (3.17)$$

where g_i are coupling constants that depend on the theory.

Lastly, we consider the Yukawa amplitude i.e., interaction between two massive fermions with mass m_f and a massless scalar. This amplitude can be written as follows:

$$\mathcal{M}_3^{IJ} = g \lambda_1^{AI} \lambda_2^{BJ} M_{AB} \quad (3.18)$$

Noting that the only possibilities of M_{AB} are p_{iAB} , we get:

$$\mathcal{M}_3^{IJ} = g (\langle 1^I 2^J \rangle + [1^I 2^J]) \quad (3.19)$$

Note that this amplitude does not include any x factors.

3.3 3-Massive Particles

Now, let us consider three massive particles that transform in the representations R_1 , R_2 and R_3 of $Sp(4)$ respectively and we denote them as $1^{\{I_1 \dots I_{N_1}\}_{R_1}}$, $2^{\{J_1 \dots J_{N_2}\}_{R_2}}$ and $3^{\{K_1 \dots K_{N_3}\}_{R_3}}$. Then the scattering amplitude can be written as follows:

$$\mathcal{M}_3^{\{I_1 \dots I_{N_1}\}_{R_1} \{J_1 \dots J_{N_2}\}_{R_2} \{K_1 \dots K_{N_3}\}_{R_3}} = \sum_{\alpha} g_{\alpha} \tilde{\lambda}_{1A_1}^{I_1} \dots \tilde{\lambda}_{1A_{N_1}}^{I_{N_1}} \tilde{\lambda}_{2B_1}^{J_1} \dots \tilde{\lambda}_{2B_{N_2}}^{J_{N_2}} \tilde{\lambda}_{3C_1}^{K_1} \dots \tilde{\lambda}_{3C_{N_3}}^{K_{N_3}} \times M^{\{A_1 \dots A_{N_1}\}_{R_1} \{B_1 \dots B_{N_2}\}_{R_2} \{C_1 \dots C_{N_3}\}_{R_3}} \quad (3.20)$$

As before, sum over α is to indicate that the coupling constant can be different for different terms in M . The available $SU(4)$ tensors to construct M are as follows:

$$\delta_B^A; \epsilon_{ABCD}; \epsilon^{ABCD}; \frac{1}{f_1(m_i)} p_1^{AB}; \frac{1}{f_2(m_i)} p_2^{AB}; \frac{1}{f_3(m_i)} p_3^{AB} \quad (3.21)$$

where f_i 's are functions of mass dimension 1. Even though it is a straightforward exercise, the construction of M can get quite tedious as the number of indices increase.

Let us now give an example where all the three particles transform as a symmetric 2-tensor of $Sp(4)$ i.e., the particles we consider are $1^{(I_1 I_2)}$, $2^{(J_1 J_2)}$ and $3^{(K_1 K_2)}$ where the parentheses denote symmetrization. Also, for simplicity, let us take the masses of all the particles to be same i.e., $m_1 = m_2 = m_3 = m$. The scattering amplitude can be written as follows:

$$\mathcal{M}_3^{(I_1 I_2)(J_1 J_2)(K_1 K_2)} = \sum_{\alpha} g_{\alpha} \tilde{\lambda}_{1A_1}^{I_1} \tilde{\lambda}_{1A_2}^{I_2} \tilde{\lambda}_{2B_1}^{J_1} \tilde{\lambda}_{2B_2}^{J_2} \tilde{\lambda}_{3C_1}^{K_1} \tilde{\lambda}_{3C_2}^{K_2} M^{(A_1 A_2)(B_1 B_2)(C_1 C_2)} \quad (3.22)$$

Some of the possible terms in M are as follows:

$$1. \frac{1}{m^3} p_1^{A_1 B_1} p_2^{A_2 C_1} p_3^{B_2 C_2} + (\text{symmetrizing in } \{A_1, A_2\}, \{B_1, B_2\} \text{ and } \{C_1, C_2\}) \quad (3.23)$$

$$2. \frac{1}{m^3} p_1^{A_1 C_1} p_2^{A_2 B_1} p_3^{B_2 C_2} + (\text{symmetrizing in } \{A_1, A_2\}, \{B_1, B_2\} \text{ and } \{C_1, C_2\}) \quad (3.24)$$

$$3. \frac{1}{m^3} p_1^{B_1 C_1} p_2^{A_1 C_2} p_3^{A_2 B_2} + (\text{symmetrizing in } \{A_1, A_2\}, \{B_1, B_2\} \text{ and } \{C_1, C_2\}) \quad (3.25)$$


4 Four Point Scattering Amplitudes

In order to construct four point tree amplitudes, we start by noting the fact that the four point function has at most simple poles in the Mandelstam variables s, t, u . We construct four point function such that its residues R_s, R_t, R_u in s, t, u channels is equal to product of the left and right three point functions.


Before we start, we set the notation for various channels. Define $s_{ij} := (p_i + p_j)^2$ which in 4-particles processes yield the following:

$$s = s_{12}, t = s_{24}, u = s_{23} \quad (4.1)$$

The first example we consider is the Yukawa interaction. More precisely, we consider the amplitude of two massless scalars and two massive fermions with mass m_f . The possible channels are s and u and are given as follows:



The residue in s -channel is given by the product of the following three point functions:




Using the result of the Yukawa three point amplitudes, we get the residue as

$$\mathcal{R}_s = (g\langle 1^I 5^K \rangle + g'[1^I 5^K]) (g\langle 3^J 5_K \rangle + g'[3^J 5_K]) \quad (4.2)$$

Using $p_1 + p_2 = p_5$ and simplifying, we get:

$$\mathcal{R}_s = m_f[1^I 3^J](g^2 + gg') + m_f[3^J 1^I](g'^2 + gg') + g^2\langle 1^I | 2 | 3^J \rangle + g'^2[1^I | 2 | 3^J] \quad (4.3)$$

Similarly, the residue in the u -channel is given by the following:




Using $p_1 + p_4 = p_5$, we get:

$$\mathcal{R}_u = m_f [1^I 3^J] (g^2 + gg') + m_f [3^J 1^I] (g'^2 + gg') + g^2 \langle 1^I | 4 | 3^J \rangle + g'^2 [1^I | 4 | 3^J] \quad (4.4)$$

Since there are no spurious poles in these residues, the four amplitude is simply given as:

$$\mathcal{M}_4^{IJ} = \frac{\mathcal{R}_s}{s - m_f^2} + \frac{\mathcal{R}_u}{u - m_f^2} \quad (4.5)$$


Next example is $f\bar{f} \rightarrow \bar{f}f$ which again concerns with Yukawa interaction. In the s -channel, Feynman diagram is given as follows:



Scalar propagator is massless here and particles 1, 2, 3, 4 are massive with all having equal mass m . Then the residue in the s -channel is given as follows:

$$\mathcal{R}_s = g^2 ([1^I 2^J] + \langle 1^I 2^J \rangle) (\langle 4^L 3^K \rangle + [4^L 3^K]) \quad (4.6)$$

Similarly in the t -channel, we have the following Feynman diagram:




Residue in this channel is given as follows:

$$\mathcal{R}_t = g^2 ([4^L 2^J] + \langle 4^L 2^J \rangle) (\langle 1^I 3^K \rangle + [1^I 3^K]) \quad (4.7)$$

Since there are no spurious poles appearing, the four point scattering amplitude is given as follows:

$$\mathcal{M}_4^{IJKL} = \frac{\mathcal{R}_s}{s} + \frac{\mathcal{R}_t}{t} \quad (4.8)$$

As a third example, we consider analogue of Compton scattering for scalars in $6D$ i.e., we consider $\phi\gamma \rightarrow \phi\gamma$. This process includes s and u channels and are given as:



Here particles 1 and 3 are massive scalars along with a massive scalar propagator (all of equal mass m) and particles 2 and 4 are massless vectors. Then the product of the residues in the above s -channel is as follows:

$$\mathcal{R}_s = g^2 m^2 x_{12}^{a\dot{a}} x_{34}^{b\dot{b}} \quad (4.9)$$

To simplify this expression, we start by writing down the following equations:

$$p_{IAB} \lambda_2^{Aa} = m x_{12}^{a\dot{a}} \tilde{\lambda}_{2B\dot{a}}; \quad p_I^{CB} \lambda_{4C}^{\dot{b}} = -m x_{34}^{b\dot{b}} \lambda_{4b}^B \quad (4.10)$$

where p_I denotes the momentum of the scalar propagator. Multiplying these two equations, we get:

$$\langle 2^a 4^{\dot{b}} \rangle = x_{12}^{a\dot{a}} x_{34}^{b\dot{b}} \langle 4_b 2_{\dot{a}} \rangle \quad (4.11)$$

Using the following identity:

$$\langle 2^a 4^{\dot{b}} \rangle \langle 2^c 4^{\dot{d}} \rangle^{-1} - \langle 2_c 4^{\dot{b}} \rangle \langle 2_a 4^{\dot{d}} \rangle^{-1} = -\delta_c^a \delta_{\dot{d}}^{\dot{b}} \quad (4.12)$$

we find the residue in s -channel to be:


$$\mathcal{R}_s = g^2 \frac{m^2 \langle 4^b 2^{\dot{a}} \rangle \langle 2^a 4^{\dot{b}} \rangle - \langle 4^b | 1 | 2^a \rangle [4^{\dot{b}} | 3 | 2^{\dot{a}}]}{t} \quad (4.13)$$

Note that the s -channel residue contains a pole in t . Similarly, we can compute the residue along u -channel and the four point amplitude that reproduces these residues is:

$$\mathcal{M}^{a\dot{a};b\dot{b}} = g^2 \frac{m^2 \langle 4^b 2^{\dot{a}} \rangle \langle 2^a 4^{\dot{b}} \rangle - \langle 4^b | 1 | 2^a \rangle [4^{\dot{b}} | 3 | 2^{\dot{a}}]}{(s - m^2)(u - m^2)} \quad (4.14)$$

While computing residues, we need to use the relation $s + t + u = 2m^2$. For instance, we can see that in the s -channel, the pole is at $s = m^2$ and near this pole, we have $m^2 - u = t$. Using this, we recover the correct residue.

As a fourth example, we consider the interaction $\phi\phi^* \rightarrow \phi\phi^*$ which is mediated by a photon. This process has s and u channels and are given as following:



Here particles 1, 2, 3, 4 are scalars with each of them having mass m while the propagator is massless vector. We start with the residue in the s -channel that is given by:

$$\mathcal{R}_s = g^2 m^2 x_{12}^{a\dot{a}} x_{34\dot{a}a} = -g^2 \frac{\langle \zeta^c | 2 | 5^a \rangle [\chi^{\dot{c}} | 3 | 5_{\dot{a}}]}{\langle \zeta^c 5_{\dot{a}} \rangle [\chi^{\dot{c}} 5^a]} \quad (4.15)$$

where ζ and χ are reference spinors and g is a dimensionless constant. To simplify this expression, we need the following identity:

$$p_{AB} q^{CD} = -2(p \cdot q) (\delta_A^C \delta_B^D - \delta_A^D \delta_B^C) + p^{CD} q_{AB} + (\delta_A^C q_{BF} p^{FD} - \delta_A^D q_{BF} p^{FC} - (A \leftrightarrow B)) \quad (4.16)$$

Using this identity along¹² with $\langle 5^a | 2.3 | 5_{\dot{a}} \rangle = 0$, we obtain the residue as:

$$\mathcal{R}_s = 2g^2 (p_2 \cdot p_3) \quad (4.18)$$

Note that contrary to previous examples, there are no poles in t or u even though there are x -factors. Further, if we have started the computation of \mathcal{R}_s by writing $x_{12}^{a\dot{a}} = -\frac{\langle \zeta^c | 1 | 5^a \rangle}{\langle \zeta^c 5_{\dot{a}} \rangle}$, then we get $\mathcal{R}_s = -2g^2 (p_1 \cdot p_3)$. So, more symmetrically, we write the residue as follows:

$$\mathcal{R}_s = g^2 (p_2 - p_1) \cdot p_3 \quad (4.19)$$

The residue in the u -channel is obtained by replacing $1 \leftrightarrow 3$ in \mathcal{R}_s . Thus, the four point amplitude of this process is given by:

$$\mathcal{M}_4 = g^2 \left(\frac{(p_2 - p_1) \cdot p_3}{s} + \frac{(p_2 - p_3) \cdot p_1}{u} \right) \quad (4.20)$$

5 Higgs Mechanism

In this section, we discuss how Higgs mechanism can be understood by demanding that the massive amplitudes reduces to massless amplitudes in the high energy limit. We take the three point amplitude of massive vectors as working example. In the high energy limit, we demand that it should reduce to sum of amplitudes of the following diagrams:

¹²Note that the following identity can be obtained by multiplying the following two equations:

$$p_{2AB} \lambda_5^{Aa} = m x_{12}^{a\dot{a}} \tilde{\lambda}_{5B\dot{a}}; \quad p_3^{CB} \tilde{\lambda}_{5C}^{\dot{a}} = m y_{34\dot{a}b} \lambda_5^{Ba} \quad (4.17)$$




Figure 1: High energy limit of 3–point massive vector amplitude

Note that since a, b, \dots are $SU(2)$ indices, $2^{[ab]}$ and $3^{[cd]}$ are scalar particles with one independent component each.

Before proceeding ahead, we mention that as opposed to $4D$ case, the general three point amplitudes in $6D$ have not been constructed and in particular as far as we know, the amplitude of the second diagram have not been constructed explicitly anywhere. Also, the formalism developed in this paper needs at least one of the particles to be massive and is not applicable to obtain massless amplitudes.

Ideally, our strategy would be to retain those terms in massive 3-vector amplitude which would exactly reproduce the massless amplitudes of both the diagrams on RHS of figure 1 in the high energy limit. But as we do not know the amplitude of second diagram, we can only check the high energy limit of certain components and it is a possibility that to match the second diagram in high energy limit, we might have to retain extra terms in the massive amplitude as compared to (5.4).

We start by constructing the amplitude of three massive vectors that are labelled¹³ as $1^{[I_1 I_2]}$, $2^{[J_1 J_2]}$ and $3^{[K_1 K_2]}$ using the strategy presented in the section 3. The massive scattering amplitude of $1^{[I_1 I_2]}$, $2^{[J_1 J_2]}$ and $3^{[K_1 K_2]}$ is given as follows:

$$\mathcal{M}_3^{[I_1 I_2]; [J_1 J_2]; [K_1 K_2]} = \sum_{\alpha} \frac{g_{\alpha}}{m^2} \lambda_1^{A_1 I_1} \lambda_1^{A_2 I_2} \lambda_2^{B_1 J_1} \lambda_2^{B_2 J_2} \lambda_3^{C_1 K_1} \lambda_3^{C_2 K_2} M_{[A_1 A_2]; [B_1 B_2]; [C_1 C_2]} \quad (5.1)$$

where a factor of m^2 is used to make the coupling constants g_{α} dimensionless¹⁴. Using various tensor structures, we obtain the above amplitude as a function of following Lorentz ($SU(4)$) invariant and the little group ($Sp(4)$) covariant quantities:

$$\begin{aligned} &\langle 1^{I_1} 1^{I_2} \rangle; \langle 2^{J_1} 2^{J_2} \rangle; \langle 3^{K_1} 3^{K_2} \rangle \\ &\langle 1^{I_1} 2^{J_2} \rangle; \langle 2^{J_1} 3^{K_2} \rangle; \langle 3^{K_1} 1^{I_2} \rangle; \langle 1^{I_1} 3^{K_2} \rangle; \langle 2^{J_1} 1^{I_2} \rangle; \langle 3^{K_1} 2^{J_2} \rangle \\ &\langle 1^{I_2} 2^{J_1} \rangle; \langle 2^{J_2} 3^{K_1} \rangle; \langle 3^{K_2} 1^{I_1} \rangle; \langle 1^{I_2} 3^{K_1} \rangle; \langle 2^{J_2} 1^{I_1} \rangle; \langle 3^{K_2} 2^{J_1} \rangle \end{aligned}$$

¹³We are taking the antisymmetric representation of the little group as it has five independent components that match with the degrees of freedom of a massive vector in $6D$.

¹⁴Note that we have used the fact that the dimension of three vector amplitude is one. See [4] for example.

$$\begin{aligned}
& \langle 1^{I_1} 2^{J_1} \rangle; \langle 2^{J_1} 3^{K_1} \rangle; \langle 3^{K_1} 1^{I_1} \rangle; \langle 1^{I_1} 3^{K_1} \rangle; \langle 2^{J_1} 1^{I_1} \rangle; \langle 3^{K_1} 2^{J_1} \rangle \\
& \langle 1^{I_2} 2^{J_2} \rangle; \langle 2^{J_2} 3^{K_2} \rangle; \langle 3^{K_2} 1^{I_2} \rangle; \langle 1^{I_2} 3^{K_2} \rangle; \langle 2^{J_2} 1^{I_2} \rangle; \langle 3^{K_2} 2^{J_2} \rangle \\
& \langle 1^{I_1} | 2 | 1^{I_2} \rangle; \langle 2^{J_1} | 3 | 2^{J_2} \rangle; \langle 3^{K_1} | 1 | 3^{K_2} \rangle; [1^{I_1} | 2 | 1^{I_2}]; [2^{J_1} | 3 | 2^{J_2}]; [3^{K_1} | 1 | 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_1} 2^{J_2} \rangle; \langle 1^{I_1} 1^{I_2} 3^{K_1} 3^{K_2} \rangle; \langle 2^{J_1} 2^{J_2} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_1} 2^{J_2}]; [1^{I_1} 1^{I_2} 3^{K_1} 3^{K_2}]; [2^{J_1} 2^{J_2} 3^{K_1} 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_1} 3^{K_1} \rangle; \langle 1^{I_1} 2^{J_1} 2^{J_2} 3^{K_1} \rangle; \langle 1^{I_1} 2^{J_1} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_1} 3^{K_1}]; [1^{I_1} 2^{J_1} 2^{J_2} 3^{K_1}]; [1^{I_1} 2^{J_1} 3^{K_1} 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_2} 3^{K_1} \rangle; \langle 1^{I_2} 2^{J_1} 2^{J_2} 3^{K_1} \rangle; \langle 1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_2} 3^{K_1}]; [1^{I_2} 2^{J_1} 2^{J_2} 3^{K_1}]; [1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_1} 3^{K_2} \rangle; \langle 1^{I_1} 2^{J_1} 2^{J_2} 3^{K_2} \rangle; \langle 1^{I_1} 2^{J_1} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_1} 3^{K_2}]; [1^{I_1} 2^{J_1} 2^{J_2} 3^{K_2}]; [1^{I_1} 2^{J_1} 3^{K_1} 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_2} 3^{K_2} \rangle; \langle 1^{I_2} 2^{J_1} 2^{J_2} 3^{K_2} \rangle; \langle 1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_2} 3^{K_2}]; [1^{I_2} 2^{J_1} 2^{J_2} 3^{K_2}]; [1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2}] \\
& \langle 1^{I_1} 1^{I_2} 2^{J_2} 3^{K_2} \rangle; \langle 1^{I_2} 2^{J_1} 2^{J_2} 3^{K_2} \rangle; \langle 1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2} \rangle; [1^{I_1} 1^{I_2} 2^{J_2} 3^{K_2}]; [1^{I_2} 2^{J_1} 2^{J_2} 3^{K_2}]; [1^{I_2} 2^{J_1} 3^{K_1} 3^{K_2}]
\end{aligned} \tag{5.2}$$

where we used the following notation:

$$\langle i^I j^J \rangle = \lambda_i^{AI} \tilde{\lambda}_{jA}^J; \langle i^I | p | j^J \rangle = \lambda_i^{AI} \lambda_j^{BJ} p_{AB}; \langle i^I j^J k^K l^L \rangle = \epsilon_{ABCD} \lambda_i^{AI} \lambda_j^{BJ} \lambda_k^{CK} \lambda_l^{DL} \tag{5.3}$$

The other possible structures that we did not mention here can be written down using the listed quantities.

Now, we demand that the above three point amplitude of massive vectors matches with the appropriate amplitudes in the high energy limit. As a result of this constraint, we need to keep only the following terms in the massive amplitude:

$$\begin{aligned}
& \mathcal{M}_3^{[I_1 I_2]; [J_1 J_2]; [K_1 K_2]} \\
& = \frac{1}{m^3} (\langle 3^{[K_1} | p_1 - p_2 | 3^{K_2]} \rangle) (\langle 1^{I_1} 2^{J_1} \rangle \langle 1^{I_2} 2^{J_2} \rangle - \langle 1^{I_1} 2^{J_2} \rangle \langle 1^{I_2} 2^{J_1} \rangle) \\
& + \frac{1}{m^3} (\langle 2^{[J_1} | p_3 - p_1 | 2^{J_2]} \rangle) (\langle 1^{I_1} 3^{K_1} \rangle \langle 1^{I_2} 3^{K_2} \rangle - \langle 1^{I_1} 3^{K_2} \rangle \langle 1^{I_2} 3^{K_1} \rangle) \\
& + \frac{1}{m^3} (\langle 1^{[I_1} | p_2 - p_3 | 1^{I_2]} \rangle) (\langle 2^{J_1} 3^{K_1} \rangle \langle 2^{J_2} 3^{K_2} \rangle - \langle 2^{J_1} 3^{K_2} \rangle \langle 2^{J_2} 3^{K_1} \rangle) \\
& - \text{terms containing } J^{I_1 I_2} \text{ and/or } J^{J_1 J_2} \text{ and/or } J^{K_1 K_2}
\end{aligned} \tag{5.4}$$

The last line is needed to remove the trace terms from the amplitude. We did not explicitly write them down as they vanish while choosing the component $e_a^{+I_1} e_a^{-I_2} e_b^{+J_1} e_b^{-J_2} e_c^{+K_1} e_c^{-K_2}$. As will be explained later, this component is relevant while taking the high-energy limit to match it with the 3-gluon amplitude of [4].

This amplitude can also be obtained from the interaction of the form $(\partial_\mu A_\nu) A^\mu A^\nu$ using Feynman rules as follows:

$$V^{[I_1 I_2]; [J_1 J_2]; [K_1 K_2]} = \left(\varepsilon_1^{[I_1 I_2]} \cdot (p_2 - p_3) \right) (\varepsilon_2^{[J_1 J_2]} \cdot \varepsilon_3^{[K_1 K_2]}) + \text{cyclic in } 1, 2, 3 \text{ particles} \tag{5.5}$$

where the massive polarization vectors are given as follows:

$$\begin{aligned}
(\varepsilon_i^{[IJ]})^{AB} &= \frac{1}{m} \left(\lambda_i^{AI} \lambda_i^{BJ} - \lambda_i^{BI} \lambda_i^{AJ} + \frac{1}{2} p_i^{AB} J^{IJ} \right) \\
(\varepsilon_i^{[IJ]})_{AB} &= \frac{1}{m} \left(\tilde{\lambda}_{iA}^I \tilde{\lambda}_{iB}^J - \tilde{\lambda}_{iB}^I \tilde{\lambda}_{iA}^J - \frac{1}{2} p_{iAB} J^{IJ} \right) \\
(\varepsilon_{[IJ]})^{AB} &= \frac{1}{m} \left(\lambda_I^A \lambda_J^B - \lambda_I^B \lambda_J^A - \frac{1}{2} p^{AB} J_{IJ} \right) \\
(\varepsilon_{[IJ]})_{AB} &= \frac{1}{m} \left(\tilde{\lambda}_{iAI} \tilde{\lambda}_{iBJ} - \tilde{\lambda}_{iBI} \tilde{\lambda}_{iAJ} + \frac{1}{2} p_{iAB} J_{IJ} \right)
\end{aligned} \tag{5.6}$$

The second form can be obtained¹⁵ by using the relation $\varepsilon_{AB} = \frac{1}{2} \epsilon_{ABCD} \varepsilon^{CD}$ and third one by contracting the first equation with $J_{KI} J_{LJ}$. Note that this choice of polarization vectors satisfy the following necessary relations:

$$\begin{aligned}
\varepsilon_{iAB} p_i^{AB} &= \varepsilon_i^{AB} p_{iAB} = 0 \\
(\varepsilon_i^{[IJ]})^{AB} (\varepsilon_i^{[KL]})_{AB} &= 2 (J^{IK} J^{JL} - J^{JL} J^{IK}) - J^{IJ} J^{KL} \\
(\varepsilon_i^{[IJ]})^{AB} (\varepsilon_{i[IJ]})_{CD} &= 2 (\delta_C^A \delta_D^B - \delta_D^A \delta_C^B) + \frac{1}{m^2} p^{AB} p_{CD}
\end{aligned} \tag{5.8}$$

We now show that in the high energy limit this massive amplitude indeed reduces to the appropriate amplitudes as given in (1). First, we consider the high energy limit of the component $e_a^{+I_1} e_{\dot{a}}^{-I_2} e_b^{+J_1} e_{\dot{b}}^{-J_2} e_c^{+K_1} e_{\dot{c}}^{-K_2}$. This should reproduce the three point amplitude of gluons considered in [4]. The corresponding amplitude of the above component is as follows:

$$\mathcal{M}^{a\dot{a};b\dot{b};c\dot{c}} = \frac{1}{p_1 p_2 p_3} \left[\langle 3^c | 1 | \eta_3^{\dot{c}} \rangle \left(\langle 1^a \tilde{\eta}_2^b | \langle \eta_1^{\dot{a}} 2^b \rangle - \langle 1^a 2^b | \langle \eta_1^{\dot{a}} \tilde{\eta}_2^b \rangle \right) + \text{cyclic in } 1,2,3 \text{ particles} \right] \tag{5.9}$$

where $p_i^2 = E_i^2 - m^2$. This expression is same as the three gluon amplitude in equation (52) of [4] if we choose the reference spinors in the polarization vectors of i^{th} particle to be η_i . As the final answer should not depend on the choice of polarization vectors, we see that we have obtained the 3-gluon amplitude by taking the high energy limit of the massive amplitude (5.4). Using KLT relations [7] we accordingly obtain the 3-point amplitude in linearized gravity.

Note that this can also be obtained using the Feynman rules computation of this particular component coming from the three point vertex similar to gluons. More explicitly, this

¹⁵The following identity is useful in deriving this:

$$\epsilon_{ABCD} \lambda^{CI} \lambda^{DJ} = \tilde{\lambda}_A^I \tilde{\lambda}_B^J - \tilde{\lambda}_A^J \tilde{\lambda}_B^I - p_{AB} J^{IJ} \tag{5.7}$$

component of three point vertex can be computed using Feynman rules as follows:

$$V^{a\dot{a};bb;\dot{c}\dot{c}} = (\varepsilon_1^{a\dot{a}} \cdot (p_2 - p_3)) (\varepsilon_2^{b\dot{b}} \cdot \varepsilon_3^{\dot{c}\dot{c}}) + (\varepsilon_2^{b\dot{b}} \cdot (p_3 - p_1)) (\varepsilon_3^{\dot{c}\dot{c}} \cdot \varepsilon_1^{a\dot{a}}) + (\varepsilon_3^{\dot{c}\dot{c}} \cdot (p_1 - p_2)) (\varepsilon_1^{a\dot{a}} \cdot \varepsilon_2^{b\dot{b}}) \quad (5.10)$$

where the corresponding components of massive polarization vectors are as follows:


$$\begin{aligned} (\varepsilon_i^{a\dot{a}})^{AB} &= \frac{1}{p_i} [\lambda_i^{Aa} \eta_i^{B\dot{a}} - \lambda_i^{Ba} \eta_i^{A\dot{a}}] \\ (\varepsilon_i^{a\dot{a}})_{AB} &= \frac{1}{p_i} [\tilde{\eta}_{iB}^a \tilde{\lambda}_{iA}^{\dot{a}} - \tilde{\eta}_{iA}^a \tilde{\lambda}_{iB}^{\dot{a}}] \end{aligned} \quad (5.11)$$

Using these expressions, we can show that $V^{a\dot{a};bb;\dot{c}\dot{c}}$ is same as the terms we retained in (5.9) up to a numerical factor.

6 Massive Recursion Relations

In this section, we discuss BCFW recursion relations [8]. For concreteness, we stick to the discussion of computing four point amplitudes using BCFW. The discussion presented here is preliminary and incomplete due to non-vanishing ‘‘boundary’’ term at infinity in the shift parameter z .

Let the four point function schematically be given as follows:



All the momenta are incoming and we have $p_1 + p_2 + p_3 + p_4 = 0$. The masses of the particles are m_1, m_2, m_3, m_4 . To obtain BCFW recursion relations, we start by shifting two momenta that are on either¹⁶ side of the diagram as follows:

$$\hat{p}_1^\mu(z) = p_1^\mu + zq^\mu, \quad \hat{p}_2^\mu(z) = p_2^\mu - zq^\mu \quad (6.1)$$

where z is a complex number and q^μ is independent of z . Note that shift ensures that the momentum conservation still holds i.e., we have $\hat{p}_1 + \hat{p}_2 + p_3 + p_4 = 0$. Also, we demand that the shifted momenta are on-shell. That is, we need $\hat{p}_1^2 = m_1^2$ and $\hat{p}_2^2 = m_2^2$. This imposes the following constraints on q :

$$p_1 \cdot q = p_2 \cdot q = q \cdot q = 0 \quad (6.2)$$

¹⁶That is, we can shift the momenta of particles $\{1, 2\}$ or $\{1, 4\}$ or $\{2, 3\}$ or $\{3, 4\}$. In the following discussion, we shift the particles 1 and 2.

The four point amplitude that can be computed using the shifted momenta is denoted as $\mathcal{A}_4(z)$. Note that the original four point amplitude is simply $\mathcal{A}_4(z = 0)$. Equivalently, this statement can be written as follows:

$$\mathcal{A}_4(0) = \oint \frac{dz}{z} \mathcal{A}_4(z) \quad (6.3)$$

where the contour is a circle around the origin. Now, we specialize to tree-level diagrams and hence the amplitude $\mathcal{A}_4(z)$ has no branch cuts and can be shown using $q^2 = 0$ that it has only simple poles. These poles are at $\hat{p}_5^2 = m_5^2$ where 5 denotes the propagator. Simplifying this expression shows that the poles are at $z = \frac{m_5^2 - p_5^2}{2p_5 \cdot q}$ where $p_5 = p_1 + p_3$. With this knowledge, we deform the contour so that we can write:

$$\mathcal{A}_4(0) = - \sum_{z_\alpha \neq 0} \mathcal{A}(z_\alpha) + B \quad (6.4)$$

where z_α are all¹⁷ the simple poles and $\mathcal{A}(z_\alpha)$ is the residue at these poles. B is the residue as $z \rightarrow \infty$. Ideally, we want the contribution at infinity to go to zero.

To obtain the BCFW recursion relations, we start by noting that as $\hat{p}_5^2 - m_5^2 = 0$, the propagator goes on-shell and hence the 4-point amplitude separates into a product of three point amplitudes i.e.,

$$\mathcal{A}_4(0) = - \sum_{z_\alpha \neq 0} \mathcal{A}_L(\hat{p}_1(z_\alpha), p_3, \hat{p}_5(z_\alpha)) \frac{1}{p_5^2 - m_5^2} \mathcal{A}_R(-\hat{p}_5(z_\alpha), p_4, \hat{p}_2(z_\alpha)) + B \quad (6.5)$$

Hence, starting from three point amplitudes, we can obtain four point amplitudes and these recursion relations can be obtained for higher point functions following the discussion presented here.

To compute the four point amplitude using the above relation, we generally need the explicit forms of the shifted helicity spinors $\hat{\lambda}_{1,2}; \hat{\tilde{\lambda}}_{1,2}$ corresponding to the shifted momenta $\hat{p}_{1,2}$. These shifted spinors are related to the explicit solutions of q^μ that satisfy the properties (6.2). In 4D, if at least one of the particles 1 and 2 is massless, then q^μ and (hence) $\hat{\lambda}_{1,2}; \hat{\tilde{\lambda}}_{1,2}$ can be determined easily. See [9] for the explicit forms. One of the advantages in 4D is that as long as one of the particles is massless, we can work with the little group $U(1)$ which simplifies various computations related to BCFW recursion relations. So, the authors of [9] consider shifts only in the massless legs while computing various amplitudes using their formalism.

In 6D, as mentioned in the earlier sections, the massless little group is $SO(4) \sim SU(2) \times SU(2)$ and an extra factor of the form $M^{a\dot{a}}$ needs to be included in q^μ to account for the


¹⁷Note that in the present case of 4-point amplitude, there is only one such simple pole.

invariance of momentum under the little group. For instance, one of the choices¹⁸ of q^μ is:

$$q^\mu = \varepsilon_{1a\dot{a}}^\mu M^{a\dot{a}} \quad (6.6)$$

where $\varepsilon_{1a\dot{a}}^\mu$ is the massless polarization vector of the first particle and we need $\det M = 0$ to ensure $q^2 = 0$. Also, following [4], we choose the reference spinor in the polarization vector to be λ_2^{Aa} .

We now apply similar shifts as above to compute the six dimensional analogue of Compton scattering in scalar QED. That is, we consider $\phi\gamma^{a\dot{a}} \rightarrow \phi\gamma^{b\dot{b}}$. The diagrams that contribute to this amplitude are:



We shift the momenta of massless particles as follows:

$$\hat{p}_2^\mu = p_2^\mu + \frac{z}{\sqrt{2}} M^{a\dot{a}} \varepsilon_{2a\dot{a}}^\mu; \quad \hat{p}_4^\mu = p_4^\mu - \frac{z}{\sqrt{2}} M^{a\dot{a}} \varepsilon_{2a\dot{a}}^\mu \quad (6.7)$$

We choose the reference spinor in ε_2 to be the helicity spinors of particle 4. For this choice, the polarization vector is given as follows:

$$\varepsilon_{a\dot{a}}^{AB} = \sqrt{2} \frac{\lambda_{4c}^A \lambda_{2a}^B - \lambda_{4c}^B \lambda_{2a}^A}{\langle 4_c 2_{\dot{a}} \rangle}; \quad \varepsilon_{2AB}^{a\dot{a}} = \sqrt{2} \frac{\tilde{\lambda}_{2A}^{\dot{a}} \tilde{\lambda}_{4B\dot{b}} - \tilde{\lambda}_{2B}^{\dot{a}} \tilde{\lambda}_{4A\dot{b}}}{[4_{\dot{b}} 2_a]} \quad (6.8)$$

From these expressions, we can compute the shifted helicity spinors to be:

$$\hat{\lambda}_2^{Aa} = \lambda_2^{Aa} + z m^a n^b \lambda_{4b}^A; \quad \hat{\lambda}_{2A}^{\dot{a}} = \tilde{\lambda}_{2A}^{\dot{a}} - z \tilde{m}^{\dot{a}} \tilde{n}^{\dot{b}} \tilde{\lambda}_{4Ab} \quad (6.9)$$

$$\hat{\lambda}_4^{Aa} = \lambda_4^{Aa} + z n^a m^b \lambda_{2b}^A; \quad \hat{\lambda}_{4A}^{\dot{a}} = \tilde{\lambda}_{4A}^{\dot{a}} - z \tilde{n}^{\dot{a}} \tilde{m}^{\dot{b}} \tilde{\lambda}_{2Ab} \quad (6.10)$$

where we have used the fact that $\det M = 0$ to write $M^{a\dot{a}} = m^a \tilde{n}^{\dot{a}}$ and we have defined n 's as follows:

$$n^a = \tilde{m}^{\dot{b}} [2_{\dot{b}} 4_a]^{-1}; \quad \tilde{n}^{\dot{a}} = m^b [2^b 4_{\dot{a}}]^{-1} \quad (6.11)$$

Before applying the recursion relations, note that the poles are at $z_s = -\sqrt{2} \frac{p_2 \cdot p_1}{M^{a\dot{a}} \varepsilon_{2a\dot{a}} \cdot p_1}$ in the s -channel and $z_u = -\sqrt{2} \frac{p_3 \cdot p_1}{M^{a\dot{a}} \varepsilon_{2a\dot{a}} \cdot p_3}$ in the u -channel. The s -channel contribution of the amplitude is given by:

$$\mathcal{M}_s = m^2 \hat{x}_{12}^{a\dot{a}} \frac{1}{(p_1 + p_2)^2 - m^2} \hat{x}_{34}^{b\dot{b}} \quad (6.12)$$

¹⁸Note that it can be argued at least for gauge theory amplitudes that the following is one of the choices for which $\mathcal{A}(z) \rightarrow 0$ as $z \rightarrow \infty$. See the discussion in the beginning of section five of [4] for instance.

where the definitions of \hat{x} 's following section 3.2.2 are:

$$m\hat{x}_{12}^{aa} = \frac{\langle \zeta^c | 1 | \hat{2}^a \rangle}{\langle \zeta^c \hat{2}_a \rangle}; \quad m\hat{x}_{34}^{bb} = \frac{[\zeta'^c | 3 | \hat{4}^b]}{\langle \hat{4}_b \zeta'^c \rangle} \quad (6.13)$$

For convenience, we choose $\zeta = 4$ and $\zeta' = 2$. With these choices along with the explicit forms of λ 's, we get:

$$\mathcal{M}_s = \left(\frac{\langle 4^c | 1 | 2^a \rangle}{\langle 4^c 2_a \rangle} + z_s m^a n^d \frac{\langle 4^c | 1 | 4_d \rangle}{\langle 4^c 2_a \rangle} \right) \frac{1}{2p_1 \cdot p_2} \left(\frac{[2^c | 3 | 4^b]}{\langle 4_b 2^c \rangle} - z_s \tilde{n}^b \tilde{m}^d \frac{[2^c | 3 | 2_d]}{\langle 4_b 2^c \rangle} \right) \quad (6.14)$$

where we have used $\hat{p}_5^\mu = \hat{p}_1^\mu + \hat{p}_2^\mu$. Similarly the u -channel contribution can be computed to be as follows:

$$\mathcal{M}_u = \left(\frac{\langle 4^c | 3 | 2^a \rangle}{\langle 4^c 2_a \rangle} + z_u m^a n^d \frac{\langle 4^c | 3 | 4_d \rangle}{\langle 4^c 2_a \rangle} \right) \frac{1}{2p_3 \cdot p_2} \left(\frac{[2^c | 1 | 4^b]}{\langle 4_b 2^c \rangle} - z_u \tilde{n}^b \tilde{m}^d \frac{[2^c | 1 | 2_d]}{\langle 4_b 2^c \rangle} \right) \quad (6.15)$$

We can in principle simplify these expressions further by using the expressions for z_s and z_u . But before doing that let us analyze the behaviour of $\mathcal{A}(z)$ as $z \rightarrow \infty$.

We start by recalling the expression of scalar Compton scattering amplitude (see (4.14)). The corresponding 4-point amplitude is given as follows:

$$\mathcal{M}^{a\dot{a};b\dot{b}} = g^2 \frac{m^2 \langle 4^b 2^{\dot{a}} \rangle \langle 2^a 4^{\dot{b}} \rangle - \langle 4^b | 1 | 2^a \rangle [4^{\dot{b}} | 3 | 2^{\dot{a}}]}{(p_1 \cdot p_2)(p_3 \cdot p_2)} \quad (6.16)$$

As we shift 2 and 4 in this expression, we see that it goes like z^0 as $z \rightarrow \infty$ and hence BCFW is not applicable in this situation. That is, the ‘naive’ shift that was used in the massless case¹⁹ does not work in the massive case even though we restricted our shifts to massless vector particles. There are some works in the literature which discuss on how to proceed with BCFW recursion relations if one has ‘boundary’ term at infinity and we leave such endeavours for future work.

Acknowledgments

We thank Avinash Raju for early collaboration and emphasizing to us that $SO(5) \sim Sp(4)$.

A Four Dimensions

There are typos in [2], so we have solved and reproduced here the results of massive spinor helicity formalism in 4D in different conventions. We have also calculated some of the results from a different perspective using Lorentz boost.

¹⁹Note that those shifts were used to compute gauge theory amplitudes. So, it might be possible that those shifts work only for gauge theories.

A.1 Results in the Convention $x^\mu = (t, -\vec{x})$ as in arXiv:1709.04891

The conventions adopted in [2] are as follows (natural units are employed everywhere):

- $g_{\mu\nu} = (+, -, -, -)$
- $x^\mu = (t, -\vec{x}) \Rightarrow p_\mu = (E, \vec{p})$ ²⁰
- $\epsilon_{ij} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \Rightarrow \epsilon^{ij} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This is the convention adopted for both $\epsilon_{\alpha\dot{\alpha}}$ and ϵ_{IJ} where α and $\dot{\alpha}$ are Lorentz indices while I and J are little group indices, used as raising/lowering operators.
- 4- vector Pauli matrices are $\sigma^\mu = (1, \vec{\sigma})$ and $\bar{\sigma}^\mu = (1, -\vec{\sigma})$ where $\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ and $\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Corresponding to these conventions, we have the momentum bispinor matrix as follows:

$$p_{\alpha\dot{\alpha}} = \sigma \cdot p = (\sigma^\mu)_{\alpha\dot{\alpha}} p_\mu = \begin{pmatrix} E + p \cos(\theta) & p \sin(\theta) e^{-i\phi} \\ p \sin(\theta) e^{i\phi} & E - p \cos(\theta) \end{pmatrix} \quad (\text{A.1})$$

and

$$p^{\dot{\alpha}\alpha} = \bar{\sigma} \cdot p = (\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} p_\mu = \begin{pmatrix} E - p \cos(\theta) & -p \sin(\theta) e^{-i\phi} \\ -p \sin(\theta) e^{i\phi} & E + p \cos(\theta) \end{pmatrix} \quad (\text{A.2})$$

A.1.1 Massless Case

The Lorentz group in this case is $\text{Spin}(3, 1) \cong SL(2, \mathbb{C})$ and the little group is $SO(2) \cong U(1)$. The massless spinors λ_α and $\tilde{\lambda}_{\dot{\alpha}}$ satisfy the following equations:

$$p_{\alpha\dot{\alpha}} = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}} \quad (\text{A.3})$$

The forms of λ and $\tilde{\lambda}$ that satisfy this equation are as follows:

$$\lambda_\alpha = \sqrt{2E} \begin{pmatrix} c \\ s \end{pmatrix}, \quad \tilde{\lambda}_{\dot{\alpha}} = \sqrt{2E} \begin{pmatrix} c \\ s^* \end{pmatrix} \quad (\text{A.4})$$

Here $c := \cos(\frac{\theta}{2})$, $s := \sin(\frac{\theta}{2})e^{i\phi}$ and $s^* := \sin(\frac{\theta}{2})e^{-i\phi}$. They happen to satisfy the following massless Dirac equations (or Weyl equations):

$$p_{\alpha\dot{\alpha}} \lambda^\alpha = 0, \quad p_{\alpha\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}} = 0 \quad (\text{A.5})$$

²⁰For maximal compatibility with the results in [2], we find that we need this convention. Results in the standard convention $x^\mu = (t, \vec{x})$ are dealt in sub-appendix A.2.

A.1.2 Massive Case

In the massive case, the Lorentz group remains the same but the little group changes to $SO(3) \cong SU(2)$. The latter contributes an index to both the spinors λ_α and $\tilde{\lambda}_{\dot{\alpha}}$. The spinors satisfy the following equation:

$$p_{\alpha\dot{\alpha}} = \lambda_\alpha^I \tilde{\lambda}_{\dot{\alpha}I} \quad (\text{A.6})$$

This is a little group invariant quantity as little group index is summed over. This has to be true by the very definition of the little group. The forms of λ_α^I and $\tilde{\lambda}_{\dot{\alpha}}^I$ that satisfy this equation are as follows²¹:

$$\lambda_\alpha^I = \begin{pmatrix} \sqrt{E+pc} & -\sqrt{E-ps^*} \\ \sqrt{E+ps} & \sqrt{E-pc} \end{pmatrix}, \quad \tilde{\lambda}_{\dot{\alpha}}^I = \begin{pmatrix} \sqrt{E-ps} & \sqrt{E+pc} \\ -\sqrt{E-pc} & \sqrt{E+ps^*} \end{pmatrix} \quad (\text{A.7})$$

Determinants of both of these spinors are equal to m . They happen to satisfy the following massive Dirac equations:

$$p_{\alpha\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}I} = -m \lambda_\alpha^I, \quad p_{\alpha\dot{\alpha}} \lambda^{\alpha I} = m \tilde{\lambda}_{\dot{\alpha}}^I \quad (\text{A.8})$$

Now to evaluate the high energy limit, it is preferable to decompose the aforementioned λ_α^I and $\tilde{\lambda}_{\dot{\alpha}}^I$ as follows:

$$\lambda_\alpha^I = \sqrt{E+p}\zeta_\alpha^+ \zeta^{-I} + \sqrt{E-p}\zeta_\alpha^- \zeta^{+I} \quad (\text{A.9})$$

and

$$\tilde{\lambda}_{\dot{\alpha}}^I = \sqrt{E+p}\tilde{\zeta}_{\dot{\alpha}}^- \zeta^{+I} + \sqrt{E-p}\tilde{\zeta}_{\dot{\alpha}}^+ \zeta^{-I} \quad (\text{A.10})$$

where $\zeta^{\mp I}$ are the basis vectors spanning two dimensional spinor space, chosen to be $\zeta^{-I} = \begin{pmatrix} 1 & 0 \end{pmatrix}$ and $\zeta^{+I} = \begin{pmatrix} 0 & 1 \end{pmatrix}$ so that they satisfy the normalization condition $\epsilon_{IJ}\zeta^{+I}\zeta^{-J} = 1$. With this choice of basis vectors, we have the following expressions for ζ_α^\pm and $\tilde{\zeta}_{\dot{\alpha}}^\mp$:

$$\zeta_\alpha^+ = \begin{pmatrix} c \\ s \end{pmatrix}, \quad \tilde{\zeta}_{\dot{\alpha}}^- = \begin{pmatrix} c \\ s^* \end{pmatrix}, \quad \zeta_\alpha^- = \begin{pmatrix} -s^* \\ c \end{pmatrix}, \quad \tilde{\zeta}_{\dot{\alpha}}^+ = \begin{pmatrix} s \\ -c \end{pmatrix} \quad (\text{A.11})$$

Here we define $\lambda_\alpha := \sqrt{E+p}\zeta_\alpha^+$, $\eta_\alpha := \sqrt{E-p}\zeta_\alpha^-$, $\tilde{\lambda}_{\dot{\alpha}} := \sqrt{E+p}\tilde{\zeta}_{\dot{\alpha}}^-$ and $\tilde{\eta}_{\dot{\alpha}} := \sqrt{E-p}\tilde{\zeta}_{\dot{\alpha}}^+$. This enables us to write equations (A.9) and (A.10) as follows:

$$\lambda_\alpha^I = \lambda_\alpha \zeta^{-I} + \eta_\alpha \zeta^{+I} \quad (\text{A.12})$$

and

$$\tilde{\lambda}_{\dot{\alpha}}^I = \tilde{\lambda}_{\dot{\alpha}} \zeta^{+I} + \tilde{\eta}_{\dot{\alpha}} \zeta^{-I} \quad (\text{A.13})$$

²¹The expression for $\tilde{\lambda}_{\dot{\alpha}}^I$ has a typo in [2] (eq. (C.2)) following which a systematic error is carried forward in there in further results from this point onwards. Their calculations of scattering amplitudes remain unaffected throughout their paper because the explicit forms are not required anywhere in those calculations.

Making identifications with angle and square spinor notations $\lambda_\alpha \rightarrow |p\rangle_\alpha \Rightarrow \lambda^\alpha \rightarrow \langle p|^\alpha$ and $\tilde{\lambda}_{\dot{\alpha}} \rightarrow [p]_{\dot{\alpha}} \Rightarrow \tilde{\lambda}^{\dot{\alpha}} \rightarrow [p]^{\dot{\alpha}}$, we obtain the following results:

$$\langle \lambda \eta \rangle = \lambda^\alpha \eta_\alpha = \epsilon^{\alpha\beta} \lambda_\beta \eta_\alpha = -m = -\langle \eta \lambda \rangle \quad (\text{A.14})$$

and

$$[\tilde{\lambda} \tilde{\eta}] = \tilde{\lambda}_{\dot{\alpha}} \tilde{\eta}^{\dot{\alpha}} = \tilde{\lambda}_{\dot{\alpha}} \epsilon^{\dot{\alpha}\dot{\beta}} \tilde{\eta}_{\dot{\beta}} = -m = -[\tilde{\eta} \tilde{\lambda}] \quad (\text{A.15})$$

Rewriting $\eta_\alpha = \frac{\sqrt{E+p}}{\sqrt{E+p}} \sqrt{E-p} \zeta_\alpha^- = m \hat{\eta}_\alpha$ where $\hat{\eta}_\alpha = \frac{\zeta_\alpha^-}{\sqrt{E+p}}$ and similarly $\tilde{\eta}_{\dot{\alpha}} = m \hat{\tilde{\eta}}_{\dot{\alpha}}$ where $\hat{\tilde{\eta}}_{\dot{\alpha}} = \frac{\tilde{\zeta}_{\dot{\alpha}}^+}{\sqrt{E+p}}$, equations (A.14) and (A.15) become the following²²:

$$\langle \lambda \hat{\eta} \rangle = -1, \quad [\tilde{\lambda} \hat{\tilde{\eta}}] = -1 \quad (\text{A.16})$$

A.2 Results in the Standard Convention $x^\mu = (t, +\vec{x})$

The conventions adopted in all our calculations are as follows (again, natural units are employed everywhere):

- $g_{\mu\nu} = (+, -, -, -)$
- $x^\mu = (t, +\vec{x}) \Rightarrow p_\mu = (E, -\vec{p})$
- $\epsilon_{ij} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \Rightarrow \epsilon^{ij} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This is the convention adopted for both $\epsilon_{\alpha\dot{\alpha}}$ and ϵ_{IJ} where α and $\dot{\alpha}$ are Lorentz indices while I and J are little group indices, used as raising/lowering operators.
- 4- vector Pauli matrices are $\sigma^\mu = (1, \vec{\sigma})$ and $\bar{\sigma}^\mu = (1, -\vec{\sigma})$ where $\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ and $\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

These are the standard convention that is found in literature. In this convention we get the momentum bispinor matrix as follows:

$$p_{\alpha\dot{\alpha}} = \sigma \cdot p = (\sigma^\mu)_{\alpha\dot{\alpha}} p_\mu = \begin{pmatrix} E - p \cos(\theta) & -p \sin(\theta) e^{-i\phi} \\ -p \sin(\theta) e^{i\phi} & E + p \cos(\theta) \end{pmatrix} \quad (\text{A.17})$$

²²In order to make $\langle \lambda \eta \rangle = [\tilde{\lambda} \tilde{\eta}] = m \Rightarrow \langle \lambda \hat{\eta} \rangle = [\tilde{\lambda} \hat{\tilde{\eta}}] = 1$, we must chose basis vectors $\zeta^{\mp I}$ such that $\epsilon_{IJ} \zeta^{+I} \zeta^{-J} = -1$, for example $\zeta^{-I} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\zeta^{+I} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. With this choice of basis vectors, eq. (A.11) gets modified accordingly as $\tilde{\zeta}_{\dot{\alpha}}^- = \begin{pmatrix} -c \\ -s^* \end{pmatrix}$, $\zeta_\alpha^- = \begin{pmatrix} s^* \\ -c \end{pmatrix}$ while ζ_α^+ and $\tilde{\zeta}_{\dot{\alpha}}^+$ remain unchanged. Equations (A.9) and (A.10) continue to hold by making these adjustments.

and

$$p^{\dot{\alpha}\alpha} = \bar{\sigma} \cdot p = (\bar{\sigma}^\mu)^{\dot{\alpha}\alpha} p_\mu = \begin{pmatrix} E + p \cos(\theta) & p \sin(\theta) e^{-i\phi} \\ p \sin(\theta) e^{i\phi} & E - p \cos(\theta) \end{pmatrix} \quad (\text{A.18})$$

A.2.1 Massless Case

Again we have $p_{\alpha\dot{\alpha}} = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}}$. The forms of λ and $\tilde{\lambda}$ that satisfy this equation are as follows:

$$\lambda_\alpha = \sqrt{2E} \begin{pmatrix} -s^* \\ c \end{pmatrix}, \quad \tilde{\lambda}_{\dot{\alpha}} = \sqrt{2E} \begin{pmatrix} -s \\ c \end{pmatrix} \quad (\text{A.19})$$

They happen to satisfy the following massless Dirac equations (or Weyl equations):

$$p_{\alpha\dot{\alpha}} \lambda^\alpha = 0, \quad p_{\alpha\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}} = 0 \quad (\text{A.20})$$

A.2.2 Massive Case

Again we have $p_{\alpha\dot{\alpha}} = \lambda_\alpha^I \tilde{\lambda}_{\dot{\alpha}I}$. The forms of λ_α^I and $\tilde{\lambda}_{\dot{\alpha}I}$ that satisfy this equation are as follows:

$$\lambda_\alpha^I = \begin{pmatrix} -\sqrt{E+ps^*} & -\sqrt{E-pc} \\ \sqrt{E+pc} & -\sqrt{E-ps} \end{pmatrix}, \quad \tilde{\lambda}_{\dot{\alpha}I} = \begin{pmatrix} \sqrt{E-pc} & -\sqrt{E+ps} \\ \sqrt{E-ps^*} & \sqrt{E+pc} \end{pmatrix} \quad (\text{A.21})$$

Determinants of both of these spinors are equal to m . They happen to satisfy the following massive Dirac equations:

$$p_{\alpha\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}I} = -m \lambda_\alpha^I, \quad p_{\alpha\dot{\alpha}} \lambda^{\alpha I} = m \tilde{\lambda}_{\dot{\alpha}I} \quad (\text{A.22})$$

Now to evaluate the high energy limit, it is preferable to decompose the aforementioned λ_α^I and $\tilde{\lambda}_{\dot{\alpha}I}$ as follows:

$$\lambda_\alpha^I = \sqrt{E+pc} \zeta_\alpha^+ \zeta^{-I} + \sqrt{E-pc} \zeta_\alpha^- \zeta^{+I} \quad (\text{A.23})$$

and

$$\tilde{\lambda}_{\dot{\alpha}I} = \sqrt{E+pc} \tilde{\zeta}_{\dot{\alpha}}^- \zeta^{+I} + \sqrt{E-pc} \tilde{\zeta}_{\dot{\alpha}}^+ \zeta^{-I} \quad (\text{A.24})$$

where $\zeta^{\mp I}$ are the basis vectors spanning two dimensional spinor space, chosen to be $\zeta^{-I} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\zeta^{+I} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ so that they satisfy the normalization condition $\epsilon_{IJ} \zeta^{+I} \zeta^{-J} = -1$. With this choice of basis vectors, we have the following expressions for ζ_α^\pm and $\tilde{\zeta}_{\dot{\alpha}}^\mp$:

$$\zeta_\alpha^+ = \begin{pmatrix} -s^* \\ c \end{pmatrix}, \quad \tilde{\zeta}_{\dot{\alpha}}^- = \begin{pmatrix} s \\ -c \end{pmatrix}, \quad \zeta_\alpha^- = \begin{pmatrix} c \\ s \end{pmatrix}, \quad \tilde{\zeta}_{\dot{\alpha}}^+ = \begin{pmatrix} c \\ s^* \end{pmatrix} \quad (\text{A.25})$$

Here we define $\lambda_\alpha := \sqrt{E+pc} \zeta_\alpha^+$, $\eta_\alpha := \sqrt{E-pc} \zeta_\alpha^-$, $\tilde{\lambda}_{\dot{\alpha}} := \sqrt{E+pc} \tilde{\zeta}_{\dot{\alpha}}^-$ and $\tilde{\eta}_{\dot{\alpha}} := \sqrt{E-pc} \tilde{\zeta}_{\dot{\alpha}}^+$. This enables us to write equations (A.23) and (A.24) as follows:

$$\lambda_\alpha^I = \lambda_\alpha \zeta^{-I} + \eta_\alpha \zeta^{+I} \quad (\text{A.26})$$

and

$$\tilde{\lambda}_\alpha^I = \tilde{\lambda}_\alpha \zeta^{+I} + \tilde{\eta}_\alpha \zeta^{-I} \quad (\text{A.27})$$

Making identifications with angle and square spinor notations $\lambda_\alpha \rightarrow |p\rangle_\alpha \Rightarrow \lambda^\alpha \rightarrow \langle p|^\alpha$ and $\tilde{\lambda}_{\dot{\alpha}} \rightarrow [p]_{\dot{\alpha}} \Rightarrow \tilde{\lambda}^{\dot{\alpha}} \rightarrow [p]^{\dot{\alpha}}$, we obtain the following results:

$$\langle \lambda \eta \rangle = \lambda^\alpha \eta_\alpha = \epsilon^{\alpha\beta} \lambda_\beta \eta_\alpha = +m = -\langle \eta \lambda \rangle \quad (\text{A.28})$$

and

$$[\tilde{\lambda} \tilde{\eta}] = \tilde{\lambda}_{\dot{\alpha}} \tilde{\eta}^{\dot{\alpha}} = \tilde{\lambda}_{\dot{\alpha}} \epsilon^{\dot{\alpha}\dot{\beta}} \tilde{\eta}_{\dot{\beta}} = +m = -[\tilde{\eta} \tilde{\lambda}] \quad (\text{A.29})$$

Rewriting $\eta_\alpha = \frac{\sqrt{E+p}}{\sqrt{E+p}} \sqrt{E-p} \zeta_\alpha^- = m \hat{\eta}_\alpha$ where $\hat{\eta}_\alpha = \frac{\zeta_\alpha^-}{\sqrt{E+p}}$ and similarly $\tilde{\eta}_{\dot{\alpha}} = m \hat{\eta}_{\dot{\alpha}}$ where $\hat{\eta}_{\dot{\alpha}} = \frac{\zeta_{\dot{\alpha}}^+}{\sqrt{E+p}}$, equations (A.28) and (A.29) become the following:

$$\langle \lambda \hat{\eta} \rangle = +1 = -\langle \hat{\eta} \lambda \rangle, \quad [\tilde{\lambda} \hat{\eta}] = +1 = -[\hat{\eta} \tilde{\lambda}] \quad (\text{A.30})$$

A.3 Obtaining Massive Spinors via Lorentz Boost

We can also construct massive helicity spinors in the following way using Lorentz boost but we are not going to use this approach and the corresponding explicit results elsewhere in the paper.

Consider a moving Dirac spinor in 'chiral' basis $\Psi = (\phi_R(v), \chi_L(v))^T$ where v is the speed, that takes the form $\phi_R(0) = \chi_L(0)$ (which is true for massive Dirac spinor at rest) and transforms as the following:

$$\Psi \rightarrow \begin{pmatrix} \Lambda(v) & 0 \\ 0 & \Lambda(-v) \end{pmatrix} \Psi \quad (\text{A.31})$$

This is a 4×4 matrix where $\Lambda(v)$ and $\Lambda(-v)$ are defined as follows:

$$\Lambda(v) = \exp \left(-\frac{\vec{\sigma} \cdot \vec{\rho}}{2} + i \frac{\vec{\sigma} \cdot \vec{\theta}}{2} \right) \quad (\text{A.32})$$

and

$$\Lambda(-v) = (\Lambda(v)^\dagger)^{-1} = \exp \left(\frac{\vec{\sigma} \cdot \vec{\rho}}{2} + i \frac{\vec{\sigma} \cdot \vec{\theta}}{2} \right) \quad (\text{A.33})$$

In the spinor space, the rest frame momentum bispinor is given by the matrix $M = \begin{pmatrix} m & 0 \\ 0 & m \end{pmatrix}$. Boosting this gives the momentum bispinor having a general momentum \vec{p} and energy E as shown in the following:

$$M \rightarrow N M N^\dagger \quad (\text{A.34})$$

where

$$N = \sqrt{\frac{E+m}{2m}} \left(1 - \frac{\vec{\sigma} \cdot \vec{p}}{E+m} \right) = \sqrt{\frac{E+m}{2m}} \begin{pmatrix} \left(1 - \frac{p \cos(\theta)}{E+m} \right) & -\frac{p \sin(\theta) e^{-i\phi}}{(E+m)} \\ -\frac{p \sin(\theta) e^{i\phi}}{(E+m)} & \left(1 + \frac{p \cos(\theta)}{E+m} \right) \end{pmatrix} \quad (\text{A.35})$$

N and N^\dagger have determinants equal to 1 as expected since N belongs to $SL(2, \mathbb{C})$ group.

This gives what we call $p_{\alpha\dot{\alpha}}$ as follows:

$$p_{\alpha\dot{\alpha}} = N M N^\dagger = \begin{pmatrix} E - p \cos(\theta) & -p \sin(\theta) e^{-i\phi} \\ -p \sin(\theta) e^{i\phi} & E + p \cos(\theta) \end{pmatrix} \quad (\text{A.36})$$

Similarly to obtain $p^{\dot{\alpha}\alpha}$, we use $M \rightarrow N' M N'^\dagger$ where N' is given by the following:

$$N' = \sqrt{\frac{E+m}{2m}} \left(1 + \frac{\vec{\sigma} \cdot \vec{p}}{E+m} \right) = \sqrt{\frac{E+m}{2m}} \begin{pmatrix} \left(1 + \frac{p \cos(\theta)}{E+m} \right) & \frac{p \sin(\theta) e^{-i\phi}}{(E+m)} \\ \frac{p \sin(\theta) e^{i\phi}}{(E+m)} & \left(1 - \frac{p \cos(\theta)}{E+m} \right) \end{pmatrix} \quad (\text{A.37})$$

N' and N'^\dagger have determinants equal to 1 as expected since N' belongs to $SL(2, \mathbb{C})$ group.

This gives the following result upon boosting from the rest frame:

$$p^{\dot{\alpha}\alpha} = N' M N'^\dagger = \begin{pmatrix} E + p \cos(\theta) & p \sin(\theta) e^{-i\phi} \\ p \sin(\theta) e^{i\phi} & E - p \cos(\theta) \end{pmatrix} \quad (\text{A.38})$$

Note that the conventions used by us are: $g_{\mu\nu} = (+, -, -, -)$, $\epsilon_{12} = -1$ where $\epsilon_{\alpha\beta}/\epsilon^{\alpha\beta}$ and $\epsilon_{\dot{\alpha}\dot{\beta}}/\epsilon^{\dot{\alpha}\dot{\beta}}$ are the lowering/raising operators and $x^\mu = (t, \vec{x}) \Rightarrow p^\mu = (E, \vec{p}) \Rightarrow p_\mu = (E, -\vec{p})$.

We can write the rank 2 matrix (eq. (A.36)) as $p_{\alpha\dot{\alpha}} = \lambda_\alpha^I \tilde{\lambda}_{\dot{\alpha}I}$. Writing the rest frame momentum bispinor matrix as $p_{o,\alpha\dot{\alpha}} = p_o^{\dot{\alpha}\alpha} = \begin{pmatrix} m & 0 \\ 0 & m \end{pmatrix} = \begin{pmatrix} \sqrt{m} \\ 0 \end{pmatrix} \begin{pmatrix} \sqrt{m} & 0 \end{pmatrix} + \begin{pmatrix} 0 \\ \sqrt{m} \end{pmatrix} \begin{pmatrix} 0 & \sqrt{m} \end{pmatrix}$, we get the following results for λ_α^I and $\tilde{\lambda}_{\dot{\alpha}I}$ obtained via boosting through N and N' respectively:

$$\lambda_\alpha^I = \frac{1}{\sqrt{2(E+m)}} \begin{pmatrix} E + m - p \cos(\theta) & -p \sin(\theta) e^{-i\phi} \\ -p \sin(\theta) e^{i\phi} & E + m + p \cos(\theta) \end{pmatrix} \quad (\text{A.39})$$

and

$$\tilde{\lambda}_{\dot{\alpha}I} = \frac{1}{\sqrt{2(E+m)}} \begin{pmatrix} p \sin(\theta) e^{i\phi} & E + m - p \cos(\theta) \\ -(E + m + p \cos(\theta)) & -p \sin(\theta) e^{-i\phi} \end{pmatrix} \quad (\text{A.40})$$

These satisfy:

- i. $p_{\alpha\dot{\alpha}} = \lambda_\alpha^I \tilde{\lambda}_{\dot{\alpha}I}$
- ii. $p_{\alpha\dot{\alpha}} \tilde{\lambda}^{\dot{\alpha}I} = -m \lambda_\alpha^I$
- iii. $p_{\alpha\dot{\alpha}} \lambda^{\alpha I} = m \tilde{\lambda}_{\dot{\alpha}I}$
- iv. $\det(\lambda) = \det(\tilde{\lambda}) = m$

B Six Dimensions

B.1 Conventions

The conventions adopted are as follows:

- $g_{\mu\nu} = (+, -, -, -)$
- $x^\mu = (t, +\vec{x}) \Rightarrow p_\mu = (E, -\vec{p})$
- $\epsilon_{ij} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \Rightarrow \epsilon^{ij} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This is the convention adopted for both $\epsilon_{\dot{\alpha}\dot{\beta}}$ and $\epsilon_{\alpha\beta}$ where $\dot{\alpha}$ and α belongs to the first and the second $SU(2)$ respectively of the $6D$ massless little group $SO(4) \cong SU(2) \times SU(2)$.
- The invariant tensor of the group $Sp(4)$ is taken to be $J_{IJ} = \begin{pmatrix} 0_{2 \times 2} & 1_{2 \times 2} \\ -1_{2 \times 2} & 0_{2 \times 2} \end{pmatrix}$.
- 4- vector Pauli matrices are $\sigma^\mu = (1, \vec{\sigma})$ and $\bar{\sigma}^\mu = (1, -\vec{\sigma})$ where $\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ and $\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.
- $6D$ -Pauli matrices are of the form $\Sigma^\mu = (\gamma_0, \gamma_1, \gamma_2, \gamma_3, \gamma_4, \gamma_5)$ and $\bar{\Sigma}^\mu = (\tilde{\gamma}_0, \tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3, \tilde{\gamma}_4, \tilde{\gamma}_5)$. They are taken to be as follows:
 - a. $\gamma_0 = i\sigma^1 \otimes \sigma^2, \tilde{\gamma}_0 = -i\sigma^1 \otimes \sigma^2$
 - b. $\gamma_1 = i\sigma^2 \otimes \sigma^3, \tilde{\gamma}_1 = i\sigma^2 \otimes \sigma^3$
 - c. $\gamma_2 = -\sigma^2 \otimes \sigma^0, \tilde{\gamma}_2 = \sigma^2 \otimes \sigma^0$
 - d. $\gamma_3 = -i\sigma^2 \otimes \sigma^1, \tilde{\gamma}_3 = -i\sigma^2 \otimes \sigma^1$
 - e. $\gamma_4 = -\sigma^3 \otimes \sigma^2, \tilde{\gamma}_4 = \sigma^3 \otimes \sigma^2$
 - f. $\gamma_5 = i\sigma^0 \otimes \sigma^2, \tilde{\gamma}_5 = i\sigma^0 \otimes \sigma^2$

In this convention we get the momentum bispinor matrix as follows:

$$p_{AB} = \Sigma \cdot p = (\Sigma^\mu)_{AB} p_\mu = \begin{pmatrix} 0 & -ip_4 - p_5 & -p_1 - ip_2 & E + p_3 \\ ip_4 + p_5 & 0 & p_3 - E & p_1 - ip_2 \\ p_1 + ip_2 & E - p_3 & 0 & ip_4 - p_5 \\ -E - p_3 & ip_2 - p_1 & p_5 - ip_4 & 0 \end{pmatrix} \quad (\text{B.1})$$

and

$$p^{AB} = \bar{\Sigma} \cdot p = (\bar{\Sigma}^\mu)^{AB} p_\mu = \begin{pmatrix} 0 & ip_4 - p_5 & -p_1 + ip_2 & p_3 - E \\ -ip_4 + p_5 & 0 & p_3 + E & p_1 + ip_2 \\ p_1 - ip_2 & -E - p_3 & 0 & -ip_4 - p_5 \\ E - p_3 & -p_1 - ip_2 & p_5 + ip_4 & 0 \end{pmatrix} \quad (\text{B.2})$$

where $\vec{p} = (p_1, p_2, p_3, p_4, p_5)$ is used.

B.2 Angle/Square Brackets and Lorentz Invariant Objects in 6D

We introduce the angle and square brackets notation using the following equations:

$$p_{AB} = \tilde{\lambda}_A^I \tilde{\lambda}_{BI} = |p\rangle_A^I [p]_{BI}, \quad p^{AB} = \lambda_I^A \lambda^{BI} = |p\rangle_I^A \langle p|^{BI} \quad (\text{B.3})$$

As we can see, the same spinor solutions are used to construct each type of momentum matrix, hence the following results must hold:

$$[p]_{AI} J^{IJ} = [p]_A^J = |p\rangle_A^J, \quad |p\rangle_I^A J^{IJ} = |p\rangle^{AJ} = \langle p|^{AJ} \quad (\text{B.4})$$

where J^{IJ} is the inverse of J_{IJ} which is the tensor left invariant under $Sp(4)$ group transformation.

Let particles be labelled by lower-case Latin indices. With all the aforementioned conventions, various Lorentz invariant quantities can be constructed that transform covariantly under the little group transformation. They are as follows:

- i. $\lambda_i^{AI} \tilde{\lambda}_{jA}^J = \tilde{\lambda}_{jA}^I \lambda_i^{AJ} = \langle i|j\rangle$ where $\langle i|j\rangle$ has little group indices as $\langle i^I|j^J\rangle$.
- ii. $\lambda_{iI}^A \tilde{\lambda}_{jAJ} = \tilde{\lambda}_{jAI} \lambda_{iJ}^A = [j|i]$ but due to (B.4), this is essentially same as (i) above. Here too the little group indices are $[j_J|i_I]$.
- iii. $\epsilon_{ABCD} \lambda_i^{AI} \lambda_j^{BJ} \lambda_k^{CK} \lambda_l^{DL} = \langle i^I j^J k^K l^L\rangle$ that can be simply written as $\langle ijkl\rangle$.
- iv. $\epsilon^{ABCD} \tilde{\lambda}_{iAI} \tilde{\lambda}_{jBJ} \tilde{\lambda}_{kCK} \tilde{\lambda}_{lDL} = [i_I j_J k_K l_L]$ that can be simply written as $[ijkl]$.

Using (B.3) we see that the mass dimensions of each angle and square spinor is \sqrt{m} . Constructing the scalar product $p \cdot q$ using these and $Tr[\Sigma^\mu \bar{\Sigma}^\nu] = Tr[\bar{\Sigma}^\mu \Sigma^\nu] = 4\eta^{\mu\nu}$, we get the following results:

$$4p \cdot q = 4\eta^{\mu\nu} p_\mu q_\nu = \langle q^{AJ} | p_A^I \rangle [p_{BI} | q_J^B] \quad (\text{B.5})$$

The momentum conservation $\sum_i p_i = 0$ takes the following form using (B.3):

$$\sum_i \langle q|i\rangle [i|k] = 0 \quad (\text{B.6})$$

or

$$\sum_i [q|i] \langle i|k \rangle = 0 \quad (\text{B.7})$$

where q and k are reference spinors. Eqs. (B.6) and (B.7) follow from first and second equations of (B.3) respectively.

The explicit form of $[p|q\rangle$ calculated using the spinor solutions obtained earlier is as follows:

$$[p|q\rangle = m J_{IJ} \Rightarrow \text{Det}([p|q\rangle) = (m^2)^2 \quad (\text{B.8})$$

In massless case, determinant becomes 0 thereby rendering $[p|q\rangle$ to be rank 1 matrix whereas in the massive case here, the rank is 2. So there is no problem in taking inverses of these Lorentz invariant, little group covariant objects in the massive case unlike the massless case.

B.3 $p^{AB} = \lambda^{AI} \lambda_I^B$

Solving the massless Dirac equation $p_{AB} \lambda^{Aa} = 0$, where a is the second $SU(2)$ little group index of the $SO(4) \cong SU(2) \times SU(2)$, gives the massless spinor solution λ^{Aa} in the limit $E \rightarrow p$ where p denotes the norm of \vec{p} . Similarly another spinor solution $\eta^{A\dot{a}}$ is obtained by taking the massless limit $E \rightarrow -p$. The latter (unphysical) massless limiting condition is important in order to systematically construct the massive spinor as shown in eq. (B.11). Both of these solutions correspond to zero eigenvalue of the corresponding momentum bispinor matrix. They satisfy the following equations:

$$p^{AB} = \lambda^{Aa} \lambda_a^B, p'^{AB} = \eta^{A\dot{a}} \eta_{\dot{a}}^B \quad (\text{B.9})$$

where $\vec{p} \rightarrow -\vec{p}$ in p'^{AB} . The forms of both the spinors are as follows:

$$\lambda_a^A = \frac{1}{\sqrt{-ip_4 - p_5}} \begin{pmatrix} p_1 - ip_2 & p_3 - p \\ -(p + p_3) & p_1 + ip_2 \\ 0 & -ip_4 - p_5 \\ -ip_4 - p_5 & 0 \end{pmatrix}, \eta^{A\dot{a}} = \frac{1}{\sqrt{ip_4 + p_5}} \begin{pmatrix} -(p + p_3) & p_1 - ip_2 \\ -(p_1 + ip_2) & p - p_3 \\ ip_4 + p_5 & 0 \\ 0 & -(ip_4 + p_5) \end{pmatrix} \quad (\text{B.10})$$

The massive case in $D = 6$ has the same Lorentz group $SU^*(4)$ but the little group changes to $SO(5) \cong Sp(4)$. Then the massive spinor λ^{AI} (index I runs over the group $Sp(4)$) is constructed using the above calculated massless spinors as follows:

$$\lambda^{AI} = \sqrt{\frac{E+p}{2p}} \lambda_a^A e^{+Ia} + \sqrt{\frac{E-p}{2p}} \eta^{A\dot{a}} e_{\dot{a}}^{-I} \quad (\text{B.11})$$

where e^{+Ia} and $e_{\dot{a}}^{-I}$ are the basis vectors in the spinor space explicitly given as follows:

$$e^{+Ia} = \begin{pmatrix} 0 & +i & 0 & 0 \\ 0 & 0 & 0 & +i \end{pmatrix} \quad (\text{B.12})$$

and

$$e_{\dot{a}}^{-I} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & +1 & 0 \end{pmatrix} \quad (\text{B.13})$$

where the basis vectors satisfy the following normalization conditions:

$$e^{+Ia} J_{IJ} e^{+Jb} = e_{\dot{a}}^{-I} J_{IJ} e_{\dot{b}}^{-J} = \begin{pmatrix} 0 & -1 \\ +1 & 0 \end{pmatrix}, e^{+Ia} J_{IJ} e_{\dot{b}}^{-J} = e_{\dot{a}}^{-I} J_{IJ} e^{+Jb} = 0 \quad (\text{B.14})$$

The coefficients in eq. (B.11) are chosen in order to satisfy the determinant condition $\det(\lambda^{AI}) = m^2$ which is natural consequence of the equation $p^{AB} = \lambda^{AI} \lambda_I^A$ where $\det(p^{AB}) = (m^2)^2$. Also the coefficients conveniently ensure that the massless condition in the limit $E \rightarrow p$ is recovered smoothly. Only the first term survives and the coefficient goes to 1.

Plugging eqs. (B.10), (B.12) and (B.13) in eq. (B.11) give the following massive spinor form:

$$\lambda_I^A = \frac{1}{\sqrt{-ip_4 - p_5}} \begin{pmatrix} \sqrt{\frac{E-p}{2p}}(ip_1 + p_2) & -i\sqrt{\frac{E+p}{2p}}(p - p_3) & -i\sqrt{\frac{E-p}{2p}}(p + p_3) & -\sqrt{\frac{E+p}{2p}}(ip_1 + p_2) \\ i\sqrt{\frac{E-p}{2p}}(p - p_3) & \sqrt{\frac{E+p}{2p}}(ip_1 - p_2) & \sqrt{\frac{E-p}{2p}}(p_2 - ip_1) & i\sqrt{\frac{E+p}{2p}}(p + p_3) \\ 0 & i\sqrt{\frac{E+p}{2p}}(p_4 - ip_5) & \sqrt{\frac{E-p}{2p}}(-p_4 + ip_5) & 0 \\ \sqrt{\frac{E-p}{2p}}(p_4 - ip_5) & 0 & 0 & -i\sqrt{\frac{E+p}{2p}}(p_5 - ip_4) \end{pmatrix} \quad (\text{B.15})$$

This satisfies the following equations:

- i. $\text{Det}(\lambda^{AI}) = (E - p)(E + p) = m^2$
- ii. $p^{AB} = \lambda^{AI} \lambda_I^B$. Note that the indices are written in the same order as the massless case, namely $p^{AB} = \lambda^{Aa} \lambda_a^B$.
- iii. Massless limit $E \rightarrow p$ matches with the known massless spinor helicity formalism where the spinor satisfies the massless Dirac equation $p_{AB} \lambda_a^A = 0$ or $p_{AB} \lambda^{Aa} = 0$.
- iv. Actions of momentum bispinor acting on massless spinors are:

$$p_{AB} \lambda_a^A = (E - p) \tilde{\eta}_{Ba}, \quad p_{AB} \eta^{A\dot{a}} = (E + p) \tilde{\lambda}_{\dot{B}}^{\dot{a}} \quad (\text{B.16})$$

B.4 $p_{AB} = \tilde{\lambda}_{AI} \tilde{\lambda}_B^I$

We can write massive spinor $\tilde{\lambda}_A^I$ in terms of massless spinors $\tilde{\lambda}_A^{\dot{a}}$ and $\tilde{\eta}_{Aa}$ as follows:

$$\tilde{\lambda}_A^I = \sqrt{\frac{E+p}{2p}} \tilde{\lambda}_A^{\dot{a}} e_a^{-I} + \sqrt{\frac{E-p}{2p}} \tilde{\eta}_{Aa} e^{+Ia} \quad (\text{B.17})$$

Here \dot{a} belongs to the first $SU(2)$ of the massless little group $SO(4) \cong SU(2) \times SU(2)$ and I belongs to the massive little group $Sp(4)$ which is congruent to $SO(5)$. The massless spinors are obtained by solving the massless Dirac equation $p^{AB} \tilde{\lambda}_A^{\dot{a}} = 0$ and $p'^{AB} \tilde{\eta}_{Aa} = 0$ where p'^{AB} has $\vec{p} \rightarrow -\vec{p}$. Both these spinor solutions correspond to zero eigenvalue of the corresponding momentum bispinor matrices. They satisfy $p_{AB} = \tilde{\lambda}_{A\dot{a}} \tilde{\lambda}_B^{\dot{a}}$ and $p'_{AB} = \tilde{\eta}_{Aa} \tilde{\eta}_B^a$ where again p'_{AB} means that the substitution $\vec{p} \rightarrow -\vec{p}$ is made in p_{AB} . Again, the unphysical massless limiting condition $E \rightarrow -p$ is required to systematically construct the massive spinor as clear in eq. (B.17).

The basis vectors in the spinor space, e_a^{-I} and e^{+Ia} , are the same objects as used in eq. (B.11). Their explicit forms are exactly the same as in eqs. (B.13) and (B.12) respectively. This makes the normalization of the basis vectors same as in eq. (B.14).

The coefficients in eq. (B.17) are decided by the condition $\det(\tilde{\lambda}_A^I) = m^2$ which is a natural consequence of the relation $p_{AB} = \tilde{\lambda}_{AI} \tilde{\lambda}_B^I$ where $\det(p_{AB}) = (m^2)^2$. Also the coefficients ensure that the massless limit is recovered smoothly when $E \rightarrow p$. In this limit, the second term dies and the coefficient of first term goes to 1, thereby leaving only $\tilde{\lambda}_A^{\dot{a}}$ which was obtained in the first place by solving the massless Dirac equation in the limit $E \rightarrow p$.

Plugging everything in eq. (B.17) gives the massive spinor $\tilde{\lambda}_A^I$ as follows:

$$\tilde{\lambda}_{AI} = \frac{1}{\sqrt{-ip_4 - p_5}} \begin{pmatrix} \sqrt{\frac{E+p}{2p}}(-p_4 + ip_5) & 0 & 0 & \sqrt{\frac{E-p}{2p}}(p_4 - ip_5) \\ 0 & \sqrt{\frac{E-p}{2p}}(p_4 - ip_5) & \sqrt{\frac{E+p}{2p}}(-p_4 + ip_5) & 0 \\ -i\sqrt{\frac{E+p}{2p}}(p - p_3) & \sqrt{\frac{E-p}{2p}}(p_2 - ip_1) & i\sqrt{\frac{E+p}{2p}}(p_1 + ip_2) & -i\sqrt{\frac{E-p}{2p}}(p + p_3) \\ i\sqrt{\frac{E+p}{2p}}(p_1 - ip_2) & -i\sqrt{\frac{E-p}{2p}}(p - p_3) & -i\sqrt{\frac{E+p}{2p}}(p + p_3) & -\sqrt{\frac{E-p}{2p}}(ip_1 + p_2) \end{pmatrix} \quad (\text{B.18})$$

This satisfies the following equation:

- i. $\text{Det}(\tilde{\lambda}_A^I) = (E - p)(E + p) = m^2$
- ii. $p_{AB} = \tilde{\lambda}_{AI} \tilde{\lambda}_B^I$. Note that the indices are written in the same order as the massless case, namely $p_{AB} = \tilde{\lambda}_{A\dot{a}} \tilde{\lambda}_B^{\dot{a}}$.
- iii. Massless limit $E \rightarrow p$ matches with the known massless spinor helicity formalism satisfying the massless Dirac equation $p^{AB} \tilde{\lambda}_{A\dot{a}} = 0$ or $p^{AB} \tilde{\lambda}_A^{\dot{a}} = 0$.

iv. Actions of momentum bispinor acting on massless spinors are:

$$p^{AB}\tilde{\lambda}_A^{\dot{a}} = (E - p)\eta^{B\dot{a}}, \quad p^{AB}\tilde{\eta}_{A\dot{a}} = (E + p)\lambda_a^B \quad (\text{B.19})$$

B.5 Massive Dirac Equation

The spinors λ^{AI} as constructed in eq. (B.11) and explicitly shown in eq. (B.15) as well as $\tilde{\lambda}_A^I$ in eqs. (B.17) and (B.18) satisfy the following massive Dirac equation in $6D$:

$$p_{AB}\lambda^{AI} = m\tilde{\lambda}_B^I, \quad p^{AB}\tilde{\lambda}_A^I = m\lambda^{BI} \quad (\text{B.20})$$

C Summary of Important Conventions/Definitions in $6D$

To summarize, we are using the following conventions/definitions:

$$p^{AB}p_{BC} = m^2\delta_C^A; \quad p_{AB}p^{BC} = m^2\delta_A^C \quad (\text{C.1})$$

$$p_{AB}\lambda^{AI} = m\tilde{\lambda}_B^I; \quad p^{AB}\tilde{\lambda}_A^I = m\lambda^{BI} \quad (\text{C.2})$$

$$p^{AB} = \lambda^{AI}\lambda_I^B; \quad p_{AB} = \tilde{\lambda}_{AI}\tilde{\lambda}_B^I \quad (\text{C.3})$$

$$\lambda_I^A\tilde{\lambda}_{AJ} = mJ_{IJ}; \quad \lambda^{AI}\tilde{\lambda}_A^J = -mJ^{IJ} \quad (\text{C.4})$$

$$\lambda^{AI}\tilde{\lambda}_{AJ} = m\delta_J^I; \quad \lambda_I^A\tilde{\lambda}_A^J = -m\delta_I^J \quad (\text{C.5})$$

$$\lambda^{AI}\tilde{\lambda}_{BI} = m\delta_B^A; \quad \lambda_I^A\tilde{\lambda}_B^I = -m\delta_B^A \quad (\text{C.6})$$

$$J_{IJ}\lambda^{AJ} = \lambda_I^A; \quad J^{IJ}\lambda_J^A = \lambda^{AI} \quad (\text{C.7})$$

$$J_{IJ}J^{JK} = \delta_I^K \quad (\text{C.8})$$

$$e_{\dot{a}}^{-I}J_{IJ}e_{\dot{b}}^{-J} = \epsilon_{\dot{a}\dot{b}}; \quad e^{+I\dot{a}}J_{IJ}e^{+J\dot{b}} = -\epsilon^{ab} \quad (\text{C.9})$$

$$p_{AB}\lambda^{Aa} = (E - p)\tilde{\eta}_B^a; \quad p_{AB}\eta_{\dot{a}}^A = (E + p)\tilde{\lambda}_{B\dot{a}} \quad (\text{C.10})$$

$$p^{AB}\tilde{\eta}_A^a = (E + p)\lambda^{Ba}; \quad p^{AB}\tilde{\lambda}_{A\dot{a}} = (E - p)\eta_{\dot{a}}^B \quad (\text{C.11})$$

$$\lambda^{Ab}\epsilon_{ab} = \lambda_a^A; \quad \tilde{\lambda}_{Ab}\epsilon^{\dot{a}\dot{b}} = \tilde{\lambda}_{\dot{a}}^A \quad (\text{C.12})$$

$$\lambda^{Aa}\tilde{\eta}_A^b = -2p\epsilon^{ab}; \quad \eta_{\dot{a}}^A\tilde{\lambda}_{Ab} = -2p\epsilon_{\dot{a}b} \quad (\text{C.13})$$

$$p^{AB} = \frac{1}{2}\epsilon^{ABCD}p_{CD} \quad (\text{C.14})$$

$$p_{AB}q^{AB} = -4p.q \quad (\text{C.15})$$

$$[p_{\dot{b}}q_{\dot{e}}]^{-1}\langle q_{\dot{e}}p_{\dot{a}} \rangle = \delta_{\dot{a}}^{\dot{b}}; \quad \langle q_{\dot{b}}p_{\dot{a}} \rangle^{-1}[p_{\dot{a}}q_{\dot{c}}] = \delta_c^{\dot{b}} \quad (\text{C.16})$$

$$\langle p_a q^b \rangle^{-1} = -\frac{\langle p^a q_b \rangle}{2p \cdot q} \quad (\text{C.17})$$

$$\langle p^a | q | p^b \rangle = -2\epsilon^{ab}(p \cdot q); \quad [p^a | q | p^b] = +2\epsilon^{ab}(p \cdot q) \quad (\text{C.18})$$

The identity involving Levi-Civita tensor of $SU(4)$ are as follows:

$$\epsilon_{ABEF} \epsilon^{CDGH} = \begin{vmatrix} \delta_A^C & \delta_B^C & \delta_E^C & \delta_F^C \\ \delta_A^D & \delta_B^D & \delta_E^D & \delta_F^D \\ \delta_A^G & \delta_B^G & \delta_E^G & \delta_F^G \\ \delta_A^H & \delta_B^H & \delta_E^H & \delta_F^H \end{vmatrix} \quad (\text{C.19})$$

$$\begin{aligned} &= \delta_A^C [\delta_B^D (\delta_E^G \delta_F^H - \delta_E^H \delta_F^G) - \delta_E^D (\delta_B^G \delta_F^H - \delta_B^H \delta_F^G) + \delta_F^D (\delta_B^G \delta_E^H - \delta_B^H \delta_E^G)] \\ &\quad - \delta_B^C [\delta_A^D (\delta_E^G \delta_F^H - \delta_E^H \delta_F^G) - \delta_E^D (\delta_A^G \delta_F^H - \delta_A^H \delta_F^G) + \delta_F^D (\delta_A^G \delta_E^H - \delta_A^H \delta_E^G)] \\ &\quad + \delta_E^C [\delta_A^D (\delta_B^G \delta_F^H - \delta_B^H \delta_F^G) - \delta_B^D (\delta_A^G \delta_F^H - \delta_A^H \delta_F^G) + \delta_F^D (\delta_A^G \delta_B^H - \delta_A^H \delta_B^G)] \\ &\quad - \delta_F^C [\delta_A^D (\delta_B^G \delta_E^H - \delta_B^H \delta_E^G) - \delta_B^D (\delta_A^G \delta_E^H - \delta_A^H \delta_E^G) + \delta_E^D (\delta_A^G \delta_B^H - \delta_A^H \delta_B^G)] \end{aligned} \quad (\text{C.20})$$

References

- [1] C. Cheung, ‘‘TASI Lectures on Scattering Amplitudes,’’ arXiv:1708.03872 [hep-ph].
- [2] N. Arkani-Hamed, T. C. Huang and Y. t. Huang, ‘‘Scattering Amplitudes For All Masses and Spins,’’ arXiv:1709.04891 [hep-th].
- [3] H. Elvang and Y. t. Huang, ‘‘Scattering Amplitudes,’’ arXiv:1308.1697 [hep-th].
- [4] C. Cheung and D. O’Connell, ‘‘Amplitudes and Spinor-Helicity in Six Dimensions,’’ JHEP **0907**, 075 (2009) [arXiv:0902.0981 [hep-th]].
- [5] H. Georgi, ‘‘Lie Algebras In Particle Physics. From Isospin To Unified Theories,’’ Front. Phys. **54**, 1 (1982).
- [6] J. Plefka, T. Schuster and V. Verschinin, ‘‘From Six to Four and More: Massless and Massive Maximal Super Yang-Mills Amplitudes in $6D$ and $4D$ and their Hidden Symmetries,’’ JHEP **1501**, 098 (2015) [arXiv:1405.7248 [hep-th]].
- [7] H. Kawai, D. C. Lewellen and S. H. H. Tye, ‘‘A Relation Between Tree Amplitudes of Closed and Open Strings,’’ Nucl. Phys. B **269**, 1 (1986).
- [8] R. Britto, F. Cachazo, B. Feng and E. Witten, ‘‘Direct proof of tree-level recursion relation in Yang-Mills theory,’’ Phys. Rev. Lett. **94**, 181602 (2005) [arXiv:hep-th/0501052].

- [9] S. D. Badger, E. W. N. Glover, V. V. Khoze and P. Svrcek, “Recursion relations for gauge theory amplitudes with massive particles,” *JHEP* **0507**, 025 (2005) [arXiv:hep-th/0504159].