

DETERMINANTAL HYPERSURFACES AND REPRESENTATIONS OF COXETER GROUPS

Ž. ČUČKOVIČ, M. I. STESSIN, AND A. B. TCHERNEV

ABSTRACT. Given a finite generating set $T = \{g_0, \dots, g_n\}$ of a group G , and a representation ρ of G on a Hilbert space V , we investigate how the geometry of the set $D(T, \rho) = \{[x_0 : \dots : x_n] \in \mathbb{C}\mathbb{P}^n \mid \sum x_i \rho(g_i) \text{ not invertible}\}$ reflects the properties of ρ . When V is finite-dimensional this is an algebraic hypersurface in $\mathbb{C}\mathbb{P}^n$. In the special case $T = G$ and $\rho =$ the left regular representation of G , this hypersurface is defined by the *group determinant*, an object studied extensively in the founding work of Frobenius that lead to the creation of representation theory. We focus on the classic case when G is a finite Coxeter group, and make T by adding the identity element 1_G to a Coxeter generating set for G . Under these assumptions we show in our first main result that if ρ is the left regular representation, then $D(T, \rho)$ determines the isomorphism class of G . Our second main result is that if G is not of exceptional type, and ρ is any finite dimensional representation, then $D(T, \rho)$ determines ρ .

1. INTRODUCTION

Given a tuple of linear operators A_0, \dots, A_n , on a complex N -dimensional vector space V , the determinant

$$\mathcal{P}(x_0, \dots, x_n) = \det [x_0 A_0 + \dots + x_n A_n]$$

is a homogeneous polynomial in x_0, \dots, x_n of degree N . Zeros of this polynomial form an algebraic closed subscheme of complex projective space $\mathbb{C}\mathbb{P}^n$ called a **determinantal hypersurface**, and we denote it by $\sigma(A_0, \dots, A_n)$.

Determinantal hypersurfaces are objects with a long history in algebraic geometry. A major direction of research related to their study is to determine which hypersurfaces in $\mathbb{C}\mathbb{P}^n$ are determinantal. This is a classical avenue, see e.g. [4, 12, 13, 23, 24, 30, 34, 39].

In this paper we follow another approach to determinantal hypersurfaces, where one analyzes what the geometry of $\sigma(A_0, \dots, A_n)$ can say about the mutual relations between the operators A_0, \dots, A_n . While this kind of question is natural from the perspective of operator theory, until recently it seems to have attracted much less attention in geometry.

One of the few relevant instances of work along these lines is the result of Motzkin and Taussky [32] that a real curve in $\mathbb{C}\mathbb{P}^2$ having a self-adjoint determinantal representation with one of the three operators being invertible (and, therefore, assumed to be the identity) satisfies the condition: the operators commute if and only if

2010 *Mathematics Subject Classification*. Primary: 20F55, 20C33, 14J70, 47A25; Secondary: 47A67.

Key words and phrases. determinantal hypersurfaces, Coxeter groups, joint spectrum, group representations.

this determinantal curve is a union of projective lines (in [32] the result is stated in equivalent but different form).

Another — and the most relevant to the subject of this paper — instance where the geometry of a determinantal hypersurface has been extensively studied from our point of view comes from classic work of Frobenius that goes back to the origins of representation theory. Specifically, when $G = \{g_0, \dots, g_n\}$ is a finite group and A_i represents the action of g_i by left multiplication on the group ring $\mathbb{C}[G]$ (this is usually called the *left regular representation* of G), the defining equation of $\sigma(A_0, \dots, A_n)$ is called the *group determinant* of G . The papers of Frobenius [17, 18, 19] show, in modern language, that the irreducible components of $\sigma(A_0, \dots, A_n)$ are in bijective correspondence with the irreducible representations of G , and the multiplicity of each component equals the dimension of the corresponding irreducible representation. For more on the group determinant and related problems we refer the reader to [8, 9, 10, 11, 16, 26, 27, 29] and the references there. In particular, it was shown by Formanek and Sibley [16] that the group determinant determines the isomorphism class of the group as well.

A substantial drawback when working with group determinants is that, as the size of the group rises, the resulting increase in complexity makes the group determinant essentially inaccessible. It is natural to ask if it is possible, instead of all elements, to take only a suitable generating set for G , thus computing a much simpler determinantal hypersurface while still retaining all the salient features of the group determinant. Furthermore, it is desirable to have analogous results also for finitely generated groups that are not necessarily finite. To do this one would need a generalization for the notion of a determinantal hypersurface to the setting of operators on an infinite dimensional space. This naturally leads us to the notion of **projective joint spectrum** for operators acting on a Hilbert space V , which was introduced by Yang [40]. If A_0, \dots, A_n are bounded linear operators on V , their joint spectrum is the set

$$\sigma(A_0, \dots, A_n) = \{[x_0 : \dots : x_n] \in \mathbb{C}\mathbb{P}^n : x_0 A_0 + \dots + x_n A_n \text{ is not invertible}\}.$$

When V is finite dimensional, the joint spectrum is a determinantal hypersurface. Since in this paper we predominantly deal with the finite dimensional case, we will use the terms “determinantal hypersurface” and “joint spectrum” interchangeably. The relation between the geometry of the joint spectrum (or its complement, the **joint resolvent set**) and the properties of the tuple A_0, \dots, A_n was investigated in [2, 3, 5, 14, 15, 21, 22, 25, 37, 36, 40]. In [22] joint spectrum was used in relation to the Gelfand-Naimark-Siegel representation of the infinite dihedral group.

Let $T = \{g_0, \dots, g_n\}$ be a set of generators of a group G , and let $\rho : G \rightarrow GL(V)$ be a homomorphism into the group of bounded invertible linear operators on a Hilbert space V . Let

$$D(T, \rho) = \sigma(\rho(g_0), \dots, \rho(g_n))$$

be the corresponding joint spectrum. If ρ is the left regular representation of G we will write just $D(T)$ (when G is infinite its left regular representation is obtained by the left action of G on the completion of $\mathbb{C}[G]$ with respect to its inner product norm, see Section 2). The results of Frobenius and of Formanek and Sibley naturally prompt us to ask:

- (1) When does $D(T)$ determine G ?
- (2) When does $D(T, \rho)$ determine ρ ?

- (3) If ρ is finite dimensional and irreducible, when is $D(T, \rho)$ reduced and irreducible?

We address these fundamental structural questions in the classical case when G is a finitely generated Coxeter group. We consider the generating set T obtained by adding the identity element 1_G to a Coxeter generating set for G ; see Section 2 for definitions. Our first main result concerns question (1):

Theorem 1.1. *Let G be a Coxeter group with Coxeter generating set $\{g_1, \dots, g_n\}$, and let $T = \{1_G, g_1, \dots, g_n\}$. Let G' be a group, and let $T' = \{1_{G'}, g'_1, \dots, g'_n\}$ be a generating set for G' . We have:*

- (1) *If $D(T) \supseteq D(T')$ as subsets of $\mathbb{C}\mathbb{P}^n$, then there is an epimorphism of groups $f : G \rightarrow G'$ such that $f(g_i) = g'_i$ for each $1 \leq i \leq n$. In particular, if G is finite then so is G' .*
- (2) *If G is finite and $D(T) = D(T')$ as subschemes of $\mathbb{C}\mathbb{P}^n$, then the homomorphism f from (1) is an isomorphism.*

Next, we consider question (2) for finite dimensional representations. It is easy to see that, when V is finite dimensional, $D(T, \rho)$ is completely determined by the multiset of isomorphism classes of irreducible representations that appear as factors in a composition series for V when considered as a $\mathbb{C}[G]$ -module via ρ . Thus whenever $\mathbb{C}[G]$ is not a semisimple ring, there will always exist non-equivalent representations of G having the same determinantal hypersurface. Note however, that this kind of obstruction does not arise if G is finite, or if one considers either only unitary or only irreducible representations.

In our second main result we present a positive answer to question (2) when G is a finite Coxeter group of regular type. The exceptional types will be treated in a separate paper.

Theorem 1.2. *Let G and T be as in Theorem 1.1. Suppose G is finite of type either I (dihedral group), or A , or B , or D . If for two finite dimensional complex linear representations ρ_1 and ρ_2 of G we have*

$$(1.3) \quad D(T, \rho_1) = D(T, \rho_2)$$

as subschemes of $\mathbb{C}\mathbb{P}^n$, then the representations ρ_1 and ρ_2 are equivalent.

In our proofs we make a conscious attempt to emphasize the direct connection between properties of the generating set T and the geometry of the corresponding determinantal hypersurface. Thus, in the case when G is dihedral, even though there are alternative arguments using known results from representation theory, we chose to rely on the tools developed in the recent paper [37], where the relation between the appearance of algebraic curves of finite multiplicity in the joint spectrum of a tuple of operators and decomposability of the tuple is investigated. In the case of unitary self-adjoint operators the technique developed there can be applied to finding commutation relations between the operators, and, therefore, be applied to representations of Coxeter groups. This method is quite general, and proved to be useful also for other, non-Coxeter groups. The upcoming paper [38] will contain the details.

To handle the cases A , B , and D in Theorem 1.2, we demonstrate that in those types $D(T, \rho)$ encodes the character of the representation ρ . We accomplish this by leveraging an explicit combinatorial algorithm that works uniformly in all three regular types, and transforms an element g of G into what we call **echelon form** –

an element with good combinatorial properties that belongs to the same conjugacy class, and is, in a precise sense, no bigger than g . The resulting combinatorics seems to be new and may be of independent interest.

The structure of the paper is as follows. Section 2 provides definitions and basic results about group representations and Coxeter groups. Section 3 contains the needed background material and results regarding joint spectra.

In Section 4 we analyze joint spectra of involution pairs. The computations reveal an interesting connection to Tchebyshev's polynomials. We use these results and give the proof of Theorem 1.1.

In Section 5 we work with finite dimensional unitary representations of dihedral groups. We demonstrate how one can use the tools from [37] to construct explicitly the decomposition of such a representation into irreducible factors directly from the defining equation of the joint spectrum. We use that to give a proof of Theorem 1.2 for the case when G is finite dihedral, and prove an analogous result for unitary representations of the infinite dihedral group. We also address question (3) for that setting. Of course, since the irreducible representations of dihedral groups are well known, these results can also be obtained by direct analysis of the determinantal hypersurfaces of these irreducible representations.

Sections 6 and 7 are devoted to developing the combinatorics needed for the proof of Theorem 1.2 for G of types A , B , or D . Section 6 considers words in the alphabet of the Coxeter generators of G , and introduces a partial ordering and *admissible transformations* on these words. We show that admissible transformations preserve conjugacy classes, and are non-increasing with respect to the partial ordering. Section 7 introduces echelon forms. The main result there is the “Ordering Theorem”, Theorem 7.5, which states that every word can be transformed into echelon form via a sequence of admissible transformations. This result is the key to our proof of Theorem 1.2, which we present in Section 8.

The combinatorics of admissible transformations can be extended to other Coxeter groups. In Section 9 we give an example how this can be used to study representations of the affine Coxeter group \tilde{C}_2 via determinantal hypersurfaces.

2. PRELIMINARIES

Let G be group. Recall that a (complex) representation of G is a group homomorphism $\rho : G \rightarrow GL(V)$ from G to the group of bounded invertible linear operators on a Hilbert space V . The representation ρ is called *unitary* if $\rho(w)$ is unitary for all $w \in G$; it is called *finite dimensional* if V is finite dimensional; and it is called *faithful* if ρ is injective. Two representations $\rho_1 : G \rightarrow GL(V_1)$ and $\rho_2 : G \rightarrow GL(V_2)$ are *equivalent* if there is a bounded linear isomorphism $C : V_1 \rightarrow V_2$ such that $\rho_2(w) = C\rho_1(w)C^{-1}$ for every $w \in G$.

The group ring $\mathbb{C}[G]$ has inner product $\langle \sum_{g \in G} a_g g, \sum_{g \in G} b_g g \rangle = \sum_{g \in G} a_g \bar{b}_g$ and corresponding induced norm $\|\sum a_g g\| = \sum |a_g|^2$. We write $\mathbb{C}[G]^\vee$ for the Hilbert space obtained by completing $\mathbb{C}[G]$ with respect to this norm. Of course, if G is finite then $\mathbb{C}[G]^\vee = \mathbb{C}[G]$. For each $g \in G$ the left multiplication by g on $\mathbb{C}[G]$ induces a bounded invertible linear operator $\lambda(g)$ on $\mathbb{C}[G]^\vee$, and the resulting map $\lambda : G \rightarrow GL(\mathbb{C}[G]^\vee)$ is a faithful unitary representation of G called the *left regular representation*.

When dealing with representations of finite groups we will rely heavily on the following basic fact (see [35]).

3. MORE ON JOINT SPECTRA

Recall that for an algebraic hypersurface in $\mathbb{C}\mathbb{P}^n$ or in \mathbb{C}^n defined by a polynomial $F = F_1^{r_1} \dots F_m^{r_m}$ (with each polynomial F_i irreducible and F_i not associate with F_j for $i \neq j$), the *components* of that hypersurface are defined by the polynomials $F_i^{r_i}$ (thus they are irreducible but not necessarily reduced), the *reduced components* are defined by the polynomials F_i , and the exponent r_i is called the *multiplicity* of the reduced component defined by F_i . We say that a point on our hypersurface is *regular (with multiplicity r)* if it belongs to only one reduced component (of multiplicity r), and is a regular point on that reduced component.

Let A be a bounded linear operator on a Hilbert space V . We consider its norm given by $\|A\| = \sup_{|u|=1} |A(u)|$. The *spectrum* of A is the set

$$\sigma(A) = \{\lambda \in \mathbb{C} \mid A - \lambda I \text{ is not invertible}\}.$$

Note that when A is self-adjoint we have $\sigma(A) \subseteq \mathbb{R}$ and $\|A\| = \sup_{\lambda \in \sigma(A)} |\lambda|$. We will use the following elementary consequences of the Spectral Theorem and the Spectral Mapping Theorem, see [31].

Proposition 3.1. *Let A be a bounded linear operator on a Hilbert space V .*

- (a) *If $p(x)$ is any polynomial then $\sigma(p(A)) = p(\sigma(A)) = \{p(\lambda) \mid \lambda \in \sigma(A)\}$.*
- (b) *If A is normal and $\sigma(A) = \{1\}$ then $A = I$, the identity operator.*

Now let A_1, A_2, \dots, A_n be bounded linear operators on a Hilbert space V . We will be considering the projective joint spectrum $\sigma(-I, A_1, \dots, A_n)$ of the tuple $-I, A_1, A_2, \dots, A_n$, where I is the identity operator, and its intersection with the chart $\{[x_0 : \dots : x_n] \mid x_0 \neq 0\}$. By taking $x_0 = 1$ we identify this intersection with a closed subset of \mathbb{C}^n called the *proper joint spectrum* of the tuple A_1, \dots, A_n , denoted by $\sigma_p(A_1, \dots, A_n)$; and when V is finite dimensional its defining polynomial is

$$\mathcal{F}(x_1, \dots, x_n) = \det(-I + x_1 A_1 + \dots + x_n A_n).$$

In particular, a point $x = (x_1, \dots, x_n) \in \mathbb{C}^n$ belongs to the proper joint spectrum if and only if the operator $A(x) = x_1 A_1 + \dots + x_n A_n$ has 1 in its spectrum. We note that when V is finite dimensional and $\sigma_p(A_1, \dots, A_n)$ is not empty (this latter is the case, for example, if A_i is invertible for some i), the projective joint spectrum $\sigma(-I, A_1, \dots, A_n)$ is the closure of the determinantal hypersurface $\sigma_p(A_1, \dots, A_n)$, and its defining polynomial is the homogenization of $\mathcal{F}(x_1, \dots, x_n)$.

We will be using the following basic observation multiple times:

Proposition 3.2. *Suppose V is finite dimensional, let $u = (u_1, u_2)$ be a regular point of $\sigma_p(A_1, A_2)$ of multiplicity r , and let Γ be the corresponding unique reduced component containing u .*

If the tangent line to Γ at u does not pass through the origin $(0, 0)$ then the operator $A(u) = u_1 A_1 + u_2 A_2$ has eigenvalue 1 of multiplicity exactly r .

Proof. Indeed, let $\mathcal{F}(x_1, x_2) = \det(-I + x_1 A_1 + x_2 A_2) = F^r H$ be the defining polynomial of $\sigma_p(A_1, A_2) \neq \emptyset$, where F is a defining irreducible polynomial of Γ , and H is not divisible by F . Then the defining polynomial $\mathcal{P}(x_0, x_1, x_2) = \det(-x_0 I + x_1 A_1 + x_2 A_2) = \tilde{F}^r \tilde{H}$ of $\sigma(-I, A_1, A_2)$ is the homogenization of \mathcal{F} , where \tilde{F} and \tilde{H} are the homogenizations of F and H . Note \mathcal{P} is also (as a polynomial in x_0) the characteristic polynomial of the operator $A(x)$. Thus we have $\mathcal{P}(x_0, u_1, u_2) = \tilde{F}^r(x_0, u_1, u_2) \tilde{H}(x_0, u_1, u_2) = (1 - x_0)^m \prod_{i=1}^k (\lambda_i - x_0)^{m_i}$ where the

λ_i are the eigenvalues of $A(u)$ different from 1, and the m_i their multiplicities. Since u is only on Γ , it follows that $\tilde{H}(1, u_1, u_2) = H(u) \neq 0$, hence $(1 - x_0)^m$ is a factor of $\tilde{F}^r(x_0, u_1, u_2)$. Thus it suffices to check that $x_0 = 1$ is not a double root of $\tilde{F}(x_0, u_1, u_2)$ when considered as a polynomial in x_0 . Since for the corresponding partial derivative in x_0 we have $\tilde{F}_{x_0}(1, u_1, u_2) = -u_1 F_{x_1}(u_1, u_2) - u_2 F_{x_2}(u_1, u_2)$, and the equation for the tangent line to Γ at u is $(x_1 - u_1)F_{x_1}(u) + (x_2 - u_2)F_{x_2}(u) = 0$, the desired conclusion is immediate from our assumption that the tangent line does not contain the origin. \square

It was shown in [37] that substantial information regarding mutual relations between the A_i s is captured by the geometry of their proper joint spectrum, and we review some of these results below, in the case of two operators A_1 and A_2 .

First we note the behaviour of the joint spectrum under change of coordinates. Let

$$\mathbf{C} = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$$

be an invertible complex-valued matrix. Write

$$(3.3) \quad B_1 = c_{11}A_1 + c_{12}A_2, \quad B_2 = c_{21}A_1 + c_{22}A_2.$$

Then it is immediate from the definitions that

$$(3.4) \quad \sigma_p(A_1, A_2) = \mathbf{C}^T \sigma_p(B_1, B_2).$$

Next, suppose V is finite dimensional and A_1 is self-adjoint. Let $\lambda \neq 0$ be an eigenvalue of A_1 , and suppose that $(1/\lambda, 0)$ is a regular point of the determinantal hypersurface $\sigma_p(A_1, A_2)$. Let $\mathcal{R}(x_1, x_2)$ be a defining polynomial for the reduced component of $\sigma_p(A_1, A_2)$ that contains $(1/\lambda, 0)$; thus one of the partial derivatives, say, $\mathcal{R}_{x_1} = \frac{\partial \mathcal{R}}{\partial x_1}$, does not vanish at $(1/\lambda, 0)$. Then the implicit function theorem implies that in a neighborhood of $(1/\lambda, 0)$ the equation $\{\mathcal{R} = 0\}$ determines x_1 as an implicit analytic function of x_2 defined in a neighborhood of the origin:

$$(3.5) \quad x_1 = x_1(x_2), \quad x_1(0) = \frac{1}{\lambda}.$$

For each eigenvalue w in the spectrum $\sigma(A_1)$ of A_1 let P_w be the orthogonal projection onto the w -eigensubspace of A_1 . We also set $P = P_\lambda$. Since A_1 is self-adjoint, the spectral decomposition for A_1 is

$$(3.6) \quad A_1 = \lambda P + \sum_{w \in \sigma(A_1) \setminus \{\lambda\}} w P_w.$$

We also introduce the following operator:

$$(3.7) \quad T(A_1) = T = \sum_{w \in \sigma(A_1) \setminus \{\lambda\}} \frac{\lambda}{w - \lambda} P_w.$$

The following result was proved (in significantly greater generality) during the course of [37, Proof of Theorem 7.3].

Theorem 3.8. *Suppose that V is finite dimensional, that A_1 and A_2 are self-adjoint, and that $\lambda \neq 0$ is an eigenvalue of A_1 such that*

- $(1/\lambda, 0)$ is a regular point of $\sigma_p(A_1, A_2)$; and
- $\mathcal{R}_{x_1}(1/\lambda, 0) \neq 0$, where $\mathcal{R}(x_1, x_2)$ is a defining polynomial for the reduced component of $\sigma_p(A_1, A_2)$ containing $(1/\lambda, 0)$.

Then we have

$$(3.9) \quad PA_2P = -x_1'(0)P$$

and

$$(3.10) \quad PA_2TA_2P = -\frac{x_1''(0)}{2}P,$$

where we regard x_1 as an implicit analytic function (3.5) of x_2 defined via the equation $\mathcal{R} = 0$.

4. JOINT SPECTRA AND COXETER GROUPS

Here we study properties of joint spectra of involution pairs A_1, A_2 , and use these to prove Theorem 1.1.

Lemma 4.1. *Let A_1 and A_2 be bounded linear operators on a Hilbert space V such that $A_1^2 = A_2^2 = I$. Then:*

- (1) *The set $\sigma_p(A_1, A_2) \cup -\sigma_p(A_1, A_2)$ is the union of all the “complex ellipses” $\mathcal{E}_\alpha = \{x^2 + \alpha xy + y^2 = 1\}$ with $\alpha \in \sigma(A_1A_2 + A_2A_1)$.*
- (2) *When $\sigma(A_1A_2 + A_2A_1)$ is a finite set then each connected component of $\sigma_p(A_1, A_2) \setminus \{(\pm 1, 0) (0, \pm 1)\}$ is either $L \setminus \{(\pm 1, 0) (0, \pm 1)\}$ with L one of the lines $x \pm y = \pm 1$, or $\mathcal{E}_\alpha \setminus \{(\pm 1, 0) (0, \pm 1)\}$ for some $\alpha \in \sigma(A_1A_2 + A_2A_1)$.*
- (3) *When V is finite dimensional each reduced component of $\sigma_p(A_1, A_2)$ is either a line of the form $x \pm y = \pm 1$, or a “complex ellipse” \mathcal{E}_α with $\alpha \in \sigma(A_1A_2 + A_2A_1) \setminus \{-2, 2\}$.*

Proof. Suppose that $(x, y) \in \sigma_p(A_1, A_2) \cup -\sigma_p(A_1, A_2)$. This is if and only if $1 \in \sigma(xA_1 + yA_2)$ or $-1 \in \sigma_p(xA_1 + yA_2)$, which, by Proposition 3.1, happens precisely when $1 \in \sigma((xA_1 + yA_2)^2)$. That in turn happens if and only if the operator

$$(4.2) \quad (xA_1 + yA_2)^2 - I = xy[A_1A_2 + A_2A_1] - (1 - x^2 - y^2)I$$

is not invertible. We see that when $xy \neq 0$ this is equivalent to $\alpha = \frac{1-x^2-y^2}{xy}$ being in the spectrum of $A_1A_2 + A_2A_1$, and (x, y) being a point on \mathcal{E}_α (and this last fact is true for trivial reasons also when $xy = 0$). This completes the proof of part (1).

When $\sigma(A_1A_2 + A_2A_1)$ is a finite set equation (4.2) implies that $\frac{1-x^2-y^2}{xy}$ is constant on every connected component of $\sigma_p(A_1, A_2) \setminus \{(\pm 1, 0) (0, \pm 1)\}$, which yields (2). Therefore, if V is finite dimensional, then each reduced component of $\sigma_p(A_1, A_2)$ is a reduced component of an ellipse \mathcal{E}_α for some $\alpha \in \sigma(A_1A_2 + A_2A_1)$. When $\alpha \neq \pm 2$ the ellipse is irreducible, so our reduced component coincides with the ellipse. When $\alpha = \pm 2$ the ellipse is the union of two of the lines $x \pm y = \pm 1$, hence our reduced component is one of these lines. \square

Lemma 4.3. *Let A_1 and A_2 be as in Lemma 4.1, and let $m \geq 2$ be an integer. If*

$$(1) \quad (A_1A_2)^m = I$$

then

$$(2) \quad \sigma(A_1A_2 + A_2A_1) \subseteq \{2 \cos(2\pi k/m) \mid k = 0, \dots, m-1\}.$$

If both A_1 and A_2 are unitary, then conditions (1) and (2) are equivalent.

Proof. For each $n \geq 0$ set $R_n = (1/2)[(A_1A_2)^n + (A_2A_1)^n]$. It is straightforward to check that

$$\begin{aligned} R_0 &= I, \\ R_1 &= (1/2)(A_1A_2 + A_2A_1), \quad \text{and} \\ R_n &= 2R_1R_{n-1} - R_{n-2} \quad \text{for } n \geq 2. \end{aligned}$$

It follows immediately by induction that for each $n \geq 0$ we have

$$R_n = T_n(R_1),$$

where $T_n(z)$ are Tchebyshev's polynomials of the first kind defined by

$$\begin{aligned} T_0(z) &= 1, \\ T_1(z) &= z, \quad \text{and} \\ T_n(z) &= 2zT_{n-1}(z) - T_{n-2}(z) \quad \text{for } n \geq 2. \end{aligned}$$

It is well known that for each real $z \in [-1, 1]$ one has $T_n(z) = \cos(n \cos^{-1}(z))$, cf. Rivlin [33], in particular the polynomial $T_n(z) - 1$ is of degree n and has for its set of roots the set $\{\cos(2\pi k/n) \mid k = 0, \dots, n-1\}$.

Now, suppose $(A_1A_2)^m = I$. Thus $(A_2A_1)^m = I$ as well, hence $R_m = T_m(R_1) = I$. Since by Proposition 3.1(a) we have $\sigma(R_m) = T_m(\sigma(R_1))$, we must have $T_m(\alpha) = 1$ for each $\alpha \in \sigma(R_1)$. Therefore $\sigma(R_1) \subseteq \{\cos(2\pi k/m) \mid k = 0, \dots, m-1\}$, which implies (2) as desired.

Next, suppose A_1 and A_2 are both unitary and (2) holds. Then R_n is self-adjoint for each n . In particular R_m is self-adjoint and has $\sigma(R_m) = T_m(\sigma(R_1)) = \{1\}$. It follows from Proposition 3.1(b) that $R_m = I$. Thus $(A_1A_2)^m + (A_1A_2)^{-m} = 2I$ hence $(A_1A_2)^{2m} - 2(A_1A_2)^m + I = 0$. Proposition 3.1(a) now yields that $(\lambda-1)^2 = 0$ for every $\lambda \in \sigma((A_1A_2)^m)$. So $(A_1A_2)^m$ is a unitary hence normal operator whose spectrum is the singleton $\{1\}$, and therefore (1) holds by Proposition 3.1(b). \square

Remark 4.4. The above proofs show that the spectral decomposition of the operator $A_1A_2 + A_2A_1$ determines the decomposition of the pair (A_1, A_2) and the decomposition of the joint spectrum of A_1 and A_2 into irreducible components. The fact that $A_1A_2 + A_2A_1$ commutes with both A_1 and A_2 , and, therefore, belongs to the center of the group-algebra, plays a key role here.

Now we are ready to present the proof of Theorem 1.1.

Proof of Theorem 1.1. We have $T = \{1_G, g_1, \dots, g_n\}$ where $\{g_1, \dots, g_n\}$ is a system of Coxeter generators of for G . Also, we have $T' = \{1_{G'}, g'_1, \dots, g'_n\}$ where $\{g'_1, \dots, g'_n\}$ is a generating set for the group G' . Let A_i (resp. A'_i) be the operator on $\mathbb{C}[G]^\vee$ (resp. $\mathbb{C}[G']^\vee$) induced via left multiplication by g_i (resp. g'_i) on the group ring $\mathbb{C}[G]$ (resp. $\mathbb{C}[G']$). Suppose $D(T) \supseteq D(T')$. We also have $\sigma_p(A_1, \dots, A_n) \supseteq \sigma_p(A'_1, \dots, A'_n)$. Furthermore, for all i and all $k < l$ intersecting with the projective line $\{x_j = 0 \mid 0 \neq j \neq i\}$, and with the projective plane $\{x_j = 0 \mid j \notin \{0, k, l\}\}$ shows that $\sigma(-I, A_i) \supseteq \sigma(-I, A'_i)$ and $\sigma(-I, A_i, A_j) \supseteq \sigma(-I, A'_i, A'_j)$, hence also that $\sigma_p(A_i, A_j) \supseteq \sigma_p(A'_i, A'_j)$. In particular, for each i we have $\{-1, 1\} = \sigma(A_i) \supseteq \sigma(A'_i)$. Since left regular representations are unitary, A'_i is normal and hence an involution by Proposition 3.1.

Next, suppose that the Coxeter exponent m_{ij} is finite. Then by Lemma 4.3 the spectrum of $A_iA_j + A_jA_i$ is finite and contained in the set $\{2 \cos(2\pi k/m_{ij}) \mid k =$

$0, \dots, m_{ij} - 1\}$. Hence by Lemma 4.1 it is completely determined by $\sigma_p(A_i, A_j)$, which is a finite union of ellipses. Since $\sigma_p(A'_i, A'_j)$ is contained inside $\sigma_p(A_i, A_j)$, again Lemma 4.1 tells us that the spectrum of $A'_i A'_j + A'_j A'_i$ is finite and contained in the spectrum of $A_i A_j + A_j A_i$. Therefore by Lemma 4.3 we obtain $(A'_1 A'_2)^{m_{ij}} = I$. Since the left regular representation is faithful, the generators g'_i of G' satisfy the defining relations on the generators g_i of G .

Now suppose G is finite and $D(T) = D(T')$ as subschemes of $\mathbb{C}\mathbb{P}^n$. But then $|G| = \deg D(T) = \deg D(T') = |G'|$, hence the epimorphism f has to be an isomorphism. \square

5. REPRESENTATIONS OF DIHEDRAL GROUPS

We investigate finite dimensional unitary representations of dihedral groups.

The following result was for us the first indication that joint spectra can be employed to study Coxeter groups. Observe how the proof uses the equation of the circle to explicitly construct the desired invariant subspace.

Theorem 5.1. *Let A_1, A_2 be self-adjoint linear operators on an N -dimensional Hilbert space V , and suppose that A_1 is invertible and that $\|A_2\| = 1$. Further suppose that the “complex unit circle” $\{(x, y) \in \mathbb{C}^2 : x^2 + y^2 = 1\}$ is a reduced component of both $\sigma_p(A_1, A_2)$ and $\sigma_p(A_1^{-1}, A_2)$, of multiplicity n in $\sigma_p(A_1, A_2)$, that the points $(\pm 1, 0)$ do not belong to any other component of either $\sigma_p(A_1, A_2)$ or $\sigma_p(A_1^{-1}, A_2)$, and that the points $(0, \pm 1)$ do not belong to any other component of $\sigma_p(A_1, A_2)$. Then:*

- (1) A_1 and A_2 have a common $2n$ -dimensional invariant subspace L ;
- (2) The pair of restrictions $A_1|_L$ and $A_2|_L$ is unitary equivalent to the following pair of $2n \times 2n$ involutions C_1 and C_2 , each block-diagonal with n equal 2×2 blocks along the diagonal:

$$C_1 = \begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & -1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & -1 \end{bmatrix}, \quad C_2 = \begin{bmatrix} 0 & 1 & \dots & 0 & 0 \\ 1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & \dots & 1 & 0 \end{bmatrix}.$$

- (3) The group generated by C_1 and C_2 represents the Coxeter group B_2 .

Proof. Since the coordinate axes are not tangent to the unit circle at any of the points $(\pm 1, 0)$ and $(0, \pm 1)$, and these are regular points of the unit circle, Proposition 3.2 shows that both 1 and -1 are eigenvalues of A_1 and of A_2 with multiplicity n . Let e_1, \dots, e_n and e_{n+1}, \dots, e_{2n} be a pair of orthonormal eigenbases for the eigenspaces of A_1 with eigenvalues 1 and -1 respectively, and ξ_1, \dots, ξ_n and $\xi_{n+1}, \dots, \xi_{2n}$ be a similar pair of orthonormal eigenbases for A_2 . The spectral decomposition of A_1 in our case looks like

$$A_1 = P_1 - P_2 + \sum_{j=3}^k \lambda_j P_j,$$

where P_1 and P_2 are the orthogonal projections on the spaces $\text{span}\{e_1, \dots, e_n\}$ and $\text{span}\{e_{n+1}, \dots, e_{2n}\}$ respectively, λ_j for $j = 3, \dots, k$ are all other eigenvalues of A_1 ,

and P_j is the orthogonal projection on the λ_j -eigensubspace of A_1 . In our situation equations (3.9) and (3.10) applied to the projection P_1 give

$$(5.2) \quad P_1 A_2 P_1 = 0$$

$$(5.3) \quad P_1 A_2 T A_2 P_1 = -\frac{1}{2} P_1,$$

where T is given by (3.7), which applied to our case turns into

$$T = -\frac{1}{2} P_2 + \sum_{j=3}^k \frac{1}{\lambda_j - 1} P_j.$$

Equations (5.2) and (5.3) imply that for every $j = 1, \dots, n$ we have

$$\begin{aligned} P_1 A_2 T A_2 e_j &= -\frac{1}{2} e_j, \\ -P_1 A_2 \left(\frac{1}{2} P_2 A_2 e_j + \sum_{s=3}^k \frac{1}{1 - \lambda_s} P_s A_2 e_j \right) &= -\frac{1}{2} e_j, \\ P_1 \left(\frac{1}{2} A_2 P_2 A_2 e_j + \sum_{s=3}^k \frac{1}{1 - \lambda_s} A_2 P_s A_2 e_j \right) &= \frac{1}{2} e_j, \\ \sum_{l=1}^n \left[\frac{1}{2} \langle A_2 P_2 A_2 e_j, e_l \rangle + \sum_{s=3}^k \frac{1}{1 - \lambda_s} \langle A_2 P_s A_2 e_j, e_l \rangle \right] e_l &= \frac{1}{2} e_j, \\ \sum_{l=1}^n \left[\frac{1}{2} \langle P_2 A_2 e_j, A_2 e_l \rangle + \sum_{s=3}^k \frac{1}{1 - \lambda_s} \langle P_s A_2 e_j, A_2 e_l \rangle \right] e_l &= \frac{1}{2} e_j, \\ \sum_{l=1}^n \left[\frac{1}{2} \langle P_2 A_2 e_j, P_2 A_2 e_l \rangle + \sum_{s=3}^k \frac{1}{1 - \lambda_s} \langle P_s A_2 e_j, P_s A_2 e_l \rangle \right] e_l &= \frac{1}{2} e_j, \end{aligned}$$

The last relation implies

$$(5.4) \quad \frac{1}{2} \langle P_2 A_2 e_j, P_2 A_2 e_l \rangle + \sum_{s=3}^k \frac{1}{1 - \lambda_s} \langle P_s A_2 e_j, P_s A_2 e_l \rangle = \frac{1}{2} \delta_j^l$$

for all $1 \leq j, l \leq n$, where δ_j^l is the Kronecker symbol. Putting $l = j$ in (5.4) yields

$$(5.5) \quad \frac{1}{2} |P_2 A_2 e_j|^2 + \sum_{s=3}^k \frac{|P_s(A_2 e_j)|^2}{1 - \lambda_s} = \frac{1}{2}, \quad 1 \leq j \leq n.$$

Since the spectral resolution for A_1^{-1} is

$$A_1^{-1} = P_1 - P_2 + \sum_{j=3}^k \frac{1}{\lambda_j} P_j,$$

the operator $T(A_1^{-1})$ is given by

$$T(A_1^{-1}) = -\frac{1}{2} P_2 + \sum_{j=3}^k \frac{\lambda_j}{1 - \lambda_j} P_j.$$

Now, equation (3.10) applied to $\sigma_p(A_1^{-1}, A_2)$ yields

$$(5.6) \quad P_1 A_2 T(A_1^{-1}) A_2 P_1 = -\frac{1}{2} P_1,$$

A similar argument applied to (5.6) gives

$$(5.7) \quad \frac{1}{2} |P_2 A_2 e_j|^2 - \sum_{s=3}^k \frac{\lambda_s |P_s A_2 e_j|^2}{1 - \lambda_s} = \frac{1}{2}.$$

Adding (5.7) and (5.5) we obtain

$$(5.8) \quad \sum_{s=2}^k |P_s A_2 e_j|^2 = 1$$

Now, since (5.2) means $P_1(A_2 e_j) = 0$, the displayed equation above yields

$$(5.9) \quad |A_2 e_j| = 1.$$

Since $\|A_2\| = 1$, it follows from (5.9) that for every $1 \leq j \leq n$ we have $e_j \in \text{span}\{\xi_1, \dots, \xi_{2n}\}$. A similar analysis shows that for each $n+1 \leq j \leq 2n$ we have $e_j \in \text{span}\{\xi_1, \dots, \xi_{2n}\}$. Thus,

$$L = \text{span}\{\xi_1, \dots, \xi_{2n}\} = \text{span}\{e_1, \dots, e_{2n}\}$$

is a common invariant subspace of A_1 and A_2 , which completes the proof of (1).

Next, we remark that the restrictions of A_1 and A_2 to L are involutions, and, since $P_1 A_2 e_j = 0$ for $j = 1, \dots, n$ and $P_2 A_2 e_j = 0$ for $j = n+1, \dots, 2n$, that

$$(5.10) \quad A_2 e_j \in \text{span}\{e_{n+1}, \dots, e_{2n}\}, \text{ for } j = 1, \dots, n;$$

$$(5.11) \quad A_2 e_j \in \text{span}\{e_1, \dots, e_n\}, \text{ for } j = n+1, \dots, 2n.$$

Therefore, in the basis e_1, \dots, e_{2n} the restrictions of A_1 and A_2 to L look like

$$(5.12) \quad A_1|_L = \begin{bmatrix} I_n & 0_n \\ 0_n & -I_n \end{bmatrix}, \quad A_2|_L = \begin{bmatrix} 0_n & D_n \\ D_n^* & 0_n \end{bmatrix},$$

where I_n and 0_n are the identity and zero $n \times n$ matrices respectively, and D_n is a unitary $n \times n$ matrix. Write

$$U = \begin{bmatrix} D_n^* & 0_n \\ 0_n & I_n \end{bmatrix},$$

then U is unitary and

$$(5.13) \quad U \left(\begin{bmatrix} I_n & 0_n \\ 0_n & -I_n \end{bmatrix}, \begin{bmatrix} 0_n & D_n \\ D_n^* & 0_n \end{bmatrix} \right) U^* = \left(\begin{bmatrix} I_n & 0_n \\ 0_n & -I_n \end{bmatrix}, \begin{bmatrix} 0_n & I_n \\ I_n & 0_n \end{bmatrix} \right).$$

Now, the interchange of coordinate vectors $e_{2j} \leftrightarrow e_{2j-1+n}$ for $j = 1, \dots, \lfloor \frac{n}{2} \rfloor$ finishes the proof of (2).

To prove (3) note that the relations $C_1^2 = C_2^2 = (C_1 C_2)^4 = 1$ are straightforward to verify. \square

We now prove a result that builds upon Lemma 4.1.

Theorem 5.14. *Let A_1 and A_2 be unitary self-adjoint linear operators on a finite-dimensional Hilbert space V . Then:*

- (1) *Every reduced component of $\sigma_p(A_1, A_2)$ is either a line $\{x \pm y = \pm 1\}$ or an "ellipse" $\{x^2 + 2xy \cos(2\pi\theta) + y^2 = 1\}$ for some $0 < \theta < 1/2$.*

- (2) If a line $\{x \pm y = \pm 1\}$ is a reduced component of multiplicity r of the joint spectrum $\sigma_p(A_1, A_2)$ then A_1 and A_2 have a corresponding common eigenspace of dimension r .
- (3) If an “ellipse” $\{x^2 + 2xy \cos(2\pi\theta) + y^2 = 1\}$ with $0 < \theta < 1/2$ is a reduced component of the proper joint spectrum $\sigma_p(A_1, A_2)$ of multiplicity r , then A_1 and A_2 have a corresponding common invariant subspace of dimension $2r$ that is a direct sum of r two-dimensional common invariant subspaces.

Proof. Since A_1 and A_2 are unitary involutions, the operator $R = A_1A_2 + A_2A_1$ is self-adjoint of norm ≤ 2 . Part (1) now is an immediate consequence of Lemma 4.1.

We give the proof of (2) for the case when our line is $\{x + y = 1\}$. The proof of the other cases is analogous.

Choose a point $(\gamma, \beta) \in \{x + y = 1\}$ with positive real coordinates. The spectrum of the operator $\gamma A_1 + \beta A_2$ is the set $\{\lambda \in \mathbb{R} : (1/\lambda)(\gamma, \beta) \in \sigma_p(A_1, A_2)\}$. Since by part (1) all nonlinear reduced components of $\sigma_p(A_1, A_2)$ are convex inside the positive real quadrant and intersect $\{x + y = 1\}$ at $(1, 0)$ and $(0, 1)$, we have that $\max\{|\lambda| : (1/\lambda)(\gamma, \beta) \in \sigma_p(A_1, A_2)\}$ corresponds to the point of the intersection of the lines $\{x + y = 1\}$ and $\{\beta x - \gamma y = 0\}$. Thus $\max\{|\lambda| : (1/\lambda)(\gamma, \beta) \in \sigma_p(A_1, A_2)\} = 1/(\gamma + \beta) = 1$. Since the operator $\gamma A_1 + \beta A_2$ is self-adjoint, this implies $\|\gamma A_1 + \beta A_2\| = 1$. Also, this point of intersection is a regular point of $\sigma_p(A_1, A_2)$. By (3.4) the joint spectrum of $\gamma A_1 + \beta A_2$ and A_2 also contains the line $\{x + y = 1\}$ (with the same multiplicity r), and $(1, 0)$ is a regular point of $\sigma_p(\gamma A_1 + \beta A_2, A_2)$. Since $\|\gamma A_1 + \beta A_2\| = \|A_2\| = 1$, by [5, Proofs of Lemma 5 and Lemma 9] we have that A_2 and $\gamma A_1 + \beta A_2$ have a common eigensubspace corresponding to the eigenvalue 1. By Proposition 3.2 this eigenspace has dimension equal to the multiplicity of the line $\{x + y = 1\}$ in $\sigma_p(\gamma A_1 + \beta A_2, A_2)$, hence is r -dimensional. Clearly it is an eigensubspace of eigenvalue 1 for A_1 as well.

Now we proceed with the proof of (3). In view of part (2), by restricting to orthogonal complements if necessary, we may assume without loss of generality that $\sigma_p(A_1, A_2)$ contains none of the lines $\{x \pm y = 1\}$ or $\{x \pm y = -1\}$. By part (1) and its proof, all reduced components of the proper joint spectrum are the nonsingular “ellipses”

$$(5.15) \quad \mathcal{E}_i = \{x^2 + 2xy \cos(2\pi\theta_i) + y^2 = 1\}, \quad \text{with } 1/2 > \theta_1 > \dots > \theta_s > 0,$$

their intersection points are $(\pm 1, 0)$ and $(0, \pm 1)$, and $\alpha_i = 2 \cos(2\pi\theta_i)$ is an eigenvalue of $R = A_1A_2 + A_2A_1$ for each i . Note that if M_i is the corresponding eigenspace of R , then for every $(x, y) \in \mathbb{R}^2$ on the ellipse \mathcal{E}_i the space M_i is invariant under the action of $xA_1 + yA_2$. Indeed, let $\xi \in M_i$. Then

$$(xA_1 + yA_2)^2 \xi = (x^2 + y^2)\xi + xyR_1 \xi = \xi.$$

Thus, the restriction of $(xA_1 + yA_2)^2$ to M_i is the identity of M_i . Since the eigenvalues of $(xA_1 + yA_2)^2$ are just the squares of the eigenvalues of $xA_1 + yA_2$ with the same eigenvectors, every eigenvector of $(xA_1 + yA_2)^2$ with eigenvalue 1 is a linear combination of eigenvectors of $(xA_1 + yA_2)$ with eigenvalues ± 1 . Conversely, the displayed above equality shows also that when $xy \neq 0$ each eigenvector ξ of $(xA_1 + yA_2)$ with eigenvalue ± 1 , is an eigenvector of R with eigenvalue α_i . Thus, when $xy \neq 0$ the eigenspace M_i is the direct sum of the eigenspaces of $(xA_1 + yA_2)$ with eigenvalues ± 1 , and, therefore, is invariant under $(xA_1 + yA_2)$. In particular, since when $xy \neq 0$ the point (x, y) belongs to only one reduced component (of multiplicity n_i) of $\sigma_p(A_1, A_2)$, is a nonsingular point on that component, and no lines

through the origin are tangent to any of the reduced components of $\sigma_p(A_1, A_2)$, by Proposition 3.2 the eigenspace of $xA_1 + yA_2$ of eigenvalue 1 has dimension n_i . Since our components are invariant under the transformation $(x, y) \mapsto (-x, -y)$, when $xy \neq 0$ the eigenspace of $xA_1 + yA_2$ of eigenvalue -1 is also of dimension n_i ; hence the dimension of M_i is $2n_i$.

Next, the projective joint spectrum $\sigma(I, A_1, A_2)$ is the union of the homogenizations of the “ellipses” (5.15), with the same multiplicities. Furthermore, since there are no real solutions to the equation $x^2 + 2xy \cos(2\pi\theta_i) + y^2 = 0$, there are no real points of the form $[0 : x : y]$ in $\sigma(I, A_1, A_2)$, i.e. no real points on the “infinite” line $\{x_0 = 0\}$; and hence 0 is not in the spectrum of $(xA_1 + yA_2)$. Therefore for every $(x, y) \in \mathbb{R}^2$ with $(x, y) \neq (0, 0)$ the operator $(xA_1 + yA_2)$ is invertible. Also, if $(x, y) \in \mathbb{R}^2$ belongs to \mathcal{E}_i , then

$$(5.16) \quad (xA_1 + yA_2)^{-1}|_{M_i} = (xA_1 + yA_2)|_{M_i}.$$

Indeed, let $\xi \in M_i$. We have

$$(xA_1 + yA_2)^2\xi = (x^2 + xy\alpha_i + y^2)\xi.$$

Thus, $(xA_1 + yA_2)^{-1}$ is defined by

$$(5.17) \quad (xA_1 + yA_2)^{-1}|_{M_i} = \frac{1}{x^2 + xy\alpha_i + y^2}(xA_1 + yA_2)|_{M_i},$$

which, of course, implies (5.16).

Next we note that

$$x^2 + 2xy \cos(2\pi\theta_i) + y^2 = \frac{1 + \cos(2\pi\theta_i)}{2}(x + y)^2 + \frac{1 - \cos(2\pi\theta_i)}{2}(x - y)^2,$$

and, therefore, each “ellipse” (5.15) is centered at the origin, has axes along lines $\{x = y\}$ and $\{x = -y\}$, the lengths of its semiaxes $\sqrt{\frac{1}{1 + \cos(2\pi\theta_i)}}$ and $\sqrt{\frac{1}{1 - \cos(2\pi\theta_i)}}$ respectively, and recall that we write n_i for its multiplicity.

Set

$$B_1 = \frac{1}{\sqrt{2(1 + \cos(2\pi\theta_1))}}(A_1 + A_2)$$

$$B_2 = \frac{1}{\sqrt{2(1 - \cos(2\pi\theta_1))}}(A_1 - A_2).$$

It is easy to check that the joint spectrum of B_1 and B_2 is the union of the “ellipses”

$$(5.18) \quad \mathcal{E}'_i = \left\{ \frac{1 + \cos(2\pi\theta_i)}{1 + \cos(2\pi\theta_1)}x^2 + \frac{1 - \cos(2\pi\theta_i)}{1 - \cos(2\pi\theta_1)}y^2 = 1 \right\}$$

for $i = 1, \dots, s$, each with multiplicity n_i . In particular, this implies that the unit circle $\{x^2 + y^2 = 1\}$ is in $\sigma_p(B_1, B_2)$, has multiplicity n_1 , and the spectrum of B_2 consists of numbers $\{\pm\sqrt{\frac{1 - \cos(2\pi\theta_i)}{1 - \cos(2\pi\theta_1)}}\}$. Hence, the norm of B_2 is equal to one. It also follows from (5.18) that the points $(\pm 1, 0)$ and $(0, \pm 1)$ are regular points of $\sigma_p(B_1, B_2)$, all of them having multiplicity n_1 .

Finally, if $(x, y) \in \mathbb{R}^2$ belongs to \mathcal{E}'_i , then

$$\begin{aligned} xB_1 + yB_2 &= \left(\frac{1}{\sqrt{2(1 + \cos(2\pi\theta_1))}}x + \frac{1}{\sqrt{2(1 - \cos(2\pi\theta_1))}}y \right) A_1 \\ &\quad + \left(\frac{1}{\sqrt{2(1 + \cos(2\pi\theta_1))}}x - \frac{1}{\sqrt{2(1 - \cos(2\pi\theta_1))}}y \right) A_2 = u_1A_1 + u_2A_2. \end{aligned}$$

It is easy to check that

$$u_1^2 + 2u_1u_2 \cos(2\pi\theta_i) + u_2^2 = 1$$

which, of course, means that $(u_1, u_2) \in \mathcal{E}_i$. According to (5.16)

$$\begin{aligned} (xB_1 + yB_2)^{-1}|_{M_i} &= (u_1A_1 + u_2A_2)^{-1}|_{M_i} \\ &= (u_1A_1 + u_2A_2)|_{M_i} = (xB_1 + yB_2)|_{M_i}. \end{aligned}$$

Let $\lambda_i = \sqrt{\frac{1+\cos(2\pi\theta_1)}{1+\cos(2\pi\theta_i)}}$. It follows that $(\lambda_i B_1)^{-1}|_{M_i} = \lambda_i B_1|_{M_i}$, hence

$$\begin{aligned} (xB_1^{-1} + yB_2)|_{M_i} &= xB_1^{-1}|_{M_i} + yB_2|_{M_i} \\ &= x\lambda_i^2 B_1|_{M_1} + yB_2|_{M_i} = (x\lambda_i^2 B_1 + yB_2)|_{M_i}. \end{aligned}$$

Therefore, except for finitely many points, when $(\lambda_i^2 x, y)$ is on \mathcal{E}'_i the space M_i is the direct sum of eigenspaces of eigenvalues ± 1 for the operator $(xB_1^{-1} + yB_2)$. Thus each ellipse $\mathcal{E}''_i = \left\{ \frac{1+\cos(2\pi\theta_1)}{1+\cos(2\pi\theta_i)}x^2 + \frac{1-\cos(2\pi\theta_1)}{1-\cos(2\pi\theta_i)}y^2 = 1 \right\}$ is a reduced component of $\sigma_p(B_1^{-1}, B_2)$, and by Proposition 3.2 it has multiplicity $\geq n_i$. Since the sum of the multiplicities equals $(1/2) \dim V$, we must have that the multiplicity of \mathcal{E}''_i equals n_i for each i , and therefore these are all reduced components of $\sigma_p(B_1^{-1}, B_2)$. In particular, the “unit circle” $\{x^2 + y^2 = 1\}$ is a reduced component of $\sigma_p(B_1^{-1}, B_2)$, of multiplicity n_1 , and the points $(\pm 1, 0)$ do not belong to any other of its components. It follows from Theorem 5.1 that M_1 is a direct sum of n_1 two-dimensional common invariant subspaces for B_1 and B_2 and the restriction of the pair B_1, B_2 to each of these subspaces is unitary equivalent to the pair

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Of course, each of these subspaces is also invariant under A_1 and A_2 , and the restrictions on these subspaces of the pair A_1, A_2 is unitary equivalent to

$$\begin{bmatrix} \cos(\pi\theta_1) & \sin(\pi\theta_1) \\ \sin(\pi\theta_1) & -\cos(\pi\theta_1) \end{bmatrix}, \begin{bmatrix} \cos(\pi\theta_1) & -\sin(\pi\theta_1) \\ -\sin(\pi\theta_1) & -\cos(\pi\theta_1) \end{bmatrix}.$$

Denote by \mathcal{N}_1 the orthocomplement to M_1 . Then \mathcal{N}_1 is invariant under A_1 and A_2 and the proper joint spectrum $\sigma_p(A_1|_{\mathcal{N}_1}, A_2|_{\mathcal{N}_1})$ is the union of the ellipses \mathcal{E}_j with $j = 2, \dots, s$, each with multiplicity n_j . A similar argument applied to the subspace M_2 of \mathcal{N}_1 shows that M_2 is a direct sum of n_2 two-dimensional common invariant subspaces for A_1 and A_2 , and that their restrictions to those subspaces are unitary equivalent to

$$(5.19) \quad \begin{bmatrix} \cos(\pi\theta_2) & \sin(\pi\theta_2) \\ \sin(\pi\theta_2) & -\cos(\pi\theta_2) \end{bmatrix}, \begin{bmatrix} \cos(\pi\theta_2) & -\sin(\pi\theta_2) \\ -\sin(\pi\theta_2) & -\cos(\pi\theta_2) \end{bmatrix}.$$

Since $\theta = \theta_j$ for some j , iterating the above procedure j times yields the claimed result. \square

As an immediate consequence of Theorem 5.14 and its proof we have:

Theorem 5.20. *Let G be a Coxeter group of type $I(n)$ for some $2 \leq n \leq \infty$, let $\{g_1, g_2\}$ be a set of Coxeter generators, and let $\rho : G \rightarrow GL(V)$ be a finite dimensional unitary representation of G .*

- (1) *If $\rho' : G \rightarrow GL(V')$ is another finite dimensional unitary representation of G such that $\sigma_p(\rho(g_1), \rho(g_2)) = \sigma_p(\rho'(g_1), \rho'(g_2))$ as subschemes of \mathbb{C}^2 , then ρ and ρ' are equivalent.*
- (2) *If ρ is irreducible then $\sigma_p(\rho(g_1), \rho(g_2))$ is reduced and irreducible.* \square

6. ADMISSIBLE TRANSFORMATIONS

For the rest of this paper $n \geq 2$, the group G is a finite Coxeter group of type A_n , B_n , or D_{n+1} , and $\{g_1, g_2, \dots\}$ is a set of Coxeter generators for G .

For a word $w = g_{i_1} \dots g_{i_N}$, where the same letter g_k might occur more than once, we write $|w| = N$. Let a_k be the number of times g_k occurs in w . The *signature* of w is the sequence

$$\text{sig}(w) = (a_1, a_2, \dots),$$

and the *content* of w is the sequence

$$\text{ct}(w) = (|w|, a_1, \dots, a_{n-1}) \in \mathbb{N}^n.$$

Since $|w| = a_1 + a_2 + \dots$, in types A_n and B_n the content and the signature of w carry the same information, while in type D_{n+1} it is clearly possible to have words with the same content but different signatures. Let \leq_{lex} be the lexicographic ordering on \mathbb{N}^n , that is, $c = (c_0, \dots, c_{n-1}) <_{lex} (d_0, \dots, d_{n-1}) = d$ precisely when $c \neq d$ and $c_m < d_m$ for $m = \min\{i \mid c_i \neq d_i\}$. We always consider our words partially ordered $<$ by

$$w_1 < w_2 \quad \text{if and only if} \quad \text{ct}(w_1) <_{lex} \text{ct}(w_2).$$

For example, $g_2 g_1 g_2 < g_1 g_2 g_1$ whenever $n \geq 2$.

Definition 6.1. (a) We introduce *admissible transformations* on words. They are:

- (1) Cancelling transformations. That is, for each i the transformation

$$w' g_i g_i w'' \mapsto w' w''$$

is admissible.

- (2) Commuting transformations. This means that if g_i and g_j commute, then the transformation

$$w' g_i g_j w'' \mapsto w' g_j g_i w''$$

is admissible.

- (3) Circular transformations. Those are:

$$g_{i_1} \dots g_{i_k} g_{i_{k+1}} \dots g_{i_N} \mapsto g_{i_{k+1}} \dots g_{i_N} g_{i_1} \dots g_{i_k}.$$

- (4) Replacement transformations. These replace a certain subword consisting of a successive string of letters by another representation of this subword. Specifically,

$$\begin{aligned} w'g_i g_{i+1} g_i w'' &\mapsto w'g_{i+1} g_i g_{i+1} w'' && \text{for } 1 \leq i \leq n-2; \\ w'g_{n-1} g_n g_{n-1} w'' &\mapsto w'g_n g_{n-1} g_n w'' && \text{in } A_n \text{ and } D_{n+1}; \\ w'g_{n-1} g_{n+1} g_{n-1} w'' &\mapsto w'g_{n+1} g_{n-1} g_{n+1} w'' && \text{in } D_{n+1}; \end{aligned}$$

and in addition the “tent commuting” replacements

$$\begin{aligned} w't_k g_j w'' &\mapsto w'g_j t_k w'' && \text{for } k < j \leq n-1 \text{ in } B_n \text{ and } D_{n+1}; \\ w't_k g_n w'' &\mapsto w'g_{n+1} t_k w'' && \text{and} \\ w't_k g_{n+1} w'' &\mapsto w'g_n t_k w'' && \text{for } k \leq n-1 \text{ in } D_{n+1}; \\ w't_k g_n w'' &\mapsto w'g_n t_k w'' && \text{for } k \leq n-1 \text{ in } B_n \end{aligned}$$

are admissible transformations. Here the k th “tent word” t_k is defined for $1 \leq k \leq n-1$ as

$$t_k = \begin{cases} g_k \cdots g_{n-1} g_n g_{n-1} \cdots g_k & \text{in } B_n; \\ g_k \cdots g_{n-1} g_n g_{n+1} g_{n-1} \cdots g_k & \text{in } D_{n+1}. \end{cases}$$

(b) We write $w_1 \rightsquigarrow w_2$ if there is a sequence of admissible transformations that maps w_1 into w_2 . Taking the equivalence closure of this transitive relation on words, we obtain the notion of *c-equivalence*. We write $w_1 \sim w_2$ if the words w_1 and w_2 are *c-equivalent*.

Remark 6.2. (a) Note that in B_n and D_{n+1} we have the following “tent commuting” equalities of elements. First,

$$\begin{aligned} t_k g_j &= u g_{j-1} g_j t_{j+1} g_j g_{j-1} (v g_j) \\ &= u g_{j-1} g_j t_{j+1} (g_j g_{j-1} g_j) v \\ &= u g_{j-1} g_j (t_{j+1} g_{j-1}) g_j g_{j-1} v \\ &= u (g_{j-1} g_j g_{j-1}) t_{j+1} g_j g_{j-1} v \\ &= (u g_j) g_{j-1} g_j t_{j+1} g_j g_{j-1} v \\ &= g_j (u g_{j-1} g_j t_{j+1} g_j g_{j-1} v) \\ &= g_j t_k \end{aligned}$$

whenever $k < j \leq n-1$, as u and v only involve letters g_i with $i \leq j-2$ hence they commute with g_j , while t_{j+1} only involves g_i s with $i \geq j+1$, hence it commutes with g_{j-1} . For trivial reasons, the same equality also holds when $j+1 < k \leq n-1$.

In a similar manner, when $k \leq n - 1$ we have in D_{n+1} the equalities

$$\begin{aligned}
t_k g_n &= u g_{n-1} (g_n g_{n+1}) g_{n-1} (v g_n) \\
&= u g_{n-1} g_{n+1} (g_n g_{n-1} g_n) v \\
&= u (g_{n-1} g_{n+1} g_{n-1}) g_n g_{n-1} v \\
&= (u g_{n+1}) g_{n-1} (g_{n+1} g_n) g_{n-1} v \\
&= g_{n+1} (u g_{n-1} g_n g_{n+1} g_{n-1} v) \\
&= g_{n+1} t_k
\end{aligned}$$

and $t_k g_{n+1} = g_n t_k$. Finally, in B_n we have the equalities

$$\begin{aligned}
t_k g_n &= u g_{n-1} g_n g_{n-1} (v g_n) \\
&= u (g_{n-1} g_n g_{n-1} g_n) v \\
&= (u g_n) g_{n-1} g_n g_{n-1} v \\
&= g_n (u g_{n-1} g_n g_{n-1} v) \\
&= g_n t_k
\end{aligned}$$

for all $k \leq n - 1$.

(b) Combining the above yields in B_n and D_{n+1} also the tent commuting equalities

$$t_k t_j = t_j t_k$$

whenever $1 \leq k, j \leq n - 1$.

(c) It is immediate from (a) and the defining relations of G that if $w_1 \rightsquigarrow w_2$ and no circular transformations were used, then w_1 and w_2 represent the same element of the group G .

(d) It is also clear from (a) that if w_1 is c -equivalent to w_2 then these words represent elements of G that belong to the same conjugacy class. The converse to this is also true, but since we will not need it here, we leave the easy though somewhat cumbersome proof as an exercise for the interested reader.

We conclude this section with a simple but crucially important observation.

Lemma 6.3. *If $w_1 \rightsquigarrow w_2$ then $\text{ct}(w_1) \geq_{lex} \text{ct}(w_2)$.*

Proof. Indeed every listed admissible transformation either preserves the content of the word, or lowers it. \square

7. ECHELON FORMS

Now we will show that words in our Coxeter group G have conjugates in a special form that we call **echelon form**.

Definition 7.1. We say that a word w is in *echelon form* if it is of the form

$$w = \delta_1 \delta_2 \dots \delta_n \delta_{n+1}$$

where for each i the word δ_i satisfies

$$(7.2) \quad \delta_i \in \begin{cases} \{1, g_i\} & \text{if } i \leq n \text{ and } G = A_n; \\ \{1\} & \text{if } i = n + 1 \text{ and } G = A_n \text{ or } G = B_n; \\ \{1, g_n\} & \text{if } i = n \text{ and } G = B_n; \\ \{1, g_i\} & \text{if } i \geq n \text{ and } G = D_{n+1}; \\ \{1, g_i, t_i\} & \text{if } i \leq n - 1 \text{ and } G = B_n \text{ or } G = D_{n+1}. \end{cases}$$

Remark 7.3. Note that when w is in echelon form, it can be recovered from $\text{ct}(w)$ in types A_n and B_n , while in type D_{n+1} there are at most two possibilities for w given its content. More precisely, in all types, given $\text{ct}(w) = (|w|, a_1, \dots, a_{n-1})$, the tuple (a_1, \dots, a_i) determines the tuple $(\delta_1, \dots, \delta_i)$ whenever $1 \leq i \leq n-1$. In types A_n and B_n the tuple $(\delta_1, \dots, \delta_{n-1})$ together with $|w|$ determines δ_n and hence w . In type D_{n+1} the tuple $(\delta_1, \dots, \delta_{n-1})$ together with $|w|$ leaves at most two possibilities for (δ_n, δ_{n+1}) , depending on the parity of $a_n + a_{n+1}$. If this parity is even, there is no choice. If the parity is odd, then there is a choice between either $(g_n, 1)$ or $(1, g_{n+1})$ for (δ_n, δ_{n+1}) .

We can extend the last remark even further.

Proposition 7.4. *Let w_1 and w_2 be two words in D_{n+1} , both in echelon form, and suppose that $\text{ct}(w_1) = \text{ct}(w_2) = (W, a_1, \dots, a_{n-1})$.*

If $W - a_1 - \dots - a_{n-1} \geq 3$ then w_1 and w_2 belong to the same conjugacy class as elements of G .

Proof. By Remark 7.3 we may assume that $a_n + a_{n+1} = W - a_1 - \dots - a_{n-1}$ is odd and that

$$w_1 = \delta_1 \dots \delta_{n-1} g_n \quad \text{and} \quad w_2 = \delta_1 \dots \delta_{n-1} g_{n+1},$$

where each δ_i satisfies (7.2). Since $a_n + a_{n+1} \geq 3$, we must have $\delta_i = t_i$ for at least one index i . Let k be the biggest such index, and let j be the second biggest (if such exists). Then $\delta_i \in \{1, g_i\}$ for each $j < i < n$ with $i \neq k$, hence the tent commuting equalities from Remark 6.2 yield

$$\begin{aligned} w_1 &= \delta_1 \dots \delta_{j-1} t_j (\delta_{j+1} \dots \delta_{k-1}) t_k (\delta_{k+1} \dots \delta_{n-1}) g_n \\ &= \delta_1 \dots \delta_{j-1} t_j (\delta_{k+1} \dots \delta_{n-1}) (\delta_{j+1} \dots \delta_{k-1}) t_k g_n \\ &= \delta_1 \dots \delta_{j-1} t_j (\delta_{k+1} \dots \delta_{n-1}) (\delta_{j+1} \dots \delta_{k-1}) g_{n+1} t_k \\ &= \delta_1 \dots \delta_{j-1} t_j (\delta_{k+1} \dots \delta_{n-1}) g_{n+1} (\delta_{j+1} \dots \delta_{k-1}) t_k. \end{aligned}$$

Now applying a circular transformation we obtain

$$\sim (\delta_{j+1} \dots \delta_{k-1}) t_k \delta_1 \dots \delta_{j-1} t_j (\delta_{k+1} \dots \delta_{n-1}) g_{n+1}$$

and using again the tent commuting equalities we get

$$= \delta_1 \dots \delta_{j-1} t_j (\delta_{j+1} \dots \delta_{k-1}) t_k (\delta_{k+1} \dots \delta_{n-1}) g_{n+1} = w_2$$

as desired. \square

The following key theorem is the main result of this section.

Theorem 7.5 (Ordering Theorem). *Let w be a word. There exists a word \tilde{w} in echelon form such that $w \rightsquigarrow \tilde{w}$.*

The proof relies on the following lemma.

Lemma 7.6. *Let i be the smallest index of a letter in w . Then $w \rightsquigarrow w' \delta_i w''$ where the words w' and w'' can contain only letters with index $> i$, and δ_i satisfies (7.2). Furthermore, the sequence of admissible transformations can be chosen so that no circular transformations are used.*

Proof. The case when $i \geq n$ is trivial, so we may assume that $i \leq n-1$, hence without loss of generality also that $i = 1$. If $a_1 = 1$ then no transformations are

needed, so we may assume that $a_1 \geq 2$. Thus we have $w = ug_1vg_1s$ where u and v do not involve g_1 .

Consider now the case $n = 2$. We prove this case by induction on a_1 . Suppose first that $a_1 = 2$. Then s does not involve g_1 . Since v can involve only g_2 and/or g_3 , a sequence of commuting and/or cancelling transformations yields $w \rightsquigarrow ug_1g_2g_1s$ or, in case of D_3 , either $w \rightsquigarrow ug_1g_2g_3g_1s$ or $w \rightsquigarrow ug_1g_3g_1s$. When $G = A_2$ or $G = D_3$ a replacement transformation gives $ug_1g_2g_1s \rightsquigarrow ug_2g_1g_2s$ which is in the desired form. When $G = B_2$ we already have $ug_1g_2g_1s = ut_1s$ is in the desired form. When $G = D_3$ then $ug_1g_2g_3g_1s = ut_1s$ is also in the desired form, and a replacement transformation yields $ug_1g_3g_1s \rightsquigarrow ug_3g_1g_3s$ which is also in the desired form. Suppose next that $a_1 \geq 3$. Then $s = s'g_1s''$ where s'' does not involve g_1 . By induction we have $ug_1vg_1s' \rightsquigarrow u'\delta_1v'$ where u' and v' do not involve g_1 and no circular transformations were used, hence the same sequence of transformations yields $w \rightsquigarrow u'\delta_1v'g_1s''$. Since v' can involve only g_2 and/or g_3 , a sequence of cancelling and/or commuting transformations yields $w \rightsquigarrow u'\delta_1g_2g_1s''$ or, in case of D_3 , also either $w \rightsquigarrow u'\delta_1g_2g_3g_1s''$ or $w \rightsquigarrow u'\delta_1g_3g_1s''$. In case A_2 we have $\delta_1 = g_1$ hence a replacement transformation brings us to $w \rightsquigarrow u'g_2g_1g_2s''$ which is in the desired form. In case B_2 and $\delta_1 = g_1$ then we are already have $u'g_1g_2g_1s'' = u't_1s''$ which is in the desired form. In case B_2 and $\delta_1 = t_1 = g_1g_2g_1$ then a tent commuting transformation followed by cancelling yields $u't_1g_2g_1s'' \rightsquigarrow u'g_2t_1g_1s'' = u'g_2g_1g_2g_1g_1s'' \rightsquigarrow u'g_2g_1g_2s''$ which is again in the desired form. In the case D_3 and $\delta_1 = g_1$ we handle the possibilities $u'g_1g_2g_1s''$ and $u'g_1g_3g_1s''$ just as in the case A_2 , and the possibility $u'g_1g_2g_3g_1s'' = u't_1s''$ is already in the desired form. Finally, we look at the case D_3 and $\delta_1 = t_1 = g_1g_2g_3g_1$. Then using tent commuting transformations yields $u't_1v'g_1s'' \rightsquigarrow u'v''t_1g_1s''$ where v'' is obtained by exchanging the letters g_2 and g_3 in v' . Then, a cancelling transformation produces $u'v''t_1g_1s'' = u'v''g_1g_2g_3g_1g_1s'' \rightsquigarrow u'v''g_1g_2g_3s''$ which is again in the desired form and completes the proof of the case $n = 2$.

Next, suppose $n \geq 3$. By induction on n we have that $v \rightsquigarrow v'\delta_2v''$ where v' and v'' can only involve letters with index 3 or higher, and no circular transformations are used. Therefore the same sequence of admissible transformations yields $w \rightsquigarrow ug_1v'\delta_2v''g_1t$. Since g_1 commutes with all the letters appearing in v' and v'' , a sequence of commuting transformations yields $w \rightsquigarrow uv'g_1\delta_2g_1v''t$. If $\delta_2 = 1$ then a cancelling transformation yields $w \rightsquigarrow uv'v''t$; if $\delta_2 = g_2$ then a replacement transformation gives $w \rightsquigarrow uv'g_2g_1g_2v''t$; and if $\delta_2 = t_2$ then we have $uv'g_1t_2g_1v''t = uv't_1v''t$. If $a_1 = 2$ then all three cases above are in the desired form. If $a_1 \geq 3$ then the first two cases have a smaller a_1 and in them we are done by induction on a_1 . In the last case, induction on a_1 gives $v''t \rightsquigarrow z'\delta_1z''$ where z' and z'' do not involve g_1 , and no circular transformations are used. Therefore the same sequence of transformations yields $w \rightsquigarrow uv't_1z'\delta_1z''$, hence a sequence of tent commuting replacement transformations produces $w \rightsquigarrow u''t_1\delta_1z''$. Now, if $\delta_1 = g_1$ then $t_1\delta_1 = g_1t_2g_1g_1$ hence a cancelling transformation yields $w \rightsquigarrow u''g_1t_2z''$ which is in the desired form. Finally, if $\delta_1 = t_1$ we get $t_1\delta_1 = g_1t_2g_1g_1t_2g_1$ hence a cancelling transformation yields $w \rightsquigarrow u''g_1t_2t_2g_1z''$ which has smaller a_1 hence we are done by induction on a_1 in this case as well. This completes the proof of the lemma. \square

Proof of Theorem 7.5. Let i_1 be the smallest index of a letter in w . By Lemma 7.6 we get $w \rightsquigarrow w'\delta_{i_1}w''$ hence a circular transformation produces $w \rightsquigarrow \delta_{i_1}w''w' = \delta_1 \dots \delta_{i_1}w_{i_1}$ where w_{i_1} only involves letters with index $> i_1$. Suppose for some

$k \leq n - 1$ we have already obtained $w \rightsquigarrow \delta_1 \dots \delta_k w_k$ where w_k can only involve letters with index $> k$. If $k = n - 1$ then the only possible letters involved in w_{n-1} are g_n and g_{n+1} , so we are done after a sequence of commuting and/or cancelling transformations. If $k < n - 1$ then by Lemma 7.6 we have $w_k \rightsquigarrow w'_k \delta_{k+1} w''_k$ where no circular transformations are used, and w'_k and w''_k can involve only letters with index $> k + 1$. Thus the same sequence of transformations yields $w \rightsquigarrow \delta_1 \dots \delta_k w'_k \delta_{k+1} w''_k$. Therefore a sequence of commuting and/or tent commuting transformations produces $w \rightsquigarrow z \delta_1 \dots \delta_{k+1} w''_k$ where z can involve only letters with index $> k + 1$. Now a circular transformation gives $w \rightsquigarrow \delta_1 \dots \delta_{k+1} w_{k+1}$ where $w_{k+1} = w''_k z$ can involve only letters of index $> k + 1$. Iterating this process we arrive at the desired conclusion. \square

8. THE PROOF OF THEOREM 1.2

We are now ready to present the proof of our second main result.

Proof of Theorem 1.2. By Theorem 2.1 we may assume that both ρ_1 and ρ_2 are unitary. Also, the equality $D(T, \rho_1) = D(T, \rho_2)$ implies $\dim V_1 = \deg D(T, \rho_1) = \deg D(T, \rho_2) = \dim V_2$, and

$$(8.1) \quad \sigma_p(\rho_1(g_1), \dots, \rho_1(g_n)) = \sigma_p(\rho_2(g_1), \dots, \rho_2(g_n)).$$

In particular, the case when G is finite of type $I(n)$ is an immediate consequence from Theorem 5.20.

In the remaining cases for G it suffices by Theorem 2.2 to show that the equality of subschemes $D(T, \rho_1) = D(T, \rho_2)$ implies that the characters χ_{ρ_1} and χ_{ρ_2} are the same.

Let $N = \dim V_1 = \dim V_2$. For each i let $A_i = \rho_1(g_i)$ and let $B_i = \rho_2(g_i)$. For every $(x_1, \dots, x_n) \in \mathbb{C}^n$ the characteristic polynomials for the operators $x_1 A_1 + \dots + x_n A_n$ and $x_1 B_1 + \dots + x_n B_n$ are equal. Therefore the spectra of $x_1 A_1 + \dots + x_n A_n$ and $x_1 B_1 + \dots + x_n B_n$ are the same counting multiplicities. This implies that the traces of these operators are the same, in other words

$$x_1 \operatorname{Tr}(A_1) + \dots + x_n \operatorname{Tr}(A_n) = x_1 \operatorname{Tr}(B_1) + \dots + x_n \operatorname{Tr}(B_n).$$

Since this is true for all x_1, \dots, x_n we obtain

$$(8.2) \quad \operatorname{Tr}(A_j) = \operatorname{Tr}(B_j), \quad j = 1, \dots, n.$$

Similarly, since for each m the eigenvalues of $(x_1 A_1 + \dots + x_n A_n)^m$ are just the m th powers of the eigenvalues of $x_1 A_1 + \dots + x_n A_n$, we get that for every $m \in \mathbb{N}$ the spectra of $(x_1 A_1 + \dots + x_n A_n)^m$ and $(x_1 B_1 + \dots + x_n B_n)^m$ are the same counting multiplicities. Since this is true for arbitrary $(x_1, \dots, x_n) \in \mathbb{C}^n$, we obtain that for every m and $\alpha = (a_1, \dots, a_n) \in \mathbb{N}^n$ with $a_1 + \dots + a_n = m$ one has

$$(8.3) \quad \sum_{\operatorname{sig}(u)=\alpha} \operatorname{Tr}(\rho_1(u)) = \sum_{\operatorname{sig}(u)=\alpha} \operatorname{Tr}(\rho_2(u))$$

To complete the proof it is enough to show by induction on the partial ordering of words that for each word w in G we have

$$\operatorname{Tr}(\rho_1(w)) = \operatorname{Tr}(\rho_2(w)).$$

When $w = 1$ this statement is trivial since both sides equal N . Suppose $w \neq 1$ and we have proved our statement for all words $< w$. Let $m = |w|$ and $\operatorname{sig}(w) = \alpha = (a_1, a_2, \dots)$. For each word u with $\operatorname{sig}(u) = \alpha$ let \tilde{u} be a word in echelon form

such that $u \rightsquigarrow \tilde{u}$, as per the Ordering Theorem. Thus, by Lemma 6.3 we have $\text{ct}(w) = \text{ct}(u) \geq_{lex} \text{ct}(\tilde{u})$. Let

$$M = |\{u \mid \text{sig}(u) = \alpha \text{ and } \text{ct}(\tilde{u}) = \text{ct}(w)\}|.$$

Since u and \tilde{u} belong to the same conjugacy class in G , we can rewrite equation (8.3) as follows:

$$(8.4) \quad \sum_{\substack{\text{sig}(u)=\alpha \\ \tilde{u} < w}} \text{Tr}(\rho_1(\tilde{u})) + \sum_{\substack{\text{sig}(u)=\alpha \\ \text{ct}(\tilde{u})=\text{ct}(w)}} \text{Tr}(\rho_1(\tilde{u})) \\ = \sum_{\substack{\text{sig}(u)=\alpha \\ \tilde{u} < w}} \text{Tr}(\rho_2(\tilde{u})) + \sum_{\substack{\text{sig}(u)=\alpha \\ \text{ct}(\tilde{u})=\text{ct}(w)}} \text{Tr}(\rho_2(\tilde{u})).$$

If $\tilde{u} < w$ then by induction hypothesis we get

$$\text{Tr}(\rho_1(w)) = \text{Tr}(\rho_1(\tilde{w})) = \text{Tr}(\rho_2(\tilde{w})) = \text{Tr}(\rho_2(w)),$$

so we may assume that $\text{ct}(w) = \text{ct}(\tilde{w})$. Also, note that, in type D_{n+1} , if $a_n = 0$ (resp. $a_{n+1} = 0$) and $\text{sig}(u) = \alpha$, then by the nature of the possible admissible transformations, \tilde{u} does not involve g_n (resp. g_{n+1}). Therefore, in view of Remark 7.3 and Proposition 7.4, in all possible cases for u with $\text{sig}(u) = \alpha$ and $\text{ct}(\tilde{u}) = \text{ct}(w)$ we must have \tilde{u} in the same conjugacy class as w . Therefore $\text{Tr}(\rho_i(\tilde{u})) = \text{Tr}(\rho_i(w))$ hence

$$\sum_{\substack{\text{sig}(u)=\alpha \\ \text{ct}(\tilde{u})=\text{ct}(w)}} \text{Tr}(\rho_i(\tilde{u})) = M \text{Tr}(\rho_i(w))$$

for $i = 1, 2$. Since by the induction hypothesis $\text{Tr}(\rho_1(\tilde{u})) = \text{Tr}(\rho_2(\tilde{u}))$ whenever $\tilde{u} < w$, equation (8.4) reduces to

$$M \text{Tr}(\rho_1(w)) = M \text{Tr}(\rho_2(w))$$

and the desired conclusion is immediate. \square

We also note the following result which was obtained during the course of establishing equality (8.3) in the proof above.

Theorem 8.5. *Let G be any finitely generated group with generators g_1, \dots, g_n , and let ρ_1 and ρ_2 be two finite dimensional representations of G . If the equality (8.1) holds, then for every $(k_1, \dots, k_n) \in \mathbb{N}^n$ we have*

$$(8.6) \quad \sum_{\text{sig}(w)=(k_1, \dots, k_n)} \chi_{\rho_1}(w) = \sum_{\text{sig}(w)=(k_1, \dots, k_n)} \chi_{\rho_2}(w),$$

where w denotes a word in the alphabet given by the set $\{g_1, \dots, g_n\}$.

9. EXAMPLE: \tilde{C}_2

Recall that \tilde{C}_2 is the Coxeter group with three generators b_1, b_2, b_3 that satisfy the relations:

$$(9.1) \quad b_j^2 = 1, \quad j = 1, 2, 3; \quad (b_1 b_2)^4 = (b_2 b_3)^4 = 1; \quad b_1 b_3 = b_3 b_1.$$

It is well known that this group is *affine*, that is, it contains an abelian normal subgroup such that the quotient group is finite. In this particular case it is easy to see that this normal abelian subgroup is the following: let

$$r_1 = b_1 b_2 b_3 b_2, \quad r_2 = b_2 b_1 b_2 b_3.$$

Let us check that the group generated by r_1 and r_2 is an abelian normal subgroup of \tilde{C}_2 . First we observe that the relations (9.1) imply

$$(9.2) \quad \begin{aligned} b_1 r_1 &= r_1^{-1} b_1 \\ b_2 r_1 &= r_2 b_2, \\ b_3 r_1 &= r_1 b_3, \\ b_1 r_2 &= r_2 b_1 \\ b_2 r_2 &= r_1 b_2, \\ b_3 r_2 &= r_2^{-1} b_3 \\ b_3 r_2^{-1} &= r_2 b_3 \end{aligned}$$

We now can use these last relations to establish that r_1 and r_2 commute:

$$r_1 r_2 = r_1 b_2 b_1 b_2 b_3 = b_2 r_2 b_1 b_2 b_3 = b_2 b_1 r_2 b_2 b_3 = b_2 b_1 b_2 r_1 b_3 = b_2 b_1 b_2 b_3 r_1 = r_2 r_1.$$

Now, relations (9.2) imply that

$$\begin{aligned} b_1 r_1 b_1 &= r_1^{-1} \\ b_2 r_1 b_2 &= r_2 \\ b_3 r_1 b_3 &= r_1 \\ b_1 r_2 b_1 &= r_2 \\ b_2 r_2 b_2 &= r_1 \\ b_3 r_2 b_3 &= r_2^{-1} \end{aligned}$$

These last relations show that the subgroup \mathcal{N} of \tilde{C}_2 generated by r_1 and r_2 is a normal subgroup. We further remark that $b_1 = b_2 b_3 b_2 r_1^{-1}$, and, therefore, the cosets of b_2 and b_3 generate \tilde{C}_2/\mathcal{N} . We will now show:

Theorem 9.3. *If for two finite dimensional unitary representations ρ_1 and ρ_2 of the affine Coxeter group \tilde{C}_2 we have*

$$\begin{aligned} &\sigma_p(\rho_1(b_2), \rho_1(b_3), \rho_1(r_1), \rho_1(r_2), \rho_1(r_1^{-1}), \rho_1(r_2^{-1})) \\ &= \sigma_p(\rho_2(b_2), \rho_2(b_3), \rho_2(r_1), \rho_2(r_2), \rho_2(r_1^{-1}), \rho_2(r_2^{-1})), \end{aligned}$$

then for these representations $\chi_{\rho_1} = \chi_{\rho_2}$.

Proof. First we note that the subgroup generated by b_2 and b_3 is the dihedral group $I(4) = B_2$. The only words in echelon form here are 1, b_2 , b_3 , $b_2 b_3$, $b_2 b_3 b_2$, and $b_2 b_3 b_2 b_3$. Note that b_3 and $b_2 b_3 b_2$ belong to the same conjugacy class. Since every element of G can be written as $w r_1^{m_1} r_2^{m_2}$ for some word w in b_2 and b_3 , formulae (9.2) and Theorem 7.5 imply that every element in \tilde{C}_2 has in its conjugacy class a word in the form

$$b_2^{k_1} b_3^{l_1} b_2^{k_2} b_3^{l_2} r_1^{m_1} r_2^{m_2}$$

where $k_1, k_2, l_1 \in \{0, 1\}$, $k_1 \geq k_2$, $l_1 \geq k_2$, and $m_1, m_2 \in \mathbb{Z}$. In other words, each word in \tilde{C}_2 has in its conjugacy class one of the words

$$\begin{aligned} w_0(m_1, m_2) &= r_1^{m_1} r_2^{m_2}, \\ w_1(m_1, m_2) &= b_2 r_1^{m_1} r_2^{m_2}, \\ w_2(m_1, m_2) &= b_3 r_1^{m_1} r_2^{m_2}, \\ w_3(m_1, m_2) &= b_2 b_3 r_1^{m_1} r_2^{m_2}, \quad \text{or} \\ w_4(m_1, m_2) &= b_2 b_3 b_2 b_3 r_1^{m_1} r_2^{m_2}. \end{aligned}$$

Thus, we are to show that if

$$\begin{aligned} \sigma(-I, \rho_1(b_2), \rho_1(b_3), \rho_1(r_1), \rho_1(r_2), \rho_1(r_1^{-1}), \rho_1(r_2^{-1})) \\ = \sigma(-I, \rho_2(b_2), \rho_2(b_3), \rho_2(r_1), \rho_2(r_2), \rho_2(r_1^{-1}), \rho_2(r_2^{-1})), \end{aligned}$$

then for every $m_1, m_2 \in \mathbb{Z}$ and every $j = 0, 1, 2, 3, 4$

$$(9.4) \quad \chi_{\rho_1}(w_j(m_1, m_2)) = \chi_{\rho_2}(w_j(m_1, m_2)).$$

A similar argument to the one in Theorem 8.5 shows that for every $\eta \in \mathbb{N}^6$ with $\eta = (n_1, n_2, n_3, n_4, n_5, n_6)$

$$(9.5) \quad \sum_{\text{sig}(w)=\eta} \chi_{\rho_1}(w) = \sum_{\text{sig}(w)=\eta} \chi_{\rho_2}(w).$$

where the sums are taken over all words comprised of n_1 b_2 -s, n_2 b_3 -s, n_3 r_1 -s, n_4 r_2 -s, n_5 r_1^{-1} -s, and n_6 r_2^{-1} -s.

In particular, using circular transformation if necessary, we see that in the cases $n_1 = 1, n_2 = n_5 = n_6 = 0$ all terms in each of the sums (9.5) are the same. Therefore, for all $m_1, m_2 \geq 0$

$$(9.6) \quad \chi_{\rho_1}(w_1(m_1, m_2)) = \chi_{\rho_2}(w_1(m_1, m_2)).$$

A similar argument shows that the same equality holds for $m_1 \geq 0, m_2 \leq 0$, $m_1 \leq 0, m_2 \geq 0$, $m_1, m_2 \leq 0$, and, hence, (9.6) holds for all $m_1, m_2 \in \mathbb{Z}$.

Applying the same reasoning to the case $n_1 = 0, n_2 = 1$ we obtain

$$(9.7) \quad \chi_{\rho_1}(w_j(m_1, m_2)) = \chi_{\rho_2}(w_j(m_1, m_2)).$$

holds for $j = 2$ as well.

Let us show that (9.7) holds for $j = 3$. We will prove it by induction in $s = |m_1| + |m_2|$. If $s = 0$, $w_3(0, 0)$ belongs to the subgroup generated by b_2 and b_3 . Since the joint spectrum of $\rho_j(b_2)$, $\rho_j(b_3)$ and the identity is the intersection of $\sigma(-I, \rho_1(b_2), \rho_1(b_3), \rho_1(r_1), \rho_1(r_2), \rho_1(r_1^{-1}), \rho_1(r_2^{-1}))$ with the plane $\{x_3 = x_4 = x_5 = x_6 = 0\}$, we see that

$$\sigma(-I, \rho_1(b_2), \rho_1(b_3)) = \sigma(-I, \rho_2(b_2), \rho_2(b_3)),$$

so by our previous results

$$\chi_{\rho_1}(b_2 b_3) = \chi_{\rho_2}(b_2 b_3).$$

Now, let us fix $m_1, m_2 \geq 0$ with $m_1 + m_2 > 0$, and consider $n_1 = n_2 = 1$, $n_3 = m_1$, $n_4 = m_2$, $n_5 = n_6 = 0$. In each side of (9.5) there are characters of words in one of two forms:

$$(9.8) \quad r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3}, \quad \text{or} \quad r_1^{q_1} r_2^{t_1} b_3 r_1^{q_2} r_2^{t_2} b_2 r_1^{q_3} r_2^{t_3},$$

where $q_j, t_j \geq 0$, $q_1 + q_2 + q_3 = m_1$, $t_1 + t_2 + t_3 = m_2$ (we observe that the order of r_1 and r_2 does not matter since they commute).

Using circular transformations shows that each word that appears in the second form in (9.8) is in the same conjugacy class as the first one, of course with different q_j and t_j which still sum up to m_1 and m_2 respectively. Thus, it suffices to show the equality of characters for words in the first form in (9.8).

We use circular transformation to get

$$(9.9) \quad \begin{aligned} r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3} &\sim b_2 r_1^{q-2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3} r_1^{q_1} r_2^{t_1} \\ &= b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_1+q_3} r_2^{t_1+t_3}. \end{aligned}$$

By (9.2) $r_2 b_3 = b_3 r_2^{-1}$, so we can get from (9.9)

$$r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3} \sim b_2 r_1^{q_2} b_3 r_1^{q_1+q_3} r_2^{t_1-t_2+t_3}.$$

Since by (9.2) r_1 and b_3 commute, we finally obtain

$$(9.10) \quad r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3} \sim b_2 b_3 r_1^{q_1+q_2+q_3} r_2^{t_1-t_2+t_3} = b_2 b_3 r_1^{m_1} r_2^{t_1-t_2+t_3}.$$

If $|t_1 - t_2 + t_3| < m_2$, by induction

$$(9.11) \quad \chi_{\rho_1}(r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3}) = \chi_{\rho_2}(r_1^{q_1} r_2^{t_1} b_2 r_1^{q_2} r_2^{t_2} b_3 r_1^{q_3} r_2^{t_3}).$$

Of course, for non-negative t_1, t_2, t_3

$$(9.12) \quad t_1 + t_2 + t_3 = |t_1 - t_2 + t_3| \implies t_2 = 0, \text{ or } t_1 = t_3 = 0.$$

We will now show that every term in (9.5) that are not conjugates of words in the form (9.8) with a smaller $m_1 + m_2$ is in the same conjugacy class as $b_2 b_3 r_1^{m_1+m_2}$. Of course, this together with (9.5) and (9.11) will imply that (9.4) holds for $j = 3$.

By (9.12) for such a word either $t - 2 = 0$, that is the word is in the form

$$(9.13) \quad \begin{aligned} r_1^{q_1} r_2^t b_2 r_1^{q_2} b_3 r_1^{q_3} r_2^{m_2-t} &= r_1^{q_1} r_2^t b_2 b_3 r_1^{m_1-q_1} r_2^{m_2-t} \\ &\sim b_2 b_3 r_1^{m_1} r_2^{m_2}, \end{aligned}$$

(here we used the fact that b_3 and r_1 commute and a circular transformation), or $t_1 = t_3 = 0$, so the word is

$$(9.14) \quad \begin{aligned} r_1^{q_1} b_2 r_1^{q_2} r_2^{m_2} b_3 r_1^{q_3} &= r_1^{q_1} b_2 r_2^{m_2} r_1^{q_1} b_3 r_1^{q_3} \\ &= r_1^{q_1} b_2 r_2^{m_2} b_3 r_1^{q_2+q_3} \\ &\sim b_2 r_2^{m_2} b_3 r_1^{m_1}, \end{aligned}$$

here we again we used the commuting of b_3 and r_1 and a circular transformation that moved $r_1^{q_1}$ to the end of the word.

Now, by (9.2) $b_2 r_2 = r_1 b_2$ and $r_2 b_2 = b_2 r_1$, therefore, in (9.13) we have

$$b_2 b_3 r_1^{m_1} r_2^{m_2} \sim r_2^{m_2} b_2 b_3 r_1^{m_1} = b_2 r_1^{m_2} b_3 r_1^{m_1} = b_2 b_3 r_1^{m_1+m_2}.$$

For (9.14) we have

$$b_2 r_2^{m_2} b_3 r_1^{m_1} = r_1^{m_2} b_2 b_3 r_1^{m_1} \sim b_2 b_3 r_1^{m_1+m_2}.$$

This finishes the proof of (9.4) for $j = 3$ and $m_1, m_2 \geq 0$. The proofs in the cases when one of, or both m_1 and m_2 are negative are practically identical with the only difference being that for $m_1 < 0$ r_1 is replaced with r_1^{-1} , and r_2 is replaced with r_2^{-1} when $m_2 < 0$.

We now prove that (9.4) holds for $j = 4$. Again, for every pair of representations with the same joint spectrum of $b_2, b_3, r_1, r_2, r_1^{-1}, r_2^{-1}$ the restrictions of these representations to the subgroup generated by b_2 and b_3 are unitary equivalent so

$$\chi_{\rho_1}(b_2 b_3 b_2 b_3) = \chi_{\rho_2}(b_2 b_3 b_2 b_3).$$

Next we observe that relations (9.2) imply

$$b_2 r_1 b_2 = r_2, \quad b_2 r_2 b_2 = r_1, \quad b_2 r_1^{-1} b_2 = r_2^{-1}, \quad b_2 r_2^{-1} b_2 = r_1^{-1},$$

and, therefore,

$$(9.15) \quad b_2 r_1^k b_2 = r_2^k, \quad b_2 r_2^k b_2 = r_1^k, \quad k \in \mathbb{Z}.$$

Relations (9.15) imply

$$(9.16) \quad b_2 b_3 b_2 b_3 r_1^k r_2^l \sim b_2 b_3 b_2 b_3 r_1^l r_2^k, \quad k, l \in \mathbb{Z}.$$

Indeed,

$$\begin{aligned} b_2 b_3 b_2 b_3 r_1^k r_2^l &= b_2 b_3 b_2 b_3 b_2 r_2^k b_2 b_2 r_1^l b_2 = b_2 b_3 b_2 b_3 b_2 r_2^k r_1^l b_2 \\ &= b_3 b_2 b_3 r_2^k r_1^l b_2 \sim b_2 b_3 b_2 b_3 r_1^l r_2^k. \end{aligned}$$

We also remark that

$$(9.17) \quad b_2 b_3 b_2 b_3 r_1^k r_2^l \sim b_2 b_3 b_2 b_3 r_1^{k-2} r_2^l$$

and, similarly

$$(9.18) \quad b_2 b_3 b_2 b_3 r_1^k r_2^l \sim b_2 b_3 b_2 b_3 r_1^k r_2^{l-2}.$$

Verification of (9.17) goes as follows:

$$\begin{aligned} b_2 b_3 b_2 b_3 r_1^k r_2^l &= b_2 b_3 b_2 b_3 r_1^{k-1} r_2^l r_1 \\ &\sim r_1 b_2 b_3 b_2 b_3 r_1^{k-1} r_2^l \\ &= b_2 r_2 b_3 b_2 b_3 r_1^{k-1} r_2^l \\ &= b_2 b_3 r_2^{-1} b_2 b_3 r_1^{k-1} r_2^l \\ &= b_2 b_3 b_2 r_1^{-1} b_3 r_1^{k-1} r_2^l \\ &= b_2 b_3 b_2 b_3 r_1^{k-2} r_2^l. \end{aligned}$$

Relation (9.18) is an immediate consequence of (9.17) and (9.16). Now, (9.17) and (9.18) show that if $|k| + |l| \geq 2$, and $(k, l) \neq (\pm 1, \pm 1)$, then

$$(9.19) \quad b_2 b_3 b_2 b_3 r_1^k r_2^l \sim b_2 b_3 b_2 b_3 r_1^{k_1} r_2^{l_1} \quad \text{with} \quad |k_1| + |l_1| < |k| + |l|.$$

Suppose that $m_1, m_2 \geq 0$. Each term in (9.5) with $n_1 = 2, n_2 = 2, n_3 = m_1, n_4 = m_2, n_5 = n_6 = 0$ is in the form

$$(9.20) \quad r_1^{q_1} r_2^{t_1} b_{j_1} r_1^{q_2} r_2^{t_2} b_{j_2} r_1^{q_3} r_2^{t_3} b_{j_3} r_1^{q_4} r_2^{t_4} b_{j_4} r_1^{q_5} r_2^{t_5},$$

where $q_k, t_k \geq 0$, $\sum q_k = m_1$, $\sum t_k = m_2$, $j_k = \{2, 3\}$ and there are two b_2 among b_{j_k} and two b_3 . Observe that, if for some $k = 1, 2, 3$ $j_k = j_{k+1}$, then formulae (9.2) show that the word is a conjugate of either $w_3(l_1, l_2)$, or of $w_0(l_1, l_2)$ for some l_1, l_2 . Since for such words we have already proven the equality of the characters, the sums of terms in (9.5) with $j_k \neq j_{k+1}$ on the left and on the right are the same. Each of such terms is in the form (9.20) with $b_{j_1} = b_2, b_{j_2} = b_3, b_{j_3} = b_2, b_{j_4} = b_3$, or with $b_{j_1} = b_3, b_{j_2} = b_2, b_{j_3} = b_3, b_{j_4} = b_2$. Using circular transformations we easily see that the latter one is a conjugate of the former (possibly with different

q -s and t -s). As it was mentioned above, relations (9.2) shows that each such term is a conjugate of

$$b_2 b_3 b_2 b_3 r_1^k r_2^l$$

for some k and l .

A similar argument shows that the same conjugacy relation holds for arbitrary $m_1, m_2 \in \mathbb{Z}$.

Now we use an induction argument in $|m_1| + |m_2|$. We have already proven that (9.4) holds for $w_4(0, 0)$. Relation (9.19) shows that it suffices to prove it for $w_4(\pm 1, 0)$, $w_4(0, \pm 1)$ and $w_4(\pm 1, \pm 1)$.

Obviously, (9.16)–(9.18) imply

$$(9.21) \quad w_4(1, 0) \sim w_4(0, 1) \sim w_4(-1, 0) \sim w_4(0, -1),$$

$$(9.22) \quad w_4(1, 1) \sim w_4(-1, 1) \sim w_4(1, -1) \sim w_4(-1, -1).$$

All terms in relation (9.5) with $n_1 = 2, n_2 = 2, n_3 = 1, n_4 = n_5 = n_5 = 0$ that are not conjugates of $w_3(k, l)$ or $w_0(k, l)$ are

$$\begin{aligned} r_1 b_2 b_3 b_2 b_3, & \quad r_1 b_3 b_2 b_3 b_2, \\ b_2 r_1 b_3 b_2 b_3, & \quad b_3 r_1 b_2 b_3 b_2, \\ b_2 b_3 r_1 b_2 b_3, & \quad b_3 b_2 r_1 b_3 b_2, \\ b_2 b_3 b_2 r_1 b_3, & \quad b_3 b_2 b_3 r_1 b_2, \\ b_2 b_3 b_2 b_3 r_1, & \quad b_3 b_2 b_3 b_2 r_1. \end{aligned}$$

It is easy to see using (9.2) that each of them is a conjugate of one of $w_4(\pm 1, 0)$ or $w_4(0, \pm 1)$. By (9.21) all of them are in the same conjugate class and, therefore, (9.4) implies

$$\chi_{\rho_1}(w_4(\pm 1, 0)) = \chi_{\rho_2}(w_4(\pm 1, 0)), \quad \chi_{\rho_1}(w_4(0, \pm 1)) = \chi_{\rho_2}(w_4(0, \pm 1)).$$

Similarly, we can show that each term in (9.5) corresponding to $n_1 = 2, n_2 = 2, n_3 = n_4 = 1, n_5 = n_6 = 0$ is either a conjugate of $w_4(0, 0)$, or of $w_4(\pm 2, 0) \sim w_4(0, 0)$, or of $w_4(0, \pm 2) \sim w_4(0, 0)$, or $w_4(\pm 1, \pm 1)$. For the first three the equality of characters has been established. For the rest this equality follows from (9.4) and (9.22). \square

REFERENCES

- [1] A. Bjorner and F. Brenti, *Combinatorics of Coxeter Groups*, Springer, 2005.
- [2] J. P. Bannon, P. Cade, and R. Yang, On the spectrum of Banach algebra-valued entire functions, *Ill. J. Math.*, **55** (4), (2011), 1455–1465.
- [3] P. Cade and R. Yang, Projective spectrum and cyclic cohomology, *J. Funct. Anal.*, **265** (9) (2013), 1916–1933.
- [4] F. Catanese, Babbage’s conjecture, contact surfaces, symmetric determinantal varieties and applications, *Invent. Math.*, **63** (1981), 433–465.
- [5] I. Chagouel, M. Stessin, and K. Zhu, Geometric spectral theory for compact operators, *Trans. Amer. Math. Soc.*, **368** (3) (2016), 1559–1582.
- [6] H. M. S. Coxeter, Discrete groups generated by reflections, *Ann. of Math. (2)*, **35** (3) (1934), 588–621.
- [7] H. M. S. Coxeter, The complete enumeration of finite groups of the form $r_i^2 = (r_i r_j)^{k_{ij}} = 1$, *J. London Mat. Soc.*, **10** (1) (1935), 21–25.
- [8] L. E. Dickson, An elementary exposition of Frobenius’ theory of group characters and group determinants, *Ann. of Math. (2)*, **4** (1902), 25–49; also in *Mathematical papers, Vol. II*, Chelsea, New York, 1975, 737–761.
- [9] L. E. Dickson, On the group defined for any given field by the multiplication table of any given finite group, *Trans. Amer. Math. Soc.*, **3** (1902), 377–382; also in *Mathematical Papers, Vol. II*, Chelsea, New York, 1975, 75–91.

- [10] L. E. Dickson, Modular theory of group-matrices, *Trans. Amer. Math. Soc.*, **8** (1907), 389–398; also in *Mathematical Papers, Vol. II*, Chelsea, New York, 1975, 251–260.
- [11] L. E. Dickson, Modular theory of group-characters, *Bull. Amer. Math. Soc.*, **13** (1907), 477–488; also in *Mathematical Papers, Vol. IV*, Chelsea, New York, 1975, 535–546.
- [12] L. E. Dickson, Determination of all general homogeneous polynomials expressible as determinants with linear elements, *Trans. Amer. Math. Soc.*, **22** (1921), 167–179.
- [13] I. Dolgachev, *Classical Algebraic Geometry: A modern View*, Cambridge University Press, 2012.
- [14] R. G. Douglas and R. Yang, Hermitian geometry on resolvent set (I), *Proceedings of the International Workshop on Operator Theory (IWOTA 2016)*, to appear.
- [15] R. G. Douglas and R. Yang, Hermitian geometry on resolvent set (II), preprint.
- [16] E. Formanek and D. Sibley, The group determinant determines the group, *Proc. Amer. Math. Soc.*, **112** (1991), 649–656.
- [17] F. G. Frobenius, Über vertauschbare Matrizen, *Sitz. Kön. Preuss. Akad. Wiss. Berlin* (1896) 601–614; also in *Gesammelte Abhandlungen, Band II*, Springer-Verlag, New York, 1968, 705–718.
- [18] F. G. Frobenius, Über Gruppencharaktere, *Sitz. Kön. Preuss. Akad. Wiss. Berlin* (1896) 985–1021; also in *Gesammelte Abhandlungen, Band III*, Springer-Verlag, New York, 1968, 1–37.
- [19] F. G. Frobenius, Über die Primfactoren der Gruppendeterminante. *Sitz. Kön. Preuss. Akad. Wiss. Berlin* (1896) 1343–1382; also in *Gesammelte Abhandlungen, Band III*, Springer-Verlag, New York, 1968, 38–77.
- [20] M. Geck and G. Pfeiffer, *Characters of Finite Coxeter Groups and Iwahori-Hecke Algebras*, Calderon Press, Oxford, 2000.
- [21] P. Gonzalez-Vera and M.I. Stessin, Joint spectra of Toeplitz operators and optimal recovery of analytic functions, *Constructive Approximation*, **36** (1) (2012), 53–82.
- [22] R. Grigorchuk and R. Yang, Joint spectrum and infinite dihedral group, *Proc. Steklov Inst. Math.*, **297** (2017), 145–178.
- [23] J. W. Helton, S. A. McCulloch, and V. Vinnikov, Non-commutative convexity arises from linear matrix inequalities, *J. Funct. Anal.*, **240** (1) (2006), 105–191.
- [24] J. W. Helton and V. Vinnikov, Linear matrix inequality representation set, *Comm. Pure and Appl. Math.*, **60**, (2007), 654–674.
- [25] W. He, X. Wang, and R. Yang, Projective spectrum and kernel bundle (II), *J. Oper. Theory*, **297** (2017), 417–433.
- [26] H.-J. Hoehnke and K. W. Johnson, The 1-, 2-, and 3-characters determine a group, *Bull. Amer. Math. Soc.*, **27** (1992), 243–245.
- [27] H.-J. Hoehnke and K. W. Johnson, k -Characters and group invariants, *Comm. in Algebra*, **26** (1998), 1–27.
- [28] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, *Cambridge Studies in Advanced Mathematics* **10**, Cambridge University Press, 1992.
- [29] K. W. Johnson, On the group determinant, *Math. Proc. Cambridge Philos. Soc.*, **109** (1991), 299–311.
- [30] D. Kerner and V. Vinnikov, Determinantal representations of singular hypersurfaces in \mathbb{P}^n , *Adv. Math.*, **231** (2012), 1619–1654.
- [31] P. D. Lax, *Functional Analysis*, Wiley, 2002.
- [32] T. S. Motzkin and O. Taussky, Pairs of matrices with property L, *Trans. Amer. Math. Soc.*, **73** (1952), 108–114.
- [33] T. Rivlin, *The Chebyshev polynomials*, *Pure and Applied Mathematics*, Wiley-Interscience, New York-London-Sydney, 1974.
- [34] T. G. Room, Self-transformations of determinantal quadric surfaces I, *Proc. London Mat. Soc.*, **51** (2) (1950), 348–361.
- [35] J. P. Serre, *Linear Representations of Finite Groups*, *Graduate Texts in Mathematics* **42**, Springer-Verlag, New York-Heidelberg, 1977.
- [36] M. Stessin, R. Yang, K. Zhu, Analyticity of a joint spectrum and multivariable analytic Fredholm theorem, *New York J. Math.*, **17a** (2011), 39–44.
- [37] M. I. Stessin and A. B. Tchertnev, Spectral algebraic curves and decomposable operator tuples, [arxiv:1509.06274](https://arxiv.org/abs/1509.06274).
- [38] M. I. Stessin and A. B. Tchernev, in preparation.

- [39] V. Vinnikov, Complete description of determinantal representations of smooth irreducible curves, *Lin. Algebra and Appl.*, **125** (1989), 103–140.
- [40] R. Yang, Projective spectrum in Banach algebras, *J. Topol. Anal.*, **1** (2009), 289–306.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TOLEDO, TOLEDO, OH 43606, USA

E-mail address: `zcuckovi@math.utoledo.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY AT ALBANY, SUNY, ALBANY, NY 12222, USA

E-mail address: `mstessin@albany.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY AT ALBANY, SUNY, ALBANY, NY 12222, USA

E-mail address: `atchernev@albany.edu`