

# A brief note about the Scott entanglement measure

Saeed Haddadi\*

*Department of Physics, Payame Noor University,*

*P.O. Box 19395-3697, Tehran, Iran.*

## Abstract

Scott studied various interesting aspects of multipartite entanglement measures. He has defined a class of related multipartite entanglement measures in the obvious manner. Recently, Jafarpour *et al.* (Int. J. Quantum Inform. **13**, 1550047 2015) have calculated the entanglement quantity of two-dimensional 5-site spin system by means of Scott measure (or generalized Meyer-Wallach measure). As it was presented in their calculation  $Q_2 > Q_3$ . In this paper we would like to point out that in 5-qudit system,  $Q_2$  does not provide a stronger entanglement than  $Q_3$ , while there is a simple relationship between them.

PACS numbers: 03.67.Mn, 03.67.-a

Keywords: Scott measure; Meyer-Wallach measure; 5-site spin system.

---

\*Electronic address: haddadi@physicist.net

In 2004, Scott [1] has investigated the average bipartite entanglement, over all possible divisions of a multipartite system, as a useful multipartite entanglement measure [2]. The Scott measure for  $N$ -qudit system ( $d$ -dimensional) is defined as follows [1, 3]

$$Q_m(|\psi\rangle) \equiv \frac{d^m}{d^m - 1} \left( 1 - \frac{m!(N-m)!}{N!} \sum_{|S|=m} \text{Tr}[\rho_S^2] \right), \quad m = 1, \dots, \lfloor N/2 \rfloor \quad (1)$$

where  $S \subset \{1, \dots, N\}$  and  $\rho_S = \text{Tr}_S(|\psi\rangle\langle\psi|)$  is the reduced density matrix for qudits  $S$  after tracing out the rest ( $d=2$  for two-dimensional spin system). The  $Q_m$  quantities correspond to the average entanglement between subsystems which consists  $m$  qudits and the remaining  $N - m$  qudits [4].

Recently, Jafarpour *et al.* [5] based on Scott measure have calculated the entanglement quantity in two-dimensional 5-site spin system. The authors of the original paper [5] have calculated  $Q_2$  and  $Q_3$  in Table 1 and Table 2, whereas there is a relation between them fixed by the Schmidt decomposition [6]; namely the eigenvalues of the reduced density matrix of qudits 12, will be the same as the eigenvalues of the reduced density matrix of qudits 345. In other words,  $\text{Tr}[\rho_{ij}^2] = \text{Tr}[\rho_{klm}^2]$  via  $i, j, k, l, m \in \{1, \dots, 5\}$  for all 5-qudit pure states [7], which this implies a simple relationship between  $Q_2$  and  $Q_3$ . We emphasize that in a 5-qudit system,  $Q_2$  and  $Q_3$  both capture entanglement in partition 2 qudits vs 3 qudits. We have also shown in Ref [8] that for 5-qudit system,  $Q_2$  and  $Q_3$  both refer to 3-2 partitions of the same state. Therefore, Jafarpour *et al.* [5] could not say that  $Q_2 > Q_3$  for all states in a two-dimensional 5-site spin system.

In conclusion, the above analysis shows that the authors [5] did not use the Scott measure properly. Since  $Q_2$  and  $Q_3$  are proportional to each other by a numerical factor (as seen from (1)). In fact for 5-qudit system  $Q_1$  refers to 1-4 partitions and  $Q_2$  or  $Q_3$  both refer to 3-2 partitions of the same state. Hence, we conclude that  $Q_2$  does not provide a stronger entanglement than  $Q_3$  in 5-qudit system.

## References

---

- [1] A. J. Scott, *Phys. Rev. A* **69** (2004) 052330.
- [2] S. Haddadi and M. Bohloul, *Int. J. Theor. Phys.* **57** (2018) 3912.
- [3] P. J. Love *et al.*, *Quantum. Inf. Process.* **6** (2007) 187.
- [4] A. Borras *et al.*, *J. Phys. A: Math. Theor.* **40** (2007) 13407.
- [5] M. Jafarpour, S. Ghanavati and D. Afshar, *Int. J. Quantum Inform.* **13** (2015) 1550047.
- [6] A. Ekert and P. L. Knight, *Am. J. Phys.* **63** (1995) 415.
- [7] XN. Zhu and SM. Fei, *Quantum. Inf. Process.* **13** (2014) 815.
- [8] S. Haddadi, *Int. J. Theor. Phys.* **56** (2017) 2811.