

PERIODICITY INDUCED BY NOISE AND INTERACTION IN THE KINETIC MEAN-FIELD FITZHUGH-NAGUMO MODEL

ERIC LUÇON AND CHRISTOPHE POQUET

ABSTRACT. We consider the long-time behavior of a population of mean-field oscillators modeling the activity of interacting excitable neurons in large population. Each neuron is represented by its voltage and recovery variables, which are solution to a FitzHugh-Nagumo system, and interacts with the rest of the population through a mean-field linear coupling, in the presence of noise. The aim of the paper is to study the emergence of collective oscillatory behaviors induced by noise and interaction on such a system. The main difficulty of the present analysis is that we consider the kinetic case, where interaction and noise are only imposed on the voltage variable. We prove the existence of a stable cycle for the infinite population system, in a regime where the local dynamics is small.

1. INTRODUCTION

1.1. A mean-field model of interacting FitzHugh-Nagumo neurons. We are interested in this paper in the behavior as $t \rightarrow \infty$ of the following McKean-Vlasov process

$$\begin{cases} dX_t = \delta \left(X_t - \frac{X_t^3}{3} - Y_t \right) dt - K (X_t - \mathbf{E}[X_t]) dt + \sqrt{2}\sigma dB_t \\ dY_t = \frac{\delta}{c} (X_t + a - bY_t) dt \end{cases}, \quad t \geq 0, \quad (1.1)$$

where $(X_t, Y_t) \in \mathbb{R}^2$, B_t is a standard Brownian motion on \mathbb{R} , $a \in \mathbb{R}$ and b, c, δ, K, σ are positive parameters. The evolution (1.1) is a prototype of a nonlinear stochastic differential equation (the nonlinearity coming from the fact that X_t interacts with its own law through its expectation $\mathbf{E}[X_t]$). It is named kinetic by analogy to the classical kinetic interacting particle systems, noise and interactions are only applied on the "momentum" X_t , and not on the "position" Y_t .

The system (1.1) is the natural macroscopic limit (as $n \rightarrow \infty$) of the following system of coupled mean-field diffusions $(X_{i,t}, Y_{i,t})$, $i = 1, \dots, n$, $n \geq 1$

$$\begin{cases} dX_{i,t} = \delta \left(X_{i,t} - \frac{X_{i,t}^3}{3} - Y_{i,t} \right) dt - K \left(X_{i,t} - \frac{1}{n} \sum_{j=1}^n X_{j,t} \right) dt + \sqrt{2}\sigma dB_{i,t} \\ dY_{i,t} = \frac{\delta}{c} (X_{i,t} + a - bY_{i,t}) dt \end{cases}, \quad t \geq 0, \quad (1.2)$$

where (B_1, \dots, B_n) are i.i.d. standard Brownian motions. The motivation comes from neuroscience: (1.2) models the evolution of n neurons of FitzHugh-Nagumo type, each represented by its voltage X_i and recovery variable Y_i , that are coupled through a linear mean-field interaction (this corresponds to a coupling via electrical synapses, see [6]).

Date: June 24, 2019.

2010 Mathematics Subject Classification. 60K35, 35K55, 35Q84, 37N25, 82C26, 82C31, 92B20.

Key words and phrases. FitzHugh-Nagumo model, McKean-Vlasov process, nonlinear Fokker-Planck equation, mean-field systems, excitable systems, slow-fast dynamics, noise-induced dynamics, Wasserstein distance.

1.2. Emergence of collective structured dynamics for excitable systems. In (1.2), the intrinsic dynamics of each neuron is of FitzHugh-Nagumo type [14, 25, 2]: when $K = \sigma = 0$, $\delta = 1$, the system (1.2) reduces to a collection of copies of the isolated system

$$d(X_t, Y_t) = F(X_t, Y_t)dt, \quad (1.3)$$

where F is given by

$$F(x, y) := \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} := \begin{pmatrix} x - \frac{x^3}{3} - y \\ \frac{1}{c}(x + a - by) \end{pmatrix}, \quad x, y \in \mathbb{R}. \quad (1.4)$$

Although the transitions in the FitzHugh-Nagumo model are complex in general (see [25]), two main dynamical patterns emerge for (1.3): a resting state (corresponding to a unique stable point for (1.3)) and a spiking regime (corresponding to a limit cycle for (1.3), see [25] and Figure 1 below). In this sense, the system (1.3) is a prototype of an excitable dynamics [19]: it is possible to choose appropriately the parameters a, b, c (and we will do so in the following) so that the unperturbed system (1.3) is in a resting state but such that the addition of a small perturbation makes the system fall into a oscillatory regime (spiking activity).




FIGURE 1. Phase diagrams for the system (1.3) for two choices of parameters a, b and c (the voltage X is represented along the x -axis and the recovery variable Y on the y -axis). Stable (resp. unstable) points and limit cycles are represented in blue (resp. red). The nullclines of the FitzHugh-Nagumo system (1.3) are represented in dashed lines.

Our aim is to analyse the joint influence of noise and interaction on the emergence of collective periodic behaviors for such a system of coupled excitable units. The point of the paper is to address this issue at the level of (1.1), that is for an infinite population $n = \infty$. Note that this problem can be equivalently considered at the level of the Fokker-Planck PDE associated to (1.1): the law $\mu_t = \mathcal{L}(X_t, Y_t)$ of the McKean-Vlasov process (1.1) is a weak solution to the following nonlinear kinetic Fokker-Planck PDE

$$\begin{aligned} \partial_t \mu_t = \sigma^2 \partial_{x^2}^2 \mu_t - \partial_x \left(\left[\delta \left(x - \frac{x^3}{3} - y \right) - K \left(x - \int_{\mathbb{R}^2} z_1 \mu_t(dz_1, dz_2) \right) \right] \mu_t \right) \\ - \frac{\delta}{c} \partial_y ((x + a - by) \mu_t), \quad t \geq 0, \quad (1.5) \end{aligned}$$

whose solution $t \mapsto \mu_t(dx, dy)$ takes its values in the set of probability measures on \mathbb{R}^2 . Equivalently the unique solution μ to (1.5) such that $\mu|_{t=0} = \mu_0$ is the law of the process (1.1) with $\mu_0 = \mathcal{L}(X_0)$. Well-posedness results concerning both (1.5) and (1.1) (in appropriate L^2 -spaces with exponential weights) are addressed in [23, 21].

Remark 1.1. *Observe here that we have in (1.1), (1.2) and (1.5) an interplay between three parameters: the strength of interaction $K > 0$, the intensity of noise $\sigma > 0$ and the scaling parameter $\delta > 0$ of the local dynamics. In fact, a simple time change in (1.1) shows that only two of these parameters are really relevant: we analyse below the long-time dynamics of (1.1) in terms of δ and $\frac{\sigma^2}{K}$.*

Our aim is to analyse the emergence of synchronicity in the mean-field system (1.1) (or equivalently (1.5)) under the joint influence of noise and interaction: the main result of the paper (see Theorem 1.4 below) concerns the existence of a stable invariant cycle for (1.5) in a regime where the interaction and noise are nontrivial and large w.r.t. to the intrinsic dynamics :

$$0 < \frac{\sigma^2}{K} < \infty \text{ and } \delta \ll 1. \quad (1.6)$$

The first condition of (1.6) is a natural regime for the emergence of collective oscillations for (1.1): informally, when $K = 0$ (that is in absence of interaction) for every isolated units (1.3) in an excitable state (first frame of Figure 1), the addition of noise alone make them leave the resting state (leading to a variety of uncorrelated stochastic dynamical patterns, e.g. canard-type excursion, mixed-mode oscillations, etc., see [19, 4] for further references). It is only when one adds further some nontrivial interaction $K > 0$ that these excursions may happen collectively, leading in the $n \rightarrow \infty$ limit to global oscillations of the system. Note that our main result is sufficiently versatile to track carefully how much noise (for a given K) one has to put in the system in order to see oscillations: we refer to Section 1.4 where we describe bifurcations of (1.1) in terms of σ (from $\sigma = 0$ to $\sigma \rightarrow \infty$).

Different asymptotics have been considered in previous works for the same model. We refer to Section 1.6 below for more details.

1.3. Slow-fast dynamics approach. The second hypothesis $\delta \ll 1$ in (1.6) comes from the fact that our approach relies on a perturbative slow-fast analysis. We prove namely that when $\delta \ll 1$, the system (1.1) admits solutions that are close to having Gaussian distributions. We present here the heuristic of this reduction, which will be made rigorous in the next sections.

For the variable X_t in (1.1), when δ is small (and K and σ remain of order 1), the interaction and noise terms constitute a fast part of the dynamics, while the intrinsic dynamics term δF constitutes a slow one. When $\delta = 0$, the fast dynamics part of (1.1) simply reduces an Ornstein Ulhenbeck process (of constant expectation) with Gaussian invariant measure. Hence, when δ is now positive but small, it is natural to approximate at first order in δ , the distribution of X_t by a Gaussian distribution $\mathcal{N}(\mathbf{E}[X_t], \sigma^2/K)$, where $\mathbf{E}[X_t]$ evolves slowly in time.

Now if X_t is at first order Gaussian, so is Y_t since its dynamics is linear. So, at first order in δ , (X_t, Y_t) should have a Gaussian distribution $\mathcal{N}(m_t, \Gamma_\delta)$, where Γ_δ is a symmetric covariance matrix, and $m_t = (\mathbf{E}[X_t], \mathbf{E}[Y_t])$ satisfies

$$\dot{m}_t = \delta \begin{pmatrix} \mathbf{E}[X_t] - \frac{\mathbf{E}[X_t^3]}{3} - \mathbf{E}[Y_t] \\ \frac{1}{c} (\mathbf{E}[X_t] + a - b\mathbf{E}[Y_t]) \end{pmatrix}. \quad (1.7)$$

Considering that $X_t \approx \mathcal{N}(x_t, \sigma^2/K)$, we obtain $\mathbf{E}[X_t^3] \approx \mathbf{E}[X_t]^3 + 3\frac{\sigma^2}{K}$, which leads to the approximation

$$\dot{m}_t \approx \delta F_{\frac{\sigma^2}{K}}(m_t), \quad (1.8)$$

with

$$F_u(x, y) := \begin{pmatrix} f_u(x, y) \\ g(x, y) \end{pmatrix} := \begin{pmatrix} (1-u)x - \frac{x^3}{3} - y \\ \frac{1}{c}(x+a-by) \end{pmatrix}, \quad x, y \in \mathbb{R}. \quad (1.9)$$

Note that (1.9) is once again of FitzHugh-Nagumo type, only modified by the prefactor $(1-u)$ in front of the x variable.

To compute Γ_δ , we denote $Z_t = (Z_t^x, Z_t^y)$ a first order approximation of the centered process $(X_t - \mathbf{E}[X_t], Y_t - \mathbf{E}[Y_t])$, defined by the following system of equations:

$$\begin{cases} dZ_t^x = -KZ_t^x dt + \sqrt{2}\sigma dB_t \\ dZ_t^y = \frac{\delta}{c}(Z_t^x - bZ_t^y) dt \end{cases}. \quad (1.10)$$

Remark that due to the fact that this dynamics is linear, if the distribution of Z_0 is a Gaussian, then the distribution of Z_t remains Gaussian, and straightforward calculations lead to

$$\frac{d}{dt} \mathbf{Var}[Z_t^y] = \frac{2\delta}{c} \mathbf{Cov}[Z_t^x, Z_t^y] - \frac{2b\delta}{c} \mathbf{Var}[Z_t^y], \quad (1.11)$$

$$\frac{d}{dt} \mathbf{Cov}[Z_t^x, Z_t^y] = \frac{\delta}{c} \mathbf{Var}[Z_t^x] - \left(K + \frac{b\delta}{c}\right) \mathbf{Cov}[Z_t^x, Z_t^y]. \quad (1.12)$$

The equilibrium solution of this system of equations is given by $\mathbf{Var}[Z_t^x] = \frac{\sigma^2}{K}$, $\mathbf{Var}[Z_t^y] = \frac{\sigma^2}{K} \frac{\delta}{b(Kc+b\delta)}$ and $\mathbf{Cov}[Z_t^x, Z_t^y] = \frac{\sigma^2}{K} \frac{\delta}{Kc+b\delta}$, and we thus define

$$\Gamma_\delta = \frac{\sigma^2}{K} \begin{pmatrix} 1 & \frac{\delta}{Kc+b\delta} \\ \frac{\delta}{Kc+b\delta} & \frac{\delta}{b(Kc+b\delta)} \end{pmatrix}. \quad (1.13)$$

Remark 1.2. *A similar slow-fast analysis was made in [21] for the elliptic case (that is with nontrivial interaction $K_2 > 0$ and noise $\sigma_2 > 0$ on the Y -variable)*

$$\begin{cases} dX_t = \delta f(X_t, Y_t) dt - K_1(X_t - \mathbf{E}[X_t]) dt + \sqrt{2}\sigma_1 dB_{1,t} \\ dY_t = \delta g(X_t, Y_t) dt - K_2(Y_t - \mathbf{E}[Y_t]) dt + \sqrt{2}\sigma_2 dB_{2,t} \end{cases}, \quad (1.14)$$

where $K_1, K_2, \sigma_1, \sigma_2 > 0$ and $B_{1,t}, B_{2,t}$ independent. For this elliptic case both X_t and Y_t have fast terms in their dynamics, and thus the distribution of (X_t, Y_t) is at first order a

Gaussian $\mathcal{N}(m_t, \Gamma)$, with $\Gamma = \begin{pmatrix} \frac{\sigma_1^2}{K_1} & 0 \\ 0 & \frac{\sigma_2^2}{K_2} \end{pmatrix}$ and $\dot{m}_t \approx F_{\frac{\sigma_2^2}{K_1}}(m_t)$, with the same function

F_u , defined by (1.9). In particular, the fact that $F_{\frac{\sigma_2^2}{K_1}}$ does not depend on (K_2, σ_2) in the

elliptic case is again a strong argument in favor of the validity of the approximation (1.8) in the kinetic case. However, it is not possible to make $K_2, \sigma_2 \rightarrow 0$ in the arguments of [21], as they rely strongly on the non-degeneracy of the noise on both variables. Secondly, contrary to [21], where the slow-fast reduction is based on the geometric properties of the PDE (1.5) (in a suitable L^2 -space with exponential weights), we focus mostly here on the properties of the system (1.1), relying on Wasserstein type estimates.

1.4. Collective oscillations in the FitzHugh-Nagumo model under noise and interaction. Supposing that the analysis made above is rigorous, i.e. that the coupled mean-field system (1.1) admits solutions (X_t, Y_t) that are at first order Gaussian distribution $\mathcal{N}(m_t, \Gamma_\delta)$, then (at least at first order) the analysis of (1.1) can be reduced to the analysis the dynamics of its expectation m_t , for which we provided the first order approximation (1.8). Since (1.8) is nothing else than another FitzHugh-Nagumo system

(slowed-down by a factor δ), everything reads now in terms of the bifurcations of the system (recall (1.9))

$$\begin{pmatrix} \dot{x}_t \\ \dot{y}_t \end{pmatrix} = F_u(x_t, y_t) \quad (1.15)$$

as u increases from $u = 0$ (isolated system (1.3)) to $u = \frac{\sigma^2}{K} > 0$ (coupled system with noise and interaction). In particular, as we will see below, a crucial observation is that it is possible to choose carefully the parameters a, b, c so that (1.15) has a unique stable state for $u = 0$, whereas the same system exhibits oscillations for values of u chosen in a bounded interval (see Figure 2 below): this is the signature of the emergence of periodic dynamics due to noise and interaction in (1.1).

The analysis of the bifurcations of the system (1.15) was already made in [21], Section 3, since it is also the dynamics obtained by slow-fast reduction of the elliptic case. We present this analysis here, for the sake of completeness. With no loss of generality (Remark 1.1), we can suppose $K = 1$. It is then possible to read from Figure 2 the behavior of the system (1.1) as the noise intensity $u = \sigma^2$ increases: in absence of noise ($u = 0$), each neuron has a fixed-point dynamics. As u increases, we observe the emergence (through a saddle-node bifurcation of cycles) of a stable cycle coexisting with an unstable cycle and stable fixed-point (see the case $u = 0.086$) for (1.15). As u increases further, the unstable cycle collides with the stable fixed-point, resulting in only one stable limit cycle coexisting with an unstable fixed-point (see the case $u = 0.2$). For large noise, oscillations disappear (see $u = 0.8$). We refer to [21], Section 3.4 for more details on these transitions, in terms of Hopf bifurcations, and also for a study of the bistable case.

As an illustration of the accuracy of this analysis for the particle system (1.2), we reproduce in Figure 3 the dynamics of the empirical measure of (1.2), for $\delta = 0.2$, $a = \frac{1}{3}$, $b = 1$, $c = 10$, $K = 1$, $\sigma^2 = 0.2$ (which corresponds to $u = 0.2$ in Figure 2) and $n = 10^5$: the mean-value of the empirical density follows precisely the limit cycle given by (1.15) in the case $u = 0.2$. One notable difference between the present simulation in the kinetic case and the simulations in [21] in the elliptic case (see [21], Figure 7) is the shape of the empirical density in both cases: due to the absence of noise on the y -coordinate, the variance along the y -direction is here significantly smaller than in the elliptic case.

1.5. Main results. In the remaining of the paper, we fix the parameters (a, b, c, σ, K) such that the system

$$\frac{1}{\delta} \begin{pmatrix} \dot{x}_t \\ \dot{y}_t \end{pmatrix} = F_{\frac{\sigma^2}{K}}(x_t, y_t), \quad (1.16)$$

admits a limit cycle (recall that this is possible even if it is not the case for the system (1.3), see Section 1.4 and [21], Section 3.4).

Remark 1.3. *Note that the existence of such limit cycle does not depend on δ : if, in the case $\delta = 1$, we denote this stable periodic solution by $(\gamma_t)_{t \in [0, T_\gamma]}$ (T_γ being the period of the limit cycle for $\delta = 1$), the behavior of (1.16) for general $\delta > 0$ may be deduced by a simple time change: the corresponding periodic orbit becomes $(\gamma_t^\delta) := (\gamma_{\delta t})_{t \in [0, T_\gamma/\delta]}$, with period $\frac{T_\gamma}{\delta}$. When no confusion is possible, we will drop the superscript δ in γ^δ in the following. We adopt also the definition*

$$\mathbb{S}_\delta := \mathbb{R}/((T_\gamma/\delta)\mathbb{Z}) \quad (1.17)$$




FIGURE 2. Phase diagrams for the system (1.15) for parameters $a = \frac{1}{3}$, $b = 1$, $c = 10$ for different choices of u . Stable (resp. unstable) points and limit cycles are represented in blue (resp. red). The nullclines of the FitzHugh-Nagumo system (1.15) are represented in dashed lines.

endowed with the quotient topology induced by the euclidean distance on \mathbb{R} . Such topology can be generated by the following metric: for all $\bar{\varphi} \equiv \varphi \left[\frac{T_\gamma}{\delta} \right]$, $\bar{\psi} \equiv \psi \left[\frac{T_\gamma}{\delta} \right]$,

$$d_{\mathbb{S}}(\bar{\varphi}, \bar{\psi}) := \max \left(|\varphi - \psi|, \frac{T_\gamma}{\delta} - |\varphi - \psi| \right). \quad (1.18)$$

We denote q_m the distribution given by

$$q_m(z) = q_m^\delta(z) := \frac{1}{2\pi \det(\Gamma_\delta)} \exp \left(\frac{1}{2} (z - m) \cdot \Gamma_\delta^{-1} (z - m) \right), \quad z \in \mathbb{R}^2, \quad (1.19)$$

and

$$\mathcal{G}_\delta := \left\{ q_{\gamma_t}^\delta : t \geq 0 \right\}. \quad (1.20)$$

Below, W_2 is a distance of Wasserstein-type that is precisely defined in Definition 2.2. The main result of the paper is the following:

Theorem 1.4. *Choose parameters a, b, c, K and σ^2 such that (1.16) admits an stable limit cycle. Then there exists a $\delta_c > 0$ such that for all $\delta \leq \delta_c$ there exists a periodic solution ν_t^{per} to (1.5), defining an invariant cycle \mathcal{C}_δ which satisfies $\sup_t \text{dist}_{W_2}(\nu_t^{per}, \mathcal{G}_\delta) = O(\delta)$.*

Moreover there exist positive constants C_1 and C_2 that do not depend on δ , a positive constant $C(\delta)$ and a positive rate $\lambda(\delta)$ such that if $\mu_0 \in \mathcal{P}_2$ satisfies

$$\int_{\mathbb{R}^2} z^6 d\mu_0(z) \leq C_1, \quad \text{and} \quad \text{dist}_{W_2}(\mu_0, \mathcal{C}_\delta) \leq C_2\delta, \quad (1.21)$$




FIGURE 3. Time evolution of the empirical density of the particle system (1.2), in the case $\delta = 0.2$, $a = \frac{1}{3}$, $b = 1$, $c = 10$, $K = 1$, $\sigma^2 = 0.2$ and $n = 100000$. The corresponding evolution of the empirical mean-value is represented.

then for $t \mapsto \mu_t$ the solution to (1.5) with initial condition μ_0 we have:

$$\text{dist}_{W_2}(\mu_t, \mathcal{C}_\delta) \leq C(\delta) e^{-\lambda(\delta)t} \text{dist}_{W_2}(\mu_0, \mathcal{C}_\delta). \quad (1.22)$$

Remark 1.5. The constants $\lambda(\delta)$ and $C(\delta)$ obtained in the proof of Theorem 1.4 satisfy

$$\lambda(\delta) \xrightarrow{\delta \rightarrow 0} \min \left\{ \lambda, \frac{b}{c} \right\}, \quad \text{and} \quad C(\delta) \xrightarrow{\delta \rightarrow 0} \infty, \quad (1.23)$$

where λ is the exponential rate of attraction of the limit cycle of (1.16), in the case $\delta = 1$.

Remark 1.6. We focus in this work on the proof of the existence of a stable cycle for (1.5) when (1.16) admits one, but it is clear that one can prove the existence of a stable fixed point for (1.5) when (1.16) admits one by following the same arguments (in fact simpler arguments, no need of Floquet theory in that case).

1.6. Comments and existing literature.

On the mean-field FitzHugh-Nagumo model. The structure of the proof we provide in this paper, relying on a fixed-point theorem, is inspired from the classical proofs of the theory of persistence of normally hyperbolic invariant manifold [12, 13, 18, 3, 28, 32], which we could not apply directly here due to the singularity of our problem (when $\delta = 0$ the dynamics of Y_t is trivial and thus there can not be any stable compact invariant manifold for (1.5)).

Several recent other works have analyzed the long-time behavior of the FitzHugh-Nagumo Fokker-Planck PDE (1.5). The paper [23] considers the situation of small interaction (i.e. $\delta > 0$ and $\sigma > 0$ fixed with $K \rightarrow 0$). Along with some well-posedness

estimates concerning (1.5), the main result of [23] concerns the existence of stationary states for (1.5) in the limit of small interaction. The case where $K \rightarrow \infty$ with δ and σ fixed is analyzed in [24]. The authors prove concentration in large time around singular solutions (clamping) in such a case. In a regime similar to ours, one should finally mention the recent [31] which analyzes the possible microscopic mechanisms responsible for the emergence of collective oscillations.

As already said, this paper addresses the case of an interaction modeling electrical synapses. A common framework in neuronal models concerns interactions through chemical synapses [6], which is not covered by this work. The question of the possibility of extension of the present results to this case is open. In this direction, a recent work [7] addresses synchronization issues of interacting neurons with Hodgkin-Huxley dynamics with chemical synapses.

On the dynamics of the particle system (1.2). Obtaining a rigorous proof of the existence of periodic behaviors in infinite population limit of mean-field interacting particle models is a problem that has been studied in different situations, as chemical reaction models, rotors models, spin models, Hawkes processes, etc., see for example [26, 27, 16, 15, 17, 9, 8, 10], each proof relying, as in the present work, on a drastic phase reduction of the model.

Transposing the dynamics of the nonlinear process (1.1) to the dynamics of the particle system (1.2) is a nontrivial task, that we do not address here (see nonetheless Figure 3 for a numerical evidence of the accuracy of the present analysis w.r.t. the particle system (1.2)). A standard way to couple i.i.d. copies $(X_t^{(i)}, Y_t^{(i)})$ ($i = 1, \dots, n$) of (1.1) with the particles $(X_{i,t}, Y_{i,t})$ in (1.2) is to choose identical initial condition and noise B_i so that the following standard propagation of chaos estimates [29, 23, 22, 21] holds:

$$\sup_{i=1, \dots, n} \mathbf{E} \left[\sup_{0 \leq t \leq T} \left| (X_{i,t}, Y_{i,t}) - (X_t^{(i)}, Y_t^{(i)}) \right|^2 \right] \leq \frac{C e^{CT}}{n}. \quad (1.24)$$

This propagation of chaos result may be equivalently expressed in terms of the convergence of the empirical measure $\mu_{n,t} := \frac{1}{n} \sum_{j=1}^n \delta_{(X_{j,t}, Y_{j,t})}$ of the particle system (1.2) to the solution of (1.5), on any time interval $[0, T]$. From (1.24), we see that (1.1) gives a correct approximation of (1.2) at least up to times T of order $c \ln n$, for $c > 0$ small enough. The question of the relevance of this mean-field approximation for times $T \gg \ln n$ is a long-standing issue in the literature (see e.g. [5, 11] and references therein). In the case of collective periodic behaviors that is of particular interest in neuroscience, one should mention in particular [5, 20] (for phase oscillators), [10] (for point processes with Hawkes dynamics) and [1] (for Curie-Weiss dynamics modeling social interactions).

1.7. Organization of the paper. In Section 2, we prove some controls on the moments of (1.1) as well as some estimates of proximity of solutions of (1.5) to the Gaussian manifold (1.20). Section 3 gathers the main estimates (relying in particular on Floquet theory) concerning (1.1). The main result (Theorem 1.4) is proven in Section 4. A technical lemma is postponed to the appendix.

2. MOMENT ESTIMATES AND PROXIMITY TO GAUSSIAN DISTRIBUTIONS

Take $R_0 > 0$ such that the periodic solution γ^δ of (1.16) is strictly included in the open euclidean ball $B(0, R_0)$. Note that, by Remark 1.3, R_0 does not depend on δ . For any initial condition to (1.1) such that $(\mathbf{E}[X_0], \mathbf{E}[Y_0]) \in B(0, R_0)$, define the exit time

$$T_e(R_0) = T_e^\delta(R_0) = \inf\{t \geq 0 : (\mathbf{E}[X_t], \mathbf{E}[Y_t]) \notin B(0, R_0)\}. \quad (2.1)$$

Here, we note that $T_e(R_0)$ is independent of the particular coupling of (X_0, Y_0) , provided its marginals are fixed.

Lemma 2.1. *There exist positive constants δ_0, κ_0^x and κ_0^y such that if $\delta \leq \delta_0$ the following is true: for any initial condition (X_0, Y_0) to (1.1) such that*

$$\mathbf{E}[X_0^6] \leq \kappa_0^x, \quad \text{and} \quad \mathbf{E}[Y_0^6] \leq \kappa_0^y, \quad (2.2)$$

then the solution (X_t, Y_t) of (1.1) satisfies

$$\sup_{t \in [0, T_e(R_0)]} \mathbf{E}[X_t^6] \leq \kappa_0^x, \quad \text{and} \quad \sup_{t \in [0, T_e(R_0)]} \mathbf{E}[Y_t^6] \leq \kappa_0^y. \quad (2.3)$$

Proof. For $n > 0$ denote the stopping time $\tau_n = \inf\{t \geq 0 : X_t^2 + Y_t^2 = n^2\}$, and denote $X_t^{(n)} = X_{t \wedge \tau_n}$ and $Y_t^{(n)} = Y_{t \wedge \tau_n}$. By Itô formula we get

$$\begin{aligned} \frac{1}{6}(X_t^{(n)})^6 &= \frac{1}{6}(X_0^{(n)})^6 + \int_0^{t \wedge \tau_n} \left[\delta (X_s^{(n)})^5 \left(X_s^{(n)} - \frac{(X_s^{(n)})^3}{3} - Y_s^{(n)} \right) + K \mathbf{E}[X_s] (X_s^{(n)})^5 \right. \\ &\quad \left. - K (X_s^{(n)})^6 + 5\sigma^2 (X_s^{(n)})^2 \right] ds + \sigma \int_0^{t \wedge \tau_n} (X_s^{(n)})^5 dB_s. \end{aligned} \quad (2.4)$$

Using the following inequalities, for positive constant c_1, \dots, c_4 that we do not give explicitly to keep notations simple,

$$\begin{aligned} x^6 - \frac{x^8}{3} &\leq c_1 - \frac{x^8}{4}, \quad x^5 y \leq \frac{x^8}{4} + c_2 y^6, \quad K x^5 \mathbf{E}[X_t] - K x^6 \leq c_3 \mathbf{E}[X_t] - \frac{K x^6}{3}, \\ 5\sigma^2 x^4 - \frac{2K x^6}{3} &\leq c_4 - \frac{K x^6}{3}, \end{aligned} \quad (2.5)$$

we obtain,

$$\frac{1}{4} \frac{d}{dt} \mathbf{E}[(X_t^{(n)})^6] \leq c_1 \delta + c_4 + c_3 \mathbf{E}[X_t] + c_2 \delta \mathbf{E}[(Y_t^{(n)})^6] - \frac{K}{3} \mathbf{E}[(X_t^{(n)})^6]. \quad (2.6)$$

On the other hand,

$$\frac{c}{6\delta} (Y_t^{(n)})^6 = \frac{c}{6\delta} (Y_0^{(n)})^6 + \int_0^{\tau_n} (Y_s^{(n)})^5 \left(X_s^{(n)} + a - b Y_s^{(n)} \right) ds, \quad (2.7)$$

and using the inequalities, for positive constants c_6, c_7 ,

$$x y^5 \leq c_6 x^6 + \frac{b}{3} y^6, \quad \text{and} \quad a y^5 - \frac{2b}{3} y^6 \leq c_7 - \frac{b}{3} y^6, \quad (2.8)$$

we get

$$\frac{c}{6\delta} \frac{d}{dt} \mathbf{E}[(Y_t^{(n)})^6] \leq c_7 + c_6 \mathbf{E}[(X_t^{(n)})^6] - \frac{b}{3} \mathbf{E}[(Y_t^{(n)})^6]. \quad (2.9)$$

Using (2.6) and (2.9) and the Grönwall inequality we deduce that in fact $\mathbf{E}[X_t^6] + \mathbf{E}[Y_t^6] < \infty$ and the estimates made above are in fact valid for $n = \infty$. In particular, for $t \in [0, T_e(R_0)]$ we have

$$\frac{1}{4} \frac{d}{dt} \mathbf{E}[X_t^6] \leq c_1 \delta + c_4 + c_3 R_0 + c_2 \delta \mathbf{E}[Y_t^6] - \frac{K}{3} \mathbf{E}[X_t^6], \quad (2.10)$$

$$\frac{c}{6\delta} \frac{d}{dt} \mathbf{E}[Y_t^6] \leq c_7 + c_6 \mathbf{E}[X_t^6] - \frac{b}{3} \mathbf{E}[Y_t^6]. \quad (2.11)$$

Define now $\kappa_0^x = \frac{4}{K}(c_1 + c_3 R_0 + c_2)$, and $\kappa_0^y = \frac{4}{b}(c_7 + c_6 \kappa_0^x)$ and

$$t_0 = \inf\{t \geq 0 : \mathbf{E}[(X_t^{(n)})^6] \geq \kappa_0^x \text{ or } \mathbf{E}[Y_t^6] \geq \kappa_0^y\}. \quad (2.12)$$

For all $t \in [0, t_0 \wedge T_e(R_0)]$ supposing that $\mathbf{E}[X_0^6] \leq \kappa_0$ and $\mathbf{E}[Y_0^6] \leq \kappa_0$, we get from (2.10) and (2.11) that

$$\mathbf{E}[X_t^6] \leq \max \left\{ \kappa_0^x, \frac{3}{K} (c_1 \delta + c_4 + c_3 R_0 + c_2 \delta \kappa_0^y) \right\}, \quad (2.13)$$

$$\mathbf{E}[Y_t^6] \leq \max \left\{ \kappa_0^y, \frac{3}{b} (c_7 + c_6 \kappa_0^x) \right\}. \quad (2.14)$$

and these maxima are equal to κ_0^x and κ_0^y respectively as soon as $\delta \leq \min\{1, 1/R_0\}$. Moreover it is clear with this choice of κ_0^x that if X_t satisfies $\mathbf{E}[X_t^6] = \kappa_0^x$, then $\frac{d}{dt} \mathbf{E}[X_t^6] < 0$, and the same applies for Y_t . So $t_0 \wedge T_e(R_0) = T_e(R_0)$, which concludes the proof. \square

Definition 2.2. For a probability distribution ν on \mathbb{R}^2 , $m = \int_{\mathbb{R}^2} z d\nu(z)$ its expectation, we denote $\tilde{\nu}$ the centered version of ν : $\int_{\mathbb{R}^2} \phi(z) d\nu(z) = \int_{\mathbb{R}^2} \phi(z + m) d\tilde{\nu}(z)$ (for all test function ϕ). For two distributions ν_1 and ν_2 on \mathbb{R}^2 , denote by $\mathcal{C}(\nu_1, \nu_2)$ the set of all couplings π of ν_1 and ν_2 , that is the set of all probability measures $\pi(d(x_1, y_1), d(x_2, y_2))$ on $\mathbb{R}^2 \times \mathbb{R}^2$ with marginals $\nu_1(d(x_1, y_1))$ and $\nu_2(d(x_2, y_2))$. If $\pi \in \mathcal{C}(\nu_1, \nu_2)$, we denote by $\tilde{\pi} \in \mathcal{C}(\tilde{\nu}_1, \tilde{\nu}_2)$ the corresponding coupling of the centered measures:

$$\int_{\mathbb{R}^2 \times \mathbb{R}^2} \phi(z_1, z_2) \pi(dz_1, dz_2) = \int_{\mathbb{R}^2 \times \mathbb{R}^2} \phi(z_1 + m_1, z_2 + m_2) \tilde{\pi}(dz_1, dz_2)$$

For $\beta \in (0, 1)$, we consider the following Wasserstein-type distance $W = W(\delta, b, \beta)$ on $\mathcal{P}(\mathbb{R}^2)$,

$$W(\nu_1, \nu_2) := \inf \{ \Lambda(\pi), \pi \in \mathcal{C}(\nu_1, \nu_2) \}, \quad (2.15)$$

where for a fixed $\pi \in \mathcal{C}(\nu_1, \nu_2)$, $z_i = (x_i, y_i) \in \mathbb{R}^2$, $i = 1, 2$

$$\Lambda(\pi) := \max \left(\delta^\beta \left| \int_{\mathbb{R}^2 \times \mathbb{R}^2} (z_1 - z_2) \pi(dz_1, dz_2) \right|, \sqrt{\int_{\mathbb{R}^2 \times \mathbb{R}^2} |x_1 - x_2|^2 \tilde{\pi}(dz_1, dz_2)}, b \sqrt{\int_{\mathbb{R}^2 \times \mathbb{R}^2} |y_1 - y_2|^2 \tilde{\pi}(dz_1, dz_2)} \right). \quad (2.16)$$

The factor δ^β in the definition of the distance W (and later the definition of the distances W_θ , see section 4) will be useful to obtain the contraction property, at the end of the proof of Lemma 4.2.

Remark 2.3. (1) The definition (2.16) is of course equivalent to

$$\Lambda(\pi) = \max \left(\delta^\beta |\mathbf{E}[(X_1, Y_1)] - \mathbf{E}[(X_2, Y_2)]|, \sqrt{\mathbf{E}[|\tilde{X}_1 - \tilde{X}_2|^2]}, b \sqrt{\mathbf{E}[|\tilde{Y}_1 - \tilde{Y}_2|^2]} \right), \quad (2.17)$$

where $\{(X_1, Y_1), (X_2, Y_2)\} \sim \pi$, for $\tilde{X} = X - \mathbf{E}[X]$.

(2) Note that the first term in (2.16) is independent of the coupling $\pi \in \mathcal{C}(\nu_1, \nu_2)$, so that we can also write

$$W(\nu_1, \nu_2) = \max \left\{ \delta^\beta \left| \int_{\mathbb{R}^2} z \nu_1(dz) - \int_{\mathbb{R}^2} z \nu_2(dz) \right|, W(\tilde{\nu}_1, \tilde{\nu}_2) \right\}. \quad (2.18)$$

(3) Remark that W is equivalent to the standard Wasserstein-2 distance W_2 : for δ small enough

$$\min(1, b^{-1}) W \leq W_2 \leq 3\delta^{-\beta} W. \quad (2.19)$$

Denote by $(\tilde{X}_t, \tilde{Y}_t)$ the centered version of (X_t, Y_t) , that satisfies

$$\begin{cases} d\tilde{X}_t = \delta \left(X_t - \frac{X_t^3}{3} - Y_t \right) dt - \dot{x}_t dt - K \tilde{X}_t dt + \sqrt{2}\sigma dB_t \\ d\tilde{Y}_t = \frac{\delta}{c} \left(\tilde{X}_t - b\tilde{Y}_t \right) dt \end{cases}, \quad (2.20)$$

where $m_t = (x_t, y_t) = (\mathbf{E}[X_t], \mathbf{E}[Y_t])$. Observe that, by (1.1), m_t solves

$$\begin{cases} \dot{x}_t = \delta \left(x_t - \frac{\int x^3 d\mu_t}{3} - y_t \right) \\ \dot{y}_t = \frac{\delta}{c} (x_t + a - by_t) \end{cases}. \quad (2.21)$$

Lemma 2.4. *There exists positive constants δ_1 and κ_1 such that, for all initial condition μ_0 such that $m_0 = (x_0, y_0) = (\int x \mu_0(dx, dy), \int y \mu_0(dx, dy)) \in B(0, R_0)$ and $\int x^6 \mu_0(dx, dy) \leq \kappa_0^x$ and $\int y^6 \mu_0(dx, dy) \leq \kappa_0^y$ (recall Lemma 2.1), the following is true: if $\delta \leq \delta_1$ and $W(\tilde{\mu}_0, q_0) \leq \kappa_1 \delta$, then*

$$\sup_{t \in [0, T_e(R_0)]} W(\tilde{\mu}_t, q_0) \leq \kappa_1 \delta. \quad (2.22)$$

Proof. We suppose that $\delta \leq \delta_0$ where δ_0 is given by Lemma 2.1. For an arbitrary $\varepsilon > 0$, consider a coupling $\tilde{\pi}_0 \in \mathcal{C}(\tilde{\mu}_0, q_0)$ such that

$$\Lambda(\tilde{\pi}_0)^2 < W(\tilde{\mu}_0, q_0)^2 + \varepsilon. \quad (2.23)$$

In the following, we consider $\left\{ (\tilde{X}_0, \tilde{Y}_0), (Z_0^x, Z_0^y) \right\}$ with law $\tilde{\pi}_0$. For this initial condition, we consider both (X_t, Y_t) solution to (1.1) with initial condition $(X_0, Y_0) = (\tilde{X}_0 + x_0, \tilde{Y}_0 + y_0)$ and $(\tilde{X}_t, \tilde{Y}_t)$, solution to (2.20) with initial condition $(\tilde{X}_0, \tilde{Y}_0)$. Consider also the process (Z_t^x, Z_t^y) solution to the system (1.10) with initial condition (Z_0^x, Z_0^y) with distribution q_0 . The calculations made in Section 1.3 show that (Z_t^x, Z_t^y) has also distribution q_0 , so that, by construction, the law $\tilde{\pi}_t$ of $\left\{ (\tilde{X}_t, \tilde{Y}_t), (Z_t^x, Z_t^y) \right\}$ belongs to $\mathcal{C}(\tilde{\mu}_t, q_0)$. We have

$$\frac{1}{2} d(\tilde{X}_t - Z_t^x)^2 = -K(X_t - Z_t^x)^2 dt + (\tilde{X}_t - Z_t^x) \left(\delta \left(X_t - \frac{X_t^3}{3} - Y_t \right) - \dot{x}_t \right) dt. \quad (2.24)$$

By Lemma 2.1 and (2.21), there exists a positive constant C_0 such that for all $t \in [0, T_e(R_0)]$,

$$\mathbf{E} \left[\left(\delta \left(X_t - \frac{X_t^3}{3} - Y_t \right) - \dot{x}_t \right)^2 \right]^{\frac{1}{2}} \leq C_0 \delta. \quad (2.25)$$

Note that the constant C_0 only depends on the parameters (a, b, c) of the model and on R_0 through the constants κ_0^x and κ_0^y defined in Lemma 2.1. In particular, C_0 does not depend on ε . We then have, for all $t \in [0, T_e(R_0)]$,

$$\frac{1}{2} \frac{d}{dt} \mathbf{E}[(\tilde{X}_t - Z_t^x)^2] \leq C_0 \delta (\mathbf{E}[(\tilde{X}_t - Z_t^x)^2])^{\frac{1}{2}} - K \mathbf{E}[(\tilde{X}_t - Z_t^x)^2]. \quad (2.26)$$

If now we choose the constant κ_1 to be $\kappa_1 := \frac{C_0}{K}$ and assume that $W(\tilde{\mu}_0, q_0) \leq \kappa_1 \delta$, then by (2.23), we have $\mathbf{E}[(\tilde{X}_0 - Z_0^x)^2] \leq \frac{C_0^2}{K^2} \delta^2 + \varepsilon$. Consequently, by Lemma A.1,

$$\sup_{t \in [0, T_e(R_0)]} \mathbf{E}[(\tilde{X}_t - Z_t^x)^2] \leq \frac{C_0^2}{K^2} \delta^2 + \varepsilon. \quad (2.27)$$

In a same way,

$$\frac{c}{2\delta} d(\tilde{Y}_t - Z_t^y)^2 = (\tilde{Y}_t - Z_t^y)(\tilde{X}_t - Z_t^x) dt - b(\tilde{Y}_t - Z_t^y)^2 dt, \quad (2.28)$$

and from the estimate (2.27) we have just obtained above, we get for $t \in [0, T_e(R_0)]$

$$\frac{c}{2\delta} \frac{d}{dt} \mathbf{E}[(\tilde{Y}_t - Z_t^y)^2] \leq \left(\frac{C_0^2 \delta^2}{K^2} + \varepsilon \right)^{\frac{1}{2}} (\mathbf{E}[(\tilde{Y}_t - Z_t^y)^2])^{\frac{1}{2}} - b \mathbf{E}[(\tilde{Y}_t - Z_t^y)^2]. \quad (2.29)$$

From $W(\tilde{\mu}_0, q_0) \leq \kappa_1 \delta$, we obtain that $\mathbf{E}[(\tilde{Y}_0 - Z_0^y)^2] \leq \frac{1}{b^2} \left(\frac{C_0^2}{K^2} \delta^2 + \varepsilon \right)$, so that by Lemma A.1, we have

$$\sup_{t \in [0, T_e(R_0)]} \mathbf{E}[(\tilde{Y}_t - Z_t^y)^2] \leq \frac{1}{b^2} \left(\frac{C_0^2}{K^2} \delta^2 + \varepsilon \right). \quad (2.30)$$

Estimates (2.27) and (2.30) are a fortiori true for the infimum over all couplings $W(\tilde{\mu}_t, q_0)$. Letting $\varepsilon \searrow 0$, this concludes the proof, taking $\delta_1 = \delta_0$. \square

3. FLOQUET THEORY AND CONTRACTION CLOSE TO γ

We introduce in this paragraph the minimal notions from Floquet theory [30] that are necessary for our purpose. For the simplicity of exposition, this is first done in the case $\delta = 1$ from which the general case $\delta > 0$ may be deduced up to minor modifications (see section 3.2 below).

3.1. The case $\delta = 1$. Recall the definition of F_u in (1.9). Denote by $\Pi(s, t)$ the principal matrix solution associated to the periodic solution $\gamma = \gamma^1$ of (1.16), when $\delta = 1$, that is the solution to

$$\partial_t \Pi(t, s) = DF_{\frac{\sigma^2}{K}}(\gamma_t) \Pi(t, s), \quad \Pi(s, s) = I_d. \quad (3.1)$$

$\Pi(t, s)$ is invertible, with

$$\Pi(t, s) \Pi(s, t) = I_d, \quad (3.2)$$

and satisfies (recall that $t \mapsto \gamma_t$ has period T_γ):

$$\Pi(t, u) \Pi(u, s) = \Pi(t, s), \quad \Pi(t + T_\gamma, s + T_\gamma) = \Pi(t, s). \quad (3.3)$$

Denote $Q(0)$ the matrix such that

$$\Pi(T_\gamma, 0) = e^{-T_\gamma Q(0)}. \quad (3.4)$$

Since

$$F_{\frac{\sigma^2}{K}}(\gamma_t) = \Pi(t, s) F_{\frac{\sigma^2}{K}}(\gamma_s), \quad (3.5)$$

one of the eigenvalues of $Q(s)$ is simply 0, with corresponding eigenvector $F_{\frac{\sigma^2}{K}}(\gamma_0)$. By hypothesis γ is stable and thus the other eigenvalue is positive (see [30], Chapter 12), we denote it λ and e_0 a corresponding eigenvector. Define $N(t, 0) = \Pi(t, 0) e^{tQ(0)}$, so that $P(t, 0) = N(t, 0) e^{-tQ(0)}$. $N(t, 0)$ is clearly periodic: $N(t + T_\gamma, 0) = N(t, 0)$.

We can now define

$$Q(s) = \Pi(s, 0) Q(0) \Pi(0, s), \quad \text{and} \quad N(t, s) = \Pi(t, t-s) N(t-s, 0) \Pi(0, s), \quad (3.6)$$

so that

$$\Pi(t, s) = N(t, s) e^{-(t-s)Q(s)}, \quad (3.7)$$

and N satisfies $N(t + T_\gamma, s) = N(t, s)$ and $Q(s)$ has the same spectral decomposition as $Q(0)$. In particular $e_s = N(s, 0) e_0$ is an eigenvector associated with the eigenvalue λ for $Q(s)$. It is easy to see that $Q(t) = \Pi(t, s) Q(s) \Pi(s, t)$ and $N(t, u) N(u, s) = N(t, s)$.

We consider p_s and p_s^\perp the linear form associated to the projection on $F_{\frac{\sigma^2}{K}}$ and e_s respectively, i.e. $p_s, p_s^\perp : \mathbb{R}^2 \rightarrow \mathbb{R}$ and satisfies $p_s(e_s) = p_s^\perp(F_{\frac{\sigma^2}{K}}(\gamma_s)) = 0$ and $p_s(F_{\frac{\sigma^2}{K}}(\gamma_s)) = p_s^\perp(e_s) = 1$. It is easy to see that $p_t^\perp(N(t, s)\cdot) = p_s^\perp(\cdot)$ and that, for all $t > s$,

$$\left| p_t^\perp(\Pi(t, s)u) \right| \leq e^{-\lambda(t-s)} p_s^\perp(u). \quad (3.8)$$

We consider also a constant C_Π such that $|\Pi(t, s)u| \leq C_\Pi|u|$ for all $t > s$.

For $\alpha > 0$ and all $s \in [0, T_\gamma)$, we define $E_s(\alpha)$ as

$$E_s(\alpha) := \{u \in \gamma_s + \text{span}(e_s) : \left| p_s^\perp(u - \gamma_s) \right| \leq \alpha\}. \quad (3.9)$$

3.2. The general case $\delta > 0$. All that have been exposed in the previous paragraph for the limit cycle γ^1 in the case $\delta = 1$ can be easily transposed for γ^δ when $\delta > 0$, with the following definitions: $\Pi^\delta(t, s) := \Pi^1(\delta t, \delta s)$, $Q^\delta(s) := \delta Q^1(\delta s)$, $N^\delta(t, s) = N^1(\delta t, \delta s)$, $e_s^\delta := e_{\delta s}^1$, $p_s^\delta := p_{\delta s}^1$, $p_s^{\delta, \perp} := p_{\delta s}^{1, \perp}$ and

$$E_s^\delta(\alpha) := \{u \in \gamma_s^\delta + \text{span}(e_s^\delta) : \left| p_s^{\delta, \perp}(u - \gamma_s^\delta) \right| \leq \alpha\}, \quad s \in [0, T_\gamma/\delta). \quad (3.10)$$

Note in particular the spectral gap λ when $\delta = 1$ is changed into $\delta\lambda$: (3.8) becomes

$$\left| p_t^{\delta, \perp}(\Pi^\delta(t, s)u) \right| \leq e^{-\lambda\delta(t-s)} p_s^{\delta, \perp}(u). \quad (3.11)$$

We now give a classical result of projection γ . For a similar result in a more general situation, see for example [28].

Lemma 3.1. *There exist $\alpha_0 > 0$ such that for all z in the α_0 -neighborhood of γ^δ , there exists a unique $\theta =: \text{proj}^\delta(z) \in \mathbb{S}_\delta$ such that $z \in E_\theta^\delta(1)$. Moreover, for all z, h such that z and $z + h$ are in the α_0 -neighborhood of γ^δ , if $\theta^\delta = \text{proj}^\delta(z)$,*

$$\text{proj}^\delta(z + h) = \theta^\delta + \frac{1}{\delta} \frac{p_{\theta^\delta}^\delta(h)}{1 + p_{\theta^\delta}^\delta \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_{\theta^\delta}^\delta) + \frac{1}{\delta} Q^\delta(\theta^\delta) \right) (z - \gamma_{\theta^\delta}^\delta) \right)} + \frac{1}{\delta} O(h^2), \quad (3.12)$$

where the rest $O(h^2)$ is uniform in $\delta \leq 1$. In particular, under the previous hypotheses, there exists a constant $C_{\text{proj}} > 0$, independent of δ , such that

$$d_{\mathbb{S}_\delta} \left(\text{proj}^\delta(z + h), \text{proj}^\delta(z) \right) \leq \frac{C_{\text{proj}}}{\delta} |h|. \quad (3.13)$$

Proof. Assume first that the result holds in the case $\delta = 1$. Then it is easy to see that the unique candidate for the projection in the case $\delta > 0$ is given by $\text{proj}^\delta(z) := \frac{\text{proj}^1(z)}{\delta} \in \mathbb{S}_\delta$. In particular, one deduces (3.12) and (3.13) from the case $\delta = 1$ and the change of variables formulas in § 3.2. We now prove the result for $\delta = 1$, which basically follows from the Implicit Functions Theorem. Let

$$(z, \theta) \in \mathbb{R}^2 \times \mathbb{S}_\delta \mapsto f(z, \theta) := p_\theta(z - \gamma_\theta). \quad (3.14)$$

Relying on the identities $p_\theta(z - \gamma_\theta) = p_0(N(0, \theta)(z - \gamma_\theta))$ and

$$\begin{aligned} \frac{d}{d\theta} N(0, \theta) &= \frac{d}{d\theta} (N(\theta, 0)^{-1}) = -N(0, \theta) \left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) N(\theta, 0) + N(\theta, 0) Q(0) \right) N(0, \theta) \\ &= -N(0, \theta) \left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) + Q(\theta) \right), \end{aligned} \quad (3.15)$$

we obtain

$$\begin{aligned} \partial_\theta f(z, \theta) &= -p_\theta \left(F_{\frac{\sigma^2}{k}}(\gamma_\theta) \right) - p_\theta \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) + Q(\theta) \right) (z - \gamma_\theta) \right) \\ &= -1 - p_\theta \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) + Q(\theta) \right) (z - \gamma_\theta) \right). \end{aligned} \quad (3.16)$$

Taking $z_0 = \gamma_\theta + \alpha e_\theta$ with α small enough, we have $f(z_0, \gamma_\theta) = 0$ and $\partial_\theta f(z_0, \gamma_\theta) \neq 0$, so that the existence, local uniqueness and smoothness of the projection $\text{proj}(z)$ in a tubular neighborhood of γ follows from the Implicit functions Theorem and the compactness of γ . Moreover, denoting $\theta = \text{proj}(z)$ and $\theta_h = \text{proj}(z + h) - \text{proj}(z)$, we have $\theta_h = O(h)$ by smoothness of the projection, and

$$\begin{aligned} p_{\theta+\theta_h}(z + h - \gamma_{\theta+\theta_h}) &= 0 \\ &= p_\theta(z + h - \gamma_\theta) - \theta_h p_\theta(\dot{\gamma}_\theta) - \theta_h p_\theta \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) + Q(\theta) \right) (z - \gamma_\theta) \right) + O(h^2) \\ &= p_\theta(h) - \theta_h \left(1 + p_\theta \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_\theta) + Q(\theta) \right) (z - \gamma_\theta) \right) \right) + O(h^2), \end{aligned} \quad (3.17)$$

which implies (3.12). \square

In the following $\gamma = \gamma^\delta$ stands for the limit cycle for $\delta > 0$. We will denote

$$\text{dist}_\Pi^\delta(z, \gamma) := \left| p_{\text{proj}(z)}^{\delta, \perp} (z - \gamma_{\text{proj}(z)}) \right|. \quad (3.18)$$

Now a variation of constants (see [30], page 84) shows that the solution to

$$\dot{z}_t = \delta DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+t}) z_t + g(t), \quad (3.19)$$

for a smooth mapping $g(t)$, satisfies

$$z_t = \Pi^\delta(\theta_0 + t, \theta_0) z_t + \int_0^t \Pi^\delta(\theta_0 + t, \theta_0 + s) g(s) \, ds. \quad (3.20)$$

So, since

$$\dot{m}_t - \dot{\gamma}_{\theta_0+t} = \delta \int_{\mathbb{R}^2} F(z + m_t) \, d\tilde{\mu}_t(z) - \delta \int_{\mathbb{R}^2} F(z + \gamma_{\theta_0+t}) q_0(z) \, dz, \quad (3.21)$$

we have

$$m_t - \gamma_{\theta_0+t} = \Pi^\delta(\theta_0 + t, \theta_0) (m_0 - \gamma_0) + \delta \int_0^t \Pi^\delta(\theta_0 + t, \theta_0 + s) g(s), \quad (3.22)$$

with

$$\begin{aligned} g(s) &= F_{\frac{\sigma^2}{K}}(m_s) - F_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+s}) - DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+s})(m_s - \gamma_{\theta_0+s}) \\ &\quad + \int_{\mathbb{R}^2} F(z + m_s) \, d\tilde{\mu}_s(z) - \int_{\mathbb{R}^2} F(z + m_s) q_0(z) \, dz. \end{aligned} \quad (3.23)$$

For an initial condition μ_0 of (1.5) with $m_0 = \int_{\mathbb{R}^2} z \, d\mu_0(z) \in E_\theta^\delta(1)$, we define $T_{r,\theta}(\mu_0)$ the return time

$$T_{r,\theta}(\mu_0) = \inf\{t > 0 : m_t \in E_\theta^\delta(1)\}. \quad (3.24)$$

Lemma 3.2. *There exist positive constants κ_2 and δ_2 such that for any μ_0 satisfying the hypotheses of Lemma 2.1 and Lemma 2.4, the following is true: if $\delta \leq \delta_2$ and $\text{dist}_{\Pi}^{\delta}(m_0, \gamma) \leq \kappa_2 \delta$, then*

$$\sup_{t \geq 0} \text{dist}_{\Pi}^{\delta}(m_t, \gamma) \leq \kappa_2 \delta. \quad (3.25)$$

In particular $T_e(R_0) = \infty$.

Proof. First remark that, by regularity of F and compactness of the limit cycle γ , there exists positive constants C_1 and α_F such that for all $s \in [0, T_{\gamma}/\delta]$ and m such that $|m - \gamma_s| \leq \alpha_F$, we have

$$\left| F_{\frac{\sigma^2}{K}}(m) - F_{\frac{\sigma^2}{K}}(\gamma_s) - DF_{\frac{\sigma^2}{K}}(\gamma_s)(m - \gamma_s) \right| \leq C_1 |m - \gamma_s|^2. \quad (3.26)$$

Consider a $\kappa_2 > 0$ and suppose that δ is small enough such that $(1 + C_{\Pi})\kappa_2 \delta < \alpha_F$. Suppose then that $m_0 \in E_{\theta_0}^{\delta}(\kappa_2 \delta)$ for $\theta_0 = \text{proj}(m_0) \in \mathbb{S}_{\delta}$ and consider the time

$$t_1 := \inf\{t \geq 0 : |m_t - \gamma_{\theta_0+t}| \geq (1 + C_{\Pi})\kappa_2 \delta\}. \quad (3.27)$$

Since γ is strictly included in $B(0, R_0)$, for δ taken small enough we have $t_1 \leq T_e(R_0)$. For $t \leq t_1$ we obtain

$$|m_t - \gamma_{\theta_0+t}| \leq C_{\Pi} |m_0 - \gamma_{\theta_0}| + C_{\Pi} \delta \int_0^t |g(s)| \, ds, \quad (3.28)$$

and

$$|g(s)| \leq C_1 |m_s - \gamma_{\theta_0+s}|^2 + \left| \int_{\mathbb{R}^2} F(z + m_s) \, d\tilde{\mu}_s(z) - \int_{\mathbb{R}^2} F(z + m_s) q_0(z) \, dz \right|. \quad (3.29)$$

Now, recalling the notations used in the proof of Lemma 2.4, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} F(z + m_s) \, d\tilde{\mu}_s(z) - \int_{\mathbb{R}^2} F(z + m_s) q_0(z) \, dz \right|^2 \\ &= \left(\mathbf{E} \left[X_s - \frac{X_s^3}{3} - Y_s \right] - \mathbf{E} \left[Z_s^x + x_s - \frac{(Z_s^x + x_s)^3}{3} - Z_s^y - y_s \right] \right)^2 \\ & \quad + \left(\mathbf{E} [X_s + a - bY_s] - \mathbf{E} [Z_s^x + x_s + a - Z_s^y - y_s] \right)^2, \end{aligned} \quad (3.30)$$

where $\{(X_t, Y_t), (Z_t^x, Z_t^y)\}$ is the ε -coupling introduced in Lemma 2.4. First, note that, by (2.27), we have for $t \leq t_1 \leq T_e(R_0)$, $\mathbf{E}[X_t - x_t - Z_t^x]^2 \leq \mathbf{E} \left[\left| \tilde{X}_t - Z_t^x \right|^2 \right] \leq \kappa_1^2 \delta^2 + \varepsilon$. Moreover, using Lemma 2.1 and Lemma 2.4,

$$\begin{aligned} \mathbf{E}[X_t^3 - (Z_t^x + x_t)^3] &= \mathbf{E} \left[\tilde{X}_t^3 - (Z_t^x)^3 \right] + 3x_t \mathbf{E} \left[\tilde{X}_t^2 - (Z_t^x)^2 \right], \\ &\leq C_{R_0, \sigma^2/K} \mathbf{E} \left[\left| \tilde{X}_t - Z_t^x \right|^2 \right]^{\frac{1}{2}} \leq C_2 (\delta^2 + \varepsilon)^{\frac{1}{2}}, \end{aligned}$$

for some constant $C_2 > 0$, independent of ε . This altogether implies that, for some constant $C_3 > 0$,

$$|g(s)| \leq C_1 |m_s - \gamma_{\theta_0+s}|^2 + C_3 (\delta^2 + \varepsilon)^{\frac{1}{2}}. \quad (3.31)$$

Taking $\varepsilon \searrow 0$ ($g(s)$ does not depend on ε), one obtains

$$|g(s)| \leq C_1 |m_s - \gamma_{\theta_0+s}|^2 + C_3 \delta. \quad (3.32)$$

We deduce that, for $t \leq t_1$, since $m_0 - \gamma_{\theta_0} = p_{\theta_0}^\perp(m_0 - \gamma_{\theta_0})$,

$$|m_t - \gamma_{\theta_0+t}| \leq C_\Pi \kappa_2 \delta \left(1 + \delta t \left(C_1(1 + C_\Pi)^2 \kappa_2 \delta + \frac{C_3}{\kappa_2} \right) \right), \quad (3.33)$$

and thus, choosing $\kappa_2 \geq 4C_3C_\Pi T_\gamma$ and $\delta \leq \frac{1}{4T_\gamma C_1 \kappa_2 C_\Pi (1+C_\Pi)^2}$ we obtain that

$$t_1 \geq \frac{2T_\gamma}{\delta}. \quad (3.34)$$

Note that choosing also $\delta \leq \frac{\alpha_0}{(1+C_\Pi)\kappa_2}$, (recall the definition of α_0 in Lemma 3.1) ensures that m_t stays in the α_0 -neighborhood of the limit cycle γ . In particular, the projection

$$\theta_t = \theta_t^\delta = \text{proj}^\delta(m_t) \quad (3.35)$$

is well-defined for all $t \in [0, \frac{2T_\gamma}{\delta}]$.

Let us now prove that $\dot{\theta}_t^\delta = 1 + O(\delta)$ for $t \leq t_1$, which implies in particular that $T_{r,\theta}(\mu_0) \in \left[\frac{T_\gamma}{\delta} - c_1, \frac{T_\gamma}{\delta} + c_1 \right]$ for some $c_1 > 0$ that does not depend on δ (but may depend on κ_2). Now, recalling Lemma 3.1, for $s \leq t_1$ we have $\dot{\theta}_s = \frac{1}{\delta} p_{\theta_s}(\dot{m}_s) + O(\delta)$, with

$$p_{\theta_s}(\dot{m}_s) = \delta + p_{\theta_s}(\dot{m}_s) - p_{\theta_s} \left(\delta F_{\frac{\sigma^2}{K}}(\gamma_{\theta_s}) \right), \quad (3.36)$$

and

$$\begin{aligned} \frac{1}{\delta} \left(\dot{m}_s - \delta F_{\frac{\sigma^2}{K}}(\gamma_{\theta_s}) \right) &= F_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+s}) - F_{\frac{\sigma^2}{K}}(\gamma_{\theta_s}) + F_{\frac{\sigma^2}{K}}(m_s) - F_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+s}) \\ &\quad + \int_{\mathbb{R}^2} F(z + m_s) d\tilde{\mu}_s(z) - \int_{\mathbb{R}^2} F(z + m_s) q_0(z) dz \\ &= F_{\frac{\sigma^2}{K}}(\gamma_{\theta_0+s}) - F_{\frac{\sigma^2}{K}}(\gamma_{\theta_s}) + O(\delta), \end{aligned} \quad (3.37)$$

where we have used the fact that $t \leq t_1$ and $F_{\frac{\sigma^2}{K}}$ is smooth, and similar arguments as above to tackle the difference of integral terms. Moreover, by (3.13) and the definition of $\gamma = \gamma^\delta$, $|\gamma_{\theta_0+s} - \gamma_{\theta_s}| = |\gamma_{\delta(\theta_0+s)}^\delta - \gamma_{\delta\theta_s}^\delta| = |\gamma_{\text{proj}^1(\gamma_{\theta_0+s})}^\delta - \gamma_{\text{proj}^1(m_s)}^\delta| \leq C_{\text{proj},\gamma} |m_s - \gamma_{\theta_0+s}|$ for some constant $C_{\text{proj},\gamma}$. By (3.34), this last quantity is of order $O(\kappa_2\delta)$, uniformly in $s \in [0, \frac{2T_\gamma}{\delta}]$. We obtain $p_{\theta_s}(\dot{m}_s) = \delta + O(\delta^2)$, and thus $\dot{\theta}_s = 1 + O(\delta)$.

It remains now to prove that $|m_t - \gamma_{\text{proj}(m_t)}| \leq \kappa_2\delta$ for all $t \leq T_{r,\theta}(\mu_0)$, taking a larger value for κ_2 if needed. But we have on one hand, recalling (3.11), (3.22) and the estimates made above,

$$\left| p_{\theta_0+t}^\perp(m_t - \gamma_{\theta_0+t}) \right| \leq \kappa_2 \delta \left(e^{-\lambda\delta t} + C_\Pi \delta t \left(C_1(1 + C_\Pi)^2 \kappa_2 \delta + \frac{C_3}{\kappa_2} \right) \right). \quad (3.38)$$

Now, remarking that, for $\eta = \frac{1-e^{-2\lambda T_\gamma}}{2T_\gamma}$, $e^{-\lambda u} + \eta u \leq 1$ for $u \in [0, 2T_\gamma]$, we have

$$\left| p_{\theta_0+t}^\perp(m_t - \gamma_{\theta_0+t}) \right| \leq \left(1 - \frac{\eta}{2} \delta t \right) \kappa_2 \delta, \quad (3.39)$$

if we take $\kappa_2 \geq \frac{4C_3C_\Pi}{\eta}$ and $\delta \leq \frac{\eta}{4C_1C_\Pi(1+C_\Pi)^2\kappa_2}$.

On the other hand, using the fact that $p_{\theta_t}^\perp(\dot{\gamma}_{\theta_t}) = 0$, that for all w

$$\left| \left(p_{\theta_t}^{\delta,\perp} - p_{\theta_0+t}^{\delta,\perp} \right) (w) \right| = \left| \left(p_{\delta\theta_t}^{1,\perp} - p_{\delta\theta_0+\delta t}^{1,\perp} \right) (w) \right| \leq \delta C_{\text{proj},\gamma} |w| |\theta_0 + t - \theta_t|,$$

and the fact that $|m_t - \gamma_{\theta_0+t}| \leq (1 + C_\Pi)\kappa_2\delta$, we have

$$\begin{aligned} \left| p_{\theta_t}^\perp(m_t - \gamma_{\theta_t}) - p_{\theta_0+t}^\perp(m_t - \gamma_{\theta_0+t}) \right| &\leq \left| p_{\theta_t}^\perp(\gamma_{\theta_0+t} - \gamma_{\theta_t}) \right| + \left| (p_{\theta_t}^\perp - p_{\theta_0+t}^\perp)(m_t - \gamma_{\theta_0+t}) \right| \\ &\leq C_{\text{proj},\gamma}\delta^2 (|\theta_0 + t - \theta_t|^2 + \kappa_2|\theta_0 + t - \theta_t|). \end{aligned} \quad (3.40)$$

If $v_t := \theta_t - (\theta_0 + t)$ we have, using (3.12)

$$\delta\dot{v}_t - p_{\theta_t}(\dot{m}_t) + \delta = p_{\theta_t}(\dot{m}_t) \left(\frac{1}{1 + p_{\theta_t} \left(\left(DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_t}) + \frac{1}{\delta}Q^\delta(\theta_t) \right) (m_t - \gamma_{\theta_t}) \right)} - 1 \right) \quad (3.41)$$

The term within the brackets is controlled by $C|m_t - \gamma_{\theta_t}|$, for some universal constant $C > 0$. By the regularity of $t \mapsto \gamma_t^\delta = \gamma_{t\delta}^1$ and the Lipchitz-continuity of $\text{proj}(\cdot)$ (recall (3.13)), there is a constant $C_{\gamma,\text{proj}} > 0$

$$|\gamma_{\theta_0+t} - \gamma_{\theta_t}| = \left| \gamma_{\text{proj}(\gamma_{\theta_0+t})} - \gamma_{\text{proj}(m_t)} \right| \leq C_{\gamma,\text{proj}} |\gamma_{\theta_0+t} - m_t|.$$

Hence, $|m_t - \gamma_{\theta_t}| \leq |m_t - \gamma_{\theta_0+t}| + |\gamma_{\theta_0+t} - \gamma_{\theta_t}| \leq C'_{\gamma,\text{proj}} |m_t - \gamma_{\theta_0+t}|$. Putting things together with $|m_t - \gamma_{\theta_0+t}| \leq (1 + C_\Pi)\kappa_2\delta$ (recall (3.34)), we obtain finally that

$$|\delta\dot{v}_t - p_{\theta_t}(\dot{m}_t) + \delta| \leq C_{\text{proj},\gamma}\kappa_2\delta |p_{\theta_t}(\dot{m}_t)|. \quad (3.42)$$

So recalling that $p_{\theta_t}(\dot{m}_t) = \delta + O(\kappa_2\delta^2)$, we have $\dot{v}_t = O(\kappa_2\delta)$, and thus $v_t = tO(\kappa_2\delta)$, since $v_0 = 0$. This means that for $t \in \left[0, \frac{2T_\gamma}{\delta}\right]$ and some positive constant C_4 ,

$$\left| p_{\theta_t}^\perp(m_t - \gamma_{\theta_t}) - p_{\theta_0+t}^\perp(m_t - \gamma_{\theta_0+t}) \right| \leq C_4\kappa_2^2\delta^3 t. \quad (3.43)$$

Finally, we obtain, combining (3.39) and (3.43),

$$\text{dist}_\Pi(m_t, \gamma) \leq \left(1 - \frac{\eta}{2}\delta t + C_4\kappa_2\delta^2 t\right) \kappa_2\delta, \quad (3.44)$$

which implies that $|m_t - \gamma_{\text{proj}(m_t)}| \leq \kappa_2\delta$ for all $t \leq T_{r,\theta}(\mu_0)$, taking $\delta \leq \frac{\eta}{2C_4\kappa_2}$. This completes the proof, since $\mu_{T_{r,\theta}(\mu_0)}$ satisfies then the same hypotheses as μ_0 (recall Lemma 2.1, Lemma 2.4 and the fact that $T_{r,\theta}(\mu_0) \leq T_e(R_0)$), the result for all t is obtained by recursion. \square

For two solutions $(X_{1,t}, Y_{1,t}), (X_{2,t}, Y_{2,t})$ of (1.1) with initial conditions $(X_{1,0}, Y_{1,0})$ and $(X_{2,0}, Y_{2,0})$, with respective distributions $\mu_{1,t}$ and $\mu_{2,t}$ and expectations $m_{1,t} = (x_{1,t}, y_{1,t})$ and $m_{2,t} = (x_{2,t}, y_{2,t})$, we will use the notations

$$\begin{aligned} \Delta X_t &= X_{1,t} - X_{2,t}, & \Delta Y_t &= Y_{1,t} - Y_{2,t}, & \Delta \tilde{X}_t &= \tilde{X}_{1,t} - \tilde{X}_{2,t}, & \Delta \tilde{Y}_t &= \tilde{Y}_{1,t} - \tilde{Y}_{2,t}, \\ \Delta m_t &= m_{1,t} - m_{2,t}, & \Delta \theta_t &= \theta_{1,t} - \theta_{2,t} = \text{proj}(m_{1,t}) - \text{proj}(m_{2,t}). \end{aligned} \quad (3.45)$$

We choose T such that

$$e^{-\lambda T} \leq \frac{1}{4}, \quad e^{-\frac{b}{c}T} \leq \frac{1}{16}. \quad (3.46)$$

Lemma 3.3. *There exist $\delta_3 > 0$ and $\kappa_3 \geq 1$ such that for all $\delta \leq \delta_3$, the following is true: if for $i = 1, 2$, $\int x^6 d\mu_{i,0} \leq \kappa_0^x$, $\int y^6 d\mu_{i,0} \leq \kappa_0^y$ (recall Lemma 2.1), $W(\tilde{\mu}_{i,0}, q_0) \leq \kappa_1\delta$ and $\text{dist}_\Pi(m_{i,0}, \gamma) \leq \kappa_2\delta$, then for all $t \in \left[0, \frac{2T}{\delta}\right]$ we have*

$$W(\mu_{1,t}, \mu_{2,t}) \leq \kappa_3 W(\mu_{1,0}, \mu_{2,0}). \quad (3.47)$$

Moreover, under the same assumptions, there exist positive constants $\kappa_4, \kappa_5, \kappa_6$ such that for any coupling $\{(X_{1,0}, Y_{1,0}), (X_{2,0}, Y_{2,0})\} \sim \pi_0 \in \mathcal{C}(\mu_{1,0}, \mu_{2,0})$ of the initial condition, the solutions $(X_{1,t}, Y_{1,t}), (X_{2,t}, Y_{2,t})$ of (1.1) driven by the same Brownian motion satisfy

$$\left(\mathbf{E} \left[\left(\Delta \tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq e^{-(K-\kappa_4\delta)t} \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} + \kappa_5 \delta^{1-\beta} \Lambda(\pi_0), \quad (3.48)$$

and

$$\left(\mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq e^{-\frac{b\delta}{c}t} \left(\left(\mathbf{E} \left[\left(\Delta \tilde{Y}_0 \right)^2 \right] \right)^{\frac{1}{2}} + \kappa_6 \delta \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} \right) + \frac{\kappa_5}{b} \delta^{1-\beta} \Lambda(\pi_0). \quad (3.49)$$

Proof. Fix an arbitrary coupling $\{(X_{1,0}, Y_{1,0}), (X_{2,0}, Y_{2,0})\}$ with law $\pi_0 \in \mathcal{C}(\mu_{1,0}, \mu_{2,0})$ and consider the solutions $(X_{i,t}, Y_{i,t}), i = 1, 2$ to (1.1) with initial condition $(X_{i,0}, Y_{i,0}), i = 1, 2$ driven by the same Brownian motion. The law π_t of $\{(X_{1,t}, Y_{1,t}), (X_{2,t}, Y_{2,t})\}$ belongs to $\mathcal{C}(\mu_{1,t}, \mu_{2,t})$.

Step 1: control on Δm_t . This part is based on the a priori control obtained in Lemmas 2.1, 2.4 and 3.2. We have

$$\begin{aligned} \Delta \dot{m}_t &= \delta \int_{\mathbb{R}^2} F(z + m_{1,t}) d\tilde{\mu}_{1,t}(z) - \delta \int_{\mathbb{R}^2} F(z + m_{2,t}) d\tilde{\mu}_{2,t}(z) \\ &= \delta DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_{2,0+t}}) \Delta m_t + \delta h(t), \end{aligned} \quad (3.50)$$

where

$$\begin{aligned} h(t) &:= \left(DF_{\frac{\sigma^2}{K}}(m_{2,t}) - DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_{2,0+t}}) \right) \Delta m_t \\ &\quad + \int_{\mathbb{R}^2} DF(z + m_{2,t}) \Delta m_t d\tilde{\mu}_{1,t}(z) - \int_{\mathbb{R}^2} DF(z + m_{2,t}) \Delta m_t q_0(z) dz \\ &\quad + \int_{\mathbb{R}^2} \left(F(z + m_{1,t}) - F(z + m_{2,t}) - DF(z + m_{2,t}) \Delta m_t \right) d\tilde{\mu}_{1,t}(z) \\ &\quad + \int_{\mathbb{R}^2} F(z + m_{2,t}) d\tilde{\mu}_{1,t}(z) - \int_{\mathbb{R}^2} F(z + m_{2,t}) d\tilde{\mu}_{2,t}(z), \\ &=: h_1(t) + \dots + h_4(t). \end{aligned} \quad (3.51)$$

One obtains from (3.20) that

$$\Delta m_t = \Pi^\delta(\theta_{2,0} + t, \theta_{2,0}) \Delta m_0 + \delta \int_0^t \Pi^\delta(\theta_{2,0} + t, \theta_{2,0} + s) h(s) ds. \quad (3.52)$$

Since $m_{2,0} \in E_0^\delta(\kappa_2\delta)$ by assumption, adapting the proof of Lemma 3.2, we can find a constant $C(T)$ such that $|m_{2,t} - \gamma_{\theta_{2,0+t}}| \leq C(T)\delta$, and $|m_{2,t} + \gamma_{\theta_{2,0+t}}| \leq 2R_0$ for all $t \leq \frac{2T}{\delta}$. Hence, we deduce that

$$\begin{aligned} |h_1(t)| &\leq \left| DF_{\frac{\sigma^2}{K}}(m_{2,t}) - DF_{\frac{\sigma^2}{K}}(\gamma_{\theta_{2,0+t}}) \right| |\Delta m_t| = \left| (x_{2,t})^2 - \left(\gamma_{\theta_{2,0+t}}^x \right)^2 \right| |\Delta m_t| \\ &\leq 4C(T)R_0\delta |\Delta m_t| =: c_1\delta |\Delta m_t|. \end{aligned}$$

Moreover, using the notations of Lemma 2.4,

$$\begin{aligned}
|h_2(t)| &\leq \left| \int_{\mathbb{R}^2} DF(z + m_{2,t}) d\tilde{\mu}_{1,t}(z) - \int_{\mathbb{R}^2} DF(z + m_{2,t}) q_0(z) dz \right| |\Delta m_t| \\
&= \left| \int_{\mathbb{R}^2} (x + x_{2,t})^2 d\tilde{\mu}_{1,t}(x, y) - \int_{\mathbb{R}^2} (x + x_{2,t})^2 q_0(x, y) dx dy \right| |\Delta m_t| \\
&= \left| \mathbf{E} \left[\left(\tilde{X}_{1,t} \right)^2 - (Z_t^x)^2 \right] \right| |\Delta m_t| \\
&\leq \left(\mathbf{E} \left[(X_{1,t})^2 \right]^{\frac{1}{2}} + |x_{1,t}| + \mathbf{E} \left[(Z_t^x)^2 \right]^{\frac{1}{2}} \right) \mathbf{E} \left[\left| \tilde{X}_{1,t} - Z_t^x \right|^2 \right]^{\frac{1}{2}} |\Delta m_t| \\
&\leq \left((\kappa_0^x)^{\frac{1}{6}} + R_0 + \frac{\sigma}{K^{\frac{1}{2}}} \right) \kappa_1 \delta |\Delta m_t| \\
&=: c_2 \delta |\Delta m_t|.
\end{aligned}$$

Concerning h_3 , a Taylor expansion shows that there a numerical constant C such that $|h_3(t)| \leq C(|x_{1,t}| + |x_{2,t}|) |\Delta m_t|^2 \leq c_3 \delta |\Delta m_t|$, for some constant c_3 . It remains to treat the last term h_4 :

$$\begin{aligned}
|h_4(t)|^2 &= \left| \mathbf{E} \left[\Delta \tilde{X}_t - \left(\frac{(\tilde{X}_{1,t} + x_{2,t})^3}{3} - \frac{(\tilde{X}_{2,t} + x_{2,t})^3}{3} \right) - \Delta \tilde{Y}_t \right] \right|^2 \\
&\quad + \left| \frac{1}{c} \mathbf{E} \left[\Delta \tilde{X}_t - b \Delta \tilde{Y}_t \right] \right|^2, \quad (3.53)
\end{aligned}$$

and relying on Lemma 2.1, Lemma 2.4 and the identity

$$(x_1 + n)^3 - (x_2 + n)^3 = (x_1 - x_2)((x_1 + n)^2 + (x_1 + n)(x_2 + n) + (x_2 + n)^2), \quad (3.54)$$

we obtain that

$$|h_4(t)| \leq c_4 \left(\left(\mathbf{E} \left[(\Delta \tilde{X}_t)^2 \right] \right)^{\frac{1}{2}} + b \left(\mathbf{E} \left[(\Delta \tilde{Y}_t)^2 \right] \right)^{\frac{1}{2}} \right), \quad (3.55)$$

for some constant $c_4 > 0$. Putting everything together, we obtain, for $t \leq \frac{2T}{\delta}$,

$$\begin{aligned}
|\Delta m_t| &\leq C_{\Pi} \left(|\Delta m_0| + c_4 \delta \int_0^t \left(\left(\mathbf{E} \left[(\Delta \tilde{X}_s)^2 \right] \right)^{\frac{1}{2}} + b \left(\mathbf{E} \left[(\Delta \tilde{Y}_s)^2 \right] \right)^{\frac{1}{2}} \right) ds \right) \\
&\quad + C_{\Pi} (c_1 + c_2 + c_3) \delta^2 \int_0^t |\Delta m_s| ds. \quad (3.56)
\end{aligned}$$

With these notations at hand, for the rest of the proof, we define

$$\kappa_3 := C_{\Pi} (1 + 4c_4 T) e^{2C_{\Pi}(c_1+c_2+c_3)T} + 1 \quad (3.57)$$

The reason for this particular choice of κ_3 will become clear at Step 4 below. For this choice of κ_3 , define (recall the definition of $d(\cdot, \cdot)$ in (2.16))

$$t_2 := t_2(\kappa_3, \delta) := \inf \{ t \geq 0 : \Lambda(\pi_t) > \kappa_3 \Lambda(\pi_0) \}. \quad (3.58)$$

Note that t_2 depends on κ_3 and δ but also on the choice of the particular coupling π_0 of the initial condition. For the rest of the proof, we use the shortcut :

$$\Lambda_t := \Lambda(\pi_t). \quad (3.59)$$

Step 2: control on $\Delta\tilde{X}_t$. We have

$$\begin{aligned} \frac{1}{2} d \left(\Delta\tilde{X}_t \right)^2 &= -K \left(\Delta\tilde{X}_t \right)^2 dt \\ &\quad + \left(\Delta\tilde{X}_t \right) \left(-\Delta\dot{x}_t + \delta \left(\Delta X_t - \frac{(X_{1,t})^3 - (X_{2,t})^3}{3} - \Delta Y_t \right) \right) dt, \end{aligned} \quad (3.60)$$

so that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] &\leq -(K - \delta) \mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \\ &\quad + \left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \left(|\Delta\dot{x}_t| + \delta |\Delta x_t| + \delta |\Delta y_t| + \delta \left(\mathbf{E} \left[\left(\Delta\tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} \right) \\ &\quad - \frac{\delta}{3} \mathbf{E} \left[\Delta\tilde{X}_t \left(\left(\tilde{X}_{1,t} + x_{1,t} \right)^3 - \left(\tilde{X}_{2,t} + x_{2,t} \right)^3 \right) \right]. \end{aligned} \quad (3.61)$$

Now, remarking that, for $x_1, x_2, n_1, n_2 \in \mathbb{R}$ and $\Delta x := x_1 - x_2$, $\Delta n := n_1 - n_2$,

$$\begin{aligned} &-\Delta x \left((x_1 + n_1)^3 - (x_2 + n_2)^3 \right) \\ &= -\Delta x \left(x_1^3 - x_2^3 + 3(x_1^2 n_1 - x_2^2 n_2) + 3(x_1 n_1^2 - x_2 n_2^2) + n_1^3 - n_2^3 \right) \\ &= -(\Delta x)^2 \left(x_1^2 + x_1 x_2 + x_2^2 + 3n_1(x_1 + x_2) + 3n_1^2 \right) \\ &\quad - \Delta x \Delta n \left(3x_2^2 + 3x_2 + n_1^2 + n_1 n_2 + n_2^2 \right) \\ &\leq 12n_1^2 (\Delta x)^2 - \Delta x \Delta n \left(3x_2^2 + 3x_2 + n_1^2 + n_1 n_2 + n_2^2 \right), \end{aligned} \quad (3.62)$$

we get, recalling Lemma 2.1, for a positive constant C_1 ,

$$\begin{aligned} \mathbf{E} \left[\Delta\tilde{X}_t \left(\left(\tilde{X}_{1,t} + x_{1,t} \right)^3 - \left(\tilde{X}_{2,t} + x_{2,t} \right)^3 \right) \right] \\ \leq C_1 \left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] + |\Delta x_t| \left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \right). \end{aligned} \quad (3.63)$$

Proceeding similarly, we obtain, for some constant C_2 ,

$$\begin{aligned} \frac{1}{\delta} |\Delta\dot{x}_t| &= \left| \mathbf{E} \left[\Delta X_t - \frac{(X_{1,t})^3 - (X_{2,t})^3}{3} - \Delta Y_t \right] \right| \\ &\leq C_2 \left(\left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} + \left(\mathbf{E} \left[\left(\Delta\tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} + |\Delta m_t| \right). \end{aligned} \quad (3.64)$$

Gathering all these estimates, we get, for positive constants κ_4 and C_3 , for $t \leq t_2$,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] &\leq -(K - \kappa_4 \delta) \mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \\ &\quad + C_3 \delta \left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \left(\left(\mathbf{E} \left[\left(\Delta\tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} + |\Delta m_t| \right) \\ &\leq -(K - \kappa_4 \delta) \mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] + C_3 \kappa_3 \delta \left(b^{-1} + \delta^{-\beta} \right) \Lambda_0 \left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}}, \end{aligned} \quad (3.65)$$

where we recall the definition of Λ_t in (3.59) and the definition of t_2 in (3.58). By Lemma A.1, this implies that for all $t \leq t_2$,

$$\left(\mathbf{E} \left[\left(\Delta \tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \max \left\{ \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}}, \frac{C_3 \kappa_3 (b^{-1} + \delta^{-\beta})}{K - \kappa_4 \delta} \delta \Lambda_0 \right\}. \quad (3.66)$$

So taking δ small enough (depending on κ_3), we deduce in particular that, since $\beta \in (0, 1)$ by hypothesis, for $t \leq t_2$

$$\mathbf{E} \left[\left(\Delta \tilde{X}_t \right)^2 \right]^{\frac{1}{2}} \leq \Lambda_0. \quad (3.67)$$

Step 3: control on $\Delta \tilde{Y}_t$. A simple calculation leads to

$$\frac{c}{2\delta} d \left(\Delta \tilde{Y}_t \right)^2 = \left(\Delta \tilde{X}_t \Delta \tilde{Y}_t - b \left(\Delta \tilde{Y}_t \right)^2 \right) dt, \quad (3.68)$$

so that, for $t \leq t_2$, using (3.67),

$$\frac{c}{2\delta} \frac{d}{dt} \mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \leq -b \mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] + \Lambda_0 \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}}, \quad (3.69)$$

which implies in particular, by Lemma A.1, that for all $t \leq t_2$,

$$\left(\mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \max \left\{ \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_0 \right)^2 \right] \right)^{\frac{1}{2}}, \frac{\Lambda_0}{b} \right\} \leq \frac{\Lambda_0}{b}. \quad (3.70)$$

Step 4: proof of (3.47). Let us first prove that, for the choice of κ_3 in (3.57), we have, for $\delta \leq \delta_3$ for some δ_3 sufficiently small

$$t_2(\kappa_3, \delta) \geq \frac{2T}{\delta}. \quad (3.71)$$

Suppose that δ_3 is chosen sufficiently small so that (3.67) is true and such that $\delta_3 \leq 1$. Suppose that (3.71) does not hold for some $\delta \leq \delta_3$: $t_2 < \frac{2T}{\delta}$. For this δ , for $t \leq t_2 < \frac{2T}{\delta}$, incorporating (3.67) and (3.70) into (3.56) and applying Grönwall's Lemma gives, for $t \leq t_2 < \frac{2T}{\delta}$,

$$\begin{aligned} \delta^\beta |\Delta m_t| &\leq C_\Pi \left(\delta^\beta |\Delta m_0| + 2c_4 \delta^{1+\beta} t \Lambda_0 \right) e^{C_\Pi (c_1 + c_2 + c_3) \delta^2 t}, \\ &\leq \Lambda_0 C_\Pi (1 + 4c_4 T) e^{2C_\Pi (c_1 + c_2 + c_3) T} = (\kappa_3 - 1) \Lambda_0, \end{aligned} \quad (3.72)$$

where we have used $\delta_3 \leq 1$. Collecting (3.67), (3.70) and (3.72), we obtain that for all $t \leq t_2$, $\Lambda_t < \kappa_3 \Lambda_0$. By continuity, there exists some $\epsilon > 0$ such that $\Lambda_t < \kappa_3 \Lambda_0$ for $t \in [t_2, t_2 + \epsilon)$, which contradicts the definition of t_2 . Hence, (3.71) follows. In particular, since W is an infimum over all possible coupling, for $t \leq \frac{2T}{\delta}$, $W(\mu_{1,t}, \mu_{2,t}) \leq \kappa_3 \Lambda_0 = \kappa_3 \Lambda(\pi_0)$. Since this is true for all possible coupling of the initial condition, (3.47) follows.

Step 5: Let us now prove (3.48): from (3.65) and applying Lemma A.1, we deduce that, for all $t \leq \frac{2T}{\delta}$,

$$\left(\mathbf{E} \left[\left(\Delta \tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq e^{-(K - \kappa_4 \delta)t} \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} + \frac{C_3 \kappa_3 (b^{-1} + \delta^{-\beta})}{K - \kappa_4 \delta} \delta \Lambda_0, \quad (3.73)$$

which implies (3.48) with $\kappa_5 = \frac{2C_3\kappa_3b(1+b)}{K}$ as soon as $\delta \leq \frac{K}{2\kappa_4}$. Let us now turn to the proof of (3.49): from (3.69), we have, for all $t \leq \frac{2T}{\delta}$,

$$\begin{aligned} \frac{c}{2\delta} \frac{d}{dt} \mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] &\leq -b \mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \\ &+ \left(e^{-(K-\kappa_4\delta)t} \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} + \kappa_5 \delta^{1-\beta} \Lambda_0 \right) \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}}, \end{aligned} \quad (3.74)$$

so that, by Lemma A.1 again implies

$$\begin{aligned} \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} &\leq e^{-\frac{b\delta}{c}t} \left(\left(\mathbf{E} \left[\left(\Delta \tilde{Y}_0 \right)^2 \right] \right)^{\frac{1}{2}} + \frac{\delta}{cK - (c\kappa_4 + b)\delta} \left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} \right) \\ &+ \kappa_5 \delta^{1-\beta} \frac{\Lambda_0}{b}, \end{aligned} \quad (3.75)$$

which leads to (3.49) with $\kappa_6 = \frac{2}{cK}$, as soon as $\delta \leq \frac{cK}{2(c\kappa_4+b)}$.

This concludes the proof of Lemma 3.3. \square

Lemma 3.4. *There exists a $\delta_4 > 0$ and positive constants $\kappa_7, \kappa_8, \kappa_9$ such that if $\delta \leq \delta_4$, if the hypotheses of Lemma 3.3 are satisfied, and if moreover we have the initial bounds $W(\tilde{\mu}_{1,0}, \tilde{\mu}_{2,0}) \leq \kappa_7 \delta^2 |\Delta \theta_0|$ and $\left| p_{\theta_{1,0}}^\perp(m_{1,0} - \gamma_{\theta_{1,0}}) - p_{\theta_{2,0}}^\perp(m_{2,0} - \gamma_{\theta_{2,0}}) \right| \leq \kappa_8 \delta^{\frac{3}{2}} |\Delta \theta_0|$, then for all $t \in [0, \frac{2T}{\delta}]$ we have*

$$\left| |\Delta \theta_t| - |\Delta \theta_0| \right| \leq \kappa_9 \delta^{\frac{1}{2}} |\Delta \theta_0|, \quad (3.76)$$

and for all $t \in [\frac{T}{\delta}, \frac{2T}{\delta}]$ we have

$$W(\tilde{\mu}_{1,t}, \tilde{\mu}_{2,t}) \leq \kappa_7 \delta^2 |\Delta \theta_t|, \quad (3.77)$$

and

$$\left| p_{\theta_{1,t}}^\perp(m_{1,t} - \gamma_{\theta_{1,t}}) - p_{\theta_{2,t}}^\perp(m_{2,t} - \gamma_{\theta_{2,t}}) \right| \leq \kappa_8 \delta^{\frac{3}{2}} |\Delta \theta_t|. \quad (3.78)$$

Proof. Define $\kappa_7 = 16\kappa_5 |\dot{\gamma}|$, fix a $\kappa_8 > 0$ whose value will be chosen later, and suppose that $W(\tilde{\mu}_{1,0}, \tilde{\mu}_{2,0}) \leq \kappa_7 \delta^2 |\Delta \theta_0|$ and $\left| p_{\theta_{1,0}}^\perp(m_{1,0} - \gamma_{\theta_{1,0}}) - p_{\theta_{2,0}}^\perp(m_{2,0} - \gamma_{\theta_{2,0}}) \right| \leq \kappa_8 \delta^{\frac{3}{2}} |\Delta \theta_0|$.

In what follows, we also fix some $\varepsilon > 0$ and consider some coupling $\{(X_{1,0}, Y_{1,0}), (X_{2,0}, Y_{2,0})\} \sim \pi_0 \in \mathcal{C}(\mu_{0,1}, \mu_{2,0})$ of the initial condition such that

$$\Lambda(\pi_0) < W(\mu_{1,0}, \mu_{2,0}) + \varepsilon. \quad (3.79)$$

Remark first that, since $m_{1,0} - m_{2,0} = m_{1,0} - \gamma_{\theta_{1,0}} + \gamma_{\theta_{1,0}} - \gamma_{\theta_{2,0}} + \gamma_{\theta_{2,0}} - m_{2,0}$ and $m_{i,0} - \gamma_{\theta_{i,0}} = p_{\theta_{i,0}}^\perp(m_{i,0} - \gamma_{\theta_{i,0}}) e_{\theta_{i,0}}$ for $i = 1, 2$, we have

$$\begin{aligned} |\Delta m_0| &\leq |\gamma_{\theta_{1,0}} - \gamma_{\theta_{2,0}}| + \left| \left(p_{\theta_{1,0}}^\perp(m_{1,0} - \gamma_{\theta_{1,0}}) - p_{\theta_{2,0}}^\perp(m_{2,0} - \gamma_{\theta_{2,0}}) \right) e_{\theta_{2,0}} \right| \\ &+ \left| p_{\theta_{1,0}}^\perp(m_{1,0} - \gamma_{\theta_{1,0}}) (e_{\theta_{1,0}} - e_{\theta_{2,0}}) \right|, \end{aligned} \quad (3.80)$$

so that

$$|\Delta m_0| \leq \left(|\dot{\gamma}|_\infty + C_\Pi \left(\kappa_8 \delta^{\frac{1}{2}} + C_\Pi \kappa_2 \delta \right) \right) \delta |\Delta \theta_0| \leq 2\delta |\dot{\gamma}| |\Delta \theta_0|, \quad (3.81)$$

for δ small enough (depending on κ_8). By definition of κ_7 , this means in particular that

$$W(\mu_{1,0}, \mu_{2,0}) = \max \left(\delta^\beta |\Delta m_0|, W(\tilde{\mu}_{1,0}, \tilde{\mu}_{2,0}) \right) \leq 2|\dot{\gamma}| \delta^{1+\beta} |\Delta \theta_0|, \quad (3.82)$$

provided that $\delta \leq \left(\frac{2|\dot{\gamma}|}{\kappa_7}\right)^{\frac{1}{1-\beta}}$. Now remark that

$$||\Delta\theta_t| - |\Delta\theta_0|| \leq |\theta_{1,t} - (\theta_{1,0} + t)| + |\theta_{2,t} - (\theta_{2,0} + t)|, \quad (3.83)$$

and relying on similar arguments as in Lemma 3.2, we have $|\theta_{i,t} - (\theta_{i,0} + t)| \leq C_4\delta t$, for $t \in [0, \frac{2T}{\delta}]$, $i = 1, 2$, and some positive constant C_4 . So, for $|\Delta\theta_0| \geq \delta^{-\frac{1}{2}}$, we have $||\Delta\theta_t| - |\Delta\theta_0|| \leq 2C_4T\delta^{\frac{1}{2}}|\Delta\theta_0|$. Suppose now that $|\Delta\theta_0| \leq \delta^{-\frac{1}{2}}$. From Lemma 3.1 and Lemma 3.2 we deduce that

$$\Delta\theta_t = \frac{1}{\delta} (p_{\theta_{2,t}}(\Delta m_t) + O(\delta|\Delta m_t| + |\Delta m_t|^2)) = \frac{1}{\delta} (p_{\theta_{2,0+t}}(\Delta m_t) + O(\delta|\Delta m_t| + |\Delta m_t|^2)). \quad (3.84)$$

The first point is to be able to replace the term $p_{\theta_{2,0+t}}(\Delta m_t)$ by $p_{\theta_{2,0}}(\Delta m_0)$, namely to prove that there exists some constant C'_5 , independent of ε such that for all $t \in [0, \frac{2T}{\delta}]$

$$|p_{\theta_{2,0+t}}(\Delta m_t) - p_{\theta_{2,0}}(\Delta m_0)| \leq C'_5 (\delta^2|\Delta\theta_0| + \varepsilon). \quad (3.85)$$

Indeed, noting first that $p_{\theta_{2,0+t}}(\Pi^\delta(\theta_{2,0} + t, \theta_{2,0})\Delta m_0) = p_{\theta_{2,0}}(\Delta m_0)$, we obtain from (3.56) and (3.47) that for $t \in [0, \frac{2T}{\delta}]$

$$\begin{aligned} |p_{\theta_{2,0+t}}(\Delta m_t) - p_{\theta_{2,0}}(\Delta m_0)| &= |p_{\theta_{2,0+t}}(\Delta m_t) - p_{\theta_{2,0+t}}(\Pi^\delta(\theta_{2,0} + t, \theta_{2,0})\Delta m_0)| \\ &\leq C_{\Pi}(c_1 + c_2 + c_3)\kappa_3\delta^{2-\beta}tW(\mu_{1,0}, \mu_{2,0}) \\ &\quad + C_{\Pi}\delta \int_0^t c_4 \left(\left(\mathbf{E} \left[\left(\Delta \tilde{X}_s \right)^2 \right] \right)^{\frac{1}{2}} + b \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_s \right)^2 \right] \right)^{\frac{1}{2}} \right) ds. \end{aligned} \quad (3.86)$$

By definition of the coupling π_0 in (3.79) and the assumption $W(\tilde{\mu}_{1,0}, \tilde{\mu}_{2,0}) \leq \kappa_7\delta^2|\Delta\theta_0|$, we have $\left(\mathbf{E} \left[\left(\Delta \tilde{X}_0 \right)^2 \right] \right)^{\frac{1}{2}} \leq \kappa_7\delta^2|\Delta\theta_0| + \varepsilon$ and $b \left(\mathbf{E} \left[\left(\Delta \tilde{Y}_0 \right)^2 \right] \right)^{\frac{1}{2}} \leq \kappa_7\delta^2|\Delta\theta_0| + \varepsilon$. Hence, using (3.48), (3.49), the integral term in (3.86) can be bounded above by

$$C_5\delta t (\delta^2|\Delta\theta_0| + \varepsilon)$$

for some constant $C_5 > 0$, independent of ε . Thus, (3.85) is a direct consequence of (3.82).

Secondly, remark that

$$\Delta\theta_0 = \frac{1}{\delta} (p_{\theta_{2,0}}(\Delta m_0) + O(\delta|\Delta m_0| + |\Delta m_0|^2)), \quad (3.87)$$

and that (3.72) implies that $|\Delta m_t| \leq C_6|\Delta m_0|$ for some constant C_6 . Gathering these estimates, (3.84) and (3.85), we obtain (recall that $|\Delta m_0| \leq 2\delta|\dot{\gamma}||\Delta\theta_0|$ and $|\Delta\theta_0| \leq \delta^{-\frac{1}{2}}$):

$$\Delta\theta_t - \Delta\theta_0 = O\left(\delta^{\frac{1}{2}}|\Delta\theta_0| + \frac{\varepsilon}{\delta}\right). \quad (3.88)$$

Since obviously neither $\Delta\theta_t$ nor $\Delta\theta_0$ depend on ε , one can make $\varepsilon \searrow 0$ in the previous estimate and obtain (3.76) with a κ_9 that does not depend on κ_8 , for δ small enough (depending on κ_8).

It remains to prove (3.77) and (3.78). Using (3.79), (3.82) and the assumption on the initial condition X_0 into (3.48), we obtain

$$\left(\mathbf{E} \left[\left(\Delta \tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \left(e^{-(K-\kappa_4\delta)t}\kappa_7 + 2\kappa_5|\dot{\gamma}| \right) \delta^2|\Delta\theta_0| + \varepsilon \left(e^{-(K-\kappa_4\delta)t} + \kappa_5\delta^{1-\beta} \right).$$

Using (3.76), we have

$$|\Delta\theta_0| \leq \frac{|\Delta\theta_t|}{1 - \kappa_9\delta^{\frac{1-\beta}{2}}} \quad (3.89)$$

which gives

$$\left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \left(e^{-(K-\kappa_4\delta)t} + \frac{2\kappa_5|\dot{\gamma}|}{\kappa_7} \right) \kappa_7\delta^2 \frac{|\Delta\theta_t|}{1 - \kappa_9\delta^{\frac{1-\beta}{2}}} + \varepsilon \left(e^{-(K-\kappa_4\delta)t} + \kappa_5\delta^{1-\beta} \right). \quad (3.90)$$

Choosing $\delta \leq \frac{1}{2\kappa_9}$, we have

$$\left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \left(e^{-(K-\kappa_4\delta)t} + \frac{2\kappa_5|\dot{\gamma}|}{\kappa_7} \right) 2\kappa_7\delta^2 |\Delta\theta_t| + \varepsilon \left(e^{-(K-\kappa_4\delta)t} + \kappa_5\delta^{1-\beta} \right). \quad (3.91)$$

Recalling that $\kappa_7 = 16\kappa_5|\dot{\gamma}|$ and choosing $\delta \leq \frac{K}{2\kappa_4}$ and $\delta \leq \frac{KT}{6\ln 2}$ (so that $e^{-\frac{K}{2}\frac{T}{\delta}} \leq \frac{1}{8}$), we obtain for $t \in [\frac{T}{\delta}, \frac{2T}{\delta}]$,

$$\left(\mathbf{E} \left[\left(\Delta\tilde{X}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \frac{\kappa_7}{2}\delta^2 |\Delta\theta_t| + \varepsilon \left(e^{-(K-\kappa_4\delta)t} + \kappa_5\delta^{1-\beta} \right). \quad (3.92)$$

Concerning (3.49), by similar arguments, we have

$$b \left(\mathbf{E} \left[\left(\Delta\tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \left[e^{-\frac{b\delta}{c}t} (1 + b\kappa_6\delta) + \frac{2\kappa_5|\dot{\gamma}|}{\kappa_7} \right] 2\kappa_7\delta^2 |\Delta\theta_t| + \varepsilon \left(\kappa_5\delta^{1-\beta} + e^{-\frac{b\delta}{c}t} (1 + b\kappa_6\delta) \right), \quad (3.93)$$

which gives, by definition of κ_7 and T (recall (3.46)) and choosing $\delta \leq \frac{1}{b\kappa_6}$, for $t \in [\frac{T}{\delta}, \frac{2T}{\delta}]$:

$$b \left(\mathbf{E} \left[\left(\Delta\tilde{Y}_t \right)^2 \right] \right)^{\frac{1}{2}} \leq \frac{\kappa_7}{2}\delta^2 |\Delta\theta_t| + \varepsilon \left(\kappa_5\delta^{1-\beta} + e^{-\frac{b\delta}{c}t} (1 + b\kappa_6\delta) \right). \quad (3.94)$$

Similar estimates are *a fortiori* valid for the infimum $W(\tilde{\mu}_{1,t}, \tilde{\mu}_{2,t})$ and letting $\varepsilon \searrow 0$, we obtain the result (3.77).

Now, recalling Lemma 3.2,

$$\left| p_{\theta_{1,t}}^\perp(m_{1,t} - \gamma_{\theta_{1,t}}) - p_{\theta_{2,t}}^\perp(m_{2,t} - \gamma_{\theta_{2,t}}) \right| \leq 2C_\Pi\kappa_2\delta, \quad (3.95)$$

so that (3.78) is directly valid for $|\Delta\theta_t| \geq 2C_\Pi\kappa_2\delta^{\frac{3}{2}}$. Suppose now that $|\Delta\theta_t| \leq 2C_\Pi\kappa_2\delta^{\frac{3}{2}}$. Relying again on Lemma 3.2, and using the fact that $p_{\theta_{2,t}}^\perp(\dot{\gamma}_{\theta_{2,t}}) = 0$, we have for $t \in [0, \frac{2T}{\delta}]$,

$$p_{\theta_{1,t}}^\perp(m_{1,t} - \gamma_{\theta_{1,t}}) - p_{\theta_{2,t}}^\perp(m_{2,t} - \gamma_{\theta_{2,t}}) = p_{\theta_{2,t}}^\perp(\Delta m_t) + \delta^2 O(|\Delta\theta_t|) + \delta^2 O(|\Delta\theta_t|^2), \quad (3.96)$$

and since $|\theta_{2,t} - (\theta_{2,0} + t)| \leq C_4\delta t$ and

$$|\Delta m_t| \leq \delta^{-\beta} W(\mu_{1,t}, \mu_{2,t}) \leq \delta^{-\beta} \kappa_3 W(\mu_{1,0}, \mu_{2,0}) \leq 2\delta|\dot{\gamma}||\Delta\theta_0|, \quad (3.97)$$

we obtain

$$p_{\theta_{1,t}}^\perp(m_{1,t} - \gamma_{\theta_{1,t}}) - p_{\theta_{2,t}}^\perp(m_{2,t} - \gamma_{\theta_{2,t}}) = p_{\theta_{2,0+t}}^\perp(\Delta m_t) + O\left(\delta^{\frac{3}{2}}|\Delta\theta_t|\right). \quad (3.98)$$

Similarly,

$$p_{\theta_{1,0}}^\perp(m_{1,0} - \gamma_{\theta_{1,0}}) - p_{\theta_{2,0}}^\perp(m_{2,0} - \gamma_{\theta_{2,0}}) = p_{\theta_{2,0}}^\perp(\Delta m_0) + O\left(\delta^{\frac{3}{2}}|\Delta\theta_t|\right), \quad (3.99)$$

and using the decomposition (3.52) and relying on similar estimates as made above,

$$p_{\theta_{2,0+t}}^\perp(\Delta m_t) = e^{-\lambda\delta t} p_{\theta_{2,0}}^\perp(\Delta m_0) + O\left(\delta^{\frac{3}{2}}|\Delta\theta_t|\right). \quad (3.100)$$

So, using again (3.76), there exists a constant C_7 that does not depend on κ_8 such that

$$\left| p_{\theta_{1,t}}^\perp(m_{1,t} - \gamma_{\theta_{1,t}}) - p_{\theta_{2,t}}^\perp(m_{2,t} - \gamma_{\theta_{2,t}}) \right| \leq \left(2e^{-\lambda\delta t} + \frac{C_7}{\kappa_8} \right) \kappa_8 \delta^{\frac{3}{2}} |\Delta\theta_t|, \quad (3.101)$$

which implies (3.78), taking $\kappa_8 \geq 2C_7$, and recalling (3.46). \square

4. FIXED POINT

Recall the definitions of the constants κ_i , $i = 0, \dots, 8$ appearing in Section 2 and Section 3. For $\delta \leq \delta_4$ (recall Lemma 3.4), consider the space $\mathcal{F} = \mathcal{F}(\delta)$ composed of the functions $f : \mathbb{S}_\delta \rightarrow \mathcal{P}_2$ that satisfy

(1)

$$\int_{\mathbb{R}^2} x^6 f(\theta)(dz) \leq \kappa_0^x \quad \text{and} \quad \int_{\mathbb{R}^2} y^6 f(\theta)(dz) \leq \kappa_0^y, \quad \text{for all } \theta \in \mathbb{S}_\delta, \quad (4.1)$$

(2)

$$W(\widetilde{f(\theta)}, q_0) \leq \kappa_1 \delta, \quad \text{for all } \theta \in \mathbb{S}_\delta, \quad (4.2)$$

(3)

$$\int_{\mathbb{R}^2} z f(\theta)(dz) \in E_\theta(\kappa_2 \delta), \quad \text{for all } \theta \in \mathbb{S}_\delta, \quad (4.3)$$

(4)

$$W(\widetilde{f(\theta_1)}, \widetilde{f(\theta_2)}) \leq \kappa_7 \delta^2 |\theta_1 - \theta_2|, \quad \text{for all } \theta_1, \theta_2 \in \mathbb{S}_\delta, \quad (4.4)$$

(5)

$$\left| p_{\theta_1}^\perp \left(\int_{\mathbb{R}^2} z f(\theta_1)(dz) - \gamma_{\theta_1} \right) - p_{\theta_2}^\perp \left(\int_{\mathbb{R}^2} z f(\theta_2)(dz) - \gamma_{\theta_2} \right) \right| \leq \kappa_8 \delta^{\frac{3}{2}} |\theta_1 - \theta_2|, \quad (4.5)$$

for all $\theta_1, \theta_2 \in \mathbb{S}_\delta$.

We define, for $\theta \in \mathbb{S}_\delta$, the distance $W_\theta = W_\theta(\delta, b, \beta)$ on probability measures ν that satisfy $\int_{\mathbb{R}^2} z \nu(dz) \in E_\theta(\kappa_2 \delta)$ as follows:

$$W_\theta(\nu_1, \nu_2) := \max \left\{ \delta^\beta \left| p_\theta^\perp \left(\int_{\mathbb{R}^2} z d\nu_1(z) - \int_{\mathbb{R}^2} z d\nu_2(z) \right) \right|, W(\tilde{\nu}_1, \tilde{\nu}_2) \right\}, \quad (4.6)$$

where W is defined in (2.15). We consider the distance $\text{dist}_{\mathcal{F}}$ on \mathcal{F} defined as

$$\text{dist}_{\mathcal{F}}(f_1, f_2) = \sup_{\theta \in \mathbb{S}_\delta} W_\theta(f_1(\theta), f_2(\theta)). \quad (4.7)$$

Remark that, on \mathcal{F} , $\frac{1}{C_\Pi} W \leq W_\theta \leq C_\Pi W$ and this distance is thus equivalent to the distance $\sup_{\theta \in \mathbb{S}_\delta} W_2(f_1(\theta), f_2(\theta))$. \mathcal{F} is then a closed subspace of $C(\mathbb{S}_\delta, \mathcal{P}_2)$ (where \mathcal{P}_2 is endowed with the standard Wasserstein-2 distance), and is thus complete when endowed with $\text{dist}_{\mathcal{F}}$.

For $t \in [0, \frac{2T}{\delta}]$ we consider the map $g_{t,f} : \mathbb{S}_\delta \rightarrow \mathbb{S}_\delta$ as

$$g_{t,f}(\theta) = \text{proj} \left(\int_{\mathbb{R}^2} z \mu_t(dz) \right), \quad (4.8)$$

where μ_t is the solution of (1.5) with $\mu_0 = f(\theta)$. Lemma 3.4 shows that, for δ small enough, $g_{t,f}$ is a bijection for any $f \in \mathcal{F}$. We consider the map $\Phi_t : \mathcal{F} \rightarrow C(\mathbb{S}_\delta, \mathcal{P}_2)$ defined as

$$\Phi_t(\theta) = \mu_{\frac{t}{\delta}}, \quad (4.9)$$

where μ is the solution of (1.5) with $\mu_0 = f\left(g_{t,f}^{-1}(\theta)\right)$.

Lemma 4.1. *If $\delta \leq \delta_4$ and $f \in \mathcal{F}$, then $\Phi_t(f) \in \mathcal{F}$ for all $t \in [T, 2T]$.*

Proof. This is a direct consequence of Lemma 2.1, Lemma 2.4, Lemma 3.2, Lemma 3.3 and Lemma 3.4. \square

Lemma 4.2. *There exist $\delta_5 > 0$ and $\kappa_{10} > 0$ such that if $\delta \leq \delta_5$ and $f_1, f_2 \in \mathcal{F}$, then for all $t \in [T, 2T]$ we have*

$$\text{dist}_{\mathcal{F}}(\Phi_t(f_1), \Phi_t(f_2)) \leq \left(\max \left\{ e^{-\lambda t}, e^{-\frac{b}{c}t} \right\} + \kappa_{10} \left(\delta^\beta + \delta^{1-\beta} \right) \right) \text{dist}_{\mathcal{F}}(f_1, f_2). \quad (4.10)$$

Proof. Fix a $t \in [T, 2T]$. For $\theta \in \mathbb{S}_\delta$, consider solutions $\mu_{i,s}$ to (1.5) starting from $\mu_{i,0} = f_i(\theta_{i,0})$ and such that $\theta_{i,\frac{t}{\delta}} = \theta$, $i = 1, 2$. Our aim is to bound $W_\theta\left(\mu_{1,\frac{t}{\delta}}, \mu_{2,\frac{t}{\delta}}\right)$. To do this we consider also the solution $\mu_{3,s}$ starting from $f_1(\theta_{2,0})$. Remark first that

$$W_\theta\left(\tilde{\mu}_{1,\frac{t}{\delta}}, \tilde{\mu}_{2,\frac{t}{\delta}}\right) = W\left(\tilde{\mu}_{1,\frac{t}{\delta}}, \tilde{\mu}_{2,\frac{t}{\delta}}\right) \leq W\left(\tilde{\mu}_{1,\frac{t}{\delta}}, \tilde{\mu}_{3,\frac{t}{\delta}}\right) + W\left(\tilde{\mu}_{3,\frac{t}{\delta}}, \tilde{\mu}_{2,\frac{t}{\delta}}\right). \quad (4.11)$$

On one hand,

$$\begin{aligned} W\left(\tilde{\mu}_{1,\frac{t}{\delta}}, \tilde{\mu}_{3,\frac{t}{\delta}}\right) &= W\left(\widetilde{\Phi_t(f_1)}\left(\theta_{1,\frac{t}{\delta}}\right), \widetilde{\Phi_t(f_1)}\left(\theta_{3,\frac{t}{\delta}}\right)\right) \\ &\leq \kappa_7 \delta^2 \left| \theta_{1,\frac{t}{\delta}} - \theta_{3,\frac{t}{\delta}} \right|, \quad (\text{by Lemma 4.1}) \\ &= \kappa_7 \delta^2 \left| \theta_{2,\frac{t}{\delta}} - \theta_{3,\frac{t}{\delta}} \right| \\ &\leq C_{\text{proj}} \kappa_7 \delta \left| m_{2,\frac{t}{\delta}} - m_{3,\frac{t}{\delta}} \right|, \quad (\text{using (3.13)}) \\ &\leq C_{\text{proj}} \kappa_7 \delta^{1-\beta} W\left(\mu_{2,\frac{t}{\delta}}, \mu_{3,\frac{t}{\delta}}\right) \\ &\leq C_{\text{proj}} \kappa_3 \kappa_7 \delta^{1-\beta} W(\mu_{2,0}, \mu_{3,0}), \quad (\text{by Lemma 3.3}) \\ &\leq C_{\Pi} C_{\text{proj}} \kappa_3 \kappa_7 \delta^{1-\beta} W_\theta(f_2(\theta_{2,0}), f_1(\theta_{2,0})), \end{aligned} \quad (4.12)$$

while on the other hand, Lemma 3.3 implies that

$$\begin{aligned} W\left(\tilde{\mu}_{3,\frac{t}{\delta}}, \tilde{\mu}_{2,\frac{t}{\delta}}\right) &\leq \max\left(e^{-(K-\delta\kappa_4)\frac{t}{\delta}}, e^{-\frac{b}{c}t}(1+b\kappa_6\delta)\right) W(\tilde{\mu}_{3,0}, \tilde{\mu}_{2,0}) \\ &\quad + \kappa_5 \delta^{1-\beta} W(\mu_{3,0}, \mu_{2,0}) \\ &\leq \left(e^{-\frac{b}{c}t} + 2C_{\Pi} \kappa_5 \delta^{1-\beta}\right) W_\theta(f_1(\theta_{2,0}), f_2(\theta_{2,0})), \end{aligned} \quad (4.13)$$

where we have taken δ small enough. So, for δ small enough,

$$W_\theta\left(\tilde{\mu}_{1,\frac{t}{\delta}}, \tilde{\mu}_{2,\frac{t}{\delta}}\right) \leq \left(e^{-\frac{b}{c}t} + O\left(\delta^{1-\beta}\right)\right) \text{dist}_{\mathcal{F}}(f_1, f_2). \quad (4.14)$$

Similarly, we have the decomposition, since $\theta_{1, \frac{t}{\delta}} = \theta_{2, \frac{t}{\delta}} = \theta$,

$$\begin{aligned} \left| p_{\theta}^{\perp} \left(m_{1, \frac{t}{\delta}} - m_{2, \frac{t}{\delta}} \right) \right| &= \left| p_{\theta_{1, \frac{t}{\delta}}}^{\perp} \left(m_{1, \frac{t}{\delta}} - \gamma_{\theta_{1, \frac{t}{\delta}}} \right) - p_{\theta_{2, \frac{t}{\delta}}}^{\perp} \left(m_{2, \frac{t}{\delta}} - \gamma_{\theta_{2, \frac{t}{\delta}}} \right) \right| \\ &\leq \left| p_{\theta_{1, \frac{t}{\delta}}}^{\perp} \left(m_{1, \frac{t}{\delta}} - \gamma_{\theta_{1, \frac{t}{\delta}}} \right) - p_{\theta_{3, \frac{t}{\delta}}}^{\perp} \left(m_{3, \frac{t}{\delta}} - \gamma_{\theta_{3, \frac{t}{\delta}}} \right) \right| \\ &\quad + \left| p_{\theta_{3, \frac{t}{\delta}}}^{\perp} \left(m_{3, \frac{t}{\delta}} - \gamma_{\theta_{3, \frac{t}{\delta}}} \right) - p_{\theta_{2, \frac{t}{\delta}}}^{\perp} \left(m_{2, \frac{t}{\delta}} - \gamma_{\theta_{2, \frac{t}{\delta}}} \right) \right|. \end{aligned} \quad (4.15)$$

On one hand, using similar bounds as above, we get

$$\begin{aligned} \left| p_{\theta_{1, \frac{t}{\delta}}}^{\perp} \left(m_{1, \frac{t}{\delta}} - \gamma_{\theta_{1, \frac{t}{\delta}}} \right) - p_{\theta_{3, \frac{t}{\delta}}}^{\perp} \left(m_{3, \frac{t}{\delta}} - \gamma_{\theta_{3, \frac{t}{\delta}}} \right) \right| \\ \leq \kappa_8 \delta^{\frac{3}{2}} \left| \theta_{1, \frac{t}{\delta}} - \theta_{3, \frac{t}{\delta}} \right| \\ \leq C_{\Pi} C_{\text{proj}} \kappa_3 \kappa_8 \delta^{\frac{1}{2} - \beta} W_{\theta}(f_2(\theta_{2,0}), f_1(\theta_{2,0})). \end{aligned} \quad (4.16)$$

On the other hand, either $\left| \theta_{3, \frac{t}{\delta}} - \theta_{2, \frac{t}{\delta}} \right| \geq \delta^{-\frac{1}{2}}$, and in this case relying on Lemma 2.4 we have

$$\begin{aligned} \left| p_{\theta_{3, \frac{t}{\delta}}}^{\perp} \left(m_{3, \frac{t}{\delta}} - \gamma_{\theta_{3, \frac{t}{\delta}}} \right) - p_{\theta_{2, \frac{t}{\delta}}}^{\perp} \left(m_{2, \frac{t}{\delta}} - \gamma_{\theta_{2, \frac{t}{\delta}}} \right) \right| \\ \leq 2\kappa_2 \delta \\ \leq 2\kappa_2 \delta^{\frac{3}{2}} \left| \theta_{3, \frac{t}{\delta}} - \theta_{2, \frac{t}{\delta}} \right| \\ \leq 2C_{\Pi} C_{\text{proj}} \kappa_2 \kappa_3 \delta^{\frac{1}{2} - \beta} W_{\theta}(f_2(\theta_{2,0}), f_1(\theta_{2,0})), \end{aligned} \quad (4.17)$$

or $\left| \theta_{3, \frac{t}{\delta}} - \theta_{2, \frac{t}{\delta}} \right| \leq \delta^{-\frac{1}{2}}$, and with similar arguments as used in the proofs of Lemma 3.3 and Lemma 3.4, we have

$$\begin{aligned} \left| p_{\theta_{3, \frac{t}{\delta}}}^{\perp} \left(m_{3, \frac{t}{\delta}} - \gamma_{\theta_{3, \frac{t}{\delta}}} \right) - p_{\theta_{2, \frac{t}{\delta}}}^{\perp} \left(m_{2, \frac{t}{\delta}} - \gamma_{\theta_{2, \frac{t}{\delta}}} \right) \right| \\ = \left| p_{\theta_{2,0+t}}^{\perp} \left(m_{3, \frac{t}{\delta}} - m_{2, \frac{t}{\delta}} \right) \right| + O\left(\delta \left| \theta_{3, \frac{t}{\delta}} - \theta_{2, \frac{t}{\delta}} \right|\right) + O\left(\left| \theta_{3, \frac{t}{\delta}} - \theta_{2, \frac{t}{\delta}} \right|^2\right) \\ = \left| p_{\theta_{2,0+t}}^{\perp} \left(m_{3, \frac{t}{\delta}} - m_{2, \frac{t}{\delta}} \right) \right| + O\left(\delta^{\frac{1}{2}} W(f_2(\theta_{2,0}), f_1(\theta_{2,0}))\right). \end{aligned} \quad (4.18)$$

Relying on the decomposition (3.52) and its following estimates (considering some coupling $\{(X_{3,0}, Y_{3,0}), (X_{2,0}, Y_{2,0})\} \sim \pi_0 \in \mathcal{C}(\mu_{3,0}, \mu_{2,0})$ of the initial condition such that $\Lambda(\pi_0) < W(\mu_{3,0}, \mu_{2,0}) + \varepsilon$),

$$\begin{aligned} \left| p_{\theta_{2,0+t}}^{\perp} \left(m_{3, \frac{t}{\delta}} - m_{2, \frac{t}{\delta}} \right) \right| \\ \leq e^{-\lambda t} \left| p_{\theta_{2,0}}^{\perp} (m_{3,0} - m_{2,0}) \right| + C_{\Pi} (c_1 + c_2 + c_3) \kappa_3 t \delta^{1-\beta} W(f_1(\theta_{2,0}), f_2(\theta_{2,0})) \\ \quad + C_{\Pi} c_4 \delta \int_0^{\frac{t}{\delta}} \left(\mathbb{E} \left[\left(\tilde{X}_{3,s} - \tilde{X}_{2,s} \right)^2 \right]^{\frac{1}{2}} + b \mathbb{E} \left[\left(\tilde{Y}_{3,s} - \tilde{Y}_{2,s} \right)^2 \right]^{\frac{1}{2}} \right) ds \\ \leq \left(\delta^{-\beta} e^{-\lambda t} + C \left(1 + \delta^{1-\beta} \right) (1 + \varepsilon) \right) W_{\theta}(f_1(\theta_{2,0}), f_2(\theta_{2,0})), \end{aligned} \quad (4.19)$$

for some positive constant C , where we have relied on Lemma 3.3 to deal with the integral term. Making ε going to 0 and gathering these estimates we get for δ small enough:

$$\delta^\beta \left| p_\theta^\perp \left(m_{1, \frac{t}{\delta}} - m_{2, \frac{t}{\delta}} \right) \right| \leq \left(e^{-\lambda t} + O\left(\delta^{\frac{1}{2}} + \delta^\beta\right) \right) \text{dist}_{\mathcal{F}}(f_1, f_2), \quad (4.20)$$

which concludes the proof. \square

We now have all the ingredients needed to prove Theorem 1.4.

Proof of Theorem 1.4. By Lemma 4.1 we have $\Phi_T(\mathcal{F}) \subset \mathcal{F}$. Moreover, recalling (3.46), Lemma 4.2 implies that Φ_T admits a unique fixed-point f_0 in \mathcal{F} . The point is now to show that $\Phi_t(f_0) = f_0$ for all $t \geq 0$. It is clear that $\Phi_{kT}(f_0) = f_0$ for all $k \in \mathbb{N}$, by the semi-group property, and it remains then to prove that $\Phi_t(f_0) = f_0$ for $t \in (0, T)$. But, fixing $t \in (0, T)$, $\Phi_t(f_0) \in \mathcal{F}$ by Lemma 4.1 and we have $\Phi_t(f_0) = \Phi_t(\Phi_T(f_0)) = \Phi_T(\Phi_t(f_0))$, so that $\Phi_t(f_0) = f_0$ by uniqueness of the fixed-point of Φ_T on \mathcal{F} .

This proves that $\mathcal{C}_\delta = \{f_0(\theta) : \theta \in \mathbb{S}_\delta\}$ is a one-dimensional invariant manifold for (1.5), and the estimates made in the proof of Lemma 2.4, showing that if $\mu_0 = f_0(\theta_0)$ then $\theta_t = 1 + O(\delta)$, imply that \mathcal{C}_δ defines in fact a periodic solution for (1.5). Moreover from (4.2) and (4.3) we deduce that for all $\theta \in \mathbb{S}_\delta$ we have $W(\widetilde{f_0(\theta)}, \widetilde{q_0}) \leq \kappa_1 \delta$, and $|\int_{\mathbb{R}^2} z f_0(dz) - \gamma_\theta| = O(\delta)$, which means that $W_2(f(\theta), q_{\gamma_\theta}) = O(\delta)$.

To prove the contraction property, first remark that, by equivalence of distances, if $\text{dist}_{W_2}(\mu, \mathcal{C}_\delta) \leq c\delta$ with c small enough, then for $\theta_0 = \text{proj}(m_0)$ we have $m_0 \in E_{\theta_0}(\kappa_2 \delta)$ and $W(\widetilde{\mu_0}, q_0) \leq \kappa_1 \delta$, and for f defined as

$$\widetilde{f(\theta)} = \widetilde{\mu_0}, \quad \text{and} \quad \int_{\mathbb{R}^2} z f(\theta)(dz) = \gamma_\theta + N(\theta, \theta_0)(m_0 - \gamma_{\theta_0}), \quad (4.21)$$

we have $f \in \mathcal{F}$. Now, on one hand, for $t \in [0, T]$, Lemma 3.3 implies that if y_t^δ is the periodic solution defining \mathcal{C}_δ , with $y_0^\delta = f_0(\theta_0)$, then $W(\mu_t, y_t^\delta) \leq \kappa_3 W(\mu_0, y_0^\delta)$, and thus for $\theta \in [0, T]$:

$$\text{dist}_{W_2}(\mu_t, \mathcal{C}_\delta) \leq 3\kappa_3 \delta^{-\beta} W(\mu_0, f_0(\theta_0)) \leq \frac{3\kappa_3}{C_{\text{II}}} \delta^{-\beta} W_{\theta_0}(\mu_0, f_0(\theta_0)). \quad (4.22)$$

On the other hand, since $\mu_t = \Phi_t(f)(\theta_t)$ and relying on the proof on Lemma 4.2 and on a basic recursion, we get that for $t \geq T$:

$$\text{dist}_{W_2}(\mu_t, \mathcal{C}_\delta) \leq \frac{3\kappa_3}{C_{\text{II}}} \delta^{-\beta} W_{\theta_t}(\Phi_t(f)(\theta_t), (f_0)(\theta_t)) \leq \frac{3\kappa_3}{C_{\text{II}}} \delta^{-\beta} e^{-\lambda(\theta)t} W_{\theta_0}(\mu_0, f_0(\theta_0)), \quad (4.23)$$

where we have defined (using the bound $e^{-at} + xt \leq e^{-(a-xe^{at})t}$):

$$\lambda(\theta) := \min\left(\lambda, \frac{b}{c}\right) - \frac{\kappa_{10} e^{2T \min(\lambda, \frac{b}{c})}}{T} (\delta^\beta + \delta^{1-\beta}). \quad (4.24)$$

It remains thus to bound $W_{\theta_0}(\mu_0, f_0(\theta_0))$ by $\text{dist}_{W_2}(\mu_0, \mathcal{C}_\delta)$. To do this we show that for a constant c we have $W_{\theta_0}(\mu_0, f(\theta_0)) \leq cW(\mu_0, f_0(\theta_0))$ for all $\theta \in \mathbb{S}_\delta$. But we have on one hand, since $m_0 \in E_{\theta_0}(\kappa_2 \delta)$,

$$\left| p_{\theta_0}^\perp (m_0 - \gamma_{\theta_0}) \right| \leq C_{\text{proj}} |m_0 - \gamma_{\theta_0}| \leq C_{\text{proj}} |m_0 - \gamma_\theta| + C_{\text{proj}} |\gamma_\theta - \gamma_{\theta_0}|, \quad (4.25)$$

and an application of Lemma 3.2 leads to

$$\theta_0 - \theta = \frac{1}{\delta} p_\theta (m_0 - \gamma_\theta) + \frac{1}{\delta} O\left(|m_0 - \gamma_\theta|^2\right). \quad (4.26)$$

So, for a constant C that does not depend on δ , $|\gamma_\theta - \gamma_{\theta_0}| = |\gamma_{\delta\theta}^1 - \gamma_{\delta\theta_0}^1| \leq C|m_0 - \gamma_\theta|$, and thus

$$\left| p_{\theta_0}^\perp(m_0 - \gamma_{\theta_0}) \right| \leq C'|m_0 - \gamma_\theta|, \quad (4.27)$$

for some constant C' that does not depend on δ . On the other hand,

$$W\left(\widetilde{\mu}_0, \widetilde{f_0}(\theta_0)\right) \leq W\left(\widetilde{\mu}_0, \widetilde{f_0}(\theta)\right) + W\left(\widetilde{f_0}(\theta), \widetilde{f_0}(\theta_0)\right), \quad (4.28)$$

and $W\left(\widetilde{f_0}(\theta), \widetilde{f_0}(\theta_0)\right) \leq \kappa_7\delta^2|\theta - \theta_0| \leq C''|m_0 - \gamma_\theta|$. Gathering all these estimates, we obtain $W_{\theta_0}(\mu_0, f(\theta_0)) \leq cW(\mu_0, f_0(\theta))$, with $c = \max(C', 1 + C'')$.

Finally, for all $t \geq 0$ and $\theta \in \mathbb{S}_\delta$ we get:

$$\text{dist}_{W_2}(\mu_t, \mathcal{C}_\delta) \leq C(\delta)e^{-\lambda(\delta)t}W_2(\mu_0, f_0(\theta)), \quad (4.29)$$

with $C(\delta) = \frac{3c}{C_{\text{II}}^2 \min(1, b^{-1})} \delta^{-\beta} \max(\kappa_3 e^{\lambda(\delta)T}, 1)$, which concludes the proof. \square

APPENDIX A. TECHNICAL LEMMA

Lemma A.1. *Let v be a continuously differentiable function on $[0, +\infty)$ such that $v(t) \geq 0$ for all $t \geq 0$. Suppose that there exists $\alpha, \beta, k > 0$ and $\zeta \geq 0$ such that*

$$v'(t) \leq -\alpha v(t) + \left(e^{-kt}\zeta + \beta\right) \sqrt{v(t)}. \quad (A.1)$$

Then, for all $t \geq 0$,

$$v(t) \leq \max\left(v(0), \frac{(\zeta + \beta)^2}{\alpha^2}\right), \quad (A.2)$$

and if $2k > \alpha$ or $\zeta = 0$, then

$$v(t) \leq \left(e^{-\alpha t/2} \left(\sqrt{v(0)} + \frac{\zeta}{2k - \alpha}\right) + \frac{\beta}{\alpha}\right)^2. \quad (A.3)$$

Proof of Lemma A.1. The first is a consequence of the fact that $v(t)$ is always non-increasing unless $\sqrt{v(t)} \leq \frac{e^{-kt}\zeta + \beta}{\alpha} \leq \frac{\zeta + \beta}{\alpha}$. We now prove the second inequality: let $t \geq 0$ such that $v(t) > 0$ and consider the maximal interval $I := (t^-, t^+)$ (which is non empty by continuity of v) containing t such that $v(u) > 0$ on I . Consider the function $f(u) := \left(\alpha\sqrt{v(u)} + \frac{\alpha}{2k - \alpha}e^{-ku}\zeta - \beta\right) e^{\alpha u/2}$. For all $u \in I$ we have

$$f'(u) = \frac{\alpha e^{\alpha u/2}}{2\sqrt{v(u)}} \left(\alpha v(u) - \left(e^{-ku}\zeta + \beta\right) \sqrt{v(u)} + v'(u)\right) \leq 0, \quad (A.4)$$

so that f is nonincreasing on I . Consider now the solution w of the equation $w'(t) = -\alpha w(t) + \left(e^{-kt}\zeta + \beta\right) \sqrt{w(t)}$ such that $w(t^-) = v(t^-)$. Then, relying on the same calculations, $g := u \mapsto \left(\alpha\sqrt{w(u)} + \frac{\alpha}{2k - \alpha}e^{-ku}\zeta - \beta\right) e^{\alpha u/2}$ is constant on I , and equal to $g(t^-)$. This means that $f(u) \leq f(t^-) = g(t^-) = g(u)$ and, by definition of f and g , this implies that

$$\begin{aligned} v(u) \leq w(u) &= \left(\frac{e^{-\alpha u/2}}{\alpha} g(t^-) - \frac{e^{-ku}}{2k - \alpha} \zeta + \frac{\beta}{\alpha}\right)^2 \\ &= \left(e^{-\alpha(u-t^-)/2} \sqrt{v(t^-)} + \frac{e^{-\alpha(u-t^-)/2} e^{-kt^-} - e^{-ku}}{2k - \alpha} \zeta + \left(1 - e^{\alpha(u-t^-)/2}\right) \frac{\beta}{\alpha}\right)^2. \end{aligned} \quad (A.5)$$

We have now two possibilities: either $t^- = 0$ and then

$$v(u) \leq \left(e^{-\alpha u/2} \sqrt{v(0)} + \frac{e^{-\alpha u/2} - e^{-ku}}{2k - \alpha} \zeta + \left(1 - e^{-\alpha u/2}\right) \frac{\beta}{\alpha} \right)^2, \quad (\text{A.6})$$

or $t^- > 0$, and by continuity of v we have $v(t^-) = 0$. In this case, recalling that $2k > \alpha$ or $\zeta = 0$, we obtain

$$v(u) \leq \left(\frac{e^{-\alpha u/2}}{2k - \alpha} \zeta + \frac{\beta}{\alpha} \right)^2. \quad (\text{A.7})$$

This proves Lemma A.1. □

ACKNOWLEDGEMENTS

C. Poquet benefited from the support of the ANR-17-CE40-0030 (Entropy, Flows, Inequalities).

REFERENCES

- [1] M. Aleandri and I. G. Minelli. Opinion dynamics with Lotka-Volterra type interactions. *arXiv e-prints*, page arXiv:1811.05937, Nov 2018.
- [2] J. Baladron, D. Fasoli, O. Faugeras, and J. Touboul. Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons. *The Journal of Mathematical Neuroscience*, 2(1):10, 2012.
- [3] P. Bates, K. Lu, and C. Zeng. *Existence and persistence of invariant manifolds for semiflows in Banach space.*, volume 135. Mem. Amer. Math. Soc., 1998.
- [4] N. Berglund and B. Gentz. *Noise-induced phenomena in slow-fast dynamical systems*. Probability and its Applications (New York). Springer-Verlag London, Ltd., London, 2006. A sample-paths approach.
- [5] L. Bertini, G. Giacomin, and C. Poquet. Synchronization and random long time dynamics for mean-field plane rotators. *Probability Theory and Related Fields*, pages 1–61, 2013.
- [6] M. Bossy, O. Faugeras, and D. Talay. Clarification and complement to “Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons”. *J. Math. Neurosci.*, 5:Art. 19, 23, 2015.
- [7] M. Bossy, J. Fontbona, and H. Olivero. Synchronization of stochastic mean field networks of Hodgkin-Huxley neurons with noisy channels. 2018.
- [8] F. Collet, M. Formentin, and D. Tovazzi. Rhythmic behavior in a two-population mean-field Ising model *Physical Review E* 94, 042139, 2016.
- [9] P. Dai Pra, M. Fischer and D. Regoli. A Curie-Weiss model with dissipation. *Journal of Statistical Physics*, 152(1):37–53, 2013.
- [10] S. Ditlevsen and E. Löcherbach. Multi-class oscillating systems of interacting neurons. *Stochastic Processes and their Applications*, 127(6):1840–1869, 2017.
- [11] A. Durmus, A. Eberle, A. Guillin, and R. Zimmer. Uniform propagation of chaos for granular media equations. to appear, 2018.
- [12] N. Fenichel. Persistence and smoothness of invariant manifolds for flows. *Indiana Univ. Math. J.*, 21:193–226, 1972.
- [13] N. Fenichel. Geometric singular perturbation theory for ordinary differential equations. *Journal of Differential Equations*, 31:53–98, 1979.
- [14] R. FitzHugh. Impulses and physiological states in theoretical models of nerve membrane. *Biophysical Journal*, 1(6):445–466, 1961.
- [15] G. Giacomin, E. Luçon, and C. Poquet. Coherence stability and effect of random natural frequencies in populations of coupled oscillators. *Journal of Dynamics and Differential Equations*, 26(2):333–367, 2014.
- [16] G. Giacomin, K. Pakdaman, X. Pellegrin and C. Poquet. Transitions in active rotator systems: invariant hyperbolic manifold approach. *SIAM Journal on Mathematical Analysis*, 44(6):4165–4194, 2012.

- [17] G. Giacomin and C. Poquet. Noise, interaction, nonlinear dynamics and the origin of rhythmic behaviors. *Brazilian Journal of Probability and Statistics*, 29(2):460–493, 2015.
- [18] M. W. Hirsch, C. C. Pugh, and M. Shub. *Invariant manifolds*. Lecture Notes in Mathematics, Vol. 583. Springer-Verlag, Berlin, 1977.
- [19] B. Lindner, J. Garcia-Ojalvo, A. Neiman, and L. Schimansky-Geier. Effects of noise in excitable systems. *Physics Reports*, 392(6):321 – 424, 2004.
- [20] E. Luçon and C. Poquet. Long time dynamics and disorder-induced traveling waves in the stochastic Kuramoto model. *Ann. Inst. Henri Poincaré Probab. Stat.*, 53(3):1196–1240, 2017.
- [21] E. Luçon and C. Poquet. Emergence of oscillatory behaviors for excitable systems with noise and mean-field interaction, a slow-fast dynamics approach. *ArXiv e-prints*, Feb 2018.
- [22] E. Luçon and W. Stannat. Mean field limit for disordered diffusions with singular interactions. *Ann. Appl. Probab.*, 24(5):1946–1993, 2014.
- [23] S. Mischler, C. Quiñinao, and J. Touboul. On a Kinetic Fitzhugh–Nagumo Model of Neuronal Network. *Communications in Mathematical Physics*, 342(3):1001–1042, 2016.
- [24] C. Quiñinao and J. D. Touboul. Clamping and Synchronization in the strongly coupled FitzHugh–Nagumo model. *ArXiv e-prints*, Apr. 2018.
- [25] C. Rocsoreanu, A. Georgescu, and N. Giurgiteanu. *The FitzHugh–Nagumo model*, volume 10 of *Mathematical Modelling: Theory and Applications*. Kluwer Academic Publishers, Dordrecht, 2000. Bifurcation and dynamics.
- [26] M. Scheutzow. Noise can create periodic behavior and stabilize nonlinear diffusions. *Stochastic processes and their applications*, 20(2):323–331, 1985.
- [27] M. Scheutzow. Periodic behavior of the stochastic Brusselator in the mean-field limit. *Probability theory and related fields*, 72(3):425–462, 1986.
- [28] G. R. Sell and Y. You. *Dynamics of evolutionary equations*, volume 143 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2013.
- [29] Sznitman, A.-S. Topics in propagation of chaos. *Ecole d’été de probabilités de Saint-Flour XIX–1989*, Springer, 165–251, 1991.
- [30] G. Teschl. *Ordinary differential equations and dynamical systems*, volume 140 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [31] J. D. Touboul, C. Piette, L. Venance, and G. Bard Ermentrout. Noise-induced synchronization and anti-resonance in excitable systems; Implications for information processing in Parkinson’s Disease and Deep Brain Stimulation. *arXiv e-prints*, page arXiv:1905.01342, May 2019.
- [32] S. Wiggins. *Normally hyperbolic invariant manifolds in dynamical systems*, volume 105 of *Applied Mathematical Sciences*. Springer, 2013.

LABORATOIRE MAP5 (UMR CNRS 8145), UNIVERSITÉ PARIS DESCARTES, SORBONNE PARIS CITÉ,
75270 PARIS, FRANCE, ERIC.LUCON@PARIDESCARTES.FR.

UNIV LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN,
F-69622 VILLEURBANNE, FRANCE, POQUET@MATH.UNIV-LYON1.FR