

# On a finite trigonometric sum related to Dedekind sum

Mouloud Goubi

ABSTRACT. Finite trigonometric sums appear in various branches of Physics, Mathematics and their applications. For  $p, q$  to coprime positive integers and  $r \in \mathbb{Z}$  we consider the finite trigonometric sums involving the product of three trigonometric functions

$$S(p, q, r) = \sum_{k=1}^{p-1} \cos\left(\frac{\pi kr}{p}\right) \csc\left(\frac{\pi kq}{p}\right) \csc\left(\frac{\pi k}{p}\right).$$

In this work we establish a reciprocity law satisfied by  $S(p, q, r)$  in some particular cases. And we compute explicitly the value of the sum

$$S(p, 1, 0) = \sum_{k=1}^{p-1} \csc^2\left(\frac{\pi k}{p}\right).$$

## 1. Introduction and main results

Let us remember the cosecant function  $\csc$  defined by

$$\csc \theta = \frac{1}{\sin \theta}.$$

From which the cosecant numbers are extracted, as the rational coefficients of its power series expansion, a generalization of these numbers is given in the paper [6] of V. Kowalenko.

For  $p, q$  two coprime positive integers and  $r \in \mathbb{Z}$  we consider the following finite trigonometric sum.

$$(1.1) \quad S(p, q, r) = \sum_{k=1}^{p-1} \cos\left(\frac{\pi kr}{p}\right) \csc\left(\frac{\pi kq}{p}\right) \csc\left(\frac{\pi k}{p}\right)$$

Let  $0 \leq \bar{r} \leq 2p - 1$  the represent of the class of  $r$  in  $\mathbb{Z}/2p\mathbb{Z}$ , then

$$S(p, q, r) = S(p, q, \bar{r})$$

---

2010 *Mathematics Subject Classification.* Primary 33B10, 11L03.

*Key words and phrases.* Dedekind sum, finite trigonometric sums, cosecant function.

For example if  $r \equiv 0 \pmod{2p}$ ;

$$S(p, q, r) = \sum_{k=1}^{p-1} \csc\left(\frac{\pi k q}{p}\right) \csc\left(\frac{\pi k}{p}\right)$$

The periodicity of cosine states that there are exactly  $2p$  finite trigonometric sums of this kind for a fixed positive integers  $p, q$ .

If  $q = \bar{q} \pmod{2p}$  then

$$S(p, q, r) = S(p, \bar{q}, r)$$

and for a fixed  $p$  and  $q, r$  are arbitrary integers such that  $q$  is not multiple of  $p$  their exist exactly  $2p(2p-1)$  finite trigonometric sums  $S(p, q, r)$ .

Now if  $r \equiv 0 \pmod{p}$  and  $q \equiv 1 \pmod{2p}$ , we get

$$S(p, q, r) = \sum_{k=1}^{p-1} \csc^2\left(\frac{\pi k}{p}\right),$$

and if  $q = 1$  and  $r \equiv 1 \pmod{p}$  then

$$S(p, q, 0) = \sum_{k=1}^{p-1} (-1)^k \csc^2\left(\frac{\pi k}{p}\right).$$

Finally for  $r = q$  the considered trigonometric sum will be written in the following form

$$\sum_{k=1}^{p-1} \cot\left(\frac{\pi k q}{p}\right) \csc\left(\frac{\pi k}{p}\right).$$

The problem of evaluation of the sum

$$\sum_{k=1}^{p-1} \csc^{2m}\left(\frac{\pi k}{p}\right)$$

is still open. A partial answer is given in the work of the physician N. Gauthier and Paul S. Bruckman [5] at order  $Q$ , where  $Q = \frac{p-1}{2}$  if  $p$  odd and  $Q = \frac{p-2}{2}$  otherwise.

$$\sum_{k=1}^Q \csc^{2m}\left(\frac{\pi k}{p}\right) = \sum_{k=1}^m (2k-1)! \varphi_{k-1, m-1} J_{2k}(p), \quad m \geq 1$$

where

$$\varphi_{r, m} = \frac{s_{r, m}}{(2m+1)!}; \quad 0 \leq r \leq m; \quad m \geq 1,$$

$s_{r,m}$  is the sum of all the possible distinct products of the following numbers  $4.1^2, 4.2^2, \dots, 4.2^m$  and

$$J_{2k}(p) = \left(\frac{p}{\pi}\right)^{2k} \sum_{r=1}^Q \sum_n (r - np)^{-2k}.$$

But the value of the sum of maximal order of even integral powers of the secant  $\sum_{k=1}^{p-1} \sec^{2m} \left(\frac{\pi k}{p}\right)$  is expressed by [3, p. 1]

$$\sum_{k=1}^{p-1} \sec^{2m} \left(\frac{\pi k}{p}\right) = p \sum_{k=1}^{2m-1} (-1)^{m+k} \binom{m-1+kp}{2m-1} \sum_{j=k}^{2m-1} \binom{2m}{j+1}$$

The problem of finite sums with negative powers of cosecant or secant is completely resolved in [4] and [3, Theorem2.1 p.4], we copy here the result for cosecant

$$\sum_{k=0}^{p-1} \csc^{-2m} \left(\frac{\pi k}{p}\right) = \begin{cases} 2^{1-2m} p \left( \binom{2m-1}{m-1} + \sum_{n=1}^{\lfloor m/p \rfloor} (-1)^{pn} \binom{2m}{m-pn} \right), & \text{if } m \geq p, \\ 2^{1-2m} p \left( \binom{2m-1}{m-1} \right), & m < p. \blacksquare \end{cases}$$

Let  $m, n, l \in \mathbb{Z}$ , the last finite trigonometric sums can be extended to the general case

$$(1.2) \quad S_{n,m,l}(p, q, r) = \sum_{k=1}^{p-1} \cos^n \left(\frac{\pi kr}{p}\right) \csc^m \left(\frac{\pi kq}{p}\right) \csc^l \left(\frac{\pi k}{p}\right),$$

and then

$$S(p, q, r) = S_{1,1,1}(p, q, r).$$

Since  $p - k$  runs all the numbers from 1 to  $p - 1$  for  $k \in \{1, 2, 3, \dots, p - 1\}$  then the sum  $S_{n,m,l}(p, q, r)$  can be written in the following form

$$S_{n,m,l}(p, q, r) = \sum_{k=1}^{p-1} \cos^n \left(\pi r - \frac{\pi kr}{p}\right) \csc^m \left(\pi q - \frac{\pi kq}{p}\right) \csc^l \left(\pi - \frac{\pi k}{p}\right).$$

but  $\cos(\pi k - \theta) = (-1)^k \cos \theta$  and  $\sin(\pi k - \theta) = (-1)^{k+1} \sin \theta$  then

$$S_{n,m,l}(p, q, r) = \sum_{k=1}^{p-1} (-1)^{rn+(q+1)m} \cos^n \left(\frac{\pi kr}{p}\right) \csc^m \left(\frac{\pi kq}{p}\right) \csc^l \left(\frac{\pi k}{p}\right),$$

which means that

$$\left(1 - (-1)^{rn+(q+1)m}\right) S_{n,m,l}(p, q, r) = 0$$

and for  $rn + (q + 1)m$  odd number we get

$$S_{n,m,l}(p, q, r) = 0$$

On general, evaluating the reciprocity law of  $S_{n,m,l}(p, q, r)$  steal an open problem, for which we need more tools to resolve.

For more background about finite trigonometric sums in literature we refer to [1], [2] and [8]. The methods of computation based only on the residue theorem from complex analysis. The well-known classical Dedekind sum is

$$(1.3) \quad s(q, p) = \sum_{k=1}^{p-1} \left( \left( \frac{kq}{p} \right) \right) \left( \left( \frac{k}{p} \right) \right)$$

where  $q, p$  are coprime positive integers and

$$((x)) := \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \text{ is not an integer,} \\ 0, & \text{otherwise.} \end{cases}$$

In terms of fractional part function  $\{.\}$  (ie.  $\{x\} = x - [x]$ ) we can write

$$(1.4) \quad s(q, p) = \frac{1-p}{4} + \sum_{k=1}^{p-1} \left\{ \frac{kq}{p} \right\} \left\{ \frac{k}{p} \right\}$$

From the property that  $\{x\} + \{-x\} = 1$ , If  $0 < q < p$ , we get

$$(1.5) \quad s(p-q, p) = \frac{p-1}{2} - s(q, p).$$

$s(q, p)$  satisfies the following reciprocity law [7, p.4]

$$(1.6) \quad s(q, p) + s(p, q) = -\frac{1}{4} + \frac{1}{12} \left( \frac{q}{p} + \frac{p}{q} + \frac{1}{pq} \right)$$

The reciprocity law inducts an expression for  $s(p-q, p)$  on function of  $s(q, p)$ .

$$(1.7) \quad s(p, p-q) = s(q, p) + \frac{1}{4} + \frac{1}{12p(p-q)} (-6p^3 + 6p^2q + 2p^2 + q^2 - 2pq + 1)$$

This sum is interesting, since it can only be traduced to a finite trigonometric sum of product of cotangent functions [7, p.18]

$$(1.8) \quad s(q, p) = \frac{1}{4p} \sum_{k=1}^{p-1} \cot \left( \frac{\pi kq}{p} \right) \cot \left( \frac{\pi k}{p} \right)$$

The objective of this paper is to establish a reciprocity law for the trigonometric sums  $S(p, q, q \pm 1)$  and  $S(p, p-q, p-q \pm 1)$ , and get explicit evaluation of the sum

$$S(p, 1, 0) = \sum_{k=1}^{p-1} \csc^2 \left( \frac{\pi k}{p} \right).$$

**THEOREM 1.1.** *For  $p, q$  two coprime positive integers the following statement is true*

$$(1.9)$$

$$q[S(p, q, q+1) + S(p, q, q-1)] + p[S(q, p, p+1) + S(q, p, p-1)] = -2pq + \frac{2}{3}(p^2 + q^2 + 1) \blacksquare$$

And if  $q < p$  we get

(1.10)

$$\begin{aligned} p[S(q, p, p+1) + S(q, p, p-1)] - q[S(p, p-q, p-q+1) + S(p, p-q, p-q-1)] \\ = 2pq - 4p^2q + \frac{2}{3}(p^2 + q^2 + 1) \end{aligned}$$

COROLLARY 1.1. Let  $(p, q) = 1$  if  $q < p$  then

(1.11)

$$S(p, q, q+1) + S(p, q, q-1) + S(p, p-q, p-q+1) + S(p, p-q, p-q-1) = 4pq(p-1) \blacksquare$$

The following theorem gives another reciprocity law and compute the sum of the maximal order of the square of cosecant.

THEOREM 1.2. Let  $q \equiv 1 \pmod{p}$  then

$$(1.12) \quad S(q, p, p+1) + S(q, p, p-1) = \frac{2}{3} \left( q + \frac{p^2}{q} + \frac{1}{q} - p^2 - 2 \right)$$

and

$$(1.13) \quad \sum_{k=1}^{p-1} \csc^2 \left( \frac{\pi k}{p} \right) = \frac{p^2 - 1}{3}$$

## 2. proof of main results

LEMMA 2.1.

$$(2.1) \quad \cot q\beta \cot \beta = \frac{\cos(q+1)\beta + \cos(q-1)\beta}{2 \sin q\beta \sin \beta}$$

PROOF. From the well-known trigonometric formula

$$\cos \alpha \cos \beta = \frac{1}{2} (\cos(\alpha + \beta) + \cos(\alpha - \beta))$$

we get

$$\cot \alpha \cot \beta = \frac{\cos \alpha \cos \beta}{\sin \alpha \sin \beta} = \frac{\cos(\alpha + \beta) + \cos(\alpha - \beta)}{2 \sin \alpha \sin \beta}$$

and the result is deduced by taking  $\alpha = q\beta$ . □

LEMMA 2.2.

$$(2.2) \quad \sum_{k=1}^{p-1} \cot \left( \frac{\pi k q}{p} \right) \cot \left( \frac{\pi k}{p} \right) = \frac{1}{2} \{S(p, q, q+1) + S(p, q, q-1)\}$$

PROOF.

$$\sum_{k=1}^{p-1} \cot \left( \frac{\pi k q}{p} \right) \cot \left( \frac{\pi k}{p} \right) = \frac{1}{2} \sum_{k=1}^{p-1} \left( \cos \left( \frac{\pi k (q+1)}{p} \right) + \cos \left( \frac{\pi k (q-1)}{p} \right) \right) \csc \left( \frac{\pi k q}{p} \right) \csc \left( \frac{\pi k}{p} \right) \blacksquare$$

The decomposition of the right hand of the equality on two sums gives the result. □

Using the relation (2.2) we deduce that

$$(2.3) \quad s(q, p) = \frac{1}{8p} \{S(p, q, q+1) + S(p, q, q-1)\}$$

LEMMA 2.3. *Let  $q \equiv 1 \pmod{p}$  then*

$$(2.4) \quad s(q, p) = \frac{(p-1)(p-2)}{12p}$$

and

$$(2.5) \quad s(p, q) = \frac{1}{12} \left( \frac{q-2}{p} + \frac{p}{q} + \frac{1}{pq} - p \right)$$

PROOF. For  $q \equiv 1 \pmod{p}$ ,  $s(q, p) = s(1, p)$  is explicitly evaluated and

$$s(q, p) = \sum_{k=1}^{p-1} \left( \frac{k}{p} - \frac{1}{2} \right)^2$$

which becomes

$$s(q, p) = \frac{p-1}{4} - \frac{1}{p} \sum_{k=1}^{p-1} k + \frac{1}{p^2} \sum_{k=1}^{p-1} k^2$$

It's well-known that

$$\sum_{k=1}^{p-1} k = \frac{(p-1)p}{2}$$

and

$$\sum_{k=1}^{p-1} k^2 = \frac{p(p-1)(2p-1)}{6}.$$

and the result (2.4) follows.

The second result (2.5) is the consequence of the relation (2.4) and the reciprocity law (1.6).  $\square$

2.0.1. *Proof of Theorem 1.1.* From the relation (2.2) Lemma 2.2 and the expression (1.8) we get

$$s(q, p) = \frac{1}{8p} \{S(p, q, q+1) + S(p, q, q-1)\}$$

By symmetry we get the similar expression

$$s(p, q) = \frac{1}{8q} \{S(q, p, p+1) + S(q, p, p-1)\}.$$

The reciprocity formula (1.6) inducts

$$\begin{aligned} \frac{1}{8p} \{S(p, q, q+1) + S(p, q, q-1)\} + \frac{1}{8q} \{S(q, p, p+1) + S(q, p, p-1)\} = \\ -\frac{1}{4} + \frac{1}{12} \left( \frac{q}{p} + \frac{p}{q} + \frac{1}{pq} \right). \end{aligned}$$

Multiplying this equality by  $8pq$  we get the result (1.9) Theorem1.1.

Only we have

$$s(p-q, p) = \frac{1}{8p} \{S(p, p-q, p-q+1) + S(p, p-q, p-q-1)\}$$

and from the relation (1.5) we get

$$S(q, p) = \frac{p-1}{2} - \frac{1}{8p} \{S(p, p-q, p-q+1) + S(p, p-q, p-q-1)\},$$

and then

$$\frac{1}{8p} \{S(p, q, q+1) + S(p, q, q-1)\} = \frac{p-1}{2} - \frac{1}{8p} \{S(p, p-q, p-q+1) + S(p, p-q, p-q-1)\}, \blacksquare$$

thus

$$q \{S(p, q, q+1) + S(p, q, q-1)\} = 4pq(p-1) - q \{S(p, p-q, p-q+1) + S(p, p-q, p-q-1)\}, \blacksquare$$

and then

$$\begin{aligned} p \{S(q, p, p+1) + S(q, p, p-1)\} - q \{S(p, p-q, p-q+1) + S(p, p-q, p-q-1)\} \\ = 2pq - 4p^2q + \frac{2}{3}(p^2 + q^2 + 1) \blacksquare \end{aligned}$$

2.0.2. *Proof of Corollary1.1.* The reciprocity theorem (1.11) of Corollary1.1 is result of the relation (1.9) minus the relation (1.10)

2.0.3. *Proof of Theorem1.2.* From the reciprocity theorem

$$s(p, q) = -\frac{1}{4} + \frac{1}{12} \left( \frac{q}{p} + \frac{p}{q} + \frac{1}{pq} \right) - s(q, p)$$

using the relation (2.4) we deduce that

$$s(p, q) = -\frac{1}{4} + \frac{1}{12} \left( \frac{q}{p} + \frac{p}{q} + \frac{1}{pq} \right) - \frac{(p-1)(p-2)}{12p}$$

and

$$s(p, q) = -\frac{1}{4} + \frac{1}{12} \left( \frac{q}{p} + \frac{p}{q} + \frac{1}{pq} - p + 3 - \frac{2}{p} \right).$$

Thus

$$s(p, q) = \frac{1}{12} \left( \frac{q-2}{p} + \frac{p}{q} + \frac{1}{pq} - p \right).$$

From the relation (2.3) we get

$$\frac{1}{8p} \{S(q, p, p+1) + S(q, p, p-1)\} = \frac{1}{12} \left( \frac{q-2}{p} + \frac{p}{q} + \frac{1}{pq} - p \right)$$

and the result (1.12) follows.

Taking  $q = 1$ , and combining the relation (2.3) and equality (2.4) Lemma2.3 we get

$$S(p, 1, 2) + S(p, 1, 0) = \frac{2}{3}(p-1)(p-2),$$

then

$$\sum_{k=1}^{p-1} \frac{\cos\left(\frac{2\pi k}{p}\right) + 1}{\sin^2\left(\frac{\pi k}{p}\right)} = \frac{2p^2 - 6p + 4}{3}.$$

But

$$\cos\left(\frac{2\pi k}{p}\right) = 1 - 2\sin^2\left(\frac{\pi k}{p}\right)$$

then

$$\sum_{k=1}^{p-1} \frac{2 - 2\sin^2\left(\frac{\pi k}{p}\right)}{\sin^2\left(\frac{\pi k}{p}\right)} = \frac{2p^2 - 6p + 4}{3}.$$

and

$$2 \sum_{k=1}^{p-1} \frac{1}{\sin^2\left(\frac{\pi k}{p}\right)} = \frac{2p^2 - 6p + 4}{3} + 2(p-1).$$

Thus

$$\sum_{k=1}^{p-1} \frac{2}{\sin^2\left(\frac{\pi k}{p}\right)} = \frac{p^2 - 3p + 2}{3} + p - 1$$

Finally

$$\sum_{k=1}^{p-1} \frac{1}{\sin^2\left(\frac{\pi k}{p}\right)} = \frac{p^2 - 1}{3}.$$

### 3. Additional result

They are several different sums extracted from the classical Dedekind sum which satisfy a reciprocity law for a particular case. An example of a generalized cosecant sum (1.2) is given in the following proposition

PROPOSITION 3.1.

$$(3.1) \quad 2S_{3,1,1}(p, 2, 1) - S_{1,1,1}(p, 2, 1) = \frac{1}{6}p^2 - p + \frac{5}{2}$$

PROOF. It's trivial that  $s(p, 2) = 0$  and from the reciprocity law (1.6) we get

$$s(2, p) = -\frac{1}{4} + \frac{1}{12} \left( \frac{p}{2} + \frac{5}{2p} \right).$$

Form the relation (2.3) we get

$$s(2, p) = \frac{1}{8p} (S(p, 2, 3) + S(p, 2, 1))$$

and

$$S(p, 2, 3) + S(p, 2, 1) = \sum_{k=1}^{p-1} \left( \cos\left(\frac{3\pi k}{p}\right) + \cos\left(\frac{\pi k}{p}\right) \right) \csc\left(\frac{2\pi k}{p}\right) \csc\left(\frac{\pi k}{p}\right).$$



Using the trigonometric relation

$$\begin{aligned} \cos 3\theta &= 4 \cos^3 \theta - 3 \cos \theta \\ S(p, 2, 3) + S(p, 2, 1) &= 4 \sum_{k=1}^{p-1} \cos^3 \left( \frac{\pi k}{p} \right) \csc \left( \frac{2\pi k}{p} \right) \csc \left( \frac{\pi k}{p} \right) \\ &- 2 \sum_{k=1}^{p-1} \cos \left( \frac{\pi k}{p} \right) \csc \left( \frac{2\pi k}{p} \right) \csc \left( \frac{\pi k}{p} \right). \end{aligned}$$

We deduce that

$$4S_{3,1,1}(p, 2, 1) - 2S_{1,1,1}(p, 1, 1) = 8ps(2, p).$$

Replacing  $s(2, p)$  by it's value we get the result (3.1) Proposition 3.1.  $\square$

Exploitation of  $S(2, p) = 0$  conduct to proof of the following trigonometric formula

$$\cos \left( \frac{\pi(p+1)}{2} \right) + \cos \left( \frac{\pi(p-1)}{2} \right) = 0$$

The argument is that

$$\frac{1}{16} \{S(2, p, p+1) + S(2, p, p-1)\} = s(2, p) = 0$$

and then

$$\left\{ \cos \left( \frac{\pi(p+1)}{2} \right) + \cos \left( \frac{\pi(p-1)}{2} \right) \right\} \csc \left( \frac{\pi}{2} \right) = 0$$

### References

- [1] B. C. Berndt, B. P. Yeap and B. Pin *Explicit evaluations and reciprocity theorems for finite trigonometric sums*, Adv. Appl. Math. **29** (2002) no.3 358–385
- [2] C. Datta and P. Agrawal *A Few Finite Trigonometric Sums*, Mathematics **5** (2017) no.1, Article ID 13,11 p.
- [3] da Fonseca, Carlos M., M. Lawrance, V. Kowalenko, *Basic trigonometric power sums with applications* Ramanujan J.42 (2017) no.2, 401–428
- [4] C. M. da Fonseca and V. Kowalenko, *On a finite sum with power of cosines*, Appl. Anal. Discrete Math. ,**7**,(2013),354–377
- [5] N. Gauthier and P. S. Bruckman, *Sums of the even integral powers of the cosecant and secant*, Fibonacci Q. **44** (2006) no.3 264–273
- [6] V. Kowalenko, *Generalized cosecant numbers and the Hurwitz zeta function* arXiv: 1702.04090v1 [Math. NT] 14 Feb. 2017
- [7] H. Rademacher and E. Grosswald, *Dedekind Sums*. Carus Mathematical Monograph, **16**, Mathematical Association of America, Washington D.C. 1972.
- [8] K. S. Williams and N. Y. Zhang *Evaluation of two trigonometric sums* Mathematica Slovaca, **44**, (1994) no.5, 575–583

MOULOUD GOUBI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UMMTO RP. 15000, TIZI-OUZOU, ALGERIA, LABORATOIRE D'ALGÈBRE ET THÉORIE DES NOMBRES, USTHB ALGER  
E-mail address: mouloud.goubi@ummtto.dz