

NON-UNIFORM CONTINUOUS DEPENDENCE AND CONTINUOUS DEPENDENCE FOR THE NON-RESISTIVE MHD EQUATIONS

JINLU LI, ZHAOYANG YIN, AND WEIPENG ZHU

ABSTRACT. In this paper, we establish the continuous dependence for the non-resistive MHD equations in Sobolev spaces. Our obtained result fills considerably the recent result [6] for the continuous dependence on initial data. We also show that this result is optimal. More precisely, we proved that the data-to-solution map is not uniform continuous dependence in Sobolev spaces.

1. INTRODUCTION AND MAIN RESULT

In the paper, we consider the following Cauchy problem of the non-resistive MHD equations:

$$\begin{cases} \partial_t u + u \cdot \nabla u - \Delta u + \nabla P = b \cdot \nabla b, \\ \partial_t b + u \cdot \nabla b = b \cdot \nabla u, \\ \operatorname{div} u = \operatorname{div} b = 0, \quad (u, b)|_{t=0} = (u_0, b_0), \end{cases} \quad (1.1)$$

where the unknowns are the vector fields $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$, $b : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ and the scalar function P . Here, u and b are the velocity and magnetic, respectively, while P denotes the pressure.

In [12], Jiu and Niu were first concerned with the well-posedness of system (1.1) and established the local existence of solutions in 2D for initial data in H^s , $s \geq 3$. Fefferman et al. obtained local-in-time existence of strong solutions to (1.1) in \mathbb{R}^d , $d = 2, 3$ with the initial data $(u_0, b_0) \in H^s(\mathbb{R}^d) \times H^s(\mathbb{R}^d)$, $s > \frac{d}{2}$ in [6] and $(u_0, b_0) \in H^{s-1+\varepsilon}(\mathbb{R}^d) \times H^s(\mathbb{R}^d)$, $s > \frac{d}{2}$, $0 < \varepsilon < 1$ in [7]. Chemin et al. in [4] presented the local existence of solutions to (1.1) in \mathbb{R}^d , $d = 2, 3$ with the initial data $(u_0, b_0) \in B_{2,1}^{\frac{d}{2}-1}(\mathbb{R}^d) \times B_{2,1}^{\frac{d}{2}}(\mathbb{R}^d)$ and also proved the corresponding solution is unique in 3D case. Wan in [18] resolved the uniqueness of the solution in the 2D case by using mixed space-time Besov spaces. Recently, Li, Tan and Yin in [13] obtain the local existence and uniqueness in the homogeneous Besov spaces $(u_0, b_0) \in \dot{B}_{p,1}^{\frac{d}{p}-1}(\mathbb{R}^d) \times \dot{B}_{p,1}^{\frac{d}{p}}(\mathbb{R}^d)$ where $p \in [1, 2d]$.

For the global existence of small solutions to the system (1.1), there are many interesting results when the background magnetic field is \vec{e}_1 , i.e. in the case that the initial data (u_0, b_0) is sufficiently close to the equilibrium state $(\vec{0}, \vec{e}_1)$, we refer the reader to see [1, 11, 15, 16, 17, 19, 21]. In [22], Zhang obtained the global well-posedness result with the large initial data under the assumption that the background magnetic field is large enough. Subsequently, Zhai and Zhang [20] considered

2010 *Mathematics Subject Classification.* 76W05.

Key words and phrases. The non-resistive MHD equations, non-uniform continuous dependence, continuous dependence.

the global existence and uniqueness of solution to system (1.1) with non-equilibrium background magnetic field.

However, the continuous dependence property of solutions for the Cauchy problem of the non-resistive MHD equations with initial data $(u_0, b_0) \in H^s(\mathbb{R}^d) \times H^s(\mathbb{R}^d)$, $s > \frac{d}{2}$ has not been studied yet. Our first goal in the paper is to prove the continuous dependence property of solutions to the non-resistive MHD equations in such Sobolev spaces. Together with recent local existence and uniqueness results in [6, 7], we establish local well-posedness of the Cauchy problem of the non-resistive MHD system in the sense of Hadamard.

In order to state our main result, we first recall the following local-in-time existence of strong solutions to (1.1) in [6]:

Lemma 1.1. [6] *For $s > \frac{d}{2}$, $d = 2, 3$, and initial data $u_0, b_0 \in H^s(\mathbb{R}^d)$ with $\operatorname{div} u_0 = \operatorname{div} b_0 = 0$, there exists a time $T = T(s, \|u_0\|_{H^s}, \|b_0\|_{H^s}) > 0$ such that the system (1.1) have a unique solution (u, b) , with $u, b \in \mathcal{C}([0, T]; H^s)$ and $\nabla u \in L_T^2(H^s)$. Moreover, for all $t \in [0, T]$, there holds*

$$\begin{aligned} \|u(t)\|_{H^s}^2 + \|b(t)\|_{H^s}^2 &\leq \frac{\|u_0\|_{H^s}^2 + \|b_0\|_{H^s}^2}{1 - C_s T (\|u_0\|_{H^s}^2 + \|b_0\|_{H^s}^2)}, \\ \int_0^t \|\nabla u(\tau)\|_{H^s}^2 d\tau &\leq C_s \frac{\|u_0\|_{H^s}^2 + \|b_0\|_{H^s}^2}{1 - C_s T (\|u_0\|_{H^s}^2 + \|b_0\|_{H^s}^2)}. \end{aligned}$$

To solve the continuous dependence property of the system (1.1), the main difficulty is that the system is only partially parabolic, owing to the magnetic equation which is of hyperbolic type. That is if (u^1, b^1) and (u^2, b^2) are two solutions of the system (1.1), then we set $\delta b = b^1 - b^2$ satisfies

$$\partial_t \delta b + u^1 \cdot \nabla \delta b = \delta b \cdot \nabla u^1 + b^2 \cdot \nabla \delta u - \delta u \cdot \nabla b^2.$$

Due to the term $\nabla u^2 \in \mathcal{C}([0, T]; H^{s-1})$, this precludes any attempt to tackle with the estimate $\|\delta b\|_{H^s}$. Inspired by Bona-Smith [3] and Guo-Li-Yin [8], we can take (u^2, b^2) as the solution with initial data $S_j(u_0, b_0)$ to overcome this problem. Therefore, we first will prove the continuous dependence property of solutions to the system (1.1).

Our first main theorem can be stated as follows:

Theorem 1.2. *Let $d = 2, 3$ and $s > \frac{d}{2}$. Denote $\bar{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$. Assume that $(u^n, b^n)_{n \in \bar{\mathbb{N}}}$ be the solution to the system (1.1) with initial data $(u_0^n, b_0^n)_{n \in \bar{\mathbb{N}}}$. If (u_0^n, b_0^n) tends to (u_0^∞, b_0^∞) in H^s , then there exists a positive time T independent of n such that (u^n, b^n) tends to (u^∞, b^∞) in $\mathcal{C}([0, T]; H^s) \times \mathcal{C}([0, T]; H^s)$.*

When $b = 0$, System (1.1) becomes Navier-Stokes equations. It is well known that the solution map of Navier-Stokes equation is local lipshitz. That is, if $\|u\|_{H^s}, \|v\|_{H^s} \leq R$, we can obtain that

$$\|u(t) - v(t)\|_{H^s} \leq C \|u_0 - v_0\|_{H^s}, \quad t \in [0, T],$$

for some positive time T depending only on s and R .

However, the magnetic equation of the non-resistive MHD equations have the construction of transport diffusion equation. Therefore, motivated by Himonas-Misiolek [10], we can conclude that Cauchy problem (1.3) is not uniformly continuous. The main difficulty is to bound $u^{\omega, n}(t, x) - u^{\omega, n}(0, x)$ less than n^{-1} in Sobolev H^{s-1} norm which is the main part of $\partial_t b^{h, \omega, n} + u^{\omega, n} \cdot \nabla b^{h, \omega, n}$. To overcome it, we split $u^{\omega, n}(t, x) - u^{\omega, n}(0, x)$ into two parts: $u^{\omega, n}(t, x) - e^{t\Delta} u^{\omega, n}(0, x)$ and

$e^{t\Delta}u^{\omega,n}(0, x) - u^{\omega,n}(0, x)$. We apply the Duhamel's principle to tackle with the first part. The second part can be seen as $\Delta u^{\omega,n}(0, x)$, and its estimate adds the factor $n^{-2\delta}$ compared with $u^{\omega,n}(0, x)$.

Our second main theorem can be stated as follows:

Theorem 1.3. *Let $d = 2, 3$ and $s > \frac{d}{2}$. The data-to-solution map for the system (1.1) is not uniformly continuous from any bounded subset in H^s into $C([0, T]; H^s)$. That is, there exists two sequences of solutions (u^n, b^n) and (v^n, c^n) such that*

$$\begin{aligned} & \|u_0^n\|_{H^s} + \|b_0^n\|_{H^s} + \|v_0^n\|_{H^s} + \|c_0^n\|_{H^s} \lesssim 1, \\ & \lim_{n \rightarrow \infty} (\|u_0^n - v_0^n\|, \|b_0^n - c_0^n\|_{H^s}) = 0, \\ & \liminf_{n \rightarrow \infty} (\|u^n(t) - v^n(t)\|_{H^s} + \|b^n(t) - c^n(t)\|_{H^s}) \gtrsim |\sin t|, \quad t \in [0, T]. \end{aligned}$$

The paper is organized as follows. In Section 2, we recall the Littlewood-Paley theory and give some properties of Besov spaces. In Section 3, we introduce some lemmas to overcome the difficulty of the problem. In Section 4, we prove the continuous dependence property of the system (1.1). In Section 5, we show that the solution of the system (1.1) is non-uniform continuous dependence.

Notations. Given a Banach space X , we denote its norm by $\|\cdot\|_X$. Since all spaces of functions are over \mathbb{R}^d , for simplicity, we drop \mathbb{R}^d in our notations of function spaces if there is no ambiguity. The symbol $A \lesssim B$ denotes that there exists a constant $c > 0$ independent of A and B , such that $A \leq cB$. The symbol $A \simeq B$ represents $A \lesssim B$ and $B \lesssim A$.

2. LITTLEWOOD-PALEY ANALYSIS

In this section, we will recall some facts about the Littlewood-Paley decomposition, the nonhomogeneous Besov spaces and their some useful properties. For more details, the readers can refer to [2].

There exists a couple of smooth functions (χ, φ) valued in $[0, 1]$, such that χ is supported in the ball $\mathcal{B} \triangleq \{\xi \in \mathbb{R}^d : |\xi| \leq \frac{4}{3}\}$, and φ is supported in the ring $\mathcal{C} \triangleq \{\xi \in \mathbb{R}^d : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$. Moreover,

$$\forall \xi \in \mathbb{R}^d, \quad \chi(\xi) + \sum_{j \geq 0} \varphi(2^{-j}\xi) = 1,$$

$$\forall 0 \neq \xi \in \mathbb{R}^d, \quad \sum_{j \in \mathbb{Z}} \varphi(2^{-j}\xi) = 1,$$

$$|j - j'| \geq 2 \Rightarrow \text{Supp } \varphi(2^{-j}\cdot) \cap \text{Supp } \varphi(2^{-j'}\cdot) = \emptyset,$$

$$j \geq 1 \Rightarrow \text{Supp } \chi(\cdot) \cap \text{Supp } \varphi(2^{-j}\cdot) = \emptyset,$$

Then, we can define the nonhomogeneous dyadic blocks Δ_j and nonhomogeneous low frequency cut-off operator S_j as follows:

$$\Delta_j u = 0, \quad \text{if } j \leq -2, \quad \Delta_{-1} u = \chi(D)u = \mathcal{F}^{-1}(\chi \mathcal{F}u),$$

$$\Delta_j u = \varphi(2^{-j}D)u = \mathcal{F}^{-1}(\varphi(2^{-j}\cdot)\mathcal{F}u), \quad \text{if } j \geq 0,$$

$$S_j u = \sum_{j'=-\infty}^{j-1} \Delta_{j'} u.$$

Definition 2.1. ([2]) Let $s \in \mathbb{R}$ and $1 \leq p, r \leq \infty$. The nonhomogeneous Besov space $B_{p,r}^s$ consists of all tempered distribution u such that

$$\|u\|_{B_{p,r}^s} \triangleq \left\| (2^{js} \|\Delta_j u\|_{L^p})_{j \in \mathbb{Z}} \right\|_{\ell^r(\mathbb{Z})} < \infty.$$

Remark 2.2. ([2]) When $p = r = 2$, we have $B_{2,2}^s(\mathbb{R}^d) = H^s(\mathbb{R}^d)$. Here, $H^s(\mathbb{R}^d)$ is the standard Sobolev space with the norm

$$\|u\|_{H^s}^2 := \int_{\mathbb{R}^d} (1 + |\xi^2|)^s |\hat{u}(\xi)|^2 d\xi.$$

If $s > \frac{d}{p}$, we also have $\|u\|_{L^\infty} \lesssim \|u\|_{B_{p,r}^s}$.

Then, we have the following product laws.

Lemma 2.3. ([2]) Let $s > \frac{d}{2}$, $d \geq 2$. Then there exists a constant $C = C(d, s)$ such that

$$\begin{aligned} \|uv\|_{H^{s-1}} &\leq C(\|u\|_{L^\infty} \|v\|_{H^{s-1}} + \|v\|_{L^\infty} \|u\|_{H^{s-1}}), \\ \|uv\|_{H^s} &\leq C\|u\|_{H^s} \|v\|_{H^s}, \quad \|uv\|_{H^{s-1}} \leq C\|u\|_{H^{s-1}} \|v\|_{H^s}. \end{aligned}$$

Lemma 2.4. ([2, 14]) Let $\sigma \in \mathbb{R}$ and $1 \leq p, r \leq \infty$. Let v be a vector field over \mathbb{R}^d . Assume that $\sigma > -d \min\{1 - \frac{1}{p}, \frac{1}{p}\}$. Define $R_j = [v \cdot \nabla, \Delta_j]f$. There a constant $C = C(p, \sigma, d)$ such that

$$\|(2^{j\sigma} \|R_j\|_{L^p})_{j \geq -1}\|_{\ell^r} \leq \begin{cases} C \|\nabla v\|_{B_{p,\infty}^{\frac{d}{p}} \cap L^\infty} \|f\|_{B_{p,r}^\sigma}, & \text{if } \sigma < 1 + \frac{d}{p}, \\ C \|\nabla v\|_{B_{p,\infty}^{\frac{d}{p}+1}} \|f\|_{B_{p,r}^\sigma}, & \text{if } \sigma = 1 + \frac{d}{p}, r > 1, \\ C \|\nabla v\|_{B_{p,r}^{\sigma-1}} \|f\|_{B_{p,r}^\sigma}, & \text{otherwise.} \end{cases}$$

Lemma 2.5. ([9, 10]) Let $\phi \in S(\mathbb{R})$, $\delta > 0$ and $s \geq 0$. Then, we have for all $\alpha \in \mathbb{R}$

$$\begin{aligned} \lim_{n \rightarrow \infty} n^{-\frac{\delta}{2}} \|\phi(\frac{x}{n^\delta})\|_{H^s(\mathbb{R})} &= \|\phi\|_{L^2(\mathbb{R})}, \\ \lim_{n \rightarrow \infty} n^{-\frac{1}{2}\delta-s} \|\phi(\frac{x}{n^\delta}) \cos(nx - \alpha)\|_{H^s(\mathbb{R})} &= \frac{1}{\sqrt{2}} \|\phi\|_{L^2(\mathbb{R})}, \\ \lim_{n \rightarrow \infty} n^{-\frac{1}{2}\delta-s} \|\phi(\frac{x}{n^\delta}) \sin(nx - \alpha)\|_{H^s(\mathbb{R})} &= \frac{1}{\sqrt{2}} \|\phi\|_{L^2(\mathbb{R})}. \end{aligned}$$

3. CONTINUOUS DEPENDENCE

In this section, we will show that the solution of is continuous dependence with initial data. First, we establish some Sobolev norm estimates for smooth solutions of system (1.1), which is the key component in the proof of Theorem 1.2. In order to simplify the notation, we define $(u, b) \in X_s(T)$ if

$$(u, b) \in \mathcal{C}([0, T]; B_{2,2}^\sigma), \quad \nabla u \in L^2([0, T]; B_{2,2}^\sigma), \quad \sigma \in \mathbb{R}.$$

Lemma 3.1. Let $d = 2, 3$ and $s > \frac{d}{2}$. Suppose that $(u^1, b^1) \in X_s(T)$ and $(u^2, b^2) \in X_{s+1}(T)$ are two solutions of (1.1) with initial data (u_0^1, b_0^1) and (u_0^2, b_0^2) respectively.

Denote $\delta u = u^1 - u^2$ and $\delta b = b^1 - b^2$. Then, we have for all $t \in [0, T]$,

$$\begin{aligned} & \|\delta u(t)\|_{B_{2,2}^{s-1}}^2 + \|\delta b(t)\|_{B_{2,2}^{s-1}}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{B_{2,2}^{s-1}}^2 d\tau \\ & \leq (\|\delta u_0\|_{B_{2,2}^{s-1}}^2 + \|\delta b_0\|_{B_{2,2}^{s-1}}^2) e^{A(t)}, \end{aligned}$$

and

$$\begin{aligned} & \|\delta u(t)\|_{B_{2,2}^s}^2 + \|\delta b(t)\|_{B_{2,2}^s}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{B_{2,2}^s}^2 d\tau \\ & \leq \left(\|\delta u_0\|_{B_{2,2}^s}^2 + \|\delta b_0\|_{B_{2,2}^s}^2 + C \int_0^t \|b^2\|_{H^{s+1}}^2 (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\nabla \delta u\|_{B_{2,2}^{s-1}}^2) d\tau \right) e^{A(t)}, \end{aligned}$$

with

$$A(t) = C \int_0^t (1 + \|u^1\|_{H^{s+1}}^2 + \|u^2\|_{H^{s+1}}^2 + \|b^1\|_{H^s}^2 + \|b^2\|_{H^s}^2) d\tau.$$

Proof. It is easy to show that

$$\begin{cases} \partial_t \delta u + u^1 \cdot \nabla \delta u + \delta u \cdot \nabla u^2 - \Delta \delta u + \nabla P = b^1 \cdot \nabla \delta b + \delta b \cdot \nabla b^2, \\ \partial_t \delta b + u^1 \cdot \nabla \delta b + \delta u \cdot \nabla b^2 = b^1 \cdot \nabla \delta u + \delta b \cdot \nabla u^2, \\ \operatorname{div} \delta u = \operatorname{div} \delta b = 0, \quad (\delta u, \delta b)|_{t=0} = (\delta u_0, \delta b_0). \end{cases} \quad (3.1)$$

Now, we apply Δ_j to (3.1), and take the inner product with $(\Delta_j \delta u, \Delta_j \delta b)$ and integrate by parts to have

$$\frac{1}{2} \frac{d}{dt} (\|\Delta_j \delta u\|_{L^2}^2 + \|\Delta_j \delta b\|_{L^2}^2) + \|\Delta_j \nabla \delta u\|_{L^2}^2 \leq K_1 + K_2 + K_3 + K_4 + K_5, \quad (3.2)$$

where

$$\begin{aligned} K_1 &= - \int_{\mathbb{R}^d} [\Delta_j, u^1 \cdot \nabla] \delta u \cdot \Delta_j \delta u \, dx, & K_2 &= - \int_{\mathbb{R}^d} [\Delta_j, u^1 \cdot \nabla] \delta b \cdot \Delta_j \delta b \, dx, \\ K_3 &= \int_{\mathbb{R}^d} \Delta_j (b^1 \cdot \nabla \delta b + \delta b \cdot \nabla b^2 - \delta u \cdot \nabla u^2) \Delta_j \delta u \, dx, \\ K_4 &= \int_{\mathbb{R}^d} \Delta_j (b^1 \cdot \nabla \delta u + \delta b \cdot \nabla u^2) \Delta_j \delta b \, dx, \\ K_5 &= - \int_{\mathbb{R}^d} \Delta_j (\delta u \cdot \nabla b^2) \Delta_j \delta b \, dx. \end{aligned}$$

First, we estimate the above terms in the $B_{2,2}^{s-1}$ norm. According to Lemmas 2.3-2.4, it is easy to estimate

$$\begin{aligned} |K_1| &\lesssim \|[\Delta_j, u^1 \cdot \nabla] \delta u\|_{L^2} \|\Delta_j \delta u\|_{L^2} \lesssim 2^{-2j(s-1)} c_j^2 \|\nabla u^1\|_{B_{2,2}^s} \|\delta u\|_{B_{2,2}^{s-1}}^2, \\ |K_2| &\lesssim \|[\Delta_j, u^1 \cdot \nabla] \delta b\|_{L^2} \|\Delta_j \delta b\|_{L^2} \lesssim 2^{-2j(s-1)} c_j^2 \|\nabla u^1\|_{B_{2,2}^s} \|\delta b\|_{B_{2,2}^{s-1}}^2, \\ |K_3| &\lesssim 2^{-2j(s-1)} c_j^2 (\|b^1, b^2\|_{B_{2,2}^s} \|\nabla \delta u\|_{B_{2,2}^{s-1}} \|\delta b\|_{B_{2,2}^{s-1}} \\ &\quad + \|u^2\|_{H^s} \|\delta u\|_{B_{2,2}^{s-1}} \|\nabla \delta u\|_{B_{2,2}^{s-1}}), \\ |K_4| &\lesssim 2^{-2j(s-1)} c_j^2 (\|b^1\|_{B_{2,2}^s} \|\nabla \delta u\|_{B_{2,2}^{s-1}} \|\delta b\|_{B_{2,2}^{s-1}} + \|\nabla u^2\|_{H^s} \|\delta b\|_{B_{2,2}^{s-1}}^2), \\ |K_5| &\lesssim 2^{-2j(s-1)} c_j^2 \|b^2\|_{B_{2,2}^s} (\|\delta u\|_{B_{2,2}^{s-1}} + \|\nabla \delta u\|_{B_{2,2}^{s-1}}) \|\delta b\|_{B_{2,2}^{s-1}}. \end{aligned} \quad (3.3)$$

Multiplying the inequality above by $2^{2j(s-1)}$ and summing over $j \geq -1$, we conclude from (3.2)-(3.3) that

$$\begin{aligned} & \frac{d}{dt} (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\delta b\|_{B_{2,2}^{s-1}}^2) + \|\nabla \delta u(t)\|_{B_{2,2}^{s-1}}^2 \\ & \leq A'(t) (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\delta b\|_{B_{2,2}^{s-1}}^2) + \frac{1}{2} \|\nabla \delta u\|_{B_{2,2}^{s-1}}^2, \end{aligned}$$

which leads to

$$\begin{aligned} & \|\delta u(t)\|_{B_{2,2}^{s-1}}^2 + \|\delta b(t)\|_{B_{2,2}^{s-1}}^2 + \int_0^t \|\nabla \delta u\|_{B_{2,2}^{s-1}}^2 d\tau \\ & \leq \|\delta u_0\|_{B_{2,2}^{s-1}}^2 + \|\delta b_0\|_{B_{2,2}^{s-1}}^2 + \int_0^t A'(\tau) (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\delta b\|_{B_{2,2}^{s-1}}^2) d\tau. \end{aligned}$$

This obtain the first result of this lemma. Now, we estimate the above terms in the $B_{2,2}^s$ norm. For the terms of $K_i, i = 1, 2, 3, 4, 5$, we also infer from Lemmas 2.3-2.4 that

$$\begin{aligned} |K_1| & \lesssim \|[\Delta_j, u^1 \cdot \nabla] \delta u\|_{L^2} \|\Delta_j \delta u\|_{L^2} \lesssim 2^{-2js} c_j^2 \|\nabla u^1\|_{B_{2,2}^s} \|\delta u\|_{B_{2,2}^s}, \\ |K_2| & \lesssim \|[\Delta_j, u^1 \cdot \nabla] \delta b\|_{L^2} \|\Delta_j \delta b\|_{L^2} \lesssim 2^{-2js} c_j^2 \|\nabla u^1\|_{B_{2,2}^s} \|\delta b\|_{B_{2,2}^s}, \\ |K_3| & \lesssim 2^{-2js} c_j^2 (\|b^1, b^2\|_{B_{2,2}^s} \|\nabla \delta u\|_{B_{2,2}^s} \|\delta b\|_{B_{2,2}^s} + \|u^2\|_{H^s} \|\delta u\|_{B_{2,2}^s} \|\nabla \delta u\|_{B_{2,2}^s}), \\ |K_4| & \lesssim 2^{-2js} c_j^2 (\|b^1\|_{B_{2,2}^s} \|\nabla \delta u\|_{B_{2,2}^s} \|\delta b\|_{B_{2,2}^s} + \|\nabla u^2\|_{H^s} \|\delta b\|_{B_{2,2}^s}^2), \\ |K_5| & \lesssim 2^{-2js} c_j^2 (\|\delta u, \nabla \delta u\|_{B_{2,2}^{s-1}} \|b^2\|_{H^{s+1}}) \|\delta b\|_{B_{2,2}^s}. \end{aligned} \quad (3.4)$$

Multiplying the inequality above by 2^{2js} and summing over $j \geq -1$, we deduce from (3.2), (3.4) that

$$\begin{aligned} & \frac{d}{dt} (\|\delta u\|_{B_{2,2}^s}^2 + \|\delta b\|_{B_{2,2}^s}^2) + \|\nabla \delta u(t)\|_{B_{2,2}^s}^2 \\ & \leq A'(t) (\|\delta u\|_{B_{2,2}^s}^2 + \|\delta b\|_{B_{2,2}^s}^2) + \frac{1}{2} \|\nabla \delta u\|_{B_{2,2}^s}^2 \\ & \quad + \|b^2\|_{H^{s+1}}^2 (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\nabla \delta u\|_{B_{2,2}^{s-1}}^2), \end{aligned}$$

which implies

$$\begin{aligned} & \|\delta u(t)\|_{B_{2,2}^s}^2 + \|\delta b(t)\|_{B_{2,2}^s}^2 + \int_0^t \|\nabla \delta u(\tau)\|_{B_{2,2}^s}^2 d\tau \\ & \leq \|\delta u_0\|_{B_{2,2}^s}^2 + \|\delta b_0\|_{B_{2,2}^s}^2 + \int_0^t A'(\tau) (\|\delta u\|_{B_{2,2}^s}^2 + \|\delta b\|_{B_{2,2}^s}^2) d\tau \\ & \quad + C \int_0^t \|b^2\|_{H^{s+1}}^2 (\|\delta u\|_{B_{2,2}^{s-1}}^2 + \|\nabla \delta u\|_{B_{2,2}^{s-1}}^2) d\tau. \end{aligned}$$

This completes the proof of this lemma. \square

Then, in virtue of Lemma 3.1, we will show that the solution to the system (1.1) depend continuously on the initial data.

Proof of Theorem 1.2. First, according to Lemma 1.1, there exist a positive $T_n > 0$ such that (1.1) have a solution $(u^n, b^n) \in X_s(T_n)$. Indeed, by Lemma 1.1, we have

$$T_n \geq \frac{1}{2C_s (\|u_0^n\|_{H^s}^2 + \|b_0^n\|_{H^s}^2)} \geq \frac{1}{2C_s R} := T,$$

with $R = \sup_{n \geq 0} (\|u_0^n\|_{H^s}^2 + \|b_0^n\|_{H^s}^2)$. Moreover, we also obtain from Lemma 1.1 that for all $t \in [0, T]$,

$$\|u^n(t)\|_{B_{2,2}^s}^2 + \|b^n(t)\|_{B_{2,2}^s}^2 + \int_0^t \|\nabla u^n\|_{B_{2,2}^s}^2 d\tau \leq C(\|u_0^n\|_{B_{2,2}^s}^2 + \|b_0^n\|_{B_{2,2}^s}^2).$$

Since $s + 1 > 1 + \frac{d}{2}$, we use the similar method as in Lemma 3.1 to have for all $t \in [0, T]$,

$$\begin{aligned} & \|u^n(t)\|_{B_{2,2}^{s+1}}^2 + \|b^n(t)\|_{B_{2,2}^{s+1}}^2 + \int_0^t \|\nabla u^n\|_{B_{2,2}^{s+1}}^2 d\tau \\ & \leq (\|u_0^n\|_{B_{2,2}^{s+1}}^2 + \|b_0^n\|_{B_{2,2}^{s+1}}^2) e^{C \int_0^t (1 + \|u^n\|_{B_{2,2}^{s+1}}^2 + \|b^n\|_{B_{2,2}^{s+1}}^2) d\tau} \\ & \leq C(\|u_0^n\|_{B_{2,2}^{s+1}}^2 + \|b_0^n\|_{B_{2,2}^{s+1}}^2). \end{aligned} \quad (3.5)$$

Let $(u_j^n, b_j^n) \in \mathcal{C}([0, T]; B_{2,2}^{s+1})$ be the approximate equations of the system (1.1):

$$\begin{cases} \partial_t u_j^n + u_j^n \cdot \nabla u_j^n - \Delta u_j^n + \nabla P_{n,j} = b_j^n \cdot \nabla b_j^n, \\ \partial_t b_j^n + u_j^n \cdot \nabla b_j^n = b_j^n \cdot \nabla u_j^n, \\ \operatorname{div} u_j^n = \operatorname{div} b_j^n = 0, \quad (u_j^n, b_j^n)|_{t=0} = S_j(u_0^n, b_0^n). \end{cases}$$

Then, according to Lemma 3.1, we have for all $t \in [0, T]$,

$$\begin{aligned} & \|u_j^n(t) - u^n(t)\|_{B_{2,2}^{s-1}}^2 + \|b_j^n(t) - b^n(t)\|_{B_{2,2}^{s-1}}^2 + \int_0^t \|\nabla(u_j^n - u^n)\|_{B_{2,2}^{s-1}}^2 d\tau \\ & \leq C(\|(\operatorname{Id} - S_j)u_0^n\|_{B_{2,2}^{s-1}}^2 + \|(\operatorname{Id} - S_j)b_0^n\|_{B_{2,2}^{s-1}}^2). \end{aligned} \quad (3.6)$$

By (3.5), we have

$$\|u_j^n\|_{B_{2,2}^{s+1}}^2 + \|b_j^n\|_{B_{2,2}^{s+1}}^2 \leq C(\|S_j u_0^n\|_{B_{2,2}^{s+1}}^2 + \|S_j b_0^n\|_{B_{2,2}^{s+1}}^2) \leq C2^j. \quad (3.7)$$

Using Lemma 3.1 again and combining (3.6)-(3.7) yields

$$\begin{aligned} & \|u_j^n(t) - u^n(t)\|_{B_{2,2}^s}^2 + \|b_j^n(t) - b^n(t)\|_{B_{2,2}^s}^2 \\ & \leq C(\|(\operatorname{Id} - S_j)u_0^n\|_{B_{2,2}^s}^2 + \|(\operatorname{Id} - S_j)b_0^n\|_{B_{2,2}^s}^2 \\ & \quad + \int_0^t \|b_j^n\|_{B_{2,2}^{s+1}}^2 (\|u_j^n - u^n\|_{B_{2,2}^{s-1}}^2 + \|\nabla u_j^n - \nabla u^n\|_{B_{2,2}^{s-1}}^2) d\tau) \\ & \leq C(\|(\operatorname{Id} - S_j)u_0^n\|_{B_{2,2}^s}^2 + \|(\operatorname{Id} - S_j)b_0^n\|_{B_{2,2}^s}^2 \\ & \quad + 2^{2j}\|(\operatorname{Id} - S_j)u_0^n\|_{B_{2,2}^{s-1}}^2 + 2^{2j}\|(\operatorname{Id} - S_j)b_0^n\|_{B_{2,2}^{s-1}}^2) \\ & \leq C(\|(\operatorname{Id} - S_j)u_0^n\|_{B_{2,2}^s}^2 + \|(\operatorname{Id} - S_j)b_0^n\|_{B_{2,2}^s}^2). \end{aligned} \quad (3.8)$$

It is easy to check that

$$\begin{aligned} & \|u_j^n - u_j^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 + \|b_j^n - b_j^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 \\ & \leq \|u_j^n - u_j^\infty\|_{L_T^\infty(B_{2,2}^{s-1})} \|u_j^n - u_j^\infty\|_{L_T^\infty(B_{2,2}^{s+1})} \\ & \quad + \|b_j^n - b_j^\infty\|_{L_T^\infty(B_{2,2}^{s-1})} \|b_j^n - b_j^\infty\|_{L_T^\infty(B_{2,2}^{s+1})} \\ & \leq C2^j (\|u_0^n - u_0^\infty\|_{B_{2,2}^s} + \|b_0^n - b_0^\infty\|_{B_{2,2}^s}) \end{aligned} \quad (3.9)$$

Therefore, combing (3.8) and (3.9), we obtain

$$\begin{aligned}
& \|u^n - u^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 + \|b^n - b^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 \\
& \leq \|u_j^n - u_j^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 + \|b_j^n - b_j^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 \\
& \quad + \|u_j^n - u^n\|_{L_T^\infty(B_{2,2}^s)}^2 + \|b_j^n - b^n\|_{L_T^\infty(B_{2,2}^s)}^2 \\
& \quad + \|u_j^\infty - u^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 + \|b_j^\infty - b^\infty\|_{L_T^\infty(B_{2,2}^s)}^2 \\
& \leq C(\|(\text{Id} - S_j)u_0^n\|_{B_{2,2}^s}^2 + \|(\text{Id} - S_j)b_0^n\|_{B_{2,2}^s}^2 + \|(\text{Id} - S_j)u_0^\infty\|_{B_{2,2}^s}^2 \\
& \quad + \|(\text{Id} - S_j)b_0^\infty\|_{B_{2,2}^s}^2 + 2^j\|u_0^n - u_0^\infty\|_{B_{2,2}^s}^2 + 2^j\|b_0^n - b_0^\infty\|_{B_{2,2}^s}^2) \\
& \leq C(\|(\text{Id} - S_j)u_0^\infty\|_{B_{2,2}^s}^2 + \|(\text{Id} - S_j)b_0^\infty\|_{B_{2,2}^s}^2 + 2^j\|u_0^n - u_0^\infty\|_{B_{2,2}^s}^2 + 2^j\|b_0^n - b_0^\infty\|_{B_{2,2}^s}^2).
\end{aligned}$$

This completes the proof of Theorem 1.2.

4. NON-UNIFORM CONTINUOUS DEPENDENCE

In this section, we will give the proof of the second theorem. Motivated by Himonas and Holliman [10], we first construct two sequence approximate solutions where velocity tends to 0 and magnetic is bounded in Sobolev H^s norms. Lately, we will show that the distance between approximate solutions and actual solutions is decaying. Finally, we can conclude that Cauchy problem (1.1) is not uniformly continuous in Sobolev $B_{2,2}^s$ spaces.

Set $0 < \delta < \frac{1}{3}$. Now, we choose the magnetic having the following form (see [10]):

$$b^{\omega,n} = b^{l,\omega,n}(t, x) + b^{h,\omega,n}(t, x), \quad \omega \in \{\pm 1\}, \quad x \in \mathbb{R}^d, \quad t \in \mathbb{R},$$

where $b^{h,\omega,n}$ is the high frequency term

$$b^{h,\omega,n}(t, x) = \begin{cases} \text{rot}\phi^h(t, x) = \left(\partial_2\phi^{h,\omega,n}(t, x), -\partial_1\phi^{h,\omega,n}(t, x)\right), & d = 2, \\ \left(\partial_2\phi^{h,\omega,n}(t, x), -\partial_1\phi^{h,\omega,n}(t, x), 0\right), & d = 3, \end{cases}$$

with

$$\phi^{h,\omega,n}(t, x) = \begin{cases} n^{-\delta-s-1}\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2 - \omega t), & d = 2, \\ n^{-\delta-s-1}\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2 - \omega t)\phi(x_3) & d = 3, \end{cases} \quad n \in \mathbb{Z},$$

where $\phi \in C_c^\infty(\mathbb{R})$ satisfies $\text{supp } \phi \in [-1, 1]$ and $\phi(x) \equiv 1$ for $|x| < \frac{1}{2}$. Let $(u^{\omega,n}, b^{l,\omega,n})$ satisfies the following initial value problem:

$$\begin{cases} \partial_t u^{\omega,n} + u^{\omega,n} \cdot \nabla u^{\omega,n} - \Delta u^{\omega,n} + \nabla P^{\omega,n} = b^{l,\omega,n} \cdot \nabla b^{l,\omega,n}, \\ \partial_t b^{l,\omega,n} + u^{\omega,n} \cdot \nabla b^{l,\omega,n} = b^{l,\omega,n} \cdot \nabla u^{\omega,n}, \\ \text{div} u^{\omega,n} = \text{div} b^{l,\omega,n} = 0, \quad (u^{\omega,n}, b^{l,\omega,n})|_{t=0} = (b_0^{l,\omega,n}, b_0^{l,\omega,n}), \end{cases} \quad (4.1)$$

$$b_0^{l,\omega,n}(x) = \begin{cases} \text{rot}\phi^l(x) = \left(\partial_2\phi^{l,\omega,n}(x), -\partial_1\phi^{l,\omega,n}(x)\right), & d = 2, \\ \left(\partial_2\phi^{l,\omega,n}(x), -\partial_1\phi^{l,\omega,n}(x), 0\right), & d = 3, \end{cases}$$

with

$$\phi^{l,\omega,n}(x) = \begin{cases} -\omega n^{-1+\delta}\Phi_1\left(\frac{x_1}{n^\delta}\right)\Phi_2\left(\frac{x_2}{n^\delta}\right), & d = 2, \\ -\omega n^{-1+\delta}\Phi_1\left(\frac{x_1}{n^\delta}\right)\Phi_2\left(\frac{x_2}{n^\delta}\right)\Phi_2\left(\frac{x_3}{n^\delta}\right) & d = 3, \end{cases} \quad n \in \mathbb{Z},$$

where the localizing functions $\Phi_1, \Phi_2 \in C_c^\infty(\mathbb{R})$ are chosen such that $\Phi_1' = \Phi_2 \equiv 1$ on the support of ϕ . By the well-posedness result, the $(u^{\omega,n}, b^{l,\omega,n})$ belong to $C([0, T]; B_{2,2}^s \times B_{2,2}^s)$ and have lifespan $T \simeq 1$. Thus, we can find that $(u^{\omega,n}, b^{\omega,n})$ satisfies the following equations:

$$\begin{cases} \partial_t u^{\omega,n} + u^{\omega,n} \cdot \nabla u^{\omega,n} - \Delta u^{\omega,n} + \nabla \tilde{P}^{\omega,n} = b^{\omega,n} \cdot \nabla b^{\omega,n} + \operatorname{div} E^{\omega,n}, \\ \partial_t b^{\omega,n} + u^{\omega,n} \cdot \nabla b^{\omega,n} = b^{\omega,n} \cdot \nabla u^{\omega,n} + F^{\omega,n}, \\ \operatorname{div} u^{\omega,n} = \operatorname{div} b^{\omega,n} = 0, \quad (u^{\omega,n}, b^{\omega,n})|_{t=0} = (b_0^{l,\omega,n}, b_0^{l,\omega,n} + b_0^{h,\omega,n}), \end{cases}$$

where

$$\begin{aligned} E^{\omega,n} &= -b^{l,\omega,n} \otimes b^{h,\omega,n} - b^{h,\omega,n} \otimes b^{h,\omega,n} - b^{h,\omega,n} \otimes b^{l,\omega,n}, \\ F^{\omega,n} &= \underbrace{\partial_t b^{h,\omega,n} + u^{\omega,n} \cdot \nabla b^{h,\omega,n}}_{F_1^{\omega,n}} - \underbrace{b^{h,\omega,n} \cdot \nabla u^{\omega,n}}_{F_2^{\omega,n}}. \end{aligned}$$

First, we consider the 2D case. To estimate the terms $E^{\omega,n}$ and $F^{\omega,n}$ in Sobolev $B_{2,2}^{s-1}$ norm, we first estimate the terms $u^{\omega,n}$ and $b^{\omega,n}$. By (3.5), Lemma 1.1 and Lemma 2.5, we have for any $r \geq 0$ and $t \in [0, T]$,

$$\begin{aligned} \|u^{\omega,n}\|_{B_{2,2}^{s+1}} + \|b^{l,\omega,n}\|_{B_{2,2}^{s+1}} &\leq C \|b_0^{l,\omega,n}\|_{B_{2,2}^{s+1}} \leq C n^{-1+\delta}, \\ \|b^{h,\omega,n}\|_{H^r} &\leq C n^{r-s}, \quad \|b^{h,\omega,n}\|_{L^\infty} \leq C n^{-1}. \end{aligned} \quad (4.2)$$

For the term $E^{\omega,n}$, we also infer from Lemma 2.3 that

$$\begin{aligned} \|E^{\omega,n}\|_{B_{2,2}^{s-1}} &\leq C \|b^{l,\omega,n}\|_{B_{2,2}^s} \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} + C \|b^{h,\omega,n}\|_{L^\infty} \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} \\ &\leq C n^{-1} (n^{-1+\delta} + n^{-\delta}). \end{aligned} \quad (4.3)$$

Now, we need to estimate the term $F^{\omega,n}$. Direct calculation shows that

$$\begin{aligned} (b^{h,\omega,n})_1 &= n^{-\delta-s} \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \\ &\quad + n^{-2\delta-s-1} \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t), \end{aligned}$$

$$(b^{h,\omega,n})_2 = -n^{-2\delta-s-1} \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t),$$

$$\begin{aligned} (\partial_t b^{h,\omega,n})_1(t, x) &= \omega n^{-s-\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\ &\quad - \omega n^{-s-1-2\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t), \end{aligned}$$

$$(\partial_t b^{h,\omega,n})_2(t, x) = \omega n^{-s-1-2\delta} \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t),$$

$$\begin{aligned}
\left(u^{\omega,n} \cdot \nabla b^{h,\omega,n}\right)_1(t,x) &= -n^{-s+1-\delta} u_2^{\omega,n}(t,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad + n^{-3\delta-s-1} u_2^{\omega,n}(t,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi''\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad + 2n^{-2\delta-s} u_2^{\omega,n}(t,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \\
&\quad + n^{-2\delta-s} u_1^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \\
&\quad + n^{-3\delta-s-1} u_1^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t),
\end{aligned}$$

$$\begin{aligned}
\left(u^{\omega,n} \cdot \nabla b^{h,\omega,n}\right)_2(t,x) &= -n^{-3\delta-s-1} u_1^{\omega,n}(t,x) \phi''\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad - n^{-3\delta-s-1} u_2^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad - n^{-2\delta-s} u_2^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t).
\end{aligned}$$

Let us handle with the first component of $F^{\omega,n}$. Using the fact

$$\begin{aligned}
&\omega n^{-s-\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&= n^{-s+1-\delta} u_2^{\omega,n}(0,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t),
\end{aligned}$$

we can obtain

$$\begin{aligned}
&\left(\partial_t b^{h,\omega,n} + u^{\omega,n} \cdot \nabla b^{h,\omega,n}\right)_1(t,x) \\
&= -n^{-s+1-\delta} [u_2^{\omega,n}(t,x) - u_2^{\omega,n}(0,x)] \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad - \omega n^{-s-1-2\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \\
&\quad + n^{-3\delta-s-1} u_2^{\omega,n}(t,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi''\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad + n^{-2\delta-s} u_1^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \\
&\quad + n^{-3\delta-s-1} u_1^{\omega,n}(t,x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \\
&\quad + 2n^{-2\delta-s} u_2^{\omega,n}(t,x) \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) := \sum_{i=1}^6 F_{1,1}^{\omega,n}(i).
\end{aligned}$$

To estimate $F_{1,1}^{\omega,n}(1)$, we can deduce from the first equation of (4.1) that

$$\begin{aligned}
& \|u^{\omega,n}(t, x) - u^{\omega,n}(0, x)\|_{B_{2,2}^s} \\
& \leq \|u^{\omega,n}(t, x) - e^{t\Delta}u^{\omega,n}(0, x)\|_{B_{2,2}^s} + \|e^{t\Delta}u^{\omega,n}(0, x) - u^{\omega,n}(0, x)\|_{B_{2,2}^s} \\
& \leq \left\| \int_0^t e^{(t-\tau)\Delta} \mathbf{P}(-u^{\omega,n} \cdot \nabla u^{\omega,n} + b^{l,\omega,n} \cdot \nabla b^{l,\omega,n}) d\tau \right\|_{B_{2,2}^s} \\
& \quad + \left\| \int_0^t \frac{d}{d\tau} (e^{\tau\Delta} u^{\omega,n}(0, x)) d\tau \right\|_{B_{2,2}^s} \\
& \leq C \| -u^{\omega,n} \cdot \nabla u^{\omega,n} + b^{l,\omega,n} \cdot \nabla b^{l,\omega,n} \|_{B_{2,2}^s} + C \|\Delta u^{\omega,n}(0, x)\|_{B_{2,2}^s} \\
& \leq Cn^{2(-1+\delta)} + Cn^{-1-\delta}, \tag{4.4}
\end{aligned}$$

where $\mathbf{P} := I + \nabla(-\Delta)^{-1} \operatorname{div}$ is the $d \times d$ matrix pseudo-differential operator in \mathbb{R}^d with the symbol $(\delta_{ij} - \frac{\xi_i \xi_j}{|\xi|^2})_{i,j=1}^d$, $d=2, 3$. By Lemma 2.3 and (4.4), we have

$$\begin{aligned}
\|F_{1,1}^{\omega,n}(1)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2+2\delta} + n^{-1-\delta}, \\
\|F_{1,1}^{\omega,n}(2)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,1}^{\omega,n}(3)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,1}^{\omega,n}(4)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta}, \\
\|F_{1,1}^{\omega,n}(5)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,1}^{\omega,n}(6)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta},
\end{aligned}$$

which yield

$$\|F_{1,1}^{\omega,n}\|_{B_{2,2}^{s-1}} = \left\| \sum_{i=1}^6 F_{1,1}^{\omega,n}(i) \right\|_{B_{2,2}^{s-1}} \leq Cn^{-1}(n^{-1+2\delta} + n^{-\delta}). \tag{4.5}$$

For the second component of $F^{\omega,n}$, we also have

$$\begin{aligned}
& \left(\partial_t b^{h,\omega,n} + u^{\omega,n} \cdot \nabla b^{h,\omega,n} \right)_2(t, x) \\
& = \omega n^{-s-1-2\delta} \phi' \left(\frac{x_1}{n^\delta} \right) \phi \left(\frac{x_2}{n^\delta} \right) \cos(nx_2 - \omega t) \\
& \quad - n^{-3\delta-s-1} u_1^{\omega,n}(t, x) \phi'' \left(\frac{x_1}{n^\delta} \right) \phi \left(\frac{x_2}{n^\delta} \right) \sin(nx_2 - \omega t) \\
& \quad - n^{-3\delta-s-1} u_2^{\omega,n}(t, x) \phi' \left(\frac{x_1}{n^\delta} \right) \phi' \left(\frac{x_2}{n^\delta} \right) \sin(nx_2 - \omega t) \\
& \quad - n^{-2\delta-s} u_2^{\omega,n}(t, x) \phi' \left(\frac{x_1}{n^\delta} \right) \phi \left(\frac{x_2}{n^\delta} \right) \cos(nx_2 - \omega t) := \sum_{i=1}^4 F_{1,2}^{\omega,n}(i).
\end{aligned}$$

By Lemma 2.3, we have

$$\begin{aligned}
\|F_{1,2}^{\omega,n}(1)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,2}^{\omega,n}(2)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,2}^{\omega,n}(3)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,2}^{\omega,n}(4)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta},
\end{aligned}$$

which yield

$$\|F_{1,2}^{\omega,n}\|_{B_{2,2}^{s-1}} = \left\| \sum_{i=1}^4 F_{1,2}^{\omega,n}(i) \right\|_{B_{2,2}^{s-1}} \leq Cn^{-1-\delta}. \quad (4.6)$$

Therefore, using Lemma 2.3, (4.2) and (4.5)-(4.6), we obtain

$$\begin{aligned} \|F^{\omega,n}\|_{B_{2,2}^{s-1}} &\leq \|F_{1,1}^{\omega,n}\|_{B_{2,2}^{s-1}} + \|F_{1,2}^{\omega,n}\|_{B_{2,2}^{s-1}} + \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} \|u^{\omega,n}\|_{B_{2,2}^{s+1}} \\ &\leq Cn^{-1}(n^{-1+2\delta} + n^{-\delta}). \end{aligned} \quad (4.7)$$

Let $(u_{\omega,n}, b_{\omega,n})$ be the unique solution of 2D non-resistive MHD equations. Namely,

$$\begin{cases} \partial_t u_{\omega,n} + u_{\omega,n} \cdot \nabla u_{\omega,n} - \Delta u_{\omega,n} + \nabla P_{\omega,n} = b_{\omega,n} \cdot \nabla b_{\omega,n}, \\ \partial_t b_{\omega,n} + u_{\omega,n} \cdot \nabla b_{\omega,n} = b_{\omega,n} \cdot \nabla u_{\omega,n}, \\ \operatorname{div} u_{\omega,n} = \operatorname{div} b_{\omega,n} = 0, \quad (u_{\omega,n}, b_{\omega,n})|_{t=0} = (b_0^{l,\omega,n}, b_0^{l,\omega,n} + b^{h,\omega,n}(0, x)). \end{cases}$$

By the well-posedness result, the $(u_{\omega,n}, b_{\omega,n})$ belong to $C([0, T]; B_{2,2}^s \times B_{2,2}^s)$ and have common lifespan $T \simeq 1$. Notice that $b^{h,1,n}(0, x) = b^{h,-1,n}(0, x)$, we have

$$\begin{aligned} &\|u_{1,n}(0, x) - u_{-1,n}(0, x)\|_{B_{2,2}^s} + \|b_{1,n}(0, x) - b_{-1,n}(0, x)\|_{B_{2,2}^s} \\ &\leq C \|b_0^{l,1,n} - b_0^{l,-1,n}\|_{B_{2,2}^s} \leq Cn^{-1+\delta} \rightarrow 0, \quad n \rightarrow \infty. \end{aligned} \quad (4.8)$$

According to Lemma 3.1, we get from (3.5) that

$$\begin{aligned} &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^{s-1})}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s-1})}^2 \leq Cn^{-2+2\delta}, \\ &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^s)}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^s)}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^s)}^2 \leq C, \\ &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s+1})}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^{s+1})}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s+1})}^2 \leq Cn^2 \end{aligned} \quad (4.9)$$

Next, considering the difference $v = u_{\omega,n} - u^{\omega,n}$ and $a = b_{\omega,n} - b^{\omega,n}$, we observe that (v, a) satisfies

$$\begin{cases} \partial_t v - \Delta v + \nabla \tilde{P}_{\omega,n} = \operatorname{div} E^{\omega,n} + \operatorname{div} G, \\ \partial_t a + u_{\omega,n} \cdot \nabla a = F^{\omega,n} + J, \\ \operatorname{div} v = \operatorname{div} a = 0, \quad (v, a)|_{t=0} = (0, 0), \end{cases}$$

where

$$\begin{aligned} G &= \underbrace{b_{\omega,n} \otimes a + a \otimes b^{\omega,n}}_{G_1} - \underbrace{v \otimes u^{\omega,n} - u_{\omega,n} \otimes v}_{G_2}, \\ J &= \underbrace{b_{\omega,n} \cdot \nabla v}_{J_1} + \underbrace{a \cdot \nabla u^{\omega,n}}_{J_2} - \underbrace{v \cdot \nabla b^{\omega,n}}_{J_3}. \end{aligned}$$

Next, we shall bound the term (v, a) in Sobolev $B_{2,2}^{s-1}$ norm. According to Lemma 3.1, we have

$$\begin{aligned} &\frac{d}{dt} (\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2) + \|\nabla v\|_{B_{2,2}^{s-1}}^2 \\ &\leq C \|E^{\omega,n}\|_{B_{2,2}^{s-1}}^2 + C \|G\|_{B_{2,2}^{s-1}}^2 + C \|F^{\omega,n}\|_{B_{2,2}^{s-1}} \|a\|_{B_{2,2}^{s-1}} \\ &\quad + C \|J\|_{B_{2,2}^{s-1}} \|a\|_{B_{2,2}^{s-1}} + \frac{1}{4} \|\nabla v\|_{B_{2,2}^{s-1}}^2. \end{aligned} \quad (4.10)$$

By Lemma 2.3, (4.2) and (4.9), we get

$$\begin{aligned}
\|G_1\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}\left(\|b^{\omega,n}\|_{B_{2,2}^s} + \|b_{\omega,n}\|_{B_{2,2}^s}\right), \\
\|G_2\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}\left(\|u^{\omega,n}\|_{B_{2,2}^s} + \|u_{\omega,n}\|_{B_{2,2}^s}\right), \\
\|J_1\|_{B_{2,2}^{s-1}} &\leq C\|b_{\omega,n}\|_{B_{2,2}^s}\|\nabla v\|_{B_{2,2}^{s-1}}, \\
\|J_2\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}\|u^{\omega,n}\|_{B_{2,2}^{s+1}}, \\
\|J_3\|_{B_{2,2}^{s-1}} &\leq C\|v\|_{B_{2,2}^s}\|b^{\omega,n}\|_{B_{2,2}^s} \leq C\|v, \nabla v\|_{B_{2,2}^{s-1}}\|b^{\omega,n}\|_{B_{2,2}^s}.
\end{aligned} \tag{4.11}$$

Plugging (4.11) into (4.10) and using (4.3), (4.7), we have for all $t \in [0, T]$,

$$\begin{aligned}
&\frac{d}{dt}\left(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2\right) \\
&\leq C\left(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2\right) + \|E^{\omega,n}\|_{B_{2,2}^{s-1}}^2 + \|F^{\omega,n}\|_{B_{2,2}^{s-1}}^2 \\
&\leq C\left(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2\right) + Cn^{-2}(n^{-1+2\delta} + n^{-\delta})^2.
\end{aligned} \tag{4.12}$$

This alongs with Growall's inequality yields

$$\|v\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s-1})}^2 \leq Cn^{-2}(n^{-1+2\delta} + n^{-\delta})^2. \tag{4.13}$$

According to the interpolation inequality and using (4.2), (4.9), (4.13), we have

$$\begin{aligned}
&\|v\|_{L_T^\infty(B_{2,2}^s)}^2 + \|a\|_{L_T^\infty(B_{2,2}^s)}^2 \\
&\leq \left(\|v\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s-1})}^2\right)^{\frac{1}{2}} \left(\|v\|_{L_T^\infty(B_{2,2}^{s+1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s+1})}^2\right)^{\frac{1}{2}} \\
&\leq C(n^{-1+2\delta} + n^{-\delta}).
\end{aligned} \tag{4.14}$$

Then, combining (4.2) and (4.14), we have

$$\begin{aligned}
&\|u_{1,n}(t) - u_{-1,n}(t)\|_{B_{2,2}^s} + \|b_{1,n}(t) - b_{-1,n}(t)\|_{B_{2,2}^s} \\
&\geq \|u^{1,n}(t) - u^{-1,n}(t)\|_{B_{2,2}^s} + \|b^{1,n}(t) - b^{-1,n}(t)\|_{B_{2,2}^s} - C\varepsilon_n \\
&\geq \|b^{h,1,n}(t) - b^{h,-1,n}(t)\|_{B_{2,2}^s} - C\varepsilon'_n \\
&\geq 2|\sin t| \cdot \|n^{-\delta-s}\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2)\|_{B_{2,2}^s} - C\varepsilon'_n \\
&\geq c_0|\sin t| \cdot \|n^{-\frac{1}{2}\delta}\phi\left(\frac{x_1}{n^\delta}\right)\|_{H^s(\mathbb{R})}\|n^{-\frac{1}{2}\delta-s}\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2)\|_{H^s(\mathbb{R})} - C\varepsilon'_n,
\end{aligned}$$

where

$$\varepsilon_n = (n^{-\delta} + n^{2\delta-1})^{\frac{1}{2}}, \quad \varepsilon'_n = (n^{-\delta} + n^{2\delta-1})^{\frac{1}{2}} + n^{\delta-1}.$$

Letting n go to ∞ , then there exists some positive constant c_0 such that

$$\begin{aligned}
&\liminf_{n \rightarrow \infty} \left(\|u_{1,n}(t) - u_{-1,n}(t)\|_{B_{2,2}^s} + \|b_{1,n}(t) - b_{-1,n}(t)\|_{B_{2,2}^s}\right) \\
&\geq c_0|\sin t|.
\end{aligned} \tag{4.15}$$

Then (4.15) together with (4.8) complete the proof of the 2D case of Theorem 1.3.

Now, we consider the 3D case. Similarly, to estimate the terms $E^{\omega,n}$ and $F^{\omega,n}$ in Sobolev $B_{2,2}^{s-1}$ norm, we first estimate the terms $u^{\omega,n}$ and $b^{\omega,n}$. By (3.5), Lemma

1.1 and Lemma 2.5, we get for any $r \geq 0$ and $t \in [0, T]$,

$$\begin{aligned} \|u^{\omega,n}\|_{B_{2,2}^{s+1}} + \|b^{l,\omega,n}\|_{B_{2,2}^{s+1}} &\leq C \|b_0^{l,\omega,n}\|_{B_{2,2}^{s+1}} \leq C n^{-1+\frac{3}{2}\delta}, \\ \|b^{h,\omega,n}\|_{H^r} &\leq C n^{r-s}, \quad \|b^{h,\omega,n}\|_{L^\infty} \leq C n^{-1}. \end{aligned} \quad (4.16)$$

For $E^{\omega,n}$, by Lemma 2.3, we have

$$\begin{aligned} \|E^{\omega,n}\|_{B_{2,2}^{s-1}} &\leq C \|b^{l,\omega,n}\|_{B_{2,2}^s} \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} + C \|b^{h,\omega,n}\|_{L^\infty} \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} \\ &\leq C n^{-1} (n^{-1+\frac{3}{2}\delta} + n^{-\delta}). \end{aligned} \quad (4.17)$$

Now, we estimate $F^{\omega,n}$. Direct calculation shows that

$$\left(\partial_t b^{h,\omega,n} + u^{\omega,n} \cdot \nabla b^{h,\omega,n} \right)_3(t, x) = 0,$$

$$\begin{aligned} \left(b^{h,\omega,n} \right)_1 &= n^{-\delta-s} \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3) \\ &\quad + n^{-2\delta-s-1} \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3), \end{aligned}$$

$$\left(b^{h,\omega,n} \right)_2 = -n^{-2\delta-s-1} \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3),$$

$$\begin{aligned} \left(\partial_t b^{h,\omega,n} \right)_1(t, x) &= \omega n^{-s-\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\ &\quad - \omega n^{-s-1-2\delta} \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3), \end{aligned}$$

$$\left(\partial_t b^{h,\omega,n} \right)_2(t, x) = \omega n^{-s-1-2\delta} \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3),$$

$$\begin{aligned} \left(u^{\omega,n} \cdot \nabla b^{h,\omega,n} \right)_1(t, x) &= -n^{-s+1-\delta} u_2^{\omega,n}(t, x) \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\ &\quad + n^{-3\delta-s-1} u_2^{\omega,n}(t, x) \phi\left(\frac{x_1}{n^\delta}\right) \phi''\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\ &\quad + 2n^{-2\delta-s} u_2^{\omega,n}(t, x) \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3) \\ &\quad + n^{-2\delta-s} u_1^{\omega,n}(t, x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3) \\ &\quad + n^{-3\delta-s-1} u_1^{\omega,n}(t, x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\ &\quad + n^{-\delta-s} u_3^{\omega,n}(t, x) \phi\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \partial_3 \phi(x_3) \\ &\quad + n^{-2\delta-s-1} u_3^{\omega,n}(t, x) \phi\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \partial_3 \phi(x_3), \end{aligned}$$

$$\begin{aligned}
\left(u^{\omega,n} \cdot \nabla b^{h,\omega,n}\right)_2(t,x) &= -n^{-3\delta-s-1}u_1^{\omega,n}(t,x)\phi''\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&\quad - n^{-3\delta-s-1}u_2^{\omega,n}(t,x)\phi'\left(\frac{x_1}{n^\delta}\right)\phi'\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&\quad - n^{-2\delta-s}u_2^{\omega,n}(t,x)\phi'\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\cos(nx_2-\omega t)\phi(x_3) \\
&\quad - n^{-2\delta-s-1}u_3^{\omega,n}(t,x)\phi'\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\partial_3\phi(x_3).
\end{aligned}$$

For the first component of $F^{\omega,n}$. It is easy to see that

$$\begin{aligned}
&\omega n^{-s-\delta}\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&= n^{-s+1-\delta}u_2^{\omega,n}(0,x)\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3),
\end{aligned}$$

we can obtain

$$\begin{aligned}
&\left(\partial_t b^h + u^{\omega,n} \cdot \nabla b^{h,\omega,n}\right)_1(t,x) \\
&= -n^{-s+1-\delta}[u_2^{\omega,n}(t,x) - u_2^{\omega,n}(0,x)]\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&\quad - \omega n^{-s-1-2\delta}\phi\left(\frac{x_1}{n^\delta}\right)\phi'\left(\frac{x_2}{n^\delta}\right)\cos(nx_2-\omega t)\phi(x_3) \\
&\quad + n^{-3\delta-s-1}u_2^{\omega,n}(t,x)\phi\left(\frac{x_1}{n^\delta}\right)\phi''\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&\quad + n^{-2\delta-s}u_1^{\omega,n}(t,x)\phi'\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\cos(nx_2-\omega t)\phi(x_3) \\
&\quad + n^{-3\delta-s-1}u_1^{\omega,n}(t,x)\phi'\left(\frac{x_1}{n^\delta}\right)\phi'\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\phi(x_3) \\
&\quad + 2n^{-2\delta-s}u_2^{\omega,n}(t,x)\phi\left(\frac{x_1}{n^\delta}\right)\phi'\left(\frac{x_2}{n^\delta}\right)\cos(nx_2-\omega t)\phi(x_3) \\
&\quad + n^{-\delta-s}u_3^{\omega,n}(t,x)\phi\left(\frac{x_1}{n^\delta}\right)\phi\left(\frac{x_2}{n^\delta}\right)\cos(nx_2-\omega t)\partial_3\phi(x_3) \\
&\quad + n^{-2\delta-s-1}u_3^{\omega,n}(t,x)\phi\left(\frac{x_1}{n^\delta}\right)\phi'\left(\frac{x_2}{n^\delta}\right)\sin(nx_2-\omega t)\partial_3\phi(x_3) := \sum_{i=1}^8 F_{1,1}^{\omega,n}(i).
\end{aligned}$$

To estimate $F_{1,1}^{\omega,n}(1)$, similarly to the 2D case, we get

$$\begin{aligned}
&\|u^{\omega,n}(t,x) - u^{\omega,n}(0,x)\|_{B_{2,2}^s} \\
&\leq \|u^{\omega,n}(t,x) - e^{t\Delta}u^{\omega,n}(0,x)\|_{B_{2,2}^s} + \|e^{t\Delta}u^{\omega,n}(0,x) - u^{\omega,n}(0,x)\|_{B_{2,2}^s} \\
&\leq \left\| \int_0^t e^{(t-\tau)\Delta} \mathbf{P}(-u^{\omega,n} \cdot \nabla u^{\omega,n} + b^{l,\omega,n} \cdot \nabla b^{l,\omega,n}) d\tau \right\|_{B_{2,2}^s} \\
&\quad + \left\| \int_0^t \frac{d}{d\tau} (e^{\tau\Delta} u^{\omega,n}(0,x)) d\tau \right\|_{B_{2,2}^s} \\
&\leq C \| -u^{\omega,n} \cdot \nabla u^{\omega,n} + b^{l,\omega,n} \cdot \nabla b^{l,\omega,n} \|_{B_{2,2}^s} + C \|\Delta u^{\omega,n}(0,x)\|_{B_{2,2}^s} \\
&\leq C n^{2(-1+\frac{3}{2}\delta)} + C n^{-1-\frac{1}{2}\delta}.
\end{aligned} \tag{4.18}$$

By Lemma 2.3 and (4.18), we have

$$\begin{aligned}
\|F_{1,1}^{\omega,n}(1)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2+3\delta} + n^{-1-\frac{1}{2}\delta}, \\
\|F_{1,1}^{\omega,n}(2)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,1}^{\omega,n}(3)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,1}^{\omega,n}(4)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta}, \\
\|F_{1,1}^{\omega,n}(5)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,1}^{\omega,n}(6)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta}, \\
\|F_{1,1}^{\omega,n}(7)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2+\frac{3}{2}\delta}, \\
\|F_{1,1}^{\omega,n}(8)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta},
\end{aligned}$$

which yield

$$\|F_{1,1}^{\omega,n}\|_{B_{2,2}^{s-1}} = \left\| \sum_{i=1}^8 F_{1,1}^{\omega,n}(i) \right\|_{B_{2,2}^{s-1}} \leq Cn^{-1}(n^{-1+3\delta} + n^{-\frac{1}{2}\delta}). \quad (4.19)$$

For the second component of $F^{\omega,n}$, we also have

$$\begin{aligned}
&\left(\partial_t b^h + u^{\omega,n} \cdot \nabla b^{h,\omega,n} \right)_2(t, x) \\
&= \omega n^{-s-1-2\delta} \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3) \\
&\quad - n^{-3\delta-s-1} u_1^{\omega,n}(t, x) \phi''\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\
&\quad - n^{-3\delta-s-1} u_2^{\omega,n}(t, x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi'\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \phi(x_3) \\
&\quad - n^{-2\delta-s} u_2^{\omega,n}(t, x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \cos(nx_2 - \omega t) \phi(x_3) \\
&\quad - n^{-2\delta-s-1} u_3^{\omega,n}(t, x) \phi'\left(\frac{x_1}{n^\delta}\right) \phi\left(\frac{x_2}{n^\delta}\right) \sin(nx_2 - \omega t) \partial_3 \phi(x_3) := \sum_{i=1}^5 F_{1,2}^{\omega,n}(i).
\end{aligned}$$

By Lemma 2.3, we obtain

$$\begin{aligned}
\|F_{1,2}^{\omega,n}(1)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta}, \\
\|F_{1,2}^{\omega,n}(2)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,2}^{\omega,n}(3)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-2\delta}, \\
\|F_{1,2}^{\omega,n}(4)\|_{B_{2,2}^{s-1}} &\lesssim n^{-1-\delta}, \\
\|F_{1,2}^{\omega,n}(5)\|_{B_{2,2}^{s-1}} &\lesssim n^{-2-\delta},
\end{aligned}$$

which yield

$$\|F_{1,2}^{\omega,n}\|_{B_{2,2}^{s-1}} = \left\| \sum_{i=1}^4 F_{1,2}^{\omega,n}(i) \right\|_{B_{2,2}^{s-1}} \leq Cn^{-1-\delta}. \quad (4.20)$$

Thus, applying Lemma 2.3, (4.16) and (4.19)-(4.20), we obtain

$$\begin{aligned} \|F^{\omega,n}\|_{B_{2,2}^{s-1}} &\leq \|F_{1,1}^{\omega,n}\|_{B_{2,2}^{s-1}} + \|F_{1,2}^{\omega,n}\|_{B_{2,2}^{s-1}} + \|b^{h,\omega,n}\|_{B_{2,2}^{s-1}} \|u^{\omega,n}\|_{B_{2,2}^{s+1}} \\ &\leq Cn^{-1}(n^{-1+3\delta} + n^{-\frac{1}{2}\delta}). \end{aligned} \quad (4.21)$$

Let $(u_{\omega,n}, b_{\omega,n})$ be the unique solution of 3D non-resistive MHD equations. Namely,

$$\begin{cases} \partial_t u_{\omega,n} + u_{\omega,n} \cdot \nabla u_{\omega,n} - \Delta u_{\omega,n} + \nabla P_{\omega,n} = b_{\omega,n} \cdot \nabla b_{\omega,n}, \\ \partial_t b_{\omega,n} + u_{\omega,n} \cdot \nabla b_{\omega,n} = b_{\omega,n} \cdot \nabla u_{\omega,n}, \\ \operatorname{div} u_{\omega,n} = \operatorname{div} b_{\omega,n} = 0, \quad (u_{\omega,n}, b_{\omega,n})|_{t=0} = (b_0^{l,\omega,n}, b_0^{l,\omega,n} + b^{h,\omega,n}(0, x)). \end{cases}$$

By the well-posedness result, the $(u_{\omega,n}, b_{\omega,n})$ belong to $C([0, T]; B_{2,2}^s \times B_{2,2}^s)$ and have common lifespan $T \simeq 1$. Notice that $b^{h,1,n}(0, x) = b^{h,-1,n}(0, x)$, we obtain

$$\begin{aligned} &\|u_{1,n}(0, x) - u_{-1,n}(0, x)\|_{B_{2,2}^s} + \|b_{1,n}(0, x) - b_{-1,n}(0, x)\|_{B_{2,2}^s} \\ &\leq C \|b_0^{l,1,n} - b_0^{l,-1,n}\|_{B_{2,2}^s} \leq Cn^{-1+\frac{3}{2}\delta} \rightarrow 0, \quad n \rightarrow \infty. \end{aligned} \quad (4.22)$$

By Lemma 3.1, we get from (3.5) that

$$\begin{aligned} &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^{s-1})}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s-1})}^2 \leq Cn^{-2+3\delta}, \\ &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^s)}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^s)}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^s)}^2 \leq C, \\ &\|u_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s+1})}^2 + \|\nabla u_{\omega,n}\|_{L_T^2(B_{2,2}^{s+1})}^2 + \|b_{\omega,n}\|_{L_T^\infty(B_{2,2}^{s+1})}^2 \leq Cn^2 \end{aligned} \quad (4.23)$$

Next, considering $v = u_{\omega,n} - u^{\omega,n}$ and $a = b_{\omega,n} - b^{\omega,n}$, we observe that (v, a) satisfies

$$\begin{cases} \partial_t v - \Delta v + \nabla \tilde{P}_{\omega,n} = \operatorname{div} E^{\omega,n} + \operatorname{div} G, \\ \partial_t a + u_{\omega,n} \cdot \nabla a = F^{\omega,n} + J, \\ \operatorname{div} v = \operatorname{div} a = 0, \quad (v, a)|_{t=0} = (0, 0), \end{cases}$$

where

$$\begin{aligned} G &= \underbrace{b_{\omega,n} \otimes a + a \otimes b^{\omega,n}}_{G_1} - \underbrace{v \otimes u^{\omega,n} - u_{\omega,n} \otimes v}_{G_2}, \\ J &= \underbrace{b_{\omega,n} \cdot \nabla v}_{J_1} + \underbrace{a \cdot \nabla u^{\omega,n}}_{J_2} - \underbrace{v \cdot \nabla b^{\omega,n}}_{J_3}. \end{aligned}$$

Next, we shall bound the term (v, a) in Sobolev $B_{2,2}^{s-1}$ norm. By Lemma 3.1, we obtain

$$\begin{aligned} &\frac{d}{dt} (\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2) + \|\nabla v\|_{B_{2,2}^{s-1}}^2 \\ &\leq C \|E^{\omega,n}\|_{B_{2,2}^{s-1}}^2 + C \|G\|_{B_{2,2}^{s-1}}^2 + C \|F^{\omega,n}\|_{B_{2,2}^{s-1}} \|a\|_{B_{2,2}^{s-1}} \\ &\quad + C \|J\|_{B_{2,2}^{s-1}} \|a\|_{B_{2,2}^{s-1}} + \frac{1}{4} \|\nabla v\|_{B_{2,2}^{s-1}}^2. \end{aligned} \quad (4.24)$$

By Lemma 2.3, (4.16) and (4.23), we get

$$\begin{aligned}
\|G_1\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}(\|b^{\omega,n}\|_{B_{2,2}^s} + \|b_{\omega,n}\|_{B_{2,2}^s}), \\
\|G_2\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}(\|u^{\omega,n}\|_{B_{2,2}^s} + \|u_{\omega,n}\|_{B_{2,2}^s}), \\
\|J_1\|_{B_{2,2}^{s-1}} &\leq C\|b_{\omega,n}\|_{B_{2,2}^s}\|\nabla v\|_{B_{2,2}^{s-1}}, \\
\|J_2\|_{B_{2,2}^{s-1}} &\leq C\|a\|_{B_{2,2}^{s-1}}\|u^{\omega,n}\|_{B_{2,2}^{s+1}}, \\
\|J_3\|_{B_{2,2}^{s-1}} &\leq C\|v\|_{B_{2,2}^s}\|b^{\omega,n}\|_{B_{2,2}^s} \leq C\|v, \nabla v\|_{B_{2,2}^{s-1}}\|b^{\omega,n}\|_{B_{2,2}^s}.
\end{aligned} \tag{4.25}$$

Plugging (4.25) into (4.24) and applying (4.17), (4.21), we obtain for all $t \in [0, T]$,

$$\begin{aligned}
&\frac{d}{dt}(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2) \\
&\leq C(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2) + \|E^{\omega,n}\|_{B_{2,2}^{s-1}}^2 + \|F^{\omega,n}\|_{B_{2,2}^{s-1}}^2 \\
&\leq C(\|v\|_{B_{2,2}^{s-1}}^2 + \|a\|_{B_{2,2}^{s-1}}^2) + Cn^{-2}(n^{-1+3\delta} + n^{-\frac{1}{2}\delta})^2.
\end{aligned} \tag{4.26}$$

This alongs with Grownall's inequality yields

$$\|v\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s-1})}^2 \leq Cn^{-2}(n^{-1+3\delta} + n^{-\frac{1}{2}\delta})^2. \tag{4.27}$$

By (4.16), (4.23), (4.27) and using the interpolation inequality, we have

$$\begin{aligned}
&\|v\|_{L_T^\infty(B_{2,2}^s)}^2 + \|a\|_{L_T^\infty(B_{2,2}^s)}^2 \\
&\leq (\|v\|_{L_T^\infty(B_{2,2}^{s-1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s-1})}^2)^{\frac{1}{2}}(\|v\|_{L_T^\infty(B_{2,2}^{s+1})}^2 + \|a\|_{L_T^\infty(B_{2,2}^{s+1})}^2)^{\frac{1}{2}} \\
&\leq C(n^{-1+3\delta} + n^{-\frac{1}{2}\delta}).
\end{aligned} \tag{4.28}$$

Then, combining (4.16) and (4.28), we get

$$\begin{aligned}
&\|u_{1,n}(t) - u_{-1,n}(t)\|_{B_{2,2}^s} + \|b_{1,n}(t) - b_{-1,n}(t)\|_{B_{2,2}^s} \\
&\geq \|u^{1,n}(t) - u^{-1,n}(t)\|_{B_{2,2}^s} + \|b^{1,n}(t) - b^{-1,n}(t)\|_{B_{2,2}^s} - C\varepsilon_n \\
&\geq \|b^{h,1,n}(t) - b^{h,-1,n}(t)\|_{B_{2,2}^s} - C\varepsilon'_n \\
&\geq 2|\sin t| \cdot \|n^{-\delta-s}\phi(\frac{x_1}{n^\delta})\phi(\frac{x_2}{n^\delta})\sin(nx_2)\phi(x_3)\|_{B_{2,2}^s} - C\varepsilon'_n \\
&\geq c_0|\sin t| \cdot \|n^{-\frac{1}{2}\delta}\phi(\frac{x_1}{n^\delta})\|_{H^s(\mathbb{R})}\|n^{-\frac{1}{2}\delta-s}\phi(\frac{x_2}{n^\delta})\sin(nx_2)\|_{H^s(\mathbb{R})}\|\phi(x_3)\|_{H^s(\mathbb{R})} - C\varepsilon'_n,
\end{aligned}$$

where

$$\varepsilon_n = (n^{-\frac{1}{2}\delta} + n^{3\delta-1})^{\frac{1}{2}}, \quad \varepsilon'_n = (n^{-\frac{1}{2}\delta} + n^{3\delta-1})^{\frac{1}{2}} + n^{\frac{3}{2}\delta-1}.$$

Letting n go to ∞ , then there exists some positive constant \tilde{c}_0 such that

$$\begin{aligned}
&\liminf_{n \rightarrow \infty} (\|u_{1,n}(t) - u_{-1,n}(t)\|_{B_{2,2}^s} + \|b_{1,n}(t) - b_{-1,n}(t)\|_{B_{2,2}^s}) \\
&\geq \tilde{c}_0|\sin t|.
\end{aligned} \tag{4.29}$$

Then (4.29) together with (4.22) complete the proof of 3D case.

Acknowledgements. J. Li was partially supported by NSFC (No.11801090) and Z. Yin was partially supported by NSFC (No.11671407 and No.11271382), FDCT (No. 098/2013/A3), Guangdong Special Support Program (No. 8-2015), and the key project of NSF of Guangdong province (No. 2016A030311004).

REFERENCES

- [1] H. Abidi and P. Zhang, On the global solution of a 3-D MHD system with initial data near equilibrium, *Comm. Pure Appl. Math.* **70** (2017), 1509–1561.
- [2] H. Bahouri, J. Chemin and R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren der mathematischen Wissenschaften, 343, Springer, Heidelberg (2011).
- [3] J. L. Bona and R. Smith, The initial-value problem for the Korteweg-de Vries equation, *Philos. Trans. R. Soc. Lond. Ser. A*, **278** (1975), 555–601.
- [4] J. Chemin, D. McCormick, J. Robinson and J. Rodrigo, Local existence for the non-resistive MHD equations in Besov spaces, *Adv. Math.*, **286** (2016), 1–31.
- [5] Y. Chen and P. Zhang, The global existence of small solutions to the incompressible viscoelastic fluid system in 2 and 3 space dimensions, *Comm. Partial Differential Equations*, **31** (2006), 1793–1810.
- [6] C. Fefferman, D. McCormick, J. Robinson and J. Rodrigo, Higher order commutator estimates and local existence for the non-resistive MHD equations and related models, *J. Funct. Anal.*, **267** (2014), 1035–1056.
- [7] C. Fefferman, D. McCormick, J. Robinson and J. Rodrigo, Local existence for the non-resistive MHD equations in nearly optimal Sobolev spaces, *Arch. Ration. Mech. Anal.*, **223** (2017), 677–691.
- [8] Z. Guo, J. Li and Z. Yin, Local well-posedness of the incompressible Euler equations in $B_{\infty,1}^1$ and the inviscid limit of the Navier-Stokes equations, *J. Funct. Anal.*, (2018), <https://doi.org/10.1016/j.jfa.2018.07.004>.
- [9] A. A. Himonas and C. Holliman, The Cauchy problem for the Novikov equation, *Nonlinearity*, **25** (2012), 449–479.
- [10] A. A. Himonas and G. Misiolek, Non-uniform dependence on initial data of solutions to the Euler equations of hydrodynamics, *Comm. Math. Phys.*, **296** (2010), 285–301.
- [11] X. Hu, F. Lin, Global existence for two dimensional incompressible magnetohydrodynamic flows with zero magnetic diffusivity, [arXiv:1405.0082](https://arxiv.org/abs/1405.0082).
- [12] Q. Jiu and D. Niu, Mathematical results related to a two-dimensional magnetohydrodynamic equations, *Acta Math. Sci. Ser. B Engl. Ed.* **26**, (2006), 744–756.
- [13] J. Li, W. Tan and Z. Yin, Local existence and uniqueness for the non-resistive MHD equations in homogeneous Besov spaces, *Adv. Math.*, **317** (2017), 786–798.
- [14] J. Li and Z. Yin, Well-posedness and analytic solutions of the two-component Euler-Poincaré system, *Monatsh. Math.*, **183** (2017), 509–537.
- [15] F. Lin, L. Xu and P. Zhang, Global small solutions of 2-D incompressible MHD system, *J. Differential Equations* **259** (2015), 5440–5485.
- [16] F. Lin and T. Zhang, Global small solutions to a complex fluid model in three dimensional, *Arch. Ration. Mech. Anal.* **216** (2015), 905–920.
- [17] X. Ren, J. Wu, Z. Xiang and Z. Zhang, Global existence and decay of smooth solution for the 2-D MHD equations without magnetic diffusion, *J. Funct. Anal.* **267** (2014), 503–541.
- [18] R. Wan, On the uniqueness for the 2D MHD equations without magnetic diffusion, *Nonlinear Anal. Real World Appl.*, **30** (2016), 32–40.
- [19] L. Xu and P. Zhang, Global small solutions to three-dimensional incompressible magnetohydrodynamical system, *SIAM J. Math. Anal.* **47** (2015), 26–65.
- [20] C. Zhai, and T. Zhang, Global existence and uniqueness theorem to 2-D incompressible non-resistive MHD system with non-equilibrium background magnetic field, *J. Differential Equations* **261** (2016), 3519–3550.
- [21] T. Zhang, An elementary proof of the global existence and uniqueness theorem to 2-D incompressible non-resistive MHD system, [arXiv:1404.5681](https://arxiv.org/abs/1404.5681).
- [22] T. Zhang, Global solutions to the 2D viscous, non-resistive MHD system with large background magnetic field, *J. Differential Equations* **260** (2016), 5450–5480.

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCES, GANNAN NORMAL UNIVERSITY, GANZHOU
341000, CHINA

E-mail address: lij129@mail2.sysu.edu.cn

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, CHINA
& FACULTY OF INFORMATION TECHNOLOGY, MACAU UNIVERSITY OF SCIENCE AND TECHNOLOGY,
MACAU, CHINA

E-mail address: mcsyzy@mail.sysu.edu.cn

SCHOOL OF MATHEMATICS AND INFORMATION SCIENCE, GUANGZHOU UNIVERSITY, GUANGZHOU,
510006, CHINA

E-mail address: mathzwp2010@163.com