FUNCTION THEORETICAL APPLICATIONS OF LAGRANGIAN SPECTRAL INVARIANTS

MORIMICHI KAWASAKI

ABSTRACT. Entov and Polterovich considered the concept of heaviness and superheaviness by the Oh-Schwarz spectral invariants. The Oh-Schwarz spectral invariants are defined in terms of the Hamiltonian Floer theory. In this paper, we define heaviness and superheaviness by spectral invariants defined in terms of the Lagrangian Floer theory and provide their applications. As one of them, we define a relative symplectic capacity which measures the existence of Hamiltonian chord between two disjoint Lagrangian submanifolds and provide an upper bound of it in a special case. We also provide applications to non-degeneracy of spectral norms, a relative energy capacity inequality, fragmentation norm and non-displaceability.

1. PRINCIPAL RESULTS

1.1. **Introduction.** In a series of papers by Entov and Polterovich *et al.*, they constructed a fertile theory by asymptotic Oh-Schwarz spectral invariants. Polterovich and Rosen's book "Function theory on symplectic manifolds" [PR] is a good survey of their works.

In their theory, (abstractly) displaceable subsets play important roles in many situations. In the present paper, we consider a relative version of their theory *i.e.* we replace abstract displaceability in their theory with relative displaceability from a fixed Lagrangian submanifold. Our main tool is a Lagrangian spectral invariant whose properties are studied by Leclercq and Zapolsky [LZ].

We provide applications of our observation to non-displaceability, a relative version of energy capacity inequality, Poisson bracket invariant for open covers and existence problem of a Hamiltonian chord between two disjoint submanifolds.

1.2. Notions on Hamiltonian isotopies and monotonicity of Lagrangian submanifolds. Let (M, ω) be a symplectic manifold. For a Hamiltonian function $H: M \to \mathbb{R}$ with compact support, we define the *Hamiltonian vector field* X_H associated with H by

$$\omega(X_H, V) = -dH(V)$$
 for any $V \in \mathcal{X}(M)$,

where X(M) is the set of smooth vector fields on M.

Let S^1 denote \mathbb{R}/\mathbb{Z} . For a (time-dependent) Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ with compact support and for $t \in S^1$, we define $H_t: M \to \mathbb{R}$ by $H_t(x) = H(t, x)$. Let X_H^t denote the Hamiltonian vector field associated with H_t and $\{\phi_H^t\}_{t\in\mathbb{R}}$ denote the isotopy generated by X_H^t such that $\phi^0 = \text{id.}$ Let ϕ_H denote ϕ_H^1 and ϕ_H is called *the Hamiltonian diffeomorphism generated by* H. For $x \in M$, let $\gamma_H^x: [0, 1] \to M$ denote the path defined by $\gamma_H^x(t) = \phi_H^t(x)$.

Let *X*, *Y* be subsets of a symplectic manifold (M, ω) . *X* is *displaceable* from *Y* if there exists a Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ such that $\phi_H(X) \cap \overline{Y} = \emptyset$, where \overline{Y} is the topological closure of *Y*. *X* is *non-displaceable* from *Y* otherwise. *X* is *(abstractly) displaceable* if *X* is displaceable from *X* itself.

For Hamiltonian functions $F, G: S^1 \times M \to \mathbb{R}$, let $F \models G: S^1 \times M \to \mathbb{R}$ denote a Hamiltonian function defined by $(F \models G)(t, x) = F(t, x) + G(t, (\phi_F^t)^{-1}(x))$. For a Hamiltonian function

 $H: S^1 \times M \to \mathbb{R}$, let $\overline{H}: S^1 \times M \to \mathbb{R}$ denote a Hamiltonian function defined by $\overline{H}(t, x) = -H(t, (\phi_H^t)^{-1}(x))$. Note that $\phi_{F \models G}^t = \phi_F^t \phi_G^t$ and $\phi_{\overline{H}}^t = (\phi_H^t)^{-1}$.

A Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ is *normalized* if $\int_M H_t \omega^m = 0$ for any *t*, where $m = \frac{1}{2} \dim(M)$.

For a Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ with compact support on a symplectic manifold M, we define *the Hofer length* ||H|| of H by

$$||H|| = \int_0^1 (\max_{x \in M} H_t(x) - \min_{x \in M} H_t(x)) dt.$$

For subsets X, Y of M, we define the displacement energy E(X; Y) of X from Y by

$$E(X;Y) = \inf\{||H||; H \in C_c^{\infty}(S^1 \times M), \phi_H^1(X) \cap \overline{Y} = \emptyset\},\$$

where \overline{Y} is the topological closure of Y. If X is non-displaceable from Y, we define $E(X; Y) = +\infty$. We also define another invariant $\overline{E}(X; Y)$ by

$$\overline{E}(X;Y) = \min\{E(X;X), E(X;Y)\}$$

For a closed Lagrangian submanifold L, we have the following two homomorphisms.

$$\omega \colon \pi_2(M, L) \to \mathbb{R}$$
, the symplectic area,

 μ : $\pi_2(M, L) \rightarrow \mathbb{Z}$, the *Maslov index*.

For a positive number λ , *L* is said to be λ -monotone if

$$\omega(A) = \lambda \cdot \mu(A)$$
 for any $A \in \pi_2(M, L)$.

L is said to be *monotone* if *L* is λ -monotone for some positive number λ . Note that *L* is λ -monotone for any positive number λ if *L* is a Lagrangian submanifold with $\pi_2(M, L) = 0$.

The minimal Maslov number N_L of L is defined to be the positive generator of the subgroup $\mu(\pi_2(M, L))$ of \mathbb{Z} if it is nontrivial, and we set $N_L = \infty$ otherwise.

1.3. Lagrangian heaviness and superheaviness. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$.

For an element a of the quantum homology $QH_*(L)$ and a continuous function $H: S^1 \times M \to \mathbb{R}$, Leclercq and Zapolsky defined the Lagrangian spectral invariant $c^L(a, H) \in \mathbb{R}$ (See Section 2).

For an idempotent **a** of the quantum homology $QH_*(L)$, we define the functional ζ_a^L : $C^0(S^1 \times M) \to \mathbb{R}$ as the stabilization of $c^L(\mathbf{a}, \cdot)$;

$$\zeta_{\mathsf{a}}^{L}(H) = \lim_{k \to +\infty} \frac{c^{L}(\mathsf{a}, H^{\natural k})}{k},$$

where a function $H^{\natural k} \colon S^1 \times M \to \mathbb{R}$ is defined by

$$H^{\natural k} = \underbrace{H \natural \cdots \natural H}_{k}.$$

Definition 1.1. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$, $QH_*(L) \ne 0$ and a an idempotent of the quantum homology $QH_*(L)$. A closed subset *X* of (M, ω) is said to be (L, a)-heavy if

$$\zeta_{a}^{L}(H) \ge \inf_{S^{1} \times X} H$$
 for any continuous function $H: S^{1} \times M \to \mathbb{R}$.

X is said to be (L, a)-superheavy if

 $\zeta_{a}^{L}(H) \leq \sup_{S^{1} \times Y} H$ for any continuous function $H: S^{1} \times M \to \mathbb{R}$.

A closed subset X of (M, ω) is called *L*-(*super*)heavy in the Lagrangian sense if X is (L, a)-(super)heavy for some idempotent a of $QH_*(L)$.

By an argument similar to [EP09], we see that X is (L, a)-heavy if X is (L, a)-superheavy. By Theorem 39 of [LZ], we can prove the following theorem.

Theorem 1.2. Let L_1, L_2 be monotone Lagrangian submanifolds with same monotonicity constants of closed symplectic manifolds $(M_1, \omega_1), (M_2, \omega_2)$ with $N_{L_1} \ge 2, N_{L_2} \ge 2$, respectively. Let X_1, X_2 be a (L_1, a_1) -heavy, (L_2, a_2) -heavy subset, respectively. Then $X_1 \times X_2$ is a $(L_1 \times L_2, a_1 \times a_2)$ heavy subset.

1.4. **Examples.** We provide some examples of heavy and superheavy subsets. A priority of Entov-Polterovich's theory is that we can prove non-displaceability of singular subsets. Entov and Polterovich proved that a stem which can be singular is superheavy ([EP09]). We pose a generalization of concept of stem.

Definition 1.3. Let *L* be a subset of a closed symplectic manifold *M*. Let \mathbb{A} be a finitedimensional Poisson-commutative subspace of $C^{\infty}(M)$ and $\Phi: M \to \mathbb{A}^*$ be the moment map defined by $\langle \Phi(x), F \rangle = F(x)$. A non-empty fiber $\Phi^{-1}(p), p \in \mathbb{A}^*$ is called *an L-stem* of \mathbb{A} if each non-empty fibers $\Phi^{-1}(q)$ with $q \neq p$ is displaceable from *L* or $\Phi^{-1}(q)$ itself. If a subset *X* of *M* is an *L*-stem of a finite-dimensional Poisson-commutative subspace of $C^{\infty}(M)$, *X* is called just *an L-stem*.

For example, L itself is an L-stem if L is a closed subset of M. Any stem in the original sense of [EP09] is also an L-stem for any L.

Theorem 1.4. Let L be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Then any L-stem is (L, a)-superheavy for any non-trivial idempotent a.

It is a natural problem whether (super)heaviness with respect to some idempotent implies (super)heaviness with respect to another idempotent. We provide the following two results. In Section 3, we provide more generalized theorems.

Corollary 1.5. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let e' be an idempotent of $QH_*(L)$. Then

- Any [M]-superheavy subset X is (L, e')-superheavy,
- Any (L, e')-heavy subset Y is [M]-heavy, in particular, non-displaceable from Y itself.

Corollary 1.6. Let L be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let e' be a non-trivial idempotent of $QH_*(L)$. Then any (L, e')-heavy subset X is (L, [L])-heavy and any (L, [L])-superheavy subset Y is (L, e')-superheavy.

The proof of Corollary 1.6 is similar to Theorem 1.5 of [EP09] and thus we omit the proof.

We provide many examples of heavy or superheavy subsets when M is a 2-torus. For a positive number R, we define the 2-torus T_R^2 by $T_R^2 = \mathbb{R}/R\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$ with the coordinates (x, y). For $s \in \mathbb{R}/R\mathbb{Z}$ and $t \in \mathbb{R}/\mathbb{Z}$, let L_m^s and L_1^t be a meridian curve $\{x = s\}$ and a longitude curve $\{y = t\}$, respectively.

We have many examples of heavy, superheavy subsets of T_R^2 .

Proposition 1.7. For any positive number R, the following propositions hold on T_R^2 .

- (1) $L_{\rm m}^{s'}$ is $(L_{\rm m}^s, [L_{\rm m}^s])$ -superheavy if s = s'.
- (2) $L_{m}^{s'}$ is not $(L_{m}^{s}, [L_{m}^{s}])$ -heavy if $s \neq s'$.
- (3) L_1^t is $(L_m^s, [L_m^s])$ -heavy for any s, t.
- (4) $L_1^{\hat{t}}$ is not $(L_m^s, [L_m^s])$ -superheavy for any s, t.
- (5) $L_{\rm m}^{s'} \cup L_{\rm l}^{t}$ is $(L_{\rm m}^{s}, [L_{\rm m}^{s}])$ -superheavy for any s, s', t.

Later, we provide a generalization of (3) of Proposition 1.7 (See Proposition 10.3).

1.5. **Applications to non-displaceability.** As an application of Lagrangian spectral invariants to non-displaceability, we have the following theorem.

Theorem 1.8. Let L be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$. Let X, Y be a (L, a)-heavy subset, a (L, a)-superheavy subset of M, respectively. Then X is non-displaceable from Y.

As a corollary of Theorem 1.2, (3) of Proposition 1.7, Theorem 1.4 and Theorem 1.8, we obtain the following corollary.

Corollary 1.9. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let *X*, *Y* be *L*-stems. Then the subset $X \times L_1^t$ of $M \times T_R^2$ is non-displaceable from $Y \times L_m^s$ for any *s*, *t*.

Remark 1.10. Since both of $X \times L_m^s$ and $Y \times L_l^t$ are displaceable by symplectomorphisms which are isotopic to the identity, they are not superheavy in the original sense of Entov and Polterovich. Thus we cannot apply (iii) of Theorem 1.4 of [EP09] to Corollary 1.9.

Entov and Polterovich proved that any moment map has at least one fiber which is nondisplaceable from itself (Theorem 2.1 of [EP06], Theorem 6.1.8 of [PR]). We improve their result under an assumption that *M* admits a Lagrangian submanifold satisfying some conditions.

Corollary 1.11. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. For any moment map $\Phi: M \to \mathbb{R}^k$, there exists a point y_0 of $\Phi(L)$ such that $\Phi^{-1}(y_0)$ is non-displaceable from *L* and $\Phi^{-1}(y_0)$ itself.

Proof. Assume on the contrary that $\Phi^{-1}(y)$ is displaceable from *L* or $\Phi^{-1}(y)$ itself for any point *y* of $\Phi(L)$. Note that, for any point *y* of $\Phi(M) \setminus \Phi(L)$, $\Phi^{-1}(y)$ is disjoint from *L*, in particular displaceable from *L*. Hence, any fiber of Φ is an *L*-stem and thus, by Theorem 1.4 and Theorem 1.8, non-displaceable from *L*. This is a contradiction.

Remark 1.12. By an argument similar to Subsection 9.2 of [PR], we can regard Corollary 1.11 as a corollary of Theorem 1.18.

1.6. Non-degeneracy of spectral norms and the energy capacity inequality. In the process of proving (3) of Proposition 1.7, we prove the non-degeneracy of spectral norms.

Theorem 1.13. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Then for any idempotent **a** of the quantum homology $QH_*(L)$ and any Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ with compact support such that $\phi_H(L) \ne L$,

$$c^{L}(\mathbf{a}, H) + c^{L}(\mathbf{a}, \bar{H}) > 0.$$

On the case of the Oh-Schwarz spectral invariants, the non-degeneracy of spectral invariants is proved by Schwarz [Schw] when (M, ω) is symplectically ashperical and *H* is non-degenerate. After that, Oh [Oh05a] generalized it to the case in which (M, ω) is a general closed symplectic manifold and *H* is a general Hamiltonian function. Frauenfelder and Schlenk [FS] proved it when (M, ω) is a weakly convex symplectic manifold and *H* is a general Hamiltonian function.

Polterovich and Rosen gave another proof of Oh's result using a Poisson bracket inequality (Proposition 4.6.2 of [PR]). Similarly to Polterovich and Rosen, we use Poisson bracket inequality to prove Theorem 1.13. However, our Poisson bracket inequality is weaker than their one and we have to improve their argument.

Remark 1.14. The author tried to prove the non-degeneracy of spectral norms by an idea similar to Schwarz or Oh or Frauenfelder-Schlenk's one, but it failed because of some difficulties. Here, we explain one of them. In Schwarz and Oh's proof, it is important to prove the existence of a (broken) Floer trajectory between two orbits which represent [*M*] and [pt], where [pt] is the homology class representing a point (See Subsection 2.3 of [Schw] and Theorem 5.4 of [Oh05a]). In our case, $QH_*(L) \equiv H_*(L)$ does not hold in general, and thus we cannot take a homology class in $QH_*(L)$ representing [pt] in general.

As its application, we obtain a relative version of the energy capacity inequality which is announced by Lisi and Rieser (Theorem 4.1 of [LR]) and proved by Humilière, Leclercq and Seyfaddini for cotangent bundles (Lemma 7 of [HLS]).

Let (N, ω) be a symplectic manifold and *L* a closed subset of *N*.

Definition 1.15. An autonomous Hamiltonian function $H: N \to \mathbb{R}$ with compact support is *L-simple* if

(1): There exists a compact subset K of N such that

$$K \cap L \neq \emptyset$$
 and $H|_{N \setminus K} \equiv 0$,

(2): There exists an open subset W of N and a constant m(H) such that

$$W \cap L \neq \emptyset$$
 and $H|_W \equiv m(H)$,

(3): The only critical values of H are 0 and m(H).

A Hamiltonian function $H: N \to \mathbb{R}$ is *L*-slow if *H* is *L*-simple and all of its Hamiltonian chord of length at most 1 from *L* to *L* are constant *i.e.* for any $0 < t \le 1$ and any smooth path $z: [0, 1] \to N$ with $\dot{z}(t) = X_H^t(z(t))$ and $z(0), z(1) \in L, z(s) = z(0)$ holds for any $0 < s \le t$. Then, we define the Lisi-Rieser-type capacity $C_{LR}(N; L)$ by

$$C_{LR}(N; L) = \sup\{m(H); H: N \to \mathbb{R} \text{ is } L\text{-slow}\}.$$

Theorem 1.16. Let *L* be a Lagrangian submanifold of a closed symplectic manifold (M, ω) with $\omega(\pi_2(M, L)) = 0$, $N_L \ge 2$ and $QH_*(L) \ne 0$ (Note that $\omega(\pi_2(M, L)) = 0$ implies monotonicity of *L*). Then, for an open subset *U* of *M* with $L \cap U \ne \emptyset$,

$$C_{LR}(U;L\cap U) \le \bar{E}(U;L).$$

Remark 1.17. Our definition of simplicity follows [FGS] and it is slightly different from Lisi and Rieser's original one. In their original definition, "The only critical values of *H* are 0 and m(H)" is replaced by " $0 \le H(x) \le m(H)$ for all $x \in M$ ". Note that Theorem 1.16 does not imply Lisi and Rieser's original energy capacity inequality.

1.7. Application to Poisson bracket invariant. Let Q_N denote the cube $[0, 1]^N$. For a partition of unity $\vec{F} = \{F_1, \ldots, F_N\}$ on a symplectic manifold (M, ω) , we define the magnitude $\kappa_{cl}(\vec{F})$ of its Poisson noncommutativity by

$$\kappa_{cl}(\vec{F}) = \max_{x, y \in Q_N} ||\{\sum_i x_i F_i, \sum_j y_j F_j\}||.$$

For an open cover \mathcal{U} of M, we define the Poisson bracket invariant $pb(\mathcal{U}) = \inf \kappa_{cl}(\vec{F})$. Here the infimum is taken over all partition \vec{F} of the unity subordinated to \mathcal{U} .

For an open cover $\mathcal{U} = \{U_1, \dots, U_N\}$ of M, we define the displacement energy $\overline{E}(\mathcal{U}; L)$ of \mathcal{U} by $\overline{E}(\mathcal{U}; L) = \max_i \overline{E}(U_i; L)$.

The following theorem is a relative version of Theorem 3.1 of [P] and Theorem 9.2.2 of [PR] (see also Theorem 1.8 of [EPZ]).

Theorem 1.18. Let L be a monotone Lagrangian submanifold of a 2m-dimensional closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let $\mathcal{U} = \{U_1, \ldots, U_N\}$ be an open cover such that each U_i is displaceable from L or U_i itself. Then

$$\operatorname{pb}(\mathcal{U}) \cdot \overline{E}(\mathcal{U}; L) \ge (2N^2)^{-1}$$

In particular, $\overline{E}(\mathcal{U}; L) > 0$ and

$$\mathsf{pb}(\mathcal{U}) \ge (2N^2 \cdot \bar{E}(\mathcal{U};L))^{-1} > 0.$$

Polterovich used the Poisson bracket inequality efficiently to prove the original (abstract) version of Theorem 1.18, but our Poisson bracket inequality (Proposition 7.3) is too complicated to prove Theorem 1.18. For this difficulty, we avoid to use a partial symplectic quasi-state. Instead of this, we use a Lagrangian spectral invariant directly. The reason why Polterovich used a partial quasi-state is that it has the semi-homogeneity property. Instead of the semi-homogeneity property of a partial symplectic quasi-state, we use the Hamiltonian shift property of a Lagrangian spectral invariant.

We provide an application of Theorem 1.18. For an open cover $\mathcal{U} = \{U_i\}_i$ of a manifold M, let $\mathcal{U} \times \mathcal{U}$ denote the open cover $\{U_i \times U_i\}_{i,i}$ of $M \times M$

Example 1.19. Let $\{U_1, \ldots, U_M, V_1, \ldots, V_N\}$, $\{W_1, W_2\}$ be open cover of a circle $S^1 = \mathbb{R}/\mathbb{Z}$ such that

- $U_i \neq S^1$, $V_j \neq S^1$ and $W_k \neq S^1$ for any *i*, *j* and *k*,
- $0 \in U_i$ for any i = 1, ..., M and $0 \notin V_j$ for any j = 1, ..., N.

We define a cover \mathcal{U} of $T_1^2 = S^1 \times S^1$ by

$$\mathcal{U} = \{U_i \times W_k, V_j \times S^{\perp}\}_{i,j,k}.$$

Then, any element of \mathcal{U} is displaceable from $\{0\} \times S^1$. Thus, by Theorem 1.18, $pb(\mathcal{U})$ is positive. Similarly, we can prove that $pb(\mathcal{U} \times \mathcal{U})$ is also positive. On the other hand, we can easily confirm that $pb(\mathcal{U}') = 0$ where

$$\mathcal{U}' = \{U_i \times S^1, V_j \times S^1\}_{i,j}.$$

Remark 1.20. Kaoru Ono pointed out that we can prove positivity of $pb(\mathcal{U}) > 0$ by the original Polterovich's theorem when $\{U_1, \ldots, U_M\}$ is not an open cover of a circle $S^1 = \mathbb{R}/\mathbb{Z}$.

We note that the assumption $QH_*(L) \neq 0$ in Theorem 1.18 is essential. $QH_*(L) \neq 0$ implies non-displaceability of *L*. We provide the following proposition if *L* is displaceable.

Proposition 1.21. Let X be a closed subset of a closed symplectic manifold (M, ω) . Assume that X is displaceable from X itself. Then, there exists an open cover $\mathcal{U} = \{U_1, \ldots, U_N\}$ of M such that any U_i is displaceable from X and $pb(\mathcal{U}) = 0$.

1.8. Hamiltonian chords between two disjoint subsets. For subsets Y_0, Y_1 of an open symplectic manifold (N, ω) . For a Hamiltonian function $H: S^1 \times N \to \mathbb{R}$ with compact support and a homotopy class $\alpha \in \pi_1(M, Y_0 \cup Y_1)$, We define a set $Ch(H; Y_0, Y_1, \alpha)$ of Hamiltonian chords of H from Y_0 to Y_1 in α by

Ch(*H*; *Y*₀, *Y*₁,
$$\alpha$$
) = {*z*: [0, 1] → *N*; $\dot{z}(t) = X_H^t(z(t)), z(0) \in Y_0 \text{ and } z(1) \in Y_1, [z] = \alpha$ }.

For subsets Y_0 , Y_1 and Z of an open symplectic manifold (N, ω) and a homotopy class α of $\pi_1(N, Y_0 \cup Y_1)$, we define the relative symplectic capacity $C_{BEP}(N, Y_0, Y_1, Z, \alpha)$ by

$$C_{BEP}(N, Y_0, Y_1, Z, \alpha) = \inf\{K > 0; \forall H \in \mathcal{H}_K(N, Z), \mathsf{Ch}(H; Y_0, Y_1, \alpha) \neq \emptyset\},\$$

where

$$\mathcal{H}_{K}(N,Z) = \{ H \in C_{c}^{\infty}(S^{1} \times N); \inf_{S^{1} \times Z} H \ge K \}.$$

Remark 1.22. Buhovsky, Entov and Polterovich studied existence problem of Hamiltonian chords between two disjoint subsets under some robust restriction on the C^0 -profile of the Hamiltonian function ([BuEP]). They made a method to find a Hamiltonian chord whose length is equal to or smaller than 1. On the other hand, an upper bound of C_{BEP} provides a Hamiltonian chord whose length is equal to 1.

For $R = (R_1, ..., R_n) \in (\mathbb{R}_{>0})^n$, let $\mathbb{A}(R)$ denote an annulus $(0, R_1) \times (0, R_2) \times (0, R_n) \times (\mathbb{R}/\mathbb{Z}) \times \cdots \times (\mathbb{R}/\mathbb{Z})$ with the coordinates $(p_1, ..., p_n, q_1, ..., q_n)$. We fix a symplectic form $dp_1 \wedge dq_1 + \cdots + dp_n \wedge dq_n$ on $\mathbb{A}(R)$. For $s = (s_1, ..., s_n) \in (\mathbb{R}/\mathbb{Z})^n$ for any *i*, define $\mathbb{A}(R)_s$ by

$$\mathbb{A}(R)_s = \{(p,q) \in \mathbb{A}(R); q = s\}.$$

For $R = (R_1, ..., R_n) \in (\mathbb{R}_{>0})^n$, $s = (s_1, ..., s_n) \in \mathbb{R}^n$ and subsets *Y*, *Z* of *M*, we define a relative symplectic capacity C(M, Y, Z, R, s) by

$$C(M, Y, Z, R, s) = C_{BEP}(M \times \mathbb{A}(R), Y \times \mathbb{A}(R)_0, Y \times \mathbb{A}(R)_s, Z \times \mathbb{A}(0), \alpha_s).$$

Here α_s is the homotopy class of $\pi_1(M \times \mathbb{A}(R), Y \times \mathbb{A}(R)_0 \cup Y \times \mathbb{A}(R)_s)$ represented by the path $\gamma_s: [0, 1] \to M \times \mathbb{A}(R)$ defined by $t \mapsto (p_0, ([s_1t], \dots, [s_nt]), (0, \dots, 0))$ where p_0 is the fixed base point of *Y*.

Theorem 1.23. Let *L* be a monotone Lagrangian submanifold of a 2*m*-dimensional closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let *Z* be an (L, a)-heavy subset of *M* for some idempotent *a*. Then for any $R = (R_1, \ldots, R_n) \in (\mathbb{R}_{>0})^n$ and any $s = (s_1, \ldots, s_n) \in \mathbb{R}^n$,

$$C(M, L, Z, R, s) \le 2 \sum_{i=1}^{n} R_i \cdot |s_i| + \lambda(m+n).$$

In [K16], the author used the fact "the meridian curve is heavy in the sense of Entov and Polterovich" ([EP09]). We use (3) of Proposition 1.7 instead of that fact.

In contrast, we prove the following proposition when Z is displaceable from L (recall that (L, a)-heavy subset is non-displaceable from L).

Proposition 1.24. Let Z be a displaceable compact of a closed symplectic manifold (M, ω) . Assume $R_k \cdot s_k > E(Z; L)$ for some k, where E(Z; L) is the displacement energy. Then

$$C(M, L, Z, R, s) = +\infty.$$

The proof is also a Lagrangian analogue of [K16].

Technical remark. In the case of Oh-Schwarz spectral invariants, (abstractly) displaceable open subsets have some special property. The source of these properties is the bounded spectrum condition for (abstractly) displaceable subsets (Proposition 4.4),

In our Lagrangian spectral invariants, we prove the bounded spectrum condition for relatively displaceable subsets (Proposition 4.2) and it enables us to make relative versions of some known results.

However, it is not sufficient to replace the bounded spectrum condition for (abstractly) displaceable subsets with the one for relatively displaceable subsets. There is a large technical difference between Oh-Schwarz spectral invariants and Lagrangian spectral invariants. Oh-Schwarz spectral invariants are invariant under the action of Hamiltonian diffeomorphisms, but the same property does not hold for Lagrangian ones. This difference makes some technical difficulties.

In the author's opinion, the easiest way to understand this difficulty is comparing our Poisson bracket inequality (Proposition 7.3) with the original one (Theorem 3.1 of [P] and Proposition 4.6.1 of [PR]). For more precise explanation, see Remark 7.4.

Overview of the present paper. In Section 2, we prepare some notions and define Lagrangian spectral invariants. In Section 3, we prove Corollary 1.5.

In Section 4, we define the bounded spectrum condition for open subsets of a symplectic manifold. We prove abstractly or relatively displaceable open subsets satisfy the bounded spectrum condition (Propositions 4.2, 4.3). In Section 5, we prove Theorem 1.18 and Proposition 1.21.

In Section 6, we provide some properties of asymptotic Lagrangian spectral invariants and prove Theorem 1.8.. In Section 7, we formulate a Lagrangian version of Poisson bracket inequality (Proposition 7.3) which is necessary for proving some results in this paper.

In Section 8, we prove some properties of asymptotic Lagrangian spectral invariants and Theorem 1.4. In Section 9, we prove Theorem 1.13. In Section 10, we prove a relative version of the energy capacity inequality (Theorem 1.16) by Theorem 1.13.

In this section, we also prove Proposition 1.7. To prove heaviness of a longitude curve with respect to a meridian curve ((3) of Proposition 1.7), we use Theorem 1.13.

In Section 11, we prove Theorem 1.23 by (3) of Theorem 1.7. In Section 12, we prove Proposition 1.24.

Acknowledgment. For the author, this paper is the first work where he uses the Lagrangian Floer theory. The author thanks Professor Yong-Geun Oh and Professor Kaoru Ono for replying some elementary questions on the Lagrangian Floer theory. He also thanks Yoosik Kim, Weonmo Lee, Ryuma Orita, Professor Leonid Polterovich, Fumihiko Sanda and Frol Zapolsky for faithful discussions and comments. Especialy, Ryuma Orita checked this paper carefully and pointed out a lot of careless mistakes.

He is supported by IBS-R003-D1 and JP18J00765.

2. Preliminary

In the present paper, we follow the notion of [LZ] and we consider only the case in which the coefficient on Floer theory is $\mathbb{Z}/2\mathbb{Z}$.

For a subset *X* of *M*, we define

 $\Omega_0(X) = \{z: [0, 1] \to M, \text{ smooth map}; z(0), z(1) \in X, [z] = 0_X\},\$

where 0_X is the trivial element of $\pi_1(M, X)$.

To consider the Lagrangian Floer theory, we consider the following covering space $\tilde{\Omega}_0(L)$ of $\Omega_0(L)$. Note that our explanation is a rough sketch and omit the definitions of some words. See [LZ] and [Z] for them and details.

Let $z \in \Omega_0(L)$. Two cappings $\hat{z}, \hat{z}': D^2 \setminus \{1\} \to M$ of z are *equivalent* if the preglued map $\hat{z} \not\models (-\hat{z}')$ defines the trivial element in $\pi_2(M, L)$, where $-\hat{z}'$ is the opposite map of \hat{z}' . Two pairs $(z, \hat{z}), (z', \hat{z}')$ are *equivalent* if z = z' and \hat{z}, \hat{z}' are equivalent as cappings of z = z'. Let $[z, \hat{z}]$ denote the class represented by (z, \hat{z}) and define

$$\hat{\Omega}_0(L) = \{[z, \hat{z}]; z \in \Omega_0(L), \hat{z} \text{ is a capping of } z\}.$$

For a Hamiltonian function $H: S^1 \times M \to \mathbb{R}$, the action functional $\mathcal{R}^L_H: \tilde{\Omega}_0(L) \to \mathbb{R}$ is defined by

$$\mathcal{A}_H^L([z,\hat{z}]) = \int_0^1 H_t(z(t))dt - \int_{D^2} \hat{z}^* \omega.$$

For a connected subset X of M, let Ch(H; X) denote $Ch(H; X, X, 0_X)$. We define its covering $\tilde{Ch}(H; X)$ by

$$\tilde{\mathsf{Ch}}(H;X) = \{[z,\hat{z}] \in \tilde{\Omega}_0(X); z \in \mathsf{Ch}(H;X)\}.$$

Then we regard $\tilde{Ch}(H; L)$ as the set of critical points of \mathcal{A}_{H}^{L} . We define its spectrum Spec(H; L) as the set of critical values of \mathcal{A}_{H}^{L} *i.e.*

$$\operatorname{Spec}(H;L) = \{\mathcal{R}_{H}^{L}([z,\hat{z}]); [z,\hat{z}] \in \widetilde{\operatorname{Ch}}(H;L)\}.$$

We define the *non-degeneracy* of Hamiltonian functions as follows:

Definition 2.1. A Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ is called *non-degenerate* if for any element *z* of Ch(*H*; *L*), the induced linear map $(d\phi_H^1)_{z(0)}: T_{z(0)}M \to T_{z(1)}M$ maps the linear subspace $T_{z(0)}L$ to a linear subspace transverse to $T_{z(1)}L$.

When *H* is non-degenerate, the Floer chain complex $CF_*(H; L)$ is generated by Ch(H; L) as a module over $\mathbb{Z}/2\mathbb{Z}$. The complex $CF_*(H; L)$ is graded by the Conley-Zehnder index ind_{CZ} . Note that $\operatorname{ind}_{CZ}([z, \hat{z} \natural A]) = \operatorname{ind}_{CZ}([z, \hat{z}]) - \mu(A)$ for any map $A \in \pi_2(M)$ in our convention. Let $F: M \to \mathbb{R}$ be a function which is the pullback of a Morse function $\tilde{F}: L \to \mathbb{R}$ on some Weinstein neighborhood of *L* and *x* a critical point of \tilde{F} such that $d\tilde{F}$ is C^1 -small near *x*. Then $\operatorname{ind}_{Morse}(x) = \operatorname{ind}_{CZ}([x, c_x])$, where c_x is a trivial capping disk and $\operatorname{ind}_{Morse}$ is the Morse index. We formally obtain the boundary map of this complex by counting isolated negative gradient flow lines of \mathcal{A}^L_H and we define its homology group $HF_*(H; L)$ which is called *the Lagrangian Floer homology on contractible trajectories* of *H*.

Oh [Oh94], Biran and Cornea [BC] (see also [LZ]) defined the quantum homology $QH_*(L)$ of a monotone Lagrangian submanifold L with $N_L \ge 2$ and proved that there exists a natural isomorphism $\Phi: QH_*(L) \to HF_*(H; L)$. We call this isomorphism the PSS isomorphism ([PSS96]).

Biran and Cornea defined the quantum product * on $QH_*(L)$. $QH_*(L)$ has the fundamental class [L] which is the unit with respect to *. If $QH_*(L) \neq 0$, then $[L] \neq 0$ since [L] is the unit of $QH_*(L)$.

Given an element $A = \sum_{i} a_i [z_i, \hat{z}_i]$ of $CF_*(H; L)$, we define the action level $l_H(A)$ of A by

$$l_H(A) = \max\{\mathcal{H}_H^L([z_i, \hat{z}_i]); a_i \neq 0\}.$$

For a non-zero element a of $QH_*(L)$, we define the spectral invariant associated with H and a by

$$c^{L}(\mathbf{a}, H) = \inf\{l_{H}(A); [A] = \Phi(\mathbf{a})\}.$$

It is known that $c^{L}(a, H)$ is a finite number for any non-degenerate H and any non-trivial $a \in QH_{*}(L)$.

Let $H: S^1 \times M \to \mathbb{R}$ be a continuous function, a a non-zero element of $QH_*(L)$. We define the spectral invariant $c^L(\mathbf{a}, H)$ associated with H by

$$c^{L}(\mathbf{a},H) = \lim_{n \to \infty} c^{L}(\mathbf{a},H_{n}),$$

where $\{H_n\}_n$ is a sequence of non-degenerate Hamiltonian functions converging to H in L^{∞} -norm. It is known that this limit exists and does not depend on the choice of $\{H_n\}_n$. For a more precise argument, see Subsection 3.3 of [LZ].

In the present paper, the following proposition of Leclercq and Zapolsky plays a very important role.

Proposition 2.2 (Theorem 3 and Proposition 4 of [LZ]). The spectral invariant has the following properties for any non-trivial elements a, b of $QH_*(L)$.

(1)*Lipschitz continuity:* For any continuous functions $F, G: S^1 \times M \to \mathbb{R}$,

$$\int_{0}^{1} \min_{M} (F_{t} - G_{t}) dt \le c^{L}(\mathbf{a}, F) - c^{L}(\mathbf{a}, G) \le \int_{0}^{1} \max_{M} (F_{t} - G_{t}) dt,$$

- (2)*Homotopy invariance:* Assume that normalized smooth functions $F, G: S^1 \times M \rightarrow \mathbb{R}$ satisfy $\phi_F^1 = \phi_G^1$ and that their Hamiltonian isotopies $\{\phi_F^t\}$ and $\{\phi_G^t\}$ are homotopic relative to endpoints. Then $c^L(\mathbf{a}, F) = c^L(\mathbf{a}, G)$,
- (3)*Triangle inequality:* $c^{L}(a * b, F \natural G) \leq c^{L}(a, F) + c^{L}(b, G)$ for any continuous functions $F, G: S^{1} \times M \to \mathbb{R}$,
- (4)*Hamiltonian shift:* Let $\rho: S^1 \to \mathbb{R}$ be a continuous function. Then, for any continuous function $H: S^1 \times M \to \mathbb{R}$,

$$c^{L}(\mathbf{a}, H + \rho) = c^{L}(\mathbf{a}, H) + \int_{0}^{1} \rho(t) dt,$$

where $H + \rho \colon S^1 \times M \to \mathbb{R}$ is a Hamiltonian function defined by $(H + \rho)(t, x) = H(t, x) + \rho(t)$,

(5): $c^{L}([L], 0) = 0$,

(6) *Spectrality:* For any smooth function $H: S^1 \times M \to \mathbb{R}$,

$$c^{L}(\mathbf{a}, H) \in \operatorname{Spec}(H; L).$$

For a general Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ (*H* can be degenerate), we define the spectral invariant $c^L(a, H)$ by the Lipschitz continuity. Then the spectral invariant defined for general Hamiltonian functions also satisfies the properties in Proposition 2.2.

In this paper, we also consider spectral invariant defined in terms of the Hamiltnian Floer theory (see [Schw], [Oh05b] and [Oh06]). For a closed symplectic manifold (M, ω) , we have the quantum homology $QH_*(M)$ (For simplicity, we consider the coefficient $\mathbb{Z}/2\mathbb{Z}$). For a nontrivial element a of the quantum homology $QH_*(M)$ and a continuous function $H: S^1 \times M \to \mathbb{R}$, let c(a, H) denote the Oh-Schwarz (Hamiltonian) spectral invariant associated with a and H. There are some conventions of the Oh-Schwarz spectral invariants, We use the same convention

as [EP09]. In this paper, we use the same convention as [EP09] and [LZ] and compare our spectral invariants with the Oh-Schwarz spectral invariants.

The quantum homology $QH_*(M)$ acts on $QH_*(L)$ via the quantum module action ([BC], [Z]),

•:
$$QH_*(M) \otimes QH_*(L) \rightarrow QH_*(L)$$
.

Related to this action, we have the following convenient inequality.

Proposition 2.3 (Proposition 5 of [LZ]). *Let* e *be an element of* $QH_*(M)$ *and* e' *an element of* $QH_*(L)$.

For any Hamiltonian functions $F, G: S^1 \times M \to \mathbb{R}$ *,*

$$c^{L}(\mathbf{e} \bullet \mathbf{e}', F \natural G) \le c^{L}(\mathbf{e}', F) + c(\mathbf{e}, G).$$

3. Proof of Corollary 1.5

We provide the following theorem which generalizes Corollary 1.5.

Theorem 3.1. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let e be an idempotent of $QH_*(M)$ and e' an idempotent of $QH_*(L)$. Assume that there exists an element e'' of $QH_*(L)$ such that $e \bullet e'' = e'$. Then

- Any e-superheavy subset X is (L, e')-superheavy,
- Any (L, e')-heavy subset Y is e-heavy.

Since $[M] \bullet e' = e'$, Corollary 1.5 follows from Theorem 3.1. To prove Theorem 3.1, we prove the following lemma.

Lemma 3.2. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let e be an idempotent of $QH_*(M)$ and e' an idempotent of $QH_*(L)$. Assume that there exists an element e'' of $QH_*(L)$ such that $e \bullet e'' = e'$. Then

$$\zeta_{\mathsf{e}'}^{L}(F) \leq \zeta_{\mathsf{e}}(F).$$

Proof. By Proposition 2.3, for any integer *k*,

$$c^{L}(e', kF) \le c(e, kF) + c^{L}(e'', 0).$$

Thus, by diving by k and take the limit, we complete the proof.

By Definition 1.1, Theorem 3.1 immediately follows from Lemma 3.2. The idea of comparing Lagrangian spectral invariant and Hamiltonian spectral invariant using the module structure • comes from Section 6 of [LZ].

4. The bounded spectrum condition and Lagrangian version of Ostrover's trick

The following notion was introduced by the author.

Definition 4.1 ([K14]). Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let a be an idempotent of $QH_*(L)$.

An open subset U of M satisfies the bounded spectrum condition with respect to a if there is a positive number K such that

$$c^L(\mathsf{a}, F) < K,$$

for any Hamiltonian function $F: S^1 \times U \to \mathbb{R}$ with compact support.

A subset X of M satisfies the bounded spectrum condition with respect to a if there is an open neighborhood U of X such that U satisfies the bounded spectrum condition with respect to a.

In this section, we give some examples of open subsets with the bounded spectrum condition. By considering a Lagrangian analogue of Octrover's argument [Oc], we prove the following

By considering a Lagrangian analogue of Ostrover's argument [Os], we prove the following proposition. We use this proposition for several times in this paper.

Proposition 4.2. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let *U* be an open subset of *M* and *H* a Hamiltonian function satisfying $\phi_H(U) \cap L = \emptyset$. Then for any idempotent **a** of the quantum homology $QH_*(L)$ and any Hamiltonian function $F: S^1 \times U \to \mathbb{R}$ with compact support on $S^1 \times U$,

$$c^{L}(\mathbf{a}, F) \leq c^{L}(\mathbf{a}, H) + c^{L}(\mathbf{a}, \overline{H}).$$

Proof. We give an upper bound of the spectral invariant associated with F using the concatenation with ϕ_{H}^{t} .

We can choose a smooth function $\chi: [0, \frac{1}{2}] \to [0, 1]$ satisfying the following conditions.

- $\frac{\partial \chi}{\partial t}(t) \ge 0$ for any $t \in [0, \frac{1}{2}]$, and
- $\chi(t) = 0$ for any $t \in [0, \frac{1}{5}]$ and $\chi(t) = 1$ for any $t \in [\frac{2}{5}, \frac{1}{2}]$.

For a real number *s* with $s \in [0, 1]$, we define the new Hamiltonian function $K^s : S^1 \times M \to \mathbb{R}$ as follows:

$$K^{s}(t,x) = \begin{cases} \frac{\partial \chi}{\partial t}(t)\bar{H}(\chi(t),x) & \text{when } t \in [0,\frac{1}{2}], \\ s\frac{\partial \chi}{\partial t}(t-\frac{1}{2})F(s\chi(t-\frac{1}{2}),x) & \text{when } t \in [\frac{1}{2},1]. \end{cases}$$

Since $\frac{\partial \chi}{\partial t} = 0$ on neighborhoods of t = 0 and $t = \frac{1}{2}$, K^s is a smooth Hamiltonian function.

We claim $\text{Spec}(K^s; L) \subset \text{Spec}(\bar{H}; L)$ for a real number *s* with $s \in [0, 1]$. Let $F^s: S^1 \times \hat{W} \to \mathbb{R}$ denote the Hamiltonian function defined by

$$F^{s}(t,x) = s\frac{\partial\chi}{\partial t}(\frac{t}{2})F(s\chi(\frac{t}{2}),x).$$

Let $[z, \hat{z}]$ be an element of $\tilde{Ch}(\bar{H}; L)$ and define $x \in M$ by x = z(0). If $x \in \phi_H(U)$, then $\gamma_{K^s}^x \notin \Omega_0(L)$ and in particular, $\gamma_{K^s}^x \notin Ch(\bar{H}; L)$ since $\phi_H(U) \cap L = \emptyset$. If $x \notin \phi_H(U)$, then $\phi_{\bar{H}}(x) \notin U$. Thus $\gamma_{K^s}^x$ is equal to $\gamma_{\bar{H}}^x$ up to parameter change and $\int_0^1 H(t, \gamma_{\bar{H}}^x(t))dt = \int_0^1 K^s(t, \gamma_{K^s}^x(t))dt$. Therefore we see that there exists a natural inclusion map $\iota \colon \tilde{Ch}(K^s; L) \to \tilde{Ch}(\bar{H}; L)$ which preserves values of the action functional, and hence $\operatorname{Spec}(K^s; L) \subset \operatorname{Spec}(\bar{H}; L)$. implies $c^L(a, K^s) \in \operatorname{Spec}(\bar{H}; L)$.

By the Lipschitz continuity for spectral invariants (Proposition 2.2 (1)), $c^{L}(a, K^{s})$ depends continuously on *s*. Since Spec(\overline{H} ; *L*) is a measure-zero set (Lemma 30 of [LZ]), $c^{L}(a, K^{s})$ is a constant function of *s*. The homotopy invariance for spectral invariants (Proposition 2.2 (2)) implies

$$c^{L}(\mathbf{a}, K^{0}) = c^{L}(\mathbf{a}, \bar{H}).$$

Hence for any $s \in [0, 1]$,

$$c^{L}(\mathbf{a}, K^{s}) = c^{L}(\mathbf{a}, \bar{H}).$$

Then $c^{L}(a, F)$ is estimated as follows.

$$c^{L}(\mathbf{a}, F) \le c^{L}(\mathbf{a}, K^{1}) + c^{L}(\mathbf{a}, H)$$
$$= c^{L}(\mathbf{a}, \bar{H}) + c^{L}(\mathbf{a}, H).$$

For (abstractly) displaceable subsets, we have the following proposition.

 \Box

Proposition 4.3. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$.

Let U be an open subset of M and H a Hamiltonian function satisfying $\phi_H(U) \cap \overline{U} = \emptyset$. Then for any idempotent **a** of the quantum homology $QH_*(L)$ and any Hamiltonian function $F: S^1 \times U \to \mathbb{R}$ with compact support on $S^1 \times U$,

$$c^{L}(\mathbf{a}, F) \le c([M], H) + c([M], \bar{H}) + c^{L}(\mathbf{a}, 0).$$

To prove Proposition 4.3, we recall the original Ostrover's trick.

Proposition 4.4 (Proposition 3.1 of [U10], see also Proposition 2.1 of [FGS]). Let (M, ω) be a closed symplectic manifold. Let U be an open subset of M and H a Hamiltonian function satisfying $\phi_H(U) \cap \overline{U} = \emptyset$. Then for any idempotent a of the quantum homology $QH_*(M, \omega)$ and any Hamiltonian function $F: S^1 \times U \to \mathbb{R}$ with compact support on $S^1 \times U$,

$$c(\mathbf{a}, F) \le c(\mathbf{a}, H) + c(\mathbf{a}, \bar{H}).$$

Proof of Proposition 4.3. By Proposition 2.3,

$$c^{L}(\mathbf{a}, F) \le c([M], F) + c^{L}(\mathbf{a}, 0).$$

By Proposition 4.4,

$$c([M], F) \le c([M], H) + c([M], \bar{H}).$$

Thus,

$$c^{L}(\mathbf{a}, F) \le c([M], H) + c([M], \bar{H}) + c^{L}(\mathbf{a}, 0).$$

By the Lipschitz continuity of Lagrangian spectral invariant, the following corollary immediately follows from Proposition 4.2, 4.3 and (1), (5) of Proposition 2.2.

Corollary 4.5. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Let *U* be an open subset of *M* which is displaceable from *L* or *U* itself. Then for any Hamiltonian function $F: S^1 \times U \rightarrow \mathbb{R}$ with compact support on $S^1 \times U$,

 $c^{L}([L], F) \leq \overline{E}(U; L).$

5. Application to Poisson bracket invariant

Lemma 5.1. For any Hamiltonian functions $F, G: S^1 \times M \to \mathbb{R}$ and a non-trivial element a of $QH_*(L)$,

$$|c^{L}(\mathbf{a}, F + G) - c^{L}(\mathbf{a}, F \natural G)| \le \frac{1}{2} ||\{F, G\}||.$$

Proof. For any t > 0 and $x \in M$,

$$G(\phi_F^t x) - G(x) = \int_0^t \frac{d}{ds} G(\phi_F^s x) ds = \int_0^t \{G, F\}(\phi_F^s x) ds$$

Thus, for any *t*,

$$||(F+G) - (F \natural G)_t|| = ||G - G \circ (\phi_F^t)^{-1}|| = ||G \circ \phi_F^t - G|| \le t ||\{F, G\}||.$$

By the Lipshitz continuity,

$$c^{L}(\mathbf{a}, F + G) - c^{L}(\mathbf{a}, F \natural G)$$

$$\leq \int_{0}^{1} ||(F + G) - (F \natural G)_{t}||dt$$

$$\leq \int_{0}^{1} t ||\{F, G\}||dt$$

$$= \frac{1}{2} ||\{F, G\}||.$$

Proof of Theorem 1.18. Let \vec{F} be a partition of the unity subordinated to \mathcal{U} .

Fix a positive number R. Define functions $G_k: M \to \mathbb{R}$ (k = 0, ..., N), $H_k: \mathbb{R} \to \mathbb{R}$ (k = 1, ..., N) by $G_k(x) = \sum_{i=1}^k F_i(x)$ and $H_k(R) = c^L([L], RG_k) - c^L([L], RG_{k-1})$.

Then, by the triangle inequality,

$$c^{L}([L], R\bar{F}_{k}) \leq c^{L}([L], (RG_{k-1})\natural(RF_{k})) - c^{L}([L], RG_{k-1}) \leq c^{L}([L], RF_{k}).$$

Since $\text{Supp}(F_k) \subset U_k$ for any k = 1, ..., N, by Corollary 4.5,

$$-\bar{E}(\mathcal{U};L) \leq c^{L}([L],(RG_{k-1})\natural(RF_{k})) - c^{L}([L],RG_{k-1}) \leq \bar{E}(\mathcal{U};L)$$

Hence $|c^L([L], (RG_{k-1})\natural(RF_k)) - c^L([L], RG_{k-1})| \leq \overline{E}(\mathcal{U}; L).$

By Lemma 5.1 and the definition of $\kappa_{cl}(\vec{F})$,

$$\begin{aligned} |c^{L}([L], RG_{k-1} + RF_{k}) - c^{L}([L], (RG_{k-1})\natural(RF_{k}))| \\ &\leq \frac{1}{2} ||\{RG_{k-1}, RF_{k}\}|| \\ &= \frac{R^{2}}{2} ||\{G_{k-1}, F_{k}\}|| \\ &\leq \frac{R^{2}}{2} \kappa_{cl}(\vec{F}). \end{aligned}$$

Thus, by the triangle inequality,

$$\begin{aligned} |H_{k}(R)| &= |c^{L}([L], RG_{k}) - c^{L}([L], RG_{k-1})| \\ &\leq |c^{L}([L], RG_{k-1} + RF_{k}) - c^{L}([L], (RG_{k-1})\natural(RF_{k}))| \\ &+ |c^{L}([L], (RG_{k-1})\natural(RF_{k})) - c^{L}([L], RG_{k-1})| \\ &\leq \frac{R^{2}}{2}\kappa_{cl}(\vec{F}) + \bar{E}(\mathcal{U}; L). \end{aligned}$$

By the triangle inequality,

$$c([L], R) - c([L], 0) = H_N(R) + H_{N-1}(R) + \dots + H_1(R) \leq |H_N(R)| + |H_{N-1}(R)| + \dots + |H_1(R)| \leq N(\frac{R^2}{2}\kappa_{cl}(\vec{F}) + \bar{E}(\mathcal{U}; L)).$$

By the Hamiltonian shift property ((4) of Proposition 2.2), c([L], R) - c([L], 0) = R. Thus

$$1 \le N(\frac{R}{2}\kappa_{cl}(\vec{F}) + R^{-1} \cdot \bar{E}(\mathcal{U};L))$$

The right-hand side is minimized by $R = (2\bar{E}(\mathcal{U};L))^{\frac{1}{2}}(\kappa_{cl}(\vec{F}))^{-\frac{1}{2}}$ and hence

 $1 \leq N(2\bar{E}(\mathcal{U};L) \cdot \kappa_{cl}(\vec{F}))^{\frac{1}{2}}.$

Therefore

$$\kappa_{cl}(\vec{F}) \cdot \bar{E}(\mathcal{U};L) \ge (2N^2)^{-1}$$

By taking the infimum over all partition \vec{F} of unity subordinated to \mathcal{U} ,

$$\operatorname{pb}(\mathcal{U}) \cdot \overline{E}(\mathcal{U}; L) \ge (2N^2)^{-1}.$$

г	-	-	٦
L			I

Proof of Proposition 1.21. Let F be a smooth function such that $F^{-1}(0) = X$. Since $F^{-1}(0) = X$ is displaceable from X, there is a positive number ϵ such that $F^{-1}((-2\epsilon, 2\epsilon))$ is also displaceable from X. Define open subsets U_1, U_2, U_3 by $U_1 = F^{-1}((-\infty, -\epsilon))$ $U_2 = F^{-1}((-2\epsilon, 2\epsilon))$ $U_3 = F^{-1}((\epsilon, +\infty))$ and set $\mathcal{U} = \{U_1, U_2, U_3\}$. By the definition, U_2 is displaceable from X. U_1 and U_3 are disjoint from X and in particular, displaceable from X. Since $U_1 \cap U_2 \cap U_3 = \emptyset$, $pb(\mathcal{U}) = 0$. Thus this \mathcal{U} satisfies all of the conditions.

6. PROPERTIES OF PARTIAL QUASI-STATE

Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$.

 ζ_{a}^{L} satisfies the following properties.

Proposition 6.1. (1)*Partial quasi-additivity:* For Hamiltonian functions $F, G: M \to \mathbb{R}$ with $\{F, G\} = 0$ and Supp(G) is displaceable from Supp(G) itself, $\zeta_a^L(F + G) = \zeta_a^L(F)$.

- (1')Lagrangian partial quasi-additivity: For Hamiltonian functions $F, G: M \to \mathbb{R}$ with $\{F, G\} = 0$ and $\operatorname{Supp}(G)$ is displaceable from $L, \zeta_{a}^{L}(F+G) = \zeta_{a}^{L}(F)$.
- (2)*Normalization*: $\zeta_{a}^{L}(1) = 1$.
- (3)Semi-homogeneity: For any autonomous Hamiltonian function $F: M \to \mathbb{R}$ and any positive number s, $\zeta_a^L(sF) = s\zeta_a^L(F)$.

We prove (1) of Proposition 6.1 and (1') in Section 8. Th proofs of (2) and (3) of Proposition 6.1 are quite similar to the corresponding statements on Oh-Schwarz spectral invariants (Theorem 3.6 of [EP09]) and thus we omit the proofs.

To prove Theorem 1.8, we prepare some properties of ζ_a^L .

Proposition 6.2. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$ and a a non-trivial idempotent of $QH_*(L)$. Let *X*, *Y* be a (L, \mathbf{a}) -heavy subset, a (L, \mathbf{a}) -superheavy subset of *M*, respectively. Then $X \cap Y \ne \emptyset$.

The proof of Proposition 6.2 is quite similar to the proof of (iii) of Theorem 1.4 in [EP09] and thus we omit the proof.

Lemma 6.3. For any continuous function $F: S^1 \times M \to \mathbb{R}$ and a Hamiltonian diffeomorphism ψ ,

$$\zeta_{\mathsf{a}}^{L}(F \circ \psi) = \zeta_{\mathsf{a}}^{L}(F),$$

where $F \circ \psi$ is a function defined by

$$F \circ \psi(t, x) = F(t, \psi(x)).$$

The proof of Proposition 6.3 is quite similar to the proof of 2 of Theorem 1.8 in [MVZ] and thus we omit the proof.

By the definition of heaviness and superheaviness, Lemma 6.3 implies the following corollary.

Corollary 6.4. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$, a a non-trivial idempotent of $QH_*(L)$ and *X* a (L, a)-heavy subset. Then $\psi(X)$ is a (L, a)-heavy subset for any Hamiltonian diffeomorphism ψ .

Proof of Theorem 1.8. To prove by contradiction, suppose that there exists a Hamiltonian diffeomorphism ψ such that

$$\psi(X) \cap Y = \emptyset.$$

By Corollary 6.4, $\psi(X)$ is (L, a)-heavy. Since Y is (L, a)-superheavy, this contradicts Proposition 6.2.

7. POISSON BRACKET INEQUALITY ON LAGRANGIAN SPECTRAL INVARIANTS

A large difference between our work and Entov and Polterovich's original argument is that Lagrangian spectral invariants are not invariant under the action of Hamiltonian diffeomorphisms. We need a more complicated argument than the Entov and Polterovich's original one.

In this section, we use spectral invariants defined on the universal covering of the group of Hamiltonian diffeomorphisms. To define them, we prepare some notions.

Let (M, ω) be a symplectic manifold. Let $\operatorname{Ham}(M, \omega)$ be the universal covering of the group $\operatorname{Ham}(M, \omega)$ of Hamiltonian diffeomorphisms. Note that an element of $\operatorname{Ham}(M, \omega)$ is represented by a path in $\operatorname{Ham}(M, \omega)$ starting from the identity. For a Hamiltonian function $H: S^1 \times M \to \mathbb{R}$ with compact support, let $\tilde{\phi}_H$ denote an element of $\operatorname{Ham}(M, \omega)$ represented by the path $\{\phi_H^t\}_{t \in [0,1]}$ in $\operatorname{Ham}(M, \omega)$.

For a non-trivial element a of $QH_*(L)$ and an element $\tilde{\phi}$ of $Ham(M, \omega)$, we define its spectral invariant $c^L(\mathbf{a}, \tilde{\phi})$ by

$$c^{L}(\mathbf{a}, \tilde{\phi}) = c^{L}(\mathbf{a}, H),$$

where $H: S^1 \times M \to \mathbb{R}$ is a normalized Hamiltonian function with compact support such that $\tilde{\phi}_H = \tilde{\phi}$. By the homotopy invariance ((2) of Proposition 2.2), $c^L(\mathbf{a}, H)$ does not depend on the choice of H and hence $c^L(\mathbf{a}, \tilde{\phi})$ is well-defined. For normalized Hamiltonian functions $F, G: S^1 \times M \to \mathbb{R}, F \natural G$ is also a normalized Hamiltonian function. Thus, the following triangle inequality follows from (3) of Proposition 2.2.

Proposition 7.1. $c^{L}(a * b, \tilde{\phi}\tilde{\psi}) \leq c^{L}(a, \tilde{\phi}) + c^{L}(b, \tilde{\psi})$ for any $\tilde{\phi}, \tilde{\psi} \in Ham(M, \omega)$ and any $a, b \in QH_{*}(L)$.

For a non-trivial idempotent **a** of $QH_*(L)$, we define the homogenization σ_a^L : $\widetilde{Ham}(M, \omega) \rightarrow \mathbb{R}$ of $c^L(\mathbf{a}, \cdot)$ by

$$\sigma_{\mathsf{a}}^{L}(\tilde{\phi}) = \lim_{k \to +\infty} \frac{c^{L}(\mathsf{a}, \tilde{\phi}^{k})}{k}.$$

By Proposition 7.1, we can prove the existence of this limit.

For a non-trivial element a of $QH_*(L)$ and elements \tilde{f}, \tilde{g} of $Ham(M, \omega)$, we define the following invariant $q_{a,\tilde{f}}(\tilde{g})$ by

$$q_{\mathsf{a},\tilde{f}}(\tilde{g}) = \sup_{k \in \mathbb{Z}} \{ c^L(\mathsf{a}, \tilde{f}^{-k} \tilde{g} \tilde{f}^k) \} + \sup_{k \in \mathbb{Z}} \{ c^L(\mathsf{a}, \tilde{f}^{-k} \tilde{g}^{-1} \tilde{f}^k) \}.$$

For simplicity, let $q_{a}(\tilde{g})$ denote $q_{a,id}(\tilde{g})$. The author does not know whether $q_{a,\tilde{f}}(\tilde{g}) < +\infty$ for any $\tilde{f}, \tilde{g} \in \widetilde{\text{Ham}}(M, \omega)$.

Then we can prove the following proposition which generalizes Proposition 3.5.3 in [PR].

Proposition 7.2. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$.

Then, for any non-trivial idempotent **a** of $QH_*(L)$ and any $\tilde{f}, \tilde{g} \in \widetilde{Ham}(M, \omega)$,

$$|\sigma_{\mathsf{a}}^{L}(\tilde{f}\tilde{g}) - \sigma_{\mathsf{a}}^{L}(\tilde{f}) - \sigma_{\mathsf{a}}^{L}(\tilde{g})| \le q_{\mathsf{a},\tilde{f}}(\tilde{g})$$

In particular, $q_{a,\tilde{f}}(\tilde{g}) \ge 0$.

Proof. Note that

$$(\tilde{f}\tilde{g})^k = \tilde{h}\tilde{f}^k$$
, where $\tilde{h} = (\tilde{f}\tilde{g}\tilde{f}^{-1})(\tilde{f}^2\tilde{g}\tilde{f}^{-2})\cdots(\tilde{f}^k\tilde{g}\tilde{f}^{-k})$

Thus, by Proposition 7.1,

$$c^{L}(\mathbf{a}, (\tilde{f}\tilde{g})^{k}) \leq c^{L}(\mathbf{a}, \tilde{h}) + c^{L}(\mathbf{a}, \tilde{f}^{k}),$$

$$c^{L}(\mathbf{a}, (\tilde{f}\tilde{g})^{k}) \geq c^{L}(\mathbf{a}, \tilde{f}^{k}) - c^{L}(\mathbf{a}, \tilde{h}^{-1}).$$

Thus

$$-c^{L}(\mathbf{a},\tilde{h}^{-1}) - c^{L}(\mathbf{a},\tilde{g}^{k}) \leq c^{L}(\mathbf{a},(\tilde{f}\tilde{g})^{k}) - c^{L}(\mathbf{a},\tilde{f}^{k}) - c^{L}(\mathbf{a},\tilde{g}^{k}) \leq c^{L}(\mathbf{a},\tilde{h}) - c^{L}(\mathbf{a},\tilde{g}^{k}).$$

Since $\tilde{h}^{-1} = (\tilde{f}^k \tilde{g}^{-1} \tilde{f}^{-k}) \cdots (\tilde{f}^2 \tilde{g}^{-1} \tilde{f}^{-2}) (\tilde{f}^1 \tilde{g}^{-1} \tilde{f}^{-1})$, by Proposition 7.1 and the definition of $q_{a,\tilde{f}}(\tilde{g})$,

$$\begin{aligned} c^{L}(\mathbf{a}, \tilde{h}^{-1}) + c^{L}(\mathbf{a}, \tilde{g}^{k}) \\ &\leq c^{L}(\mathbf{a}, \tilde{f}^{k} \tilde{g}^{-1} \tilde{f}^{-k}) + \dots + c^{L}(\mathbf{a}, \tilde{f}^{2} \tilde{g}^{-1} \tilde{f}^{-2}) + c^{L}(\mathbf{a}, \tilde{f}^{1} \tilde{g}^{-1} \tilde{f}^{-1}) + kc^{L}(\mathbf{a}, \tilde{g}) \\ &\leq \sum_{i=-k}^{-1} (c^{L}(\mathbf{a}, \tilde{f}^{-i} \tilde{g}^{-1} \tilde{f}^{i}) + c^{L}(\mathbf{a}, \tilde{g})) \\ &\leq kq_{\mathbf{a}, \tilde{f}}(\tilde{g}). \end{aligned}$$

By Proposition 7.1,

$$c^{L}(\mathbf{a}, \mathrm{id}) - c^{L}(\mathbf{a}, \tilde{g}^{k}) \leq c^{L}(\mathbf{a}, \tilde{g}^{-k}) \leq kc^{L}(\mathbf{a}, \tilde{g}^{-1}).$$

Hence $-c^{L}(\mathbf{a}, \tilde{g}^{k}) \leq kc^{L}(\mathbf{a}, \tilde{g}^{-1}) - c^{L}(\mathbf{a}, \mathrm{id})$. Thus, by Proposition 7.1 and the definitions of \tilde{h} and $q_{\mathbf{a}, \tilde{f}}(\tilde{g})$.

$$c^{L}(\mathbf{a}, \tilde{h}) - c^{L}(\mathbf{a}, \tilde{g}^{k})$$

$$\leq c^{L}(\mathbf{a}, \tilde{f}^{-1}\tilde{g}\tilde{f}) + c^{L}(\mathbf{a}, \tilde{f}^{-2}\tilde{g}\tilde{f}^{2}) + \dots + c^{L}(\mathbf{a}, \tilde{f}^{-k}\tilde{g}\tilde{f}^{k}) + kc^{L}(\mathbf{a}, \tilde{g}^{-1}) - c^{L}(\mathbf{a}, \mathrm{id})$$

$$\leq \sum_{i=1}^{k} (c^{L}(\mathbf{a}, \tilde{f}^{-i}\tilde{g}\tilde{f}^{i}) + c^{L}(\mathbf{a}, \tilde{g}^{-1})) - c^{L}(\mathbf{a}, \mathrm{id})$$

$$\leq kq_{\mathbf{a},\tilde{f}}(\tilde{g}) - c^{L}(\mathbf{a}, \mathrm{id}).$$

Thus

$$-kq_{\mathsf{a},\tilde{f}}(\tilde{g}) \le c^{L}(\mathsf{a},(\tilde{f}\tilde{g})^{k}) - c^{L}(\mathsf{a},\tilde{f}^{k}) - c^{L}(\mathsf{a},\tilde{g}^{k}) \le kq_{\mathsf{a},\tilde{f}}(\tilde{g}) - c^{L}(\mathsf{a},\mathrm{id}).$$

By dividing by k and passing to the limit as $k \to +\infty$, we complete the proof.

For autonomous Hamiltonian functions $F, G: M \to \mathbb{R}$, we define D(F, G) by

 $D(F,G) = \min\{\sup_{t \in \mathbb{R}} q_{\mathtt{a},\tilde{\phi}_F}(\tilde{\phi}_{tG}), \sup_{t \in \mathbb{R}} q_{\mathtt{a},\tilde{\phi}_G}(\tilde{\phi}_{tF})\}.$

By Proposition 7.2, $D(F, G) \ge 0$ or $D(F, G) = +\infty$.

We give a generalization of Poisson bracket inequality.

Proposition 7.3. Let *L* be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$ and $QH_*(L) \ne 0$. Then, for any non-trivial idempotent a of $QH_*(L)$ and any Hamiltonian functions $F, G: M \to \mathbb{R}$ with $D(F, G) < +\infty$,

$$|\zeta_{\mathsf{a}}^{L}(F+G) - \zeta_{\mathsf{a}}^{L}(F) - \zeta_{\mathsf{a}}^{L}(G)| \le (2D(F,G) \cdot ||\{F,G\}||)^{1/2}.$$

In the present paper, we give good upper bounds of D(F, G) under some situations.

By Proposition 2.2 and Proposition 7.2, we can prove Proposition 7.3 quite similarly to Proposition 4.6.1 of [PR]. Therefore we omit the proof of Proposition 7.3.

Remark 7.4. Let U be an open subset with the bounded spectrum condition. If the equality " $q_{a,\tilde{f}}(\tilde{g}) = q_{a,id}(\tilde{g})$ " holds for any $\tilde{f}, \tilde{g} \in Ham(M, \omega)$, then we have D(F, G) < K for any Hamiltonian function $F: M \to \mathbb{R}$ with $Supp(F) \subset U$.

In the case of Oh-Schwarz spectral invariants, a similar equality actually holds and hence we obtain an upper bound of D(F, G). However, in our case, the above equality does not hold and thus it is more difficult to give an upper bound of D(F, G).

8. Proof of partial quasi-additivities and Theorem 1.4

Proposition 8.1. Let $G: M \to \mathbb{R}$ be a Hamiltonian function. Assume that Supp(G) satisfies the bounded spectrum condition with respect to some idempotent a of $QH_*(L)$, $\zeta_a^L(G) = 0$.

Proof. By the definition of the bounded spetrum condition, there is a positive number K such that

$$c^L(\mathsf{a}, kG) < K,$$

for any $k \in \mathbb{Z}$. Then, by the triangle inequality and the homotopy invariance ((2) and (3) of Proposition 2.2),

$$c^{L}(\mathbf{a}, kG) \ge -c^{L}(\mathbf{a}, -kG) + c^{L}(\mathbf{a}, 0) > -K + c^{L}(\mathbf{a}, 0),$$

and hence

$$-K + c^L(\mathbf{a}, 0) < c^L(\mathbf{a}, kG) < K,$$

for any $k \in \mathbb{Z}$. Thus

$$\zeta_{\mathsf{a}}^{L}(G) = \lim_{k \to +\infty} \frac{c^{L}(\mathsf{a}, kG)}{k} = 0.$$

To prove (1) and (1') of Proposition 6.1, we give a more general proposition.

Proposition 8.2. For Hamiltonian functions $F, G: M \to \mathbb{R}$ with $\{F, G\} = 0$ and Supp(G) satisfies the bounded spectrum condition with respect to $a, \zeta_a^L(F+G) = \zeta_a^L(F)$.

 \Box

Proof. Since $\{F, G\} = 0$, $(\tilde{\phi}_G)^{-k} \tilde{\phi}_F^t (\tilde{\phi}_G)^k = \tilde{\phi}_F^t$ for any $t \in \mathbb{R}$ and any $k \in \mathbb{Z}$. Hence $q_{a,\tilde{\phi}_G}(\tilde{\phi}_F) = q_a(\tilde{\phi}_F)$ and $D(F, G) < +\infty$. Thus, by Proposition 7.3,

$$|\zeta_{\mathsf{a}}^{L}(F+G) - \zeta_{\mathsf{a}}^{L}(F) - \zeta_{\mathsf{a}}^{L}(G)|| \le (2D(F,G) \cdot ||\{F,G\}||)^{1/2} = 0.$$

Hence, by Proposition 8.1,

$$\zeta_{\mathsf{a}}^{L}(F+G) = \zeta_{\mathsf{a}}^{L}(F) + \zeta_{\mathsf{a}}^{L}(G) = \zeta_{\mathsf{a}}^{L}(F).$$

Proof of (1') of Proposition 6.1. Since Supp(G) is displaceable from *L*, there is an open neighborhood *U* of Supp(G) which is displaceable from *L*. Thus, by Proposition 4.2, Supp(G) satisfies the bounded spectral condition with respect to any idempotent a and thus, by Proposition 8.2,

$$\zeta_{a}^{L}(F+G) = \zeta_{a}^{L}(F).$$

We can prove (1) of Proposition 6.1 by Proposition 4.3 similarly to (1') of Proposition 6.1.

The proof of Theorem 1.4 is quite similar to the one of the original stem case if we know (1) and (1') of Proposition 6.1 and we omit the proof.

9. Non-degeneracy of spectral norms

We prove the following lemmas for a non-trivial idempotent **a** of $QH_*(L)$ with $c^L(\mathbf{a}, 0) \leq 0$.

Lemma 9.1. Assume that an idempotent a of $QH_*(L)$ satisfies $c^L(a, 0) \le 0$. For any Hamiltonian function $F, H: S^1 \times M \to \mathbb{R}$ with $\text{Supp}(F_t) \cap L = \emptyset$ for any t,

$$c^{L}(\mathbf{a}, F \natural H) = c^{L}(\mathbf{a}, H \natural F) = c^{L}(\mathbf{a}, H).$$

Proof. Since Supp(*F*) is disjoint from *L*, by Proposition 4.2, $c^{L}(a, F) \leq 2c^{L}(a, 0) \leq 0$ and $c^{L}(a, \bar{F}) \leq 2c^{L}(a, 0) \leq 0$. By the triangle inequality and the homotopy invariance ((3), (2) of Proposition 2.2),

$$c^{L}(\mathbf{a}, H \natural F) \le c^{L}(\mathbf{a}, H) + c^{L}(\mathbf{a}, F),$$

$$c^{L}(\mathbf{a}, H \natural F) + c^{L}(\mathbf{a}, \overline{F}) \ge c^{L}(\mathbf{a}, H).$$

Thus

$$c^{L}(\mathbf{a}, H) - c^{L}(\mathbf{a}, \overline{F}) \le c^{L}(\mathbf{a}, H \natural F) \le c^{L}(\mathbf{a}, H) + c^{L}(\mathbf{a}, F)$$

Since $c^{L}(\mathbf{a}, F) \leq 0$ and $c^{L}(\mathbf{a}, \bar{F}) \leq 0$, $c^{L}(\mathbf{a}, F) = c^{L}(\mathbf{a}, \bar{F}) = 0$ and $c^{L}(\mathbf{a}, H \natural F) = c^{L}(\mathbf{a}, H)$. We can prove $c^{L}(\mathbf{a}, F \natural H) = c^{L}(\mathbf{a}, H)$ similarly.

Lemma 9.2. Assume that an idempotent a of $QH_*(L)$ satisfies $c^L(a, 0) \leq 0$. For any $\psi \in Ham(M, \omega)$ and any Hamiltonian function $F: S^1 \times M \to \mathbb{R}$ with $Supp(F_t) \cap L = \emptyset$ for any t,

$$q_{\mathsf{a},\tilde{\phi}_F}(\tilde{\psi}) = q_\mathsf{a}(\tilde{\psi}).$$

Proof. Take a normalized Hamiltonian H generating $\tilde{\psi}$. By (2) of Proposition 2.2,

$$c^{L}(\mathbf{a}, \tilde{\phi}_{F}^{-k} \tilde{\psi} \tilde{\phi}_{F}^{k}) = c^{L}(\mathbf{a}, \bar{F}^{\natural k} \natural H \natural F^{\natural k}),$$

 $c^{L}(\mathbf{a}, \tilde{\phi}_{F}^{-k} \tilde{\psi}^{-1} \tilde{\phi}_{F}^{k}) = c^{L}(\mathbf{a}, \bar{F}^{\natural k} \natural \bar{H} \natural F^{\natural k}).$

By Lemma 9.1, $c^{L}(\mathbf{a}, \bar{F}^{\natural k} \natural H \natural F^{\natural k}) = c^{L}(\mathbf{a}, H) = c^{L}(\mathbf{a}, \tilde{\psi})$ and $c^{L}(\mathbf{a}, \bar{F}^{\natural k} \natural \bar{H} \natural F^{\natural k}) = c^{L}(\mathbf{a}, \bar{H}) = c^{L}(\mathbf{a}, \tilde{\psi}^{-1})$. Thus, by the definition of $q_{\mathbf{a}, \tilde{\phi}_{F}}(\tilde{\phi})$, we complete the proof.

Proposition 9.3. Assume that an idempotent \mathbf{a} of $QH_*(L)$ satisfies $c^L(\mathbf{a}, 0) \leq 0$. Let \hat{U} be an open subset of M with $\hat{U} \cap L \neq \emptyset$. Then there exists an autonomous Hamiltonian function $F: M \to \mathbb{R}$ such that $\operatorname{Supp}(F) \subset \hat{U}$ and $c^L(\mathbf{a}, F) > 0$.

Proof. Fix a Riemannian metric on *M*. For a subset *X* and a positive number *r*, let $N_r(X)$, $\bar{N}_r(X)$ denote the *r*-neighborhood of *X*, its topological closure, respectively.

Since $\hat{U} \cap L \neq \emptyset$, we can take a point *x* in $\hat{U} \cap L$. Then, there exists a positive number ϵ such that $\bar{N}_{4\epsilon}(\{x\})$ is displaceable from *L* and $\bar{N}_{4\epsilon}(\{x\}) \subset \hat{U}$.

Set $L' = L \setminus \overline{N}_{3\epsilon}(\{x\})$. Since dim $L' = \dim L = \frac{1}{2} \dim M$ and L' is not compact, L' is displaceable from L. Thus there is a positive number ϵ' such that $\overline{N}_{2\epsilon'}(L')$ is also displaceable from L.

Set $\epsilon'' = \min\{\epsilon, \epsilon'\}$. We define three subsets U_P, U_S, U_K of M by $U_P = N_{4\epsilon}(\{x\}), U_S = N_{2\epsilon''}(L'), U_K = M \setminus N_{\epsilon''}(L)$. Then,

- U_S is displaceable from L,
- $U_P \cup U_S \cup U_K = M$,
- $U_K \cap L = \emptyset$.

Take a partitions $\{P, S, K\}$ of unity subordinated to the open cover $\{U_P, U_S, U_K\}$ *i.e.* $P, S, K: M \rightarrow [0, 1]$ are smooth functions, $\text{Supp}(P) \subset U_P$, $\text{Supp}(S) \subset U_S$, $\text{Supp}(K) \subset U_K$, and $P + S + K \equiv 1$. Since U_{-} is displaceable from L and (S, 1 - S) = 0, by (1') (2) of Proposition 6.1

Since U_S is displaceable from L and $\{S, 1 - S\} = 0$, by (1'), (2) of Proposition 6.1,

$$1 = \zeta_{a}^{L}(1) = \zeta_{a}^{L}(1 - S).$$

Since $U_K \cap L = \emptyset$, by Lemma 9.2,

$$q_{\mathsf{a},\tilde{\phi}_K}(\tilde{\phi}_P) = q_\mathsf{a}(\tilde{\phi}_P).$$

To prove by contradiction, we assume $c^{L}(\mathbf{a}, tP) \leq 0$ for any real number t. Since P is a timeindependent Hamiltonian function, $\overline{(tP)} = -tP$ for any t and hence $(\tilde{\phi}_{tP})^{-1} = \tilde{\phi}_{-tP}$. Thus, for any t,

$$q_{\mathbf{a}}(\tilde{\phi}_{tP}) = c^{L}(\mathbf{a}, \tilde{\phi}_{tP}) + c^{L}(\mathbf{a}, (\tilde{\phi}_{tP})^{-1})$$
$$= c^{L}(\mathbf{a}, \tilde{\phi}_{tP}) + c^{L}(\mathbf{a}, \tilde{\phi}_{-tP})$$
$$= c^{L}(\mathbf{a}, tP) + c^{L}(\mathbf{a}, -tP) \le 0.$$

By Proposition 7.2, $q_a(\tilde{\phi}_{tP}) = 0$ and therefore $q_{a,\tilde{\phi}_k}(\tilde{\phi}_{tP}) = q_a(\tilde{\phi}_{tP}) = 0$. Hence, by Proposition 7.3,

$$\zeta_{\mathsf{a}}^{L}(P+K) = \zeta_{\mathsf{a}}^{L}(P) + \zeta_{\mathsf{a}}^{L}(K).$$

Since U_P and U_K are displaceable from L, by Propositions 4.2 and 8.1,

$$\zeta_{\rm a}^L(P) = \zeta_{\rm a}^L(K) = 0,$$

and hence $\zeta_a^L(P + K) = 0$. Since $P + S + K \equiv 1$, this contradict $1 = \zeta_a^L(1 - S)$. Hence we prove that $c^L(a, tP) > 0$ for some *t*. Since $\text{Supp}(P) \subset U_P \subset \hat{U}$, we complete the proof. \Box

Proof of Theorem 1.13. If $c^{L}(\mathbf{a}, 0) > 0$, then, by (3) of Proposition 2.2, $c^{L}(\mathbf{a}, H) + c^{L}(\mathbf{a}, \overline{H}) \ge c^{L}(\mathbf{a}, 0) > 0$. Thus we may assume $c^{L}(\mathbf{a}, 0) \le 0$.

Since $\tilde{\phi}_H(L) \neq L$, there is an open subset U of M such that $\tilde{\phi}_H(U) \cap L = \emptyset$ and $U \cap L \neq \emptyset$. Then, by Lemma 9.3, there exists a Hamiltonian function $F: S^1 \times U \to \mathbb{R}$ with compact support on $S^1 \times U$ such that $c^L(\mathbf{a}, F) > 0$. Thus, by Proposition 4.2, $c^L(\mathbf{a}, H) + c^L(\mathbf{a}, \bar{H}) \ge c^L(\mathbf{a}, F) > 0$. \Box

10. The energy capacity inequality and heavy, superheavy subsets on a torus

Proposition 10.1. Let *L* be a Lagrangian submanifold of a closed symplectic manifold (M, ω) with $\omega(\pi_2(M, L)) = 0$, $N_L \ge 2$ and $QH_*(L) \ne 0$ (Note that $\omega(\pi_2(M, L)) = 0$ implies monotonicity of *L*). Assume that a Hamiltonian function $H: M \to \mathbb{R}$ with compact support is *L*-slow. Then

$$c^L([L], H) = m(H).$$

Proof. Since $\omega(\pi_2(M, L)) = 0$ and H is L-slow, Spec $(H; L) = \{m(H), 0\}$, Spec $(\bar{H}; L) = \{-m(H), 0\}$ and $\phi_H(L) \neq L$. By Theorem 1.13 and $\phi_H(L) \neq L$, $c^L([L], H) + c^L([L], \bar{H}) > 0$. Thus, by the spectrality ((6) of Proposition 2.2), $c^L([L], H) = m(H)$ and $c^L([L], \bar{H}) = 0$.

Proof of Theorem 1.16. Let *F* be an $L \cap U$ -simple and $L \cap U$ -slow Hamiltonian function with compact support on *U*. Define the Hamiltonian function $\hat{F} \colon M \to \mathbb{R}$ by

$$\hat{F}(x) = \begin{cases} F(x) & \text{if } x \in U, \\ 0 & \text{if } x \notin U. \end{cases}$$

Then, by the definition of *L*-simplicity, \hat{F} is an *L*-slow Hamiltonian function with compact support on *M*. Then, by Proposition 10.1, $c^{L}([L], \hat{F}) = m(F)$. By Corollary 4.5, $c^{L}([L], \hat{F}) \leq \bar{E}(U; L)$. Thus $m(F) \leq \bar{E}(U; L)$. By taking the supremum over all $L \cap U$ -simple and $L \cap U$ -slow functions with compact support on *U*, we complete the proof.

To prove (3) of Proposition 1.7, we use the following lemma.

Lemma 10.2. Let $f : \mathbb{R}/R\mathbb{Z} \to \mathbb{R}$ be a smooth function with only two critical value $v_{\text{max}} > v_{\text{min}}$ and define a Hamiltonian function $F : T_R^2 \to \mathbb{R}$ on T_R^2 by F(x, y) = f(y). Then $c^L([L_m^s], F) = v_{\text{max}}$.

Proof. By the definition of the spectrum, $\text{Spec}(F; L_m^s) = \{v_{\text{max}}, v_{\text{min}}\}$ and $\text{Spec}(\bar{F}; L_m^s) = \{-v_{\text{max}}, -v_{\text{min}}\}$. However, by Theorem 1.13, $c^{L_m^s}([L_m^s], F) + c^{L_m^s}([L_m^s], \bar{F}) > 0$. Thus by the spectrality ((6) of Proposition 2.2), $c^{L_m^s}([L_m^s], F) = v_{\text{max}}$ and $c^{L_m^s}([L_m^s], \bar{F}) = -v_{\text{min}}$.

Proof of (3) of Proposition 1.7. Fix a Hamiltonian function $H: S^1 \times T_R^2 \to \mathbb{R}$. Take a function $f: \mathbb{R}/R\mathbb{Z} \to \mathbb{R}$ with only two critical values $v_{\max} = \inf_{S^1 \times L_1^t} H > v_{\min}$ such that $F(x, y) \leq H(x, y)$ where $F: T_R^2 \to \mathbb{R}$ is a Hamiltonian function on T_R^2 defined by F(x, y) = f(x). Then by Lemma 10.2, $c^{L_m^s}([L_m^s], kF) = kv_{\max} = k \inf_{S^1 \times L_1^t} H$ for any positive integer k. By the monotonicity of Lagrangian spectral invariants, $\zeta_{[L_m^s]}^{L_m^s}(H) \geq \zeta_{[L_m^s]}^{L_m^s}(F) = \inf_{S^1 \times L_1^t} H$.

We can prove the following proposition similarly to (3) of Proposition 1.7.

Proposition 10.3. Let L be a monotone Lagrangian submanifold of a closed symplectic manifold (M, ω) with $N_L \ge 2$, $\omega(\pi_2(M, L)) = \{0\}$ and $QH_*(L) \ne 0$. Let L' be a compact Lagrangian submanifold of (M, ω) . Assume that there are a Weinstein coordinate $w: W \rightarrow T^*L'$ of L' (note that w(L') is the zero section of T^*L') and a point q of L' such that

$$w(W \cap L) = T_a^* L' \cap \operatorname{Im}(w).$$

Also assume that L' is diffeomorphic to a torus or a negatively curved Riemannian manifold and the map $\iota_*\pi_1(L',q) \to \pi_1(M,q) \to \pi_1(M,L,q)$ is injective, where ι is the homomorphism induced from the inclusion map $\iota: (L',q,q) \to (M,L,q)$. Then L' is (L, \mathbf{a}) -heavy for any nontrivial idempotent \mathbf{a} .

For example, we can apply Proposition 10.3 to a Riemann surface with higher genus. We give proofs of the other parts of Proposition 1.7.

Proof of (1) of Proposition 1.7. If s = s', $L_m^{s'} = L_m^s$ is an L_m^s -stem and in particular, $(L_m^s, [L_m^s])$ -superheavy.

Proof of (2) of Proposition 1.7. If $s \neq s'$, $L_m^{s'} \cap L_m^s = \emptyset$. Thus, by (1) of Proposition 1.7 and Proposition 6.2, $L_m^{s'}$ is not $(L_m^s, [L_m^s])$ -heavy.

Proof of (4) of Proposition 1.7. Take $t' \in \mathbb{R}/\mathbb{Z}$ with $t' \neq t$. Then, by (3) of Proposition 1.7, $L_1^{t'}$ is $(L_m^s, [L_m^s])$ -heavy. Since $L_1^t \cap L_1^{t'} = \emptyset$, by Proposition 6.2, L_1^t is not $(L_m^s, [L_m^s])$ -superheavy. \Box

Proof of (5) of Proposition 1.7. For any *s*, *s'*, *t*, $L_m^{s'} \cup L_1^t$ is an L_m^s -stem and in particular, $(L_m^s, [L_m^s])$ -superheavy.

11. Proof of Theorem 1.23

In order to prove Theorem 1.23, we give an upper bound of the spectral invariant associated with a Hamiltonian function $F: S^1 \times M \times \mathbb{A}(R) \to \mathbb{R}$ such that $Ch(F; L \times \mathbb{A}(R)_0, L \times \mathbb{A}(R)_s, \alpha_s) = \emptyset$. Here, for $R = (R_1, \ldots, R_n) \in (\mathbb{R}_{>0})^n$ and a positive real number ϵ with $\epsilon < \min\{R_1, \ldots, R_n\}$, let $R(\epsilon)$ denote $(R_1 - \epsilon, \ldots, R_n - \epsilon)$ and $I_{R(\epsilon)}^n$ denote $(\epsilon, R_1) \times \cdots \times (\epsilon, R_n) \subset (\mathbb{R}_{>0})^n$. For R = $(R_1, \ldots, R_n) \in (\mathbb{R}_{>0})^n$, T_R^n is defined to be $\mathbb{R}/R_1\mathbb{Z} \times \cdots \mathbb{R}/R_n\mathbb{Z}$ and we set the symplectic form $\omega_0 = dp_1 \wedge dq_1 + \cdots + dp_n \wedge dq_n$ on $T_R^n \times T^n$ with the coordinates $(p, q) = (p_1, \ldots, p_n, q_1, \ldots, q_n)$.

Proposition 11.1. Let *L* be a λ -monotone Lagrangian submanifold of a 2*m*-dimensional closed symplectic manifold (M, ω) with $N_L \geq 2$ and $QH_*(L) \neq 0$. Let $s = (s_1, \ldots, s_n)$ and $R = (R_1, \ldots, R_n)$ be elements of \mathbb{Z}^n and $(\mathbb{R}_{>0})^n$, respectively. For a positive real number ϵ with $3\epsilon < \min\{R_1, \ldots, R_n\}$, let U_{ϵ} be the open subset of $T_R^n \times T^n$ defined by

$$U_{\epsilon} = \{ (p,q) \in T_R^n \times T^n; p \in I_{R(3\epsilon)}^n \}$$

We fix the symplectic form $\operatorname{pr}_1^* \omega + \operatorname{pr}_2^* \omega_0$ on $M \times T_R^n \times T^n$, where $\operatorname{pr}_1 \colon M \times T_R^n \times T^n \to M$ and $\operatorname{pr}_2 \colon M \times T_R^n \times T^n \to T_R^n \times T^n$ are the projections defined by $\operatorname{pr}_1(x, p, q) = x$ and $\operatorname{pr}_2(x, p, q) = (p, q)$. Then for any Hamiltonian function $F \colon S^1 \times M \times U_{\epsilon} \to \mathbb{R}$ with compact support such that $\operatorname{Ch}(F; L \times (T_R^n \times \{0\}), L \times (T_R^n \times \{s\}), (0_L, \alpha_s)) = \emptyset$,

$$c^{L \times (T_R^n \times \{0\})}([L \times (T_R^n \times \{0\})], F) < 2 \sum_{i=1}^n R_i \cdot |s_i| + \lambda(m+n).$$

To prove Proposition 11.1, we use the following proposition. For a smooth path $z: [0, 1] \rightarrow M$, let ev(z) denote the point z(0).

Proposition 11.2. Let W be an open subset of a 2w-dimensional connected closed symplectic manifold (\hat{W}, ω) , \hat{Z}_0 , \hat{Z}_1 compact λ -monotone Lagrangian submanifolds of \hat{W} with $N_{\hat{Z}_0} \ge 2$ and $QH_*(\hat{Z}_0) \neq 0$ and α a homotopy class of $\pi_1(\hat{W}, \hat{Z}_0 \cup \hat{Z}_1)$. Assume that a Hamiltonian function $H: \hat{W} \to \mathbb{R}$ satisfies the following conditions.

- $\operatorname{Supp}(H) \subset W$
- $\phi_H(\hat{Z}_1) = \hat{Z}_0$ and $[\gamma_H^x] = \bar{\alpha}$ for any point x in Z_1 where $Z_1 = W \cap \hat{Z}_1$,
- Set $Z_0 = W \cap \hat{Z}_0$. On some Weinstein neighborhood of Z_0 , H is the pullback of a Morse function $\tilde{H}: Z_0 \to \mathbb{R}$,
- $ev(Ch(H; Z_0)) = Crit(H|_{Z_0}),$
- $\operatorname{ind}_{Morse}(x) = \operatorname{ind}_{CZ}([x, c_x])$ for any point x in $\operatorname{Crit}(H|_{Z_0})$.

Then for any Hamiltonian function $F: S^1 \times W \to \mathbb{R}$ with compact support such that $Ch(F; Z_0, Z_1, \alpha) = \emptyset$,

$$c^{Z_0}([\hat{Z}_0], F) \le 2 ||H||_{L^{\infty}} + \lambda w.$$

Proof. To give an upper bound of the spectral invariant associated with F, we consider the concatenation of ϕ_F^1 and a Hamiltonian diffeomorphism ϕ_H^1 with trajectories in $\bar{\alpha}$.

Let $K: S^1 \times \hat{W} \to \mathbb{R}$ be a Hamiltonian function defined by

$$K(t, x) = \begin{cases} \frac{\partial \chi}{\partial t}(t) H(\chi(t), x) & \text{when } t \in [0, \frac{1}{2}], \\ \frac{\partial \chi}{\partial t}(t - \frac{1}{2}) F(\chi(t - \frac{1}{2}), x) & \text{when } t \in [\frac{1}{2}, 1], \end{cases}$$

where $\chi: [0, \frac{1}{2}] \to [0, 1]$ is the function defined in the proof of Proposition 4.2. We claim

$$c^{Z_0}([\hat{Z}_0], K) \le ||H||_{L^{\infty}} + \lambda w.$$

Let $[z, \hat{z}] \in \tilde{\Omega}_0(\hat{Z}_0)$ and define *x* by $x = z(0) \in \hat{Z}_0$. If $x \in Z_0$, by the assumption of *H*, $[\gamma_H^x] = \bar{\alpha}$. Since the path γ_K^x is the concatenation of the paths γ_H^x and $\gamma_F^{\phi_H(x)}$ up to parameter change, $Ch(F; Z_0, Z_1, \alpha) = \emptyset$ implies $\gamma_K^x \notin \Omega_0(\hat{Z}_0)$ for any $x \in Z_0$. If $x \notin Z_0$, then $\phi_H(x) \notin Z_0$. Thus γ_K^x is equal to γ_H^x up to parameter change and $\int_0^1 H(t, \gamma_H^x(t)) dt = \int_0^1 K(t, \gamma_K^x(t)) dt$. Therefore there exists the natural inclusion map ι : $\tilde{Ch}(K; Z_0) \to \tilde{Ch}(H; Z_0)$ which preserves values of the action functionals and the Conley-Zehnder indices.

We give an upper bound of the critical value of the action functional $\mathcal{A}_{K}^{\hat{z}_{0}}$ which attains the fundamental class $[\hat{z}_{0}]$ of $QH_{*}(\hat{z}_{0})$. Since every element of $Ch(H; \hat{z}_{0})$ is a constant path, every element of $Ch(K; \hat{z}_{0})$ is also a constant path. Since $Ch(K; \hat{z}_{0})$ is a finite set and \hat{z}_{0} is monotone, $\mathcal{A}_{K}^{\hat{z}_{0}}(\tilde{Ch}(H; \hat{z}_{0}))$ is a discrete subset of \mathbb{R} . Thus $c^{\hat{z}_{0}}([\hat{z}_{0}], K)$ is attained by a 1-length trajectory of the Conley-Zehnder index *w* that is the dimension of the fundamental class. Since every element of $Ch(K; \hat{z}_{0})$ is a constant path, there exist a point *x* in $Crit(\tilde{H})$ and $A \in \pi_{2}(\hat{W}, \hat{z}_{0})$ such that $ind_{CZ}([x, c_{x} \natural A]) = w$ and $c^{\hat{z}_{0}}([\hat{z}_{0}], K) = \mathcal{A}_{K}^{\hat{z}_{0}}([x, c_{x} \natural A])$. Then, by the assumption,

$$ind_{Morse}(x) - \mu(A)$$

= $ind_{CZ}([x, c_x]) - \mu(A)$
= $ind_{CZ}([x, c_x \natural A])$
= w .

Since $0 \le \operatorname{ind}_{\operatorname{Morse}}(x) \le w$,

$$w \le \mu(A) \le 0.$$

Since ι preserves values of the action functionals,

$$\mathcal{R}_{K}^{\tilde{Z}_{0}}([x, c_{x} \natural A]) = \mathcal{R}_{H}^{\tilde{Z}_{0}}([x, c_{x} \natural A])$$
$$= H(x) - \omega(A)$$
$$= H(x) - \lambda \mu(A).$$

Thus, by $-w \leq \mu(A) \leq 0$ and $\lambda \geq 0$, $c^{\hat{Z}_0}([\hat{Z}_0], K) \leq ||H||_{L^{\infty}} + \lambda w$. By $||\bar{H}||_{L^{\infty}} = ||H||_{L^{\infty}}$, the Lipschitz continuity and the homotopy invariance for spectral invariants (Proposition 2.2 (1) and (2)) imply

$$c^{\hat{Z}_0}([\hat{Z}_0], F) \le c^{\hat{Z}_0}([\hat{Z}_0], K) + \|\bar{H}\|_{L^{\infty}}$$
$$= 2\|H\|_{L^{\infty}} + \lambda w.$$

The idea of using a Hamiltonian function H satisfying the above conditions comes from Irie's paper [I]. Seyfaddini's techniques of using the monotonicity assumption [Se] are also important in our proof.

To prove Proposition 11.1, we construct the Hamiltonian function H in Proposition 11.2 using $H^{R,\epsilon,e}$ given by the following lemma.

Lemma 11.3. Let R, ϵ be positive real numbers such that $3\epsilon < R$. Let w_1 and w_2 denote the points $(\epsilon, 0)$ and $(2\epsilon, 0)$ in $T_R^1 \times T^1$, respectively. For an integer *e*, there exists a Hamiltonian function $H^{R,\epsilon,e}: T^1_R \times T^1 \to \mathbb{R}$ satisfying the following conditions.

- $H^{R,\epsilon,e}(p,q) = -ep \text{ on } U_{\epsilon} = (3\epsilon, R) \times T^1$,
- Crit($H^{R,\epsilon,e}|_{T^1_p \times \{0\}}$) = { w_1, w_2 },
- $H^{R,\epsilon,e}|_{T_p^1 \times \{0\}}$ is a Morse function,
- $||H^{R,\epsilon,e}||_{L^{\infty}}^{\sim} < (R-\epsilon) \cdot |e|,$
- $dH^{R,\epsilon,e}$ is C^1 -small near w_1, w_2 , $ev(Ch(H^{R,\epsilon,e}; T^1_R \times \{0\})) = Crit(H^{R,\epsilon,e}|_{T^1_R \times \{0\}}).$

Here $Crit(H^{R,\epsilon,e})$ is the set of critical points of $H^{R,\epsilon,e}$.

Proof. Let $\underline{H}^{R,\epsilon,e}$: $T_R^1 \to \mathbb{R}$ be a function satisfying the following conditions.

- $\underline{H}^{R,\epsilon,e}(p) = -ep$ on $U_{\epsilon} = (3\epsilon, R)$,
- $\overline{\operatorname{Crit}}(\underline{H}^{R,\epsilon,e}) = \{\epsilon, 2\epsilon\},\$
- $H^{R,\epsilon,\overline{e}}$ is a Morse function,
- $\overline{||\underline{H}^{R,\epsilon,e}||_{L^{\infty}}} < (R \epsilon) \cdot |e|,$ $d\underline{H}^{R,\epsilon,e}$ is C^1 -small near $p = \epsilon, 2\epsilon$.

Define the Hamiltonian function $H^{R,\epsilon,e}: T^1_R \times T^1 \to \mathbb{R}$ by $H^{R,\epsilon,e}(p,q) = \underline{H}^{R,\epsilon,e}(p)$. Then, the last condition follows from C¹-smallness of $d\underline{H}^{R,\epsilon,e}$ near $p = \epsilon, 2\epsilon$. The other conditions immediately follow from the conditions of $H^{R,\epsilon,e}: \overline{T}_R^1 \to \mathbb{R}$.

Proof of Proposition 11.1. To use Proposition 11.2, we construct the Hamiltonian function H. Define the Hamiltonian function $H': T_R^n \times T^n \to \mathbb{R}$ by

$$H'(p,q) = \sum_{i=1}^n H^{R_i,\epsilon_i,e_i}(p_i,q_i).$$

Then $[\gamma_{H'}^x] = -e \in \pi_1(T_R^n \times T^n, T_R^n \times \{0\})$ for any $x \in U_{\epsilon}$. Thus we can take a neighborhood V of \bar{U}_{ϵ} such that

$$ev(Ch(H'; L \times (T_R^n \times \{0\}))) \cap \overline{V} = \emptyset.$$

In order to compute the spectral invariant associated with F, we take a perturbation of pr_1^*H' (recall that $\operatorname{pr}_1: M \times T_R^n \times T^n \to M$ is the first projection). Let $\rho: T_R^n \times T^n \to [0, 1]$ be a function such that

$$\rho(p,q) = \begin{cases} 1 & \text{for any } (p,q) \in (T_R^n \times T^n) \setminus V, \\ 0 & \text{for any } (p,q) \in U_{\epsilon}. \end{cases}$$

Let $G: M \to \mathbb{R}$ be a function satisfying the following conditions.

• On some Weinstein neighborhood of L, G is the pullback of a C^2 -small Morse function $\tilde{G}: L \to \mathbb{R},$

Define the Hamiltonian function $H: M \times T_R^n \times T^n \to \mathbb{R}$ by

$$H(y, p, q) = H'(p, q) + \rho(p, q) \cdot G(y)$$

Since \tilde{G} is sufficiently C^2 -small, then

- $ev(Ch(H; L \times (T_R^n \times \{0\}))) \cap (M \times V) = \emptyset$, and
- there exist only finitely many points y_1, \ldots, y_k in M such that $Crit(G) = ev(Ch(tG; L)) = \{y_1, \ldots, y_k\}$ for any $t \in (0, 1]$.

Since $ev(Ch(H; L \times (T_R^n \times \{0\}))) \cap (M \times V) = \emptyset$, $ev(Ch(H; L \times (T_R^n \times \{0\}))) \subset M \times ((T_R^n \times T^n) \setminus V)$. Since $\rho(p, q) = 1$ for any $(p, q) \in (T_R^n \times T^n) \setminus V$,

$$H(y, p, q) = H'(p, q) + G(y),$$

for any $(y, p, q) \in M \times ((T_R^n \times T^n) \setminus V)$. Thus

$$ev(Ch(H; L \times (T_R^n \times \{0\}))) = \{(y_i, (w_{j_1}, \dots, w_{j_n}), (0, \dots, 0))\}_{i \in \{1, \dots, k\}, j_1, \dots, j_n \in \{1, 2\}} = Crit(H|_{L \times (T_R^n \times \{0\})}).$$

By conditions of H^{R_i,ϵ,e_i} 's and G, dH is C¹-small near critical points of $H|_{L\times(T^n_b\times\{0\})}$. Thus

$$\operatorname{ind}_{\operatorname{Morse}}(x) = \operatorname{ind}_{\operatorname{CZ}}([x, c_x]),$$

for any point x in $\operatorname{Crit}(H|_{L\times(T_R^n\times\{0\})})$. Hence H satisfies the conditions of Proposition 11.2 and thus we apply Proposition 11.2.

By Proposition 11.2 and $\|\bar{H}\|_{L^{\infty}} = \|H\|_{L^{\infty}}$, the Lipschitz continuity and the homotopy invariance for spectral invariants (Proposition 2.2 (1) and (2)) imply

$$c^{L\times\mathbb{A}(R)_0}([L\times\mathbb{A}(R)_0],F) \le 2||H||_{L^{\infty}} + \lambda(m+n)$$

$$< 2(\sum_{i=1}^n (R_i - \epsilon) \cdot |e_i| + ||G||_{L^{\infty}}) + \lambda(m+n).$$

Since the Morse function G is sufficiently C^2 -small,

$$c^{L \times \mathbb{A}(R)_0}([L \times \mathbb{A}(R)_0], F) < 2 \sum_{i=1}^n R_i \cdot |e_i| + \lambda(m+n).$$

Proof of Theorem 1.23. Fix a positive real number ϵ such that $\epsilon < \min\{R_1, \ldots, R_n\}$ and take a Hamiltonian function $F: S^1 \times M \times I_{R(\epsilon)}^n \times T^n \to \mathbb{R}$ with compact support such that $F|_{S^1 \times Z \times T^n} \ge 2\sum_{i=1}^n R_i \cdot |e_i| + \max\{0, \lambda(m+n)\}$. Assume $Ch(F; L \times (T_R^n \times \{0\}), L \times (T_R^n \times \{s\}), (0_L, \alpha_s)) = \emptyset$. Then, Proposition 11.1 and the triangle inequality ((3) of Proosition 2.2) imply

$$\zeta_{[L\times\mathbb{A}(R(\epsilon))_0]}^{L\times\mathbb{A}(R(\epsilon))_0}(F) < 2\sum_{i=1}^n R_i \cdot |e_i| + \max\{0, \lambda(m+n)\}.$$

By (3) of Proposition 1.7 and Theorem 1.2, $Z \times T^n$ is a $L \times \mathbb{A}(R(\epsilon))_0$ -heavy subset. Since Corollary 1.6 implies that $Z \times T^n$ is $[L \times \mathbb{A}(R(\epsilon))_0]$ -heavy, by Definition 1.1,

$$\zeta_{[L\times\mathbb{A}(R(\epsilon))_0]}^{L\times\mathbb{A}(R(\epsilon))_0}(F) \ge 2\sum_{i=1}^n R_i \cdot |e_i| + \max\{0, \lambda(m+n)\}.$$

These two inequalities contradict. Since any Hamiltonian function $F: S^1 \times M \times I_{R(0)}^n \times T^n \to \mathbb{R}$ with compact support has support in $S^1 \times M \times I_{R(\epsilon)}^n \times T^n$ for some ϵ , we complete the proof of Theorem 1.23.

12. Flexibility result

To prove Proposition 1.24, we use the following proposition which is a chord-version of Proposition 3.3.2 of [BPS03].

Proposition 12.1. Let (N, ω) be an open symplectic manifold, Y_0, Y_1, Z subsets of N and α a homotopy class in $\pi_1(N, Y_0 \cup Y_1)$. Assume that Z is compact and there exists a Hamiltonian function $H: S^1 \times N \to \mathbb{R}$ with compact support satisfying the following conditions.

- (1) $Y_0 \cap \phi_H(Z) = \emptyset$.
- (2) $\mathsf{Ch}(H; Y_1, Y_0, \bar{\alpha}) = \emptyset.$

Then $C_{BEP}(N, Y_0, Y_1, Z, \alpha) = +\infty$.

Proof. Fix a positive number *K*. Since *Z* is compact and $Y_0 \cap \phi_H(Z) = \emptyset$, there exists an open neighborhood *U* of *Z* such that $Y_0 \cap \phi_H(U) = \emptyset$. Let $F: U \to \mathbb{R}$ be a Hamiltonian function with compact support such that $\inf_Z F + \sup_{S^1 \times N} \overline{H} \ge K$. Since $F \not\models \overline{H}(t, x) = F(t, x) + \overline{H}(t, (\phi_F^t)^{-1}(x))$ and $\inf_Z F + \sup_{S^1 \times N} \overline{H} \ge K$.

We claim that $\operatorname{Ch}(F \natural \overline{H}; Y_0, Y_1, \alpha) = \emptyset$. On the contrary, we assume that $\operatorname{Ch}(F \natural \overline{H}; Y_0, Y_1, \alpha) \neq \emptyset$, take an element $z: [0, 1] \to N$ of $\operatorname{Ch}(F \natural \overline{H}; Y_0, Y_1, \alpha)$ and set y = z(0). Since $y \in Y_0, \phi_{\overline{H}}(y) \in \phi_{\overline{H}}(Y_0) = (\phi_H)^{-1}(Y_0)$. Since $Y_0 \cap \phi_H(U) = \emptyset$, $(\phi_H)^{-1}(Y_0) \cap U = \emptyset$ and thus $\phi_{\overline{H}}(y) \notin U$. Hence, since F has support in $U, \phi_{F \natural \overline{H}}(y) = \phi_{\overline{H}}(y)$ and $\gamma_{\overline{H}}^y \in \operatorname{Ch}(\overline{H}; Y_0, Y_1, \alpha)$. $\gamma_{\overline{H}}^y \in \operatorname{Ch}(\overline{H}; Y_0, Y_1, \alpha)$ implies $\gamma_H^{(\phi_H)^{-1}(y)} \in \operatorname{Ch}(H; Y_1, Y_0, \overline{\alpha})$ and it contradicts with $\operatorname{Ch}(H; Y_1, Y_0, \overline{\alpha}) = \emptyset$. Thus we complete the proof of $\operatorname{Ch}(F \natural \overline{H}; Y_0, Y_1, \alpha) = \emptyset$.

Since *K* is any positive number, $C_{BEP}(N, Y_0, Y_1, Z, \alpha) = +\infty$.

Proof of Proposition 1.24. To use Proposition 12.1, we construct a Hamiltonian function $\hat{H}: S^{1} \times M \times \mathbb{A}(R) \to \mathbb{R}$ such that $(L \times \mathbb{A}(R)_{0}) \cap \phi_{\hat{H}}^{1}(Z \times \mathbb{A}(0)) = \emptyset$ and $Ch(\hat{H}; L \times \mathbb{A}(R)_{s}, L \times \mathbb{A}(R)_{0}, \bar{\alpha}) = \emptyset$. Fix a sufficiently small positive number ϵ . By the assumption, we can take a Hamiltonian function $H: S^{1} \times M \to \mathbb{R}$ with compact support such that $||H|| < E(Z; L) + \epsilon$ and $L \cap \phi_{H}^{1}(Z) = \emptyset$. Since $|s_{k}| \cdot R_{k} > E(Z; L)$ and ϵ is sufficiently small, we can take a function $\rho_{k}: (-R_{k}, R_{k}) \to \mathbb{R}$ with compact support and such that

- $\rho_k \equiv 1$ in a neighborhood of $\{0\}$,
- $|\dot{\rho}_k(x)| < |s_k| \cdot (E(Z;L) + \epsilon)^{-1}$ for any $x \in (-R_k, R_k)$.

For $i \neq k$, we take a function $\rho_i \in C_c^{\infty}(-R_i, R_i)$ with $\rho_i \equiv 1$ in a neighborhood of {0}. We define the Hamiltonian function $\hat{H}: S^1 \times M \times \mathbb{A}(R) \to \mathbb{R}$ by

$$\hat{H}(t, x, p, q) = \prod_{i} \rho_i(p_i) \cdot H(t, x).$$

Then

$$(X_{\hat{H}}^{t})_{(x,p,q)} = (\prod_{i} \rho_{i}(p_{i}) \cdot (X_{H}^{t})_{x}, 0, \dots, 0, \dot{\rho}_{1}(p_{1}) \cdot H(t, x), \dots, \dot{\rho}_{n}(p_{n}) \cdot H(t, x)).$$

Since $\rho_i \equiv 1$ in a neighborhood of {0}, $(L \times \mathbb{A}(0)) \cap \phi_{\hat{H}}^1(Z \times \mathbb{A}(0)) = \emptyset$. By the above comuputation of $X_{\hat{H}}, \phi_{\hat{H}}^1(Z \times \mathbb{A}(0)) \subset M \times \mathbb{A}(0)$ and thus $(L \times \mathbb{A}(R)_0) \cap \phi_{\hat{H}}^1(Z \times \mathbb{A}(0)) = \emptyset$. Since $|\dot{\rho}_k| < |s_k| \cdot (E(Z; L) + \epsilon)^{-1}$ and $\int_0^1 ||H_t||_{L^{\infty}} dt = ||H|| < E(Z; L) + \epsilon, \int_0^1 |\dot{\rho}_k(p_k)| \cdot |H(t, x)| dt = \dot{\rho}_k(p_k) \cdot \int_0^1 |H(t, x)| dt$ is smaller than $|s_k|$ and hence $Ch(\hat{H}; L \times \mathbb{A}(R)_s, L \times \mathbb{A}(R)_0, \bar{\alpha}_s) = \emptyset$. Thus Proposition 12.1 implies

$$C(M, L, Z, R, s) = C_{BEP}(M \times \mathbb{A}(R), L \times \mathbb{A}(R)_0, L \times \mathbb{A}(R)_s, Z \times \mathbb{A}(0), \alpha_s) = +\infty.$$

References

- [BC] P. Biran and O. Cornea, *Rigidity and uniruling for Lagrangian submanifolds*, Geom. Topol., **13**(5) (2009), 2881–2989.
- [BM] M. Borman and M. McLean, *Bounding Lagrangian widths via geodesic paths*, Comp. Math., **150** (12) (2014), 2143-2183.
- [BuEP] L. Buhovsky, M. Entov and L. Polterovich, *Poisson brackets and symplectic invariants*, Selecta Math. (N.S.), 1(18) (2012), 89-157.
- [BPS03] P. Biran, L. Polterovich and D. Salamon, *Propagation in Hamiltonian dynamics and relative symplectic homology*, Duke Math. J., **119**(1) (2003), 65-118.
- [EP03] M. Entov and L. Polterovich, Calabi quasimorphism and quantum homology, Int. Math. Res. Not., 30 (2003), 1635– 1676.
- [EP06] M. Entov and L. Polterovich, *Quasi-states and symplectic intersections*, Comment. Math. Helv., **81** (1) (2006), 75-99.
- [EP09] M. Entov and L. Polterovich, *Rigid subsets of symplectic manifolds*, Comp. Math., **145** (3) (2009), 773-826.
- [EPZ] M. Entov, L. Polterovich and F. Zapolsky, *Quasi-morphisms and the Poisson bracket*, Pure Appl. Math. Q., 3 (4) (2007), 1037–1055
- [FGS] U. Frauenfelder, V. Ginzburg and F. Schlenk, *Energy capacity inequalities via an action selector*, in Geometry, Spectral Theory, Groups, and Dynamics, Contemp. Math., Vol. 387 (Amer. Math. Soc., Providence, 2005), 129-152.
- [FS] U. Frauenfelder and F. Schlenk, Hamiltonian dynamics on convex symplectic manifolds, Israel J. Math., 159 (2007), 1-56.
- [HLS] V. Humilière, R. Leclercq and S. Seyfaddini, *Coisotropic rigidity and C⁰-symplectic geometry*, Duke Math. J., 164(4) (2015), 767-799.
- [I] K. Irie, Hofer-Zehnder capacity and a Hamiltonian circle action with noncontractible orbits, arXiv:1112.5247v1.
- [K14] M. Kawasaki, Superheavy Lagrangian immersions in surfaces, to appear in J. Symplectic Geom..
- [K16] M. Kawasaki, *Heavy subsets and non-contractible trajectories*, arXiv:1606.01964 (2016).
- [L] R. Leclercq, Spectral invariants in Lagrangian Floer theory, J. Mod. Dyn., 2 (2) (2008), 249-286.
- [LZ] R. Leclercq and F. Zapolsky, Spectral invariants for monotone Lagrangians, J. Topol. Anal., Online Ready, https://doi.org/10.1142/S1793525318500267.
- [LR] S. Lisi and A. Rieser, Coisotropic Hofer-Zehnder capacities and non-squeezing for relative embeddings, arXiv:1312.7334.
- [M] D. McDuff, Displacing Lagrangian toric fibers via probes. Low-dimensional and symplectic topology, from: "Lowdimensional and symplectic topology", (M Usher, editor), Proc. Sympos. Pure Math. 82, Amer. Math. Soc., Providence, RI, 2011, 131-160.
- [MVZ] A. Monzner, N. Vichery and F. Zapolsky, *Partial quasimorphisms and quasistates on cotangent bundles, and symplectic homogenization*, Journal of Modern Dynamics, **6**(2) (2012), 205-249.
- [Oh94] Y.-G. Oh, Relative Floer and quantum cohomology and the symplectic topology of Lagrangian submanifolds, from "Contact and symplectic geometry (Cambridge, 1994)", Publ. Newton Inst., 8, Cambridge Univ. Press, Cambridge, (1996), 201-267.
- [Oh05a] Y.-G. Oh, Spectral invariants, analysis of the Floer moduli space, and geometry of the Hamiltonian diffeomorphism group, Duke Math. J., **130**(2) (2005), 199-295.
- [Oh05b] Y.-G. Oh, Construction of spectral invariants of Hamiltonian paths on closed symplectic manifolds, in The Breadth of Symplectic and Poisson Geometry, eds. J. E. Marsden and T. Ratiu (Birkhäuser/Springer, 2005), 525-570.
- [Oh06] Y. -G. Oh, Lectures on Floer theory and spectral invariants of Hamiltonian flows, Morse Theoretic Methods in Nonlinear Analysis and in Symplectic Topology, NATO Sci. Ser. II Math. Phys. Chem., 217 (2006), 321-416.
- [Os] Y. Ostrover, A comparison of Hofer's metrics on Hamiltonian diffeomorphisms and Lagrangian submanifolds, Commun. Contemp. Math., 5 (5) (2003), 803-811.
- [PSS96] S. Piunikhin, D. Salamon and M. Schwarz, Symplectic Floer-Donaldson theory and quantum cohomology, Contact and symplectic geometry (Cambridge, 1994), Publ. Newton Inst, 8 (1996), 171-200.
- [P] L. Polterovich, *Quantum unsharpness and symplectic rigidity*, Lett. Math. Phys., **102**(3) (2012), 24-264.
- [PR] L. Polterovich and D. Rosen, Function theory on symplectic manifolds, CRM Monograph Series, 34. American Mathematical Society, Providence, RI, (2014).
- [Schw] M. Schwarz, On the action spectrum for closed symplectically aspherical manifolds, Pacific J. Math., **193**(2) (2000), 419–461.
- [Se] S. Seyfaddini, Spectral killers and Poisson bracket invariants, J. Mod. Dyn., 9 (2015), 51-66.
- [U10] M. Usher, *The sharp energy-capacity inequality*, Commun. Contemp. Math., **12**(3) (2010), 457-473.
- [Z] F. Zapolsky, The Lagrangian Floer-quantum-PSS package and canonical orientations in Floer theory, arXiv:1507.02253v2.

(Morimichi Kawasaki) Center for Geometry and Physics, Institute for Basic Science (IBS), Pohang 37673, Republic of Korea (previous), Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan (present) *E-mail address*: kawasaki@kurims.kyoto-u.ac.jp