

# NON-COMMUTATIVE STEIN INEQUALITY AND ITS APPLICATIONS

ALI TALEBI AND MOHAMMAD SAL MOSLEHIAN

ABSTRACT. The non-commutative Stein inequality asks whether there exists a constant  $C_{p,q}$  depending only on  $p, q$  such that

$$\left\| \left( \sum_n |\mathcal{E}_n(x_n)|^q \right)^{\frac{1}{q}} \right\|_p \leq C_{p,q} \left\| \left( \sum_n |x_n|^q \right)^{\frac{1}{q}} \right\|_p \quad (S_{p,q}),$$

for (positive) sequences  $(x_n)$  in  $L_p(\mathcal{M})$ . The validity of  $(S_{p,2})$  for  $1 < p < \infty$  and  $(S_{p,1})$  for  $1 \leq p < \infty$  are known. In this paper, we verify (i)  $(S_{p,\infty})$  for  $1 < p \leq \infty$ ; (ii)  $(S_{p,p})$  for  $1 \leq p < \infty$ ; (iii)  $(S_{p,q})$  for  $1 \leq q \leq 2$  and  $q < p < \infty$ . We also present some applications.

## 1. INTRODUCTION

Throughout this note,  $(\mathcal{M}, \tau)$  denotes a non-commutative probability space, that is, a von Neumann algebra  $\mathcal{M}$  equipped with a normal faithful finite trace  $\tau$  with  $\tau(1) = 1$ , where 1 stands for the identity of  $\mathcal{M}$ .

For  $1 \leq p < \infty$  the space  $L_p(\mathcal{M})$  is the completion of  $\mathcal{M}$  with respect to the  $p$ -norm  $\|x\|_p := \tau(|x|^p)^{1/p}$ , where  $|x| = (x^*x)^{1/2}$  is the absolute value of  $x$ . If  $p = \infty$ ,  $L_p(\mathcal{M})$  is  $\mathcal{M}$  itself with the operator norm. We denote by  $L_p^+(\mathcal{M})$  the positive cone in  $L_p(\mathcal{M})$  consisting of those elements, which are limits of sequences of positive elements in  $\mathcal{M}$ .

Assume that  $\mathcal{N}$  is a von Neumann subalgebra of  $\mathcal{M}$ . Then there exists a map  $\mathcal{E}_{\mathcal{N}} : \mathcal{M} \rightarrow \mathcal{N}$ , named the conditional expectation, which satisfies the following properties:

- (i)  $\mathcal{E}_{\mathcal{N}}$  is normal positive contractive projection from  $\mathcal{M}$  onto  $\mathcal{N}$ ;
- (ii)  $\mathcal{E}_{\mathcal{N}}(axb) = a\mathcal{E}_{\mathcal{N}}(x)b$  for every  $x \in \mathcal{M}$  and  $a, b \in \mathcal{N}$ ;
- (iii)  $\tau \circ \mathcal{E}_{\mathcal{N}} = \tau$ .

Moreover,  $\mathcal{E}_{\mathcal{N}}$  is the unique map verifying (ii), (iii). It is known that  $\mathcal{E}_{\mathcal{N}}$  can be extended to a contraction, denoted by the same  $\mathcal{E}_{\mathcal{N}}$ , from  $L_p(\mathcal{M})$  into  $L_p(\mathcal{N})$ .

By a filtration we mean an increasing sequence  $(\mathcal{M}_n)_{n \geq 0}$  of von Neumann subalgebras of  $\mathcal{M}$  such that  $\bigcup_{n \geq 0} \mathcal{M}_n$  generates  $\mathcal{M}$  in the  $w^*$ -topology. A sequence  $(x_n)_{n \geq 0}$  in  $L_p(\mathcal{M})$

---

2010 *Mathematics Subject Classification.* 46L53, 47A30, 60E15.

*Key words and phrases.* Stein inequality; non-commutative probability space; trace; conditional expectation.

is said to be adapted to  $(\mathcal{M}_n)_{n \geq 0}$  if  $x_n \in L_p(\mathcal{M}_n)$  ( $n \geq 0$ ). The reader is referred to [14, 16, 17, 18] for more information on non-commutative probability spaces.

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space,  $(\mathcal{F}_n)_{n=0}^\infty$  be a filtration on it and  $(X_n)_{n=1}^\infty$  be a stochastic process adapted to  $(\mathcal{F}_n)_{n=0}^\infty$ . The so-called Stein inequality

$$(1.1) \quad \left\| \left( \sum_{n=1}^{\infty} |\mathbb{E}_{\mathcal{F}_{n-1}} X_n|^q \right)^{\frac{1}{q}} \right\|_p \leq C_{p,q} \left\| \left( \sum_{n=1}^{\infty} |X_n|^q \right)^{\frac{1}{q}} \right\|_p \quad (S_{p,q})$$

was proved by Stein [15] for  $1 < p < \infty$  and  $q = 2$  and by Asmar and Montgomery-Smith [1] for any  $1 < p < \infty$  and  $1 \leq q \leq \infty$ . Using the duality between the martingale Hardy spaces and BMO-spaces, Lepingle [8] verified inequality (1.1) for adapted process with  $q = 2$ ,  $p = 1$  and  $C_{p,q} = 2$ . In a paper of Bourgain ([4], proposition 5) inequality (1.1) was obtained with a constant 3. It is evident that non-commutative probability theory is inspired by classical probability theory and quantum mechanics. Several mathematicians investigated and applied inequality (1.1) in the non-commutative setting.

In 1997, Pisier and Xu [11] proved a non-commutative Stein inequality for  $1 < p < \infty$  and  $q = 2$ . They indeed proved that there is a constant  $C_p$  depending on  $p$  such that for any finite sequence  $(x_n)_{n=1}^N$  in  $L_p(\mathcal{M})$ ,

$$\left\| \left( \sum_{n=1}^N |\mathcal{E}_n(x_n)|^2 \right)^{\frac{1}{2}} \right\|_p \leq C_p \left\| \left( \sum_{n=1}^N |x_n|^2 \right)^{\frac{1}{2}} \right\|_p,$$

See also [13] for another proof. In 2002, Junge [6] verified the non-commutative analogue of (1.1) for  $1 \leq p < \infty$  and  $q = 1$ .

In 2014, Qiu [12] employ a duality argument mentioned to show that if  $(\mathcal{M}_n)_{n=0}^N$  is an increasing filtration of von Neumann subalgebras and  $(x_n)_{n=1}^N$  is an adapted sequence in  $L_1(\mathcal{M})$ , then

$$\left\| \left( \sum_{n=1}^N |\mathcal{E}_{n-1}(x_n)|^2 \right)^{\frac{1}{2}} \right\|_1 \leq 2 \left\| \left( \sum_{n=1}^N |x_n|^2 \right)^{\frac{1}{2}} \right\|_1,$$

where  $\mathcal{E}_n$  denotes the conditional expectation with respect to  $\mathcal{M}_n$ .

The aim of this note is to obtain some versions of the non-commutative Stein inequality (1.1) for other values  $p$  and  $q$ . More precisely, we verify (i)  $(S_{p,\infty})$  for  $1 < p \leq \infty$  and  $q = \infty$  in Theorem 2.8; (ii)  $(S_{p,p})$  for  $1 \leq p < \infty$  in Proposition 2.11; (iii)  $(S_{p,q})$  for  $1 \leq q \leq 2$  and  $q < p < \infty$  in Corollary 2.15.

## 2. STEIN INEQUALITY IN NONCOMMUTATIVE SETTINGS

Suppose that  $(X, \mathcal{F}, \mu)$  is a measure space. The Banach space of all sequences  $f = (f_n)_{n \geq 1}$  of functions such that the norm

$$\|f\|_{p,q} := \left( \int_X \left( \sum_n |f_n(x)|^q \right)^{\frac{p}{q}} d\mu(x) \right)^{\frac{1}{p}}$$

is finite, is denoted by  $L_p(\ell_q)$ , in the case that  $1 \leq p, q < \infty$ . In the case that  $q = \infty$ , we set

$$\|f\|_{p,\infty} := \left( \int_X \left( \sup_n |f_n(x)| \right)^p d\mu(x) \right)^{\frac{1}{p}}$$

If  $p = \infty$ , we adopt the natural definition of essential supremum norm.

The behavior of  $L_p(\ell_q)$  is very similar to  $L_p(X, \mu)$ . For instance, the dual space of  $L_p(\ell_q)$  ( $1 \leq p, q < \infty$ ) is  $L_{p'}(\ell_{q'})$ , where  $\frac{1}{p} + \frac{1}{p'} = \frac{1}{q} + \frac{1}{q'} = 1$ ; cf. [15].

*Remark 2.1.* Difficulty of having a non-commutative analogue of the space  $L_p(\ell_\infty)$  is the lack of a non-commutative analogue of maximum. It is evident that there may be no maximum of two positive matrices. So we lead to the notion of  $L_p(\mathcal{M}, \ell_\infty)$ ; cf. [6].

We recall that the non-commutative spaces  $L_p(\mathcal{M}, \ell_1)$  and  $L_p(\mathcal{M}, \ell_\infty)$  playing an essential role in non-commutative analysis. The results presented in this section come from [2, 6, 18].

**Definition 2.2.** The space of all sequences  $x = (x_n)_{n \in \mathbb{N}}$  in  $L_p(\mathcal{M})$ , which can be decomposed as  $x_n = ay_nb$  for each  $n \in \mathbb{N}$ , where  $a, b \in L_{2p}(\mathcal{M})$  and  $y = (y_n)_{n \in \mathbb{N}}$  is in  $\mathcal{M}$ , is denoted by  $L_p(\mathcal{M}, \ell_\infty)$ . It is known that  $L_p(\mathcal{M}, \ell_\infty)$  is a Banach space equipped with the norm

$$\|x\|_{L_p(\mathcal{M}, \ell_\infty)} := \inf \left\{ \|a\|_{2p} \sup_{n \geq 1} \|y_n\|_\infty \|b\|_{2p} \right\},$$

where the infimum runs over all possible decompositions of  $x$  as above.

The element  $\|x\|_{L_p(\mathcal{M}, \ell_\infty)}$  is denoted by  $\|\sup_n^+ x_n\|_p$ . The reader should notice that  $\|\sup_n^+ x_n\|_p$  is a notation for  $\sup_n x_n$ . Thus we may consider the space  $L_p(\mathcal{M}, \ell_\infty)$  as a non-commutative analogue of the classical space  $L_p(\ell_\infty)$ .

• Let  $L_p(\mathcal{M}, \ell_1)$  be the space of all sequences  $x = (x_n)_{n \in \mathbb{N}}$  in  $L_p(\mathcal{M})$ , which admits a factorization of the form  $x_n = \sum_{k \in \mathbb{N}} c_{kn}^* d_{kn}$  ( $n \in \mathbb{N}$ ), where  $(c_{kn})_{k, n \geq 1}$  and  $(d_{kn})_{k, n \geq 1}$  are sequences in  $L_{2p}(\mathcal{M})$  such that  $\sum_{k, n} c_{kn}^* c_{kn}$  and  $\sum_{k, n} d_{kn}^* d_{kn}$  are in  $L_p(\mathcal{M})$ .

It is known that  $L_p(\mathcal{M}, \ell_1)$  equipped with the norm

$$\|x\|_{L_p(\mathcal{M}, \ell_1)} := \inf \left\{ \left\| \sum_{k, n} c_{kn}^* c_{kn} \right\|_p^{\frac{1}{2}} \left\| \sum_{k, n} d_{kn}^* d_{kn} \right\|_p^{\frac{1}{2}} \right\},$$

where the infimum runs over all possible decompositions of  $x$  as above, is a Banach space.

It is proved in [18] that the space  $L_p(\mathcal{M}, \ell_1)$  ( $1 \leq p < \infty$ ) is the predual of  $L_{p'}(\mathcal{M}, \ell_\infty)$  (where  $p'$  is the conjugate to  $p$ ) under the duality  $\langle x, y \rangle = \sum_{n \in \mathbb{N}} \tau(x_n y_n)$  for  $x \in L_p(\mathcal{M}, \ell_1)$  and  $y \in L_{p'}(\mathcal{M}, \ell_\infty)$ .

*Remark 2.3.* For any positive sequence  $x = (x_n)$  in  $L_p(\mathcal{M}, \ell_1)$  (i.e.,  $x_n \geq 0$  for all  $n$ ) we have  $\|x\|_{L_p(\mathcal{M}, \ell_1)} = \|\sum_{n \in \mathbb{N}} x_n\|_p$ . Therefore one may consider the space  $L_p(\mathcal{M}, \ell_1)$  as a generalization of the space  $L_p(\ell_1)$  in the non-commutative setting.

In the sequel, we present some useful properties of the spaces  $L_p(\mathcal{M}, \ell_1)$  and  $L_p(\mathcal{M}, \ell_\infty)$ .

**Theorem 2.4.** ([18, Proposition 2.12]) *Let  $1 \leq p \leq \infty$ .*

- (1) *Each element in the unit ball of  $L_p(\mathcal{M}, \ell_1)$  (resp.  $L_p(\mathcal{M}, \ell_\infty)$ ) is a sum of eight (resp. sixteen) positive elements in the same ball.*
- (2) *For any  $x \in L_p(\mathcal{M}, \ell_\infty)$  it holds that*

$$\|\sup_n^+ x_n\|_p = \sup \left\{ \sum_{n \in \mathbb{N}} \tau(x_n y_n) : y \in L_{p'}(\mathcal{M}, \ell_1) \text{ and } \|y\|_{L_{p'}(\mathcal{M}, \ell_1)} \leq 1 \right\}.$$

Moreover, if  $x$  is positive,

$$\|\sup_n^+ x_n\|_p = \sup \left\{ \sum_{n \in \mathbb{N}} \tau(x_n y_n) : y_n \in L_{p'}^+ \text{ and } \left\| \sum_{n \in \mathbb{N}} y_n \right\|_{p'} \leq 1 \right\}.$$

Some important spaces of sequences in  $L_p(\mathcal{M})$  can be formed via the row and column spaces, which are related to Burkholder–Gundy non-commutative inequalities.

**Definition 2.5.** Let  $1 \leq p \leq \infty$  and  $x = (x_n)_{n=1}^N$  be a finite sequence in  $L_p(\mathcal{M})$ . Set two norms

$$\|x\|_{L_p(\mathcal{M}; \ell_2^C)} := \left\| \left( \sum_{n=1}^N |x_n|^2 \right)^{\frac{1}{2}} \right\|_p \quad \text{and} \quad \|x\|_{L_p(\mathcal{M}; \ell_2^R)} := \left\| \left( \sum_{n=1}^N |x_n^*|^2 \right)^{\frac{1}{2}} \right\|_p.$$

The corresponding completion spaces are denoted by  $L_p(\mathcal{M}; \ell_2^C)$  and  $L_p(\mathcal{M}; \ell_2^R)$ , respectively.

*Remark 2.6.* Note that  $L_p(\mathcal{M}; \ell_2^C)$  (resp.  $L_p(\mathcal{M}; \ell_2^R)$ ) is isometric to the column (resp. row) subspace of  $L_p(\mathcal{M} \otimes B(\ell_2))$  via the following maps, where  $B(\ell_2)$  is the algebra of bounded linear maps on  $\ell_2$  with its usual trace  $Tr$  and  $\mathcal{M} \otimes B(\ell_2)$  denotes the von Neumann tensor product equipped with the (semifinite faithful) tensor trace  $\tau \otimes Tr$ :

$$x = (x_n)_{n \geq 0} \mapsto \begin{pmatrix} x_0 & 0 & \dots \\ x_1 & 0 & \dots \\ \vdots & \vdots & \end{pmatrix} \quad (\text{resp. } x = (x_n)_{n \geq 0} \mapsto \begin{pmatrix} x_0 & x_1 & \dots \\ 0 & 0 & \dots \\ \vdots & \vdots & \end{pmatrix}).$$

It is known (cf. [11, p. 670]) that  $L_p(\mathcal{M}; \ell_2^C)^* = L_q(\mathcal{M}; \ell_2^C)$  and  $L_p(\mathcal{M}; \ell_2^R)^* = L_q(\mathcal{M}; \ell_2^R)$ , whenever  $\frac{1}{p} + \frac{1}{q} = 1$  ( $1 \leq p < \infty$ ).

For notational convenience, we recall the space  $CR_p[L_p(\mathcal{M})]$ .

- For  $1 \leq p \leq \infty$ , let us define the space  $CR_p[L_p(\mathcal{M})]$  in two cases as follows:
  - If  $p \geq 2$ ,

$$CR_p[L_p(\mathcal{M})] = L_p(\mathcal{M}; \ell_2^C) \cap L_p(\mathcal{M}; \ell_2^R)$$

with the following norm:

$$\|(x_n)\|_{CR_p[L_p(\mathcal{M})]} = \max \left\{ \|(x_n)\|_{L_p(\mathcal{M}; \ell_2^C)}, \|(x_n)\|_{L_p(\mathcal{M}; \ell_2^R)} \right\}.$$

- If  $p < 2$ ,

$$CR_p[L_p(\mathcal{M})] = L_p(\mathcal{M}; \ell_2^C) + L_p(\mathcal{M}; \ell_2^R)$$

equipped with the following norm:

$$\|(x_n)\|_{CR_p[L_p(\mathcal{M})]} = \inf \left\{ \|(a_n)\|_{L_p(\mathcal{M}; \ell_2^C)} + \|(b_n)\|_{L_p(\mathcal{M}; \ell_2^R)} \right\},$$

where the infimum runs over all possible decompositions  $x_n = a_n + b_n$  with  $a_n$  and  $b_n$  in  $L_p(\mathcal{M})$ .

In [7] Junge and Xu defined an interesting complex interpolation space between  $L_p(\mathcal{M}, \ell_\infty)$  and  $L_p(\mathcal{M}, \ell_1)$  as  $L_p(\mathcal{M}, \ell_q) := [L_p(\mathcal{M}, \ell_\infty), L_p(\mathcal{M}, \ell_1)]_{\frac{1}{q}}$ . The reader is referred to [3] for interpolation theory and to [7] for some useful properties of the space  $L_p(\mathcal{M}, \ell_q)$ . Note that if  $\mathcal{M}$  is injective, this definition is a special case of Pisier's vector-valued non-commutative  $L_p$ -space theory [10].

In the sequel, let  $(\mathcal{M}_n)_{n=0}^\infty$  be an increasing sequence of von Neumann subalgebras of  $\mathcal{M}$  and  $\mathcal{E}_n$  denote the conditional expectation of  $\mathcal{M}$  with respect to  $\mathcal{M}_n$ . We consider its extension, denoted by the same  $\mathcal{E}_n$  from  $L_p(\mathcal{M})$  to  $L_p(\mathcal{M}_n)$ .

In [6], a noncommutative Doob's inequality is obtained by the following dual version of Doob's inequality:

**Lemma 2.7.** *Let  $1 \leq p < \infty$ . Then for every sequence of positive elements  $(x_n)$  in  $L_p(\mathcal{M})$*

$$(2.1) \quad \left\| \sum_{n \in \mathbb{N}} \mathcal{E}_n(x_n) \right\|_p \leq C_p \left\| \sum_{n \in \mathbb{N}} x_n \right\|_p, \quad (1 \leq p < \infty) \quad (DD_p)$$

in which  $C_p$  is a positive constant depending only on  $p$ .

By a duality argument, it is deduced from  $DD_p$  (see [6]) that

$$(2.2) \quad \left\| \sup_n |\mathcal{E}_n(x)| \right\|_p \leq C_{p'} \|x\|_p.$$

The next result is a non-commutative Stein inequality for the case when  $1 \leq p < \infty$  and  $q = \infty$ , which is a consequence of  $DD_p$ . We state it for the sake of completeness.

**Theorem 2.8** ( $S_{p,\infty}$  for  $1 < p \leq \infty$ ). *Let  $1 \leq p < \infty$  and  $x = (x_n)_{n \in \mathbb{N}}$  be an arbitrary sequence in  $L_p(\mathcal{M})$ . Then*

$$\left\| (\mathcal{E}_n(x_n))_{n \in \mathbb{N}} \right\|_{L_p(\mathcal{M}, \ell_\infty)} \leq C \left\| (x_n)_{n \in \mathbb{N}} \right\|_{L_p(\mathcal{M}, \ell_\infty)} \quad (S_{p,\infty})$$

for some positive constant  $C$  depending only on  $p$ .

*Proof.* It follows from Theorem 2.4(1) that any element in the unit ball of  $L_p(\mathcal{M}, \ell_\infty)$  is a finite linear combination of positive elements in the same ball. Hence it is enough to deduce the result for positive sequences. Using Theorem 2.4(2) we have

$$\begin{aligned} \left\| \sup_n^+ \mathcal{E}_n(x_n) \right\|_p &= \sup \left\{ \sum_{n \in \mathbb{N}} \tau(x_n \mathcal{E}_n(y_n)) : y_n \in L_{p'}^+ \text{ and } \left\| \sum_{n \in \mathbb{N}} y_n \right\|_{p'} \leq 1 \right\} \\ &= C_{p'} \sup \left\{ \sum_{n \in \mathbb{N}} \tau \left( x_n \frac{\mathcal{E}_n(y_n)}{C_{p'}} \right) : y_n \in L_{p'}^+ \text{ and } \left\| \sum_{n \in \mathbb{N}} y_n \right\|_{p'} \leq 1 \right\} \\ &\leq C_{p'} \sup \left\{ \sum_{n \in \mathbb{N}} \tau(x_n z_n) : z_n \in L_{p'}^+ \text{ and } \left\| \sum_{n \in \mathbb{N}} z_n \right\|_{p'} \leq 1 \right\} \\ &= C_{p'} \left\| \sup_n^+ x_n \right\|_p \quad (\text{by Theorem 2.4(2)}) \end{aligned}$$

To get the inequality above, we note that  $\mathcal{E}_n$  is a positive map for any  $n$  and use the inequality  $DD_{p'}$  (2.1).  $\square$

*Remark 2.9.* If we put  $x_n = x$  for all  $n$  in Theorem 2.8, then  $(S_{p,\infty})$  implies (2.2).

**Corollary 2.10.** *Let  $1 \leq p < \infty$  and  $x = (f_n)_{n \in \mathbb{N}}$  be an arbitrary sequence in  $L_p(\Omega, \mathcal{F}, \mathbb{P})$ . Then*

$$\left\| \sup_{n \in \mathbb{N}} |\mathbb{E}_n(f_n)| \right\|_p \leq C \left\| \sup_{n \in \mathbb{N}} |f_n| \right\|_p$$

for some positive constant  $C$  depending only on  $p$ .

The next Proposition provides a non-commutative Stein inequality for the case when  $1 \leq p = q < \infty$ . We should notify that the case  $p = q = \infty$  is proved in Theorem 2.8.

**Proposition 2.11** ( $S_{q,q}$  for  $1 \leq q < \infty$ ). *Let  $(x_n)_{n \in \mathbb{N}}$  be an arbitrary sequence in  $L_q(\mathcal{M})$ . Then*

$$\left\| \left( \sum_{n \in \mathbb{N}} |\mathcal{E}_{n-1}(x_n)|^q \right)^{\frac{1}{q}} \right\|_q \leq \left\| \left( \sum_{n \in \mathbb{N}} |x_n|^q \right)^{\frac{1}{q}} \right\|_q \quad (S_{q,q})$$

for every real number  $q \geq 1$ .

*Proof.* We show the desired inequality by some properties of trace and conditional expectation as follows.

$$\begin{aligned}
& \left\| \left( \sum_{n \in \mathbb{N}} |\mathcal{E}_{n-1}(x_n)|^q \right)^{\frac{1}{q}} \right\|_q^q \\
&= \tau \left( \sum_{n=1}^{\infty} |\mathcal{E}_{n-1}(x_n)|^q \right) = \tau \left( \sup_{N \in \mathbb{N}} \left( \sum_{n=1}^N |\mathcal{E}_{n-1}(x_n)|^q \right) \right) \\
&= \sup_{N \in \mathbb{N}} \sum_{n=1}^N \tau (|\mathcal{E}_{n-1}(x_n)|^q) \quad (\text{by the normality of } \tau) \\
&= \sup_{N \in \mathbb{N}} \sum_{n=1}^N \|\mathcal{E}_{n-1}(x_n)\|_q^q \leq \sup_{N \in \mathbb{N}} \sum_{n=1}^N \|x_n\|_q^q \quad (\text{by the contractivity of } \mathcal{E}_n) \\
&= \sup_{N \in \mathbb{N}} \tau \left( \sum_{n=1}^N |x_n|^q \right) = \tau \left( \sum_{n=1}^{\infty} |x_n|^q \right) \quad (\text{again by the normality of } \tau) \\
&= \left\| \left( \sum_{n \in \mathbb{N}} |x_n|^q \right)^{\frac{1}{q}} \right\|_q^q.
\end{aligned}$$

□

In [11], the non-commutative Stein inequality  $S_{p,2}$  is shown for  $1 < p < \infty$  by applying Burkholder–Gundy inequality. Next, Qiu [12] obtained inequality  $S_{1,2}$  for adapted sequences. The next corollary provides Stein inequality in the spaces  $CR_p[L_P(\mathcal{M})]$ .

**Corollary 2.12.** *Let  $1 < p < \infty$  and  $x = (x_n)_{n \in \mathbb{N}}$  be an adapted sequence in  $L_p(\mathcal{M})$ . Then*

$$\|(\mathcal{E}_{n-1}(x_n))_n\|_{CR_p[L_P(\mathcal{M})]} \leq C_p \|(x_n)_n\|_{CR_p[L_P(\mathcal{M})]},$$

where  $C_p$  is the constant appeared in  $S_{p,2}$ .

*Proof.* Suppose that  $p \geq 2$ . By replacing  $x_n$  by  $x_n^*$  in inequality  $S_{p,2}$  we get

$$\begin{aligned}
\|(\mathcal{E}_{n-1}(x_n))_n\|_{L_p(\mathcal{M}; \ell_2^R)} &= \left\| \left( \sum_{n \in \mathbb{N}} |\mathcal{E}_{n-1}(x_n^*)|^2 \right)^{\frac{1}{2}} \right\|_p = \left\| \left( \sum_{n \in \mathbb{N}} |\mathcal{E}_{n-1}(x_n^*)|^2 \right)^{\frac{1}{2}} \right\|_p \\
&\quad (\text{by the } * \text{-preserving property of } \mathcal{E}_n) \\
&\leq C_p \left\| \left( \sum_{n \in \mathbb{N}} |x_n^*|^2 \right)^{\frac{1}{2}} \right\|_p \quad (\text{by } S_{p,2}) \\
&= C_p \|(x_n)_n\|_{L_p(\mathcal{M}; \ell_2^R)}.
\end{aligned}$$

Employing again inequality  $S_{p,2}$ , we infer that

$$\|(\mathcal{E}_{n-1}(x_n))_n\|_{CR_p[L_p(\mathcal{M})]} \leq C_p \|(x_n)_n\|_{CR_p[L_p(\mathcal{M})]}.$$

Now let  $1 < p < 2$ . Assume that  $x_n = a_n + b_n$  with  $a_n$  and  $b_n$  in  $L_p(\mathcal{M})$ . Then  $\mathcal{E}_{n-1}(x_n) = \mathcal{E}_n(a_n) + \mathcal{E}_n(b_n)$  is a factorization of  $\mathcal{E}_{n-1}(x_n)$  for all  $n \in \mathbb{N}$ . By applying inequality  $S_{p,2}$  to  $(a_n)_n$  and  $(b_n)_n$ , we have

$$\|(\mathcal{E}_{n-1}(a_n))_n\|_{L_p(\mathcal{M}; \ell_2^C)} \leq C_p \|(a_n)_n\|_{L_p(\mathcal{M}; \ell_2^C)}$$

and

$$\|(\mathcal{E}_{n-1}(b_n))_n\|_{L_p(\mathcal{M}; \ell_2^R)} \leq C_p \|(b_n)_n\|_{L_p(\mathcal{M}; \ell_2^R)}.$$

Hence

$$\begin{aligned} \|(\mathcal{E}_{n-1}(x_n))_n\|_{CR_p[L_p(\mathcal{M})]} &= \inf \left\{ \|(c_n)_n\|_{L_p(\mathcal{M}; \ell_2^C)} + \|(d_n)_n\|_{L_p(\mathcal{M}; \ell_2^R)} \right\} \\ &\leq \|(\mathcal{E}_{n-1}(a_n))_n\|_{L_p(\mathcal{M}; \ell_2^C)} + \|(\mathcal{E}_{n-1}(b_n))_n\|_{L_p(\mathcal{M}; \ell_2^R)} \\ &\leq C_p \left( \|(a_n)_n\|_{L_p(\mathcal{M}; \ell_2^C)} + \|(b_n)_n\|_{L_p(\mathcal{M}; \ell_2^R)} \right), \end{aligned}$$

where the infimum runs over all possible decompositions  $(\mathcal{E}_{n-1}(x_n))_n = (c_n)_n + (d_n)_n$  with  $(c_n)_n \in L_p(\mathcal{M}; \ell_2^C)$  and  $(d_n)_n \in L_p(\mathcal{M}; \ell_2^R)$ .

Whence by taking infimum over all decompositions as above, we conclude the required inequality.  $\square$

*Remark 2.13.* Due to the dual version of Doob's inequality and noting that each element in the unit ball of  $L_p(\mathcal{M}, \ell_1)$  is a sum of eight positive elements in the same ball (see 2.4(1)), we deduce that the linear map  $\phi : L_p(\mathcal{M}; \ell_1) \rightarrow L_p(\mathcal{M}; \ell_1)$  with  $\phi((x_n)) = (\mathcal{E}_n(x_n))$  is bounded. However, Junge obtained it with the bound  $C_p$  as same as the constant, which is obtained from inequality  $DD_p$  (2.1). Moreover,  $\phi$  is a bounded linear map on  $L_p(\mathcal{M}; \ell_\infty)$  by Theorem 2.8. Applying interpolation theorem one may deduce that  $\phi : L_p(\mathcal{M}; \ell_q) \rightarrow L_p(\mathcal{M}; \ell_q)$  is bounded. It seems that there is a problem to get the general Stein inequality. The problem is whether the equality

$$\|(x_n)\|_{L_p(\mathcal{M}; \ell_q)} = \left\| \left( \sum_n x_n^q \right)^{\frac{1}{q}} \right\|_p$$

holds for any positive sequence  $(x_n)$  or not. However we establish inequality  $S_{p,q}$  for  $1 \leq q \leq 2$  and  $p \geq q$  in the following theorem.

**Theorem 2.14.** *Suppose that  $1 \leq q \leq 2$ ,  $p \geq q$ ,  $(y_n)$  is a sequence of isometries in  $\mathcal{M}$ , and  $(x_n)$  in  $L_p^+(\mathcal{M})$  is an arbitrary positive sequence. Then*

$$\left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n(y_n^* x_n y_n))^q \right)^{\frac{1}{q}} \right\|_p \leq C \left\| \left( \sum_{n \in \mathbb{N}} y_n^* x_n^q y_n \right)^{\frac{1}{q}} \right\|_p,$$



where  $C$  is a positive constant depending only on  $p$  and  $q$ .

*Proof.* We have

$$\begin{aligned}
\left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n (y_n^* x_n y_n))^q \right)^{\frac{1}{q}} \right\|_p &\leq \left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n ((y_n^* x_n y_n)^q)) \right)^{\frac{1}{q}} \right\|_p \\
&\quad \text{(by the Choi–Davis–Jensen inequality (see e.g. [5, 9])} \\
&\quad \text{and the operator monotonicity of } t^r \text{ (} 0 < r \leq 1 \text{))} \\
&\leq \left\| \sum_{n \in \mathbb{N}} \mathcal{E}_n (y_n^* x_n^q y_n) \right\|_{\frac{p}{q}}^{\frac{q}{p^2}} \\
&\quad \text{(by the Jensen operator inequality)} \\
&\leq C_{\frac{p}{q}} \left\| \sum_{n \in \mathbb{N}} y_n^* x_n^q y_n \right\|_{\frac{p}{q}}^{\frac{q}{p^2}} \quad \text{(by } DD_{\frac{p}{q}} \text{ (2.1))} \\
&= C_{\frac{p}{q}} \left\| \left( \sum_{n \in \mathbb{N}} y_n^* x_n^q y_n \right)^{\frac{1}{q}} \right\|_p.
\end{aligned}$$

□

**Corollary 2.15** ( $S_{p,q}$  for  $1 \leq q \leq 2$  and  $q < p < \infty$ ). *Suppose that  $1 \leq q \leq 2$ ,  $q < p < \infty$  and  $(x_n)$  in  $L_p^+(\mathcal{M})$  is an arbitrary positive sequence. Then*

$$(2.3) \quad \left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n (x_n))^q \right)^{\frac{1}{q}} \right\|_p \leq C \left\| \left( \sum_{n \in \mathbb{N}} (x_n)^q \right)^{\frac{1}{q}} \right\|_p,$$

where  $C$  is a positive constant depending only on  $p$  and  $q$ .

*Proof.* Apply Theorem 2.14 with  $y_n = 1$  for all  $n \in \mathbb{N}$ . □

The following results give some applications of  $S_{p,q}$  (2.3), which is interesting on its own right.

**Corollary 2.16** (Semi-noncommutative case). *Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space, and  $(\mathcal{F}_n)_{n=0}^\infty$  be a filtration on it. If  $1 \leq q \leq 2$ ,  $q < p < \infty$ , then there exists a positive constant  $C$  such that for every positive stochastic process  $(f_n)_{n=1}^\infty$  with values in  $L_p(\mathcal{M})$*

$$\left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n (f_n))^q \right)^{\frac{1}{q}} \right\|_p \leq C \left\| \left( \sum_{n \in \mathbb{N}} (f_n)^q \right)^{\frac{1}{q}} \right\|_p,$$

in which  $\mathcal{E}_n = \mathbb{E}_n \otimes id_{L_p(\mathfrak{M})}$ , where  $\mathbb{E}_n$  is the usual conditional expectation with respect to  $\mathcal{F}_n$ .

*Proof.* Set  $\mathfrak{N} := L_\infty(\Omega, \mathcal{F}, \mathbb{P}) \otimes \mathfrak{M}$  and  $\nu := \mathbb{E} \otimes \tau$ , and apply (2.3). □

**Corollary 2.17.** *If  $1 \leq q \leq 2$ ,  $q < p < \infty$ , then there exists a positive constant  $C$ , depending only on  $p$  and  $q$ , such that*

$$\left\| \left( \sum_{n \in \mathbb{N}} (\mathcal{E}_n(r_n))^q \right)^{\frac{1}{q}} \right\|_p \leq C$$

for every sequence of mutually orthogonal projections  $(r_n)_{n \geq 1}$  in  $L_p(\mathcal{M})$ .

**Corollary 2.18** (Matrix algebra case). *If  $1 \leq q \leq 2$ ,  $q < p < \infty$ , then there exists a positive constant  $C$  such that for every sequence of positive matrices  $(X_n)_n$  in  $S_p$ , the Schatten  $p$ -class on  $l_2$ , equipped with the usual trace it holds that*

$$\left\| \left( \sum_{n \in \mathbb{N}} (E_n X_n E_n)^q \right)^{\frac{1}{q}} \right\|_p \leq C \left\| \left( \sum_{n \in \mathbb{N}} (X_n)^q \right)^{\frac{1}{q}} \right\|_p,$$

where  $E_n$  is the operator projecting a sequence in  $l_2$  into its  $n$  first coordinates.

*Proof.* Regarding  $B(l_2^n)$  as a subalgebra of  $B(l_2)$ , we have an increasing filtration  $(B(l_2^n))_{n \in \mathbb{N}}$ . The corresponding conditional expectation from  $B(l_2)$  onto  $B(l_2^n)$  is  $\mathcal{E}_n$  such that  $\mathcal{E}_n(X) = E_n X E_n$  ( $X \in B(l_2)$ ), which extends to a contractive projection from  $S_p$  onto  $S_p^n$  (see [18]). Now, apply (2.3).  $\square$

## REFERENCES

- [1] N. Asmar and S. Montgomery-Smith, *Littlewood–Paley theory on solenoids*, Colloq. Math. **65** (1993), 69–82.
- [2] T. N. Bekjan, Z. Chen and A. Osekowski, *Noncommutative maximal inequalities associated with convex functions*, Trans. Amer. Math. Soc. **369** (2017), no. 1, 409–427.
- [3] J. Bergh and J. Lofstrom, *Interpolation spaces*, Springer-Verlag, Berlin, 1976.
- [4] J. Bourgain, *Embedding  $L^1$  in  $L^1/H^1$* , Trans. Amer. Math. Soc. **278** (1983), no. 2, 689–702.
- [5] M. D. Choi, *Some assorted inequalities for positive linear maps on  $C^*$ -algebras*, J. Operator Theory, **4** (1980), 271–285.
- [6] M. Junge, *Doob’s inequality for non-commutative martingales*, J. Reine Angew. Math. **549** (2002), 149–190.
- [7] M. Junge and Q. Xu, *Noncommutative Burkholder/Rosenthal inequalities II: Applications*, Israel J. Math. **167** (2008), 227–282.
- [8] D. Lepingle, *Une inégalité de martingales*, (French) Séminaire de Probabilités, XII (Univ. Strasbourg, Strasbourg, 1976/1977), pp. 134–137, Lecture Notes in Math., 649 (1978), Springer, Berlin.
- [9] M. S. Moslehian and H. Najafi, *Around operator monotone functions*, Integral Equations Operator Theory **71** (2011), no. 4, 575–582.
- [10] G. Pisier, *Non-commutative vector valued  $L_p$ -spaces and completely  $p$ -summing maps*, Astérisque No. 247, 1998.
- [11] G. Pisier and Q. Xu, *Non-commutative martingale inequalities*, Comm. Math. Phys. **189** (1997), no. 3, 667–698.
- [12] Y. Qiu, *A non-commutative version of Lepingle–Yor martingale inequality*, Statist. Probab. Lett. **91** (2014), 52–54.

- [13] N. Randrianantoanina, *Non-commutative martingale transforms*, J. Funct. Anal. **194** (2002), no. 1, 181–212.
- [14] Gh. Sadeghi and M. S. Moslehian, *Noncommutative martingale concentration inequalities*, Illinois J. Math. **58** (2014), no. 2, 561–575.
- [15] E. M. Stein, *Topics in harmonic analysis related to the Littlewood-Paley theory*, Annals of Mathematics Studies, No. 63 Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo 1970.
- [16] A. Talebi, M. S. Moslehian and Gh. Sadeghi, *Etemadi and Kolmogorov inequalities in noncommutative probability spaces*, Michigan Math. J. (to appear), arXiv:1709.08044.
- [17] A. Talebi, M. S. Moslehian and Gh. Sadeghi, *Noncommutative Blackwell–Ross martingale inequality*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. (to appear), arXiv:1705.07122.
- [18] Q. Xu, *Operator spaces and non-commutative  $L_p$* , Lectures in the Summer School on Banach spaces and Operator spaces, Nankai University China, 2007.

DEPARTMENT OF PURE MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P.O. BOX 1159, MASHHAD 91775, IRAN.

*E-mail address:* alitalebimath@yahoo.com

*E-mail address:* moslehian@um.ac.ir and moslehian@member.ams.org