Kevin Keating Department of Mathematics University of Florida Gainesville, FL 32611 USA

keating@ufl.edu

November 5, 2018

Abstract

Let K be a local field and let L/K be a totally ramified Galois extension of degree p^n . Being semistable [2] and possessing a Galois scaffold [3] are two conditions which facilitate the computation of the additive Galois module structure of L/K. In this note we show that L/K is semistable if and only if L/K has a Galois scaffold. We also give sufficient conditions in terms of Galois scaffolds for the extension L/K to be stable.

1 Introduction

Let K be a local field whose residue field is perfect with characteristic p. Let L/K be a finite Galois extension and set G = Gal(L/K). Then L is free of rank 1 over K[G]by the normal basis theorem. Let \mathcal{O}_K , \mathcal{O}_L be the integer rings of K, L, and let \mathcal{M}_K , \mathcal{M}_L be the maximal ideals of these rings. For a fractional ideal \mathcal{M}_L^h of L define the associated order

$$\mathfrak{A}_h = \{ \gamma \in K[G] : \gamma(\mathcal{M}_L^h) \subset \mathcal{M}_L^h \}.$$

of \mathcal{M}_{L}^{h} in K[G]. Suppose \mathcal{M}_{L}^{h} is free over some order \mathcal{B} of K[G]. Then $\mathcal{B} = \mathfrak{A}_{h}$, and \mathcal{M}_{L}^{h} is free of rank 1 over \mathfrak{A}_{h} . If L/K is at most tamely ramified then $\mathfrak{A}_{h} = \mathcal{O}_{K}[G]$ and \mathcal{M}_{L}^{h} is free of rank 1 over $\mathcal{O}_{K}[G]$ for all $h \in \mathbb{Z}$ [6, Th. 1].

On the other hand, if L/K is wildly ramified then in most cases \mathfrak{A}_h properly contains $\mathcal{O}_K[G]$, and hence \mathcal{M}_L^h is not free over $\mathcal{O}_K[G]$. The problem of determining when \mathcal{M}_L^h is free over \mathfrak{A}_h for a totally ramified extension L/K of degree p^n appears to be quite difficult. Two classes of extensions for which a partial answer can be obtained are semistable extensions [2] and extensions with a Galois scaffold [3]. In this note we consider the relation between semistable extensions and Galois scaffolds. In particular, we show that L/K admits a Galois scaffold if and only if L/K is semistable.

2 Galois Scaffolds

In this section we give the definition of a Galois scaffold. We also give sufficient conditions for an extension to admit a Galois scaffold. This criterion will be used in section 4.

Let L/K be a totally ramified Galois extension of degree p^n and set G = Gal(L/K). Let $b_1 \leq b_2 \leq \cdots \leq b_n$ be the lower ramification breaks of L/K, counted with multiplicity. Assume that $p \nmid b_i$ for $1 \leq i \leq n$. Set $\mathbb{S}_{p^n} = \{0, 1, \ldots, p^n - 1\}$ and write $s \in \mathbb{S}$ in base p as

$$s = s_{(0)}p^0 + s_{(1)}p^1 + \dots + s_{(n-1)}p^{n-1}$$

Define a partial order on \mathbb{S}_{p^n} by $s \leq t$ if $s_{(i)} \leq t_{(i)}$ for $0 \leq i \leq n-1$. Then by Lucas's theorem we have $s \leq t$ if and only if $p \nmid {t \choose s}$. Define $\mathfrak{b} : \mathbb{S}_{p^n} \to \mathbb{Z}$ by

$$\mathfrak{b}(s) = s_{(0)}p^0b_n + s_{(1)}p^1b_{n-1} + \dots + s_{(n-1)}p^{n-1}b_1.$$

Let $r : \mathbb{Z} \to \mathbb{S}_{p^n}$ be the function which maps $a \in \mathbb{Z}$ onto its least nonnegative residue modulo p^n . The function $r \circ (-\mathfrak{b}) : \mathbb{S}_{p^n} \to \mathbb{S}_{p^n}$ is a bijection since $p \nmid b_i$. Therefore we may define $\mathfrak{a} : \mathbb{S}_{p^n} \to \mathbb{S}_{p^n}$ to be the inverse of $r \circ (-\mathfrak{b})$. Let $v_L : L \to \mathbb{Z} \cup \{\infty\}$ denote the normalized valuation on L.

Definition 2.1 ([3], Definition 2.6). Let $\mathfrak{c} \geq 1$. A Galois scaffold $(\{\Psi_i\}, \{\lambda_t\})$ for L/K with precision \mathfrak{c} consists of elements $\Psi_i \in K[G]$ for $1 \leq i \leq n$ and $\lambda_t \in L$ for all $t \in \mathbb{Z}$ such that the following hold:

- (i) $v_L(\lambda_t) = t$ for all $t \in \mathbb{Z}$.
- (ii) $\lambda_{t_1} \lambda_{t_2}^{-1} \in K$ whenever $t_1 \equiv t_2 \pmod{p^n}$.
- (iii) $\Psi_i(1) = 0$ for $1 \le i \le n$.
- (iv) For $1 \leq i \leq n$ and $t \in \mathbb{Z}$ there exists $u_{it} \in \mathcal{O}_K^{\times}$ such that the following congruence modulo $\lambda_{t+p^{n-i}b_i}\mathcal{M}_L^{\mathfrak{c}}$ holds:

$$\Psi_i(\lambda_t) \equiv \begin{cases} u_{it}\lambda_{t+p^{n-i}b_i} & \text{if } \mathfrak{a}(r(t))_{(n-i)} \ge 1, \\ 0 & \text{if } \mathfrak{a}(r(t))_{(n-i)} = 0. \end{cases}$$

A Galois scaffold for L/K with infinite precision consists of the above data with the congruence in (iv) replaced by equality.

In [3, Th. 3.1] sufficient conditions are given for an ideal \mathcal{M}_L^h in an extension L/K with a Galois scaffold to be free over its associated order \mathfrak{A}_h . When the precision of the scaffold is sufficiently large these sufficient conditions are shown to be necessary as well.

Let $(\{\Psi_i\}, \{\lambda_t\})$ be a Galois scaffold for L/K. For $s \in \mathbb{S}_{p^n}$ set

$$\Psi^{(s)} = \Psi_n^{s_{(0)}} \Psi_{n-1}^{s_{(1)}} \dots \Psi_2^{s_{(n-2)}} \Psi_1^{s_{(n-1)}}$$

As explained on page 974 of [3], it follows from the definition of a Galois scaffold that for every $t \in \mathbb{Z}$ there is $U_{st} \in \mathcal{O}_K^{\times}$ such that the following holds modulo $\lambda_{t+\mathfrak{b}(s)}\mathcal{M}_L^{\mathfrak{c}}$:

$$\Psi^{(s)}(\lambda_t) \equiv \begin{cases} U_{st}\lambda_{t+\mathfrak{b}(s)} & \text{if } s \preceq \mathfrak{a}(r(t)), \\ 0 & \text{if } s \not\preceq \mathfrak{a}(r(t)). \end{cases}$$
(2.1)

Proposition 2.2. Let $\Phi_1, \ldots, \Phi_n \in K[G]$ and let $\{\lambda_t : t \in \mathbb{Z}\}$ be a subset of L satisfying conditions (i) and (ii) in Definition 2.1. Suppose that for all $t \in \mathbb{Z}$ and $1 \leq i \leq n$ we have

$$v_L(\Phi_i(\lambda_t)) = t + p^{n-i}b_i \text{ if } \mathfrak{a}(r(t))_{(n-i)} \ge 1,$$
 (2.2)

$$v_L(\Phi_i(\lambda_t)) > t + p^{n-i}b_i \text{ if } \mathfrak{a}(r(t))_{(n-i)} = 0.$$
 (2.3)

Then there are $\Psi_1, \ldots, \Psi_n \in K[G]$ such that $(\{\Psi_i\}, \{\lambda_t\})$ is a Galois scaffold for L/K with precision $\mathfrak{c} = 1$.

Proof. For $1 \leq i \leq n$ set $\Psi_i = \Phi_i - \Phi_i(1) \cdot \mathrm{id}_L$. It follows from (2.2) and (2.3) that $v_L(\Phi_i(1)) > p^{n-i}b_i$. Hence for all $t \in \mathbb{Z}$ we have

$$\Psi_i(\lambda_t) \equiv \Phi_i(\lambda_t) \pmod{\lambda_{t+p^{n-i}b_i}\mathcal{M}_L},$$

so (2.2) and (2.3) hold with Φ_i replaced by Ψ_i . If $\mathfrak{a}(t)_{(n-i)} \geq 1$ then there is $u_{it} \in \mathcal{O}_K^{\times}$ such that

$$\Psi_i(\lambda_t) \equiv u_{it}\lambda_{t+p^{n-i}b_i} \pmod{\lambda_{t+p^{n-i}b_i}} \mathcal{M}_L.$$

Therefore $(\{\Psi_i\}, \{\lambda_t\})$ is a Galois scaffold for L/K with precision 1.

3 Stable and semistable extensions

In this section we outline Bondarko's theory of stable and semistable extensions [1, 2]. We also present some interpretations and refinements of Bondarko's work which will be useful in the next section.

Let L/K be a totally ramified Galois extension of degree p^n and let $T = \sum_{\sigma \in G} \sigma$ denote the trace element of K[G]. Define $\phi : L \otimes_K L \to L[G]$ by

$$\phi(a \otimes b) = aTb = \sum_{\sigma \in G} a\sigma(b)\sigma.$$

By Remark 1.2 of [1], ϕ is an isomorphism of vector spaces over K. The following is proved in Proposition 1.6.1 of [1]:

Proposition 3.1. Let $\alpha, \beta \in L \otimes_K L$ and write $\phi(\alpha) = \sum_{\sigma \in G} a_\sigma \sigma$ and $\phi(\beta) = \sum_{\sigma \in G} b_\sigma \sigma$. Then $\phi(\alpha\beta) = \sum_{\sigma \in G} a_\sigma b_\sigma \sigma$. Let $H = \langle (p^n, -p^n) \rangle$ be the subgroup of $\mathbb{Z} \times \mathbb{Z}$ generated by the element $(p^n, -p^n)$. Then

$$\mathcal{F} = \{(a, b) \in \mathbb{Z} \times \mathbb{Z} : 0 \le b < p^n\}$$

is a set of coset representatives for $(\mathbb{Z} \times \mathbb{Z})/H$. For $(a, b) \in \mathbb{Z} \times \mathbb{Z}$ write [a, b] for the coset (a, b) + H. We define a partial order on $(\mathbb{Z} \times \mathbb{Z})/H$ by $[a, b] \leq [c, d]$ if and only if there is $(c', d') \in [c, d]$ with $a \leq c'$ and $b \leq d'$. We easily see that $[a, b] \not\leq [c, d]$ if and only if $[c + 1, d - p^n + 1] \leq [a, b]$ (see [2, (16)]).

Fix a uniformizer π_L for L and let \mathcal{T} be the set of Teichmüller representatives for K. Let $\beta \in L \otimes_K L$. Then there are uniquely determined $c_j \in L$ and $a_{ij} \in \mathcal{T}$ such that

$$\beta = \sum_{j=0}^{p^n-1} c_j \otimes \pi_L^j = \sum_{(i,j)\in\mathcal{F}} a_{ij} \pi_L^i \otimes \pi_L^j.$$

Set

$$R(\beta) = \{ [i, j] \in (\mathbb{Z} \times \mathbb{Z}) / H : (i, j) \in \mathcal{F}, \ a_{ij} \neq 0 \}$$

Definition 3.2. Define the diagram of β to be

$$D(\beta) = \{ [a, b] \in (\mathbb{Z} \times \mathbb{Z}) / H : [i, j] \le [a, b] \text{ for some } [i, j] \in R(\beta) \}.$$

It is observed in Remark 2.4.3 of [2] that while $R(\beta)$ can depend on the choice of uniformizer π_L for L, $D(\beta)$ does not depend on this choice. Let $G(\beta)$ denote the set of minimal elements of $D(\beta)$ with respect to \leq . Then $G(\beta)$ is also the set of minimal elements of $R(\beta)$.

Let $\delta_{L/K} = \mathcal{M}_L^d$ denote the different of L/K and set $i_0 = d - p^n + 1$. Then i_0 is the 0th index of inseparability of L/K [4, Prop. 3.18], and $\mathfrak{b}(p^n - 1) = i_0$ [5, IV, §1, Prop. 4]. The following is proved in Proposition 2.4.2 of [2]:

Theorem 3.3. Let $\xi \in K[G] \setminus \{0\}$, let $\beta \in L \otimes_K L$ satisfy $\xi = \phi(\beta)$, and let $a, b \in \mathbb{Z}$. Then the following are equivalent:

- (a) $[a,b] \in G(\beta)$.
- (b) For all $y \in L$ with $v_L(y) = -b i_0$ we have $v_L(\xi(y)) = a$.

For $\xi \in K[G]$ with $\xi \neq 0$ we define $f_{\xi} : \mathbb{Z} \to \mathbb{Z}$ by

$$f_{\xi}(a) = \min\{v_L(\xi(y)) : y \in \mathcal{M}_L^a\}.$$

Then $f_{\xi}(a+1) \ge f_{\xi}(a)$. Furthermore, for every $a \in \mathbb{Z}$ there is $z \in L$ with $v_L(z) = a$ and $v_L(\xi(z)) = f_{\xi}(a)$.

Corollary 3.4. Let $\xi \in K[G] \setminus \{0\}$, let $\beta \in L \otimes_K L$ satisfy $\xi = \phi(\beta)$, and let $a, b \in \mathbb{Z}$. Then the following are equivalent:

(a) $[a,b] \in G(\beta)$.

(b) $f_{\xi}(-b-i_0) = a \text{ and } f_{\xi}(-b-i_0+1) > a.$

Proof. Suppose $[a, b] \in G(\beta)$. It follows from Theorem 3.3 that for $y \in L$ with $v_L(y) = -b - i_0$ we have $v_L(\xi(y)) = a$. Therefore $f_{\xi}(-b - i_0) = a$. Suppose $f_{\xi}(-b - i_0 + 1) = a$. Then there is $z \in L$ with $v_L(z) = -b - i_0 + 1$ and $v_L(\xi(z)) = a$. It follows that there is $u \in \mathcal{O}_K^{\times}$ such that

$$v_L(\xi(y+uz)) = v_L(\xi(y) + u\xi(z)) > a.$$

Since $v_L(y+uz) = -b - i_0$ this contradicts Theorem 3.3. Hence $f_{\xi}(-b - i_0 + 1) > a$.

Suppose $f_{\xi}(-b-i_0) = a$ and $f_{\xi}(-b-i_0+1) > a$. Then there is $y_0 \in L$ such that $v_L(y_0) = -b - i_0$ and $v_L(\xi(y_0)) = a$. Let $y \in L$ satisfy $v_L(y) = -b - i_0$. Then there are $u \in \mathcal{O}_K^{\times}$ and $z \in \mathcal{M}_L^{-b-i_0+1}$ such that $y = uy_0 + z$. It follows that $v_L(\xi(z)) > a$, and hence that $v_L(\xi(y)) = v_L(u\xi(y_0) + \xi(z)) = a$. Since this holds for all $y \in L$ such that $v_L(y) = -b - i_0$ we get $[a, b] \in G(\beta)$ by Theorem 3.3.

The following application of Theorem 3.3 gives information about the effect that ξ has on valuations of arbitrary elements of L^{\times} . For a related result see Proposition 2.5.2 of [2].

Corollary 3.5. Let $\xi \in K[G] \setminus \{0\}$, let $\beta \in L \otimes_K L$ satisfy $\xi = \phi(\beta)$, and let $a, b \in \mathbb{Z}$. Then the following are equivalent:

(a)
$$[a,b] \in D(\beta)$$
.

(b) $f_{\xi}(-b - i_0) \le a$.

Proof. Suppose $[a, b] \in D(\beta)$. Then there is $[c, d] \in G(\beta)$ such that $[c, d] \leq [a, b]$. We may assume that $c \leq a$ and $d \leq b$. It follows from Corollary 3.4 that

$$f_{\xi}(-b-i_0) \le f_{\xi}(-d-i_0) = c \le a.$$

Suppose $[a, b] \notin D(\beta)$. Let d be the largest integer such that $d \leq b$ and there exists c such that $[c, d] \in G(\beta)$. By Corollary 3.4 we have

$$f_{\xi}(-b-i_0) \le f_{\xi}(-d-i_0) = c.$$

Suppose $f_{\xi}(-b - i_0) < c$. Then there is d' such that $d < d' \leq b$ and

$$f_{\xi}(-d'-i_0) < f_{\xi}(-d'-i_0+1).$$

Setting $c' = f_{\xi}(-d' - i_0)$ we get $[c', d'] \in G(\beta)$ by Corollary 3.4. This contradicts the maximality of d, so we must have $f_{\xi}(-b - i_0) = c$. Furthermore, since $[c, d] \not\leq [a, b]$ we have a < c. Hence $f_{\xi}(-b - i_0) > a$.

For $\beta \in L \otimes_K L$ with $\beta \neq 0$ set

$$d(\beta) = \min\{i+j : [i,j] \in D(\beta)\}$$

Define the diagonal of β to be

$$N(\beta) = \{[i, j] \in D(\beta) : i + j = d(\beta)\}.$$

Then $N(\beta) \subset G(\beta)$.

Definition 3.6. Let L/K be a totally ramified Galois extension of degree p^n . Say that L/K is a semistable extension if there is $\beta \in L \otimes_K L$ such that $\phi(\beta) \in K[G]$, $p \nmid d(\beta)$, and $|N(\beta)| = 2$. Say that L/K is a stable extension if β can be chosen to satisfy the additional condition $G(\beta) = N(\beta)$.

In [2, Th. 4.3.2] sufficient conditions are given for an ideal \mathcal{M}_L^h in a semistable extension L/K to be free over its associated order. In some cases where L/K is stable these conditions are shown to be necessary as well.

The following congruences imply that for $n \ge 2$ most totally ramified Galois extensions of degree p^n are not semistable.

Proposition 3.7. Let L/K be a semistable extension and let $b_1 \leq b_2 \leq \cdots \leq b_n$ be the lower ramification breaks of L/K, counted with multiplicity. Then $b_i \equiv -i_0 \pmod{p^n}$ for $1 \leq i \leq n$.

Proof. This follows from Propositions 3.2.2 and 4.3.2.1 in [2].

Motivated by the theory of scaffolds, we give a definition of precision for semistable extensions.

Definition 3.8. Let L/K be a totally ramified Galois extension of degree p^n and let $\mathfrak{c} \geq 1$. We say that L/K is semistable with precision \mathfrak{c} if there is $\beta \in L \otimes_K L$ such that $\phi(\beta) \in K[G], p \nmid d(\beta), |N(\beta)| = 2$, and $a + b \geq d(\beta) + \mathfrak{c}$ for all $[a, b] \in G(\beta) \setminus N(\beta)$.

If L/K is stable then we may choose β so that $G(\beta) = N(\beta)$. Hence L/K is semistable with infinite precision in this case. On the other hand, if L/K is semistable with sufficiently high precision then L/K is stable:

Proposition 3.9. Let L/K be a totally ramified Galois extension of degree p^n , and let $h \in \mathbb{S}_{p^n}$ satisfy $h \equiv i_0 \pmod{p^n}$. Suppose L/K is semistable with precision

$$\mathfrak{c} \ge \max\{h-1, p^n - h - 1\}.$$

Then L/K is a stable.

Proof. Since L/K is semistable with precision \mathfrak{c} there is $\alpha \in L \otimes_K L$ with $\phi(\alpha) \in K[G]$, $p \nmid d(\alpha), |N(\alpha)| = 2$, and $e + f \geq d(\alpha) + \mathfrak{c}$ for all $[e, f] \in G(\alpha) \smallsetminus N(\alpha)$. The proof of Proposition 3.2.1 in [2] shows that $N(\alpha) = \{[d(\alpha), 0], [0, d(\alpha)]\}$. By Proposition 4.3.2.1 in [2] we have $d(\alpha) \equiv i_0 \equiv h \pmod{p^n}$. Set $m = (d(\alpha) - h)/p^n$. Then $m \in \mathbb{Z}$, so we may define $\beta = \pi_K^{-m} \alpha$. We get $\phi(\beta) = \pi_K^{-m} \phi(\alpha) \in K[G]$ and $D(\beta) = [-mp^n, 0] + D(\alpha)$. It follows that $N(\beta) = \{[h, 0], [0, h]\}$ and $e + f \geq h + \mathfrak{c}$ for all $[e, f] \in G(\beta) \smallsetminus N(\beta)$.

Suppose there is $(e, f) \in \mathcal{F}$ such that $[e, f] \in G(\beta) \setminus N(\beta)$. Then $[h, 0] \not\leq [e, f]$ and $[0, h] \not\leq [e, f]$. It follows that

$$[e+1, f-p^n+1] \le [h, 0] \tag{3.1}$$

$$[e+1, f-p^n+1] \le [0, h]. \tag{3.2}$$

By (3.1) we have $e + 1 \le h$. Hence if $0 \le f \le h - 1$ then

$$e + f \le 2h - 2 < 2h - 1.$$

If $h \leq f \leq p^n - 1$ then by (3.2) we have $e + 1 \leq 0$, and hence

$$e + f \le -1 + p^n - 1 < p^n - 1$$

In either case we get $e + f < h + \mathfrak{c}$, a contradiction. Hence $G(\beta) \smallsetminus N(\beta) = \emptyset$, so L/K is stable.

4 Semistable extensions and Galois scaffolds

Let L/K be a totally ramified extension of degree p^n . In this section we show that L/K is semistable if and only if L/K has a Galois scaffold. We also show that if L/K has a Galois scaffold with sufficiently high precision then L/K is stable.

Theorem 4.1. Let L/K be a totally ramified Galois extension of degree p^n which is semistable. Then L/K admits a Galois scaffold with precision c = 1.

Proof. Let $h \in \mathbb{S}_{p^n}$ satisfy $h \equiv i_0 \pmod{p^n}$. Then by Propositions 3.2.1 and 4.3.2.1 of [2] there is $\beta \in L \otimes_K L$ such that $\phi(\beta) \in K[G]$ and $N(\beta) = \{[0,h], [h,0]\}$. For $1 \leq i \leq n$ set $\Theta_i = \phi(\beta^{p^n - p^{n-i} - 1})$. Since $\phi(\beta) \in K[G]$ it follows from Proposition 3.1 that $\Theta_i \in K[G]$. We clearly have $d(\beta^{p^n - p^{n-i} - 1}) = (p^n - p^{n-i} - 1)h$ and

$$N(\beta^{p^n - p^{n-i} - 1}) = \left\{ [sh, (p^n - p^{n-i} - 1 - s)h] : s \in \mathbb{S}_{p^n}, \ p \nmid \binom{p^n - p^{n-i} - 1}{s} \right\}.$$
 (4.1)

Let $\{\lambda_t : t \in \mathbb{Z}\}$ be a set of elements of L such that $v_L(\lambda_t) = t$ and $\lambda_{t'}\lambda_t^{-1} \in K$ for all $t, t' \in \mathbb{Z}$ with $t' \equiv t \pmod{p^n}$. It follows from (4.1) and Corollary 3.5 that for $t \in \mathbb{Z}$ we have

$$v_L(\Theta_i(\lambda_t)) \ge t + (p^n - p^{n-i} - 1)h + i_0.$$
 (4.2)

Since $p \nmid h$ there is a unique $s_i(t) \in \mathbb{S}_{p^n}$ such that

$$t \equiv -(p^n - p^{n-i} - 1 - s_i(t))h - i_0 \pmod{p^n}.$$
(4.3)

By applying Theorem 3.3 and Corollary 3.5 to (4.1) we see that equality holds in (4.2) if and only if $p \nmid \binom{p^n - p^{n-i} - 1}{s_i(t)}$. It follows from Lucas's theorem that this is equivalent to $s_i(t)_{(n-i)} \neq p-1$, which holds if and only if $p^{n-i} + s_i(t) < p^n$ and $(p^{n-i} + s_i(t))_{(n-i)} \geq 1$. Since $h = i \pmod{p^n}$ it follows from (4.3) that

Since $h \equiv i_0 \pmod{p^n}$ it follows from (4.3) that

$$t \equiv (p^{n-i} + s_i(t))h \pmod{p^n}$$

By Proposition 3.7 we have $h \equiv -b_j \pmod{p^n}$ for $1 \leq j \leq n$. Therefore

$$\mathfrak{a}(r(t)) \equiv p^{n-i} + s_i(t) \pmod{p^n}.$$
(4.4)

If we have equality in (4.2) then by the preceding paragraph we get $\mathfrak{a}(r(t)) = p^{n-i} + s_i(t)$ and $\mathfrak{a}(r(t))_{(n-i)} \geq 1$. Conversely, if $\mathfrak{a}(r(t))_{(n-i)} \geq 1$ then $\mathfrak{a}(r(t)) \geq p^{n-i}$, and hence $p^{n-i}+s_i(t) = \mathfrak{a}(r(t)) < p^n$ by (4.4). It follows from the preceding paragraph that equality holds in (4.2) in this case. Thus equality holds in (4.2) if and only if $\mathfrak{a}(r(t))_{(n-i)} \geq 1$.

For $1 \leq i \leq n$ set

$$v_i = \frac{(p^n - p^{n-i} - 1)h + i_0 - p^{n-i}b_i}{p^n}.$$

Then $v_i \in \mathbb{Z}$, so we may define $\Phi_i = \pi_K^{-v_i} \Theta_i \in K[G]$. Using (4.2) we get $v_L(\Phi_i(\lambda_t)) \ge t + p^{n-i}b_i$, with equality if and only if $\mathfrak{a}(r(t))_{(n-i)} \ge 1$. Hence by Proposition 2.2 there are $\Psi_i \in K[G]$ such that $(\{\Psi_i\}, \{\lambda_t\})$ is a Galois scaffold for L/K with precision 1. \Box

Theorem 4.2. Let L/K be a totally ramified Galois extension of degree p^n which has a Galois scaffold ($\{\Phi_i\}, \{\lambda_t\}$) with precision $\mathfrak{c} \geq 1$. Then L/K is semistable with precision \mathfrak{c} .

Proof. Set $\xi = \Psi^{(p^n-2)}$. It follows from (2.1) that for every $t \in \mathbb{Z}$ there is $U_{p^n-2,t} \in \mathcal{O}_K^{\times}$ such that the following holds modulo $\lambda_{t+\mathfrak{b}(p^n-2)}\mathcal{M}_L^{\mathfrak{c}}$:

$$\xi(\lambda_t) \equiv \begin{cases} U_{p^n-2,t}\lambda_{t+\mathfrak{b}(p^n-2)} & \text{if } p^n-2 \preceq \mathfrak{a}(r(t)), \\ 0 & \text{if } p^n-2 \not\preceq \mathfrak{a}(r(t)). \end{cases}$$
(4.5)

Therefore we have $v_L(\xi(\lambda_t)) \geq t + \mathfrak{b}(p^n - 2)$, with equality if and only if $p^n - 2 \leq \mathfrak{a}(r(t))$. Now let $y \in L^{\times}$ be arbitrary and set $v_L(y) = t$. It follows from the above that $v_L(\xi(y)) \geq t + \mathfrak{b}(p^n - 2)$, again with equality if and only if $p^n - 2 \leq \mathfrak{a}(r(t))$. The condition $p^n - 2 \leq \mathfrak{a}(r(t))$ is equivalent to $\mathfrak{a}(r(t)) \in \{p^n - 1, p^n - 2\}$, which is valid if and only if we one of the following congruences holds:

$$t \equiv -\mathfrak{b}(p^n - 1) \pmod{p^n} \tag{4.6}$$

$$t \equiv -\mathfrak{b}(p^n - 2) \pmod{p^n}. \tag{4.7}$$

It follows that $f_{\xi}(t) \ge t + \mathfrak{b}(p^n - 2)$, with equality if and only if either (4.6) or (4.7) holds. Let $\beta \in L \otimes_K L$ be such that $\phi(\beta) = \xi$. By Corollary 3.5 we have

$$[t + \mathfrak{b}(p^n - 2) - 1, -i_0 - t] \notin D(\beta)$$

for all $t \in \mathbb{Z}$. Furthermore, $[t + \mathfrak{b}(p^n - 2), -i_0 - t] \in D(\beta)$ if and only if either (4.6) or (4.7) holds. It follows that

$$d(\beta) = \mathfrak{b}(p^n - 2) - i_0 = -b_n$$

and $N(\beta) = \{[-b_n, 0], [0, -b_n]\}$. Since $p \nmid b_n$ this shows that L/K is semistable. Suppose $[e, f] \in G(\beta)$ and $[e, f] \notin N(\beta)$. Then $f \not\equiv 0, -b_n \pmod{p^n}$, and hence

$$-f - i_0 \not\equiv -\mathfrak{b}(p^n - 1) \pmod{p^n}$$
$$-f - i_0 \not\equiv -\mathfrak{b}(p^n - 2) \pmod{p^n}.$$

It follows from Theorem 3.3 and (4.5) that

$$e = v_L(\xi(\lambda_{-f-i_0}))$$

$$\geq -f - i_0 + \mathfrak{b}(p^n - 2) + \mathfrak{c}$$

$$= -f - b_n + \mathfrak{c}.$$

Hence $e + f \ge -b_n + \mathfrak{c} = d(\beta) + \mathfrak{c}$, so L/K is semistable with precision \mathfrak{c} .

Corollary 4.3. Let L/K be a totally ramified Galois extension of degree p^n and let $h \in \mathbb{S}_{p^n}$ satisfy $h \equiv i_0 \pmod{p^n}$. Suppose that L/K has a Galois scaffold $(\{\Phi_i\}, \{\lambda_t\})$ with precision

$$\mathfrak{c} \ge \max\{h-1, p^n - h - 1\}.$$

Then L/K is a stable extension.

Proof. This follows from Theorem 4.2 and Proposition 3.9.

Remark 4.4. The proofs of Theorem 4.2 and Corollary 4.3 don't require part (iii) of Definition 2.1, which states that Ψ_i lies in the augmentation ideal of K[G].

Remark 4.5. It would be interesting to get some sort of converse to Theorem 4.2 or Corollary 4.3. Unfortunately, when L/K is semistable with precision \mathfrak{c} the Galois scaffold for L/K produced by Theorem 4.1 does not seem to inherit this precision. Constructing a Galois scaffold with precision $\mathfrak{c} > 1$ for a stable or semistable extension L/K would require at least a refinement of Theorem 4.1, or perhaps a completely new approach.

Corollary 4.6. Let L/K be a totally ramified Galois extension of degree p^n and let G = Gal(L/K). Then the following statements are equivalent:

- (a) L/K admits a Galois scaffold with precision \mathfrak{c} for some $\mathfrak{c} \geq 1$.
- (b) L/K is semistable.
- (c) Let $\rho \in L$ satisfy $v_L(\rho) \equiv -i_0 \pmod{p^n}$. Then for all $\xi \in K[G]$ and $\lambda \in L^{\times}$ we have

$$v_L(\xi(\rho)) - v_L(\rho) \le v_L(\xi(\lambda)) - v_L(\lambda).$$

Proof. The equivalence of (a) and (b) follows from Theorems 4.1 and 4.2. The equivalence of (b) and (c) is proved in Theorem 4.4 of [2]. \Box

We conclude with some applications of our main results.

Corollary 4.7. Let L/K be a totally ramified Galois extension of degree p^n which admits a Galois scaffold with precision \mathfrak{c} for some $\mathfrak{c} \ge 1$. Then the lower ramification breaks of L/K satisfy $b_i \equiv -i_0 \pmod{p^n}$ for $1 \le i \le n$.

Proof. It follows from Theorem 4.2 that L/K is a semistable extension. Hence by Proposition 3.7 we have $b_i \equiv -i_0 \pmod{p^n}$ for all i.

Let K be a finite unramified extension of \mathbb{Q}_p , with $K \neq \mathbb{Q}_p$. In [2, §5.1] a totally ramified elementary abelian p-extension L/K is constructed which is semistable, but not stable. The following shows that there are no examples of elementary abelian pextensions in characteristic p with these properties.

Proposition 4.8. Let char(K) = p and let L/K be a totally ramified elementary abelian *p*-extension. If L/K is semistable then it is stable.

Proof. It follows from Theorem 4.1 that L/K has a Galois scaffold $(\{\Psi_i\}, \{\lambda_t\})$ with precision 1. Since char(K) = p we have $\Psi_i^p = 0$ for $1 \le i \le n$. Hence by Theorem A.1(ii) of [3] the extension L/K has a Galois scaffold with precision ∞ . Therefore by Corollary 4.3 L/K is stable.

References

- [1] M. V. Bondarko, Local Leopoldt's problem for rings of integers in abelian *p*-extensions of complete discrete valuation fields, Documenta Math. 5 (2000), 657–693.
- [2] M. V. Bondarko, Local Leopoldt's problem for ideals in totally ramified *p*-extensions of complete discrete valuation fields, Algebraic number theory and algebraic geometry, 27–57, Contemp. Math. **300**, Amer. Math. Soc. Providence, RI, 2002.
- [3] Nigel P. Byott, Lindsay N. Childs, and G. Griffith Elder, Scaffolds and generalized integral Galois module structure, Ann. Inst. Fourier (Grenoble) **68** (2018), 965–1010.
- [4] V. Heiermann, De nouveaux invariants numériques pour les extensions totalement ramifiées de corps locaux, J. Number Theory **59** (1996), 159–202.
- [5] J.-P. Serre, Corps Locaux, Hermann, Paris, 1962.
- [6] S. Ullom, Integral normal bases in Galois extensions of local fields, Nagoya Math. J. 39 (1970) 141–148.