An application of Baker's method to the Jeśmanowicz' conjecture on primitive Pythagorean triples

Maohua Le

17 October 2018

Abstract

Let m, n be positive integers such that m > n, gcd(m, n) = 1 and $m \not\equiv n \mod 2$. In 1956, L. Jeśmanowicz [6] conjectured that the equation $(m^2 - n^2)^x + (2mn)^y = (m^2 + n^2)^z$ has only the positive integer solution (x, y, z) = (2, 2, 2). This problem is not yet solved. In this paper, combining a lower bound for linear forms in two logarithms due to M. Laurent [7] with some elementary methods, we prove that if $mn \equiv 2 \mod 4$ and m > 30.8n, then Jeśmanowicz' conjecture is true.

Mathematics Subject Classification: 11D61, 11J86

Keywords: ternary purely exponential diophantine equation, primitive Pythagorean triple, Jeśmanowicz' conjecture, application of Baker's method

1 Introduction

Let \mathbb{N} be the set of positive integers. Let m, n be positive integers such that m > n, gcd(m, n) = 1 and $m \neq n \mod 2$. It is well known that the triple $(m^2 - n^2, 2mn, m^2 + n^2)$ is a primitive Pythagorean triple with $(m^2 - n^2)^2 + (2mn)^2 = (m^2 + n^2)^2$. In 1956 L. Jeśmanowicz [6] conjectured that the equation

$$(m^{2} - n^{2})^{x} + (2mn)^{y} = (m^{2} + n^{2})^{z}, x, y, z \in \mathbb{N}$$

$$(1.1)$$

has only the solution (x, y, z) = (2, 2, 2). This problem is not solved yet.

For over twenty years, many papers have investigated Jeśmanowicz' conjecture for the case that $mn \equiv 2 \mod 4$. In this respect, Jeśmanowicz' conjecture is true in the following cases:

- (i) (M.-H. Le [8]) $mn \equiv 2 \mod 4$ and $m^2 + n^2$ is an odd prime power.
- (ii) (Z.-F. Cao [1]) $(m, n) \equiv (5, 2) \mod 8$.
- (iii) (M.-J. Deng and D.-M. Huang [3]) $mn \equiv (2,3) \mod 4$ and either $m + n \not\equiv 1 \mod 16$ or y > 1.
- (iv) (M.-M. Ma and Y.-G. Chen [10]) $mn \equiv 2 \mod 4$ and y > 1.
- (v) (K. Takakuwa and Y. Asaeda [14]) $m \equiv 2 \mod 4$, n = 3 and m satisfies some conditions.
- (vi) (Y.-D. Guo and M.-H. Le [4]) $m \equiv 2 \mod 4$, n = 3 and m > 6000.

- (vii) (K. Takakuwa [13]) $m \equiv 2 \mod 4$ and $n \in \{3, 7, 11, 15\}$.
- (viii) (N. Terai [15]) n = 2.
- (ix) (M.-J. Deng and J. Guo [2]) $n \equiv 2 \mod 4$ and n < 600.
- (x) (M.-H. Le [9]) $(m, n) \equiv (2, 3) \mod 4$ and m > 81n.
- (xi) (T. Miyazaki and N. Terai [11]) $n \equiv 2 \mod 4$, m > 72n and the divisors of n satisfy some conditions.
- (xii) (P.-Z. Yuan and Q. Han [16]) $mn \equiv 2 \mod 4$, m > 72n and the divisors of m, n satisfy some conditions.

In this paper, combining a lower bound for linear forms in two logarithms due to M. Laurent [7] with some elementary methods, we improve the results of [9], [11] and [16] as follows:

Theorem 1.1. If $mn \equiv 2 \mod 4$ and m > 30.8n, then Jeśmanowicz' conjecture is true.

2 Preliminaries

Lemma 2.1 ([2]). If $n \equiv 2 \mod 4$ and n < 600 then Jeśmanowicz' conjecture is true.

Lemma 2.2 (Corollary 1.1 of [16]). If $m \equiv 2 \mod 4$, $n \not\equiv 1 \mod 8$ and n < 85, then Jeśmanowicz' conjecture is true.

Lemma 2.3 ([5]). If $mn \equiv 2 \mod 4$ and m+n has a prime divisor p with $p \not\equiv 1 \mod 16$, then Jeśmanowicz' conjecture is true.

Lemma 2.4. If $mn \equiv 2 \mod 4$, m > 30.8n and (1.1) has a solution $(x, y, z) \neq (2, 2, 2)$, then $m^2 + n^2 > m^2 - n^2 \ge 2704$.

Proof. By Lemma 2.3, if $mn \equiv 2 \mod 4$ and (1.1) has a solution (x, y, z) with $(x, y, z) \neq (2, 2, 2)$, then every prime divisor p of m + n satisfies $p \equiv 1 \mod 16$. This implies that $m + n \equiv 1 \mod 16$. Hence, we have $m + n \equiv 1 \mod 4$ and $(m, n) \equiv (2, 3)$ or $(3, 2) \mod 4$. Therefore, if $m \equiv 2 \mod 4$, then $n \equiv 3 \mod 4$ and

$$n \ge 85. \tag{2.1}$$

Similarly, by Lemma 2.1, (2.1) holds if $n \equiv 2 \mod 4$. Thus, since m > 30.8n, we get from (2.1) that

$$m^{2} + n^{2} > m^{2} - n^{2} \ge m + n > 31.8n \ge 31.8 \times 85 = 2703.$$
(2.2)

Lemma 2.5 ([15]). If $m^2 + n^2 \equiv 1 \mod 2mn$, then Jeśmanowicz' conjecture is true.

Lemma 2.6 ([10]). If $mn \equiv 2 \mod 4$ and (1.1) has a solution (x, y, z) with $(x, y, z) \neq (2, 2, 2)$, then the solution satisfies $x \equiv 0 \mod 2$, y = 1 and $z \equiv 1 \mod 2$.

Lemma 2.7. Let a_1 , a_2 , b_1 , b_2 be positive integers such that $\min\{a_1, a_2\} > 1$ and $gcd(a_1, a_2) = 1$. Further let $\Lambda = b_1 \log(a_1) - b_2 \log(a_2)$. Let ρ and μ be real numbers with $\rho > 1$ and $1/3 \le \mu \le 1$. Further let

$$\delta = \frac{1}{2}(1 + 2\mu - \mu^2), \lambda = \delta \log(\rho).$$
(2.3)

Then

$$\log|\Lambda| \ge -CA_1A_2B^2 - \sqrt{\omega\theta}B - \log(C'A_1A_2B^2), \tag{2.4}$$

where

$$A_j \ge \max\{1, (\rho+1)\log(a_j)\}, A_1A_2 \ge \lambda^2, j = 1, 2,$$
(2.5)

$$B \ge \log(\rho) + \max\left\{\frac{\log(2)}{2}, \lambda, 1.81 + \log(\lambda) + \log\left(\frac{b_1}{A_2} + \frac{b_2}{A_1}\right)\right\},$$
(2.6)

$$\omega = 2 + 2\left(1 + \frac{1}{4H^2}\right)^{1/2}, \theta = \frac{1}{2H} + \left(1 + \frac{1}{4H^2}\right)^{1/2}, H = \frac{B}{\lambda},$$
(2.7)

$$C = \frac{C_0 \mu}{\lambda^3 \delta}, C' = \frac{\sqrt{C_0 \omega \theta}}{\lambda^3}, \tag{2.8}$$

$$C_0 = \left(\frac{\omega}{6} + \frac{1}{2}\left(\frac{\omega^2}{9} + \frac{8\lambda\omega^{5/4}\theta^{1/4}}{3\sqrt{A_1A_2H}} + \frac{4}{3}\left(\frac{1}{A_1} + \frac{1}{A_2}\right)\frac{\lambda\omega}{H}\right)^{1/2}\right)^2.$$
 (2.9)

Proof. This lemma is the special case of Theorem 2 of [7] for γ_1 and γ_2 coprime positive integers. **Lemma 2.8.** Under the assumptions of Lemma 2.7, if $\min\{a_1, a_2\} \ge 2704$ and $b_1/A_2 > b_2/A_1 > 240$, then

$$\log|\Lambda| > -14.8365\log(a_1)\log(a_2)\left(1.8248 + \log\left(\frac{b_1}{\log(a_2)} + \frac{b_2}{\log(a_1)}\right)\right)^2.$$
 (2,.10)

Proof. By Lemma 2.7, we may choose parameters

$$\rho = e^{1.575}, \mu = \frac{1}{3}.$$
(2.11)

By (2.3) and (2.11), we have

$$\delta = \frac{7}{9}, \lambda = 1.225. \tag{2.12}$$

Since $\min\{a_1, a_2\} > 2703$, by (2.5), (2.11) and (2.12), we can set

$$A_j = 5.8314 \log(a_j), j = 1, 2.$$
(2.13)

Hence, by (2.13), we have

$$A_j > 46.0803, j = 1, 2. (2.14)$$

By (2.6), (2.11) and (2.12), we can set

$$B = 3.5880 + \log\left(\frac{b_1}{A_2} + \frac{b_2}{A_1}\right).$$
(2.15)

Since $b_1/A_2 > b_2/A_1 > 240$, by (2.15) and (2.7), we have

$$B > 9.7617$$
 (2.16)

and

$$H > 7.9688.$$
 (2.17)

Therefore, by (2.7) and (2.17), we get

$$\omega < 4.0040, \theta < 1.0648. \tag{2.18}$$

Further, by (2.9), (2.12), (2.14), (2.17) and (2.18), we have

$$C_0 < 1.8706,$$
 (2.19)

and by (2.8), we get

$$C < 0.4361, C' < 2.0829. \tag{2.20}$$

From (2.4) and (2.20), we have

$$\log|\Lambda| > -\left(0.4361 + \frac{\sqrt{\omega\theta}}{A_1 A_2 B} + \frac{\log(2.0829A_1 A_2 B^2)}{A_1 A_2 B^2}\right) A_1 A_2 B^2.$$
(2.21)

By (2.14), (2.16) and (2.18), we get

$$\frac{\sqrt{\omega\theta}}{A_1 A_2 B} < 1.0026 \times 10^{-4}.$$
(2.22)

Since $f(t) = \log(t)/t$ is a decreasing function for t > e, by (2.14) and (2.16), we have

$$\frac{\log(2.0829A_1A_2B^2)}{A_1A_2B^2} < \frac{\log(2.0829 \times 46.0803^2 \times 9.7617^2)}{46.0803^2 \times 9.7617^2} < 0.6401 \times 10^{-4}.$$
(2.23)

Thus, by (2.13), (2.15), (2.21), (2.22) and (2.23), we obtain

$$\log|\Lambda| > -0.4363A_1A_2B^2 > -14.8365\log(a_1)\log(a_2)\left(1.8248 + \log\left(\frac{b_1}{\log(a_2)} + \frac{b_2}{\log(a_1)}\right)\right)^2.$$
 (2.24)

The lemma is proved.

3 Proof of Theorem

We mow assume that $mn \equiv 2 \mod 4$, m > 30.8n and (1.1) has a solution (x, y, z) with $(x, y, z) \neq (2, 2, 2)$. By Lemma 2.6, the solution (x, y, z) satisfies

$$(m^2 - n^2)^x + 2mn = (m^2 + n^2)^z, x, z \in \mathbb{N}, x \equiv 0 \mod 2, z \equiv 1 \mod 2.$$
(3.1)

By Lemma 2.4, $m^2 + n^2$ and $m^2 - n^2$ satisfy (2.2). Since $x \equiv 0 \mod 2$, if $x \leq z$ then from (3.1) we get $2mn = (m^2 + n^2)^z - (m^2 - n^2)^x \geq (m^2 + n^2)^x - (m^2 - n^2)^x \geq (m^2 + n^2)^2 - (m^2 - n^2)^2 = (2mn)^2 > 2mn$, a contradiction. Therefore, we have

$$x > z \tag{3.2}$$

and x - z an odd integer.

Since $x \equiv 0 \mod 2$ and $(m^2 - n^2)^2 \equiv (m^2 + n^2)^2 \mod (2mn)$, we have $(m^2 - n^2)^x \equiv (m^2 + n^2)^x \mod (2mn)$. Further, since $gcd(2mn, m^2 + n^2) = 1$, by (3.1) and (3.2), we get

$$(m^2 + n^2)^{x-z} \equiv 1 \mod (2mn).$$
 (3.3)

Hence, by Lemma 2.5, we see from (3.3) that the case x - z = 1 can be removed. So we have

$$x - z \ge 3. \tag{3.4}$$

Let k = m/n. By (3.1) and (3.4), we have

$$(m^{2} - n^{2})^{3} \le (m^{2} - n^{2})^{x-z} < \left(\frac{m^{2} + n^{2}}{m^{2} - n^{2}}\right)^{z} = \left(1 + \frac{2}{k^{2} - 1}\right)^{z}.$$
(3.5)

Notice that $\log(1+t) < t$ for any t > 0. By (3.5), we get

$$3\log(m^2 - n^2) < \frac{2z}{k^2 - 1}.$$
(3.6)

Since k > 30.8, we see from (3.6) that

$$\frac{z}{\log(m^2 - n^2)} > \frac{3}{2}(k^2 - 1) > 1421.46.$$
(3.7)

On the other hand, by (3.1), we have

$$z \log(m^{2} + n^{2}) = \log((m^{2} - n^{2})^{x} + 2mn) = \log((m^{2} - n^{2})^{x}) + \log\left(1 + \frac{2mn}{(m^{2} - n^{2})^{x}}\right) < x \log(m^{2} - n^{2}) + \frac{2mn}{(m^{2} - n^{2})^{x}}.$$
(3.8)

Let $(a_1, a_2, b_1, b_2) = (m^2 + n^2, m^2 - n^2, z, x)$ and $\Lambda = z \log(m^2 + n^2) - x \log(m^2 - n^2)$. By (3.8), we have

$$0 < \Lambda < \frac{2mn}{(m^2 - n^2)^x}.$$
(3.9)

Further, since $(m^2 - n^2)^x > (m^2 - n^2)^2 \ge (m + n)^2 > 2mn$, we see from (3.1) that $2(m^2 - n^2)^x > (m^2 + n^2)^z$. Hence, by (3.9), we get

$$0 < \Lambda < \frac{4mn}{(m^2 + n^2)^z},\tag{3.10}$$

whence we obtain

$$\log(4mn) - \log|\Lambda| > z\log(m^2 + n^2).$$
(3.11)

Since $x \ge 4$, by (2.2), (3.7) and (3.8), we have

$$\frac{z}{\log(m^2 - n^2)} > \frac{x}{\log(m^2 + n^2)} > \frac{z}{\log(m^2 - n^2)} - \frac{2mn}{(m^2 - n^2)^x \log(m^2 + n^2)} \frac{1}{\log(m^2 - n^2)} > \frac{z}{\log(m^2 - n^2)} - \frac{1}{(m + n)^2 (\log(m + n))^2} > 1421.46 - 5 \times 10^{-8},$$
(3.12)

whence we get

$$\frac{z}{5.8314\log(m^2 - n^2)} > \frac{x}{5.8314\log(m^2 + n^2)} > 240.$$
(3.13)

Therefore, by Lemma 2.8, we see from (2.2) and (3.13) that

$$\log|\Lambda| > -14.8365\log(m^2 + n^2)\log(m^2 - n^2)\left(1.8248 + \log\left(\frac{z}{\log(m^2 - n^2)} + \frac{x}{\log(m^2 + n^2)}\right)\right)^2.$$
 (3.14)

Substituting (3.14) into (3.11), by (3.13), we have

$$\frac{\log(4mn)}{\log(m^2 + n^2)\log(m^2 - n^2)} + 14.8365\left(1.8248 + \log\left(\frac{2z}{\log(m^2 - n^2)}\right)\right)^2 > \frac{z}{\log(m^2 - n^2)}.$$
(3.15)

Since $m^2 + n^2 > 2mn$ and $m^2 - n^2 \ge m + n$, by (2.2), we have

$$\frac{\log(4mn)}{\log(m^2 + n^2)\log(m^2 - n^2)} < \frac{1}{\log(m + n)} \left(\frac{\log(2)}{\log(m + n)} + 1\right) < 0.1266.$$
(3.16)

Hence, by (3.15) and (3.16), we get

$$0.1266 + 14.8365 \left(1.8248 + \log\left(\frac{2z}{\log(m^2 - n^2)}\right) \right)^2 > \frac{z}{\log(m^2 - n^2)}.$$
(3.17)

Let $f(t) = t - 0.1266 - 14.8365(1.8248 + \log(2t))^2$. We see from (3.17) that

$$f\left(\frac{z}{\log(m^2 - n^2)}\right) < 0. \tag{3.18}$$

Since $f'(t) = 1 - 29.6730(1.8248 + \log(2t))/t$ and f'(t) > 0 for t > 250, where f'(t) is the derivative of f(t), we have f(t) > f(1420) > 0 for t > 1420. Therefore, by (3.18), we get $z/\log(m^2 - n^2) < 1420$, which contradicts (3.7). Thus, if $mn \equiv 2 \mod 4$ and m > 30.8n, then (1.1) has only the solution (x, y, z) = (2, 2, 2). The theorem is proved.

Acknowledgment: The author would like to thank Prof. R. Scott and Prof. R. Styer for reading the original manuscript carefully and giving valuable advice. Especially thanks to Prof. R. Styer for checking the calculations and providing other technical assistance.

References

- [1] Z.-F. Cao. A note on the diophantine equation $a^x + b^y = c^z$. Acta Arith., 91(1):85–93, 1999.
- [2] M.-J. Deng and J. Guo. A note on Jeśmanowicz' conjecture concerning primitive Pythagorean triples II. Acta Math. Hung., 153(2):436-448, 2017.
- [3] M.-J. Deng and D.-M. Huang. A note on Jeśmanowicz conjecture concerning primitive Pythagorean triples. Bull. Aust. Math. Soc., 95(1):5–13, 2017.
- [4] Y.-D. Guo and M.-H. Le. A note on Jeśmanowicz' conjecture concerning Pythagorean numbers. Commen. Math. Univ. St. Pauli, 44(2):225–228, 1995.
- [5] Q. Han and P.-Z. Yuan. A note on Jeśmanowicz' conjecture. Acta Math. Hung., to appear.
- [6] L. Jeśmanowicz. Several remarks on Pythagorean numbers. Wiadom. Math., 1(2):196–202, 1955/1956. (in Polish).
- [7] M. Laurent, Linear forms in two logarithms and interpolation determinants II. Acta Arith., 133(4): 325–348, 2008.

- [8] M.-H. Le. A note on Jeśmanowicz' conjecture. Colloq. Math., 69(1):47-51, 1995.
- [9] M.-H. Le. On Jeśmanowicz' conjecture concerning Pythagorean numbers. Proc. Japan Acad., Ser. A, 72A(5):97–98, 1996.
- [10] M.-M. Ma and Y.-G. Chen. Jeśmanowicz' conjecture on Pythagorean triples. Bull. Aust. Math. Soc., 96(1):30–35, 2017.
- [11] T. Miyazaki and N. Terai. On Jeśmanowicz' conjecture concerning Pythagorean triples II. Acta Math. Hung., 147(2):286–293, 2015.
- [12] T. Miyazaki, P.-Z. Yuan, and D.-Y. Wu. Generalizations of classical results on Jeśmanowicz' conjecture concerning Pythagorean triples II. J. Number Theory, 141(1):184–201, 2014.
- [13] K. Takakuwa. A remark on Jeśmanowicz' conjecture. Proc. Japan Acad., Ser. A, 72A(6):109–110, 1996.
- [14] K. Takakuwa and Y. Asaeda. On a conjecture on Pythagorean numbers II. Proc. Japan Acad., Ser. A, 69A(8):287–290, 1993.
- [15] N. Terai. On Jeśmanowicz' conjecture concerning primitive Pythagorean triples. J. Number Theory, 141(2):316–323, 2014.
- [16] P.-Z. Yuan and Q. Han. Jeśmanowicz' conjecture and related equations. Acta Arith., 184(1):37–49, 2018.

Maohua Le Institute of Mathematics Lingnan Normal College Zhangjiang, Guangdong 524048 China