QUILLEN–SUSLIN THEORY FOR CLASSICAL GROUPS: REVISITED OVER GRADED RINGS

RABEYA BASU AND MANISH KUMAR SINGH

2010 Mathematics Subject Classification: 11E57, 13A02, 13B25, 13B99, 13C10, 13C99, 15B99.

Key words: Bilinear forms, Symplectic and Orthogonal forms, Graded rings.

Abstract: In this paper we deduce a graded version of Quillen–Suslin's Local-Global Principle for the traditional classical groups, *viz.* general linear, symplectic and orthogonal groups and establish its equivalence of the normality property of the respective elementary subgroups. This generalizes previous result of Basu–Rao–Khanna; cf. [3]. Then, as an application, we establish an analogue Local-Global Principle for the commutator subgroups of the special linear and symplectic groups. Finally, by using Swan-Weibel's homotopy trick, we establish graded analogue of the Local-Global Principle for the transvection subgroups of the full automorphism groups of the linear, symplectic and orthogonal modules. This generalizes the previous result of Bak–Basu–Rao; cf. [5].

1. INTRODUCTION

In [3], the first author with Ravi A. Rao and Reema Khanna revisited Quillen-Suslin's Local-Global Principle for the linear, symplectic and orthogonal groups to show the equivalence of the Local-Global Principle and Normality property of their Elementary subgroups. A "relative version" is also established recently by the same authors; *cf.* [6]. In [3], the authors have also deduced a Local-Global Principle for the commutator subgroups of the special linear groups, and symplectic groups. In this article we aim to revisit those results over commutative \mathbb{N} -graded rings with identity. We follow the line of proof described in [3].

For the linear case, the graded version of the Local-Global Principle was studied by Chouinard in [7], and by J. Gubeladze in [8], [9]. Though the analogue results are expected for the symplectic and orthogonal groups, according to our best knowledge, that is not written explicitly in any existing literatures. By deducing the equivalence, we establish the graded version of the Local-Global Principle for the above two types of classical groups.

To generalize the existing results, from the polynomial rings to the graded rings, one has to use the line of proof of Quillen–Suslin's patching argument

Corresponding Author: Rabeya Basu – rabeya.basu@gmail.com.

(cf. [12], [13]), and Swan–Weibel's homotopy trick. For a nice exposition we refer to [8] by J. Gubeladze.

Though, we are writing this article for commutative graded rings, one may also consider standard graded algebras which are finite over its center. For a commutative N-graded ring with 1, we establish:

(1) (Theorem 3.8) Normality of the elementary subgroups is equivalent to the graded Local-Global Principle for the linear, symplectic and orthogonal groups.

(2) (Theorem 4.8) Analogue of Quillen–Suslin's Local-Global Principle for the commutator subgroups of the special linear and symplectic groups over graded rings.

In [5], the following was established by the first author with A. Bak and R.A. Rao:

(Local-Global Principle for the Transvection Subgroups) An analogue of Quillen–Suslin's Local-Global Principle for the transvection subgroup of the automorphism group of projective, symplectic and orthogonal modules of global rank at least 1 and local rank at least 3, under the assumption that the projective module has constant local rank and that the symplectic and orthogonal modules are locally an orthogonal sum of a constant number of hyperbolic planes.

In this article, we observe that by using Swan-Weibel's homotopy trick, one gets an analogue statement for the graded case. More precisely, we deduce the following fact:

(3) (**Theorem 5.4**) Let $A = \bigoplus_{i=0}^{\infty} A_i$ be a graded ring and $Q \simeq P \oplus A_0$ be a projective A_0 -module ($Q \simeq P \oplus \mathbb{H}(A_0)$ for symplectic and orthogonal modules). If an automorphism of $Q \otimes_{A_0} A$ is locally in the transvection subgroup, then it is globally in the transvection subgroup.

2. Definitions and Notations

Let us start by recalling the following well-known fact: Given a ring R and a subring R_0 , one can express R as a direct limit of subrings which are finitely generated over R_0 as rings. Considering R_0 to be the minimal subring (*i.e.* the image of \mathbb{Z}), it follows that every ring is a direct limit of Noetherian rings. Hence we may consider R to be Noetherian (*cf.* Pg 271 [11]).

Throughout this paper we assume A to be a Noetherian, commutative graded ring with identity 1. We shall write $A = A_0 \oplus A_1 \oplus A_2 \oplus \cdots$. As we know the multiplication in a graded ring satisfies the following property: For all i, j, $A_iA_j \subset A_{i+j}$. An element $a \in A$ will be denoted by $a = a_0 + a_1 + a_2 + \cdots$, where $a_i \in A_i$ for each i, and and all but finitely many a_i are zero. Let $A_+ = A_1 \oplus A_2 \oplus \cdots$. Graded structure of A induces a graded structure on $M_n(A)$ (ring of $n \times n$ matrices). Let S be a multiplicatively closed subset of A_0 . Then for a non-zero divisor $s \in S$ we shall denote the localization of a matrix $\alpha \in M_n(A)$ to $M_n(A_s)$ as α_s . Otherwise, $\alpha_i, i \in \mathbb{Z}$ will represent the *i*-th component of α . We shall use standard notations $\operatorname{GL}_n(A)$ and $\operatorname{SL}_n(A)$ to denote the group of invertible matrices and its subgroup of all invertible matrices with determinant 1 respectively.

We recall the well-known "Swan-Weibel's homotopy trick", which is the main ingredient to handle the graded case.

Definition 2.1. Let $a \in A_0$ be a fixed element. We fix an element $b = b_0 + b_1 + \cdots$ in A and define a ring homomorphism $\epsilon : A \to A[X]$ given by

 $\epsilon(b) = \epsilon(b_0 + b_1 + \cdots) = b_0 + b_1 X + b_2 X^2 + \cdots + b_i X^i + \cdots$

Then we evaluate the polynomial $\epsilon(b)(X)$ at X = a and denote the image by $b^+(a)$ i.e. $b^+(a) = \epsilon(b)(a)$. Note that $(b^+(x))^+(y) = b^+(xy)$. Observe, $b_0 = b^+(0)$. We shall use this fact frequently.

The above ring homomorphism ϵ induces a group homomorphism at the $\operatorname{GL}_n(A)$ level for every $n \geq 1$, *i.e.* for $\alpha \in \operatorname{GL}_n(A)$ we get a map

 $\epsilon : \operatorname{GL}_n(A) \to \operatorname{GL}_n(A[X])$ defined by

 $\alpha = \alpha_0 \oplus \alpha_1 \oplus \alpha_2 \oplus \dots \mapsto \alpha_0 \oplus \alpha_1 X \oplus \alpha_2 X^2 \cdots,$

where $\alpha_i \in \operatorname{GL}_n(A_i)$. As above for $a \in A_0$, we define $\alpha^+(a)$ as

$$\alpha^+(a) = \epsilon(\alpha)(a).$$

Now we are going to recall the definitions of the traditional classical groups, *viz.* the general linear groups, the symplectic and orthogonal groups (of even size) and their type subgroups, *viz.* elementary (symplectic and orthogonal elementary resp.) subgroups.

Let e_{ij} be the matrix with 1 in the *ij*-position and 0's elsewhere. The matrices of the form $\{E_{ij}(\lambda) : \lambda \in A \mid i \neq j\}$, where

$$\mathbf{E}_{ij}(\lambda) = \mathbf{I}_n + \lambda e_{ij}$$

are called the **elementary generators**.

Definition 2.2. The subgroup generated by the set $\{E_{ij}(\lambda) : \lambda \in A \mid i \neq j\}$, is called the **elementary subgroup of the general linear group** and is denoted by $E_n(A)$. Observe that $E_n(A) \subseteq SL_n(A) \subseteq GL_n(A)$.

Let σ be a permutation defined by: For $i \in \{1, \ldots, 2m\}$

$$\sigma(2i) = 2i - 1$$
 and $\sigma(2i - 1) = 2i$.

With respect to this permutation, we define two $2m \times 2m$ forms (viz.) ψ_m and $\tilde{\psi}_m$ as follows: For m > 1, let

$$\psi_1 = \begin{bmatrix} 0 & 1\\ -1 & 0 \end{bmatrix} \text{ and for } m > 1, \ \psi_m = \begin{bmatrix} \psi_{m-1} & 0\\ 0 & I_2 \end{bmatrix},$$
$$\widetilde{\psi}_1 = \begin{bmatrix} 0 & 1\\ 1 & 0 \end{bmatrix} \text{ and for } m > 1, \ \widetilde{\psi}_m = \begin{bmatrix} \widetilde{\psi}_{m-1} & 0\\ 0 & I_2 \end{bmatrix}.$$

Using above two forms, we define the following traditional classical groups:

Definition 2.3. A matrix $\alpha \in \operatorname{GL}_{2m}(A)$ is called symplectic if it fixes ψ_m under the action of conjugation, i.e.

$$\alpha^t \psi_m \alpha = \psi_m.$$

The group generated by all symplectic matrices is called the **symplectic group** and is denoted by $\operatorname{Sp}_{2m}(A) = \operatorname{Sp}_n(A)$, where n = 2m.

Definition 2.4. A matrix $\alpha \in \operatorname{GL}_{2m}(A)$ is called orthogonal if it fixes ψ_m under the action of conjugation, i.e.

$$\alpha^t \widetilde{\psi}_m \alpha = \widetilde{\psi}_m.$$

The group generated by all orthogonal matrices is called **orthogonal group** and is denoted by $O_{2m}(A) = O_n(A)$, where n = 2m.

Definition 2.5. The matrices of the form

$$\{se_{ij}(z) \in \mathrm{GL}_{2m}(A) : z \in A \mid i \neq j\},\$$

where

 $se_{ij}(z) = I_{2m} + ze_{ij}$ if $i = \sigma(j)$

or

$$se_{ij}(z) = I_{2m} + ze_{ij} - (-1)^{i+j} ze_{\sigma(j)\sigma(i)}$$
 if $i \neq \sigma(j)$ and $i < j$.

are called **symplectic elementary generators**. The subgroup generated by symplectic elementary generators is called **symplectic elementary group**.

Definition 2.6. The matrices of the form $\{oe_{ij}(z) : z \in A \mid i \neq j\}$, where

 $oe_{ij}(z) = I_{2m} + ze_{ij} - ze_{\sigma(j)\sigma(i)}, \text{ if } i \neq \sigma(j) \text{ and } i < j.$

are called orthogonal elementary generators.

The subgroup generated by orthogonal elementary generators is called **or**thogonal elementary group.

Remark 2.7. It is a well known fact that the elementary subgroups are normal subgroups of the respective classical groups; for $n \ge 3$ in the linear case, $n = 2m \ge 4$ in the symplectic case, and $n = 2m \ge 6$ in the orthogonal case. The linear case was due to A. Suslin (*cf.* [15]), symplectic was proved by V. Kopeiko (*cf.* [15]). Finally the orthogonal case was proved by Suslin–Kopeiko (*cf.* [14]).

We use the following notations to treat above three groups uniformly:

$$G(n, A)$$
 will denote $GL_n(A)$ or $Sp_{2m}(A)$ or $O_{2m}(A)$.

E(n, A) will denote $E_n(A)$ or $EO_{2m}(A)$ or $ESp_{2m}(A)$.

$$S(n, A)$$
 will denote $SL_n(A)$ or $SO_{2m}(A)$, and

 ge_{ij} will denote the elementary generators of $E_n(A)$ or $ESp_n(A)$ or $EO_n(A)$.

Here $SO_n(A)$ is the subgroup of $O_n(A)$ with determinant 1. Throughout the paper we will assume n = 2m to treat the non-linear cases. We shall assume

 $n \ge 3$ while treating the linear case, and $n \ge 6$; *i.e.* $m \ge 3$ while treating the symplectic and orthogonal cases. Let

$$\widetilde{v} = \begin{cases} v\psi_m \text{ in the symplectic case, and} \\ v\widetilde{\psi}_m \text{ in the orthogonal case.} \end{cases}$$

Definition 2.8. Let $v, w \in A^n$ be vectors of length n, then we define the inner product $\langle v, w \rangle$ as follows:

- (1) $\langle v, w \rangle = v^t w$ in the linear case,
- (2) $\langle v, w \rangle = \tilde{v}w$ in the symplectic and orthogonal cases.

Definition 2.9. Let $v, w \in A$, we define the map $M : A^n \times A^n \to M_n(A)$ as follows:

- (1) $M(v, w) = vw^t$ in the linear case,
- (2) $M(v,w) = v\widetilde{w} + w\widetilde{v}$ in the symplectic case,
- (3) $M(v,w) = v\widetilde{w} w\widetilde{v}$ in the orthogonal case.

By [G, G], we denote the commutator subgroup of G *i.e.* group generated by elements of the form $ghg^{-1}h^{-1}$, for $g, h \in G$.

Definition 2.10. A row $(v_1, \ldots, v_n) \in A^n$ is called **unimodular row of** length n if there exist $b_1, \ldots, b_n \in A$ such that $\prod_{i=1}^n v_i b_i = 1$; i.e. If I is the ideal generated by $\langle v_1, \ldots, v_n \rangle$, then (v_1, \ldots, v_n) is unimodular if and only if I = A. In the above case we say A has n-cover.

3. LOCAL-GLOBAL PRINCIPLE FOR CLASSICAL GROUPS

Before discussing the main theorem we recall following standard facts.

Lemma 3.1 (Splitting Lemma). The elementary generators of $n \times n$ matrices satisfy the following property:

$$ge_{ij}(x+y) = ge_{ij}(x) ge_{ij}(y)$$

for all $x, y \in A$ and for all i, j = 1, ..., n with $i \neq j$.

Proof. Standard. (*cf.* [3], \$3, Lemma 3.2).

Lemma 3.2. Let G be a group, and $a_i, b_i \in G$ for i = 1, ..., r. Let $J_k = \prod_{j=1}^{\kappa} a_j$. Then

$$\prod_{i=1}^{r} a_i b_i = \prod_{i=1}^{r} J_i b_i J_i^{-1} \prod_{i=1}^{r} a_i.$$

Proof. cf. [3], \$3, Lemma 3.4.

Following structural lemma pays a key role in the proof of main theorem. It is well-known for the polynomial rings; cf. [3] (\$3, Lemma 3.6). Here, we deduce the analogue for the graded rings.

Lemma 3.3. Let $G(n, A, A_+)$ denote the subgroup of G(n, A) which is equal to I_n modulo A_+ . Then the group $G(n, A, A_+) \cap E(n, A)$ is generated by the elements of the type $\epsilon ge_{ij}(A_+) \epsilon^{-1}$ for some $\epsilon \in E(n, A_0)$.

Proof. Let $\alpha \in E(n, A) \cap G(n, A, A_+)$. Then we can write

$$\alpha = \prod_{k=1}^{r} g e_{i_k j_k}(a_k)$$

for some elements $a_k \in A$, k = 1, ..., r. As $a_k = (a_0)_k + (a_+)_k$ for some $(a_0)_k \in A_0$ and $(a_+)_k \in A_+$. Using the splitting lemma (Lemma 3.1) we can rewrite the expression as:

$$\alpha = \prod_{k=1}^{r} \left(g e_{i_k j_k}(a_0)_k \right) \left(g e_{i_k j_k}(a_+)_k \right).$$

We write $\epsilon_t = \prod_{k=1}^t ge_{i_k j_k}((a_0)_k)$ for $t \in \{1, 2, \ldots, r\}$. Observe that $\epsilon_r = I_n$, as $\alpha \in G(n, A, A_+)$. Then

$$\alpha = \left(\prod_{k=1}^{r} \epsilon_k g e_{i_k j_k} \left((a_+)_k \right) \epsilon_k^{-1} \right) \left(\prod_{k=1}^{r} g e_{i_k j_k} \left((a_0)_k \right) \right) = \mathcal{AB} \text{ (say)},$$

where $\mathcal{A} = \left(\prod_{k=1}^{r} \epsilon_k g e_{i_k, j_k} ((a_+)_k) \epsilon_k^{-1}\right)$ and $\mathcal{B} = \left(\prod_{k=1}^{r} g e_{i_k j_k} ((a_0)_k)\right)$. Now go modulo A_+ . Let bar (-) denote the quotient ring modulo A_+ . Then,

$$\overline{\alpha} = \overline{I}_n = \overline{\mathcal{A}\mathcal{B}} = \overline{I}_n \overline{\mathcal{B}} = \overline{I}_n \implies \overline{\mathcal{B}} = \overline{I}_n,$$

as $\alpha \in G(n, A, A_+)$. Since entries of \mathcal{B} are in A_0 , it follows that $\mathcal{B} = I_n$.

$$\alpha = \prod_{k=1}^{r} \epsilon_k \left(g e_{i_k j_k} \left((a_+)_k \right) \right) \epsilon_k^{-1};$$

as desired.

Now we prove a variant of "Dilation Lemma" mentioned in the statement of (3) of Theorem 3.8.

Lemma 3.4. Assume the "Dilation Lemma" (Theorem 3.8 - (4)) to be true. Let $\alpha_s \in E(n, A_s)$, with $\alpha^+(0) = I_n$. Then one gets

$$\alpha^+(b+d)\alpha^+(d)^{-1} \in \mathcal{E}(n,A)$$

for some $s, d \in A_0$ and $b = s^l, l \gg 0$.

Proof. We have $\alpha_s \in E(n, A_s)$. Hence $\alpha_s^+(X) \in E(n, A_s[X])$. Let $\beta^+(X) = \alpha^+(X+d)\alpha^+(d)^{-1}$, where $d \in A_0$. Then $\beta_s^+(X) \in E(n, A_s[X])$ and $\beta^+(0) = I_n$. Hence by Theorem 3.8 - (4) their exists $\widetilde{\beta}(X) \in E(n, A[X])$ such that $\widetilde{\beta}_s(X) = \beta_s^+(bX)$. Putting X = 1, we get the required result.

Following is a very crucial result we need for our method of proof. There are many places we use this fact in a very subtle way. In particular, we mainly use this lemma for the step $(4) \Rightarrow (3)$ of 3.8.

Lemma 3.5. (cf.[10], Lemma 5.1) Let R be a Noetherian ring and $s \in R$. Then there exists a natural number k such that the homomorphism $G(n, s^k R) \rightarrow G(n, R_s)$ (induced by localization homomorphism $R \rightarrow R_s$) is injective. Moreover, it follows that the induced map $E(n, R, s^k R) \rightarrow E(n, R_s)$ is injective.

The next two lemma's will be used in intermediaries to prove the equivalent conditions mentioned in Theorem 3.8. We state it without proof.

Lemma 3.6. Let $v = (v_1, v_2, ..., v_n)$ be a unimodular row over over a commutative semilocal ring R. Then the row $(v_1, v_2, ..., v_n)$ is completable; i.e. $(v_1, v_2, ..., v_n)$ elementarily equivalent to the row (1, 0, ..., 0); i.e. their exists $\epsilon \in E_n(R)$ such that $(v_1, v_2, ..., v_n)\epsilon = (1, 0, ..., 0)$.

Proof. cf. [4], Lemma 1.2.21.

Lemma 3.7. Let A be a ring and $v \in E(n, A)e_1$. Let $w \in A^n$ be a column vector such that $\langle v, w \rangle = 0$. Then $I_n + M(v, w) \in E(n, A)$.

Proof. cf. [4], Lemma 2.2.4.

Theorem 3.8. The followings are equivalent for any graded ring $A = \bigoplus_{i=0}^{\infty} A_i$ for $n \ge 3$ in the linear case and $n \ge 6$ otherwise.

- (1) (Normality): E(n, A) is a normal subgroup of G(n, A).
- (2) If $v \in \text{Um}_n(A)$ and $\langle v, w \rangle = 0$, then $I_n + M(v, w) \in E(n, A)$.
- (3) (Local-Global Principle): Let $\alpha \in G(n, A)$ with $\alpha^+(0) = I_n$. If for every maximal ideal \mathfrak{m} of $A_0 \ \alpha_{\mathfrak{m}} \in E(n, A_{\mathfrak{m}})$, then $\alpha \in E(n, A)$.
- (4) (Dilation Lemma): Let $\alpha \in G(n, A)$ with $\alpha^+(0) = I_n$ and $\alpha_s \in E(n, A_s)$ for some non-zero divisor $s \in A_0$. Then there exists $\beta \in E(n, A)$ such that

$$\beta_s^+(b) = \alpha_s^+(b)$$

for some $b = s^l$; $l \gg 0$. i.e. $\alpha_s^+(s^l)$ will be defined over A for $l \gg 0$.

- (5) If $\alpha \in E(n, A)$, then $\alpha^+(a) \in E(n, A)$, for every $a \in A_0$.
- (6) If $v \in E(n, A)e_1$, and $\langle v, w \rangle = 0$, then $I_n + M(v, w) \in E(n, A)$.

Proof. (6) \Rightarrow (5): Let $\alpha = \prod_{k=1}^{t} (I_n + a \ M(e_{i_k}, e_{j_k}))$ where $a \in A$, and $t \ge 1$, a positive integer. Then

$$\alpha^{+}(b) = \prod_{k=1}^{t} (\mathbf{I}_{n} + a^{+}(b) \ M(e_{i_{k}}, e_{j_{k}})),$$

where $b \in A_0$. Take $v = e_i$ and $w = a^+(b)e_j$, Then $\alpha = I_n + M(v, w)$ and $\langle v, w \rangle = 0$. Indeed,

Linear case:

$$M(v, w) = vw^{t} = e_{i}(a^{+}(b)e_{j})^{t} = a^{+}(b)e_{i}e_{j}^{t} = a^{+}(b) M(e_{i}, e_{j}),$$

$$\langle v, w \rangle = v^{t}w = e_{i}^{t}(a^{+}(b)e_{j})^{t} = a^{+}(b) e_{i}^{t}e_{j} = a^{+}(b)\langle e_{i}, e_{j} \rangle = 0.$$

Symplectic case:

$$M(v,w) = vw^{t}\psi_{m} + wv^{t}\psi_{m} = e_{i}(a^{+}(b)e_{j})^{t}\psi_{m} + a^{+}(b)e_{j}e_{i}^{t}\psi_{m}$$

= $a^{+}(b)e_{i}e_{j}^{t}\psi_{m} + a^{+}(b)e_{j}e_{i}^{t}\psi_{m} = a^{+}(b) M(e_{i},e_{j}),$

$$\langle e_i, e_j \rangle = e_i^t \psi_m e_j = (e_i^t \psi_m e_i)(e_i^t e_j) = \psi_m(e_i^t e_j) = 0 \text{ hence,}$$

$$\langle v, w \rangle = v^t \psi_m w = e_i^t \psi_m a^+(b) e_j = a^+(b) e_i^t e_j = a^+(b) \langle e_i, e_j \rangle = 0.$$

Orthogonal case:

$$\begin{split} \mathbf{M}(v,w) &= v.w^t \widetilde{\psi}_m - w.v^t \widetilde{\psi}_m = e_i.(a^+(b)e_j)^t \widetilde{\psi}_m - a^+(b)e_j.e_i^t \widetilde{\psi}_m \\ &= a^+(b)e_i.e_j^t \widetilde{\psi}_m - a^+(b)e_j.e_i^t \widetilde{\psi}_m = a^+(b) \mathbf{M}(e_i,e_j), \\ \langle e_i,e_j \rangle &= e_i^t \widetilde{\psi}_m e_j = (e_i^t \widetilde{\psi}_m e_i)(e_i^t e_j) = \widetilde{\psi}_m(e_i^t e_j) = 0 \text{ hence,} \\ \langle v,w \rangle &= v^t \widetilde{\psi}_m w = e_i^t \widetilde{\psi}_m a^+(b)e_j = a^+(b)e_i^t e_j = a^+(b)\langle e_i,e_j \rangle = 0. \end{split}$$

Hence applying (6) over ring A, we have $\alpha^+(b) \in E(n, A)$. Therefore $\alpha^+(b) \in E(n, A)$ for $b \in A_0$.

(5) \Rightarrow (4): Since $\alpha_s \in E(n, A_s)$ with $(\alpha_{0s}) = I_n$, the diagonal entries of α are of the form $1 + g_{ii}$, where $g_{ii} \in (A_+)_s$ and off-diagonal entries are of the form g_{ij} , where $i \neq j$ and $g_{ij} \in (A_+)_s$.

We choose l to be large enough such that s^{l} is greater than the common denominator of all g_{ii} and g_{ij} . Then using (5), we get

$$\alpha_s^+(s^l) \in \mathcal{E}(n, A_s).$$

Since that $\alpha^+(s^l)$ permits a natural pullback (as denominators are cleared), we have $\alpha^+(s^l) \in E(n, A)$.

 $(4) \Rightarrow (3)$:

Since $\alpha_{\mathfrak{m}} \in \mathcal{E}(n, A_{\mathfrak{m}})$, we have an element $s \in A_0 - \mathfrak{m}$ such that $\alpha_s \in \mathcal{E}(n, A_s)$. Let $s_1, s_2, \ldots, s_r \in A_0$ be non-zero divisors with $s_i \in A_0 - \mathfrak{m}_i$ such that $\langle s_1, s_2, \ldots, s_r \rangle = A$. From (4) we have $\alpha^+(b_i) \in \mathcal{E}(n, A)$, for some $b_i = s_i^{l_i}$ with $b_1 + \cdots + b_r = 1$. Now consider $\alpha_{s_1s_2\ldots s_r}$, which is the image of α in $A_{s_1\cdots s_r}$. Due to Lemma 3.5, $\alpha \mapsto \alpha_{s_1s_2\cdots s_r}$ is injective and hence we can perform our calculation in $A_{s_1\cdots s_r}$ and then pull it back to A.

$$\alpha_{s_1s_2...s_r} = \alpha^+_{s_1s_2...s_r} (b_1 + b_2 \dots + b_r)$$

= $((\alpha_{s_1})_{s_2s_3...})^+ (b_1 + \dots + b_r) ((\alpha_{s_1})_{s_2s_3...})^+ (b_2 + \dots + b_r)^{-1} \dots$
 $((\alpha_{s_i})_{s_1...\hat{s_i}...s_r})^+ (b_i + \dots + b_r) ((\alpha_{s_i})_{s_1...\hat{s_i}...s_r})^+ (b_{i+1} + \dots + b_r)^{-1}$
 $\dots ((\alpha_{s_r})_{s_1s_2...s_{r-1}})^+ (b_r) ((\alpha_{s_r})_{s_1s_2...s_{r-1}})^+ (0)^{-1}$

Observing that each

$$((\alpha_{s_i})_{s_1s_2...\hat{s_i}...s_r})^+(b_i+\dots+b_r)((\alpha_{s_i})_{s_1...\hat{s_i}...s_r})^+(b_{i+1}+\dots+b_r)^{-1} \in \mathcal{E}(n,A)$$

due to Lemma 3.4 (here \hat{s}_i means we omit s_i in the product $s_1 \cdots \hat{s}_i \cdots s_r$), we have $\alpha_{s_1 \cdots s_r} \in E(n, A_{s_1 \cdots s_r})$ and hence $\alpha \in E(n, A)$.

Remark 3.9. Following is a commutative diagram (here we are assuming $\langle s_i, s_j \rangle = A$):



Let $\theta_i = \alpha_{s_i}^+(b_i + \dots + b_r)\alpha_{s_i}^+(b_{i-1} + \dots + b_r) \in A_{s_i}$ and θ_{ij} be the image of θ_i in $A_{s_is_j}$ and similarly $\pi_j = \alpha_{s_j}^+(b_j + \dots + b_r)\alpha_{s_j}^+(b_{j-1} + \dots + b_r) \in A_{s_j}$ and π_{s_i,s_j} be its image in $A_{s_is_j}$. Then due to Lemma 3.5 the product $(\theta_{s_is_j})(\pi_{s_is_j})$ can be identified with the product $\theta_i\pi_j$.

 $(3) \Rightarrow (2)$: Since polynomial rings are special case of graded rings, the result follows by using $3 \Rightarrow 2$ in [3], §3.

(2) \Rightarrow (1) \Rightarrow (6): The proof goes as in [3]. (\$3, 2 \Rightarrow 1 \Rightarrow 7 \Rightarrow 6).

4. LOCAL-GLOBAL PRINCIPLE FOR COMMUTATOR SUBGROUP

In this section we deduce the analogue of Local-Global Principle for the commutator subgroup of the linear and the symplectic group over graded rings. Unless mentioned otherwise, we assume $n \ge 3$ for the linear case and $n \ge 6$ for the symplectic case.

Let us begin with the following well-known fact for semilocal rings.

Lemma 4.1. Let A be a semilocal commutative ring with identity. Then for $n \ge 2$ in the linear case and $n \ge 4$ in the symplectic case, one gets

$$\mathbf{S}(n,A) = \mathbf{E}(n,A).$$

Proof. cf. Lemma 1.2.25 in [4].

Remark 4.2. If $\alpha = (\alpha_{ij}/1) \in G(n, A_s)$, then α has a natural pullback $\beta = (\alpha_{ij}) \in G(n, A)$ such that $\beta_s = \alpha$. If $\alpha_s \in S(n, A_s)$ such that it admits a natural pullback $\beta \in G(n, A)$, then $\beta \in S(n, A)$.

The next lemma deduces an analogue result of "Dilation Lemma" (Theorem 3.8-(3)) for S(n, A); the special linear (resp. symplectic) group.

Lemma 4.3. Let $\alpha = (\alpha)_{ij} \in S(n, A)$. Then $\alpha^+(a) \in S(n, A)$, where $a \in A_0$. Hence if $\alpha \in G(n, A)$ with $\alpha^+(0) = I_n$ and $\alpha_s \in S(n, A_s)$ then $\alpha_s^+(b) \in S(n, A)$ (after identifying $\alpha_s^+(b)$ with its pullback by using Lemma 3.5), where $s \in A_0$ is a non-zero divisor and $b = s^l$ with $l \gg 0$, for $n \ge 1$ in linear case and $n \ge 2$ in symplectic case.

Proof. Since det : $A \to A^*$ (units of A) $\subset A_0$, and $A_i A_j \subset A_{i+j}$, the nonzero component of α doesn't contribute for the value of the determinant and hence

$$\det \alpha = \det \alpha^+(0).$$

Therefore, if $\beta = \alpha^+(a)$, then

det
$$\beta = \det \beta^+(0) = \det (\alpha^+(a))^+(0) = \det \alpha^+(a \cdot 0) = \det \alpha^+(0) = 1.$$

Hence $\alpha^+(a) \in \mathcal{S}(n, A)$.

Since $\alpha_s \in S(n, A_s)$ with $\alpha^+(0) = I_n$, the diagonal entries are of the form $1+g_{ii}$, and off diagonal entries are g_{ij} , where $g_{ij} \in (A_+)_s$ for all i, j. We choose $b \in (s)$ such that $b = s^l$ with $l \gg 0$, so that b can dilute the denominator of each entries. Then $\alpha_s^+(b) \in S(n, A)$.

The next lemma gives some structural information about commutators.

Lemma 4.4. Let $\alpha, \beta \in S(n, A)$ and A_0 be a commutative semilocal ring. Then the commutator subgroup

$$[\alpha,\beta] \in [\alpha\alpha_0^{-1},\beta\beta_0^{-1}] \operatorname{E}(n,A).$$

Proof. Since A_0 is semilocal, we have $S(n, A_0) = E(n, A_0)$ by Lemma 4.1. Hence $\alpha^+(0), \beta^+(0) \in E(n, A_0)$. Let $a = \alpha \alpha^+(0)^{-1}$ and $b = \beta \beta^+(0)^{-1}$. Then

$$\begin{aligned} [\alpha,\beta] &= [\alpha\alpha^{+}(0)^{-1}\alpha^{+}(0),\beta\beta^{+}(0)^{-1}\beta^{+}(0)] \\ &= a\alpha^{+}(0)b\beta^{+}(0)\alpha^{+}(0)^{-1}a^{-1}\beta^{+}(0)^{-1}b^{-1} \\ &= (aba^{-1}b^{-1})(bab^{-1}\alpha^{+}(0)ba^{-1}b^{-1})(ba\beta^{+}(0)\alpha^{+}(0)^{-1}a^{-1}b^{-1})(b\beta^{+}(0)^{-1}b^{-1}). \end{aligned}$$

Since E(n, A) is a normal subgroup of S(n, A), the elements $bab^{-1}\alpha^+(0)ba^{-1}b^{-1}$, $ba\beta^+(0)\alpha^+(0)^{-1}a^{-1}b^{-1}$ and $b\beta^+(0)^{-1}b^{-1}$ are in E(n, A).

Corollary 4.5. Let $\alpha \in [S(n, A), S(n, A)]$ with $\alpha^+(0) = I_n$, and let A_0 be a semilocal commutative ring. Then using the normality property of the elementary (resp. elementary symplectic) subgroup α can be written as

$$\prod_{k=1}^{t} \left[\beta_k, \gamma_k\right] \epsilon,$$

for some $t \geq 1$, and $\beta_k, \gamma_k \in S(n, A)$, with $\beta_k^+(0) = \gamma_k^+(0) = I_n$, and $\epsilon \in E(n, A)$ with $\epsilon^+(0) = I_n$.

Proof. Since $\alpha \in [S(n, A), S(n, A)]$, $\alpha = \prod_{k=1}^{t} [a_k, b_k]$ for some $t \ge 1$. Using Lemma 4.4 we identify β_k with $a_k a_k^+(0)^{-1}$ and γ_k with $b_k b_k^+(0)^{-1}$. This gives

$$\alpha = \prod_{k=1}^{t} \left[\beta_k, \gamma_k \right] \epsilon$$

Then it follows that $\epsilon^+(0) = I_n$, as $\alpha^+(0) = I_n$.

The next lemma gives a variant of "Dilation Lemma" (Lemma 4.3) for S(n, A); the special linear (resp. symplectic) group.

Lemma 4.6. If $\alpha \in S(n, A_s)$ with $\alpha^+(0) = I_n$, then

$$\alpha^+(b+d)\alpha^+(d)^{-1} \in \mathcal{S}(n,A)$$

where s is a non-zero divisor and $b, d \in A_0$ with $b = s^i$ for some $i \gg 0$.

Proof. Let $\alpha^+(X) \in \mathcal{G}(n, A[X])$, then $\alpha_s^+(X) \in \mathcal{E}(n, A_s[X])$. Let $\beta^+(X) = \alpha^+(X+d)\alpha^+(d)^{-1}$. Then $\beta_s^+(X) \in \mathcal{S}(n, A_s[X])$. Hence by Lemma 4.3, $\beta^+(bX) \in \mathcal{S}(n, A)$. Putting X = 1, we get result.

The following lemma makes use of Lemma 4.3 to deduce "Dilation Lemma" for the commutator subgroup [S(n, A), S(n, A)].

Lemma 4.7. Let $\alpha_s = \prod_{i=1}^{t} [\beta_{is}, \gamma_{is}] \epsilon_s$ for some $t \ge 1$, such that β_{is} and $\gamma_{is} \in S(n, A_s)$ and $\epsilon_s \in E(n, A_s)$ with $\gamma_{is}^+(0) = \beta_{is}^+(0) = \epsilon_s^+(0) = I_n$. Then (by identifying with its pullback by using Lemma 3.5)

$$\alpha_s^+(b+d)\alpha_s^+(d)^{-1} \in [\mathcal{S}(n,A), \mathcal{S}(n,A)],$$

where $b, d \in A_0$ with $b = s^l$ for some $l \gg 0$.

Proof. Without loss of generality (as we would conclude by the end of the proof), we can assume

$$\alpha_s = [\beta_s, \gamma_s]\epsilon_s = \beta_s \gamma_s \beta_s^{-1} \gamma_s^{-1} \epsilon_s.$$

Since $\beta_s^+(0) = I_n$, by Lemma 4.3 $\beta_s^+(b), \gamma_s^+(b) \in S(n, A)$ for some $b = s^l$ with $l \gg 0$. Also, $\epsilon_s^+(b) \in E(n, A)$ by (5) of Theorem 3.8. Hence

$$\alpha_{s}^{+}(b+d)\alpha_{s}^{+}(d)^{-1} = \beta_{s}^{+}(b+d)\gamma_{s}^{+}(b+d)\beta_{s}^{+}(b+d)^{-1}\gamma_{s}^{+}(b+d)^{-1}$$
$$\epsilon_{s}^{+}(b+d)\epsilon_{s}^{+}(d)^{-1}\gamma_{s}^{+}(d)\beta_{s}^{+}(d)\gamma_{s}^{+}(d)^{-1}\beta_{s}^{+}(d)^{-1}.$$

Since $E(n, A_s)$ and $S(n, A_s)$ are normal subgroups in $G(n, A_s)$, by rearranging (using $\epsilon_s^+(b+d)\epsilon_s^+(d)^{-1}$ as intermediary) we can consider $\gamma_s^+(b+d)\gamma_s^+(d)^{-1}$ and $\beta_s^+(b+d)\beta_s^+(d)^{-1}$ together. Now by using Lemma 4.6 we get $\gamma_s^+(b+d)\gamma_s^+(d)^{-1}$

and $\beta_s^+(b+d)\beta_s^+(d)^{-1} \in \mathcal{S}(n,A)$. Hence (by identifying with its pullback by using Lemma 3.5) it follows that

$$\alpha_s^+(b+d)\alpha_s^+(d)^{-1} \in [\mathcal{S}(n,A), \mathcal{S}(n,A)].$$

Now we deduce the graded version of the Local-Global Principle for the commutators subgroups.

Theorem 4.8. Let $\alpha \in S(n, A)$ with $\alpha^+(0) = I_n$. If

$$\alpha_{\mathfrak{p}} \in [\mathcal{S}(n, A_{\mathfrak{p}}), \mathcal{S}(n, A_{\mathfrak{p}})] \text{ for all } \mathfrak{p} \in \operatorname{Spec}(\mathcal{A}_0),$$

then

$$\alpha \in [\mathcal{S}(n,A), \mathcal{S}(n,A)].$$

Proof. Since $\alpha_{\mathfrak{p}} \in [\mathcal{S}(n, A_{\mathfrak{p}}), \mathcal{S}(n, A_{\mathfrak{p}})] \subset [\mathcal{S}(n, A_{\mathfrak{m}}), \mathcal{S}(n, A_{\mathfrak{m}})]$, we have for a maximal ideal $\mathfrak{m} \supseteq \mathfrak{p}, s \in A_0 - \mathfrak{m}$ such that

$$\alpha_s \in [\mathcal{S}(n, A_s), \mathcal{S}(n, A_s)],$$

hence α_s can be decomposed as

$$\alpha_s = \prod_{i=1}^t [\beta_{i_s}, \gamma_{i_s}] \epsilon_s$$

by Corollary 4.5, where $\beta_{i_s}, \gamma_{i_s} \in \mathcal{S}(n, A_s)$ and $\epsilon_s \in \mathcal{E}(n, A_s)$. Now using the "Dilation Lemma" (Lemma 4.3 for $\mathcal{S}(n, A_s)$ and theorem 3.8 (4) for $\mathcal{E}(n, A_s)$) on each of these elements, we have for $b = s^i, i \gg 0$

$$\alpha_s^+(b) = \prod_{i=1}^t [\beta_{i_s}^+(b), \gamma_{i_s}^+(b)] \epsilon_s^+(b) \in [\mathbf{S}(n, A), \mathbf{S}(n, A)] E(n, A).$$

Since $E(n, A_s) \subseteq [S(n, A_s), S(n, A_s)]$, we have $\alpha_s \in [S(n, A_s), S(n, A_s)]$. Let $s_1, \ldots, s_r \in A_0$ non-zero divisors such that $s_i \in A_0 - \mathfrak{m}_i$ and $\langle s_1, \ldots, s_r \rangle = A_0$. Then it follows that $b_1 + \cdots + b_r = 1$ for suitable (as before) $b_i \in (s_i)$; and $i = 1, \ldots, r$. Now

$$\begin{aligned} \alpha_{s_1 s_2 \dots s_r} &= \alpha^+_{s_1 s_2 \dots s_r} (1) \\ &= \alpha^+_{s_1 s_2 \dots s_r} (b_1 + b_2 + \dots + b_r) \\ &= \left((\alpha_{s_1})_{s_2 s_3 s_4 \dots} \right)^+ (b_1 + \dots + b_r) \left((\alpha_{s_1})_{s_2 s_3 s_4 \dots} \right)^+ (b_2 + \dots + b_r)^{-1} \cdots \\ &\left((\alpha_{s_i})_{s_1 \dots \hat{s_i} \dots s_r} \right)^+ (b_i + \dots + b_r) \left((\alpha_{s_i})_{s_1 \dots \hat{s_i} \dots s_r} \right)^+ (b_{i+1} + \dots + b_r)^{-1} \\ &\cdots \left((\alpha_{s_r})_{s_1 s_2 \dots s_{r-1}} \right)^+ (b_r) \left((\alpha_{s_r})_{s_1 s_2 \dots s_{r-1}} \right)^+ (0)^{-1}. \end{aligned}$$

Using Lemma 3.5 the product is well defined (see Remark 3.9) and using Lemma 4.7, we conclude that $\alpha \in [S(n, A), S(n, A)]$.

5. Auxiliary Result for Transvection Subgroup

Let R be a commutative ring with 1. We recall that a finitely generated projective R-module Q has a unimodular element if their exists $q \in Q$ such that $qR \simeq R$ and $Q \simeq qR \oplus P$ for some projective R-module P.

In this section, we shall consider three types of (finitely generated) classical modules; *viz.* projective, symplectic and orthogonal modules over graded rings. Let $A = \bigoplus_{i=0}^{\infty} A_i$ be a commutative graded Noetherian ring with identity. For definitions and related facts we refer [5]; \$1, \$2. In that paper, the first author with A. Bak and R.A. Rao has established an analogous Local-Global Principle for the elementary transvection subgroup of the automorphism group of projective, symplectic and orthogonal modules of global rank at least 1 and local rank at least 3. In this article we deduce an analogous statement for the above classical groups over graded rings. We shall assume for every maximal ideal \mathfrak{m} of A_0 , the symplectic and orthogonal module $Q_{\mathfrak{m}} \simeq (A_0^{2m})_{\mathfrak{m}}$ with the standard form (for suitable integer m).

Remark 5.1. By definition the global rank or simply rank of a finitely generated projective *R*-module (resp. symplectic or orthogonal *R*-module) is the largest integer k such that $\oplus^k R$ (resp. $\perp^k \mathbb{H}(R)$) is a direct summand (resp. orthogonal summand) of the module. Here $\mathbb{H}(R)$ denotes the hyperbolic plane.

Let Q denote a projective, symplectic or orthogonal A_0 -module of global rank ≥ 1 , and total (or local) rank r+1 in the linear case and 2r+2 otherwise and let $Q_1 = Q \otimes_{A_0} A$.

We use the following notations to deal with the above three classical modules uniformly.

- (1) $G(Q_1) :=$ the full automorphism group of Q_1 .
- (2) $T(Q_1) :=$ the subgroup generated by transvections of Q_1 .
- (3) $ET(Q_1) :=$ the subgroup generated by elementary-transvections of Q_1 .
- (4) $\widetilde{E}(r, A) := E(r+1, A)$ for linear case and E(2r+2, A) otherwise.

In [5], the first author with Ravi A. Rao and A. Bak established the "Dilation Lemma" and the "Local-Global Principle" for the transvection subgroups of the automorphism group over polynomial rings. In this section, we generalize their results over graded rings. For the statements over polynomial rings we request the reader to look at Proposition 3.1 and Theorem 3.6 in [5].

Following is the graded version of the "Dilation Lemma".

Proposition 5.2 (Dilation Lemma). Let Q be a projective A_0 -module, and $Q_1 = Q \otimes_{A_0} A$. Let s be a non-nilpotent element in A_0 . Let $\alpha \in G(Q_1)$ with

 $\alpha^+(0) = I_n$. Suppose

$$\alpha_s \in \begin{cases} \mathbf{E}(r+1, (A_0)_s) & \text{ in the linear case,} \\ \mathbf{E}(2r+2, (A_0)_s) & \text{ otherwise.} \end{cases}$$

Then there exists $\tilde{\alpha} \in \text{ET}(Q_1)$ and $l \gg 0$ such that $\tilde{\alpha}$ localizes at $\alpha^+(b)$ for some $b = (s^l)$ and $\tilde{\alpha}^+(0) = I_n$.

Proof. Let $\alpha \in G(Q_1)$ and $\alpha(X) := \alpha^+(X) \in G(Q_1[X])$. Since $\alpha_s \in \widetilde{E}(r, A_s)$, $\alpha_s(X) \in \widetilde{E}(r, A_s[X])$. Since $\alpha(0) := \alpha^+(0) = I_n$, we can apply "Dilation Lemma" for the ring A[X] (cf.[5], Proposition 3.1) and hence there exists $\widetilde{\alpha}(X) \in \mathrm{ET}(Q_1[X])$ such that $\widetilde{\alpha}_s(X) = \alpha_s^+(bX)$ for some $b \in (s^l), l \gg 0$. Substituting X = 1, we get $\tilde{\alpha} \in \text{ET}(Q_1)$ and $\tilde{\alpha}_s := \tilde{\alpha}_s(1) = \alpha_s(b) = \alpha_s^+(b)$. This proves the proposition.

Lemma 5.3. If $\alpha \in G(Q_1)$ with $\alpha^+(0) = I_n$ such that $\alpha_s \in E(n, A_s)$, then $\alpha^+(b+d)\alpha^+(d)^{-1} \in ET(Q_1)$ for some $b \in (s^l)$ with $l \gg 0$, and $d \in A_0$.

Proof. Let $\beta = \alpha_s^+(b+d)\alpha_s^+(d)^{-1}$. Since $\alpha_s \in E(n, A_s)$ implies $\alpha^+(a) \in E(n, A_s)$ for $a \in A_s$ and hence $\beta \in E(n, A_s)$. Hence by Theorem 5.2, their exists a $\widetilde{\beta} \in \text{ET}(Q_1)$ such that $\widetilde{\beta}_s^+(b) = \beta$. Hence the lemma follows.

Theorem 5.4 (Local-Global Principle). Let Q be a projective A_0 -module, and $Q_1 = Q \otimes_{A_0} A$. Suppose $\alpha \in G(Q_1)$ with $\alpha^+(0) = I_n$. If

$$\alpha_{\mathfrak{m}} \in \begin{cases} \mathrm{E}(r+1, (A_0)_{\mathfrak{m}}) & \text{ in the linear case,} \\ \mathrm{E}(2r+2, (A_0)_{\mathfrak{m}}) & \text{ otherwise.} \end{cases}$$

for all $\mathfrak{m} \in \max(A_0)$, then $\alpha \in \mathrm{ET}(\mathrm{Q}_1) \subseteq \mathrm{T}(\mathrm{Q}_1)$.

Proof. Let $\alpha \in G(Q_1)$ with $\alpha^+(0) = I_n$. Since $\alpha_{\mathfrak{m}} \in \widetilde{E}(r, A_{\mathfrak{m}})$. Hence their exists a non-nilpotent $s \in A_0 - \mathfrak{m}$, such that $\alpha_s \in \widetilde{E}(r, A_s)$. Hence by the above "Dilation Lemma" there exists $\tilde{\alpha} \in \text{ET}(Q_1)$ such that $\tilde{\alpha}_s = \alpha_s^+(b)$, for some $b = s^l$ with $l \gg 0$. For each maximal ideal \mathfrak{m}_i we can find a suitable b_i . Since A is Noetherian, it follows that $b_1 + \cdots + b_r = 1$ for some positive integer r. Now we observe that $\alpha^+(b_i + \cdots + b_r)\alpha^+(b_{i+1} + \cdots + b_r)^{-1} \in \mathrm{ET}(Q_1)$ and hence calculating in the similar manner as we did it in Remark 3.9, we get

$$\alpha = \alpha^+(1) = \alpha^+(b_1 + \dots + b_r)\alpha^+(b_2 + \dots + b_r)^{-1} \cdots \alpha^+(b_i + \dots + b_r)$$

$$\alpha^+(b_{i+1} + \dots + b_r)^{-1} \cdots \alpha^+(0)^{-1} \in \operatorname{ET}(Q_1) \subseteq \operatorname{T}(Q_1);$$

lesired. \Box

as desired.

Acknowledgement

The second author expresses his sincere thanks to IISER Pune for allowing him to work on this project and providing him their infrastructural facilities.

References

- BAK, ANTHONY, AND NIKOLAI VAVILOV; "Structure of hyperbolic unitary groups I: elementary subgroups." In Algebra Colloquium, vol. 7, no. 2, pp. 159-196. Springer-Verlag, 2000
- [2] H. BASS; Algebraic K-Theory. W. A. Benjamin, Inc., New York, Amsterdam (1968).
- [3] RABEYA. BASU, RAVI A. RAO, REEMA KHANNA; On Quillen's Local-Global Principle, Commutative algebra and algebraic geometry, 17–30, Contemp. Math., 390, Amer. Math. Soc., Providence, RI, 2005.
- [4] RABEYA BASU; Results in Classical Algebraic K-theory, Ph.D. thesis, Tata Institute of Fundamental Research (2007), pp 70.
- [5] A. BAK, R. BASU, R.A. RAO; Local-Global Principle for transvection groups. Proceedings of The American Mathematical Society 138 (2010), no. 4, 1191–1204.
- [6] RABEYA BASU, RAVI A. RAO, REEMA KHANNA; On the equivalence of the relative Quillen-Suslin Local-Global Principle and normality of the relative elementary subgroup, (to appear) Proceeding of the conference Leavitt-path algebras and K-theory (CUSAT), Indian Statistical Institute series, Springer.
- [7] CHOUINARD L.G.; L.G., Projective modules over Krull semigroups, Michigan Math.J.29, 143-148, 1982.
- [8] J. GUBELADZE; Classical algebraic K-theory of monoid algebras, K-theory and homological algebra, Lecture Notes in Mathematics 137, 1990, 36–94.
- [9] J. GUBELADZE; Anderson's conjecture and the maximal monoid class over which projective modules are free, Math. USSR-Sb. 63 (1998), 165–180.
- [10] ROOZBEH HAZRAT, NIKOLAI VAVILOV; K_1 of Chevalley groups are nilpotent, Journal of Pure and Applied Algebra 179 (2003), no. 1-2, 99–116.
- [11] H. MATSUMURA; Commutative ring theory. Translated from the Japanese by M. Reid. Second edition. Cambridge Studies in Advanced Mathematics, 8. Cambridge University Press, Cambridge, 1989.
- [12] D. QUILLEN; Projective Modules over Polynomial Rings, Invent. Math. 36 (1976), 167– 171.
- [13] A.A. SUSLIN; On the structure of special linear group over polynomial rings, Math. USSR. Izv. 11 (1977), 221–238.
- [14] A.A. SUSLIN, V.I. KOPEIKO; Quadratic modules and Orthogonal groups over polynomial rings, Nauchn. Sem., LOMI 71 (1978), 216–250.
- [15] M.S. TULENBAEV; Schur multiplier of a group of elementary matrices of finite order, Zapiski Nauchnykh Seminarov Leningradskogo Otdeleniya Matematicheskogo Instuta im V.A. Steklova Akad. Nauk SSSR, Vol. 86 (1979), 162–169.
- [16] L.N. VASERSTEIN; On the normal subgroups of GL_n , over a ring, Algebraic K-Theory, Evanston (1980) (Proc. Conf., Northwestern Univ., Evanston III., (1980)), pp. 456–465, Lecture Notes in Math., 854, Springer, Berlin-New York, (1981).

Indian Institute of Science Education and Research (IISER Pune), India, email: rabeya.basu@gmail.com, rbasu@iiserpune.ac.in

Indian Institute of Science Education and Research (IISER Pune), India, email: manishks10100@gmail.com