# arXiv:1811.00829v1 [math.AP] 2 Nov 2018

# ON $C^{1,\alpha}$ -REGULARITY FOR CRITICAL POINTS OF A GEOMETRIC OBSTACLE-TYPE PROBLEM

# SUJIN KHOMRUTAI AND ARMIN SCHIKORRA

ABSTRACT. We consider critical points of the geometric obstacle problem on vectorial maps  $u: \mathbb{B}^2 \subset \mathbb{R}^2 \to \mathbb{R}^N$ 

$$\int_{\mathbb{B}^2} |\nabla u|^2 \quad \text{subject to } u \in \mathbb{R}^N \setminus \mathbb{B}^N(0).$$

Our main result is  $C^{1,\alpha}$ -regularity for any  $\alpha < 1$ .

Technically, we split the map  $u = \lambda v$ , where  $v : \mathbb{B}^2 \to \mathbb{S}^{N-1}$  is the vectorial component and  $\lambda = |u|$  the scalar component measuring the distance to the origin. While v satisfies a weighted harmonic map equation with weight  $\lambda^2$ ,  $\lambda$  solves the obstacle problem for

$$\int_{\mathbb{B}^2} |\nabla \lambda|^2 + \lambda^2 |\nabla v|^2, \quad \text{subject to } \lambda \ge 1.$$

where  $|\nabla v|^2 \in L^1(\mathbb{B}^2)$ . We then play ping-pong between the increases in the regularity of  $\lambda$  and v to obtain finally the  $C^{1,\alpha}$ -result.

# Contents

1.	Introduction	2
2.	Boundedness of $\lambda$	5
3.	The Euler-Lagrange equations for $v$	6
4.	Uniform a priori estimates for critical equations with elliptic $W^{1,2}$ -coefficients	7
5.	$W^{2,2-\varepsilon}$ -regularity of v	10
6.	$W^{2,2}$ -regularity for $\lambda$	14
7.	On $C^{1,\alpha}$ -regularity for $\lambda$	16
8.	Adaptations for the proof of Theorem $1.3$	18
Re	References	

### 1. INTRODUCTION

Denote by

$$D(u) := \int_{\mathbb{B}^2} |\nabla u|^2.$$

the Dirichlet energy for maps defined on the two-dimensional disk  $\mathbb{B}^2 \subset \mathbb{R}^2$ .

The classical obstacle problem for a given obstacle function  $\omega:\mathbb{B}^2\to\mathbb{R}$  analyzes the minimizer

$$\inf_{f \ge \omega} \int_{B^2} |\nabla f|^2$$

One can reformulate the obstacle problems for graphs u = (x, f(x)) as analyzing the minimizer of the problem

$$\inf_{X_{\Omega}} \int_{B^2} |\nabla u|^2,$$

where

$$\Omega = \left\{ (x,t) \in \mathbb{B}^2 \times \mathbb{R} : t < \varphi(x) \right\}$$

and the infimum is taken over the set of maps not touching  $\Omega$ .

(1.1) 
$$X_{\Omega} := \left\{ u \in H^1(\mathbb{B}^2, \mathbb{R}^3) : \ u \notin \Omega \right\}$$

It is a natural to consider this situation for sets  $\Omega$  whose boundary is smooth and compact, but which may not be a graph. In this case, u can be thought of as a soap film in threedimensional space which lives outside of a solid ball. Where the soap film intersects with the solid ball, a free boundary appears.

Geometric obstacle problems have been considered, e.g. [11] but this is quite different from our case. Much closer to our situation, considering minimizers, is the setup as in [10, 5]. Since the obstacle problem is not convex anymore, it is natural to consider not only minimizers but also critical points, which we shall do in this work.

A first observation is that the geometric setting immediately leads to regularity issues: while in the classical obstacle theory, basic  $C^{1,\alpha}$ -regularity is quite easy to obtain, already for the simplest case of round obstacles  $\Omega = \mathbb{B}^{N-1}$ , any harmonic function into  $\partial \mathbb{B}^{N-1} = \mathbb{S}^{N-1}$  is necessarily a critical point of the obstacle problem. Indeed we have,

**Proposition 1.1.** Let  $\bar{v}$  be a minimizing harmonic map from  $\mathbb{B}^n \to \mathbb{S}^{N-1}$  with respect to its own boundary values, then  $u := \bar{v}$  minimizes the Dirichlet energy in the class  $X_{\mathbb{B}^{N-1}}$  with respect to its own boundary values.

If  $\bar{v}$  is a critical (possibly non-minimizing) harmonic map from  $\mathbb{B}^2 \to \mathbb{S}^{N-1}$ , then  $\bar{v}$  is a critical map for the Dirichlet energy with respect to the class  $X_{\mathbb{B}^{N-1}}$ .

*Proof.* We split  $u = \lambda v$ , where  $\lambda = |u| \ge 1$  and  $v = \frac{u}{|u|} \in H^1(\mathbb{B}^2, \mathbb{S}^2)$ . Since  $v \cdot \nabla v \equiv 0$ , we have

$$|\nabla u|^2 = |\nabla \lambda v + \lambda \nabla v|^2 = |\nabla \lambda|^2 + |\lambda|^2 |\nabla v|^2$$

In particular,

$$\int_{\mathbb{B}^n} |\nabla u|^2 \ge \int |\nabla v|^2$$

with equality if and only if  $\lambda \equiv 1$ . The conclusion now follows.

In particular, for  $n \ge 3$  there is no hope of obtaining even mere continuity at the free boundary for the solutions of the obstacle problem: harmonic maps may only be smooth for  $n \ge 3$  on a large set (not everywhere), see [20, 5], and if we consider critical harmonic maps may be everywhere discontinuous, see [15].

This is why, for now, we shall restrict our attention to n = 2. The main result of this work is the basic regularity theory for spherical obstacles.

**Theorem 1.2.** Let  $\Omega = \mathbb{B}^{N-1} \subset \mathbb{R}^N$  be the solid unit ball. Denote the obstacle class  $X_\Omega$  as in (1.1). Then any map of which is critical in  $X_\Omega$  with respect to  $D(\cdot)$  is  $C^{1,\alpha}$ -smooth, for any  $\alpha < 1$ .

In future works we plan to analyze the free boundary, where u intersects with  $\partial\Omega$ , as well as more general obstacles.

Let us also state that as a by-product of our arguments we obtain the following regularity result for harmonic maps into the (non-compact) manifold of conformal transformations.

**Theorem 1.3.** Denote the group of conformal transformations with conformal factor bounded from below by  $\lambda_0$  as

$$CO_{\lambda_0}(N) = \left\{ \lambda \, Q \in \mathbb{R}^{N \times N} : Q \in SO(N), \lambda > \lambda_0 \right\}$$

Then for  $\lambda_0 > 0$ , any map  $P \in H^1(\mathbb{B}^2, CO_{\lambda_0}(N))$  which is a critical point of the Dirichlet energy  $D(\cdot)$  in the class of maps into  $CO_{\lambda_0}(N)$  belongs to  $C^{1,\alpha}$  for any  $\alpha < 1$ .

The proof is almost verbatim to the one of Theorem 1.2, we point out the differences in Section 8.

1.1. A reformulation of Theorem 1.2. In order to prove Theorem 1.2 we represent any point u in  $\mathbb{R}^N \setminus \mathbb{B}^N$  uniquely as

$$u = \lambda v,$$

where  $v = \frac{u}{|u|} \in \mathbb{S}^{N-1}$  and  $\lambda = |u| > 0$ .

If  $u \in H^1(\mathbb{B}^2, \mathbb{R}^n \setminus \mathbb{B}^N_1)$  then  $\lambda \in H^1(\mathbb{B}^2)$  is a scalar function and  $v \in H^1(\mathbb{B}^2, \mathbb{S}^{N-1})$ . In particular we have  $\langle v, \nabla v \rangle = 0$ , which leads to

$$|\nabla u|^2 = |\nabla \lambda v + \lambda \nabla v|^2 = |\nabla \lambda|^2 + \lambda^2 |\nabla v|^2.$$

Consequently, Theorem 1.2 can be reformulated as

**Theorem 1.4.** Let  $(\lambda, v) \in H^1(\mathbb{B}^2) \times H^1(\mathbb{B}^2, \mathbb{S}^{N-1})$  be a critical map with respect to the energy

$$E(\lambda, v) := \int |\nabla \lambda|^2 + \int \lambda^2 |\nabla v|^2$$

and subject to  $\lambda \geq \lambda_0$ . That is,

• assume that

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} E(\lambda + \varepsilon \varphi, v) \ge 0$$

holds whenever  $\varphi \in H_0^1(\mathbb{B}^2)$ , and  $\lambda + \varepsilon \varphi \geq \lambda$  almost everywhere in  $\mathbb{B}^2$  and  $(\lambda + \varepsilon \varphi)v \in H^1(\mathbb{B}^2)$  for small  $\varepsilon$ .

 $\bullet$  and

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} E\left(\lambda, \frac{v+\varepsilon\psi}{|v+\varepsilon\psi|}\right) = 0$$

holds for any  $\psi \in C_c^{\infty}(\mathbb{B}^2, \mathbb{R}^N)$ 

Then  $u = \lambda v \in C^{1,\alpha}$  for some  $\alpha > 0$ .

- **Remark 1.5.** By an easy adaptation of the proof one can show that  $\lambda \geq 1$  can be replaced by  $\lambda \geq \lambda_0$  where  $\lambda_0 \in C^{\infty}(\overline{\mathbb{B}^2}, (0, \infty))$  with  $\inf_{\mathbb{B}^2} \lambda_0 > 0$ . Observe that e.g. for starshaped obstacles the approach is much more complicated: Then one would need to assume  $\lambda \geq \lambda_0(v)$ , i.e. have to consider an obstacle depending on v, which heavily complicates the variation in v.
  - Moreover, observe that E as above is convex in λ, but not in v. That is, the only critical points in terms of λ are minimizers, but again not necessarily so v.

1.2. Outline of the proof. The proof of Theorem 1.4 is split into several parts, since we have to jump between improvements in regularity of  $\lambda$  and v. First we prove in Section 2 local boundedness of  $\lambda$ , see Proposition 2.1. Then we compute the Euler-Lagrange equations for v, in Section 3. Since by now we have shown that  $\lambda$  is locally bounded from above and below the Euler-Lagrange equations are uniformly elliptic equations with  $W^{1,2}$ -coefficients. We prove a priori  $L^p$ -estimates for such equations in Section 4, which might be interesting in their own right – see Proposition 4.1. In Section 5 we then obtain successively for v Hölder regularity, Proposition 5.1,  $W^{1,p}$ -regularity for any  $p < \infty$ , in Proposition 5.3 and finally  $W^{2,2-\varepsilon}$ -regularity in Corollary 5.5. This is the optimal regularity one can hope for without having better estimates on  $\lambda$ , see [22]. So in Section 6 we turn to improving the regularity  $\lambda$ , and the already obtained regularity for v allows us to obtain  $W^{2,2}$ -estimates for  $\lambda$  which in turn lead to  $W^{2,p}$ -estimate for v for any  $p < \infty$ , see Corollary 6.5. Lastly, with the regularity already obtained for  $\lambda$  and v we show in Section 7 that  $\lambda$  solves an elliptic inequality in viscosity sense, and we obtain  $C^{1,\alpha}$ -regularity of  $\lambda$ . With this we conclude the promised regularity of  $u = \lambda v$ .

## 2. Boundedness of $\lambda$

The scalar function  $\lambda$  is a solution to a classical (graph-)obstacle problem, however for the energy

$$\lambda \mapsto \int |\nabla \lambda|^2 + \int \lambda^2 |\nabla v|^2$$

But observe that  $|\nabla v|^2 \in L^1(\mathbb{R}^n)$ , only. In particular, a priori for general  $|\nabla v|^2 \in L^1(\mathbb{R}^n)$ , we cannot hope that  $\lambda$  is very smooth. For now we have to content ourselves with the (local) boundedness of  $\lambda$ .

**Proposition 2.1** (Boundedness of  $\lambda$ ). Let  $\lambda$ , v be as in Theorem 1.4. Then  $\lambda \in L^{\infty}_{loc}(\mathbb{B}^2)$ , that is for any compact set  $K \subset \mathbb{B}^2$  we have that  $\lambda \in L^{\infty}(K)$ .

*Proof.* We will show that  $\lambda \in L^{\infty}(B(0,r))$  for any  $r \in (0,1)$ . Fix such an r. By Fubini's theorem, there must be  $R \in (r,1)$  such that

$$\|\lambda\|_{H^1(\partial B(0,R))} \precsim \frac{1}{1-r} \|\lambda\|_{H^1(B(0,1))}$$

Since  $\partial B(0, r)$  is one-dimensional we have that  $H^1(\partial B(0, R))$  embeds in particular into  $C^0(\partial B(0, R))$ . For simplicity of notation we shall pretend that R = 1 and thus assume w.l.o.g.

$$K_1 := \|\lambda\|_{L^{\infty}(\partial B(0,1))} < \infty.$$

Let  $\eta \in C_c^{\infty}(\mathbb{B}^2, \mathbb{R}_+)$  and let  $K > K_1$ . Then for small  $\varepsilon > 0$  the following variation of  $\lambda$  is admissible

$$\lambda_{\varepsilon} := \lambda - \varepsilon \eta (\lambda - K)$$

Indeed, by convexity, whenever  $\varepsilon \|\eta\|_{\infty} < 1$ ,

$$\lambda_{\varepsilon} = (1 - \varepsilon \eta)\lambda + \varepsilon \eta K \ge 1$$
 a.e. in  $\mathbb{B}^2$ .

In particular, the Euler-Lagrange inequality for  $\lambda$  implies

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0^+} E\left(\lambda_{\varepsilon}, v\right) \ge 0,$$

that is

(2.1) 
$$\int_{\mathbb{B}^2} |\nabla \lambda|^2 \eta + \int_{\mathbb{B}^2} (\lambda - K) \nabla \lambda \cdot \nabla \eta + \int_{\mathbb{B}^2} \lambda \eta (\lambda - K) |\nabla v|^2 \le 0.$$

We would like to test this inequality with  $\eta := (\lambda - K)_+$  Then  $\eta \in H_0^1(\mathbb{B}^2)$  – the zero boundary data stems from the choice of  $K \ge K_1$ . Moreover,

$$\nabla \eta = \chi_{\{\lambda \ge K\}} \, \nabla \lambda.$$

Cf. [6, Chapter 5, Problem 18, p.308]. However  $\eta$  may not be bounded, and the resulting integrals may not converge. So instead for arbitrary k > K we test with

$$\eta_k := -(\eta - k)_- + k = \min\{\eta - k, 0\} + k \in [0, k]$$

In other words,

$$\eta_k = \begin{cases} k & \text{in } \{\lambda > K + k\} \\ \lambda - K & \text{in } \{\lambda < K + k\} \cap \{\lambda > K\} \\ 0 & \text{in } \{\lambda < K\} \end{cases}$$

Now we have  $\eta_k \in H_0^1 \cap L^\infty(\mathbb{B}^2, \mathbb{R}_+)$ , and consequently,  $\eta_k$  is admissible as testfunction in (2.1). Moreover,

(2.2) 
$$\nabla \eta_k = \chi_{\{\lambda \in (K, K+k)\}} \nabla \lambda.$$

We observe that  $(K - \lambda)\eta_k \ge 0$  and thus

$$\int_{\mathbb{B}^2} \lambda \, \eta \left( K - \lambda \right) |\nabla v|^2 \ge 0$$

Moreover, in view of (2.2),

$$\int_{\mathbb{B}^2} (\lambda - K) \nabla \lambda \cdot \nabla \eta = \int_{\mathbb{B}^2} (\lambda - K) \chi_{\lambda \in (K,k)} |\nabla \lambda|^2 \ge 0.$$

Consequently, (2.1) implies

$$\int_{\mathbb{B}^2} |\nabla \lambda|^2 \eta_k \le 0,$$

that is, since  $\eta_k \ge 0$ ,

$$|\nabla \lambda|^2 \eta_k \equiv 0.$$
$$\int_{\mathbb{B}^2} |\nabla \lambda|^2 (\lambda - K)_+ \le 0$$

0

This implies

$$|\nabla \lambda|^2 (\lambda - K)_+ \equiv 0,$$

that is

$$\left|\nabla\left((\lambda - K)_{+}\right)^{2}\right| \equiv 0,$$

But in view of (2.2) this implies

$$|\nabla(\eta_k)^2| \equiv 0,$$

which in turn gives  $\eta_k \equiv 0$  (since  $\eta_k \in H_0^1(\mathbb{B}^2)$ ). In particular  $\lambda \leq K$  almost everywhere, i.e.  $\lambda$  is bounded (recall that  $\lambda \geq \lambda_0 \in L^{\infty}(\mathbb{B})$  was assumed). 

# 3. The Euler-Lagrange equations for v

Now that  $\lambda$  is bounded, we start with computing the Euler-Lagrange equations for v, which are a weighted version of the spherical harmonic map equation. In particular we obtain a weighted version of Shatah's conservation law [23], that Hélein used in [9] to obtain regularity for harmonic maps into spheres.

**Lemma 3.1** (Euler-Lagrange equations). Let  $\lambda$  and v be as in Theorem 1.4. Then,

(3.1)  $\operatorname{div}(\lambda^2 \nabla v^i) = \Omega_{ij} \cdot \lambda^2 \nabla v^j$ 

with

$$\Omega_{ij} = v^j \nabla v^i - v^i \nabla v^j.$$

Equivalently we also have a weighted version of Shatah's conservation law [23] (3.2)  $\operatorname{div}(\lambda^2 \Omega_{ij}) = 0.$ 

*Proof.* Since  $|v| \equiv 1$  we have

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0}\frac{v+\varepsilon\psi}{|v+\varepsilon\psi|} = \psi - \langle\psi,v\rangle v,$$

and consequently, the Euler-Lagrange equations with respect to v, can be written as

$$\int_{\mathbb{B}^2} \lambda^2 \, \nabla v \cdot \nabla \psi = \int_{\mathbb{B}^2} \lambda^2 \, \nabla v \cdot \nabla (\langle \psi, v \rangle v)$$

Now,  $v \cdot \nabla v \equiv 0$  since  $|v| \equiv 1$ , so

$$\nabla v \cdot \nabla (\langle \psi, v \rangle v) = |\nabla v|^2 \langle \psi, v \rangle.$$

We thus obtain the Euler-Lagrange equation

$$\operatorname{div}(\lambda^2 \nabla v^i) = \lambda^2 v^i |\nabla v|^2.$$

Now we rewrite this equation (using again  $v^i \nabla v^i = \frac{1}{2} \nabla |v|^2 \equiv 0$ ), with he following trick

$$v^{i}|\nabla v|^{2} = v^{i} \nabla v^{j} \cdot \nabla v^{j} = \left(v^{i} \nabla v^{j} \cdot \nabla v^{j} - v^{j} \nabla v^{i}\right) \cdot \nabla v^{j}$$

This establishes (3.1). The conservation law (3.2) follows now from a direct computation.  $\Box$ 

# 4. Uniform a priori estimates for critical equations with elliptic $W^{1,2}$ -coefficients

**Proposition 4.1.** Let  $2 < p_0 < p_{\infty} < \infty$  and  $\Lambda > 1$ . Then there exists a constant  $C = C(\Lambda, p_0, p_{\infty})$ , a small  $\varepsilon = \varepsilon(\Lambda, p_0, p_{\infty}) > 0$  and a small  $\alpha = \alpha(\Lambda, p_0, p_{\infty}) > 0$  so that the following holds.

Let either p = 2 or  $p \in (p_0, p_\infty)$  and R > 0. Let  $v \in W^{1,2}(\mathbb{B}^2, \mathbb{R}^N)$  be a solution to (4.1)  $\operatorname{div}(\lambda^2 \nabla v^i) = \Omega_{ij} \cdot \lambda^2 \nabla v^j$  in  $\mathbb{B}^2$ 

where  $\lambda \in L^{\infty} \cap W^{1,2}(\mathbb{B}^2)$  satisfies

(4.2)  $\Lambda^{-1} \leq \lambda \leq \Lambda \quad almost \; everywhere \; in \; \mathbb{B}^2$ 

and  $\Omega_{ij} \in L^2_{loc}(\mathbb{B}^2, \mathbb{R}^2)$  satisfies

(4.3)  $|\Omega| \leq \Lambda |\nabla v|$  almost everywhere in  $\mathbb{B}^2$ .

If p = 2 we assume moreover

(4.4) 
$$\operatorname{div}(\lambda^2 \Omega_{ij}) = 0 \quad in \ \mathbb{B}^2.$$

Then, if  $\nabla v \in L^p_{loc}(\mathbb{B}^2)$  then for any r < R the estimate

$$\|\nabla v\|_{L^p(B(r))} \le C \left(\frac{r}{R}\right)^{\frac{\alpha}{p}} \|\nabla v\|_{L^p(B(R))}$$

holds for all balls  $B(R) \subset \mathbb{B}^2$  on which v and  $\lambda$  satisfy

$$\|\nabla\lambda\|_{L^2(B(R))} + \|\nabla v\|_{L^2(B(R))} \le \varepsilon.$$

An important ingredient for the p = 2 case is Wente's Lemma see [14, 26, 2, 24, 13, 4, 25].

**Lemma 4.2** (Wente Lemma). Let  $B \subset \mathbb{R}^2$  be a ball, and  $a, b \in W^{1,2}(B)$ . If  $w \in W^{1,2}(B(R))$  is a solution to

$$\begin{cases} \Delta w = \nabla a \cdot \nabla^{\perp} b & \text{ in } B\\ w = 0 & \text{ on } \partial B, \end{cases}$$

where  $\nabla^{\perp} = (-\partial_2, \partial_1)^T$ , then

$$||w||_{L^{\infty}(B)} + ||\nabla w||_{L^{2}(B)} \le ||\nabla a||_{L^{2}(B)} ||\nabla b||_{L^{2}(B)}.$$

The proof of Proposition 4.1 is based on the following estimate.

**Lemma 4.3.** Let  $2 < p_0 < p_{\infty} < \infty$  and  $\Lambda > 1$ . Then there exists a constant  $C = C(\Lambda, p_0, p_{\infty})$  so that the following holds.

Let either p = 2 or  $p \in (p_0, p_\infty)$  and R > 0. Let  $v \in W^{1,2}(B(R), \mathbb{R}^N)$  be a solution to (4.1) in B(R), where  $\lambda \in L^{\infty} \cap W^{1,2}(B(R))$  satisfies (4.2) in B(R) and  $\Omega_{ij} \in L^2_{loc}(B(R), \mathbb{R}^2)$ satisfies (4.3). If p = 2 we assume moreover (4.4) to hold in B(R).

Then, if  $\nabla v \in L^p(B(R))$  we have the following a priori estimate for any  $r \in (0, R]$ 

$$\|\nabla v\|_{L^{p}(B(r))} \leq C \left( \left(\frac{r}{R}\right)^{\frac{2}{p}} + \|\nabla \lambda\|_{L^{2}(B(R))} + \|\nabla v\|_{L^{2}(B(R))} \right) \|\nabla v\|_{L^{p}(B(R))}.$$

*Proof.* We use Hodge decomposition to obtain

(4.5) 
$$\lambda^2 \nabla v = \nabla a + \nabla^{\perp} b + H \quad \text{in } B(R)$$

Here  $\nabla^{\perp} = (-\partial_y, \partial_x)$ . Namely, we choose  $a, b \in W_0^{1,2}(B(R), \mathbb{R}^N)$ , and H harmonic in B(R) so that

$$\begin{cases} \Delta a = \operatorname{div}(\lambda^2 \nabla v) & \text{ in } B(R) \\ a = 0 & \text{ on } \partial B(R) \end{cases}, \quad \begin{cases} \Delta b = \operatorname{curl}(\lambda^2 \nabla v) & \text{ in } B(R) \\ b = 0 & \text{ on } \partial B(R) \end{cases}$$

From (4.1) we find that

$$\begin{cases} \Delta a = \Omega_{ij} \cdot \lambda^2 \nabla v^j & \text{in } B(R) \\ a = 0 & \text{on } \partial B(R) \end{cases}$$

From standard elliptic estimates we then obtain for any p > 2

(4.6) 
$$\|\nabla a\|_{L^{p}(B(R))} \preceq \|\lambda\|_{\infty}^{2} \|\Omega\|_{L^{2}(B(R))} \|\nabla v\|_{L^{p}(B(R))}.$$

Of course the constant may depend on p as it blows up for  $p \to \infty$  or as  $p \to 2$ . But it is uniform for  $p \in (p_0, p_\infty)$ . For p = 2, we use that by (4.4) we have a div-curl structure. Then, Wente's Lemma, Lemma 4.2, implies the same estimate (4.6) for p = 2.

For b we use compute the curl and find

$$\begin{cases} \Delta b = \nabla^{\perp}(\lambda^2) \nabla v & \text{ in } B(R) \\ b = 0 & \text{ on } \partial B(R) \end{cases}$$

Again from standard elliptic estimates for p > 2 and from Wente's Lemma and the div-curl structure for p = 2 we obtain the estimate

 $\|\nabla b\|_{L^p(B(R))} \precsim \|\lambda\|_{L^\infty(B(R))} \|\nabla \lambda\|_{L^2(B(R))} \|\nabla v\|_{L^p(B(R))}$ 

By the assumptions on  $\lambda$  and  $\Omega$  we thus get

 $\|\nabla a\|_{L^{p}(B(R))} + \|\nabla b\|_{L^{p}(B(R))} \leq C(\Lambda, p_{0}, p_{\infty}) \left(\|\nabla v\|_{L^{2}(B(R))} + \|\nabla\lambda\|_{L^{2}(B(R))}\right) \|\nabla v\|_{L^{p}(B(R))}.$ In particular we get from (4.5),

 $\|\nabla v\|_{L^{p}(B(r))} \leq C(\Lambda) \|H\|_{L^{p}(B(r))} + C(\Lambda, p_{0}, p_{\infty}) \left(\|\nabla v\|_{L^{2}(B(R))} + \|\nabla\lambda\|_{L^{2}(B(R))}\right) \|\nabla v\|_{L^{p}(B(R))},$ and

$$\|H\|_{L^{p}(B(R))} \leq C(\Lambda) \|\nabla v\|_{L^{p}(B(R))} + C(\Lambda, p_{0}, p_{\infty}) \left(\|\nabla v\|_{L^{2}(B(R))} + \|\nabla\lambda\|_{L^{2}(B(R))}\right) \|\nabla v\|_{L^{p}(B(R))},$$
  
The last ingredient is the harmonicity of  $H$ , which implies for any  $r < R$ , see, e.g. [8, Theorem 2.1, p.78],

$$||H||_{L^p(B(r))} \precsim \left(\frac{r}{R}\right)^{\frac{2}{p}} ||H||_{L^p(B(R))}.$$

Together, the last three estimates imply the claimed result.

By choosing  $r < \theta^{\frac{p}{n}} R$  for  $\theta$  small enough we obtain as a corollary

**Corollary 4.4.** Let  $2 < p_0 < p_{\infty} < \infty$  and  $\Lambda > 1$ . Then there exists a constant  $C = C(\Lambda, p_0, p_{\infty})$ , a small  $\varepsilon = \varepsilon(\Lambda, p_0, p_{\infty}) > 0$  and a small  $\theta = \theta(\Lambda, p_0, p_{\infty})$  so that the following holds.

Let either p = 2 or  $p \in (p_0, p_\infty)$  and R > 0. Let  $v \in W^{1,2}(B(R), \mathbb{R}^N)$  be a solution to (4.1) in B(R), where  $\lambda \in L^{\infty} \cap W^{1,2}(B(R))$  satisfies (4.2) in B(R) and  $\Omega_{ij} \in L^2_{loc}(B(R), \mathbb{R}^2)$ satisfies (4.3) in B(R). If p = 2 we assume moreover (4.4) to hold in B(R). Then, if  $\nabla v \in L^p(B(R))$  and if

$$\|\nabla \lambda\|_{L^{2}(B(R))} + \|\nabla v\|_{L^{2}(B(R))} < \varepsilon$$

then for  $\sigma := \theta^{\frac{p}{2}}$  we have

$$\|\nabla v\|_{L^{p}(B(\sigma R))} \leq \frac{1}{2} \|\nabla v\|_{L^{p}(B(R))}$$

Proof of Proposition 4.1. The proof now follows from Corollary 4.4 by iteration. Pick  $r \in (\sigma^{k-1}R, \sigma^k R]$  for some  $k \in \mathbb{N}$ .

For now let us assume that  $k \ge 2$ . Repeated application of Corollary 4.4 implies

$$\|\nabla v\|_{L^p(B(r))} \le 2^{1-k} \|\nabla v\|_{L^p(B(R))}$$

Since for our choice of r,

$$2^{1-k} = \sigma^{(k-1)\frac{\log 2}{-\log \sigma}} \le \left(\frac{r}{R}\right)^{\frac{\log 2}{-\log \sigma}}$$

we have found that

$$\|\nabla v\|_{L^p(B(r))} \le \left(\frac{r}{R}\right)^{\frac{\log 2}{-\log \sigma}} \|\nabla v\|_{L^p(B(R))}.$$

Since  $\sigma = \theta^{\frac{p}{2}}$  we choose (independently of p)

$$\alpha := 2 \frac{\log 2}{-\log \theta}$$

That is, we have shown

$$\|\nabla v\|_{L^p(B(r))} \le \left(\frac{r}{R}\right)^{\frac{\alpha}{p}} \|\nabla v\|_{L^p(B(R))}$$

holds for any  $r \leq \sigma^2 R$ . For  $r \in (\sigma^2 R, R)$  we use the trivial estimate

$$\|\nabla v\|_{L^{p}(B(r))} \leq \|\nabla v\|_{L^{p}(B(R))} \leq \sigma^{-2\frac{\alpha}{p}} \left(\frac{r}{R}\right)^{\frac{\alpha}{p}} \|\nabla v\|_{L^{p}(B(R))}.$$

Using again that  $\sigma = \theta^{\frac{p}{2}}$  we find for any  $r \in (\sigma^2 R, R)$ 

$$\|\nabla v\|_{L^p(B(r))} \le \|\nabla v\|_{L^p(B(R))} \le \theta^{-2\alpha} \left(\frac{r}{R}\right)^{\frac{\alpha}{p}} \|\nabla v\|_{L^p(B(R))}$$

# 5. $W^{2,2-\varepsilon}$ -regularity of v

As a consequence of our analysis in the previous section we obtain Hölder continuity of v.

**Proposition 5.1** (Initial regularity for v). Let v and  $\lambda$  be as in Theorem 1.4. Then there exists  $\alpha > 0$  such that for every compact  $K \subset \mathbb{B}^2$  we have

(5.1) 
$$\sup_{B(y_0,r)\subset K} r^{-\frac{\alpha}{2}} \|\nabla v\|_{L^2(B(y_0,r))} < \infty.$$

In particular, by Sobolev embedding in  $\mathbb{R}^2$ ,  $v \in C^{0,\alpha}_{loc}$ .

**Remark 5.2.** The proof of Hölder continuity can be found in the literature: from Lemma 3.1, more precisely (3.1) we obtain that for  $\xi^i := \lambda^2 \nabla u^i$  we have

$$\operatorname{div}(\xi^i) = \Omega_{ik}\xi^k$$

Hölder regularity now follows from a distorted version of Rivière's celebrated regularity theorem for systems with antisymmetric potential [16]. More precisely, [17, Remark 3.4.] is applicable – since  $\lambda \in L^{\infty}_{loc}$  by Proposition 2.1 and using also that by assumption  $\inf_{\mathbb{R}^2} \lambda > 0$ .

In order to obtain later higher regularity, however, we need the estimate (5.1).

Proof of Proposition 5.1. For 0 < r < R let  $B(y_0, r) \subset B(y_0, R) \subset \mathbb{B}^2$ . Since our result is away from the boundary, by Proposition 2.1 we may assume w.l.o.g. that  $\lambda$  is bounded in all of  $\mathbb{B}^2$ .

Observe that since  $\lambda, v \in W^{1,2}(\mathbb{B}^2)$ , by absolute continuity of the integral, for any  $\varepsilon > 0$  there exists a radius  $R_0 > 0$  such that

$$\sup_{B(y_0,\rho)\subset\mathbb{B}^2,\rho< R_0} \|\nabla v\|_{L^2(B(y_0,\rho))} + \|\nabla\lambda\|_{L^2(B(y_0,\rho))} < \varepsilon.$$

The claim then follows from the a priori estimates of Proposition 4.1 (for p = 2) and a covering argument.

5.1. Slightly higher integrability of the gradient of v. The next step is higher integrability of the derivative  $\nabla v$ ,

**Proposition 5.3**  $(W^{1,2+\varepsilon}$ -regularity for v). Let v and  $\lambda$  be as in Theorem 1.4. Then, there exists p > 2 such that  $v \in W^{1,p}_{loc}(\mathbb{B}^2)$ .

*Proof.* We apply Hodge decomposition on a ball B(R). Namely we split

(5.2)  $\lambda^2 \nabla v = \nabla a + \nabla^{\perp} b + H \quad \text{in } B(R)$ 

where H is harmonic in B(R) and a and b are chosen as follows (in view of Lemma 3.1):

$$\begin{cases} \Delta a = \operatorname{div}(\lambda^2 \nabla v) = \Omega \lambda^2 \nabla v & \text{in } B(R) \\ a = 0 & \text{on } \partial B(R) \end{cases}, \quad \begin{cases} \Delta b = \operatorname{curl}(\lambda^2 \nabla v) = \nabla^{\perp} \lambda^2 \nabla v & \text{in } B(R) \\ b = 0 & \text{on } \partial B(R) \end{cases}$$

With the  $\alpha$  from Proposition 5.1, the structure of  $\Omega$ , and boundedness of  $\lambda$  we obtain

$$\sup_{B(r)\subset B(R)} r^{-\alpha} \|\Delta a\|_{L^1(B(r))} < \infty.$$

but for b, since  $\nabla \lambda \in L^2$  only, we find

$$\sup_{B(r)\subset B(R)} r^{-\frac{\alpha}{2}} \|\Delta b\|_{L^1(B(r))} < \infty.$$

By (a localized version of) the Sobolev embedding for Morrey spaces, see [1], we obtain that for any  $p \in [1, \frac{2-\alpha}{1-\alpha})$ ,  $\nabla a$  and  $\nabla b$  belong to  $L^p_{loc}(B(R))$ . Since  $\alpha > 0$  we can choose p > 2, and since H is harmonic on B(R) and  $\lambda$  is bounded away from zero, from (5.2) we get  $\nabla v \in L^p_{loc}(B(R))$ .

5.2. On integrability of the gradient of v and  $W^{2,2-\varepsilon}$ -regularity. Now we can (still only assuming that  $\lambda \in W^{1,2}$ ) bootstrap the regularity for v all the way to  $W_{loc}^{1,p}$ ,  $p \in (1,\infty)$ . For this we adapt an iteration strategy by Sharp and Topping [22], see also generalizations in [21, 18]. The main technical ingredient are the uniform a priori estimates in Proposition 4.1.

**Proposition 5.4**  $(W^{1,p}$ -regularity for v for large p). Let v and  $\lambda$  be as in Theorem 1.4. Then, for any  $p \in (1, \infty)$  we have  $v \in W^{1,p}_{loc}(\mathbb{B}^2, \mathbb{R}^N)$ .

Proof of Proposition 5.4. Fix  $p_{\infty} \in (2, \infty)$ . We are going to show that  $v \in W_{loc}^{1, p_{\infty}}(\mathbb{B}^2)$ .

By Proposition 5.3 we have  $v \in W_{loc}^{1,p_1}(\mathbb{B}^2)$ . Set  $p_0 := \frac{2+p_1}{2}$ , and apply Proposition 4.1, then for some (uniform)  $\alpha$ ,

(5.3) 
$$\sup_{B(r)\subset K} r^{-\frac{\alpha}{p_1}} \|\nabla v\|_{L^{p_1}(B(r))} < \infty.$$

As in (5.2) we apply Hodge decomposition on some ball  $B(R) \subset \mathbb{B}^2$ .

(5.4) 
$$\lambda^2 \nabla v = \nabla a + \nabla^{\perp} b + H \quad \text{in } B(R)$$

where H is harmonic in B(R) and in view of Lemma 3.1 we have

$$\begin{cases} \Delta a = \Omega \lambda^2 \nabla v & \text{in } B(R) \\ a = 0 & \text{on } \partial B(R) \end{cases}, \quad \begin{cases} \Delta b = \nabla^{\perp} \lambda^2 \nabla v & \text{in } B(R) \\ b = 0 & \text{on } \partial B(R) \end{cases}$$

From (5.3) we obtain

$$\sup_{B(r)\subset B(R)} r^{-2\frac{\alpha}{p_1}} \|\Delta a\|_{L^{\frac{p_1}{2}}(B(r))} < \infty$$

and (recall that we only have  $\nabla \lambda \in L^2$ ),

$$\sup_{B(r)\subset B(R)} r^{-\frac{\infty}{p_1}} \|\Delta b\|_{L^{\frac{2p_1}{p_1+2}}(B(r))} < \infty$$

Observe that since  $p_1 > 2$  we have  $\frac{p_1}{2} > q_1 := \frac{2p_1}{2p_1+2} > 1$ . That is we have,

$$\sup_{B(r)\subset B(R)} r^{-\frac{\alpha}{p_1}} \left( \|\Delta a\|_{L^{q_1}(B(r))} + \|\Delta b\|_{L^{q_1}(B(r))} \right) < \infty$$

Again we use the Sobolev embedding on Morrey spaces, see [1]. For

$$\frac{1}{p_2} := \frac{1}{q_1} - \frac{1}{2 - \frac{\alpha}{p_1}q_1} = \frac{1}{p_1} - \frac{\alpha}{4p_1 + 4 - 2\alpha}$$

we get

$$\sup_{B(r)\subset B(R/2)} r^{-\frac{\alpha}{p_1+1}} \left( \|\nabla a\|_{L^{p_2}(B(r))} + \|\nabla b\|_{L^{p_2}(B(r))} \right) < \infty.$$

In particular, from (5.4) and harmonicity of H we get  $v \in L^{p_2}(B(R/2))$ . By a covering argument we conclude that  $v \in W^{1,p_2}_{loc}(\mathbb{B}^2)$ .

So we define a sequence  $(p_i)_i$  by

$$\frac{1}{p_{i+1}} := \frac{1}{p_i} - \frac{\alpha}{4p_i + 4 - 2\alpha}$$

By induction we obtain from Proposition 4.1  $v \in W_{loc}^{1,p_{i+1},\mathbb{R}^N}(\mathbb{B}^2)$  for every  $i \in \mathbb{N}$  such that  $p_i < p_{\infty}$ . The important point is that  $\alpha$  is uniform and does not depend on each i.

Clearly  $p_{i+1} \ge p_i$  and  $\lim_{i\to\infty} p_i = \infty$ . That is there exists  $i_0 \in \mathbb{N}$  such that  $p_i < p_\infty$  and  $p_{i+1} > p_\infty$ . That means that  $v \in W^{1,p_\infty}_{loc}(\mathbb{B}^2)$ .

As a direct corollary from Proposition 5.4 and the Euler-Lagrange equations in Lemma 3.1 we obtain  $W_{loc}^{2,2-\varepsilon}$ -regularity for v. Observe that in view of the counterexamples in [22] this is the best regularity for v one can hope for without having further improvements on the regularity of  $\lambda$ .

**Corollary 5.5.** Let v and  $\lambda$  be as in Theorem 1.4. Then, for any  $q \in (1,2)$  we have  $v \in W_{loc}^{2,q}$ .

*Proof.* From Proposition 5.4 we have that  $v \in W^{1,p}_{loc}(\mathbb{B}^2)$  for any  $p \in (1,\infty)$ . From Lemma 3.1 we thus get that for any  $q \in (1,2)$  – recall that  $\nabla \lambda \in L^2(\mathbb{B}^2)$  –

$$\operatorname{div}(\lambda^2 \nabla v) \in L^q_{loc}(\mathbb{B}^2)$$

Now

$$\Delta v = \operatorname{div}(\lambda^{-2}\lambda^2 \nabla v) = \nabla \lambda^{-2} \lambda^2 \nabla v + L^q_{loc}(\mathbb{B}^2).$$

Since  $\inf_{\mathbb{B}^2} \lambda > 0$  we have that  $\lambda^{-2} \in H^1(\mathbb{B}^2)$  and thus

$$\Delta v \in L^q_{loc}(\mathbb{B}^2).$$

Standard elliptic estimates imply now  $v \in W^{2,q}_{loc}(\mathbb{B}^2, \mathbb{R}^N)$ .

## SUJIN KHOMRUTAI AND ARMIN SCHIKORRA

# 6. $W^{2,2}$ -regularity for $\lambda$

By now, for  $\lambda$  and v as in Theorem 1.4 we have shown in Lemma 2.1 that  $\lambda \in L^{\infty}_{loc}(\mathbb{B}^2)$ and in Corollary 5.5 that  $v \in W^{2,q}_{loc}(\mathbb{B}^2)$  for any  $q \in (1, 2)$ .

Recall that we assume that  $\lambda \geq 1$ . It will be notationally convenient to work with  $\mu := \lambda - 1$ , which is a critical point of the energy

$$\int |\nabla \mu|^2 + \int (\mu^2 - 2\mu) |\nabla v|^2$$

So in the following we are going to consider the regularity of critical points  $\mu \in L^{\infty}(\mathbb{B}^2, [0, \infty))$ 

$$F(\mu) := \int |\nabla \mu|^2 + \int (\mu^2 - 2\mu) g$$
 subject to  $\mu \ge 0$ 

where  $g \in W^{1,q}_{loc}(\mathbb{B}^2)$  for any q < 2, in particular  $g \in L^2_{loc}(\mathbb{B}^2)$ .

First we observe the variational inequality.

**Lemma 6.1.** Let  $\mu$  as above, i.e. a critical point of F. Then, for any  $\varphi \in C_c^{\infty}(\mathbb{B}^2)$  such that  $\varphi \geq 0$  we have

(6.1) 
$$\int_{\mathbb{B}^2} \nabla \mu \cdot \nabla (\varphi - \mu) + \int_{\mathbb{B}^2} \mu (\varphi - \mu) g - \int_{\mathbb{B}^2} (\varphi - \mu) g \ge 0.$$

*Proof.* This follows using the variation

$$\mu_{\varepsilon} := \mu + \varepsilon(\varphi - \mu).$$

The variational inequality (6.1) for  $\mu$  is almost of the form of variational inequalities considered e.g. in [7, (2.6)], where Frehse showed how Nirenberg's method of discretely differentiating partial differential equations can be adapted to variational inequalities. Indeed, the only additional term that does not appear in [7, (2.6)] is  $\int_{\mathbb{B}^2} \mu(\varphi - \mu)g$ . So we (slightly) adapt Frehse's argument to obtain

**Proposition 6.2.** Let  $\mu \in W^{1,2}(\mathbb{B}^2) \cap L^{\infty}_{loc}(\mathbb{B}^2)$  as above, i.e. a critical point of F. If  $g \in W^{1,q}_{loc}(\mathbb{B}^2)$  for any q < 2, then  $\mu \in W^{2,2}_{loc}(\mathbb{B}^2)$ .

We now follow closely Frehse's argument in [7], and only prove the differences.

Firstly, we introduce first and second order differential quotients,

$$\delta_{i;h}\mu(x) := \frac{\mu(x + he_i) - \mu(x)}{h},$$
  
$$\delta_{i,j;h}\mu(x) := \frac{\mu(x + he_i) + \mu(x - he_j) - 2\mu(x)}{h^2}$$

The main first observation in [7, Hilfssatz 1]

**Lemma 6.3.** Let  $\eta \in C_c^{\infty}(\mathbb{B}^2)$ ,  $\eta \ge 0$ . Then for any  $h < \text{dist}(\text{supp } \eta, \partial \mathbb{B}^2)$  we have for i = 1, 2,

$$\int_{\mathbb{B}^2} \nabla \mu \cdot \nabla (\eta^2 \delta_{i,i;h} \mu) + \int_{\mathbb{B}^2} \mu (\eta^2 \delta_{i,i;h} \mu) g - \int_{\mathbb{B}^2} (\eta^2 \delta_{i,i;h} \mu) g \ge 0.$$

*Proof.* Observe that for  $\varepsilon \ll h$  we have that

 $\mu_{\varepsilon} := \mu + \varepsilon \eta^2 \delta_{i,j;h} \mu \ge 0.$ 

In particular,  $\mu_{\varepsilon}$  is a permissible variation of  $\mu$ , and the claim follows.

The only term that we have to estimate additionally to Frehse's [7] is the following:

# Lemma 6.4.

$$\int_{\mathbb{B}^2} \mu \eta^2 \,\delta_{i,i;h} \mu \, g \precsim C \, \left( 1 + \| \nabla \delta_{i,h}(\eta \mu) \|_{L^2(\mathbb{B}^2)} \right)$$

where C depends on supp  $\eta$ ,  $\|\mu\|_{L^{\infty}}$ ,  $\|\nabla\mu\|_{L^2}$ ,  $\|\nabla\eta\|_{L^{\infty}}$ ,  $\|g\|_{L^p}$ , for p sufficiently close to  $\infty$ and  $\|\nabla g\|_{W^{1,q}}$  for a q < 2 sufficiently close to 2.

*Proof.* First, a standard application of the discrete Leibniz rule,

$$\int_{\mathbb{B}^2} \mu \eta^2 \,\delta_{i,i;h} \mu \,g = -\int_{\mathbb{B}^2} \mu \,\delta_{i,-h} \delta_{i,h}(\eta \,\mu) \,g + C.$$

Thus, with the discrete integration by parts rule we obtain for any  $q \in (1, \infty)$ ,

$$\|\nabla(\eta\mu g)\|_{L^{q}} \|\delta_{i,h}(\eta\mu)\|_{L^{\frac{q}{q-1}}}$$

For q < 2 we have

$$\|\nabla(\eta\mu g)\|_{L^q} \precsim C.$$

On the other hand, since  $\delta_{i,h}(\eta\mu)$  has compact support, by Sobolev-Poincarè-embedding (since we are in two dimensions) for any  $q \in (1, 2)$ ,

$$\|\delta_{i,h}(\eta\mu)\|_{L^{\frac{q}{q-1}}} \preceq \|\nabla\delta_{i,h}(\eta\mu)\|_{L^2}.$$

*Proof of Proposition 6.* Following word-by-word the Frehse's argument in [7], using additionally the estimate Lemma 6.4 we obtain, cf. [7, p. 149],

$$\|\nabla(\delta_{i,h}(\mu\eta))\|_{L^2}^2 \le C \ (1 + \|\nabla(\delta_{i,h}(\mu\eta))\|_{L^2}) \ .$$

From Young's inequality we obtain

$$\|\nabla(\delta_{i,h}(\mu\eta))\|_{L^2}^2 \le C + 4C^2 + \frac{1}{2}\|\nabla(\delta_{i,h}(\mu\eta))\|_{L^2}^2$$

and thus we obtain a bound on  $\|\nabla(\delta_{i,h}(\mu\eta))\|_{L^2}^2$  independent of h. Letting  $h \to 0$  we get that

$$\nabla \partial_i(\mu \eta) \in L^2,$$

which readily leads to  $u \in W^{2,2}$  in the set where  $\eta \equiv 1$ . Taking  $\eta \in C_c^{\infty}(\mathbb{B}^2)$  and  $\eta \equiv 1$  on  $K \subset \mathbb{B}^2$ , K compact, we get that  $\mu \in W^{2,2}_{loc}(K)$ . This holds for any compact set  $K \subset \mathbb{B}^2$ , so the claim is proven.

**Corollary 6.5.** Let v and  $\lambda$  be as in Theorem 1.4. Then  $\lambda \in W^{2,2}_{loc}(\mathbb{B}^2)$ ,  $v \in W^{2,p}_{loc}(\mathbb{B}^2, \mathbb{R}^N)$  for any  $p < \infty$ . In particular  $\lambda \in C^{0,\alpha}$  and  $v \in C^{1,\alpha}$  for any  $\alpha < 1$ .

*Proof.* By Corollary 5.5 we have that  $|\nabla v|^2 \in W_{loc}^{1,q}$  for any q < 2. Thus Proposition 6 is applicable to  $\mu = \lambda - 1$ , and we get that  $\lambda = \mu + 1 \in W_{loc}^{2,2}(\mathbb{B}^2)$ .

To obtain  $W_{loc}^{2,p}$ -regularity for v, we consider again the equations for v, (3.1), and note that

$$\operatorname{div}(\lambda^2 \nabla v) \in L^p_{loc}(\mathbb{B}^2)$$

Moreover, since  $\lambda \geq 1$ , we compute

$$\Delta v = \operatorname{div}(\lambda^{-2}\lambda^2 \nabla v) = \lambda^{-} \nabla \lambda \nabla v + \lambda^{-2} \operatorname{div}(\lambda^2 \nabla v).$$

Since  $\nabla \lambda \in W_{loc}^{1,2}$  and  $\nabla v \in W_{loc}^{1,q}$  for any q < 2, we obtain that  $\Delta v \in L_{loc}^{p}$  and consequently standard Calderon-Zygmund theory implies that  $\nabla^{2}v \in L_{loc}^{p}$ .

# 7. On $C^{1,\alpha}$ -regularity for $\lambda$

At this stage we have that  $v \in W^{2,p}_{loc}(\mathbb{B}^2, \mathbb{R}^N)$  for all  $p \in (1, \infty)$  and  $\lambda \in W^{2,2}_{loc}(\mathbb{B}^2)$ . Observe that this implies in particular that  $\lambda$  is continuous. Since the obstacle condition  $\lambda \geq 1$  is pointwise, the theory of viscosity solutions (see e.g. [3, 12]) is more suitable now.

**Proposition 7.1.** There exists a constant  $\Lambda > 0$  such that  $\lambda$  solve in viscosity sense the inequalities

$$0 \le \Delta \lambda \le \Lambda \quad in \ \mathbb{B}^2.$$

This Proposition is a consequence of Lemma 7.3 and Lemma 7.4 below. The first observation is that  $\lambda$  is smooth in the open set  $\{\lambda > 1\}$ .

**Lemma 7.2.** We have  $\lambda, v \in C^{\infty}(\{\lambda > 1\})$  and we have

(7.1) 
$$\Delta \lambda = \lambda |\nabla v|^2 \quad pointwise \ in \ \{\lambda > 1\}$$

In view of Corollary 6.5 there exists in particular  $\Lambda > 0$  such that

$$\Delta \lambda \leq \Lambda \quad in \ \{\lambda > 1\}.$$

Proof. We revert our attention to  $u := \lambda v$  and show that  $u \in C^{\infty}(\{\lambda > 1\})$ . Let  $x_0 \in \{\lambda > 1\}$ . 1}. Then, since  $\lambda$  is continuous, there exists a ball  $B := B(x_0)$  such that  $\overline{B} \subset \{\lambda > 1\}$ . But this implies that for any  $\varphi \in C_c^{\infty}(B)$  for all suitably small

$$u_{\varepsilon} := u + \varepsilon \varphi$$

is a permissible variation of the Dirichlet energy, setting  $\lambda_{\varepsilon} := |u_{\varepsilon}| > 0$  and  $v_{\varepsilon} := \frac{u_{\varepsilon}}{|u_{\varepsilon}|^2}$ . That is,

$$\frac{d}{d\varepsilon}\Big|_{\varepsilon=0}\int |\nabla u_{\varepsilon}|^2 = 0.$$

But this implies

$$\Delta u = 0 \quad \text{in } \{\lambda > 1\},$$

so in particular  $\lambda = |u| \in C^{\infty}(\{\lambda > 1\})$  and  $v = \frac{u}{|u|} \in C^{\infty}(\{\lambda > 1\})$ .

The equation (7.1) follows now from the variation  $\lambda_{\varepsilon} := \lambda + \varepsilon \psi$  for arbitrary  $\psi \in C_c^{\infty}(\{\lambda > 1\})$ .

Lemma 7.3. We have in viscosity sense

$$\Delta \lambda \ge 0 \quad in \ \mathbb{B}^2.$$

*Proof.* Let  $x_0 \in \mathbb{B}^2$ . If  $\lambda(x_0) > 1$ , then the claim follows immediately from Lemma 7.2, since smooth solutions are viscosity solutions.

So assume that  $\lambda(x_0) = 1$ . For any test-function  $\varphi \ge \lambda$  such that  $\varphi(x_0) = 1$  we have in particular

 $\varphi \ge 1$ , and  $\varphi(x_0) = 1$ .

That is,  $\varphi$  attains its minimum at  $x_0$  and thus  $\Delta \varphi \ge 0$ .

**Lemma 7.4.** For  $\lambda$  as above we have in viscosity sense.

$$\Delta \lambda \leq \Lambda \quad in \ \mathbb{B}^2.$$

*Proof.* By the variation  $\lambda_{\varepsilon} := \lambda + \varepsilon \varphi$  for  $\varphi \in C_c^{\infty}(\mathbb{B}^2)$  and  $\varphi \ge 0$  we get the variational inequality

$$\int \nabla \lambda \cdot \nabla \varphi + \lambda \varphi |\nabla v|^2 \ge 0$$

Let  $\eta \in C_c^{\infty}(B(0,1))$  be the usual bump function,  $\eta \equiv 1$  in  $B(0,\frac{1}{2})$ ,  $\eta(-x) = \eta(x)$ ,  $\eta \geq 0$ and  $\int \eta = 1$ . We set  $\eta_{\varepsilon} := \varepsilon^{-2} \eta(\cdot/\varepsilon)$ . We denote  $\lambda_{\varepsilon} := \eta_{\varepsilon} * \lambda$  and have for any fixed testfunction  $\varphi \geq 0$  (if  $\varepsilon$  is small enough then  $\varphi * \eta_{\varepsilon} \geq 0$  is permissible as a test function)

$$\int \nabla \lambda_{\varepsilon} \cdot \nabla \varphi = \int \nabla \lambda \cdot \nabla (\varphi * \eta_{\varepsilon})$$
$$\geq -\int \lambda (\varphi * \eta_{\varepsilon}) |\nabla v|^{2}$$
$$= -\int \varphi \left(\lambda |\nabla v|^{2}\right) * \eta_{\varepsilon}$$

Since, in view of Corollary 6.5 we have

$$\|\left(\lambda|\nabla v|^{2}\right)*\eta_{\varepsilon}\|_{L^{\infty}} \leq \|\lambda|\nabla v|^{2}\|_{L^{\infty}} =: \Lambda < \infty$$

we find that

(7.2) 
$$\Delta \lambda_{\varepsilon} \leq \Lambda \quad \text{in } \mathbb{B}_{1-2\varepsilon}$$

This inequality holds in pointwise and viscosity sense, since  $\lambda_{\varepsilon}$  is smooth.

On the other hand, since  $\lambda$  is Hölder continuous, we have that  $\lambda_{\varepsilon}$  converges locally uniformly to  $\lambda$  as  $\varepsilon \to 0$ . This implies, e.g. as in [19, Lemma 2.4], that also  $\lambda$  satisfies (7.2) in viscosity sense.

As a consequence of the regularity theory of viscosity solutions to elliptic partial differential inequalities, see e.g. [19], we obtain

**Corollary 7.5.** Let be  $\lambda$  as above, then  $\lambda \in C^{1,\alpha}$  for any  $\alpha < 1$ .

# 8. Adaptations for the proof of Theorem 1.3

For matrices  $A, B \in \mathbb{R}^{N \times N}$  we denote the Hilbert-Schmidt scalar product by

$$A:B:=\sum_{i,j=1}^{\infty}A_{ij}B_{ij}.$$

Now as in the sphere case, where we have  $u \cdot \nabla u = 0$  if |u| = 1 almost everywhere, if  $P \in SO(N)$  almost everywhere then

$$\nabla P : P = P^T \nabla P : I_{N \times N} = 0,$$

since  $P^T \nabla P$  is antisymmetric and the identity matrix  $I_{N \times N}$  is symmetric.

In particular we have for  $\lambda \in H^1(\mathbb{B}^2)$  and  $P \in H^1(\mathbb{B}^2, SO(N))$ ,

$$|\nabla(\lambda P)|^2 = |\nabla\lambda|^2 + \lambda^2 |\nabla P|^2.$$

We conclude that we have to consider critical points of the energy

$$E(\lambda, P) = |\nabla \lambda|^2 + \lambda^2 |\nabla P|^2.$$

So we see that we get the analogue of Lemma 3.1. Now regularity estimates are almost verbatim of what we have here.

**Lemma 8.1** (Euler-Lagrange equations). Let  $\lambda$  and P be as in Theorem 1.3. Then,

$$\operatorname{div}(\lambda^2 \nabla P) = \lambda^2 \Omega \nabla P.$$

with

$$\Omega = -P^T \nabla P.$$

Equivalently we also have the conservation law

$$\operatorname{div}(\lambda^2 P^T \nabla P) = 0.$$

*Proof.* A permissible variation for P is  $P_{\varepsilon} := Pe^{\varepsilon \alpha \varphi}$  where  $\alpha \in so(N)$  is antisymmetric and  $\varphi \in C_c^{\infty}(\mathbb{B}^2)$ . This leads to

$$\int \lambda^2 \nabla P : \nabla (P \alpha \varphi) = 0$$

Observe that for antisymmetric  $\alpha$  we readily have

 $\nabla P: \nabla P\alpha = 0$ 

Thus, the Euler-Lagrange equations for variations in P are

$$\operatorname{div}(\lambda^2 P^T \nabla P) : \alpha = 0.$$

This holds for any antisymmetric matrix  $\alpha \in so(N)$ . Using that  $P^T \nabla P$  is also antisymmetric, we thus get

$$\operatorname{div}(\lambda^2 P^T \nabla P) = 0$$

We can equivalently rewrite this as

$$\operatorname{div}(\lambda^2 \nabla P) = \operatorname{div}(\lambda^2 P P^T \nabla P) = \lambda^2 \nabla P P^T \nabla P.$$
  
$$\nabla = \nabla (P P^T) = P^T \nabla P = -P^T \nabla P \text{ we get the claim}$$

Using that  $\nabla PP^T = \nabla (PP^T) - P^T \nabla P = -P^T \nabla P$  we get the claim.

Acknowledgment. A.S. was supported by the German Research Foundation (DFG) through grant no. SCHI-1257-3-1, by the Daimler and Benz foundation through grant no. 32-11/16, as well as the Simons foundation through grant no 579261. Part of this work was carried out while A.S. was visiting Chulalongkorn University whose hospitality is gratefully acknowledged.

### References

- [1] D. R. Adams. A note on Riesz potentials. Duke Math. J., 42(4):765–778, 1975. 12, 13
- [2] H. Brezis and J.-M. Coron. Multiple solutions of H-systems and Rellich's conjecture. Comm. Pure Appl. Math., 37(2): 149–187, 1984. 8
- [3] L. Caffarelli and X. Cabré. Fully nonlinear elliptic equations, volume 43 of AMS Colloquium Publications. AMS, Providence, RI, 1995. 16
- [4] R. Coifman, P.-L. Lions, Y. Meyer, and S. Semmes. Compensated compactness and Hardy spaces. J. Math. Pures Appl., IX. Sér., 72(3): 247–286, 1993. 8
- [5] F. Duzaar and M. Fuchs. Variational problems with nonconvex obstacles and an integral-constraint for vector-valued functions. *Math. Z.*, 191(4):585–591, 1986. 2, 3
- [6] L. C. Evans. Partial differential equations, volume 19 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition, 2010. 5
- [7] J. Frehse. Zum Differenzierbarkeitsproblem bei Variationsungleichungen höherer Ordnung. Abh. Math. Sem. Univ. Hamburg, 36:140–149, 1971. Collection of articles dedicated to Lothar Collatz on his sixtieth birthday. 14, 15
- [8] M. Giaquinta. Multiple integrals in the calculus of variations and nonlinear elliptic systems, volume 105 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1983.
- [9] F. Hélein. Régularité des applications faiblement harmoniques entre une surface et une sphère. C.R. Acad. Sci. Paris 311, Série I, pages 519–524, 1990.
- [10] S. Hildebrandt. Interior  $C^{1+\alpha}$ -regularity of solutions of two-dimensional variational problems with obstacles. *Math. Z.*, 131:233–240, 1973. 2

- [11] S. Hildebrandt and J. C. C. Nitsche. A uniqueness theorem for surfaces of least area with partially free boundaries on obstacles. Arch. Rational Mech. Anal., 79(3):189–218, 1982. 2
- [12] S. Koike. A beginner's guide to the theory of viscosity solutions. www.math.tohoku.ac.jp/~koike/evis2012version.pdf. 2012. 16
- [13] S. Müller. Higher integrability of determinants and weak convergence in L<sup>1</sup>. J. Reine Angew. Math., 412: 20–34, 1990. 8
- [14] Y. G. Reshetnyak. Stability theorems for mappings with bounded excersions. Siberian Mathematical Journal, 9(3):499–512, 1968.
- [15] T. Rivière. Everywhere discontinuous harmonic maps into spheres. Acta Math., 175(2):197–226, 1995.
  3
- [16] T. Rivière. Conservation laws for conformally invariant variational problems. Invent. Math., 168(1):1– 22, 2007. 11
- [17] A. Schikorra. A remark on gauge transformations and the moving frame method. Ann. Inst. H. Poincaré Anal. Non Linéaire, 27(2):503–515, 2010. 11
- [18] A. Schikorra. ε-regularity for systems involving non-local, antisymmetric operators. Calc. Var. Partial Differential Equations, 54(4):3531–3570, 2015. 12
- [19] A. Schikorra. A remark on  $C^{1,\alpha}$ -regularity for differential inequalities in viscosity sense. *preprint*, 2018. 18
- [20] R. Schoen and K. Uhlenbeck. Regularity of minimizing harmonic maps into the sphere. Invent. Math., 78(1):89–100, 1984. 3
- [21] B. Sharp. Higher integrability for solutions to a system of critical elliptic PDE. Methods Appl. Anal., 21(2):221–240, 2014. 12
- [22] B. Sharp and P. Topping. Decay estimates for Rivière's equation, with applications to regularity and compactness. Trans. Amer. Math. Soc., 365(5):2317–2339, 2013. 4, 12, 13
- [23] J. Shatah. Weak solutions and development of singularities of the SU(2) σ-model. Comm. Pure Appl. Math., 41(4):459–469, 1988. 6, 7
- [24] L. Tartar. Remarks on Oscillations and Stokes' Equation. Lecture Notes in Physics, 230, macroscopic Modelling of Turbulent Flows, Proceedings, Sophia-Antipolis, France, pages 24–31, 1984. 8
- [25] P. Topping. The optimal constant in Wente's L<sup>∞</sup> estimate. Comment. Math. Helv., 72(2):316–328, 1997. 8
- [26] H. C. Wente. An existence theorem for surfaces of constant mean curvature. J. Math. Anal. Appl., 26: 318–344, 1969. 8

(Sujin Khomrutai) Department of Mathematics and Computer Science, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

*E-mail address*: sujin.k@chula.ac.th

(Armin Schikorra) Department of Mathematics, University of Pittsburgh, 301 Thackeray Hall, Pittsburgh, PA 15260, USA

*E-mail address*: armin@pitt.edu