

# WEIGHTED INFINITESIMAL UNITARY BIALGEBRAS ON MATRIX ALGEBRAS AND WEIGHTED ASSOCIATIVE YANG-BAXTER EQUATIONS

YI ZHANG, XING GAO\*, AND JIA-WEN ZHENG

**ABSTRACT.** We equip a matrix algebra with a weighted infinitesimal unitary bialgebraic structure, via a construction of a suitable coproduct. Furthermore, an infinitesimal unitary Hopf algebra, under the view of Aguiar, is constructed on a matrix algebra. By exploring the relationship between weighted infinitesimal bialgebras and pre-Lie algebras, we construct a pre-Lie algebraic structure and then a new Lie algebraic structure on a matrix algebra. We also introduce the weighted associative Yang-Baxter equations (AYBEs) and obtain the relationship between solutions of weighted AYBEs and weighted infinitesimal unitary bialgebras. We give a bijection between the solutions of the associative Yang-Baxter equation of weight  $\lambda$  and Rota-Baxter operators of weight  $-\lambda$  on matrix algebras. As a consequence, weighted quasitriangular infinitesimal unitary bialgebras are constructed, which generalize the results studied by Aguiar. Finally, We show that any weighted quasitriangular infinitesimal unitary bialgebra can be made into a dendriform algebra.

## CONTENTS

|   |    |
|---|----|
| 1. Introduction   | 2  |
| 2. Weighted infinitesimal unitary bialgebras and examples                             | 5  |
| 2.1. Weighted infinitesimal unitary bialgebras  | 5  |
| 2.2. An infinitesimal unitary bialgebra on a matrix algebra                           | 6  |
| 2.3. An infinitesimal unitary Hopf algebra on a matrix algebra                        | 8  |
| 3. Pre-Lie algebras on matrix algebras  | 10 |
| 3.1. Pre-Lie algebras and weighted infinitesimal unitary bialgebras                   | 11 |
| 3.2. Two pre-Lie and a new Lie algebraic structures on a matrix algebra               | 13 |
| 4. Weighted AYBEs and weighted $\epsilon$ -unitary bialgebras                         | 14 |
| 4.1. Weighted AYBEs   | 14 |
| 4.2. A solution of a homogeneous AYBE for matrix algebras                             | 16 |
| 4.3. Weighted AYBEs and Rota-Baxter operators   | 17 |
| 4.4. A bijection between the solutions of weighted AYBEs and Rota-Baxter operators    | 18 |
| 5. Weighted quasitriangular $\epsilon$ -unitary bialgebras and dendriform algebras    | 20 |
| 5.1. Weighted quasitriangular $\epsilon$ -unitary bialgebras                          | 21 |
| 5.2. Dendriform algebras from weighted quasitriangular $\epsilon$ -unitary bialgebras | 22 |
| References  | 22 |

---

\* Corresponding author.

*Date:* February 28, 2022.

*2010 Mathematics Subject Classification.* 15A30, 16W99, 16T10, 17B60, 17D25 .

*Key words and phrases.* matrix algebra; infinitesimal bialgebra; pre-Lie algebra; associative Yang-Baxter equation.

## 1. INTRODUCTION

The interaction between studies in pure mathematics and mathematical physics has long been a rich source of inspirations that benefited both fields. This paper arose from an attempt to connect these two fields by establishing connections among weighted infinitesimal unitary bialgebras, weighted associative Yang-Baxter equations, pre-Lie algebras, Rota-Baxter algebras and dendri-form algebras.

Infinitesimal bialgebras, introduced by Joni and Rota [37], are in order to give an algebraic framework for the calculus of Newton divided differences. Namely, an infinitesimal bialgebra is a module  $A$  which is simultaneously an algebra (possibly without a unit) and a coalgebra (possibly without a counit) such that the coproduct  $\Delta$  is a derivation of  $A$  in the sense:

$$\Delta(ab) = a \cdot \Delta(b) + \Delta(a) \cdot b \text{ for } a, b \in A.$$

If an infinitesimal bialgebra has an antipode  $S$ , then it will be called an infinitesimal Hopf algebra [1]. The basic theory of infinitesimal bialgebras and infinitesimal Hopf algebras was developed by Aguiar [1, 3, 4, 5], which has proven useful not only in combinatorics [4], but in other areas of mathematics as well, such as associative Yang-Baxter equations, Drinfeld's doubles and pre-Lie algebras [1]. Recently, Wang [50] generalized Aguiar's result by developing the Drinfeld's double for braided infinitesimal Hopf algebras in Yetter-Drinfeld categories.

We emphasize that another version of infinitesimal bialgebras and infinitesimal Hopf algebras was defined by Loday and Ronco [43] and brought new life on rooted trees by Foissy [27, 28] in the sense that

$$\Delta(ab) = a \cdot \Delta(b) + \Delta(a) \cdot b - a \otimes b \text{ for } a, b \in A.$$

In [31], the authors combined the two versions of infinitesimal bialgebras by defining the coproduct of  $A$  to be the following compatibility:

$$\Delta(ab) = a \cdot \Delta(b) + \Delta(a) \cdot b + \lambda(a \otimes b) \text{ for } a, b \in A,$$

where  $\lambda \in \mathbf{k}$  is a fixed constant. This leads to the born of weighted infinitesimal (unitary) bialgebras, that is, the infinitesimal (unitary) bialgebras of weight  $\lambda$ . See Definition 2.1 below.

In the present paper, we equip a matrix algebra with a weighted infinitesimal unitary bialgebraic structure by a construction of a suitable coproduct. Moreover, we also equip an infinitesimal unitary bialgebra of weight zero on matrix algebras with an antipode such that it is further an infinitesimal unitary Hopf algebra, under the view of Aguiar [1].

Pre-Lie algebras, also called Vinberg algebras, first appeared in the work of Vinberg [49] under the name left-symmetric algebras on convex homogeneous cones and also appeared independently at the same time in the study of affine structures on manifolds [32]. Its study has a broad applications in mathematics and mathematical physics, such as classical and quantum Yang-Baxter equations [8, 9, 14, 25, 36], pre-Poisson algebras [2] and Poisson brackets [15], quantum field theory [16, 17, 35, 39, 40, 47] and operads [18, 46], Lie group and Lie algebras [38, 45, 49],  $\mathcal{O}$ -operators [7, 8, 10] and Rota-Baxter algebras [6, 11, 22, 24, 29, 30]. In [7], Bai pointed out that "Due to the nonassociativity of pre-Lie algebra, there is not a suitable (and computable) representation theory and not a complete (and good) structure theory of pre-Lie algebras". It is nature to consider how to construct them from some algebraic structures (especially associative algebras) which we have known. Our results may give a new and elementary method to construct pre-Lie algebraic structures on some associative algebras, especially on matrix algebras.

It should be pointed out that Bai [7] gave two approaches to construct pre-Lie algebras from associative algebras, see [7, Corollary 3.15] and [7, Proposition 3.16] for more details. Our

consideration of pre-Lie algebras on matrix algebras has motivations beyond a simple pursuit of weighted infinitesimal unitary bialgebras on matrix algebras. In the algebraic framework of Aguiar [5] for infinitesimal bialgebra, a pre-Lie algebraic structure is constructed from an arbitrary infinitesimal bialgebra. Motivated by Aguiar's construction, we previously derive a pre-Lie algebra from an arbitrary weighted infinitesimal bialgebra. As applications, two pre-Lie algebras on matrix algebras are built in the present paper.

Due to the construction of an infinitesimal unitary bialgebra of weight  $\lambda$  arising from a matrix algebra in this paper, there is a close relationship among the matrix algebras, pre-Lie algebras, Lie algebras and weighted infinitesimal unitary bialgebras. This situation can be summarized in the sense of following commutative diagram of categories

$$\begin{array}{ccc}
 \text{Matrix algebras} & \rightarrow & \text{Weighted infinitesimal (unitary) bialgebras} \\
 \downarrow & & \downarrow \\
 \text{Lie algebras} & \longleftarrow & \text{Pre-Lie algebras}
 \end{array}$$

Let  $\mathfrak{g}$  be a Lie algebra and  $r \in \mathfrak{g} \otimes \mathfrak{g}$ . A well-known result about classical Yang-Baxter equation (CYBE) studied by Drinfeld [20] is that the principal derivation  $\delta_r : \mathfrak{g} \rightarrow \mathfrak{g} \otimes \mathfrak{g}$  is coassociative if and only if the element

$$[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] \in \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g}$$

is  $\mathfrak{g}$ -invariant. The solutions of classical Yang-Baxter equation give rise to Lie bialgebras and quantum groups [20]. Parallel to classical Yang-Baxter equation, Aguiar [1] introduced associative Yang-Baxter equation (AYBE)

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = 0.$$

Aguiar [3] shown that any solution  $r$  of AYBE in an algebra  $A$  is a solution of CYBE in  $A^{lie}$  provided that  $r + \tau(r)$  is  $A$ -invariant. Here  $A^{lie}$  denotes the Lie algebra obtained by endowing  $A$  with the commutator bracket and  $\tau$  is the switch map. In particular, any skew-symmetric solution of AYBE is a skew-symmetric solution of CYBE in  $A^{lie}$ .

Let  $A$  be a unitary algebra. For each solution  $r \in A \otimes A$ , the principle derivation

$$\Delta_r : A \rightarrow A \otimes A, \quad a \mapsto a \cdot r - r \cdot a$$

endows  $A$  with an infinitesimal unitary bialgebra of weight zero. In this paper, we generalize Aguiar's result by the following result.

**Theorem 1.1.** (=Theorem 4.5) Let  $A$  be a unitary algebra and  $r = \sum_i u_i \otimes v_i \in A \otimes A$ . Then the weighted principle derivation  $\Delta_r(a) = a \cdot r - r \cdot a - \lambda(a \otimes 1)$  is coassociative if and only if the element  $r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} - \lambda r_{13} \in A \otimes A \otimes A$  is  $A$ -invariant.

Then we call the equation

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = \lambda r_{13}$$

an associative Yang-Baxter equation of weight  $\lambda$ . Let us emphasize that the weighted AYBEs considered here are very general, which include the AYBEs [1], the modified AYBEs [21] and the non-homogeneous AYBEs in [10]. See Remark 4.2 below. Moreover, a surprising phenomenon shows that a solution of a weighted AYBE induces a Rota-Baxter operator of weight  $-\lambda$  (Theorem 4.10). As a consequence, we give a bijection between the set of the solutions of weighted AYBE and the set of Rota-Baxter operators on matrix algebras.

**Theorem 1.2.** (=Theorem 4.12) Let  $r$  be a solution of an AYBE of weight  $\lambda$  in  $M_n(\mathbf{k})$ . Then the map  $r \rightarrow P_r$  is a bijection between the set of the solutions of AYBE of weight  $\lambda$  in  $M_n(\mathbf{k})$  and the set of Rota-Baxter operators of weight  $-\lambda$  on  $M_n(\mathbf{k})$ .

The solutions of an AYBE of weight  $\lambda$  give rise to an infinitesimal unitary bialgebra of weight  $\lambda$  and we call this infinitesimal unitary bialgebra the weighted quasitriangular infinitesimal unitary bialgebra. As an application, we can derive a dendriform algebra from a weighted quasitriangular infinitesimal unitary bialgebra and obtain a commutative diagram:

$$\begin{array}{ccccc}
 \text{Weighted quasitriangular } \epsilon\text{-unitary bialgebras} & \rightarrow & \text{Weighted } \epsilon\text{-unitary bialgebras} & & \\
 \downarrow & & & \searrow & \\
 \text{Rota-Baxter algebras} & \longrightarrow & \text{Dendriform algebras} & \longrightarrow & \text{Pre-Lie algebras}
 \end{array}$$

**Structure of the Paper.** In Section 2, we start by recalling the concept of an infinitesimal (unitary) bialgebra of weight  $\lambda$  (Definition 2.1). Then we proceed to construct two different coproducts  $\Delta_L$  and  $\Delta_\lambda$  on matrix algebra  $M_n(\mathbf{k})$  to equip them with two infinitesimal unitary bialgebraic structures (Theorems 2.8 and 2.12). At the end of this section, we equip an infinitesimal unitary Hopf algebra on matrix algebras (Theorem 2.21), under the view of Aguiar [1].

In Section 3, by investigating the relationship between weighted infinitesimal bialgebras and pre-Lie algebras (Theorem 3.5), we equip  $M_n(\mathbf{k})$  with two pre-Lie bialgebraic structures and two Lie algebraic structures (Theorems 3.6 and 3.8), beyond a construction of coproducts. It should be pointed out that the new Lie bracket on  $M_n(\mathbf{k})$  induced by  $\Delta_L$  is different from the classical one (Example 3.7), and the Lie bracket derived from  $\Delta_\lambda$  is precisely the classical Lie bracket on matrix algebra when  $\lambda = 1$  (Remark 3.9).

In Section 4, we introduce the concept of a weighted associative Yang-Baxter equation (Definition 4.1), which generalizes the concept of associative Yang-Baxter equation studied in [1, Section 5]. We propose the concept of a weighted principle derivation  $\Delta_r$  and characterize  $\Delta_r$  to be coassociative. We show that if  $r$  is a solution of a weighted AYBE in  $A$ , then the quadruple  $(A, m, 1, \Delta_r)$  is an  $\epsilon$ -unitary bialgebra of weight  $\lambda$  (Theorem 4.6). As an application, a solution of a homogeneous associative Yang-Baxter equation for matrix algebras is also given (Theorem 4.7). We also derive a Rota-Baxter operator of weight  $-\lambda$  from a solution of an AYBE of weight  $\lambda$  (Theorem 4.10). We end this section by giving a one-to-one correspondence between the solutions of the associative Yang-Baxter equation of weight  $\lambda$  and Rota-Baxter operators of weight  $-\lambda$  on matrix algebras (Theorem 4.12).

In Section 5, we first propose the concept of weighted quasitriangular  $\epsilon$ -unitary bialgebras (Definition 5.1). Similar to the classical quasitriangular bialgebras, we then give some properties of weighted quasitriangular  $\epsilon$ -unitary bialgebras (Proposition 5.3). We finally derive a dendriform algebra from a weighted quasitriangular  $\epsilon$ -unitary bialgebra (Theorem 5.6).

**Notation.** Throughout this paper, let  $\mathbf{k}$  be a unitary commutative ring unless the contrary is specified, which will be the base ring of all modules, algebras, coalgebras, bialgebras, tensor products, as well as linear maps. By an algebra we mean an associative  $\mathbf{k}$ -algebra (possibly without unit) and by a coalgebra we mean a coassociative  $\mathbf{k}$ -coalgebra (possibly without counit). For an algebra  $A$ , we view  $A \otimes A$  as an  $A$ -bimodule via

$$(1) \quad a \cdot (b \otimes c) := ab \otimes c \text{ and } (b \otimes c) \cdot a := b \otimes ca,$$

where  $a, b, c \in A$ .

## 2. WEIGHTED INFINITESIMAL UNITARY BIALGEBRAS AND EXAMPLES

In this section, we first recall the concept of a weighted infinitesimal (unitary) bialgebra [31], which generalise simultaneously the one introduced by Joni and Rota [37] and the one initiated by Loday and Ronco [43]. Then we proceed to equip a matrix algebra with a weighted infinitesimal unitary bialgebraic structure, in terms of a construction of a suitable coproduct.

**2.1. Weighted infinitesimal unitary bialgebras.** The following is the concept of a weighted infinitesimal (unitary) bialgebra proposed in [31].

**Definition 2.1.** [31] Let  $\lambda$  be a given element of  $\mathbf{k}$ . An **infinitesimal bialgebra** (abbreviated  $\epsilon$ -**bialgebra**) of weight  $\lambda$  is a triple  $(A, m, \Delta)$  consisting of an algebra  $(A, m)$  (possibly without unit) and a coalgebra  $(A, \Delta)$  (possibly without counit) that satisfies

$$(2) \quad \Delta(ab) = a \cdot \Delta(b) + \Delta(a) \cdot b + \lambda(a \otimes b) \text{ for } a, b \in A.$$

If further  $(A, m, 1)$  is a unitary algebra, then the quadruple  $(A, m, 1, \Delta)$  is called an **infinitesimal unitary bialgebra** (abbreviated  $\epsilon$ -**unitary bialgebra**) of weight  $\lambda$ .

The concept of an  $\epsilon$ -bialgebra morphism is given as usual.

**Definition 2.2.** [31] Let  $A$  and  $B$  be two  $\epsilon$ -bialgebras of weight  $\lambda$ . A map  $\phi : A \rightarrow B$  is called an **infinitesimal bialgebra morphism** (abbreviated  $\epsilon$ -bialgebra morphism) if  $\phi$  is an algebra morphism and a coalgebra morphism. The concept of an **infinitesimal unitary bialgebra morphism** can be defined in the same way.

**Remark 2.3.** (a) Let  $(A, m, 1, \Delta)$  be an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ . Then  $\Delta(1) = -\lambda(1 \otimes 1)$  by

$$\Delta(1) = \Delta(1 \cdot 1) = 1 \cdot \Delta(1) + \Delta(1) \cdot 1 + \lambda(1 \otimes 1) = 2\Delta(1) + \lambda(1 \otimes 1).$$

(b) Aguiar [1] pointed out that there is no non-zero  $\epsilon$ -bialgebra of weight zero which is both unitary and counitary. Indeed, it follows from the counicity that

$$1 \otimes 1_{\mathbf{k}} = (\text{id} \otimes \epsilon)\Delta(1) = 0,$$

and so  $1 = 0$ .

**Example 2.4.** Here are some examples of  $\epsilon$ -(unitary) bialgebras.

- (a) Any unitary algebra  $(A, m, 1)$  is an  $\epsilon$ -unitary bialgebra of weight zero by taking  $\Delta = 0$ .
- (b) [1, Example 2.3.5] The **polynomial algebra**  $\mathbf{k}\langle x_1, x_2, x_3, \dots \rangle$  is an  $\epsilon$ -unitary bialgebra of weight zero with the coproduct  $\Delta$  given by Eq. (2) and

$$\Delta(x_n) = \sum_{i=0}^{n-1} x_i \otimes x_{n-1-i} = 1 \otimes x_{n-1} + x_1 \otimes x_{n-2} + \cdots + x_{n-1} \otimes 1,$$

where we set  $x_0 = 1$ .

- (c) [1, Example 2.3.2] Let  $Q$  be a quiver. The **path algebra** of  $Q$  is the associative algebra  $\mathbf{k}Q = \bigoplus_{n=0}^{\infty} \mathbf{k}Q_n$  whose underlying  $\mathbf{k}$ -module has its basis the set of all paths  $a_1 a_2 \cdots a_n$  of length  $n \geq 0$  in  $Q$ . The multiplication  $*$  of two paths  $a_1 a_2 \cdots a_n$  and  $b_1 b_2 \cdots b_m$  is defined by

$$(a_1 a_2 \cdots a_n) * (b_1 b_2 \cdots b_m) := \delta_{t(a_n), s(b_1)} a_1 a_2 \cdots a_n b_1 b_2 \cdots b_m,$$

where  $\delta_{t(a_n), s(b_1)}$  is the Kronecker delta. The path algebra  $(\mathbf{k}Q, *, \Delta)$  is an  $\epsilon$ -bialgebra of weight zero with the coproduct defined by

$$\begin{aligned}\Delta(e) &:= 0 \text{ for } e \in Q_0 \\ \Delta(a) &:= s(a) \otimes t(a) \text{ for } a \in Q_1, \text{ and}\end{aligned}$$

$$\Delta(a_1 a_2 \cdots a_n) := s(a_1) \otimes a_2 \cdots a_n + a_1 \cdots a_{n-1} \otimes t(a_n) + \sum_{i=1}^{n-2} a_1 \cdots a_i \otimes a_{i+2} \cdots a_n \text{ for } n \geq 2.$$

(d) [43, Section 2.3] Let  $V$  denote a vector space. Recall that the **tensor algebra**  $T(V)$  over  $V$  is the tensor module,

$$T(V) = \mathbf{k} \oplus V \oplus V^{\otimes 2} \oplus \cdots \oplus V^{\otimes n} \oplus \cdots,$$

equipped with the associative multiplication  $m_T$  called concatenation defined by

$$v_1 \cdots v_i \otimes v_{i+1} \cdots v_n \mapsto v_1 \cdots v_i v_{i+1} \cdots v_n \quad \text{for } 0 \leq i \leq n,$$

with the convention that  $v_1 v_0 = 1$  and  $v_{n+1} v_n = 1$ . It is a well-known free associative algebra. The tensor algebra  $T(V)$  is an  $\epsilon$ -unitary bialgebra of weight  $-1$  with the coproduct defined by

$$\Delta(v_1 \cdots v_n) := \sum_{i=0}^n v_1 \cdots v_i \otimes v_{i+1} \cdots v_n.$$

(e) [31, Proposition 2.6] The **polynomial algebra**  $\mathbf{k}[x]$  is an  $\epsilon$ -unitary bialgebra of weight  $\lambda$  with the coproduct defined by

$$\Delta(1) := -\lambda(1 \otimes 1) \text{ and } \Delta(x^n) := \sum_{i=0}^{n-1} x^i \otimes x^{n-1-i} + \lambda \sum_{i=1}^{n-1} x^i \otimes x^{n-i} \quad \text{for } n \geq 1.$$

(f) [26, Section 1.4] Let  $(A, m, 1, \Delta, \epsilon, c)$  be a braided bialgebra with  $A = \mathbf{k} \oplus \ker \epsilon$  and the braiding  $c : A \otimes A \rightarrow A \otimes A$  given by

$$c : \begin{cases} 1 \otimes 1 \mapsto 1 \otimes 1, \\ a \otimes 1 \mapsto 1 \otimes a, \\ 1 \otimes b \mapsto b \otimes 1, \\ a \otimes b \mapsto 0, \end{cases}$$

where  $a, b \in \ker \epsilon$ . Then  $(A, m, 1, \Delta, \epsilon)$  is an  $\epsilon$ -unitary bialgebra of weight  $-1$ .

**2.2. An infinitesimal unitary bialgebra on a matrix algebra.** In this subsection, we construct an  $\epsilon$ -unitary bialgebra of weight  $\lambda$  arising from a matrix algebra.

**Definition 2.5.** [41, Chapter 17] A **matrix algebra**  $M_n(\mathbf{k})$  is a collection of  $n \times n$  matrices over  $\mathbf{k}$  that form a unitary associative algebra under matrix addition and matrix multiplication.

The multiplication on  $M_n(\mathbf{k})$  will be denoted by  $m$ . We now define a coproduct on matrix algebra  $M_n(\mathbf{k})$  to equip it with a coalgebra structure, with an eye toward constructing an  $\epsilon$ -unitary bialgebra of weight  $\lambda$  on it.

2.2.1. *The case of  $\lambda = 0$ .* Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$ . For  $M \in M_n(\mathbf{k})$ , define

$$(3) \quad \Delta_L(M) := ML \otimes L - L \otimes LM.$$

Note that  $\Delta_L(E) = 0$ , where  $E \in M_n(\mathbf{k})$  is the identity matrix. We observe that  $M_n(\mathbf{k})$  is closed under the coproduct  $\Delta_L$ .

**Lemma 2.6.** Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$ . Then for  $M, N \in M_n(\mathbf{k})$ ,

$$\Delta_L(MN) = M \cdot \Delta_L(N) + \Delta_L(M) \cdot N.$$

*Proof.* We have

$$\begin{aligned} M \cdot \Delta_L(N) + \Delta_L(M) \cdot N &= M \cdot (NL \otimes L - L \otimes LN) + (ML \otimes L - L \otimes LM) \cdot N \quad (\text{by Eq. (3)}) \\ &= MNL \otimes L - ML \otimes LN + ML \otimes LN - L \otimes LMN \quad (\text{by Eq. (1)}) \\ &= MNL \otimes L - L \otimes LMN \\ &= (MN)L \otimes L - L \otimes L(MN) \\ &= \Delta_L(MN) \quad (\text{by Eq. (3)}). \end{aligned}$$

This completes the proof.  $\square$

**Lemma 2.7.** Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$ . Then the pair  $(M_n(\mathbf{k}), \Delta_L)$  is a coalgebra (without counit).

*Proof.* It is enough to show the coassociative law:

$$(4) \quad (\text{id} \otimes \Delta_L)\Delta_L(M) = (\Delta_L \otimes \text{id})\Delta_L(M) \text{ for } M \in M_n(\mathbf{k}).$$

Applying Eq. (3) and  $L^2 = 0$ , we have

$$(5) \quad \Delta_L(L) = L^2 \otimes L - L \otimes L^2 = 0.$$

On the one hand,

$$\begin{aligned} (\text{id} \otimes \Delta_L)\Delta_L(M) &= (\text{id} \otimes \Delta_L)(ML \otimes L - L \otimes LM) \quad (\text{by Eq. (3)}) \\ &= ML \otimes \Delta_L(L) - L \otimes \Delta_L(LM) \\ &= -L \otimes \Delta_L(LM) \quad (\text{by Eq. (5)}) \\ &= -L \otimes (LML \otimes L - L \otimes L^2M) \quad (\text{by Eq. (3)}) \\ &= -L \otimes LML \otimes L \quad (\text{by } L^2 = 0). \end{aligned}$$

On the other hand,

$$\begin{aligned} (\Delta_L \otimes \text{id})\Delta_L(M) &= (\Delta_L \otimes \text{id})(ML \otimes L - L \otimes LM) \quad (\text{by Eq. (3)}) \\ &= \Delta_L(ML) \otimes L - \Delta_L(L) \otimes LM \\ &= \Delta_L(ML) \otimes L \quad (\text{by Eq. (5)}) \\ &= (ML^2 \otimes L - L \otimes LML) \otimes L \quad (\text{by Eq. (3)}) \\ &= -L \otimes LML \otimes L \quad (\text{by } L^2 = 0), \end{aligned}$$

whence Eq. (4) holds.  $\square$

Now we arrive at our first main result in this subsection.

**Theorem 2.8.** Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$ . Then the quadruple  $(M_n(\mathbf{k}), m, E, \Delta_L)$  is an  $\epsilon$ -unitary bialgebra of weight zero.



*Proof.* It follows from Lemmas 2.6 and 2.7.  $\square$

**Remark 2.9.** (a) It should be pointed out that Theorem 2.8 generalizes the case of  $n = 2$  studied in [1, Example 2.3.7].

(b) By Remark 2.3 (b), it can't add a counit to  $(M_n(\mathbf{k}), m, E, \Delta_L)$ .

2.2.2. *The case of  $\lambda \neq 0$ .* Let  $\lambda \in \mathbf{k} \setminus \{0\}$  be given. For any  $M \in M_n(\mathbf{k})$ , define

$$(6) \quad \Delta_\lambda(M) := M \otimes (-\lambda E),$$

where  $E \in M_n(\mathbf{k})$  is the identity matrix. Note that  $\Delta_\lambda(E) = -\lambda(E \otimes E)$  and  $M_n(\mathbf{k})$  is closed under the coproduct  $\Delta_\lambda$ .

**Lemma 2.10.** Let  $M, N \in M_n(\mathbf{k})$ . Then

$$\Delta_\lambda(MN) = M \cdot \Delta_\lambda(N) + \Delta_\lambda(M) \cdot N + \lambda(M \otimes N).$$

*Proof.* We have

$$\begin{aligned} M \cdot \Delta_\lambda(N) + \Delta_\lambda(M) \cdot N &= M \cdot (N \otimes (-\lambda E)) + (M \otimes (-\lambda E)) \cdot N \quad (\text{by Eq. (6)}), \\ &= MN \otimes (-\lambda E) - \lambda(M \otimes N) \quad (\text{by Eq. (1)}) \\ &= \Delta_\lambda(MN) - \lambda(M \otimes N). \end{aligned}$$

Thus

$$\Delta_\lambda(MN) = M \cdot \Delta_\lambda(N) + \Delta_\lambda(M) \cdot N + \lambda(M \otimes N).$$

This completes the proof.  $\square$

**Lemma 2.11.** The pair  $(M_n(\mathbf{k}), \Delta_\lambda)$  is a coalgebra (without counit).

*Proof.* It suffices to show the coassociative law:

$$(7) \quad (\text{id} \otimes \Delta_\lambda)\Delta_\lambda(M) = (\Delta_\lambda \otimes \text{id})\Delta_\lambda(M) \text{ for } M \in M_n(\mathbf{k}).$$

By Eq. (6), we have

$$\begin{aligned} (\text{id} \otimes \Delta_\lambda)\Delta_\lambda(M) &= (\text{id} \otimes \Delta_\lambda)(M \otimes (-\lambda E)) = M \otimes (-\lambda)\Delta_\lambda(E) = \lambda^2 M \otimes E \otimes E \\ &= M \otimes (-\lambda E) \otimes (-\lambda E) = \Delta_\lambda(M) \otimes (-\lambda E) \\ &= (\Delta_\lambda \otimes \text{id})(M \otimes (-\lambda E)) = (\Delta_\lambda \otimes \text{id})\Delta_\lambda(M), \end{aligned}$$

as desired.  $\square$

**Theorem 2.12.** Let  $\lambda \in \mathbf{k} \setminus \{0\}$  be given. Then the quadruple  $(M_n(\mathbf{k}), m, E, \Delta_\lambda)$  is an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ .

*Proof.* It follows from Lemmas 2.10 and 2.11.  $\square$

**2.3. An infinitesimal unitary Hopf algebra on a matrix algebra.** In this subsection, we equip the  $\epsilon$ -unitary bialgebra  $(M_n(\mathbf{k}), m, E, \Delta_\lambda)$  of weight zero with an antipode such that it is further an  $\epsilon$ -unitary Hopf algebra, under the view of Aguiar [1]. Denote by  $\text{Hom}_{\mathbf{k}}(A, B)$  the module consisting of linear maps from  $A$  to  $B$  throughout the remainder of this section.



**Definition 2.13.** [19, Chapter 4.1] Let  $A = (A, m, 1, \Delta, \varepsilon)$  be a classical bialgebra. Then the convolution product  $*$  on  $\text{Hom}_{\mathbf{k}}(A, A)$  is defined to be the composition:

$$f * g := m(f \otimes g)\Delta \text{ for } f, g \in \text{Hom}_{\mathbf{k}}(A, A),$$

and the triple  $(\text{Hom}_{\mathbf{k}}(A, A), *, 1 \circ \varepsilon)$  is called a convolution algebra, where  $1 \circ \varepsilon$  is the unit with respect to  $*$ . The antipode  $S$  is defined to be the inverse of the identity map with respect to the convolution product.

**Remark 2.14.** The question facing us is whether we can define the antipode for an  $\varepsilon$ -unitary bialgebra as one does for classical bialgebras  $A$ . Aguiar [1, Remark 2.2] answers this question ‘No’ due to the lack of the unit  $1 \circ \varepsilon$  with respect to  $*$ , see Remark 2.3 (b).

However, Aguiar [1] provided the perfect notion of antipode  $S$  for an  $\varepsilon$ -bialgebra.

**Definition 2.15.** [1, Section 3] Let  $A = (A, m, \Delta)$  be an  $\varepsilon$ -bialgebra. Then the **circular convolution**  $\otimes$  on  $\text{Hom}_{\mathbf{k}}(A, A)$  is defined by

$$f \otimes g := f * g + f + g, \text{ that is, } (f \otimes g)(a) := \sum_{(a)} f(a_{(1)})g(a_{(2)}) + f(a) + g(a) \text{ for } a \in A.$$

Note that  $f \otimes 0 = f = 0 \otimes f$  and so  $0 \in \text{Hom}_{\mathbf{k}}(A, A)$  is the unit with respect to the circular convolution  $\otimes$ .

Further Aguiar introduced the concept of an infinitesimal Hopf algebra via circular convolution [1, Definition 3.1]. Inspired by this, we set the following definition in [31].

**Definition 2.16.** An infinitesimal unitary bialgebra  $(A, m, 1, \Delta)$  of weight  $\lambda$  is called an **infinitesimal unitary Hopf algebra** (abbreviated  **$\varepsilon$ -unitary Hopf algebra**) of **weight**  $\lambda$  if the identity map  $\text{id} \in \text{Hom}_{\mathbf{k}}(A, A)$  is invertible with respect to the circular convolution. In this case, the inverse  $S \in \text{Hom}_{\mathbf{k}}(A, A)$  of  $\text{id}$  is called the **antipode** of  $A$ . It is characterized by the equations

$$(8) \quad \sum_{(a)} S(a_{(1)})a_{(2)} + S(a) + a = 0 = \sum_{(a)} a_{(1)}S(a_{(2)}) + a + S(a) \text{ for } a \in A,$$

where  $\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}$ .

The  $\varepsilon$ -unitary Hopf algebra satisfies many properties analogous to those of a classical Hopf algebra [1, Propositions 3.7, 3.12].

**Remark 2.17.** (a) Let  $A$  be an  $\varepsilon$ -unitary Hopf algebra of weight zero with antipode  $S$ . Then

$$S(xy) = -S(x)S(y) \text{ and } \sum_{(x)} S(x_{(1)}) \otimes S(x_{(2)}) = - \sum_{(S(x))} S(x)_{(1)} \otimes S(x)_{(2)} = -\Delta S(x).$$

(b) If  $(A, m, 1, \Delta)$  is an  $\varepsilon$ -unitary Hopf algebra of weight zero with the antipode  $S$ , then so is  $(A, m^{\text{op}}, 1, \Delta^{\text{cop}})$  with the same antipode  $S$ .

(c) It follows from Eq. (8) that  $S(1) = -1$  by taking  $a = 1$ .

**Definition 2.18.** [1, Section 4] Let  $A$  be an algebra and  $C$  a coalgebra. The map  $f : C \rightarrow A$  is called **locally nilpotent** with respect to convolution  $*$  if for each  $c \in C$  there is some  $n \geq 1$  such that

$$(9) \quad f^{*n}(c) := \sum_{(c)} f(c_{(1)})f(c_{(2)}) \cdots f(c_{(n+1)}) = 0,$$

where  $c_{(1)}, \dots, c_{(n+1)}$  are from the Sweedler notation  $\Delta^n(c) = \sum_{(c)} c_{(1)} \otimes \dots \otimes c_{(n+1)}$  and  $f^{*n}$  is defined inductively by

$$f^{*1}(c) := \sum_{(c)} f(c_{(1)})f(c_{(2)}) \text{ and } f^{*(n)} := f^{*(n-1)} * f.$$

**Lemma 2.19.** Let  $(M_n(\mathbf{k}), m, E, \Delta_L)$  be the  $\epsilon$ -unitary bialgebra of weight zero in Theorem 2.8 and

$$D := m\Delta_L : M_n(\mathbf{k}) \rightarrow M_n(\mathbf{k}).$$

Then for each  $k \geq 0$  and  $M \in M_n(\mathbf{k})$ ,  $D^{*(k+1)}(M) = 0$  and so  $D$  is locally nilpotent.

*Proof.* It suffices to prove the first statement by induction on  $k \geq 0$ . Using Sweedler notation, we may write

$$\Delta_L(M) = \sum_{(M)} M_{(1)} \otimes M_{(2)} \text{ for } M \in M_n(\mathbf{k}).$$

For the initial step of  $k = 0$ , we have

$$D^{*(1)}(M) = \sum_{(M)} D(M_{(1)})D(M_{(2)}) = 0,$$

where the last step follows from

$$D(M_{(1)}) = m\Delta_L(M_{(1)}) = M_{(1)}L^2 - L^2M_{(1)} = 0$$

by Eq. (3) and  $L^2 = 0$ . Assume the result is true for  $k = \ell$  for an  $\ell \geq 1$ , and consider the case when  $k = \ell + 1$ . Then

$$D^{*(\ell+1)}(M) = (D^{*\ell} * D)(M) = m(D^{*\ell} \otimes D)\Delta_L(M) = \sum_{(M)} D^{*\ell}(M_{(1)})D(M_{(2)}) = 0,$$

where the last step employs the induction hypothesis. This completes the proof.  $\square$

Denote by  $\mathbb{R}$  and  $\mathbb{C}$  the field of real numbers and the field of complex numbers, respectively.

**Lemma 2.20.** [1, Proposition 4.5] Let  $(A, m, \Delta)$  be an  $\epsilon$ -bialgebra and  $D := m\Delta$ , and let  $\mathbf{k}$  be a field. Suppose that either

- (a)  $\mathbf{k} = \mathbb{R}$  or  $\mathbb{C}$  and  $A$  is finite dimensional, or
- (b)  $D$  is locally nilpotent and  $\text{char}(\mathbf{k}) = 0$ .

Then  $A$  is an  $\epsilon$ -Hopf algebra with bijective antipode  $S = -\sum_{n=0}^{\infty} \frac{1}{n!}(-D)^n$ .

Indeed, in the above lemma, we can replace the condition that  $\mathbf{k}$  is a field by  $\mathbb{Q} \subseteq \mathbf{k}$ .

**Theorem 2.21.** Let  $\mathbb{Q} \subseteq \mathbf{k}$ . Then the quadruple  $(M_n(\mathbf{k}), m, E, \Delta_L)$  is an  $\epsilon$ -unitary Hopf algebra of weight zero with the bijective antipode  $S = -\sum_{n=0}^{\infty} \frac{1}{n!}(-D)^n$ .

*Proof.* By Theorem 2.8,  $(M_n(\mathbf{k}), m, E, \Delta_L)$  is an  $\epsilon$ -unitary bialgebra of weight zero. From Lemmas 2.19, 2.20 and Definition 2.16,  $(M_n(\mathbf{k}), m, E, \Delta_L)$  is an  $\epsilon$ -unitary Hopf algebra of weight zero with bijective antipode  $S = -\sum_{n=0}^{\infty} \frac{1}{n!}(-D)^n$ .  $\square$

### 3. PRE-LIE ALGEBRAS ON MATRIX ALGEBRAS

This section is devoted to recall that an  $\epsilon$ -unitary bialgebra of an arbitrary weight  $\lambda$  gives rise to a pre-Lie algebra [31], which generalizes the construction of the pre-Lie algebra from an infinitesimal bialgebra [5]. As a consequence, we equip  $M_n(\mathbf{k})$  with two pre-Lie bialgebraic structures and two Lie algebraic structures

**3.1. Pre-Lie algebras and weighted infinitesimal unitary bialgebras.** In this subsection, we first recall the concept of pre-Lie algebras and then show the connection from weighted  $\epsilon$ -unitary bialgebras to pre-Lie algebras.

**Definition 3.1.** [44] A (left) **pre-Lie algebra** is a  $\mathbf{k}$ -module  $A$  together with a binary linear operation  $\triangleright : A \otimes A \rightarrow A$  satisfying

$$(10) \quad (a \triangleright b) \triangleright c - a \triangleright (b \triangleright c) = (b \triangleright a) \triangleright c - b \triangleright (a \triangleright c) \text{ for } a, b, c \in A.$$

**Example 3.2.** Here are two well-known pre-Lie algebras on dendriform dialgebras and Rota-Baxter algebras, respectively.

- (a) Let  $(A, <, >)$  be a dendriform dialgebra. Then the multiplication  $\star$  defined by  $a \star b = a < b + a > b$  gives an associative algebra [5]. In addition, define

$$\triangleright : A \otimes A \rightarrow A, a \otimes b \mapsto a > b - a < b \text{ for } a, b \in A.$$

Then  $A$  together with  $\triangleright$  is a pre-Lie algebra [5].

- (b) Let  $(A, P)$  be a Rota-Baxter algebra of weight  $\lambda$ . If the weight  $\lambda = 0$ , then the binary operation

$$\triangleright : A \otimes A \rightarrow A, a \otimes b \mapsto P(a) \cdot b - b \cdot P(a) \text{ for } a, b \in A,$$

defines a pre-Lie algebra. If the weight  $\lambda = -1$ , then the binary operation

$$\triangleright : A \otimes A \rightarrow A, a \otimes b \mapsto P(a) \cdot b - b \cdot P(a) - x \cdot y \text{ for } a, b \in A,$$

defines a pre-Lie algebra [6].

Let  $(A, \triangleright)$  be a pre-Lie algebra. For any  $a \in A$ , let

$$L_a : A \rightarrow A, b \mapsto a \triangleright b$$

be the left multiplication operator. Let

$$L : A \rightarrow \text{Hom}_{\mathbf{k}}(A, A), a \mapsto L_a.$$

The close relation between pre-Lie algebras and Lie algebras is characterized by the following two fundamental properties.

**Lemma 3.3.** (a) [32, Theorem 1] Let  $(A, \triangleright)$  be a pre-Lie algebra. Define for elements in  $A$  a new multiplication by setting

$$[a, b] := a \triangleright b - b \triangleright a \text{ for } a, b \in A.$$

Then  $(A, [-, -])$  is a Lie algebra.

- (b) [7, Proposition 1.2] Eq. (10) rewrites as

$$L_{[a,b]} = L_a \circ L_b - L_b \circ L_a = [L_a, L_b],$$

which implies that  $L : (A, [-, -]) \rightarrow \text{Hom}_{\mathbf{k}}(A, A)$  with  $a \mapsto L_a$  gives a representation of the Lie algebra  $(A, [-, -])$ .

**Remark 3.4.** By Lemma 3.3, a pre-Lie algebra induces a Lie algebra whose left multiplication operators give a representation of the associated commutator Lie algebra.

Let  $(A, m, 1, \Delta)$  be an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ . Define

$$(11) \quad \triangleright : A \otimes A \rightarrow A, a \otimes b \mapsto a \triangleright b := \sum_{(b)} b_{(1)} a b_{(2)},$$

where  $b_{(1)}$  and  $b_{(2)}$  are from the Sweedler notation  $\Delta(b) = \sum_{(b)} b_{(1)} \otimes b_{(2)}$ . The following result captures the connection from weighted  $\epsilon$ -unitary bialgebras to pre-Lie algebras [31]. For the completeness, we record the proof here.

**Theorem 3.5.** *Let  $(A, m, 1, \Delta)$  be an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ . Then  $A$  equipped with the  $\triangleright$  in Eq. (11) is a pre-Lie algebra.*

*Proof.* For any  $a, b, c \in A$ , using the Sweedler notation  $\Delta(c) = \sum_{(c)} c_{(1)} \otimes c_{(2)}$ , we have

$$\begin{aligned}
\Delta\left(\sum_{(c)} c_{(1)}bc_{(2)}\right) &= \sum_{(c)} \Delta(c_{(1)}bc_{(2)}) = \sum_{(c)} \left( c_{(1)}b \cdot \Delta(c_{(2)}) + \Delta(c_{(1)}b) \cdot c_{(2)} + \lambda c_{(1)}b \otimes c_{(2)} \right) \quad (\text{by Eq. (2)}) \\
&= \sum_{(c)} \left( c_{(1)}b \cdot \Delta(c_{(2)}) + (c_{(1)} \cdot \Delta(b) + \Delta(c_{(1)}) \cdot b + \lambda(c_{(1)} \otimes b)) \cdot c_{(2)} + \lambda c_{(1)}b \otimes c_{(2)} \right) \\
&= \sum_{(c)} \left( c_{(1)}b \cdot \Delta(c_{(2)}) + c_{(1)} \cdot \Delta(b) \cdot c_{(2)} + \Delta(c_{(1)}) \cdot bc_{(2)} + \lambda c_{(1)} \otimes bc_{(2)} + \lambda c_{(1)}b \otimes c_{(2)} \right) \\
&= \sum_{(c)} \left( \Delta(c_{(1)}) \cdot bc_{(2)} + c_{(1)} \cdot \Delta(b) \cdot c_{(2)} + c_{(1)}b \cdot \Delta(c_{(2)}) + \lambda c_{(1)} \otimes bc_{(2)} + \lambda c_{(1)}b \otimes c_{(2)} \right) \\
&= \sum_{(c)} \left( \sum_{(c_{(1)})} c_{(1)(1)} \otimes c_{(1)(2)}bc_{(2)} + \sum_{(b)} c_{(1)}b_{(1)} \otimes b_{(2)}c_{(2)} + \sum_{(c_{(2)})} c_{(1)}bc_{(2)(1)} \otimes c_{(2)(2)} \right. \\
&\quad \left. + \lambda c_{(1)} \otimes bc_{(2)} + \lambda c_{(1)}b \otimes c_{(2)} \right) \\
&= \sum_{(e)} c_{(1)} \otimes c_{(2)}bc_{(3)} + \sum_{(e),(b)} c_{(1)}b_{(1)} \otimes b_{(2)}c_{(2)} + \sum_{(e)} c_{(1)}bc_{(2)} \otimes c_{(3)} \\
&\quad + \lambda \sum_{(e)} c_{(1)} \otimes bc_{(2)} + \lambda \sum_{(e)} c_{(1)}b \otimes c_{(2)} \quad (\text{by the coassociative law}).
\end{aligned}$$

Together this with Eq. (11), we obtain

$$\begin{aligned}
a \triangleright (b \triangleright c) &= a \triangleright \left( \sum_{(c)} c_{(1)}bc_{(2)} \right) \\
&= \sum_{(c)} c_{(1)}ac_{(2)}bc_{(3)} + \sum_{(e),(b)} c_{(1)}b_{(1)}ab_{(2)}c_{(2)} + \sum_{(e)} c_{(1)}bc_{(2)}ac_{(3)} \\
&\quad + \lambda \sum_{(e)} c_{(1)}abc_{(2)} + \lambda \sum_{(e)} c_{(1)}bac_{(2)}.
\end{aligned}$$

Moreover

$$(a \triangleright b) \triangleright c = \left( \sum_{(b)} b_{(1)}ab_{(2)} \right) \triangleright c = \sum_{(e),(b)} c_{(1)}b_{(1)}ab_{(2)}c_{(2)}.$$

Thus

(12)

$$(a \triangleright b) \triangleright c - a \triangleright (b \triangleright c) = - \sum_{(e)} c_{(1)}ac_{(2)}bc_{(3)} - \sum_{(e)} c_{(1)}bc_{(2)}ac_{(3)} - \lambda \sum_{(e)} \left( c_{(1)}abc_{(2)} + c_{(1)}bac_{(2)} \right).$$

Observe that Eq. (12) is symmetric in  $a$  and  $b$ . Hence

$$(a \triangleright b) \triangleright c - a \triangleright (b \triangleright c) = (b \triangleright a) \triangleright c - b \triangleright (a \triangleright c),$$

and so  $(A, \triangleright)$  is a pre-Lie algebra.  $\square$

**3.2. Two pre-Lie and a new Lie algebraic structures on a matrix algebra.** In this subsection, we proceed to give two pre-Lie algebraic structures on a matrix algebra. Consequently, a new Lie algebraic structure on a matrix algebra is induced by Lemma 3.3 (a).

3.2.1. *The case of  $\lambda = 0$ .* By Theorem 2.8,  $(M_n(\mathbf{k}), \mathfrak{m}, E, \Delta_L)$  is an  $\epsilon$ -unitary bialgebra of weight zero. Applying Eq. (11), we define

$$(13) \quad \triangleright_L : M_n(\mathbf{k}) \otimes M_n(\mathbf{k}) \rightarrow M_n(\mathbf{k}), \quad M \triangleright_L N := \sum_{(N)} N_{(1)} M N_{(2)},$$

where  $N_{(1)}$  and  $N_{(2)}$  are from  $\Delta_L(N) = \sum_{(N)} N_{(1)} \otimes N_{(2)}$ .

**Theorem 3.6.** *Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$ . Then the pair  $(M_n(\mathbf{k}), \triangleright_L)$  is a pre-Lie algebra and so  $(M_n(\mathbf{k}), [-, -]_L)$  is a Lie algebra, where*

$$[M, N]_L := NLML + LNLM - MLNL - LMLN.$$

*Proof.* By Theorems 2.8 and 3.5,  $(M_n(\mathbf{k}), \triangleright_L)$  is a pre-Lie algebra. The remainder follows from Lemma 3.3 (a) and

$$\begin{aligned} [M, N]_L &= M \triangleright_L N - N \triangleright_L M = \sum_{(N)} N_{(1)} M N_{(2)} - \sum_{(M)} M_{(1)} N M_{(2)} \quad (\text{by Eq. (13)}) \\ &= NLML - LMLN - (MLNL - LNLM) \quad (\text{by Eq. (3)}) \\ &= NLML + LNLM - MLNL - LMLN. \end{aligned}$$

This completes the proof.  $\square$

The following example exposes that the Lie bracket  $[-, -]_L$  is different from the one given by commutator.

**Example 3.7.** Consider the matrix algebra  $M_2(\mathbf{k})$ . Let

$$L = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, N = \begin{pmatrix} e & f \\ g & h \end{pmatrix}.$$

Then  $L^2 = 0$  and by Theorem 3.6,

$$[M, N]_L = \begin{pmatrix} 0 & ec + gd - ag - ch \\ 0 & 0 \end{pmatrix},$$

which is different from the classical Lie bracket

$$[M, N] := MN - NM = \begin{pmatrix} bg - fc & af + bh - eb - fd \\ ce + dg - ga - hc & cf - gb \end{pmatrix}.$$

3.2.2. *The case of  $\lambda \neq 0$ .* By Theorem 2.12,  $(M_n(\mathbf{k}), \mathfrak{m}, E, \Delta_\lambda)$  is an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ . Using Eq. (11), we define

$$\triangleright_\lambda : M_n(\mathbf{k}) \otimes M_n(\mathbf{k}) \rightarrow M_n(\mathbf{k}), \quad M \triangleright_\lambda N := \sum_{(N)} N_{(1)} M N_{(2)},$$

where  $N_{(1)}$  and  $N_{(2)}$  are from  $\Delta_\lambda(N) = \sum_{(N)} N_{(1)} \otimes N_{(2)}$ .

**Theorem 3.8.** *Let  $\lambda \in \mathbf{k} \setminus \{0\}$  be given. Then the pair  $(M_n(\mathbf{k}), \triangleright_\lambda)$  is a pre-Lie algebra and so  $(M_n(\mathbf{k}), [-, -]_\lambda)$  is a Lie algebra, where*

$$[M, N]_\lambda := \lambda(MN - NM).$$

*Proof.* By Theorems 2.12 and 3.5,  $(M_n(\mathbf{k}), \triangleright_\lambda)$  is a pre-Lie algebra. The remainder follows from Lemma 3.3 (a) and Eq. (6).  $\square$

**Remark 3.9.** Taking  $\lambda = 1$  in Theorem 3.8, a surprising phenomenon shows that the Lie bracket derived from  $\Delta_\lambda$  is precisely the classical Lie bracket given by commutator on matrix algebras.

#### 4. WEIGHTED AYBES AND WEIGHTED $\epsilon$ -UNITARY BIALGEBRAS

In this section, we first introduce the concept of a weighted associative Yang-Baxter equation, which generalize the results studied in [1, Section 5]. As an application, we give a solution of a (homogeneous) associative Yang-Baxter equation for matrix algebras. We finally derive a Rota-Baxter operator of weight  $-\lambda$  from a solution of a weighted associative Yang-Baxter equation.

**4.1. Weighted AYBEs.** Let us first propose the concept of weighted associative Yang-Baxter equations. Let  $A$  be a unitary algebra. For an element  $r = \sum_i u_i \otimes v_i \in A \otimes A$ , we write

$$r_{12} = \sum_i u_i \otimes v_i \otimes 1, \quad r_{13} = \sum_i u_i \otimes 1 \otimes v_i, \quad r_{23} = \sum_i 1 \otimes u_i \otimes v_i.$$

**Definition 4.1.** Let  $\lambda$  be a given element of  $\mathbf{k}$  and  $A$  a unitary algebra.

(a) The equation

$$(14) \quad r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = \lambda r_{13}$$

is called the **associative Yang-Baxter equation (AYBE for short) of weight  $\lambda$** .

(b) An element  $r = \sum_i u_i \otimes v_i \in A \otimes A$  is called a **solution of the associative Yang-Baxter equation of weight  $\lambda$**  in  $A$  if it satisfies Eq. (14).

**Remark 4.2.** (a) When  $\lambda = 0$ , the equation

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = 0$$

is called a **homogeneous associative Yang-Baxter equation**, initiated by Aguiar [1] and further studied in [10].

(b) When  $\lambda \neq 0$ , the equation

$$(15) \quad r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = \lambda r_{13},$$

is called a **non-homogeneous associative Yang-Baxter equation**.

(c) When  $\lambda = -1$ , the equation

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = -r_{13}$$

is called a **modified associative Yang-Baxter equation** studied by Ebrahimi-Fard [21].

We now give some basic definitions and notations that will be used in this section. We refer to [1, Section 5] for the classical results in the case of  $\lambda = 0$ .

**Definition 4.3.** Let  $\lambda$  be a given element of  $\mathbf{k}$ , and let  $A$  be a unitary algebra and  $W$  an  $A$ -bimodule.

(a) A linear map  $\Delta : A \rightarrow A \otimes A$  is called a **derivation of weight  $\lambda$**  if it satisfies Eq. (2), that is,

$$\Delta(ab) = a \cdot \Delta(b) + \Delta(a) \cdot b + \lambda(a \otimes b) \text{ for } a, b \in A.$$

(b) A linear map  $\Delta_r : A \rightarrow W$  associated to an element  $r \in W$  is called **principal** if it satisfies

$$(16) \quad \Delta_r(a) := a \cdot r - r \cdot a - \lambda(a \otimes 1) \text{ for } a \in A.$$

(c) An element  $r \in W$  is called **A-invariant** if it satisfies

$$a \cdot r = r \cdot a \text{ for } a \in A.$$

For an algebra  $A$ ,  $A \otimes A \otimes A$  is viewed as an  $A$ -bimodule via

$$a \cdot (b \otimes c \otimes d) = ab \otimes c \otimes d \text{ and } (b \otimes c \otimes d) \cdot a = b \otimes c \otimes da,$$

where  $a, b, c, d \in A$ .

**Proposition 4.4.** *The  $\Delta_r$  defined by Eq. (16) is a derivation of weight  $\lambda$ .*

*Proof.* For  $a, b \in A$ , it follows from Eq. (16) that

$$\begin{aligned} a \cdot \Delta_r(b) + \Delta_r(a) \cdot b &= a \cdot (b \cdot r - r \cdot b - \lambda(b \otimes 1)) + (a \cdot r - r \cdot a - \lambda(a \otimes 1)) \cdot b \\ &= ab \cdot r - a \cdot r \cdot b - \lambda ab \otimes 1 + a \cdot r \cdot b - r \cdot ab - \lambda(a \otimes b) \\ &= ab \cdot r - r \cdot ab - \lambda ab \otimes 1 - \lambda(a \otimes b) \\ &= \Delta_r(ab) - \lambda(a \otimes b), \end{aligned}$$

as desired. □

The following result plays a crucial role to obtain a relationship between solutions of weighted AYBEs and weighted  $\epsilon$ -unitary bialgebras.

**Theorem 4.5.** *Let  $A$  be a unitary algebra and  $r = \sum_i u_i \otimes v_i \in A \otimes A$ . Then the principle derivation of weight  $\lambda$*

$$\Delta_r : A \rightarrow A \otimes A, \quad a \mapsto a \cdot r - r \cdot a - \lambda(a \otimes 1)$$

*is coassociative if and only if the element*

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} - \lambda r_{13} \in A \otimes A \otimes A$$

*is A-invariant.*

*Proof.* On the one hand,

$$\begin{aligned} (\Delta_r \otimes \text{id}) \circ \Delta_r(a) &= (\Delta_r \otimes \text{id})(a \cdot r - r \cdot a - \lambda a \otimes 1) \quad (\text{by Eq. (16)}) \\ &= (\Delta_r \otimes \text{id}) \left( \sum_i (au_i \otimes v_i - u_i \otimes v_i a) - \lambda a \otimes 1 \right) \\ &= \sum_i \Delta_r(au_i) \otimes v_i - \sum_i \Delta_r(u_i) \otimes v_i a - \lambda \Delta_r(a) \otimes 1 \\ &= \sum_i (au_i \cdot r - r \cdot au_i - \lambda au_i \otimes 1) \otimes v_i - \sum_i (u_i \cdot r - r \cdot u_i - \lambda u_i \otimes 1) \otimes v_i a \\ &\quad - \lambda(a \cdot r - r \cdot a - \lambda a \otimes 1) \otimes 1 \quad (\text{by Eq. (16)}) \\ &= \sum_{i,j} (au_i u_j \otimes v_j \otimes v_i - u_j \otimes v_j a u_i \otimes v_i) - \lambda \sum_i au_i \otimes 1 \otimes v_i \\ &\quad - \sum_{i,j} (u_i u_j \otimes v_j \otimes v_i a - u_j \otimes v_j u_i \otimes v_i a) + \lambda \sum_i u_i \otimes 1 \otimes v_i a \\ &\quad - \lambda \sum_i (au_i \otimes v_i \otimes 1 - u_i \otimes v_i a \otimes 1) + \lambda^2(a \otimes 1 \otimes 1) \\ &= a \cdot r_{13}r_{12} - r_{12}(1 \otimes a \otimes 1)r_{23} - \lambda a \cdot r_{13} - r_{13}r_{12} \cdot a + r_{12}r_{23} \cdot a + \lambda r_{13} \cdot a \end{aligned}$$



$$- \lambda a \cdot r_{12} + \lambda r_{12}(1 \otimes a \otimes 1) + \lambda^2(a \otimes 1 \otimes 1).$$

On the other hand,

$$\begin{aligned} (\text{id} \otimes \Delta_r) \circ \Delta_r(a) &= (\text{id} \otimes \Delta_r)(a \cdot r - r \cdot a - \lambda a \otimes 1) \quad (\text{by Eq. (16)}) \\ &= (\text{id} \otimes \Delta_r) \left( \sum_i (au_i \otimes v_i - u_i \otimes v_i a) - \lambda a \otimes 1 \right) \\ &= \sum_i au_i \otimes \Delta_r(v_i) - \sum_i u_i \otimes \Delta_r(v_i a) - \lambda a \otimes \Delta_r(1) \\ &= \sum_i au_i \otimes (v_i \cdot r - r \cdot v_i - \lambda v_i \otimes 1) - \sum_i u_i \otimes (v_i a \cdot r - r \cdot v_i a - \lambda v_i a \otimes 1) \\ &\quad + \lambda^2 a \otimes 1 \otimes 1 \quad (\text{by Eq. (16)}) \\ &= \sum_{i,j} (au_i \otimes v_i u_j \otimes v_j - au_i \otimes u_j \otimes v_j v_i) - \lambda \sum_i au_i \otimes v_i \otimes 1 \\ &\quad - \sum_{i,j} (u_i \otimes v_i a u_j \otimes v_j - u_i \otimes u_j \otimes v_j v_i a) + \lambda \sum_i u_i \otimes v_i a \otimes 1 + \lambda^2(a \otimes 1 \otimes 1) \\ &= a \cdot r_{12} r_{23} - a \cdot r_{23} r_{13} - \lambda a \cdot r_{12} - r_{12}(1 \otimes a \otimes 1) r_{23} + r_{23} r_{13} \cdot a + \lambda r_{12}(1 \otimes a \otimes 1) \\ &\quad + \lambda^2(a \otimes 1 \otimes 1). \end{aligned}$$

Comparing the two hands,  $\Delta_r$  is coassociative if and only if

$$a \cdot r_{13} r_{12} - \lambda a \cdot r_{13} - r_{13} r_{12} \cdot a + r_{12} r_{23} \cdot a + \lambda r_{13} \cdot a = a \cdot r_{12} r_{23} - a \cdot r_{23} r_{13} + r_{23} r_{13} \cdot a$$

if and only if

$$a \cdot (r_{13} r_{12} - r_{12} r_{23} + r_{23} r_{13} - \lambda r_{13}) = (r_{13} r_{12} - r_{12} r_{23} + r_{23} r_{13} - \lambda r_{13}) \cdot a$$

if and only if

$$r_{13} r_{12} - r_{12} r_{23} + r_{23} r_{13} - \lambda r_{13}$$

is  $A$ -invariant. □

Now we arrive at our main result in this subsection.

**Theorem 4.6.** *Let  $(A, m, 1)$  be a unitary algebra and  $r \in A \otimes A$ . If  $r$  is a solution of an AYBE of weight  $\lambda$  in  $A$ , then the quadruple  $(A, m, 1, \Delta_r)$  is an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ .*

*Proof.* Suppose that  $r$  is a solution of an AYBE of weight  $\lambda$  in  $A$ . By Theorem 4.5,  $\Delta_r : A \rightarrow A \otimes A$  is coassociative. Further by Proposition 4.4,

$$\Delta_r(ab) = a \cdot \Delta_r(b) + \Delta_r(a) \cdot b + \lambda(a \otimes b) \text{ for } a, b \in A.$$

Thus the result holds by Definition 2.1. □

**4.2. A solution of a homogeneous AYBE for matrix algebras.** In this subsection, we give a solution of a homogeneous AYBE for matrix algebras.

**Theorem 4.7.** *Let  $L$  be a matrix of  $M_n(\mathbf{k})$  such that  $L^2 = 0$ . Then  $r = L \otimes L$  is a solution of a homogeneous AYBE for matrix algebras.*

*Proof.* Since  $L^2 = 0$  and  $r = L \otimes L$ , we have

$$\begin{aligned} r_{13}r_{12} &= (L \otimes 1 \otimes L)(L \otimes L \otimes 1) = L^2 \otimes L \otimes L = 0, \\ r_{12}r_{23} &= (L \otimes L \otimes 1)(1 \otimes L \otimes L) = L \otimes L^2 \otimes L = 0, \\ r_{23}r_{13} &= (1 \otimes L \otimes L)(L \otimes 1 \otimes L) = L \otimes L \otimes L^2 = 0, \end{aligned}$$

and so

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = 0.$$

This completes the proof.  $\square$

The following result gives another way to construct an  $\epsilon$ -unitary bialgebra of weight zero on the matrix algebra  $M_n(\mathbf{k})$  obtained in Subsection 2.2.

**Corollary 4.8.** Let  $L \in M_n(\mathbf{k})$  such that  $L^2 = 0$  and  $r = L \otimes L \in M_n(\mathbf{k}) \otimes M_n(\mathbf{k})$ . Define the linear map  $\Delta_r : M_n(\mathbf{k}) \rightarrow M_n(\mathbf{k}) \otimes M_n(\mathbf{k})$  by setting

$$\Delta_r(M) := M \cdot r - r \cdot M \text{ for } M \in M_n(\mathbf{k}).$$

Then the quadruple  $(M_n(\mathbf{k}), \mathfrak{m}, E, \Delta_L)$  is an  $\epsilon$ -unitary bialgebra of weight zero.

*Proof.* It follows from Theorems 4.6 and 4.7.  $\square$

**4.3. Weighted AYBEs and Rota-Baxter operators.** In this subsection, we derive a Rota-Baxter operator of weight  $-\lambda$  from an AYBE of weight  $\lambda$ , which generalizes the result studied in [2]. See also [10, Theorem 1.3].

**Definition 4.9.** [13, 34] Let  $A$  be a unitary algebra and  $\lambda$  a given element of  $\mathbf{k}$ . A linear operator  $P : A \rightarrow A$  is called a **Rota-Baxter operator of weight  $\lambda$**  if it satisfies the Rota-Baxter equation

$$(17) \quad P(a)P(b) = P(aP(b)) + P(P(a)b) + \lambda P(ab) \text{ for } a, b \in A.$$

Then the pair  $(A, P)$  is called a **Rota-Baxter algebra of weight  $\lambda$** .

The following result captures the relation between an AYBE of weight  $\lambda$  and a Rota-Baxter operator of weight  $-\lambda$ .

**Theorem 4.10.** Let  $r = \sum_i u_i \otimes v_i$  be a solution of an AYBE of weight  $\lambda$  in  $A$ . Then the linear operator

$$(18) \quad P_r : A \rightarrow A, \quad a \mapsto P_r(a) := \sum_i u_i a v_i$$

is a Rota-Baxter operator of weight  $-\lambda$ .

*Proof.* Define a  $\mathbf{k}$ -trilinear map

$$p : A \times A \times A \rightarrow A, \quad (u, v, w) \mapsto uxvwy \text{ for } u, v, w, x, y \in A,$$

which induces a  $\mathbf{k}$ -linear map

$$h : A \otimes A \otimes A \rightarrow A, \quad u \otimes v \otimes w \mapsto uxvwy.$$

Since  $r = \sum_i u_i \otimes v_i$  is a solution of a AYBE of weight  $\lambda$ , we have

$$r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = \lambda r_{13},$$

that is,

$$(19) \quad \sum_{i,j} u_i u_j \otimes v_j \otimes v_i - \sum_{i,j} u_i \otimes v_i u_j \otimes v_j + \sum_{i,j} u_j \otimes u_i \otimes v_i v_j = \lambda \sum_i u_i \otimes 1 \otimes v_i.$$

Applying  $h$  on both sides of Eq. (19),

$$\sum_{i,j} u_i u_j x v_j y v_i - \sum_{i,j} u_i x v_i u_j y v_j + \sum_{i,j} u_j x u_i y v_i v_j = \lambda \sum_i u_i x y v_i.$$

Thus

$$P_r(P_r(x)y) - P_r(x)P_r(y) + P_r(xP_r(y)) = \lambda P_r(xy),$$

that is,

$$P_r(x)P_r(y) = P_r(P_r(x)y) + P_r(xP_r(y)) - \lambda P_r(xy),$$

which implies that  $P_r$  is a Rota-Baxter operator of weight  $-\lambda$ .  $\square$

**Corollary 4.11.** Let  $A$  be a unitary algebra.

- (a) [1] If  $r = \sum_i u_i \otimes v_i$  be a solution of a homogeneous AYBE of in  $A$ , then the linear operator  $P : A \rightarrow A$  given by Eq. (18) is a Rota-Baxter operator of weight zero.
- (b) [21] If  $r = \sum_i u_i \otimes v_i$  be a solution of a modified AYBE of in  $A$ , then the linear operator  $P : A \rightarrow A$  given by Eq. (18) is a Rota-Baxter operator of weight  $-1$ .

*Proof.* It follows from Theorem 4.10 by taking  $\lambda = 0$  and  $\lambda = -1$ , respectively.  $\square$

**4.4. A bijection between the solutions of weighted AYBEs and Rota-Baxter operators.** In this subsection, we shall give a bijection between the solutions of weighted AYBEs and Rota-Baxter operators on matrix algebra  $M_n(\mathbf{k})$ , which generalize the results studied in [33, Section 3].

Let  $E_{ij} \in M_n(\mathbf{k})$ ,  $1 \leq i, j \leq n$ , be the matrix whose entry in the  $i$ -th row,  $j$ -th column is 1, and zero in all other entries. Note that  $E_{ij}E_{kl} = \delta_{jk}E_{il}$  and  $E_{ij}$ ,  $1 \leq i, j \leq n$ , are a linear basis of  $M_n(\mathbf{k})$ . Recall that  $P_r$  is defined in Eq. (18).

**Theorem 4.12.** *Let  $r$  be a solution of an AYBE of weight  $\lambda$  in  $M_n(\mathbf{k})$ . Then the map  $r \rightarrow P_r$  is a bijection between the set of the solutions of AYBE of weight  $\lambda$  in  $M_n(\mathbf{k})$  and the set of Rota-Baxter operators of weight  $-\lambda$  on  $M_n(\mathbf{k})$ .*

*Proof.* On the one hand, a linear operator

$$(20) \quad P : M_n(\mathbf{k}) \rightarrow M_n(\mathbf{k}), \quad E_{pq} \mapsto \sum_{i,l} t_{ip}^{ql} E_{il}, \quad \text{where } t_{ip}^{ql} \in \mathbf{k}.$$

is a Rota-Baxter operator of weight  $-\lambda$  on  $M_n(\mathbf{k})$  if and only if

$$(21) \quad P(E_{ab})P(E_{cd}) = P(P(E_{ab})E_{cd}) + P(E_{ab}P(E_{cd})) - \lambda P(E_{ab}E_{cd}) \text{ for } E_{ab}, E_{cd} \in M_n(\mathbf{k}).$$

Further, it follows from Eq. (20) that

$$P(E_{ab})P(E_{cd}) = \sum_{i,j} t_{ia}^{bj} E_{ij} \sum_{k,l} t_{kc}^{dl} E_{kl} = \sum_{i,j,l} t_{ia}^{bj} t_{jc}^{dl} E_{il}.$$

Similarly,

$$P(P(E_{ab})E_{cd}) = \sum_{i,j,l} t_{ja}^{bc} t_{ij}^{dl} E_{il}, \quad P(E_{ab}P(E_{cd})) = \sum_{i,j,l} t_{bc}^{dj} t_{ia}^{jl} E_{il}$$

and

$$P(E_{ab}E_{cd}) = \begin{cases} P(E_{ad}) = \sum_{i,l} t_{ia}^{dl} E_{il} & \text{if } b = c, \\ 0 & \text{if } b \neq c. \end{cases}$$

We have two cases to consider.

**Case 1.**  $b = c$ . In this case, Eq. (21) holds if and only if

$$(22) \quad \sum_j (t_{ia}^{bj} t_{jb}^{dl} - t_{ja}^{bb} t_{ij}^{dl} - t_{bb}^{dj} t_{ia}^{jl}) = -\lambda t_{ia}^{dl}.$$

On the other hand, an element

$$r = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} \otimes E_{kl} \in M_n(\mathbf{k}) \otimes M_n(\mathbf{k})$$

is a solution of the AYBE of weight  $\lambda$  if and only if

$$(23) \quad r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = \lambda r_{13}$$

where

$$r_{12} = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} \otimes E_{kl} \otimes 1, r_{13} = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} \otimes 1 \otimes E_{kl} \text{ and } r_{23} = \sum_{i,j,k,l} t_{ij}^{kl} 1 \otimes E_{ij} \otimes E_{kl}.$$

Now we have

$$(24) \quad \begin{aligned} r_{13}r_{12} &= \sum_{i,j,k,l,p,s,t} t_{pi}^{st} t_{ij}^{kl} E_{pj} \otimes E_{kl} \otimes E_{st} = \sum_{i,j,k,l,p,s,t} t_{ij}^{kl} t_{pi}^{st} E_{pj} \otimes E_{kl} \otimes E_{st}, \\ &= \sum_{i,j,k,l,p,s,t} t_{pj}^{kl} t_{ip}^{st} E_{ij} \otimes E_{kl} \otimes E_{st} \quad (\text{by exchanging the index } i \text{ and } p). \end{aligned}$$

Similarly,

$$(25) \quad r_{12}r_{23} = \sum_{i,j,k,l,p,s,t} t_{ij}^{kl} t_{lp}^{st} E_{ij} \otimes E_{kp} \otimes E_{st} \text{ and } r_{23}r_{13} = \sum_{i,j,k,l,p,q,t} t_{ij}^{kl} t_{pq}^{st} E_{pq} \otimes E_{ij} \otimes E_{kt}.$$

By substituting the summands from Eqs. (24) and (25) in Eq. (23) and gathering the coefficient of the tensor  $E_{ij} \otimes E_{kk} \otimes E_{st}$ , we obtain

$$(26) \quad \sum_p (t_{ij}^{lp} t_{pl}^{st} - t_{pj}^{ll} t_{ip}^{st} - t_{ll}^{sp} t_{ij}^{pt}) = -\lambda t_{ij}^{st}.$$

So Eq. (23) is equivalent to Eq. (26). Note that Eq. (22) and Eq. (26) coincide up to exchange of index by the following permutation

$$\begin{pmatrix} a & b & d & l \\ j & l & s & t \end{pmatrix}.$$

In summary, a linear operator  $P$  is a Rota-Baxter operator of weight  $-\lambda$  on  $M_n(\mathbf{k})$  if and only if Eq. (22) holds. An element  $r$  is a solution of the AYBE of weight  $\lambda$  if and only if Eq. (26) is valid. Eq. (22) is equivalent to Eq. (26). So in this case, there is a bijection, namely  $\phi$ , from the set of the solutions of AYBE of weight  $\lambda$  in  $M_n(\mathbf{k})$  to the set of Rota-Baxter operators of weight  $-\lambda$  on  $M_n(\mathbf{k})$ .

**Case 2.**  $b \neq c$ . In this case,  $P(E_{ab}E_{cd}) = 0$  and we can only consider  $P$  is a Rota-Baxter operator of weight zero on  $M_n(\mathbf{k})$ . Then, with the condition  $\lambda = 0$ , Eq. (21) holds if and only if

$$(27) \quad \sum_j \left( t_{ia}^{bj} t_{jc}^{dl} - t_{ja}^{bc} t_{ij}^{dl} - t_{bc}^{dj} t_{ia}^{il} \right) = 0.$$

On the other hand, an element

$$r = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} \otimes E_{kl} \in M_n(\mathbf{k}) \otimes M_n(\mathbf{k})$$

is a solution of the AYBE of weight zero in  $M_n(\mathbf{k})$  if and only if

$$(28) \quad r_{13}r_{12} - r_{12}r_{23} + r_{23}r_{13} = 0$$

Substituting the summands from Eqs. (24) and (25) in Eq. (28) and gathering the coefficient of the tensor  $E_{ij} \otimes E_{kl} \otimes E_{st}$ , we obtain

$$(29) \quad \sum_p \left( t_{ij}^{kp} t_{pl}^{st} - t_{pj}^{kl} t_{ip}^{st} - t_{kl}^{sp} t_{ij}^{pt} \right) = 0.$$

Observe that Eq. (27) and Eq. (29) coincide up to the following permutation of indexes

$$\begin{pmatrix} a & b & c & d & l \\ j & l & l & s & t \end{pmatrix}.$$

So in this case, there is also a bijection  $\phi$  from the set of the solutions of AYBE of weight zero in  $M_n(\mathbf{k})$  to the set of Rota-Baxter operators of weight zero on  $M_n(\mathbf{k})$ .

Finally, the map  $\phi$  acts as

$$\phi \left( r = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} \otimes E_{kl} \right) = P \text{ such that } P(E_{pq}) = \sum_{i,l} t_{ip}^{ql} E_{il} = \sum_{i,j,k,l} t_{ij}^{kl} E_{ij} E_{pq} E_{kl} = P_r(E_{pq}).$$

So the linear operator  $P$  is exactly the linear operator  $P_r$  defined in Eq. (18). Therefore, the bijection  $\phi$  is precisely the map  $r \rightarrow P_r$ . This completes the proof.  $\square$

**Example 4.13.** Consider the matrix algebra  $M_2(\mathbb{C})$ . Aguiar [1, Example 5.4.5] showed that all nonzero solutions of AYBE of weight zero in  $M_2(\mathbb{C})$  are

$$r_1 = E_{12} \otimes E_{12}, r_2 = E_{22} \otimes E_{12}, r_3 = (E_{11} + E_{22}) \otimes E_{12} \text{ and } r_4 = E_{11} \otimes E_{12} - E_{12} \otimes E_{11},$$

up to conjugation, transpose and scalar multiple. By Theorem 4.12, all nonzero Rota-Baxter operators (viewed as matrices) of weight zero in  $M_2(\mathbb{C})$  up to conjugation, transpose and scalar multiple are the following:

- (a)  $P_{r_1}(E_{21}) = E_{12}, P_{r_1}(E_{11}) = P_{r_1}(E_{12}) = P_{r_1}(E_{22}) = 0;$
- (b)  $P_{r_2}(E_{21}) = E_{22}, P_{r_2}(E_{11}) = P_{r_2}(E_{12}) = P_{r_2}(E_{22}) = 0;$
- (c)  $P_{r_3}(E_{11}) = E_{12}, P_{r_3}(E_{21}) = E_{22}, P_{r_3}(E_{12}) = P_{r_3}(E_{22}) = 0;$
- (d)  $P_{r_4}(E_{11}) = E_{12}, P_{r_4}(E_{21}) = -E_{11}, P_{r_4}(E_{12}) = P_{r_4}(E_{22}) = 0.$

See also in [12, 33, 48].

## 5. WEIGHTED QUASITRIANGULAR $\epsilon$ -UNITARY BIALGEBRAS AND DENDRIFORM ALGEBRAS

In this section, we introduce the concept of weighted quasitriangular  $\epsilon$ -unitary bialgebras, which generalize the quasitriangular  $\epsilon$ -bialgebras initiated by Aguiar [1]. We show that any weighted quasitriangular  $\epsilon$ -unitary bialgebra can be made into a dendriform algebra.

### 5.1. Weighted quasitriangular $\epsilon$ -unitary bialgebras.

**Definition 5.1.** Let  $(A, m, 1)$  be a unitary algebra. A **quasitriangular infinitesimal unitary bialgebra** (abbreviated **quasitriangular  $\epsilon$ -unitary bialgebra**) of weight  $\lambda$  is a quadruple  $(A, m, 1, r)$  consisting of a unitary algebra  $(A, m, 1)$  and a solution  $r \in A \otimes A$  of an associative Yang-Baxter equation of weight  $\lambda$ .

Recall that  $\Delta_r$  is defined in Eq. (16) for a  $r \in A \otimes A$ .

**Remark 5.2.** (a) By Theorem 4.6, the quadruple  $(A, m, 1, \Delta_r)$  is indeed an  $\epsilon$ -unitary bialgebra of weight  $\lambda$ .

(b) A quasitriangular  $\epsilon$ -bialgebra studied in [1] is a quasitriangular  $\epsilon$ -unitary bialgebra of weight zero. In this case, the quadruple  $(A, m, 1, \Delta_r)$  is an  $\epsilon$ -unitary bialgebra of weight zero.

Strongly motivated by Aguiar [1], we record some properties of weighted quasitriangular  $\epsilon$ -unitary bialgebras.

**Proposition 5.3.** *Let  $(A, m, 1, r)$  be a quasitriangular  $\epsilon$ -unitary bialgebra of weight  $\lambda$  and  $\Delta := \Delta_r$ . Then*

$$(30) \quad \Delta(a) = a \cdot r - r \cdot a - \lambda(a \otimes 1),$$

$$(31) \quad (\Delta \otimes \text{id})(r) = -r_{23}r_{13}, \text{ and}$$

$$(32) \quad (\text{id} \otimes \Delta)(r) = r_{13}r_{12} + \lambda(r_{13} - r_{12}).$$

Conversely, if an  $\epsilon$ -unitary bialgebra  $(A, m, 1, \Delta)$  of weight  $\lambda$  satisfies Eqs. (30), (31) and (32) for some  $r = \sum_i u_i \otimes v_i \in A \otimes A$ , then  $(A, m, 1, r)$  is a quasitriangular  $\epsilon$ -unitary bialgebra of weight  $\lambda$  and  $\Delta = \Delta_r$ .

*Proof.* Eq. (30) follows directly from Eq. (16). Having Eq. (30) in hand, we have

$$\begin{aligned} (\Delta \otimes \text{id})(r) &= \sum_i \Delta(u_i) \otimes v_i = \sum_i (u_i \cdot r - r \cdot u_i - \lambda u_i \otimes 1) \otimes v_i \\ &= \sum_{i,j} (u_i u_j \otimes v_j \otimes v_i - u_j \otimes v_j u_i \otimes v_i) - \lambda \sum_i u_i \otimes 1 \otimes v_i \\ &= r_{13}r_{12} - r_{12}r_{23} - \lambda r_{13} = -r_{23}r_{13} \quad (\text{by Eq. (14)}). \end{aligned}$$

With a similar argument,

$$\begin{aligned} (\text{id} \otimes \Delta)(r) &= \sum_i u_i \otimes \Delta(v_i) = \sum_i u_i \otimes (v_i \cdot r - r \cdot v_i - \lambda v_i \otimes 1) \\ &= \sum_{i,j} (u_i \otimes v_i u_j \otimes v_j - u_i \otimes u_j \otimes v_j v_i) - \lambda \sum_i u_i \otimes v_i \otimes 1 \\ &= r_{12}r_{23} - r_{23}r_{13} - \lambda r_{12} = r_{13}r_{12} + \lambda(r_{13} - r_{12}) \quad (\text{by Eq. (14)}). \end{aligned}$$

Conversely, if an  $\epsilon$ -unitary bialgebra  $(A, m, 1, \Delta)$  of weight  $\lambda$  satisfies Eqs. (30), (31) and (32) for some  $r \in A \otimes A$ , then the same calculation shows that  $r$  is a solution of an AYBE of weight  $\lambda$ , and so  $(A, m, 1, r)$  is a quasitriangular  $\epsilon$ -bialgebra of weight  $\lambda$ . We note finally that  $A \otimes A$  is an  $A$ -bimodule by Eq. (1), then  $\Delta = \Delta_r$  follows from Eqs. (16) and (30).  $\square$

**5.2. Dendriform algebras from weighted quasitriangular  $\epsilon$ -unitary bialgebras.** In this subsection, we derive a dendriform algebra from a weighted quasitriangular  $\epsilon$ -bialgebra.

**Definition 5.4.** [42] A **dendriform algebra** is a  $\mathbf{k}$ -module  $D$  together with two binary operations  $<: D \otimes D \rightarrow D$  and  $>: D \otimes D \rightarrow D$  that satisfy the following relations:

$$\begin{aligned}(a < b) < c &= a < (b < c + b > c), \\ (a > b) < c &= a > (b < c), \\ a > (b > c) &= (a < b + a > b) > c \text{ for } a, b, c \in D.\end{aligned}$$

We record the following lemma as a preparation.

**Lemma 5.5.** [21, 23] Let  $(A, P)$  be a Rota-Baxter algebra of weight  $\lambda$ . Define two binary operations  $>, <$  on  $A$  by

$$a > b := P(a)b \text{ and } a < b := aP(b) + \lambda ab \text{ for } a, b \in A.$$

Then the triple  $(A, >, <)$  is a dendriform algebra.

The following result captures a relationship between weighted quasitriangular  $\epsilon$ -unitary bialgebras and dendriform algebras.

**Theorem 5.6.** Let  $(A, m, 1, r)$  be a quasitriangular  $\epsilon$ -unitary bialgebra of weight  $\lambda$  with  $r = \sum_i u_i \otimes v_i \in A \otimes A$ . Define two binary operations  $>, <$  on  $A$  by

$$a > b := \sum_i u_i a v_i b \text{ and } a < b := \sum_i a u_i b v_i - \lambda ab.$$

Then the triple  $(A, >, <)$  is a dendriform algebra.

*Proof.* The result follows from Theorem 4.10 and Lemma 5.5.  $\square$

**Acknowledgments:** This work was supported by the National Natural Science Foundation of China (Grant No. 11771191 and 11501267), Fundamental Research Funds for the Central Universities (Grant No. lzujbky-2017-162), the Natural Science Foundation of Gansu Province (Grant No. 17JR5RA175).

## REFERENCES

- [1] M. Aguiar, Infinitesimal Hopf algebras, Contemporary Mathematics, New trends in Hopf algebra theory (La Falda, 1999), 1-29, *Contemp. Math.*, **267**, Amer. Math. Soc., Providence, RI, (2000). [2](#), [3](#), [4](#), [5](#), [8](#), [9](#), [10](#), [14](#), [18](#), [20](#), [21](#)
- [2] M. Aguiar, Pre-Poisson algebras, *Lett. Math. Phys.* **54** (2000), 263-277. [2](#), [17](#)
- [3] M. Aguiar, On the associative analog of Lie bialgebras, *J. Algebra* **244** (2001), 492-532. [2](#), [3](#)
- [4] M. Aguiar, Infinitesimal Hopf algebras and the cd-index of polytopes. Geometric combinatorics (San Francisco, CA/Davis, CA, 2000), *Discrete Comput. Geom.* **27** (2002), 3-28. [2](#)
- [5] M. Aguiar, Infinitesimal bialgebras, pre-Lie and dendriform algebras. Hopf algebras, 1-33, *Lecture Notes in Pure and Appl. Math.* **237**, Dekker, New York, 2004. [2](#), [3](#), [10](#), [11](#)
- [6] H. H. An and C. M. Bai, From Rota-Baxter algebras to pre-Lie algebras, *J. Phys. A: Math. Theor.* **41** (2008), 01520. [2](#), [11](#)
- [7] C. M. Bai, Introduction to pre-Lie algebras, Preprint, available at <http://einspem.upm.edu.my/equals8/CSS/pre-Lie.pdf> [2](#), [11](#)
- [8] C. M. Bai, A unified algebraic approach to the classical Yang-Baxter equation, *J. Phy. A: Math. Theor.* **40** (2007), 11073-11082. [2](#)



- [9] C. M. Bai, Left-symmetric bialgebras and an analogue of the classical Yang-Baxter equation, *Comm. Contemp. Math.* **10** (2008), 221-260. [2](#)
- [10] C. M. Bai, L. Guo and X. Ni,  $\mathcal{O}$ -operators on associative algebras and associative Yang-Baxter equations, *Pacific J. Math.* **256** (2012), 257-289. [2](#), [3](#), [14](#), [17](#)
- [11] C. M. Bai, O. Bellier, L. Guo and X. Ni, Splitting of operations, Manin products, and Rota-Baxter operators, *Int. Math. Res. Not. IMRN* (2013), 485-524. [2](#)
- [12] P. Benito, V. Gubarev, A. Pozhidaev, Rota-Baxter operators on quadratic algebras, *Mediterr. J. Math.* (to appear), DOI: 10.1007/s00009-018-1234-5. [20](#)
- [13] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* **10** (1960), 731-742. [17](#)
- [14] M. Bordemann, Generalized Lax pairs, the modified classical Yang-Baxter equation, and affine geometry of Lie groups, *Comm. Math. Phys.* **135** (1990), 201-216. [2](#)
- [15] A. A. Balinskii and S. P. Novikov, Poisson brackets of hydrodynamic type, Frobenius algebras and Lie algebras, *Dokl. Akad. Nauk SSSR* **283** (1985), 1036-1039. [2](#)
- [16] A. Connes and D. Kreimer, Hopf algebras, renormalization and non-commutative geometry, *Comm. Math. Phys.* **199** (1998), 203-242. [2](#)
- [17] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* **210** (2000), 249-273. [2](#)
- [18] F. Chapoton and M. Livernet, Pre-Lie algebras and the rooted trees operad, *Int. Math. Res. Not. IMRN* **8** (2001), 395-408. [2](#)
- [19] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, Cambridge, 1994. [9](#)
- [20] V. G. Drinfeld, Quantum groups, Proceedings of the International Congress of Mathematicians, Berkeley, (1986), 798-820. [3](#)
- [21] K. Ebrahimi-Fard, Loday-type algebras and the Rota-Baxter relation, *Lett. math. phys.* **61** (2002), 139-147. [3](#), [14](#), [18](#), [22](#)
- [22] K. Ebrahimi-Fard, J. M. Gracia-Bondia and F. Patras, Rota-Baxter algebras and new combinatorial identities, *Lett. Math. Phys.* **81** (2007), 61-75. [2](#)
- [23] K. Ebrahimi-Fard and L. Guo, Rota-Baxter algebras and dendriform algebras. *J. Pure Appl. Algebra* **212** (2008), 320-339. [22](#)
- [24] K. Ebrahimi-Fard, L. Guo and D. Kreimer, Integrable renormalization II: The general case, *Ann. Henri Poincaré* **6** (2005), 369-395. [2](#)
- [25] P. Etingof and A. Soloviev, Quantization of geometric classical  $r$ -matrices, *Math. Res. Lett.* **6** (1999), 223-228. [2](#)
- [26] L. Foissy, Quantization of the Hopf algebras of decorated planar rooted trees, preprint, 2008. [6](#)
- [27] L. Foissy, The infinitesimal Hopf algebra and the poset of planar forests, *J. Algebraic Combin.* **30** (2009), 277-309. [2](#)
- [28] L. Foissy, The infinitesimal Hopf algebra and the operads of planar forests, *Int. Math. Res. Not. IMRN* **3** (2010), 395-435. [2](#)
- [29] X. Gao and L. Guo, Rota's Classification Problem, rewriting systems and Gröbner-Shirshov bases, *J. Algebra* **470** (2017), 219-253. [2](#)
- [30] X. Gao, L. Guo and M. Rosenkranz, Free integro-differential algebras and Gröbner-Shirshov bases, *J. Algebra* **442** (2015), 354-396. [2](#)
- [31] X. Gao and Y. Zhang, Planar rooted forests and weighted infinitesimal Hopf algebras: an operated algebra approach, submitted. [2](#), [5](#), [6](#), [9](#), [10](#), [12](#)
- [32] M. Gerstenhaber, The cohomology structure of an associative ring, *Ann. of Math. (2)* **78** (1963), 267-288. [2](#), [11](#)
- [33] V. Gubarev, Rota-Baxter operators on unital algebras, ArXiv:1805.00723 [18](#), [20](#)
- [34] L. Guo, An Introduction to Rota-Baxter Algebra, International Press, 2012. [17](#)
- [35] L. Guo, S. Paycha and B. Zhang, Algebraic Birkhoff factorization and the Euler-Maclaurin formula on cones, *Duke Math. J.* **166** (2017), 537-571. [2](#)
- [36] I. Z. Golubchik and V. V. Sokolov, Generalized operator Yang-Baxter equations, integrable ODEs and nonassociative algebras, *J. Nonlinear Math. Phys.* **7** (2000), 184-197. [2](#)

- [37] S. Joni and G.-C. Rota, Coalgebras and bialgebras in combinatorics, *Stud. Appl. Math.* **61** (1979), 93-139. [2](#), [5](#)
- [38] H. Kim, Complete left-invariant affine structures on nilpotent Lie groups, *J. Differ. Geom.* **24** (1986), 373-94. [2](#)
- [39] D. Kreimer, On the Hopf algebra structure of perturbative quantum field theories, *Adv. Theor. Math. Phys.* **2** (1998), 303-334. [2](#)
- [40] D. Kreimer, New mathematical structures in renormalizable quantum field theories, *Ann. Physics* **303** (2003), 179-202. [2](#)
- [41] T. Y. Lam, Lectures on modules and rings. Graduate Texts in Mathematics, Springer-Verlag, New York, 1999. [6](#)
- [42] J.-L. Loday, Dialgebras, in: Dialgebras and Related Operads, in: Lecture Notes in Math., Vol. 1763, Springer-Verlag, 2001, pp. 7-66. [22](#)
- [43] J.-L. Loday and M. O. Ronco, On the structure of cofree Hopf algebras, *J. Reine Angew. Math.* **592** (2006), 123-155. [2](#), [5](#), [6](#)
- [44] D. Manchon, A short survey on pre-Lie algebras. Noncommutative geometry and physics: renormalisation, motives, index theory, 89-102, *ESI Lect. Math. Phys.*, Eur. Math. Soc., Zürich, 2011. [11](#)
- [45] A. Medina, Flat left-invariant connections adapted to the automorphism structure of a Lie group, *J. Diff. Geom.* **16** (1981), 445-474 [2](#)
- [46] J. Pei, C. M. Bai and L. Guo, Splitting of operads and Rota-Baxter operators on operads, *Appl. Categ. Structures* **25** (2017), 505-538. [2](#)
- [47] A. Tanasa and D. Kreimer, Combinatorial Dyson-Schwinger equations in noncommutative field theory, *J. Noncommut. Geom.* **7** (2013), 255-289. [2](#)
- [48] X. Tang, Y. Zhang and Q. Sun, Rota-Baxter operators on 4-dimensional complex simple associative algebras, *Appl. Math. Comp.* **229** (2014), 173-186. [20](#)
- [49] E.B. Vinberg, The theory of homogeneous convex cones, *Transl. Moscow Math. Soc.* **2** (1963), 303-358. [2](#)
- [50] S. X. Wang and S. H. Wang, Drinfeld double for braided infinitesimal Hopf algebras, *Comm. Algebra* **42** (2014), 2195-2212. [2](#)

DEPARTMENT OF MATHEMATICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA  
*E-mail address:* zhangy2016@lzu.edu.cn

SCHOOL OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF APPLIED MATHEMATICS AND COMPLEX SYSTEMS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA  
*E-mail address:* gaoxing@lzu.edu.cn

DEPARTMENT OF MATHEMATICS, LANZHOU UNIVERSITY, LANZHOU, GANSU 730000, P. R. CHINA  
*E-mail address:* zhengjw16@lzu.edu.cn