

Attainable values for the Assouad dimension of projections

Jonathan M. Fraser & Antti Käenmäki

Abstract

We prove that for an arbitrary upper semi-continuous function $\phi: G(1, 2) \rightarrow [0, 1]$ there exists a compact set F in the plane such that $\dim_{\text{A}} \pi F = \phi(\pi)$ for all $\pi \in G(1, 2)$, where πF is the orthogonal projection of F onto the line π . In particular, this shows that the Assouad dimension of orthogonal projections can take on any finite or countable number of distinct values on a set of projections with positive measure. It was previously known that two distinct values could be achieved with positive measure. Recall that for other standard notions of dimension, such as the Hausdorff, packing, upper or lower box dimension, a single value occurs almost surely.

Mathematics Subject Classification 2010: primary: 28A80.

Key words and phrases: Assouad dimension, orthogonal projection

1 Dimensions of Projections

Starting with the seminal work of Marstrand [M54] from the 1950s, the behaviour of dimension under orthogonal projection has received a lot of attention in the fractal geometry and geometric measure theory literature. The general principle has been that the dimension of the orthogonal projection of a Borel set in \mathbb{R}^d onto a k -dimensional subspace almost surely does not depend on the specific choice of subspace. This has been verified for many different notions of dimension and in more general settings, see for example the surveys [FFJ15, M14] and the recent advance [KOV17]. The Marstrand-Mattila projection theorem for Hausdorff dimension states that, given a Borel set $F \subseteq \mathbb{R}^d$, we have

$$\dim_{\text{H}} \pi F = \min\{k, \dim_{\text{H}} F\}$$

for almost every $\pi \in G(k, d)$, where \dim_{H} denotes the Hausdorff dimension, and $G(k, d)$ is the Grassmanian manifold consisting of k -dimensional subspaces of \mathbb{R}^d which we identify with orthogonal projections π onto a given subspace. It was recently proved by Fraser and Orponen [FO17] in the planar case and Fraser [F18] in the general case that for any set $F \subseteq \mathbb{R}^d$, we have

$$\dim_{\text{A}} \pi F \geq \min\{k, \dim_{\text{A}} F\}$$

for almost every $\pi \in G(k, d)$, where \dim_{A} denotes the Assouad dimension, see below for the definition. The surprising additional result from [FO17] being that one cannot improve this result to an almost sure equality. Indeed, $\dim_{\text{A}} \pi F$ can take on two distinct values with positive measure. The purpose of this article is to push this observation further: *how wild can $\dim_{\text{A}} \pi F$ be as a function of π ?*

To finish the introduction, we recall the definition of the Assouad dimension. For any non-empty bounded set $E \subset \mathbb{R}^d$ and $r > 0$, let $N_r(E)$ be the smallest number of open sets with diameter less than or equal to r required to cover E . The *Assouad dimension* of a non-empty set $F \subseteq \mathbb{R}^d$ is defined by

$$\dim_{\text{A}} F = \inf \left\{ s \geq 0 : (\exists C > 0) (\forall R > 0) (\forall r \in (0, R)) \right. \\ \left. \sup_{x \in F} N_r(B(x, R) \cap F) \leq C \left(\frac{R}{r} \right)^s \right\}$$

where $B(x, R)$ denotes the closed ball centred at x with radius R . It is well-known that the Assouad dimension is always an upper bound for the Hausdorff dimension and upper box dimension; for example, see [L98].

2 Results

Our main result allows one to construct compact sets in the plane whose projections behave in a predefined manner with respect to the Assouad dimension.

Theorem 2.1. *Given an upper semi-continuous function $\phi: G(1, 2) \rightarrow [0, 1]$, there exists a compact set F in the plane with $\dim_{\text{A}} F = 0$ such that $\dim_{\text{A}} \pi F = \phi(\pi)$ for all $\pi \in G(1, 2)$.*

We remark that the assumption on upper semi-continuity is only needed to prove the upper bound $\dim_{\text{A}} \pi F \leq \phi(\pi)$ in the claim. In the context of Marstrand's projection theorem, that is, the approach of describing the dimensions of generic projections, we obtain the following corollaries, which are in stark contrast to what is possible for other standard notions of dimension, such as the Hausdorff, packing or box dimensions.

Corollary 2.2. *Let $E \subseteq [0, 1]$ be any finite or countable set. There exists a compact set F in the plane such that for all $s \in E$, we have $\dim_{\text{A}} \pi F = s$ for a set of $\pi \in G(1, 2)$ of positive measure.*

Proof. This follows immediately from Theorem 2.1 by letting $\{I_x\}_{x \in E}$ be a family of pairwise disjoint compact subsets of $G(1, 2)$ each with non-empty interior and then choosing ϕ to satisfy $\phi(\pi) = x$ for all $\pi \in I_x$ and $\phi(\pi) = 0$ otherwise. \square

This answers [FO17, Question 2.2] which asked how many distinct values the Assouad dimension of πF can assume for a set of π with positive measure. This corollary also answers [FO17, Question 2.7], which asked if there exists a compact planar set for which the Assouad dimension of the projection takes different values on two sets with non-empty interior. Finally, we can answer [FO17, Question 2.6] which asked if, given $0 \leq s < \log_5 3$, there exists a compact planar set with Hausdorff dimension s , for which the Assouad dimension of the projections is not almost surely constant. This was motivated by the fact that the examples constructed in [FO17] necessarily had Hausdorff dimension at least $\log_5 3$. The examples constructed in Theorem 2.1 provide such a set when $s = 0$,

which was the most difficult case, and to adapt this for $s > 0$ one can simply add to F a self-similar set of Hausdorff and Assouad dimension s which is contained in a line. The projections of this set in all but one direction will also have dimension s and therefore will only influence the Assouad dimensions of projections where $\dim_{\text{A}} \pi F < s$.

In addition to finding many values which appear as the Assouad dimension of projections with positive probability, we can also provide the following complementary result which *avoids* all values almost surely.

Corollary 2.3. *There exists a compact set F in the plane such that, for all $s \in [0, 1]$, there is at most one π such that $\dim_{\text{A}} \pi F = s$.*

Proof. This follows immediately from Theorem 2.1 by letting ϕ be given by $\phi(x) = x/\pi$ where we identify $G(1, 2)$ with $(0, \pi]$ in the natural way. \square

3 Proof of Theorem 2.1

We require the following technical lemma which we state in more generality than we need since it may be of interest in its own right and we could not find it in the literature. The result should be thought of as a separability property for functions, and we note that for continuous functions any choice of countable dense set Q will suffice.

Lemma 3.1. *Let X and Y be separable metric spaces. For any function $\phi: X \rightarrow Y$ there exists a countable dense subset Q of X such that, given $x \in X$, there exists a sequence $(x_k)_k$ of points in Q for which $x_k \rightarrow x$ and $\phi(x_k) \rightarrow \phi(x)$ as $k \rightarrow \infty$.*

Proof. Consider the set $\text{Graph}(\phi) := \{(x, \phi(x)) : x \in X\} \subseteq X \times Y$, where the product is equipped with the sup metric. Since X and Y are separable, so is $X \times Y$ and also $\text{Graph}(\phi)$. Therefore we may extract a countable dense subset $Q' \subseteq \text{Graph}(\phi)$. If $p_X: X \times Y \rightarrow X$ is the canonical projection, then the set $Q = p_X(Q')$ clearly satisfies the requirements of the lemma. \square

For the rest of the section, we assume that $\phi: G(1, 2) \rightarrow [0, 1]$ is an upper semi-continuous function. Let Π be a countable dense subset of $G(1, 2)$ satisfying the requirements of Lemma 3.1. For each $\pi \in \Pi$ and integer $n \geq 1$, we construct a finite set $F_{\pi, n}$ associated to π as follows. Let $s = \phi(\pi)$ and choose $c \in (0, 1/2]$ such that $\log 2 / \log(1/c) = s$ provided $s > 0$ and $c = 0$ otherwise. Let $S_0, S_1: [0, 1] \rightarrow [0, 1]$ be defined by $S_0(x) = cx$ and $S_1(x) = cx + 1 - c$ and let

$$E_n = \{S_{i_1} \circ \cdots \circ S_{i_n}(0) : i_j \in \{0, 1\}\} \subset [0, 1]$$

which is a finite approximation of the self-similar set generated by $\{S_0, S_1\}$. We assume a basic familiarity with self-similar sets; see for example [F15, Chapter 9]. Note that E_n consists of 2^n points, provided $c > 0$, and 2 points if $c = 0$. If $c > 0$, let

$$Z_n = \{2^{-i} : i = 1, 2, \dots, 2^n\}$$

and otherwise let $Z_n = \{0, 1\}$. Then let $F'_n \subseteq E_n \times Z_n$ be such that $\pi_1 F'_n = E_n$ and $\pi_2 F'_n = Z_n$, where π_i denotes projection onto the i th coordinate. There are clearly many

ways to choose F'_n and so for definiteness we assume that when F'_n is viewed as the graph of a function from E_n to $[0, 1]$ it is decreasing, which uniquely determines it. The idea here is that $\pi_1 F'_n = E_n$ is uniformly approximating a self-similar set of dimension s and $\pi_2 F'_n = Z_n$ is approximating the set $Z = \{0\} \cup \bigcup_n Z_n = \{0\} \cup \{2^{-i} : i = 1, 2, \dots\}$ which has Assouad dimension 0. We now want to force the projections of F'_n to also appear 0-dimensional in directions distinct from π_1 and π_2 , which is achieved by ‘stretching’, and to re-align F'_n such that the direction of the large projection is π instead of π_1 , which is achieved by rotating. Finally, we shall place the stretched and rotated versions of F'_n on the graph of $x \mapsto x^2$ in such a way that their projections (in any direction) do not overlap with each other too much. To this end, let $g: \Pi \times \mathbb{N} \rightarrow \mathbb{N}$ be a bijection and

$$F_{\pi,n} = R_\pi \circ A_{g(\pi,n)}(F'_n) + t_{g(\pi,n)} \quad (3.1)$$

where A_i is the linear map which scales by $h(i)$ in the horizontal direction and $v(i)$ in the vertical direction, R_π is the rotation about the origin so that the base of $R_\pi([0, 1]^2)$ is parallel with π , and $t_i = (2^{-i}, 4^{-i})$ is a translation. Here h and v are decreasing functions from \mathbb{N} to $(0, 1)$ chosen such that $h(i) < v(i) < 10^{-i}$ for all $i \geq 1$ and

$$\frac{h(i)}{v(i)} 2^{2i} \rightarrow 0 \quad \text{and} \quad \frac{v(i)}{10^{-i}} \rightarrow 0.$$

Finally, let

$$F = \{(0, 0)\} \cup \bigcup_{(\pi,n) \in \Pi \times \mathbb{N}} F_{\pi,n}$$

and notice that, by construction, the set F is clearly compact.

Before showing that the set F satisfies the properties claimed in Theorem 2.1, we provide the reader with some preliminaries.

Lemma 3.2. *For all $\varepsilon > 0$ there exists a constant $C_\varepsilon \geq 1$ such that for every $\pi \in \Pi$, $n \geq 1$, $x \in \mathbb{R}^2$, and $0 < r < R$ we have $N_r(B(x, R) \cap F_{\pi,n}) \leq C_\varepsilon (R/r)^\varepsilon$.*

Proof. Fix $\varepsilon > 0$, and let $C'_\varepsilon \geq 1$ be such that for all $z \in \mathbb{R}$ and $0 < r < R$ we have

$$N_r(B(z, R) \cap Z) \leq C'_\varepsilon (R/r)^\varepsilon. \quad (3.2)$$

This can be achieved since $\dim_A Z = 0$. Fix $\pi \in \Pi$ and $n \geq 1$, and let π^\perp denote the orthogonal complement of π . For distinct $x, y \in F_{\pi,n}$, note that

$$|\pi^\perp(x) - \pi^\perp(y)| \geq 2^{-2g(\pi,n)} v(g(\pi,n))$$

and

$$|\pi(x) - \pi(y)| \leq h(g(\pi,n)),$$

and therefore, since $2^{-2i} v(i)/h(i) \rightarrow \infty$, we observe that π^\perp is a bi-Lipschitz map on $F_{\pi,n}$ with Lipschitz constants which can be bounded away from 0 and ∞ independently of π and n . In fact, the Lipschitz constants approach 1 as $g(\pi,n) \rightarrow \infty$ or $n \rightarrow \infty$. Therefore there is a uniform constant $C_0 \geq 1$ such that for all $r > 0$ and any subset $F' \subseteq F_{\pi,n}$ we have $N_r(F') \leq C_0 N_r(\pi^\perp F')$. Note that $\pi^\perp F_{\pi,n}$ is a subset of Z scaled

down by a factor of $v(g(\pi, n))$. It follows that for arbitrary $x \in \mathbb{R}^2$ and $0 < r < R$ we have

$$\begin{aligned} N_r(B(x, R) \cap F_{\pi, n}) &\leq C_0 N_r(B(\pi^\perp(x), R) \cap \pi^\perp F_{\pi, n}) \\ &\leq C_0 \sup_{z \in \mathbb{R}} N_{r/v(g(\pi, n))}(B(z, R/v(g(\pi, n)))) \cap Z \\ &\leq C_0 C'_\varepsilon (R/r)^\varepsilon \quad \text{by (3.2)} \end{aligned}$$

which proves the claim. \square

Let $\mathcal{K}(\mathbb{R})$ denote the set of all compact subsets of \mathbb{R} , which is a complete metric space when equipped with the *Hausdorff metric* $d_{\mathcal{H}}$ defined by

$$d_{\mathcal{H}}(A, B) = \inf\{\delta : A \subseteq B_\delta \text{ and } B \subseteq A_\delta\}$$

where, for any $C \in \mathcal{K}(\mathbb{R})$,

$$C_\delta = \{x \in \mathbb{R} : |x - y| < \delta \text{ for some } y \in C\}$$

denotes the open δ -neighbourhood of C . We write $B(0, 1) \subset \mathbb{R}$ to denote the closed unit ball. One of the most effective ways to bound the Assouad dimension of a set from below is to use *weak tangents*; an approach going back to Mackay and Tyson [MT10] and Keith and Laakso [KL04]. Let $F \subset \mathbb{R}$ and $E \subseteq B(0, 1)$ be compact. If there exists a sequence of similarity maps T_k on \mathbb{R} such that $d_{\mathcal{H}}(E, T_k(F) \cap B(0, 1)) \rightarrow 0$ as $k \rightarrow \infty$, then E is called a *weak tangent* to F .

Proposition 3.3. *If $F \subseteq \mathbb{R}$ is compact and E is a weak tangent to F , then $\dim_{\mathbb{A}} F \geq \dim_{\mathbb{A}} E$.*

The proof of this proposition can be found in [MT10, Proposition 6.1.5]. We also have the following dual result which shows that the Assouad dimension can always be achieved as the Hausdorff dimension of a weak tangent.

Proposition 3.4. *If $F \subseteq \mathbb{R}$ is compact, then there exists a weak tangent E to F such that $\dim_{\mathbb{H}} E = \dim_{\mathbb{A}} E = \dim_{\mathbb{A}} F$.*

This statement follows from [F08, Theorem 5.1] and [KR16, Proposition 3.13], or, alternatively, directly from [KOR18, Proposition 5.7]. We are now ready to prove the main theorem.

Proof of Theorem 2.1. Let us first show that $\dim_{\mathbb{A}} F = 0$. Fix $x \in F$ and $0 < r < R \leq 1$, and let $m := \min\{g(\pi, n) : B(x, R) \cap F_{\pi, n} \neq \emptyset\}$. Note that we may assume $R \leq 1$ since F is bounded. Suppose first that $B(x, R)$ only intersects one of the sets $F_{\pi, n}$, which is therefore necessarily the one associated to m . It follows from Lemma 3.2 that

$$N_r(B(x, R) \cap F) \leq N_r(B(x, R) \cap F_{\pi, n}) \leq C_\varepsilon (R/r)^\varepsilon.$$

Now suppose that $B(x, R)$ intersects more than one of the sets $F_{\pi, n}$. In particular, it must intersect $F_{\pi, n}$ for some $g(\pi, n) > m$. This forces

$$2^{-m-3} \leq R \leq 2^{-m+2}$$

and therefore

$$\frac{-\log R}{\log 2} - 3 \leq m \leq \frac{-\log R}{\log 2} + 2. \quad (3.3)$$

We have

$$\begin{aligned} N_r(B(x, R) \cap F) &= N_r \left(B(x, R) \cap \bigcup_{g(\pi, n)=m}^{\infty} F_{\pi, n} \right) \\ &\leq N_r \left(B(x, R) \cap \bigcup_{g(\pi, n)=\lceil \frac{-\log r}{\log 2} \rceil}^{\infty} F_{\pi, n} \right) + \sum_{g(\pi, n)=m}^{\lfloor \frac{-\log r}{\log 2} \rfloor} N_r(B(x, R) \cap F_{\pi, n}) \\ &\leq 2 + \sum_{g(\pi, n)=m}^{\lfloor \frac{-\log r}{\log 2} \rfloor} C_{\varepsilon} \left(\frac{R}{r} \right)^{\varepsilon} \quad \text{by Lemma 3.2} \\ &\leq 2 + \left(\frac{\log R/r}{\log 2} + 4 \right) C_{\varepsilon} \left(\frac{R}{r} \right)^{\varepsilon} \quad \text{by (3.3)} \\ &\leq (6C_{\varepsilon} + \tilde{C}_{\varepsilon}) \left(\frac{R}{r} \right)^{2\varepsilon} \end{aligned}$$

where \tilde{C}_{ε} is a constant depending only on ε , such that $C_{\varepsilon} \log x / \log 2 \leq \tilde{C}_{\varepsilon} x^{\varepsilon}$ for all $x \geq 1$. We have proved that $\dim_{\Lambda} F \leq 2\varepsilon$ and since $\varepsilon > 0$ is arbitrary, we get $\dim_{\Lambda} F = 0$ as required.

We now prove that $\dim_{\Lambda} \pi F \geq \phi(\pi)$ for all $\pi \in G(1, 2)$. Fix $\pi \in G(1, 2)$ and let $s = \phi(\pi)$ which we may assume is strictly positive since otherwise there is nothing to prove. Let $c \in (0, 1/2]$ be such that $s = -\log 2 / \log c > 0$. Let $(\pi_k)_k$ be a sequence of points in Π given by Lemma 3.1 such that $\pi_k \rightarrow \pi$ and $\phi(\pi_k) \rightarrow \phi(\pi)$ as $k \rightarrow \infty$. Let $F_k := F_{\pi_k, k}$ denote the set consisting of 2^k points which is associated to π_k ; see (3.1) for the precise definition. Further, let $c_k \in (0, 1/2]$ be the constant associated with F_k such that $-\log 2 / \log c_k = \phi(\pi_k)$. As $\phi(\pi_k) \rightarrow \phi(\pi)$, it follows that $c_k \rightarrow c$ as $k \rightarrow \infty$.

Let $I = [0, 1]$ and for $\lambda \in (0, 1/2]$, let $E(\lambda) \subseteq I$ be the self-similar set associated with the maps $x \mapsto \lambda x$ and $x \mapsto \lambda x + (1 - \lambda)$. For each $k \geq 1$ let $T_k: \mathbb{R} \rightarrow \mathbb{R}$ be the similarity given by

$$T_k(x) = |\pi F_k|^{-1} (x - \inf \pi F_k).$$

Here we identify orthogonal projections of \mathbb{R}^2 and \mathbb{R} in the natural way. We claim that

$$T_k(\pi F_k) \cap I \rightarrow E(c) \quad (3.4)$$

in the Hausdorff metric as $k \rightarrow \infty$. Since the map $(\pi', A) \mapsto \pi' A \in \mathcal{K}(\mathbb{R})$ defined on $G(1, 2) \times \mathcal{K}(B(0, 2))$ is uniformly continuous by the Heine-Cantor Theorem, we see that

$$d_{\mathcal{H}}(T_k(\pi F_k) \cap I, T_k(\pi_k F_k) \cap I) \rightarrow 0$$

as $k \rightarrow \infty$. Moreover, since $T_k(\pi_k F_k) = E_k$ is a finite level k approximation of the self-similar set $E(c_k)$, we see that

$$d_{\mathcal{H}}(T_k(\pi_k F_k) \cap I, E(c_k)) \leq c_k^k \leq 2^{-k} \rightarrow 0$$

as $k \rightarrow \infty$. Moreover, since $c_k \rightarrow c$ we have

$$d_{\mathcal{H}}(E(c_k), E(c)) \rightarrow 0$$

as $k \rightarrow \infty$. It therefore follows, by applying the triangle inequality twice, that

$$d_{\mathcal{H}}(T_k(\pi F_k) \cap I, E(c)) \rightarrow 0$$

as $k \rightarrow \infty$, proving (3.4).

Since $\mathcal{K}(I)$ is compact we may extract a subsequence of the sequence $T_k(\pi F) \cap I$ which converges to a set $\widehat{\pi F}$ which is necessarily a weak tangent to πF . Since $T_k(\pi F_k) \cap I \subseteq T_k(\pi F) \cap I$ for all k by the definition of F and $T_k(\pi F_k) \cap I \rightarrow E(c)$ as $k \rightarrow \infty$ by (3.4), it follows that $E(c) \subseteq \widehat{\pi F}$. Therefore Proposition 3.3 yields

$$\dim_{\mathbb{A}} \pi F \geq \dim_{\mathbb{A}} \widehat{\pi F} \geq \dim_{\mathbb{A}} E(c) = s$$

as required.

Finally, we prove that $\dim_{\mathbb{A}} \pi F \leq \phi(\pi)$ for all $\pi \in G(1, 2)$. For a given $\pi \in G(1, 2)$ (which we fix from now on) we first argue that the projections $\pi F_{\pi', n}$ ($\pi' \in \Pi$, $n \geq 1$) are exponentially separated. First suppose that π does not correspond to the vertical projection. Since

$$\frac{|F_{\pi', n}|}{2^{-g(\pi', n)}} \leq \frac{\sqrt{2}v(g(\pi', n))}{2^{-g(\pi', n)}} \rightarrow 0$$

we have that the sets $\pi(F_{\pi', n})$ are pairwise disjoint for large enough $g(\pi', n)$ (or n), and even separated by a distance of at least a constant times $2^{-g(\pi', n)}$. More precisely, there exists a constant $K \geq 1$ such that

$$\begin{aligned} \inf\{|x - y| : x \in \pi F_{\pi'', m}, y \in \pi F_{\pi', n}, \text{ and } g(\pi'', m) > g(\pi', n) \geq K\} \\ \geq 2^{-1} |\cos(\theta)| 2^{-g(\pi', n)} \end{aligned}$$

where θ is the angle π makes with the horizontal axis. Note that $\cos(\theta) \neq 0$ since π is not the vertical projection. The angle π makes with the horizontal axis is relevant because the translations t_i become tangent to the horizontal axis since they lie on the graph of $x \mapsto x^2$. Secondly, suppose that π is the vertical projection. Similar to above, since

$$\frac{|F_{\pi', n}|}{4^{-g(\pi', n)}} \leq \frac{\sqrt{2}v(g(\pi', n))}{4^{-g(\pi', n)}} \rightarrow 0$$

we have that the sets $\pi F_{\pi', n}$ are pairwise disjoint for large enough $g(\pi', n)$ (or n), and even separated by a distance of at least a constant times $4^{-g(\pi', n)}$. We are forced to replace $2^{-g(\pi', n)}$ with $4^{-g(\pi', n)}$ here because the vertical projection of the set of translations t_i is the sequence 4^{-i} , rather than (asymptotic to) $\cos(\theta)2^{-i}$. Since each of the sets $F_{\pi', n}$ are finite and the Assouad dimension is finitely stable, we may assume without loss of generality that these separation properties holds for all $F_{\pi', n}$. That is, we may assume that for all $(\pi', n) \neq (\pi'', m)$ with $g(\pi'', m) > g(\pi', n)$ we have

$$\inf\{|x - y| : x \in \pi F_{\pi'', m}, y \in \pi F_{\pi', n}\} \geq C(\pi) 4^{-g(\pi', n)} \quad (3.5)$$

for a constant $C(\pi)$ depending only on π .

Suppose E is a weak tangent to πF . By Proposition 3.4, it suffices to show that $\dim_{\mathbb{H}} E \leq \phi(\pi)$. Since E is a weak tangent, there exists a sequence of similarity maps T_l on \mathbb{R} such that $T_l(\pi F) \cap B(0, 1) \rightarrow E$ in the Hausdorff metric as $l \rightarrow \infty$. We may assume that there are arbitrarily large k such that $T_l(\pi F_{\pi', k}) \cap B(0, 1) \neq \emptyset$ for some l and some $\pi' \in \Pi$. Otherwise E is a weak tangent of a finite set and would therefore be finite itself. We may also assume that $\|T_l\| \rightarrow \infty$ since otherwise E is countable. Therefore, by taking a subsequence of $(T_l)_l$ we may assume that

$$\min\{k : T_l(\pi F_{\pi', k}) \cap B(0, 1) \neq \emptyset \text{ for some } \pi'\} \rightarrow \infty \quad (3.6)$$

as $l \rightarrow \infty$.

For a given $l \geq 1$, suppose for some large $k \geq 1$ we have $T_l(\pi F_{\pi', k}) \cap B(0, 1) \neq \emptyset$ for some π' and $|T_l(\pi F_{\pi', k})| \geq 2^{-g(\pi', k)}$. It follows that

$$\|T_l\| \sqrt{2} v(g(\pi', k)) \geq \|T_l\| |\pi F_{\pi', k}| = |T_l(\pi F_{\pi', k})| \geq 2^{-g(\pi', k)}$$

and therefore

$$\|T_l\|^{-1} \leq 2^{g(\pi', k)} v(g(\pi', k)) \sqrt{2} < 5^{-g(\pi', k)} \sqrt{2}.$$

It follows from the separation property (3.5) that $T_l(\pi F_{\pi'', i}) \cap B(0, 1) = \emptyset$ for all $(\pi'', i) \neq (\pi', k)$, provided that k is sufficiently large, depending on π .

On the other hand, suppose that for all large enough l we have $|T_l(\pi F_{\pi', k})| < 2^{-g(\pi', k)}$ for all k and π' such that $T_l(\pi F_{\pi', k}) \cap B(0, 1) \neq \emptyset$. This means that the contribution from individual $\pi(F_{\pi', k})$ vanishes to a single point in the limit and E must be a weak tangent of the projection of the closure of the set of translations, $\{\pi(t_i)\}_i \cup \{0\}$, and therefore has Hausdorff dimension 0. Therefore we may assume (by taking a subsequence if necessary) that for all l there exists a unique pair $(\pi(l), k(l))$ such that $T_l(\pi F_{\pi(l), k(l)}) \cap B(0, 1) \neq \emptyset$ and, moreover, that $|T_l(\pi F_{\pi(l), k(l)})| \geq 2^{-g(\pi(l), k(l))}$. Therefore,

$$T_l(\pi F_{\pi(l), k(l)}) \cap B(0, 1) \rightarrow E. \quad (3.7)$$

Using compactness of $G(1, 2)$, we may assume that $\pi(l) \rightarrow \pi^* \in G(1, 2)$ as $l \rightarrow \infty$. First suppose that $\pi^* \neq \pi$. It follows that E is a weak tangent of $Z = \{0\} \cup \{2^{-i} : i = 1, 2, \dots\}$ and therefore has dimension 0. On the other hand, suppose $\pi^* = \pi$. This case is more delicate and is interestingly the only point in the proof where we use upper semi-continuity of ϕ . Consider $F_l := F_{\pi(l), k(l)}$ which, for large l , is contained in a very long and thin rectangle with long side almost orthogonal to π . Let

$$E_l' = R_{\pi(l)} \circ A_{g(\pi(l), k(l))}(E_{k(l)} \times \{0\}) + t_{g(\pi(l), k(l))}$$

which, by recalling (3.1), is simply the projection of F_l onto the base of the rectangle containing it. Similarly, let

$$Z_l' = R_{\pi(l)} \circ A_{g(\pi(l), k(l))}(\{0\} \times Z_{k(l)}) + t_{g(\pi(l), k(l))}$$

which is simply the projection of F_l onto the long side of the rectangle containing it. Order the points in F_l by writing $F_l = \{x_1, x_2, \dots, x_{2k(l)}\}$ in the natural ‘from top to

bottom' order. For each $x_n \in F_l$ label the corresponding point in E'_l by e_n and the corresponding point in Z'_l by z_n . Assume without loss of generality that $k(l)$ is even and group the points in F_l into $2^{k(l)/2}$ groups of size $2^{k(l)/2}$ in increasing manner, that is,

$$F_l = \bigcup_{n=1}^{2^{k(l)/2}} \{x_i : (n-1)2^{k(l)/2} + 1 \leq i \leq n2^{k(l)/2}\}.$$

The corresponding grouping of the points in E'_l arranges the points into the level $k(l)/2$ cylinders of the self-similar set which $E_{k(l)}$ is approximating. We may assume without loss of generality that $T_l(\pi F_l) \cap B(0, 1) = \{\pi(x_i) : (n-1)2^{k(l)/2} + 1 \leq i \leq n2^{k(l)/2}\}$ for some n which we fix from now on and that $\|T_l\|h(k(l)) \leq |B(0, 1)| = 2$. The easiest case to keep in mind is if $n = 1$ which corresponds to all of F_l contributing to the tangent and not just part of it.

Observe that for $(n-1)2^{k(l)/2} + 1 \leq i \leq n2^{k(l)/2}$ we have

$$|T_l(\pi(x_i)) - T_l(\pi(z_i) + t(l))| \leq \|T_l\|h(k(l))2^{-k(l)/2} \leq 2^{-k(l)/2+1} \quad (3.8)$$

where $t(l)$ is (asymptotic to) the translation associated with the similarity corresponding to the common level $k(l)/2$ cylinder in the construction of E'_l which the corresponding points e_i all belong to. In the easiest case where $n = 1$, $t(l) = 0$.

Similarly, for $n2^{k(l)/2} + 1 \leq i \leq 2^{k(l)}$ we have

$$|T_l(\pi(x_i)) - T_l(\pi(e_i))| \leq 2^{-2^{k(l)/2}+1}. \quad (3.9)$$

This is because the distance between $\pi(x_i)$ and $\pi(e_i)$ halves every time i increases by 1 and $|T_l(\pi(x_M)) - T_l(\pi(e_M))| \leq 2$, where $M = (n-1)2^{k(l)/2} + 1$ since $T_l(\pi(x_M)), T_l(\pi(e_M)) \in B(0, 1)$. Using (3.8), (3.9), and (3.7), it follows that

$$T_l(\pi E'_l \cup (\pi Z'_l + t(l))) \cap B(0, 1) \rightarrow E.$$

However, $T_l(\pi E'_l) \rightarrow \eta_E E' + t_E$ where $\eta_E \geq 0$ is a constant, t_E is a translation and E' is a self-similar set of dimension at most $\limsup_{l \rightarrow \infty} \phi(\pi(l)) \leq \phi(\pi)$, using upper semi-continuity of ϕ . Moreover, $T_l(\pi Z'_l + t(l)) \rightarrow \eta_Z Z + t_Z$ where $\eta_Z \geq 0$ is a constant, t_Z is a translation, and $Z = \{0, 2^{-1}, 2^{-2}, \dots\}$. It follows that $\dim_{\mathbb{H}} E \leq \phi(\pi)$ as required. \square

4 Final remarks

We proved that for any upper semi-continuous function $\phi: G(1, 2) \rightarrow [0, 1]$ there exists a compact set $F \subset \mathbb{R}^2$ such that $\dim_{\mathbb{A}} \pi F = \phi(\pi)$ for all but at most one $\pi \in G(1, 2)$. Notice, however, that the function $\pi \mapsto \dim_{\mathbb{A}} \pi F$ need not be upper semi-continuous in general since a line segment satisfies $\dim_{\mathbb{A}} \pi F = 1$ for all but one direction where the dimension is 0. It is therefore natural to ask what sort of functions can be realised as $\pi \mapsto \dim_{\mathbb{A}} \pi F$. We observe that the set

$$\{\phi: G(1, 2) \rightarrow [0, 1] : \phi(\pi) = \dim_{\mathbb{A}} \pi F \text{ for some compact } F \subset \mathbb{R}^2\}$$

has cardinality \aleph_1 : Theorem 2.1 implies that it is at least \aleph_1 but it cannot be more than \aleph_1 since there is a natural surjection from the set of compact subsets of \mathbb{R}^2 onto this set.

In particular, there must be functions which *cannot* be realised as $\pi \mapsto \dim_{\mathbb{A}} \pi F$ for a compact set F .

Finally, we remark that it should be possible to generalise our proof of Theorem 2.1 to higher dimensions for an upper semi-continuous function $\phi: G(k, d) \rightarrow [0, k]$. We do not pursue the details.

Acknowledgements

JMF was financially supported by a *Leverhulme Trust Research Fellowship* (RF-2016-500) and an *EPSRC Standard Grant* (EP/R015104/1). AK was financially supported by the *Finnish Center of Excellence in Analysis and Dynamics Research* and the aforementioned EPSRC grant, which covered the local costs of the visit of AK to the University of St Andrews in May 2018 where this research began. The authors thank Eino Rossi for pointing out a simple improvement which got rid of a minor additional assumption used in an earlier version of the paper. The authors also thank Tuomas Orponen for many helpful discussions on the topic.

References

- [F15] K. J. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*, John Wiley & Sons, Hoboken, NJ, 3rd. ed., 2014.
- [FFJ15] K. J. Falconer, J. M. Fraser and X. Jin. Sixty Years of Fractal Projections, *Fractal geometry and stochastics V*, (Eds. C. Bandt, K. J. Falconer and M. Zähle), Birkhäuser, Progress in Probability, 2015.
- [F18] J. M. Fraser. *Distance sets, orthogonal projections, and passing to weak tangents*, Israel J. Math., **226**, (2018), 851–875.
- [FO17] J. M. Fraser and T. Orponen. *The Assouad dimensions of projections of planar sets*, Proc. Lond. Math. Soc., **114**, (2017), 374–398.
- [F08] H. Furstenberg. *Ergodic fractal measures and dimension conservation*, Ergodic Theory Dynam. Systems, **28**, (2008), 405–422.
- [KOV17] A. Käenmäki, T. Orponen and L. Venieri. *A Marstrand-type restricted projection theorem in \mathbb{R}^3* , preprint, available at arXiv:1708.04859.
- [KOR18] A. Käenmäki, T. Ojala, and E. Rossi. *Rigidity of quasisymmetric mappings on self-affine carpets*, Int. Math. Res. Not. IMRN, **12**, (2018), 3769–3799.
- [KR16] A. Käenmäki and E. Rossi. *Weak separation condition, Assouad dimension, and Furstenberg homogeneity*, Ann. Acad. Sci. Fenn. Math., **41**, (2016), 465–490.
- [KL04] S. Keith and T. Laakso. *Conformal Assouad dimension and modulus*, Geom. Funct. Anal., **14**, (2004), 1278–1321.
- [L98] J. Luukkainen. *Assouad dimension: antifractal metrization, porous sets, and homogeneous measures*, J. Korean Math. Soc., **35**, (1998), 23–76.
- [MT10] J. M. Mackay and J. T. Tyson. Conformal dimension. Theory and application, University Lecture Series, 54. *American Mathematical Society, Providence, RI*, 2010.
- [M54] J. M. Marstrand. *Some fundamental geometrical properties of plane sets of fractional dimensions*, Proc. London Math. Soc.(3), **4**, (1954), 257–302.

- [M14] P. Mattila. Recent progress on dimensions of projections, in *Geometry and Analysis of Fractals*, D.-J. Feng and K.-S. Lau (eds.), pp 283–301, *Springer Proceedings in Mathematics & Statistics*. **88**, Springer-Verlag, Berlin Heidelberg, 2014.

Jonathan M. Fraser
School of Mathematics and Statistics
The University of St Andrews
St Andrews, KY16 9SS, Scotland

Email: jmf32@st-andrews.ac.uk

Antti Käenmäki
Department of Physics and Mathematics
University of Eastern Finland
P.O. Box 111, FI-80101 Joensuu, Finland

Email: antti.kaenmaki@uef.fi