

# PROOF OF A RATIONAL RAMANUJAN-TYPE SERIES FOR $1/\pi$ . THE FASTEST ONE IN LEVEL 3

JESÚS GUILLERA

*To Bruce Berndt, in admiration of his inspirational work on Ramanujan's Notebooks*

ABSTRACT. Using a modular equation of level 3 and degree 23 due to Chan and Liaw, we prove the fastest convergent rational Ramanujan-type series for  $1/\pi$  of level 3.

## 1. THE METHOD

A prerequisite to understand this paper well is familiarity with the notation and the method developed in [9]. This method is based on an original idea of Wan [10], whose paper was in turn influenced by some ideas of [8]. Using the notation

$$F_s(\alpha) = {}_2F_1\left(\frac{1}{s}, 1 - \frac{1}{s} \mid \alpha\right), \quad s \in \{6, 4, 3, 2\}, \quad \ell = 4 \sin^2 \frac{\pi}{s},$$

we proved in [9] the following result:

**Theorem 1.** *Let*

$$F_s(\alpha) = m(\alpha, \beta)F_s(\beta), \quad A(\alpha, \beta) = 0,$$

*be a transformation of modular origin and degree  $1/d$ , and let  $\beta_0 = 1 - \alpha_0$  be a solution of  $A(\alpha_0, \beta_0) = 0$ , and  $m_0 = m(\alpha_0, \beta_0)$ . Then, if*

$$m_0 = \frac{1}{\sqrt{d}}, \quad \text{or} \quad m_0 = \frac{\sqrt{4d - \ell}}{2d} + \frac{\sqrt{\ell}}{2d} i,$$

*we have*

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{s}\right)_n \left(1 - \frac{1}{s}\right)_n}{(1)_n^3} (a + bn) z^n = \frac{1}{\pi},$$

*where*

$$(1) \quad z = 4\alpha_0\beta_0, \quad b = 2(1 - 2\alpha_0) \sqrt{\frac{d}{\ell}}, \quad a = -2\alpha_0\beta_0 \frac{m'_0}{\alpha'_0} \frac{d}{\sqrt{\ell}}.$$

*or*

$$(2) \quad z = 4\alpha_0\beta_0, \quad b = 2(1 - 2\alpha_0) \sqrt{\frac{d}{\ell} - \frac{1}{4}}, \quad a = -2\alpha_0\beta_0 \frac{m'_0}{\alpha'_0} \frac{d}{\sqrt{\ell}},$$

*respectively. The ' means differentiation with respect to the variable that we choose as independent.*

2. THE FORMULA  $(A, 3, 23)$ 

In this paper we prove the Ramanujan-type formula

$$(3) \quad \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n^3} (14151n + 827) \frac{(-1)^n}{500^{2n}} = \frac{1500\sqrt{3}}{\pi}.$$

It is the formula  $(A, 3, 23)$  in the notation of [9], we will also refer to it as 3A23, and is the fastest convergent rational Ramanujan-type series for  $1/\pi$  of level  $\ell = 3$ . It was discovered by Chan, Liaw and Tan [5, eq. 1.19]. As  $z_0 = 4\alpha_0\beta_0$  where  $\beta_0 = 1 - \alpha_0$ , we get

$$\alpha_0 = \frac{1}{2} - \frac{53\sqrt{89}}{1000}, \quad \beta_0 = 1 - \alpha_0 = \frac{1}{2} + \frac{53\sqrt{89}}{1000},$$

Then, with a numerical approximation of 20 digits, we have

$$m_0 = \frac{F_3(\alpha_0)}{F_3(\beta_0)} \approx 0.20508654634905660459 + 0.037653278425410375946 i,$$

which we identify as

$$m_0 = \frac{\sqrt{89}}{46} + \frac{\sqrt{3}}{46} i.$$

Observe that  $m_0$  is of the form

$$m_0 = \frac{\sqrt{4d - \ell}}{2d} + \frac{\sqrt{\ell}}{2d} i, \quad |m_0| = \frac{1}{\sqrt{d}},$$

with  $d = 23$ . Hence, for proving (3) with our method, we need a transformation of degree  $1/d$  with  $d = 23$  for the level  $\ell = 3$ , and with such a transformation we can prove it rigorously. This is done in next section using a modular equation in which the algebraic relation  $A(\alpha, \beta) = 0$  is written using two auxiliary variables  $u$  and  $v$ , in the following way:

$$(4) \quad u^k = \alpha\beta, \quad v^k = (1 - \alpha)(1 - \beta), \quad P(u, v) = 0,$$

where  $P(u, v)$  is a polynomial in  $u$  and  $v$ , and  $k$  is a positive integer.

3. THE PROOF OF  $(A, 3, 23)$ 

We use a modular equation of level 3 and degree 23 due to Heng Huat Chan and Wen-Chin Liaw [4, Corollary 3.7], in which we have replaced  $u$  and  $v$  with  $u^2$  and  $v^2$  respectively. It is

$$(5) \quad u^{12} = \alpha\beta, \quad v^{12} = (1 - \alpha)(1 - \beta), \quad P(u, v) = 0,$$

where

$$(6) \quad P(u, v) = (u^8 + v^8) - 12\sqrt{3}(u^7v + uv^7) - 87(u^6v^2 + u^2v^6) - 84\sqrt{3}(u^5v^3 + u^3v^5) \\ - 160(u^4v^4) - 2(u^4 + v^4) - 15\sqrt{3}(u^3v + uv^3) - 48(u^2v^2) + 1.$$

If we let  $\beta = 1 - \alpha$ , then we see that  $u^{12} = v^{12}$ . If we choose

$$(7) \quad v = \left( \frac{\sqrt{3}}{2} - \frac{1}{2}i \right) u,$$

and replace it in  $P(u, v) = 0$ , we have the equation

$$(8) \quad 250(1 + \sqrt{3}i)u^8 - \frac{95}{2}(1 - \sqrt{3}i)u^4 + 1 = 0,$$

which factors as

$$\frac{1 + \sqrt{3}i}{16}(10u^4 + 1 + \sqrt{3}i)(20u^2 - \sqrt{3} - i)(20u^2 + \sqrt{3} + i) = 0.$$

One solution is

$$u_0 = \frac{\sqrt{15} + \sqrt{5}}{20} + \frac{\sqrt{15} - \sqrt{5}}{20}i, \quad v_0 = \frac{\sqrt{15} + \sqrt{5}}{20} - \frac{\sqrt{15} - \sqrt{5}}{20}i.$$

Then, from

$$\alpha_0(1 - \alpha_0) = u_0^{12} = -10^{-6}, \quad \beta_0 = 1 - \alpha_0,$$

we get

$$\alpha_0 = \frac{1}{2} - \frac{53}{1000}\sqrt{89}, \quad \beta_0 = \frac{1}{2} + \frac{53}{1000}\sqrt{89}.$$

Differentiating  $P(u, v) = 0$  with respect to  $u$  at  $u = u_0$ , we find

$$v'_0 = \frac{-294573\sqrt{3} + 82573i}{516854}.$$

Then, differentiating  $P(u, v) = 0$  twice with respect to  $u$  at  $u = u_0$ , we get

$$v''_0 = \frac{8674041040500000\sqrt{5}(1 - i) + 3034180783431000\sqrt{15}(1 + i)}{17258921684500483}.$$

From (5), we see that

$$(9) \quad u^{12} = \alpha\beta, \quad u^{12} - v^{12} + 1 = \alpha + \beta.$$

Differentiating (9) with respect to  $u$  at  $u = u_0$ , we obtain  $\alpha'_0$  and  $\beta'_0$ . Then, differentiating (9) twice with respect to  $u$  at  $u = u_0$  we obtain  $\alpha''_0$  and  $\beta''_0$ . As the multiplier is given by

$$(10) \quad m^2 = \frac{1}{d} \frac{\beta(1 - \beta)}{\alpha(1 - \alpha)} \frac{\alpha'}{\beta'},$$

see [9], replacing the already known values at  $u = u_0$ , we get

$$(11) \quad m_0 = \sqrt{\frac{1}{23} \frac{\alpha'_0}{\beta'_0}} = \frac{\sqrt{89}}{46} + \frac{\sqrt{3}}{46}i = \frac{\sqrt{4d - \ell}}{2d} + \frac{\sqrt{\ell}}{2d}i.$$

Taking logarithms in (10), differentiating with respect to  $u$ , and dividing by  $\alpha'$ , we get

$$\frac{m'}{\alpha'} = \frac{m}{2\alpha'} \left( \frac{\beta'}{\beta} - \frac{\beta'}{1 - \beta} - \frac{\alpha'}{\alpha} + \frac{\alpha'}{1 - \alpha} + \frac{\alpha''}{\alpha'} - \frac{\beta''}{\beta'} \right),$$

and from it, we obtain

$$\frac{m'_0}{\alpha'_0} = \frac{827000}{69}.$$

Finally, using the formulas

$$(12) \quad z = 4\alpha_0\beta_0, \quad b = 2(1 - 2\alpha_0)\sqrt{\frac{d}{\ell} - \frac{1}{4}}, \quad a = -2\alpha_0\beta_0\frac{m'_0}{\alpha'_0}\frac{d}{\sqrt{\ell}},$$

we obtain

$$z = \frac{-1}{500^2}, \quad b = \frac{4717}{1500}\sqrt{3}, \quad a = \frac{827}{4500}\sqrt{3},$$

and we are done.

#### 4. RAMANUJAN-TYPE SERIES FOR $1/\pi$ AND MODULAR EQUATIONS

In a completely similar way we can prove other Ramanujan-type series for  $1/\pi$  using modular equations [1, 2, 3, 4], written in the form (4). For example, for proving (A, 2, 7), we can use the modular equation

$$u^8 = \alpha\beta, \quad v^8 = (1 - \alpha)(1 - \beta), \quad P(u, v) = 0,$$

where

$$P(u, v) = (u^4 + v^4) + 8\sqrt{2}(u^3v + uv^3) + 20u^2v^2 - 1,$$

see [3, 2, Theorem 10.3]. For proving (A, 3, 11), we can use

$$u^{12} = \alpha\beta, \quad v^{12} = (1 - \alpha)(1 - \beta), \quad P(u, v) = 0,$$

where

$$P(u, v) = (u^4 + v^4) + 3\sqrt{3}(u^3v + uv^3) + 6u^2v^2 - 1.$$

see [3, 2, Theorem 7.8], and we can get the proofs of the formulas (A, 3, 5) and (P, 3, 5), with the modular equation

$$u^6 = \alpha\beta, \quad v^6 = (1 - \alpha)(1 - \beta), \quad u^2 + v^2 + 3uv - 1 = 0.$$

see [3, 2, Theorem 7.6]. Note that all the above examples correspond to rational series. Of course one can use the same method to prove irrational instances.

#### 5. CONCLUSION

As we pointed out in [9], the main aspect of our method is that the modular equations used to prove the Ramanujan-type alternating series for  $1/\pi$  have a much lower degree than those in other methods. It would be interesting to analyze in depth this phenomenon. The reader is invited to compare the values of  $d$  in the tables in [9] with the respective values of  $N$  given in the tables in [7]. We hope that the method developed here will attract others to find, for example, new modular equations of levels  $\ell \geq 5$  and apply them to prove non-hypergeometric series of Ramanujan-Sato type for  $1/\pi$  [6, 7]. Of course a generalization of the method would be needed for that.

## REFERENCES

- [1] B.C. BERNDT, *Ramanujan's Notebooks, Part III* (Springer-Verlag, New York, 1991).
- [2] B.C. BERNDT, *Ramanujan's Notebooks, Part V* (Springer-Verlag, New York, 1998).
- [3] B.C. BERNDT, S. BHARGAVA AND F.G. GARVAN *Ramanujan's theories of elliptic functions to alternative bases*, Trans. Amer. Soc. **347**, 4163–4244 (1995).
- [4] H.H. CHAN AND W.-C. LIAW, *On Russell-Type Modular Equations*, Canad. J. Math. **52**, 31–36 (2000).
- [5] H.H. CHAN, W.-C. LIAW AND V. TAN, *Ramanujan's class invariant  $\lambda_n$  and a new class of series for  $1/\pi$* , J. London Math Soc. (2) **64** (2001), 93–106.
- [6] H.H. CHAN, S.H. CHAN AND ZHIGUO LIU, *Domb's numbers and Ramanujan–Sato type series for  $1/\pi$* , Adv. Math. **186**, 396–410, (2004).
- [7] H.H. CHAN AND S. COOPER, *Rational analogues of Ramanujan's series for  $1/\pi$* , Math. Proc. Camb. Phil. Soc., 1–23, (2012).
- [8] H.H. CHAN, J. WAN AND W. ZUDILIN, *Legendre polynomials and Ramanujan-type series for  $1/\pi$* , Israel J. Math. **194**, 183–207, (2013).
- [9] J. GUILLERA, *A method for proving Ramanujan's series for  $1/\pi$* , Ramanujan J., (to appear).
- [10] J. WAN, *Series for  $1/\pi$  Using Legendre's Relation*, Integr. Transforms Spec. Funct. **25** (2014), 1–14.

UNIVERSITY OF ZARAGOZA, DEPARTMENT OF MATHEMATICS, 50009 ZARAGOZA (SPAIN)  
E-mail address: jguillera@gmail.com