

# Percolation on the product graph of a regular tree and a line does not satisfy the triangle condition at the uniqueness threshold

Kohei Yamamoto

Tohoku university

## Abstract

We consider Bernoulli bond percolation on the product graph of a regular tree and a line. We show that the triangle condition does not hold at the uniqueness threshold.

## 1 Introduction

Let  $G = (V, E)$  be a connected, quasi-transitive and infinite graph, where  $V$  is the set of vertices,  $E$  is the set of edges. In Bernoulli bond percolation, each edge will be open with probability  $p$ , and closed with probability  $1 - p$  independently, where  $p \in [0, 1]$  is a fixed parameter. Let  $\Omega = \{0, 1\}^E$  be the set of samples, where  $\omega(e) = 1$  means  $e$  is open. Each  $\omega \in \Omega$  is regarded as a subgraph of  $G$  consisting of all open edges. The connected components of  $\omega$  are referred to as clusters. Let  $p_c = p_c(G)$  be the critical probability for Bernoulli bond percolation on  $G$ , that is,

$$p_c = \inf \{p \in [0, 1] \mid \text{there exists an infinite cluster almost surely}\},$$

and let  $p_u = p_u(G)$  be the uniqueness threshold for Bernoulli bond percolation on  $G$ , that is,

$$p_u = \inf \{p \in [0, 1] \mid \text{there exists a unique infinite cluster almost surely}\}.$$

For  $p \in [0, 1]$  and  $x, y \in V$ , let  $\tau_p(x, y)$  be the probability that  $x$  and  $y$  are connected in  $\omega$ , that is,  $x$  and  $y$  belong to the same cluster. Let  $\chi_p(v)$  be the expected volume of the cluster containing  $v$  which is defined by

$$\chi_p(v) = \sum_{x \in V} \tau_p(v, x).$$

This expected volume is a monotone increasing function of  $p$ , and diverges at  $p_c$ . Aizenman and Newman [1] introduced the triangle condition. They analyzed the critical behavior of  $\chi_p(v)$  if  $G = \mathbb{Z}^d$  and the triangle condition holds at  $p_c$ . Let  $\nabla_p(v)$  be the triangle diagram which is defined by

$$\nabla_p(v) = \sum_{x, y \in V} \tau_p(v, x) \tau_p(x, y) \tau_p(y, v).$$

We say that  $G$  satisfies the triangle condition at  $p$  if  $\nabla_p(v) < \infty$  for every  $v$ . When  $G = \mathbb{Z}^d$ , Hara and Slade [3] showed that the triangle condition holds at  $p_c$  for all  $d \geq 19$ . This result was improved by Fitzner and van der Hofstadt [2] for  $d \geq 11$ . It is known  $p_c = p_u$  if  $G = \mathbb{Z}^d$ . Then above result also says that the triangle condition holds at  $p_u$ . When  $G$  is the product graphs, Kozma [6] showed

that the product graph of two  $d$ -regular trees  $T_d \square T_d$  for  $d \geq 3$  holds the triangle condition at  $p_c$ . In 2017, Hutchcroft [5] showed more general cases,  $G$  is the product graph of finitely many regular trees  $T_{d_1} \square T_{d_2} \square \cdots \square T_{d_N}$  for  $d_i \geq 3$ . Hutchcroft [4] also showed that the triangle condition holds at  $p_c$  if  $G$  is nonunimodular. Furthermore, Hutchcroft showed that  $p_c < p_u$  holds if  $G$  is nonunimodular. Then we does not have the result as to whether the triangle condition holds at  $p_u$ . A nonunimodular class contains  $T_d \square \mathbb{Z}$  for  $d \geq 3$ . Therefore, we consider percolation on  $T_d \square \mathbb{Z}$  and focus on the triangle condition holds or does not. This graph is a vertex transitive graph. Then we only consider  $v = o$  where  $o$  is a fixed origin. Our main result is the following theorem.

**Theorem 1.1.** *Let  $G = T_d \square \mathbb{Z}$  for  $d \geq 3$ . Then we have*

$$\nabla_p(o) \begin{cases} < \infty & (p < p_u) \\ = \infty & (p = p_u). \end{cases}$$

To lead this result, we use a certain function  $\alpha(p)$  which is defined by

$$\alpha(p) = \alpha_d(p) = \lim_{n \rightarrow \infty} \tau_p(o, (v_n, 0))^{\frac{1}{n}},$$

where  $v_n$  is a vertex on  $T_d$  with  $n$  distance from the origin. From a homogeneity of  $T_d$ ,  $\alpha(p)$  does not depend on a choice of  $v_n$ . We abbreviate  $v_n$  as  $n$ . We check on the existence of a limit. From FKG inequality, we have

$$\tau_p(o, (n+l, 0)) \geq \tau_p(o, (n, 0))\tau_p(o, (l, 0))$$

for all  $n, l \geq 0$ . By using Fekete's subadditive lemma, the existence of the limit is ensured, and we have

$$\alpha(p) = \lim_{n \rightarrow \infty} \tau_p(o, (n, 0))^{\frac{1}{n}} = \sup_{n \geq 1} \tau_p(o, (n, 0))^{\frac{1}{n}}.$$

This function was introduced by Schonmann [7], who showed the following inequality.

$$(1.1) \quad \alpha(p_u) \leq \frac{1}{\sqrt{b}}$$

where  $b = d - 1$ . By using this inequality, Schonmann showed that there exists a.s. no unique infinite cluster at  $p_u$ . We will show that the equality is established, that is,

$$(1.2) \quad \alpha(p_u) = \frac{1}{\sqrt{b}}.$$

We introduce a example of the triangle condition, let  $G = T_d$ , it is easy to check that

$$\nabla_p(v) \begin{cases} < \infty & (p < \frac{1}{\sqrt{b}}) \\ = \infty & (p = \frac{1}{\sqrt{b}}). \end{cases}$$

We already know that  $\alpha(p)$  is strictly increasing on  $[0, p_u]$  in [8], that is,  $\alpha(p) < 1/\sqrt{b}$  for all  $p < p_u$ . We will make an upper bound and a lower bound of  $\nabla_p(v)$  by using  $\alpha(p)$ , and to lead Theorem 1.1 similar to  $T_d$ .

## 2 Proof

We define the level difference function  $L(x, y)$  from  $T_d \times T_d$  to  $\mathbb{Z}$ . Let  $\xi$  be a fixed end of  $T_d$ . The parent of a vertex  $x \in T_d$  is the unique neighbor of  $x$  that is closer to  $\xi$  than  $x$  is. We call the other vertices of  $x$  its children. If  $y$  is parent of  $x$ , then we define  $L(x, y) = 1$ . If  $y$  is child of  $x$ , then we define  $L(x, y) = -1$ . In general cases, for any  $x, y$ , there exists a unique geodesic  $\{x_i\}_{i=0}^n$  such that  $x_0 = x$  and  $x_n = y$ , then we define

$$L(x, y) = \sum_{i=1}^n L(x_{i-1}, x_i).$$

Note that  $L(x, z) = L(x, y) + L(y, z)$  and  $L(y, x) = -L(x, y)$  for any  $x, y, z \in T_d$ . This function is extended to  $T_d \square \mathbb{Z}$  naturally. Let  $\pi$  be a natural projection from  $T_d \square \mathbb{Z}$  to  $T_d$ . Then we extend  $L(x, y)$  as  $L(x, y) = L(\pi(x), \pi(y))$ . Similarly, we have  $L(x, z) = L(x, y) + L(y, z)$  and  $L(y, x) = -L(x, y)$  for any  $x, y, z \in T_d \square \mathbb{Z}$ . We define  $\Delta(x, y)$  by

$$\Delta(x, y) = b^{L(x, y)}$$

for all  $x, y \in T_d \square \mathbb{Z}$  where  $b = d - 1$ . Note that  $\Delta(x, z) = \Delta(x, y)\Delta(y, z)$  and  $\Delta(y, x) = \Delta(x, y)^{-1}$  for any  $x, y, z$ . We define the tilted susceptibility by

$$\chi_{p, 1/2}(o) = \sum_{x \in T_d \square \mathbb{Z}} \tau_p(o, x) \Delta(o, x)^{1/2}.$$

Our method is based on [4], if you would like to know more detail of the tilted susceptibility, then please refer to [4]. Hutchcroft showed the following inequality.

$$(2.1) \quad \nabla_p(o) \leq (\chi_{p, 1/2}(o))^3.$$

Therefore we will show that  $\chi_{p, 1/2}(o) < \infty$  for  $p < p_u$  to prove the first half of Theorem 1.1. Similar to  $\alpha(p)$ , the function  $\beta(p)$  is defined by

$$\beta(p) = \lim_{m \rightarrow \infty} \tau_p(o \leftrightarrow (0, m))^{\frac{1}{m}} = \sup_{m \geq 1} \tau_p(o \leftrightarrow (0, m))^{\frac{1}{m}}.$$

By FKG inequality and the homogeneity of  $T_d \square \mathbb{Z}$ , we have

$$\tau_p(o, (n, m)) \leq \alpha(p)^n, \quad \tau_p(o, (n, m)) \leq \beta(p)^{|m|}$$

for each  $(n, m)$ . For  $x \in T_d$ , we define  $I_x(p)$  by

$$I_x(p) = \sum_{m \in \mathbb{Z}} \tau_p(o, (x, m)).$$

**Lemma 2.1** ([8]). *If  $\alpha(p) < 1/\sqrt{b}$ , then we have  $\beta(p) < 1$ .*

By this lemma, the function  $I_x(p)$  is well-defined for  $p < p_u$ .

**Lemma 2.2** ([8]). *For any  $p$  such that  $\alpha(p) < 1/\sqrt{b}$ , we have*

$$\lim_{|x| \rightarrow \infty} I_x(p)^{\frac{1}{|x|}} = \alpha(p).$$

By this lemma, for any  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that

$$I_x(p) \leq (\alpha(p) + \epsilon)^{|x|}$$

for any  $x \in T_d$  such that  $|x| \geq N$ . For any  $p < p_u$ , we choose  $\epsilon$  such that  $\alpha(p) + \epsilon < 1/\sqrt{b}$ . Then we have

$$\begin{aligned} \chi_{p,1/2}(o) &= \sum_{x \in T_d} I_x(p) \Delta(o, x)^{1/2} \\ &\leq \sum_{|x| < N} I_x(p) \Delta(o, x)^{1/2} + \sum_{|x| \geq N} (\alpha(p) + \epsilon)^{|x|} \Delta(o, x)^{1/2}. \end{aligned}$$

For  $r > 0$  and  $z \in \mathbb{R}_{>0}$ , we define the function  $J(r, z)$  by

$$J(r, z) = \sum_{x \in T_d} r^{|x|} z^{L(o, x)}.$$

**Lemma 2.3** ([8]). *For any  $r < 1/\sqrt{b}$  and  $z \in (br, 1/r)$ , we have  $J(r, z) < \infty$ .*

**Remark 2.4.** *In [8], using level function based on the origin  $o$ , it equal to  $-L(o, x)$ . Then  $z$  appeared in [8] means  $z^{-1}$  in this paper.*

By this lemma, let  $r = \alpha(p) + \epsilon$  and  $z = \sqrt{b}$ . Then we have

$$\sum_{|x| \geq N} (\alpha(p) + \epsilon)^{|x|} \Delta(o, x)^{1/2} \leq J(r, z) < \infty.$$

Therefore, we have  $\nabla_p(o) < \infty$  for all  $p < p_u$ . If  $p > p_u$ , then there exists a constant  $C(p) > 0$  such that  $\tau_p(x, y) \geq C(p)$  for all  $x, y$ . Hence, we have  $\chi_{p,1/2}(o) = \infty$  for  $p > p_u$ . Hutchcroft [4] showed that the set  $\{p \in [0, 1] \mid \chi_{p,1/2}(o) < \infty\}$  is open in  $[0, 1]$ . Then we have  $\chi_{p_u,1/2}(o) = \infty$ . That means  $\alpha(p)$  must equal to  $1/\sqrt{b}$ . Then we have the equation 1.2. By using this result, we will show that  $\nabla_{p_u}(o) = \infty$ . Similar to  $I_x(p)$ , we define the function  $\mathbb{I}_x(p)$  by

$$\mathbb{I}_x(p) = \sum_{m \in \mathbb{Z}} \tau_p(o, x)^2.$$

From a homogeneity of  $T_d \square \mathbb{Z}$ , we have  $\mathbb{I}_x(p) = \mathbb{I}_y(p)$  for any  $x, y$  such that  $|x| = |y|$ . For  $|x| = n$ , we denote  $\mathbb{I}_x(p)$  as  $\mathbb{I}_n(p)$ . By using BK inequality, we have

$$\mathbb{I}_{n+l}(p) \leq \sum_{m \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} \tau_p(o, (n, k))^2 \tau_p(o, (l, m - k))^2 = \mathbb{I}_n(p) \mathbb{I}_l(p)$$

for all  $n, l \geq 1$ . Then by using Fekete's subadditive lemma, we have

$$\inf_{n \geq 1} \mathbb{I}_n(p)^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \mathbb{I}_n(p)^{\frac{1}{n}}.$$

Since  $\mathbb{I}_n(p) \geq \tau_p(o, (n, 0))^2$ , we have

$$\mathbb{I}_n(p)^{\frac{1}{n}} \geq \tau_p(o, (n, 0))^{\frac{2}{n}}.$$

By taking the limit, we have

$$\lim_{n \rightarrow \infty} \mathbb{I}_n(p)^{\frac{1}{n}} \geq \alpha(p)^2$$

By above equation and inequality, we obtain

$$(2.2) \quad \mathbb{I}_n(p) \geq \alpha(p)^{2n}$$

for all  $n \geq 0$ . From FKG inequality, we have

$$(2.3) \quad \nabla_p(o) \geq \sum_{x,y} \tau_p(o,x)\tau_p(x,o)\tau_p(o,y)\tau_p(y,o) = \left( \sum_x \tau_p(o,x)^2 \right)^2.$$

If  $\nabla_{p_u}(o) < \infty$ , then also  $\sum \tau_{p_u}(o,x)^2 < \infty$ . Hence  $\mathbb{I}_x(p_u)$  is well-defined. On the other hand, by inequality 2.2 and equation 1.2, we have

$$\nabla_{p_u}(o) = \sum_{x \in T_d} \mathbb{I}_x(p_u) \geq \left( \sum_{x \in T_d} \alpha(p_u)^{2|x|} \right)^2 \geq \left( \sum_{n \geq 1} b^n \cdot \frac{1}{b^n} \right)^2 = \infty$$

Therefore, we have a contradiction, That means  $\nabla_{p_u}(o) = \infty$ . It ends the proof of 1.1.

## References

- [1] Aizenman, M.; Newman, C. M. Tree graph inequalities and critical behavior in percolation models. *J. Statist. Phys.* **36**, no.1-2, 107–143. (1984).
- [2] Fitzner, R.; van der Hofstad, R. Nearest-neighbor percolation function is continuous for  $d \geq 10$ . *arXiv preprint arXiv:1506.07977*. (2015).
- [3] Hara, T.; Slade, G. Mean-field behaviour and the lace expansion. *Probability and phase transition*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. **420**, 87–122. (1993). *Kluwer Acad. Publ., Dordrecht*. (1994).
- [4] Hutchcroft, T. Non-uniqueness and mean-field criticality for percolation on nonunimodular transitive graphs. *arXiv preprint arXiv:1711.02590*. (2017).
- [5] Hutchcroft, T. Statistical physics on a product of trees. *arXiv preprint arXiv:1712.04911*. (2017).
- [6] Kozma, G. Percolation on a product of two trees. *Ann. Probab.* **39**, no. 5, 1864–1895. (2011).
- [7] Schonmann, R. H. Percolation in  $\infty + 1$  dimensions at the uniqueness threshold. *Perplexing problems in probability. Progress in Probability.* **44**, 53–67. Birkhuser, Boston. (1999).
- [8] Yamamoto, K. Critical probability on the product graph of a regular tree and a line. *arXiv preprint arXiv:1810.07162*. (2018).

Mathematical Institute  
 Tohoku University  
 Sendai 980-8578  
 Japan  
 E-mail:kohei.yamamoto.t1@dc.tohoku.ac.jp