Parabolic-elliptic chemotaxis model with space-time dependent logistic sources on \mathbb{R}^N . III. Transition fronts

Rachidi B. Salako Department of Mathematics The Ohio State University Columbus OH, 43210-1174 and Wenxian Shen* Department of Mathematics and Statistics Auburn University

Auburn University, AL 36849

Abstract

The current work is the third of a series of three papers devoted to the study of asymptotic dynamics in the following parabolic-elliptic chemotaxis system with space and time dependent logistic source,

$$\begin{cases} \partial_t u = \Delta u - \chi \nabla \cdot (u \nabla v) + u(a(x,t) - b(x,t)u), & x \in \mathbb{R}^N, \\ 0 = \Delta v - \lambda v + \mu u, & x \in \mathbb{R}^N, \end{cases}$$
(0.1)

where $N \geq 1$ is a positive integer, χ, λ and μ are positive constants, and the functions a(x,t) and b(x,t) are positive and bounded. In the first of the series [45], we studied the phenomena of pointwise and uniform persistence for solutions with strictly positive initials, and the asymptotic spreading for solutions with compactly supported or front like initials. In the second of the series [46], we investigate the existence, uniqueness and stability of strictly positive entire solutions of (0.1). In particular, in the case of space homogeneous logistic source (i.e. $a(x,t) \equiv a(t)$ and $b(x,t) \equiv b(t)$), we proved in [46] that the unique spatially homogeneous strictly positive entire solution $(u^*(t), v^*(t))$ of (0.1) is uniformly and exponentially stable with respect to strictly positive perturbations when $0 < 2\chi\mu < \inf_{t \in \mathbb{R}} b(t)$.

In the current part of the series, we discuss the existence of transition front solutions of (0.1) connecting (0,0) and $(u^*(t), v^*(t))$ in the case of space homogeneous logistic source. We show that for every $\chi > 0$ with $\chi \mu \left(1 + \frac{\sup_{t \in \mathbb{R}} a(t)}{\inf_{t \in \mathbb{R}} a(t)}\right) < \inf_{t \in \mathbb{R}} b(t)$, there is a positive constant c_{χ}^* such that for every $\underline{c} > c_{\chi}^*$ and every unit vector ξ , (0.1) has a transition front solution of the form $(u(x,t), v(x,t)) = (U(x \cdot \xi - C(t), t), V(x \cdot \xi - C(t), t))$ satisfying that $C'(t) = \frac{a(t) + \kappa^2}{\kappa}$ for some positive number κ , $\liminf_{t-s \to \infty} \frac{C(t) - C(s)}{t-s} = \underline{c}$, and

$$\lim_{x \to -\infty} \sup_{t \in \mathbb{R}} |U(x,t) - u^*(t)| = 0 \quad \text{and} \quad \lim_{x \to \infty} \sup_{t \in \mathbb{R}} |\frac{U(x,t)}{e^{-\kappa x}} - 1| = 0.$$

Furthermore, we prove that there is no transition front solution $(u(x,t), v(x,t)) = (U(x \cdot \xi - C(t), t), V(x \cdot \xi - C(t), t))$ of (0.1) connecting (0,0) and $(u^*(t), v^*(t))$ with least mean speed less than $2\sqrt{\underline{a}}$, where $\underline{a} = \liminf_{t-s \to \infty} \frac{1}{t-s} \int_s^t a(\tau) d\tau$.

^{*}Partially supported by the NSF grant DMS-1645673

Key words. Parabolic-elliptic chemotaxis system, logistic source, classical solution, local existence, global existence, asymptotic stability, transition front.

2010 Mathematics Subject Classification. 35B35, 35B40, 35K57, 35Q92, 92C17.

1 Introduction and the Statements of the Main Results

The current work is the third part of a series of three papers on the asymptotic dynamics in the following parabolic-elliptic chemotaxis system with space-time dependent logistic source,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + u(a(x,t) - b(x,t)u), & x \in \mathbb{R}^N, \\ 0 = \Delta v - \lambda v + \mu u, & x \in \mathbb{R}^N, \end{cases}$$
(1.1)

where u(x,t) and v(x,t) denote mobile species density and chemical density functions, respectively, χ is a positive constant which measures the sensitivity with respect to chemical signals, a(x,t) and b(x,t) are positive functions and measure the growth and self limitation of the mobile species, respectively. The constant μ is positive and the term $+\mu u$ in the second equation of (1.1) indicates that the mobile species produces the chemical substance over time. The positive constant λ measures the degradation rate of the chemical substance. System (1.1) is a type of the celebrated parabolic-elliptic Keller-Segel chemotaxis systems (see [22, 23]) with space-time dependent logistic source.

The objective of the series of the three papers is to study the asymptotic dynamics in the chemotaxis system (1.1) on the whole space with space and/or time dependent logistic source. In the first of the series, [45], we studied the phenomena of pointwise and uniform persistence for solutions with strictly positive initials, and the asymptotic spreading in (1.1) for solutions with compactly supported or front like initials. In the second part of the series, [46], we investigated the existence, uniqueness and stability of strictly positive entire solutions of (1.1). In particular it was shown in [46] that, if the logistic source is space homogeneous, in which case (1.1) becomes,

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v) + u(a(t) - b(t)u), & x \in \mathbb{R}^N, \\ 0 = \Delta v - \lambda v + \mu u, & x \in \mathbb{R}^N, \end{cases}$$
(1.2)

then for every $\chi > 0$ with $0 < 2\chi\mu < \inf_{t \in \mathbb{R}} b(t)$ there is positive number $\alpha_{\chi} > 0$, such that for every positive initial function $u_0 \in C^b_{\text{unif}}(\mathbb{R}^N) = \{u \in C(\mathbb{R}^N) | u(x) \text{ is bounded and uniformly} continuous on <math>\mathbb{R}^M\}$ with $\inf_{x \in \mathbb{R}^N} u_0(x) > 0$, there is M > 0 such that

$$||u(\cdot, t+t_0, t_0, u_0) - u^*(t)||_{\infty} \le M e^{-\alpha_{\chi} t} \text{ and } ||v(\cdot, t+t_0; t_0, u_0) - v^*(t)||_{\infty} \le \frac{\mu}{\lambda} M e^{-\alpha_{\chi} t}$$

for all $t \ge 0$ and $t_0 \in \mathbb{R}$, where $(u(x, t; t_0, u_0, v_0), v(x, t; t_0, u_0))$ denote the unique classical solution of (1.2) with

$$\lim_{t \to 0^+} \|u(\cdot, t + t_0; t_0, u_0) - u_0(\cdot)\|_{\infty} = 0,$$

and $u^*(t)$ is the unique strictly positive entire solution of the Fisher-KKP equation

$$u_t = \Delta u + u(a(t) - b(t)u), \tag{1.3}$$

and $v^*(t) = \frac{\mu}{\lambda}u^*(t)$. Hence, when $0 < 2\chi\mu < \inf_{t \in \mathbb{R}} b(t)$, the unique spatially homogeneous strictly positive entire solution $(u^*(t), v^*(t))$ of (1.5) is uniformly and exponentially stable with respect to strictly positive perturbations.

Observe that with N = 1, $a(t) \equiv 1$ and $b(t) \equiv 1$, (1.3) becomes

$$u_t = u_{xx} + u(1-u), \quad x \in \mathbb{R}.$$

$$(1.4)$$

Equation (1.4) is called in literature Fisher-KPP equation due to the pioneering works of Fisher [10] and Kolmogorov, Petrowsky, Piskunov [24] on traveling wave solutions and take-over properties of (1.4). Fisher in [10] found traveling wave solutions $u(t, x) = \phi(x - ct)$ of (1.4) $(\phi(-\infty) = 1, \phi(\infty) = 0)$ of all speeds $c \ge 2$ and showed that there are no such traveling wave solutions of slower speed. He conjectured that the take-over occurs at the asymptotic speed 2. This conjecture was proved in [24] for some special initial distribution and was proved in [4] for the general case. More precisely, it is proved in [24] that for the nonnegative solution u(t, x) of (1.4) with u(0, x) = 1 for x < 0 and u(0, x) = 0 for x > 0, $\lim_{t\to\infty} u(t, ct)$ is 0 if c > 2 and 1 if c < 2. It is proved in [4] that for any nonnegative solution u(t, x) of (1.4), if at time t = 0, u is 1 near $-\infty$ and 0 near ∞ , then $\lim_{t\to\infty} u(t, ct)$ is 0 if c > 2 and 1 if c < 2 is called the spreading speed for (1.4).

A huge amount of research has been carried out toward various extensions of traveling wave solutions and take-over properties of (1.4) to general time and space independent as well as time and/or space dependent Fisher-KPP type equations. See, for example, [3], [4], [9], [15], [21], [47], [53], etc., for the extension to general time and space independent Fisher-KPP type equations; see [5, 6, 11, 20, 26, 27, 28, 31, 38, 39, 44, 54, 55], and references therein for the extension to time and/or space periodic Fisher-KPP type equations; and see [7, 8, 16, 19, 30, 32, 33, 34, 35, 37, 49, 50, 51, 52, 57, 58], and references therein for the extension to quite general time and/or space dependent Fisher-KPP type equations. It should be pointed out that the so called periodic traveling wave solutions or pulsating traveling fronts to time and/or space periodic reaction diffusion equations are natural extension of the notion of traveling waves to general time and/or space dependent reaction equations are the natural extension of the notion of traveling waves to general time and/or space dependent reaction equations are the natural extension of the notion of traveling wave solutions in the classical sense (see [7, 8] for the introduction of the notion of traveling wave solutions in the classical sense (see [7, 8] for the introduction of the notion of traveling wave solutions in the classical sense (see [7, 8] for the introduction of the notion of traveling wave solutions in the classical sense (see [7, 8] for the introduction of the notion of traveling wave solutions in the classical sense (see [7, 8] for the introduction of the notion of transition fronts or generalized traveling waves in the general case, and [30, 48, 50, 51] for the time almost periodic or space almost periodic cases).

Considering a chemotaxis model on the whole space, it is important to study the spatial spreading and propagating properties of the mobile species in the model. Transition front solutions or generalized traveling wave solutions and spatial spread speeds are among those used to characterize such properties. There are many studies on traveling wave solutions of various types of chemotaxis models, see, for example, [1, 2, 13, 18, 25, 29, 36, 41, 42, 56], etc.. It should be mentioned that spreading speeds and traveling wave solutions of (1.1) with a(x,t) and b(x,t) being constant functions are studied in [14, 36, 41, 42]. When a(x,t) and b(x,t) depend on x and t, as it is mentioned in the above there are many studies on spreading speeds and transition front solutions of (1.1) with $\chi = 0$, but there is little study on transition front solutions of (1.1) with $\chi \neq 0$.

The objective of this third part of the series is to study the existence of transition front solutions of (1.2), i.e., (1.1) in the case that $a(x,t) \equiv a(t)$ and $b(x,t) \equiv b(t)$, connecting (0,0) and $(u^*(t), v^*(t))$. To be more precise, we study the existence of positive entire solutions of (1.2) with the form $(u(x,t), v(x,t)) = (U(x \cdot \xi - C(t), t), V(x \cdot \xi - C(t), t))$ for some $\xi \in S^{N-1} = \{\xi \in \mathbb{R}^N \mid ||\xi|| = 1\}$ and some C(t), where $(U(-\infty,t), V(-\infty,t)) = (u^*(t), v^*(t))$ and $(U(\infty,t), V(\infty,t)) = (0,0)$. It is not difficult to see that, if $(u(x,t), v(x,t)) = (U(x \cdot \xi - C(t), t), V(x \cdot \xi - C(t), t))$ ($x \in \mathbb{R}^N$) is an entire solution of (1.2), then (u(x,t), v(x,t)) =

(U(x - C(t), t), V(x - C(t), t)) $(x \in \mathbb{R})$ is an entire solution of

$$\begin{cases} u_t = u_{xx} - \chi(uv_x)_x + u(a(t) - b(t)u), & x \in \mathbb{R}, \\ 0 = v_{xx} - \lambda v + \mu u, & x \in \mathbb{R}. \end{cases}$$
(1.5)

We will then study the existence of transition front solutions of (1.5) connecting (0,0) and $(u^{*}(t), v^{*}(t)).$

In the rest of the introduction, we introduce notations and standing assumptions, and state the main results of the current paper.

1.1 Notations and standing assumptions

For every function $w : \mathbb{R} \times I \to \mathbb{R}$, where $I \subset \mathbb{R}$, we set $w_{\inf}(t) = \inf_{x \in \mathbb{R}} w(x, t), w_{\sup}(t) =$ $\sup_{x \in \mathbb{R}} w(x,t), w_{\inf} = \inf_{(x,t) \in \mathbb{R} \times I} w(x,t), \text{ and } w_{\sup} = \sup_{(x,t) \in \mathbb{R} \times I} w(x,t).$ Let

 $C^b_{\mathrm{unif}}(\mathbb{R}) = \{ u \in C(\mathbb{R}) \, | \, u(x) \text{ is uniformly continuous in } x \in \mathbb{R} \text{ and } \sup_{x \in \mathbb{R}} |u(x)| < \infty \}$

equipped with the norm $||u||_{\infty} = \sup_{x \in \mathbb{R}} |u(x)|$. For any $0 \leq \nu < 1$, let

$$C_{\text{unif}}^{b,\nu}(\mathbb{R}) = \{ u \in C_{\text{unif}}^{b}(\mathbb{R}) \mid \sup_{x,y \in \mathbb{R}, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^{\nu}} < \infty \}$$

with norm $\|u\|_{C^{b,\nu}_{\text{unif}}} = \sup_{x \in \mathbb{R}} |u(x)| + \sup_{x,y \in \mathbb{R}, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^{\nu}}$. Hence $C^{b,0}_{\text{unif}}(\mathbb{R}) = C^b_{\text{unif}}(\mathbb{R})$. For given $f \in L^1_{\text{loc}}(\mathbb{R})$, let

$$\underline{f} = \liminf_{t-s \to \infty} \frac{1}{t-s} \int_s^t f(\tau) d\tau \quad \text{and} \quad \overline{f} = \limsup_{t-s \to \infty} \frac{1}{t-s} \int_s^t f(\tau) d\tau.$$

f and \overline{f} are called the *least mean* and *greatest mean* of f, respectively.

Throughout the remaining of this paper, we shall always suppose that the following standing assumption holds.

(H) $a(x,t) \equiv a(t)$ and $b(x,t) \equiv b(t)$ are uniformly Hölder continuous in $t \in \mathbb{R}$ with exponent $0 < \nu_0 < 1$ and

$$0 < \inf_{t \in \mathbb{R}} \min\{a(t), b(t)\} \le \sup_{t \in \mathbb{R}} \max\{a(t), b(t)\} < \infty$$

Observe from (\mathbf{H}) that

$$0 < a_{\inf} \le \underline{a} \le \overline{a} \le a_{\sup} < \infty.$$

$$(1.6)$$

1.2Main results

For given $u_0 \in C^b_{\text{unif}}(\mathbb{R})$ and $t_0 \in \mathbb{R}$, let $(u(x,t;t_0,u_0),v(x,t;t_0,u_0))$ be the classical solution of (1.5) with $u(x, t_0; t_0, u_0) = u_0(x)$ for every $x \in \mathbb{R}$ (see [40, Theorem 1.1] for the existence of $(u(x,t;t_0,u_0),v(x,t;t_0,u_0)))$. Note that if $u_0(x) \ge 0$ then $u(x,t;t_0,u_0) \ge 0$ and $v(x,t;t_0,u_0) \ge 0$ for every $x \in \mathbb{R}$ and $t \in [t_0, t_0 + T_{\max})$, where $[t_0, t_0 + T_{\max})$ denotes the maximal interval of existence of $(u(x,t;t_0,u_0),v(x,t;t_0,u_0))$. A classical solution (u(x,t),v(x,t)) of (1.5) is said to be an *entire solution* of (1.5) if it is defined for every $t \in \mathbb{R}$. Note that (u(x,t), v(x,t)) = (0,0)

is an equilibrium solution of (1.5). Throughout this work we shall denote by $u^*(t)$ the unique strictly positive entire solution of (1.3) and $v^*(t) = \frac{\mu}{\lambda}u^*(t)$. Then $(u^*(t), v^*(t))$ is a positive entire solution of (1.5).

An entire solution (u(x,t), v(x,t)) of (1.5) is called a *transition front solution* connecting (0,0) and $(u^*(t), v^*(t))$ if

$$(u(x,t), v(x,t)) = (U(x - C(t), t), V(x - C(t), t))$$
(1.7)

for some $U(\cdot, \cdot)$ and C(t) satisfying

$$\lim_{z \to -\infty} |U(z,t) - u^*(t)| = \lim_{z \to -\infty} |V(z,t) - v^*(t)| = 0, \quad \text{uniform in } t \in \mathbb{R},$$
(1.8)

and

$$\lim_{z \to \infty} U(z,t) = \lim_{z \to \infty} V(z,t) = 0, \quad \text{uniform in } t \in \mathbb{R}.$$
 (1.9)

Let

$$\underline{\mathbf{c}} = \liminf_{t-s \to \infty} \frac{C(t) - C(s)}{t-s}.$$

The function $(U(\cdot, \cdot), V(\cdot, \cdot))$ and <u>c</u> are called the *profile* and *least mean speed*, respectively, of the transition front solution (u(x,t), v(x,t)) = (U(x - C(t), t), V(x - C(t), t)). If we suppose that C(t) is of class C^1 and set c(t) = C'(t), then $(U(\cdot, \cdot), V(\cdot, \cdot))$ and $c(\cdot)$ satisfy

$$\begin{cases} U_t = U_{xx} + c(t)U_x - \chi(UV_x)_x + U(a(t) - b(t)U), & x \in \mathbb{R}, \ t \in \mathbb{R}, \\ 0 = V_{xx} - \lambda V + \mu U, & x \in \mathbb{R}, \ t \in \mathbb{R}. \end{cases}$$
(1.10)

Note that the function $\eta(\kappa) := \frac{\kappa(\sqrt{\lambda-\kappa^2}+\kappa)}{\lambda-\kappa^2}$ is strictly increasing on $(0,\sqrt{\lambda})$. For given $\chi > 0$ with $b_{\inf} > \chi\mu$, let $\kappa_{\chi} \in (0,\sqrt{\lambda})$ be such that

$$\frac{b_{\inf} - \chi \mu}{\chi \mu} = \frac{\kappa_{\chi}(\sqrt{\lambda - \kappa_{\chi}^2} + \kappa_{\chi})}{\lambda - \kappa_{\chi}^2}.$$
(1.11)

Then for any $0 < \kappa \leq \kappa_{\chi}$,

$$\frac{b_{\inf} - \chi \mu}{\chi \mu} \ge \frac{\kappa (\sqrt{\lambda - \kappa^2} + \kappa)}{\lambda - \kappa^2}.$$
(1.12)

Define

$$c_{\chi}^* = \frac{\underline{a} + \kappa_{\chi^*}^2}{\kappa_{\chi^*}},\tag{1.13}$$

where $\kappa_{\chi}^* = \min\{\kappa_{\chi}, \sqrt{\underline{a}}\}$. Let **(H1)** be the following standing assumption. **(H1)** $b_{\inf} > \chi \mu (1 + \frac{a_{\sup}}{a_{\inf}})$.

The main results on the existence of transition front solutions of (1.5) are stated in the following theorem.

Theorem 1.1. Suppose that (H1) holds.

(1) For every $\underline{c} > c_{\chi}^*$, (1.5) has a transition front solution (u(x,t), v(x,t)) = (U(x-C(t),t), V(x-C(t),t)) connecting (0,0) and $(u^*(t), v^*(t))$ with least mean speed \underline{c} . Furthermore, it holds that $C(t) = \int_0^t \frac{a(s)+\kappa^2}{\kappa} ds$ and

$$\lim_{x \to -\infty} \sup_{t \in \mathbb{R}} |U(x,t) - u^*(t)| = 0, \quad and \quad \lim_{x \to \infty} \sup_{t \in \mathbb{R}} |\frac{U(x,t)}{e^{-\kappa x}} - 1| = 0, \tag{1.14}$$

where $\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{\underline{a}}\})$ is such that $\underline{c} = \frac{\underline{a} + \kappa^2}{\kappa}$.

(2) If a(t) and b(t) are periodic in t with period T, then for every $c > c_{\chi}^*$, (1.5) has a periodic transition front solution (u(x,t), v(x,t)) = (U(x - ct, t), V(x - ct, t)) connecting (0,0) and $(u^*(t), v^*(t))$, and satisfying

$$U(x,t+T) = U(x,t), \quad V(x,t+T) = V(x,t),$$
(1.15)

and

$$\lim_{x \to -\infty} \sup_{t \in \mathbb{R}} |U(x,t) - u^*(t)| = 0, \quad and \quad \lim_{x \to \infty} \sup_{t \in \mathbb{R}} |\frac{U(x,t)}{e^{-\kappa(x + ct - \int_0^t c_\kappa(s)ds)}} - 1| = 0, \quad (1.16)$$

where $\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{\hat{a}}\})$ $(\hat{a} = \frac{1}{T} \int_0^T a(t) dt)$ satisfies $c = \frac{\hat{a} + \kappa^2}{\kappa}$, and $c_{\kappa}(s) = \frac{a(s) + \kappa^2}{\kappa}$.

(3) If $a(t) \equiv a$ and $b(t) \equiv b$ are independent of t, then for every $c > c_{\chi}^*$, (1.5) has a traveling wave solution (u(x,t), v(x,t)) = (U(x-ct), V(x-ct)) connecting (0,0) and $(u^*, v^*) = (\frac{a}{b}, \frac{\mu}{\lambda} \frac{a}{b})$, and satisfying

$$\lim_{x \to -\infty} |U(x) - u^*| = 0, \quad and \quad \lim_{x \to \infty} |\frac{U(x)}{e^{-\kappa x}} - 1| = 0, \tag{1.17}$$

where
$$\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{a}\})$$
 satisfies $c = \frac{a+\kappa^2}{\kappa}$.

We have the following theorem on the nonexistence of transition front solutions of (1.5).

Theorem 1.2. Suppose that **(H1)** holds. For every $\underline{c} < 2\sqrt{\underline{a}}$, (1.5) has no transition front solution (u(x,t), v(x,t)) = (U(x - C(t), t), V(x - C(t), t)) connecting (0,0) and $(u^*(t), v^*(t))$ with least mean speed \underline{c} .

- **Remark 1.3.** (1) The results in Theorem 1.1(1), (2), and Theorem 1.2 are new. The result in Theorem 1.1(3) extends the results in [41] for the case $\lambda = \mu = 1$ (see [41, Theorem A and Remark 1.1]).
 - (2) By Theorem 1.1(1), for every $\xi \in S^{N-1} = \{\xi \in \mathbb{R}^N \mid \|\xi\| = 1\}$ and every $\underline{c} > c_{\chi}^*$, (1.1) has a transition front solution $(u(x,t), v(x,t)) = (U(x \cdot \xi - C(t), t), V(x \cdot \xi - C(t), t))$ connecting (0,0) and $(u^*(t), v^*(t))$ with least mean speed \underline{c} , where $C(t) = \int_0^t \frac{a(s) + \kappa^2}{\kappa} ds$, $\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{\underline{a}}\})$ is such that $\underline{c} = \frac{\underline{a} + \kappa^2}{\kappa}$, and (U(x, t), V(x, t)) satisfies (1.14).
 - (3) Let $c_0^* = 2\sqrt{\underline{a}}$. It is proved in [33, Theorem 2.3] that c_0^* is the minimal least mean speed of transition front solutions of (1.3), i.e., (1.5) in the absence of the chemotaxis, in the sense that for any $\underline{c} > c_0^*$, (1.3) has a transition front solution connecting 0 and $u^*(t)$ with least mean speed \underline{c} , and (1.3) has no transition front solutions connecting 0 and $u^*(t)$ with least mean speed smaller than c_0^* .

(4) For fixed $\chi > 0$ with $b_{inf} > \chi\mu$, when the degradation rate λ of the chemical substance is sufficiently large, we have $\kappa_{\chi} \ge \sqrt{\underline{a}}$ and $\kappa_{\chi}^* = \sqrt{\underline{a}}$, hence $c_{\chi}^* = 2\sqrt{\underline{a}}$, which is the minimal least mean speed of transition front solutions of (1.3). Indeed, since the function $\lambda \mapsto \frac{\sqrt{\underline{a}}(\sqrt{\lambda-\underline{a}}+\sqrt{\underline{a}})}{\lambda-\underline{a}}$ is strictly decreasing and satisfies

$$\lim_{\lambda \to \underline{a}+} \frac{\sqrt{\underline{a}} \left(\sqrt{\lambda - \underline{a}} + \sqrt{\underline{a}}\right)}{\lambda - \underline{a}} = \infty \text{ and } \lim_{\lambda \to \infty} \frac{\sqrt{\underline{a}} \left(\sqrt{\lambda - \underline{a}} + \sqrt{\underline{a}}\right)}{\lambda - \underline{a}} = 0,$$

there is a unique $\lambda_{\chi} > \underline{a}$ such that

$$\frac{\sqrt{\underline{a}}\left(\sqrt{\lambda_{\chi}-\underline{a}}+\sqrt{\underline{a}}\right)}{\lambda_{\chi}-\underline{a}} = \frac{b_{\text{inf}}-\chi\mu}{\chi\mu}$$

This implies that for any $\lambda > \lambda_{\chi}$,

$$\frac{\sqrt{\underline{a}}\left(\sqrt{\lambda - \underline{a}} + \sqrt{\underline{a}}\right)}{\lambda - \underline{a}} < \frac{b_{\inf} - \chi\mu}{\chi\mu},$$

and hence $\sqrt{\underline{a}} < \kappa_{\chi} < \sqrt{\lambda}$, where κ_{χ} is such that (1.11) holds. It then follows that, when $\lambda \geq \lambda_{\chi}(>\underline{a}), c_{\chi}^* = c_0^* = 2\sqrt{\underline{a}}$. Hence, Theorem 1.1(1) and Theorem 1.2 imply that the minimal least mean speed $c_0^* = 2\sqrt{\underline{a}}$ of transition front solutions to the time heterogeneous Fisher-KPP equation (1.3) is also the minimal least mean speed of the transition front solutions to the chemotaxis model (1.5) with $\lambda > \lambda_{\chi}(>\underline{a})$.

- (5) For fixed $\lambda > \underline{a}$, when $\chi > 0$ is sufficiently small, we also have $c_{\chi}^* = 2\sqrt{\underline{a}}$. Indeed, for any given $\lambda > \underline{a}$, we have that $c_{\chi}^* = 2\sqrt{\underline{a}}$ whenever $\chi \mu < \min\left\{\frac{a_{\inf}b_{\inf}}{a_{\inf}+a_{\sup}}, \frac{b_{\inf}(\lambda-\underline{a})}{\lambda-\underline{a}+\sqrt{\underline{a}}(\sqrt{\lambda-\underline{a}}+\sqrt{\underline{a}})}\right\}$. Again, Theorem 1.1(1) and Theorem 1.2 imply that the minimal least mean speed $c_0^* = 2\sqrt{\underline{a}}$ of transition front solutions to the time heterogeneous Fisher-KPP equation (1.3) is also the minimal least mean speed of the transition front solutions to the chemotaxis model (1.5) with χ sufficiently small.
- (6) It remains open whether for fixed $0 < \lambda \leq \underline{a}$, when $\chi > 0$ is sufficiently small, for any $\underline{c} > c_0^* = 2\sqrt{\underline{a}}$, (1.5) has a transition front solution connecting (0,0) and $(u^*(t), v^*(t))$ with least mean speed \underline{c} .

The rest of the paper is organized as follows. In section 2, we construct proper sub-solutions and super-solutions of some equations related to (1.10) with certain c(t), which will be of great use in the proofs of the main results. We prove Theorem 1.1 and Theorem 1.2 in sections 3 and 4, respectively.

2 Sub- and super-solutions

In this section, we construct proper sub-solutions and super-solutions of some equations related to (1.10) with certain c(t).

For any fixed
$$0 < \kappa < \min\{\kappa_{\chi}, \sqrt{\underline{a}}\} (\leq \min\{\sqrt{\lambda}, \sqrt{\underline{a}}\}), \text{ let } c_{\kappa}(t) = \frac{a(t) + \kappa^2}{\kappa},$$

 $\phi_{\kappa}(x) = e^{-\kappa x}, \quad \forall x \in \mathbb{R},$

$$(2.1)$$

and

$$\phi_{\kappa}^{+}(x) = \min\{\phi_{\kappa}(x), \frac{a_{\sup}}{b_{\inf} - \chi\mu}\}, \quad \forall \ x \in \mathbb{R}.$$
(2.2)

It is not difficult to see that

$$\frac{d^2}{dx^2}\phi_{\kappa} + c_{\kappa}(t)\frac{d}{dx}\phi_{\kappa} + a(t)\phi_{\kappa} = \left(\kappa^2 - \kappa c(t) + a(t)\right)\phi_{\kappa} = 0, \quad \forall \ x \in \mathbb{R}, \ t \in \mathbb{R}.$$
(2.3)

Note that

$$\underline{c}_{\kappa} := \liminf_{t-s \to \infty} \frac{1}{t-s} \int_{s}^{t} c_{\kappa}(y) dy = \frac{\underline{a} + \kappa^{2}}{\kappa}, \quad \forall \ \kappa > 0,$$
(2.4)

and

$$\overline{c}_{\kappa} := \limsup_{t-s \to \infty} \frac{1}{t-s} \int_{s}^{t} c_{\kappa}(y) dy = \frac{\overline{a} + \kappa^{2}}{\kappa}, \quad \forall \kappa > 0.$$
(2.5)

Hence,

$$\frac{a_{\inf} + \kappa^2}{\kappa} \le \underline{c}_{\kappa} \le \overline{c}_{\kappa} \le \frac{a_{\sup} + \kappa^2}{\kappa}, \quad \forall \ \kappa > 0.$$

Fix $0 < \kappa < \min\{\kappa_{\chi}, \sqrt{\underline{a}}\}\ \text{and}\ 0 < \beta_0 < \frac{1}{3}$. Let

$$\mathcal{E}_{\kappa,\beta_0,M} = \{ \phi \in C(\mathbb{R}, C^b_{\text{unif}}(\mathbb{R})) \mid 0 \le \phi(x,t) \le \phi^+_\kappa(x) \text{ and } |\phi(x+h,t) - \phi(x,t)| \le M |h|^{\beta_0}, \\ |\phi(x,t+h) - \phi(x,t)| \le M |h|^{\beta_0} \,\,\forall x,t,h \in \mathbb{R}, \, |h| \le 1 \},$$
(2.6)

where M is a positive constant to be determined later. For every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, consider

$$\mathcal{L}_{\kappa,\phi}(u) = 0, \tag{2.7}$$

where

$$\mathcal{L}_{\kappa,\phi}(u) = \partial_t u - \partial_{xx} u - (c_\kappa(t) - \chi \partial_x \psi(\cdot, \cdot; \phi)) \partial_x u - (a(t) - \chi \lambda \psi(\cdot, \cdot; \phi) - (b(t) - \chi \mu) u) u,$$

and $\psi(\cdot, \cdot; \phi)$ is given by

$$\psi(x,t;\phi) = \mu \int_0^\infty \frac{e^{-\lambda s}}{\sqrt{4\pi s}} \Big[\int_{\mathbb{R}} e^{-\frac{|x-y|^2}{4s}} \phi(y,t) dy \Big] ds$$
$$= \frac{\mu}{\sqrt{\pi}} \int_0^\infty \int_{\mathbb{R}} e^{-\lambda s - |y|^2} \phi(x + 2\sqrt{s}y,t) dy ds.$$
(2.8)

It is not difficult to prove that $\psi(\cdot,\cdot;\phi)$ solves

$$\partial_{xx}\psi(x,t;\phi) - \lambda\psi(x,t;\phi) + \mu\psi(x,t;\phi) = 0, \quad x \in \mathbb{R}, \ t \in \mathbb{R}.$$

Note that, for every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, it holds that

$$\partial_x \psi(x,t;\phi) = \mu \int_0^\infty \frac{e^{-\lambda s}}{\sqrt{4\pi s}} \Big[\int_{\mathbb{R}} \frac{(y-x)e^{-\frac{|x-y|^2}{4s}}}{2s} \phi(y,t)dy \Big] ds$$
$$= \frac{\mu}{\sqrt{\pi}} \int_0^\infty \frac{e^{-\lambda s}}{\sqrt{s}} \Big[\int_{\mathbb{R}} ze^{-z^2} \phi(x+2z\sqrt{s},t)dz \Big] ds.$$
(2.9)

Note also that, for given $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, if $u(x,t) = \phi(x,t)$ is an entire solution of (2.7), then $(U(x,t), V(x,t)) = (\phi(x,t), \psi(x,t;\phi))$ is an entire solution of (1.10) with $c(t) = c_{\kappa}(t)$. In the following, for given $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, we construct proper sub- and super-solutions of (2.7).

Definition 2.1. For each given $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, a function $u \in C^{2,1}(D)$, where $D \subset \mathbb{R} \times \mathbb{R}$, is called a super-solution (resp. sub-solution) of (2.7) on D if

$$\mathcal{L}_{\kappa,\phi}(u)(x,t) \geq 0 \ (resp. \ \mathcal{L}_{\kappa,\phi}(u)(x,t) \leq 0) \ for \ (x,t) \in D.$$

The following Lemma provides some useful estimates on $\psi(\cdot, \cdot, \phi)$ and $\partial_x \psi(\cdot, \cdot, \phi)$ for each $\phi \in \mathcal{E}_{\kappa}$.

Lemma 2.2. (i) For every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, we have that

$$0 \le \psi(x,t;\phi) \le \min\{\frac{\mu a_{\sup}}{\lambda(b_{\inf} - \chi\mu)}, \ \frac{\mu}{\lambda - \kappa^2}\phi_{\kappa}(x)\}, \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}$$
(2.10)

and

$$|\partial_x \psi(x,t;\phi)| \le \frac{\mu(\sqrt{\lambda-\kappa^2}+\kappa)}{\lambda-\kappa^2} \phi_\kappa(x), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}.$$
(2.11)

(ii) Let $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$ and $\{\phi_n\}_{n\geq 1} \subset \mathcal{E}_{\kappa,\beta_0,M}$ such that $\phi_n(x,t) \to \phi(x,t)$ as $n \to \infty$ uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$. Then

 $\psi(x,t;\phi_n) \to \psi(x,t;\phi) \quad and \quad \psi_x(x,t;\phi_n) \to \psi_x(x,t;\phi) \ as \ n \to \infty$

uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$.

Proof. (i) The following arguments are inspired from the proofs of [41, Lemmas 2.2 & 2.3]. So we refer the reader to [41] for more details on the estimates. Let $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$ be given. Since $0 \leq \phi(x,t) \leq \phi_{\kappa}(x)$, it follows from (2.8) that

$$\psi(x,t;\phi) \le \mu \int_0^\infty \frac{e^{-\lambda s}}{\sqrt{4\pi s}} \Big[\int_{\mathbb{R}} e^{-\frac{|x-y|^2}{4s}} e^{-\kappa y} dy \Big] ds = \frac{\mu}{\lambda - \kappa^2} \phi_\kappa(x).$$
(2.12)

On the other hand, we have

$$0 \le \psi(x,t;\phi) \le \mu \Big[\int_0^\infty \frac{e^{-\lambda s}}{\sqrt{4\pi s}} \Big[\int_{\mathbb{R}} e^{-\frac{|x-y|^2}{4s}} dy \Big] ds \Big] \sup_{x \in \mathbb{R}} \phi(x,t) = \frac{\mu}{\lambda} \sup_{x \in \mathbb{R}} \phi(x,t).$$
(2.13)

Inequality (2.10) follows from (2.12) and (2.13).

Using (2.9) we have

$$\left|\partial_x \psi(x,t;\phi)\right| \le \frac{\mu e^{-\kappa x}}{\sqrt{\pi}} \int_0^\infty \frac{e^{-(\lambda-\kappa^2)s}}{\sqrt{s}} \Big[\int_{\mathbb{R}} (|z|+\kappa\sqrt{s})e^{-z^2}dz \Big] ds = \frac{\mu(\sqrt{\lambda-\kappa^2}+\kappa)}{\lambda-\kappa^2} \phi_\kappa(x). \quad (2.14)$$

Inequality (2.11) then follows from (2.14). This completes the proof of (i).

(ii) Let $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$ and $\{\phi_n\}_{n\geq 1} \subset \mathcal{E}_{\kappa,\beta_0,M}$ be such that $\phi_n(x,t) \to \phi(x,t)$ as $n \to \infty$ uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$. Then

$$\lim_{n \to \infty} \sup_{(x,t) \in [-K,K]^2} \int_0^R \int_{-R}^R e^{-\lambda s} e^{-|y|^2} |\phi_n(x + 2\sqrt{sy}, t) - \phi(x + 2\sqrt{sy}, t)| dy ds = 0, \forall K > 0, \ R > 0.$$

On the other hand, observe that

$$\lim_{R \to \infty} \sup_{n \ge 1, x \in \mathbb{R}, \ t \in \mathbb{R}} \int_{\{s \ge R \text{ or } |y| \ge R\}} e^{-\lambda s} e^{-|y|^2} \phi_n(x + 2\sqrt{s}y, t) dy ds = 0.$$

Therefore, it follows from (2.8) that $\psi(x,t;\phi_n) \to \psi(x,t;\phi)$ as $n \to \infty$ uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$.

Similarly, using (2.9), the similar arguments to the above yield that $\psi_x(x,t;\phi_n) \to \psi_x(x,t;\phi)$ as $n \to \infty$ uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$.

By (2.4), we have that

$$\underline{c}_{\kappa} - 2\kappa \ge \frac{\underline{a} - \kappa^2}{\kappa} > 0, \quad \forall \ 0 < \kappa < \min\{\kappa_{\chi}, \sqrt{\underline{a}}\}.$$
(2.15)

Hence, since (H1) holds, there is $\varepsilon > 0$ such that

$$0 < \varepsilon < \min\{\kappa, \frac{\underline{a} - \kappa^2}{\kappa}\} \quad \text{and} \quad \frac{b_{\inf} - \chi\mu}{\chi\mu} > \frac{(\kappa + \varepsilon)(\sqrt{\lambda - \kappa^2} + \kappa) - \lambda}{\lambda - \kappa^2}.$$
(2.16)

With this choice of ε , it readily follows that $\underline{c}_{\kappa} - 2\kappa - \varepsilon = \frac{1}{\kappa}(\underline{a} - \kappa(\kappa + \varepsilon)) > 0.$

Lemma 2.3. Fix an $\epsilon > 0$ satisfying (2.16). There is $A \in W^{1,\infty}_{loc}(\mathbb{R}) \cap L^{\infty}(\mathbb{R})$ such that

$$A'(t) + \varepsilon(c_{\kappa}(t) - 2\kappa - \varepsilon) > A_0 := \frac{\varepsilon}{2}(\underline{c}_{\kappa} - 2\kappa - \varepsilon) = \frac{\varepsilon}{2\kappa}(\underline{a} - \kappa(\kappa + \varepsilon)) > 0, \quad a. \quad e. \quad t \in \mathbb{R}.$$
(2.17)

Moreover, there exist $\{t_n\}_{n\in\mathbb{Z}}$ such that $t_n < t_{n+1}, t_n \to \pm \infty$ as $n \to \pm \infty$ and $A \in C^1(t_n, t_{n+1})$ for every $n \in \mathbb{Z}$.

Proof. Note that $0 < \underline{c}_{\kappa} \leq \overline{c}_{\kappa} < \infty$. The lemma follows from [43, Lemma 2.2] and its proof.

We introduce the following expressions

$$A_1 = b_{\sup} - \chi \mu + \frac{\chi \mu \left(\kappa (\sqrt{\lambda^2 - \kappa^2} + \kappa) + \lambda\right)}{\lambda - \kappa^2}$$
(2.18)

and

$$A_2 = b_{\inf} - \chi \mu + \frac{\chi \mu \left(\lambda - (\kappa + \varepsilon)(\sqrt{\lambda - \kappa^2} + \kappa)\right)}{\lambda - \kappa^2}.$$
(2.19)

It follows from (2.16) that $A_1 > 0$ and $A_2 > 0$. Finally, let us take

$$d = e^{\|A\|_{\infty}} \left(1 + \frac{A_0}{A_1}\right) \tag{2.20}$$

and define

$$\phi_{\kappa,\varepsilon,d,A}(x,t) = \phi_{\kappa}(x) - de^{A(t)}\phi_{\kappa+\varepsilon}(x).$$
(2.21)

We introduce the following functions

$$x_{\kappa,\varepsilon,d,A}^{-}(t) := \frac{\ln(d) + A(t)}{\varepsilon} = \frac{\|A\|_{\infty} + A(t) + \ln(1 + \frac{A_0}{A_1})}{\varepsilon} > 0, \qquad (2.22)$$

and

$$x_{\kappa,\varepsilon,d,A}^{+}(t) := \frac{\ln(\frac{\kappa+\varepsilon}{\kappa}d) + A(t)}{\varepsilon}, \quad \forall t \in \mathbb{R}.$$
(2.23)

It is clear from the definition of $x^+_{\kappa,\varepsilon,d,A}(t)$ and $x^-_{\kappa,\varepsilon,d,A}(t)$ that

$$x^{+}_{\kappa,\varepsilon,d,A}(t) - x^{-}_{\kappa,\varepsilon,d,A}(t) = \frac{\ln(\frac{\kappa+\varepsilon}{\kappa})}{\varepsilon}, \quad \forall t \in \mathbb{R}.$$
(2.24)

The next result provides some useful information on the relationship between the functions $\phi_{\kappa,\varepsilon,d,A}(x,t), x^+_{\kappa,\varepsilon,d,A}(t)$, and $x^-_{\kappa,\varepsilon,d,A}(t)$.

Lemma 2.4. Let $\phi_{\kappa,\varepsilon,d,A}(x,t)$, $x^+_{\kappa,\varepsilon,d,A}(t)$, and $x^-_{\kappa,\varepsilon,d,A}(t)$ be given by (2.21), (2.23), and (2.22), respectively. Then, the following hold.

(i) For every $t \in \mathbb{R}$, we have that

$$\phi_{\kappa,\varepsilon,d,A}(\bar{x_{\kappa,\varepsilon,d,A}}(t),t) = 0 \text{ and } (x - \bar{x_{\kappa,\varepsilon,d,A}}(t))\phi_{\kappa,\varepsilon,d,A}(x,t) > 0, \ \forall x \neq \bar{x_{\kappa,\varepsilon,d,A}}(t).$$
(2.25)

(ii) For every $t \in \mathbb{R}$, we have that

$$0 < \frac{\ln(d) - \|A\|_{\infty}}{\varepsilon} \le x_{\kappa,\varepsilon,d,A}^{-}(t) < x_{\kappa,\varepsilon,d,A}^{+}(t) \le \frac{\ln(\frac{\kappa+\varepsilon}{\kappa}d) + \|A\|_{\infty}}{\varepsilon} < \infty.$$
(2.26)

(iii) For every $t \in \mathbb{R}$, the function $\mathbb{R} \ni x \mapsto \phi_{\kappa,\varepsilon,d,A}(x,t)$ is strictly increasing on the interval $(-\infty, x^+_{\kappa,\varepsilon,d,A}(t)]$ and is strictly decreasing on the interval $[x^+_{\kappa,\varepsilon,d,A}(t),\infty)$. Hence, we have that $\max_{x\in\mathbb{R}}\phi_{\kappa,\varepsilon,d,A}(x,t) = \phi_{\kappa,\varepsilon,d,A}(x^+_{\kappa,\varepsilon,d,A}(t),t)$ for each $t\in\mathbb{R}$. Moreover, it holds that

$$0 < \inf_{t \in \mathbb{R}} \phi_{\kappa,\varepsilon,d,A}(x_{\kappa,\varepsilon,d,A}^+(t), t).$$
(2.27)

Proof. Lemma 2.4 (i) and (ii) follow from (2.22), (2.23) and (2.24). Observe that

$$\partial_x \phi_{\kappa,\varepsilon,d,A}(x,t) = -\kappa \Big(e^{\varepsilon x} - \frac{(\kappa + \varepsilon) de^{A(t)}}{\kappa} \Big) e^{-(\kappa + \varepsilon)x}, \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R},$$

and

$$\partial_x \phi_{\kappa,\varepsilon,d,A}(x^+_{\kappa,\varepsilon,d,A}(t),t) = 0, \quad \forall \ t \in \mathbb{R}.$$

Hence the first statement in Lemma 2.4 (iii) follows from the first derivative test. We have that

$$\phi_{\kappa,\varepsilon,d,A}(x_{\kappa,\varepsilon,d,A}^{+}(t),t) = \frac{\varepsilon}{\kappa} e^{A(t) - (\kappa + \varepsilon)x_{\kappa,\varepsilon,d,A}^{+}(t)} \ge \frac{\varepsilon}{\kappa} e^{-\frac{(\kappa + \varepsilon)}{\varepsilon} \left(2\|A\|_{\infty} + \ln(d\frac{\kappa + \varepsilon}{\kappa})\right)} > 0, \quad \forall t \in \mathbb{R}$$

This completes the proof of the Lemma.

The next result provides us with sub- and super-solutions of (2.7) for every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$.

Theorem 2.5. Suppose that **(H1)** holds and let $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$. Then the following hold.

- (i) The function $u(x,t) = \phi_{\kappa}(x)$ is a super-solution of (2.7) on $\mathbb{R} \times \mathbb{R}$.
- (ii) The constant function $u(x,t) = \frac{a_{\sup}}{b_{\inf} \chi \mu}$ is a super-solution of (2.7) on $\mathbb{R} \times \mathbb{R}$.
- (iii) The function $\phi_{\kappa,\varepsilon,d,A}(x,t)$ is a sub-solution of (2.7) on the set $D_{\kappa,\varepsilon,d,A}$ defined by

$$D_{\kappa,\varepsilon,d,A} := \{ (x,t) \in \mathbb{R} \times \mathbb{R} : x \ge x_{\kappa,\varepsilon,d,A}^{-}(t) \}.$$

(iv) For every $0 < \delta \leq \frac{b_{\inf} - \chi \mu \left(1 + \frac{a_{\sup}}{a_{\inf}}\right)}{(b_{\inf} - \chi \mu)(b_{\sup} - \chi \mu)}$, the constant function $u(x, t) = \delta$ is a sub-solution of (2.7) on $\mathbb{R} \times \mathbb{R}$.

Proof. Let $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$ be given. The theorem can be proved by the similar arguments as those in [41, Theorem 2.1]. For the completeness, we provide a proof in the following.

(i) Using (2.3) and (2.11), for every $(x,t) \in \mathbb{R} \times \mathbb{R}$, we have

$$\mathcal{L}_{\kappa,\phi}(\phi_{\kappa})(x,t) = -\kappa\chi\partial_{x}\psi(x,t;u)\phi_{\kappa} + (\chi\lambda\psi(x,t;\phi) + (b(t) - \chi\mu)\phi_{\kappa})\phi_{\kappa}$$

$$\geq \left(b_{\inf} - \chi\mu - \kappa\chi\frac{\mu(\sqrt{\lambda - \kappa^{2}} + \kappa)}{\lambda - \kappa^{2}}\right)\phi_{\kappa}^{2} + \chi\lambda v(\cdot,\cdot;u)\phi_{\kappa} \geq 0.$$
(2.28)

Hence (i) follows.

(ii) Since $0 \le \psi(x, t; \phi)$, we have that

$$\mathcal{L}_{\kappa,\phi}(\frac{a_{\sup}}{b_{\inf}-\chi\mu}) = -\left(a(t)-\chi\lambda\psi(\cdot,\cdot;\phi)-(b(t)-\chi\mu)\frac{a_{\sup}}{b_{\inf}-\chi\mu}\right)\frac{a_{\sup}}{b_{\inf}-\chi\mu}$$
$$\geq \left((b(t)-\chi\mu)\frac{a_{\sup}}{b_{\inf}-\chi\mu}-a(t)\right)\frac{a_{\sup}}{b_{\inf}-\chi\mu} \geq 0.$$

Hence (ii) follows.

(iii) We first note that x > 0 whenever $(x,t) \in D_{\kappa,\varepsilon,d,A}$. For $(x,t) \in D_{\kappa,\varepsilon,d,A}$, let $\tilde{A}(t,x) = A(t) - (\kappa + \varepsilon)x$ and using Lemma 2.2 and Lemma 2.3, we have

$$\begin{split} \mathcal{L}_{\kappa,\phi}(\phi_{\kappa,\varepsilon,d,A})(x,t) &= -dA'e^{\tilde{A}(t,x)} - \left[\kappa^2 e^{-\kappa x} - d(\kappa+\varepsilon)^2 e^{\tilde{A}(t,x)}\right] - (c_\kappa(t) - \chi \partial_x \psi(\cdot,\cdot;\phi)) \left[-\kappa e^{-\kappa x} + d(\kappa+\varepsilon) e^{\tilde{A}(t,x)}\right] \\ &- \left[a(t) - \chi \lambda \psi(\cdot,\cdot;\phi) - (b(t) - \chi \mu) \phi_{\kappa,\varepsilon,d,A}\right] \phi_{\kappa,\varepsilon,d,A} \\ &= -d\left(A' + \frac{\varepsilon}{\kappa} \left(a(t) - \kappa(\kappa+\varepsilon)\right)\right) e^{\tilde{A}(t,x)} + \chi \partial_x \psi(\cdot,\cdot;\phi) \left(-\kappa e^{-\kappa x} + d(\kappa+\varepsilon) e^{\tilde{A}(t,x)}\right) \\ &+ \chi \lambda \psi(\cdot,\cdot;\phi) \phi_{\kappa,\varepsilon,d,A} + (b(t) - \chi \mu) e^{-2\kappa x} - d(b(t) - \chi \mu) \left(e^{-\kappa x} + \phi_{\kappa,\varepsilon,d,A}\right) e^{\tilde{A}(t,x)} \\ &\leq -d\left(A' + \varepsilon(c_\kappa(t) - 2\kappa - \varepsilon)\right) e^{\tilde{A}(t,x)} + \frac{\chi \lambda \mu}{\lambda - \kappa^2} \phi_\kappa(\phi_\kappa - de^{\tilde{A}(t,x)}) \\ &+ \frac{\chi \mu(\sqrt{\lambda - \kappa^2} + \kappa)}{\lambda - \kappa^2} \phi_\kappa \left(\kappa e^{-\kappa x} + d(\kappa+\varepsilon) e^{\tilde{A}(t,x)}\right) \\ &+ (b(t) - \chi \mu) e^{-2\kappa x} - d(b(t) - \chi \mu) \left(\phi_\kappa + \phi_{\kappa,\varepsilon,d,A}\right) e^{\tilde{A}(t,x)} \\ &\leq -\left(\frac{d\varepsilon}{2\kappa} \left(\underline{a} - \kappa(\kappa+\varepsilon)\right) e^{A(t) + (\kappa-\varepsilon)x} - (b(t) - \chi \mu) - \frac{\chi \mu(\lambda + \kappa(\sqrt{\lambda - \kappa^2} + \kappa)))}{\lambda - \kappa^2}\right) e^{-2\kappa x} \\ &- d\left((b(t) - \chi \mu) + \chi \mu \left(\frac{\lambda - (\kappa+\varepsilon)(\sqrt{\lambda - \kappa^2} + \kappa)}{\lambda - \kappa^2}\right)\right)\right) \phi_\kappa e^{\tilde{A}(t,x)} - d(b(t) - \chi \mu) \phi_{\kappa,\varepsilon,d,A} e^{\tilde{A}(t,x)} \\ &\leq -A_0 e^{-\|A\|_{\infty}} \left(d - \frac{A_1 e^{\|A\|_{\infty}}}{A_0}\right) e^{-2\kappa x} - dA_2 \phi_\kappa e^{\tilde{A}(t,x)} - d(b_{inf} - \chi \mu) \phi_{\kappa,\varepsilon,d,A} e^{\tilde{A}(t,x)} \end{aligned}$$

where A_0 , A_1 , and A_2 are given by (2.17), (2.18) and (2.19) respectively. Thus, (iii) follows.

(iv) Let
$$0 < \delta \leq \frac{b_{\inf} - \chi \mu \left(1 + \frac{a_{\sup}}{a_{\inf}} \right)}{(b_{\inf} - \chi \mu)(b_{\sup} - \chi \mu)}$$
. We have that

$$\mathcal{L}_{\kappa,\phi}(\delta) = -\left(a(t) - \chi \lambda \psi(\cdot, \cdot; \phi) - (b(t) - \chi \mu) \delta \right) \delta$$

$$\leq -\left(a_{\inf} - \frac{\chi \mu a_{\sup}}{b_{\inf} - \chi \mu} - (b_{\sup} - \chi \mu) \delta \right) \delta$$

$$= -a_{\inf}(b_{\sup} - \chi \mu) \left(\frac{b_{\inf} - \chi \mu \left(1 + \frac{a_{\sup}}{a_{\inf}} \right)}{(b_{\inf} - \chi \mu)(b_{\sup} - \chi \mu)} - \delta \right)$$

$$\leq 0.$$

(iv) then follows.

We recall from Lemma 2.4 (iv) that $\inf_{t\in\mathbb{R}}\phi_{\kappa,\varepsilon,d,A}(x^+_{\kappa,\varepsilon,d,A}(t),t) > 0$, where $x^+_{\kappa,\varepsilon,d,A}(t)$ is defined by (2.23). Moreover, Lemma 2.4 guarantees that for each given $t \in \mathbb{R}$ and every $\delta \in (0, \inf_{t\in\mathbb{R}}\phi_{\kappa,\varepsilon,d,A}(x^+_{\kappa,\varepsilon,d,A}(t)))$, there is a unique $x_{\kappa,\varepsilon,d,A}(t;\delta) \in (x^-_{\kappa,\varepsilon,d,A}(t), x^+_{\kappa,\varepsilon,d,A}(t))$ such that

$$\phi_{\kappa,\varepsilon,d,A}(x_{\kappa,\varepsilon,d,A}(t;\delta),t) = \delta_{\lambda}$$

Let
$$0 < \delta_0 < \min\{\frac{b_{\inf} - \chi \mu \left(1 + \frac{a_{\sup}}{a_{\inf}}\right)}{(b_{\inf} - \chi \mu)(b_{\sup} - \chi \mu)}, \inf_{t \in \mathbb{R}} \phi_{\kappa, \varepsilon, d, A}(x_{\kappa, \varepsilon, d, A}^+(t), t)\}$$
 be fixed and define

$$\phi_{\kappa}^-(x, t) = \begin{cases} \phi_{\kappa, \varepsilon, d, A}(x, t), & \text{if } x \ge x_{\kappa, \varepsilon, d, A}(t; \delta_0) \\ \delta_0, & \text{if } x \le x_{\kappa, \varepsilon, d, A}(t; \delta_0). \end{cases}$$
(2.29)

It should be noted that the function $\mathbb{R} \ni t \mapsto x_{\kappa,\varepsilon,d,A}(t;\delta_0)$ is Lipschitz continuous.

3 Existence of transition front solutions

In this section, we prove the existence of transition front solutions of (1.5). We suppose that **(H1)** holds, and $\mathcal{E}_{\kappa,\beta_0,M}$ is defined as in the previous section. The functions ϕ_{κ}^+ and ϕ_{κ}^- are given by (2.2) and (2.29), respectively. Our main idea to prove the existence of transition front solutions of (1.5) is to prove that there is $\phi(\cdot, \cdot) \in \mathcal{E}_{\kappa,\beta_0,M}$ such that $(U(x,t), V(x,t)) = (\phi(x,t), \psi(x,t;\phi))$ is an entire solution of (1.10) with $c(t) = c_{\kappa}(t)$ and that $(u(x,t), v(x,t)) = (U(x - \int_0^t c_{\kappa}(s)ds, t), V(x - \int_0^t c_{\kappa}(s)ds, t))$ is a transition front solution of (1.5). To do so, we first prove some lemmas.

For every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, $t_0 \in \mathbb{R}$ and $u_0 \in C^b_{\text{unif}}(\mathbb{R})$, let $\Phi(x,t;t_0,u_0,\phi)$, $x \in \mathbb{R}$, $t \ge t_0$ be the solution of

$$\begin{cases} \mathcal{L}_{\kappa,\phi}(\Phi) = 0, & x \in \mathbb{R}, \ t > t_0 \\ \Phi(x, t_0; t_0, u_0, \phi) = u_0(x), & \forall \ x \in \mathbb{R}, \end{cases}$$
(3.1)

where $\mathcal{L}_{\kappa,\phi}$ is given by (2.7) (the existence of $\Phi(x,t;t_0,u_0,\phi)$ follows from general semigroup theory).

Lemma 3.1. For every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$ and $t_0 \in \mathbb{R}$, let $\Phi(x,t;t_0,\phi_{\kappa}^+,\phi)$ be given by (3.1). Then the following hold.

(i) For any $t_2 < t_1$, $0 \le \Phi(x, t; t_2, \phi_{\kappa}^+, \phi) \le \Phi(x, t; t_1, \phi_{\kappa}^+, \phi) \le \phi_{\kappa}^+(x)$ for every $x \in \mathbb{R}$ and $t \ge t_1$.

(ii) For any $t_0 \in \mathbb{R}$, $\phi_{\kappa}^-(x,t) \leq \Phi(x,t;t_0,\phi_{\kappa}^+,\phi)$ for all $x \in \mathbb{R}$ and $t \geq t_0$, where $\phi_{\kappa,\varepsilon,d,A}$ is given by (2.21).

Proof. (i) Since $\Phi(x, t_0; t_0, \phi_{\kappa}^+, \phi) \leq \phi_{\kappa}(x)$ for every $x \in \mathbb{R}$, it follows from comparison principle for parabolic equations and Theorem 2.5 (i) that

$$\Phi(x,t;t_0,\phi_{\kappa}^+,\phi) \le \phi_{\kappa}(x), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \ge t_0.$$

On the other hand, since $\phi_{\kappa}^+(x) \leq \frac{a_{\sup}}{b_{\inf}-\chi\mu}$, it follows from Theorem 2.5 (ii) and comparison principle for parabolic equations that

$$\Phi(x,t;t_0,\phi_{\kappa}^+,\phi) \le \frac{a_{\sup}}{b_{\inf} - \chi\mu}, \quad \forall \ x \in \mathbb{R}, \ \forall \ t \ge t_0.$$

Thus

$$\Phi(x,t;t_0,\phi_{\kappa}^+,\phi) \le \phi_{\kappa}^+(x) = \Phi(x,t_0;t_0,\phi_{\kappa}^+,\phi) \quad \forall \ x \in \mathbb{R}, \ \forall \ t \ge t_0.$$

Hence for any $t_2 < t_1$,

$$\Phi(x, t_1; t_2, \phi_{\kappa}^+, \phi) \le \phi_{\kappa}^+(x) = \Phi(x, t_1; t_1, \phi_{\kappa}^+, \phi).$$

Then, by comparison principle for parabolic equations, we have

$$\Phi(x,t;t_2,\phi_{\kappa}^+,\phi) \le \Phi(x,t;t_1,\phi_{\kappa}^+,\phi) \quad \forall \ x \in \mathbb{R}, \ \forall \ t \ge t_1.$$

This completes the proof of (i).

(ii) Note that if $\phi_{\kappa,\varepsilon,d,A}(x,t) \leq 0$ then we are done. Observe from (i) and Lemma 2.4 (i) that $\phi_{\kappa,\varepsilon,d,A}(x_{\kappa,\varepsilon,d}^-(t),t) = 0 \leq \Phi(x_{\kappa,\varepsilon,d,A}^-(t),t;t_0,\phi_{\kappa}^+,\phi)$ for every $t \geq t_0$. Since $\phi_{\kappa,\varepsilon,d,A}(x,t_0) \leq \phi_{\kappa}^+(x)$ for every $x \geq x_{\kappa,\varepsilon,d,A}^-(t_0)$, hence it follows from Theorem 2.5 (iii) and comparison principle for parabolic equations that

$$\phi_{\kappa,\varepsilon,d,A}(x,t) \le \Phi(x,t;t_0,\phi_{\kappa}^+,\phi), \quad \forall \ x \ge x_{\kappa,\varepsilon,d,A}^-(t), \quad \forall t \ge t_0.$$

In particular, we have that

$$\delta_0 = \phi_{\kappa,\varepsilon,d,A}(x_{\kappa,\varepsilon,d,A}(t;\delta_0),t) \le \Phi(x_{\kappa,\varepsilon,d,A}(t;\delta_0),t;t_0,\phi_{\kappa}^+,\phi), \quad \forall t \ge t_0.$$

Observe that $\delta_0 \leq \phi_{\kappa}^+(x) = \phi(x, t_0; t_0, \phi_{\kappa}^+, \phi)$ for every $x \leq x_{\kappa,\varepsilon,d,A}(t; \delta_0), t \in \mathbb{R}$ and $t_0 \in \mathbb{R}$. We also note that Lemma 2.4 implies that $u(x, t) = \delta_0$ is a sub-solution of (2.7) on $(-\infty, x_{\kappa,\varepsilon,d,A}(\delta_0, t_0)]$. Hence, it follows from comparison principle for parabolic equations that

$$\delta_0 \le \Phi(x,t;t_0,\phi_{\kappa}^+,\phi), \quad \forall x \le x_{\kappa,\varepsilon,d,A}(\delta_0,t_0), \ \forall t \ge t_0.$$

This completes the proof of (ii).

It follows from Lemma 3.1 that for every $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, $\Phi(\cdot,\cdot;t_0,\phi_{\kappa}^+,\phi)$ is non-decreasing in $t_0 \in \mathbb{R}$. Then for each $\phi \in \mathcal{E}_{\kappa,\beta_0,M}$, there is $\Phi(x,t;\phi)$ such that

$$\Phi(x,t;\phi) := \lim_{t_0 \to -\infty} \Phi(x,t;t_0,\phi_{\kappa}^+,\phi) \quad \forall x,t \in \mathbb{R}.$$
(3.2)

Moreover, we have

$$\phi_{\kappa}^{+}(x) \ge \Phi(x,t;\phi) \ge \phi_{\kappa}^{-}(x,t), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}.$$
(3.3)

Hence, we introduce the following set,

$$\tilde{\mathcal{E}}_{\kappa,\beta_0,M} = \{ \phi \in \mathcal{E}_{\kappa,\beta_0,M} \, | \, \phi_{\kappa}^-(x,t) \le \phi(x,t) \le \phi_{\kappa}^+(x), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R} \}.$$
(3.4)

Lemma 3.2. For every $\phi \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$, $\Phi(\cdot,\cdot;\phi) \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ when M is sufficiently large and satisfies the parabolic equation

$$\partial_t \Phi = \partial_{xx} \Phi + (c_\kappa(t) - \chi \partial_x \psi(\cdot, \cdot; \phi)) \partial_x \Phi + (a(t) - \lambda \chi \psi(\cdot, \cdot; \phi) - (b(t) - \chi \mu) \Phi) \Phi, \ \forall (x, t) \in \mathbb{R} \times \mathbb{R}.$$
(3.5)

Proof. We will apply the similar arguments as those in [41, Lemma 3.5] to prove this lemma.

First of all, let $\{T(t)\}_{t\geq 0}$ denotes the C_0 -semigroup generated by $\Delta - I$ on $C^b_{\text{unif}}(\mathbb{R}^N)$. Using the fact that $\partial_{xx}\psi(\cdot,\cdot,\phi) = \lambda\psi(\cdot,\cdot,\phi) - \mu\phi$, the variation of constant formula yields that

$$\Phi(x,t;t_{0},\phi_{\kappa}^{+},\phi) = T(t-t_{0})\phi_{\kappa}^{+} + \int_{t_{0}}^{t} T(t-s)((c_{\kappa}(s) - \chi\partial_{x}\psi(\cdot,s,\phi))\partial_{x}\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi))ds \\ + \int_{t_{0}}^{t} T(t-s)((a(s) + 1 - \lambda\chi\psi(\cdot,s;\phi) - (b(s) - \chi\mu)\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi))\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi))ds \\ = \underbrace{T(t-t_{0})\phi_{\kappa}^{+}}_{I_{t_{0}}^{t}(\cdot,t)} + \underbrace{\int_{t_{0}}^{t} T(t-s)\partial_{x}((c_{\kappa}(s) - \chi\partial_{x}\psi(\cdot,s,\phi))\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi))ds}_{I_{t_{0}}^{2}(\cdot,t)} \\ + \underbrace{\int_{t_{0}}^{t} T(t-s)(a(s) + 1 - \chi\mu\phi(\cdot,s))\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)ds}_{I_{t_{0}}^{2}(\cdot,t)} \\ - \underbrace{\int_{t_{0}}^{t} T(t-s)(b(s) - \chi\mu)\Phi^{2}(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)ds}_{I_{t_{0}}^{4}(\cdot,t)} \end{cases}$$
(3.6)

Next, choose $\frac{\beta_0}{2} < \beta < \frac{1}{2} - \beta_0$ (such β exists for $0 < \beta_0 < \frac{1}{3}$). Let X^{β} denotes the fractional power space associated with $\Delta - I$ on $C^b_{\text{unif}}(\mathbb{R}^N)$. We claim that there is a constant \tilde{C}_{β} independent of t, u, and t_0 such that

$$\|\Phi(\cdot, t; t_0, \phi_{\kappa}^+, \phi)\|_{X^{\beta}} \le \tilde{C}_{\beta} \Big(\frac{e^{-(t-t_0)}}{(t-t_0)^{\beta}} + 1\Big), \quad \forall t > t_0.$$
(3.7)

In fact, by [40, Lemma 3.2], there is a constant $C_{\beta} > 0$ such that

$$\|I_{t_0}^1(\cdot,t)\|_{X^{\beta}} \le C_{\beta}(t-t_0)^{-\beta} e^{-(t-t_0)} \|\phi_{\kappa}^+\|_{\infty} = \frac{C_{\beta} e^{-(t-t_0)} a_{\sup}}{(t-t_0)^{\beta} (b_{\inf} - \chi \mu)},$$
(3.8)

$$\|I_{t_{0}}^{2}(\cdot,t)\|_{X^{\beta}} \leq C_{\beta} \int_{t_{0}}^{t} (t-s)^{-(\frac{1}{2}+\beta)} e^{-(t-s)} (\|c_{\kappa}\|_{\infty} + \chi \|\partial_{x}\psi(\cdot,s,\phi)\|_{\infty}) \|\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)\|_{\infty} ds$$

$$\leq C_{\beta} \int_{t_{0}}^{t} (t-s)^{-(\frac{1}{2}+\beta)} e^{-(t-s)} (\frac{a_{\sup}+\kappa^{2}}{\kappa} + \chi \mu \|\phi(\cdot,s)\|_{\infty}) \|\phi_{\kappa}^{+}\|_{\infty} ds$$

$$\leq C_{\beta} \Big(\frac{a_{\sup}+\kappa^{2}}{\kappa} + \frac{\chi \mu a_{\sup}}{b_{\inf}-\chi \mu}\Big) \frac{a_{\sup}\Gamma(\frac{1}{2}-\beta)}{b_{\inf}-\chi \mu}, \qquad (3.9)$$

$$\|I_{t_{0}}^{3}(\cdot,t)\|_{X^{\beta}} \leq C_{\beta} \int_{t_{0}}^{t} (t-s)^{-\beta} e^{-(t-s)} \Big(a_{\sup} + 1 + \chi \mu \|\phi(\cdot,s)\|_{\infty} \Big) \|\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)\|_{\infty} ds$$

$$\leq C_{\beta} \Big(a_{\sup} + 1 + \frac{\chi \mu a_{\sup}}{b_{\inf} - \chi \mu} \Big) \frac{a_{\sup} \Gamma(1-\beta)}{b_{\inf} - \chi \mu},$$
(3.10)

and

$$\|I_{t_0}^4(\cdot, t)\|_{X^\beta} \le C_\beta (b_{\sup} - \chi\mu) \frac{a_{\sup}^2 \Gamma(1-\beta)}{(b_{\inf} - \chi\mu)^2}.$$
(3.11)

The claim then follows from (3.8)–(3.11).

Now, we claim that there is a constant $\bar{C}_{\beta} > 0$ independent of t_0 , ϕ , and t such that

$$\|\Phi(\cdot, t+h; t_0, \phi_{\kappa}^+, \phi) - \Phi(\cdot, t; t_0, \phi_{\kappa}^+, \phi)\|_{X^{\beta}} \le \bar{C}_{\beta}(h^{1-\beta} + h^{\frac{1}{2}-\beta}) \Big(\frac{e^{-(t-t_0)}}{(t-t_0)^{\beta}} + 1\Big), \ \forall h > 0, \ t > t_0.$$

$$(3.12)$$

In fact, for h > 0 and $t > t_0$, we have

$$\|I_{t_0}^1(\cdot, t+h) - I_{t_0}^1(\cdot, t)\|_{X^\beta} = \|(T(h) - I)T(t+n)\phi_\kappa^+\|_\infty \le \frac{C_\beta h^\beta e^{-(t-t_0)}}{(t-t_0)^\beta} \|\phi_\kappa^+\|_\infty,$$
(3.13)

$$\begin{split} \|I_{t_{0}}^{2}(\cdot,t+h) - I_{t_{0}}^{2}(\cdot,t)\|_{X^{\beta}} \\ &\leq C_{\beta}h^{\beta}\int_{t_{0}}^{t} \frac{e^{-(t-s)}}{(t-s)^{\frac{1}{2}+\beta}} \Big(\frac{a_{\sup}+\kappa^{2}}{\kappa} + \chi\mu\|\phi(\cdot,s)\|_{\infty}\Big)\|\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)\|_{\infty}ds \\ &+ C_{\beta}\int_{t}^{t+h} \frac{e^{-(t+h-s)}}{(t+h-s)^{\frac{1}{2}+\beta}} \Big(\frac{a_{\sup}+\kappa^{2}}{\kappa} + \chi\mu\|\phi(\cdot,s)\|_{\infty}\Big)\|\Phi(\cdot,s;t_{0},\phi_{\kappa}^{+},\phi)\|_{\infty}ds \\ &\leq C_{\beta}\Big(h^{\beta}\Gamma(\frac{1}{2}-\beta) + \frac{h^{\frac{1}{2}-\beta}}{\frac{1}{2}-\beta}\Big)\Big(\frac{a_{\sup}+\kappa^{2}}{\kappa} + \frac{\chi\mu a_{\sup}}{b_{\inf}-\chi\mu}\Big)\frac{a_{\sup}}{b_{\inf}-\chi\mu}, \end{split}$$
(3.14)

$$\|I_{t_0}^3(\cdot, t+h) - I_{t_0}^3(\cdot, t)\|_{X^{\beta}} \le C_{\beta} \Big(h^{\beta} \Gamma(1-\beta) + \frac{h^{1-\beta}}{1-\beta}\Big) \Big(a_{\sup} + 1 + \frac{\chi \mu a_{\sup}}{b_{\inf} - \chi \mu}\Big) \frac{a_{\sup}}{b_{\inf} - \chi \mu}, \quad (3.15)$$

and

$$\|I_{t_0}^4(\cdot, t+h) - I_{t_0}^4(\cdot, t)\|_{X^\beta} \le C_\beta \Big(h^\beta \Gamma(1-\beta) + \frac{h^{1-\beta}}{1-\beta}\Big) \Big(b_{\sup} - \chi\mu\Big) \frac{a_{\sup}}{b_{\inf} - \chi\mu}.$$
 (3.16)

(3.12) then follows from (3.13)-(3.16).

Letting $t_0 \to -\infty$ in (3.12), we obtain

$$\|\Phi(\cdot, t+h;\phi) - \Phi(\cdot, t;\phi)\|_{X^{\beta}} \le \bar{C}_{\beta}(h^{1-\beta} + h^{\frac{1}{2}-\beta}), \ \forall h > 0, \ t \in \mathbb{R}.$$
 (3.17)

Note that X^{β} is continuously embedded in $C_{\text{unif}}^{2\nu}(\mathbb{R})$ for $0 < \nu < 2\beta$ ([17]). This together with (3.7) implies that there is $\hat{C}_{\beta} > 0$ such that

$$|\Phi(x+h,t;\phi) - \Phi(x,t;\phi)| \le \hat{C}_{\beta}|h|^{\beta_0} \quad \forall x,t,h \in \mathbb{R}, \ |h| \le 1.$$
(3.18)

By (3.3), (3.17), and (3.18), we have that $|\Phi(\cdot, \cdot; \phi) \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ for every $\phi \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ provided that $M \geq \max\{2\tilde{C}_{\beta}, \hat{C}_{\beta}\}.$

Finally, by (3.7), (3.12), Arzela-Ascoli Theorem, and [12, Chapter 3 - Theorem 15], we have that $\Phi(x,t;t_0,\phi) \to \Phi(x,t;\phi)$ in $C^{2,1}(\bar{D})$ as $t_0 \to -\infty$ for any bounded domain $D \subset \mathbb{R} \times \mathbb{R}$. This implies that $\Phi(\cdot, \cdot, \phi)$ satisfies (3.5).

In the following, we fix $\frac{\beta_0}{2} < \beta < \frac{1}{2} - \beta_0$ and $M \ge \max\{2\tilde{C}_{\beta}, \hat{C}_{\beta}\}$.

Lemma 3.3. For every $\phi \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$, the function $\Phi(\cdot,\cdot;\phi)$ given by (3.2) is the only solution of (3.5) in $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$.

Proof. We apply the similar arguments as those in [41, Lemma 3.6] to prove this lemma.

Let $\phi \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ be fixed and $U_1(x,t), U_2(x,t)$ be solutions of (3.5) in $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$. Let $\tau > 0$ be given. Since $\phi_{\kappa}^-(x,t) \leq U_i(x,t) \leq \phi_{\kappa}^+(x)$ for every $x \in \mathbb{R}$ and $t \in \mathbb{R}$, and $\lim_{x\to\infty} \frac{\phi_{\kappa}^+(x)}{\phi_{\kappa}^-(x,t)} = 1$ uniformly in $t \in \mathbb{R}$, then there is $\bar{x}_{\tau} \gg 1$ such that

$$\frac{1}{1+\tau} \le \frac{U_1(x,t)}{U_2(x,t)} \le 1+\tau, \quad \forall \ x \ge \bar{x}_{\tau}, \ \forall \ t \in \mathbb{R}.$$
(3.19)

Note that $\inf\{\phi_{\kappa}^{-}(x,t) \mid x \leq R, t \in \mathbb{R}\} > 0$ for every $R \in \mathbb{R}$ and $\phi_{\kappa}^{+}(x) \leq \frac{a_{\sup}}{b_{\inf} - \chi \mu}$. This together with (3.19) and the fact that $\phi_{\kappa}^{-}(x,t) \leq U_{i}(x,t) \leq \phi_{\kappa}^{+}(x)$ for every $(x,t) \in \mathbb{R} \times \mathbb{R}$ imply that there is $\alpha \in \mathbb{R}$ such that

$$\alpha := \inf\{\tilde{\alpha} \mid \tilde{\alpha} \ge 1 \text{ and } \frac{1}{\tilde{\alpha}} \le \frac{U_1(x,t)}{U_2(x,t)} \le \tilde{\alpha}, \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}\}.$$
(3.20)

It is clear $\alpha \ge 1$, and to prove the lemma, it suffices to prove that $\alpha = 1$. We prove this in the following by contradiction.

Assume that $\alpha > 1$. It is clear from the definition of α that

$$\begin{cases} U_1(x,t) \le \alpha U_2(x,t), & \forall x \in \mathbb{R}, t \in \mathbb{R}, \text{ and} \\ U_2(x,t) \le \alpha U_1(x,t), & \forall x \in \mathbb{R}, t \in \mathbb{R}. \end{cases}$$
(3.21)

Let $\tau \in (0, \alpha - 1)$ and \bar{x}_{τ} be given by (3.19). Note that

$$U_i(x,t) = \Phi(x,t;t_0,U_i(\cdot,t_0),\phi), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \ge t_0,$$

where $\Phi(x,t;t_0,U_i,\phi)$ is given by (3.1) for each i = 1,2. Since $\alpha > 1$ and $U_i(x,t) > 0$ for every $(x,t) \in \mathbb{R} \times \mathbb{R}$, then

$$\mathcal{L}_{\kappa,\phi}(\alpha U_i) = (\alpha - 1)(b(t) - \chi\mu)U_i(\alpha U_i) > 0, \quad \forall x \in \mathbb{R}, \ t \in \mathbb{R}, \ i = 1, 2.$$
(3.22)

Hence, since $\alpha > 1$, comparison principle for parabolic equations and (3.21) imply that

$$\begin{cases} U_1(x,t+t_0) < \Phi(x,t+t_0;t_0,\alpha U_2,\phi) < \alpha U_2(x,t+t_0), & \forall \ x \in \mathbb{R}, \ \forall t > 0, \ \forall t_0 \in \mathbb{R} \\ U_2(x,t+t_0) < \Phi(x,t+t_0;t_0,\alpha U_1,\phi) < \alpha U_1(x,t+t_0), & \forall \ x \in \mathbb{R}, \ \forall t > 0, \ \forall t_0 \in \mathbb{R}. \end{cases}$$
(3.23)

Note that (3.23) implies that

$$\min_{\frac{1}{2} \le t \le 1} (\alpha U_i(\bar{x}_\tau, t+t_0) - \Phi(\bar{x}_\tau, t+t_0; t_0, \alpha U_i, \phi)) > 0, \quad \forall t_0 \in \mathbb{R}, \quad i = 1, 2.$$

We claim that for each i = 1, 2, we have that

$$\inf_{t_0 \in \mathbb{R}} \min_{\frac{1}{2} \le t \le 1} (\alpha U_i(\bar{x}_\tau, t+t_0) - \Phi(\bar{x}_\tau, t+t_0; t_0, \alpha U_i, \phi)) > 0.$$
(3.24)

Assume that (3.24) does not hold. Without loss of generality, we may suppose that there are sequences $\{t_{0n}\}_{n\geq 1} \subset \mathbb{R}$ and $\{t_n\}_{n\geq 1} \subset [\frac{1}{2}, 1]$ such that

$$\lim_{n \to \infty} \alpha U_1(\bar{x}_\tau, t_n + t_{0n}) - \Phi(\bar{x}_\tau, t_n + t_{0n}; t_{0n}, \alpha U_1, \phi) = 0.$$
(3.25)

For each $n \geq 1$, let $u_n(x,t) = U_1(x+\tilde{x}_{\tau},t+t_n+t_{0n})$ and $\tilde{u}_n(x,t) = \Phi(x+\tilde{x}_{\tau},t+t_n+t_{0n};t_{0n},\alpha U_1,\phi)$ for $x \in \mathbb{R}$ and $t \geq -\frac{1}{2}$. Hence, by a priori estimates for parabolic equations and Arzela-Ascoli, we may suppose that there exist $u^*(x,t), \tilde{u}^*(x,t) \in C^{2,1}(\mathbb{R} \times [-\frac{1}{2},\infty)), \phi^* \in C^{\delta}(\mathbb{R} \times \mathbb{R})$, and $a^*(t), b^*(t) \in C^{\delta}(\mathbb{R})$ such that

$$(u_n(x,t), \tilde{u}_n(x,t), \phi(t+t_n+t_{0n})) \to (u^*(x,t), \tilde{u}^*(x,t), \phi^*(x,t))$$

and

$$(a(t + t_n + t_{0n}), b(t + t_n + t_{0n})) \rightarrow (a^*(t), b^*(t))$$

as $n \to \infty$ in the compact-open topology. Moreover, $u^*(x,t)$ and $\tilde{u}^*(x,t)$ solve the PDE

$$u_{t} = u_{xx} + (c^{*}(t) - \chi \partial_{x} \psi(\cdot, \cdot; \phi^{*})) u_{x} + (a^{*}(t) - \chi \lambda \psi(\cdot, \cdot; \phi^{*}) - (b^{*}(t) - \chi \mu) u) u, \ x \in \mathbb{R}, \ t \ge -\frac{1}{2}$$
(3.26)

with $c^*(t) = \frac{\kappa^2 + a^*(t)}{\kappa}$. Observe that

$$0 < \inf_{s \in \mathbb{R}} \phi_{\kappa}^{-}(x, s) \le \min\{u^{*}(x, t), \tilde{u}^{*}(x, t)\}, \quad \forall x \ll -1, \ t \ge -\frac{1}{2}$$

and that

$$\tilde{u}^*(x,t) \le \alpha u^*(x,t), \quad \forall x \in \mathbb{R}, \ t \ge -\frac{1}{2}$$

Then $0 < u^*(x,t)$ for all $x \in \mathbb{R}$ and $t > \frac{1}{2}$. Multiplying (3.26) by α , we obtain that

$$(\alpha u^{*})_{t} > u_{xx} + (c^{*}(t) - \chi \partial_{x} \psi(\cdot, \cdot))(\alpha u^{*})_{x} + (a^{*}(t) - \chi \lambda \psi(\cdot, \cdot; \phi^{*}) - (b^{*}(t) - \chi \mu)(\alpha u^{*}))(\alpha u^{*}).$$

Hence, comparison principle for parabolic equations yields that

$$u^*(x,t) < \alpha \tilde{u}^*(x,t), \quad \forall \ x \in \mathbb{R}, \ \forall \ t > -\frac{1}{2}$$

In particular

$$u^*(0,0) < \alpha \tilde{u}^*(0,0). \tag{3.27}$$

On the other hand, (3.25) implies that

$$u^*(0,0) = \alpha \tilde{u}^*(0,0),$$

which contradicts to (3.27). Thus (3.24) holds.

We note from (3.22) that

$$\mathcal{L}_{\kappa,\phi}(\alpha U_i) = (\alpha - 1)(b(t) - \chi\mu)U_i(\alpha U_i) \geq (\alpha - 1)(b_{\inf} - \chi\mu)m_{\tau}(\alpha U_i), \quad \forall \ x \leq \bar{x}_{\tau}, \ \forall \ t \in \mathbb{R}.$$
(3.28)

where $m_{\tau} = \inf\{\phi_{\kappa}^{-}(x,t) : x \leq \bar{x}_{\tau}, t \in \mathbb{R}\} > 0.$

Let $0 < \tilde{\delta} \ll 1$ such that

$$\begin{cases} \tilde{\delta} + (b_{\sup} - \chi\mu)(1 - e^{-\tilde{\delta}})e^{\tilde{\delta}}\frac{\alpha a_{\sup}}{b_{\inf} - \chi\mu} \leq (\alpha - 1)(b_{\inf} - \chi\mu)m_{\tau},\\ \inf_{t_0 \in \mathbb{R}} \min_{\frac{1}{2} \leq t \leq 1}(\alpha U_i(\bar{x}_{\tau}, t + t_0) - \Phi(\bar{x}_{\tau}, t + t_0; t_0, \alpha U_i, \phi)) > \frac{(e^{\tilde{\delta}} - 1)\alpha a_{\sup}}{b_{\inf} - \chi\mu}\\ (1 + \tau)e^{\tilde{\delta}} < \alpha. \end{cases}$$

Let $t_0 \in \mathbb{R}$ be given. For every $t \geq 0$ and i = 1, 2, let $W_i(x, t) = e^{\tilde{\delta}t} \Phi(x, t + t_0; t_0, \alpha U_i(\cdot, t_0), \phi)$. We have

$$\mathcal{L}_{\kappa,\phi}(W_i)(x,t) \leq \left(\tilde{\delta} + (b(t) - \chi\mu)(1 - e^{-\tilde{\delta}t})e^{\tilde{\delta}t}W_i(x,t)\right)W_i(x,t)$$

$$\leq \left(\tilde{\delta} + (b_{\sup} - \chi\mu)(1 - e^{-\tilde{\delta}t})e^{\tilde{\delta}t}\frac{\alpha a_{\sup}}{b_{\inf} - \chi\mu}\right)W_i(x,t)$$

$$\leq (\alpha - 1)(b_{\inf} - \chi\mu)m_{\tau}W_i(x,t), \quad \forall 0 \leq t \leq 1.$$
(3.29)

Since $\Phi(x, t + t_0; t_0, \alpha U_i, \phi) \leq \alpha U_i(x, t + t_0) \leq \frac{\alpha a_{\sup}}{b_{\inf} - \chi \mu}$ for every $t \geq 0, t_0 \in \mathbb{R}$ and $x \in \mathbb{R}$, then

$$\frac{(e^{\delta}-1)\alpha a_{\sup}}{b_{\inf}-\chi\mu} + \Phi(x,t+t_0;t_0,\alpha U_i,\phi) \ge e^{\tilde{\delta}t}\Phi(x,t+t_0;t_0,\alpha U_i,\phi), \quad \forall \ x \in \mathbb{R}, \ t_0 \in \mathbb{R}, \ 0 \le t \le 1.$$

Therefore, it follows from the choice of $\tilde{\delta}$ that

$$W_i(x,t) = e^{\tilde{\delta}t} \Phi(\tilde{x}_{\tau}, t+t_0; t_0, \alpha U_i, \phi) \le \alpha U_i(\tilde{x}_{\tau}, t+t_0), \qquad \forall t \in [\frac{1}{2}, 1], \ \forall t_0 \in \mathbb{R}.$$

Combine this with (3.28), (3.29), the comparison principle for parabolic equations yield that

$$e^{\tilde{\delta}t}\Phi(x,t+t_0;t_0;\alpha U_i,\phi) \le \alpha U_i(x,t+t_0), \quad \forall x \le \tilde{x}_{\tau}, \ \forall t \in [\frac{1}{2},1], \ \forall t_0 \in \mathbb{R}.$$

This combined with (3.23) give

$$U_1(x,t) \le \alpha e^{-\tilde{\delta}} U_2(x,t)$$
 and $U_2(x,t) \le \alpha e^{-\tilde{\delta}} U_1(x,t), \quad x \le \tilde{x}_{\tau}, \ t \in \mathbb{R}.$

Which is together with (3.19) and the fact that $(1 + \tau)e^{\tilde{\delta}} \leq \alpha$ yield that

$$\frac{1}{\alpha e^{-\tilde{\delta}}} \le \frac{U_1(x,t)}{U_2(x,t)} \le \alpha e^{-\tilde{\delta}}, \ \forall \ t \in \mathbb{R}, \forall \ x \in \mathbb{R}.$$

Hence by definition of α , we must have $\alpha \leq \alpha e^{-\tilde{\delta}}$, which is impossible. Thus $\alpha = 1$ and the lemma is proved.

Now we present the proof of Theorem 1.1

Proof of Theorem 1.1(1). For any given $0 < \underline{c} < c_{\chi}^*$, let $0 < \kappa < \kappa_{\chi}^*$ be such that $\underline{c} = \frac{\underline{a} + \kappa^2}{\kappa}$. We prove that (1.5) has a transition front solution $(u(x,t), v(x,t)) = (U(x - C_{\kappa}(t), t), V(x - C_{\kappa}(t), t))$ with $C_{\kappa}(t) = \int_0^t c_{\kappa}(s) ds$ and $c_{\kappa}(t) = \frac{\underline{a}(t) + \kappa^2}{\kappa}$. To prove this, we prove the following claim.

Claim 1. The mapping $\Phi : \tilde{\mathcal{E}}_{\kappa,\beta_0,M} \to \tilde{\mathcal{E}}_{\kappa,\beta_0,M}, \phi \mapsto \Phi(\cdot,\cdot,\phi)$, has a fixed point ϕ^* satisfying $\Phi(\cdot,\cdot,\phi^*) = \phi^*$. Furthermore, any element $\phi^* \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ with the property $\Phi(\cdot,\cdot,\phi^*) = \phi^*$ also satisfies

$$\lim_{x \to -\infty} |\phi^*(x,t) - u^*(t)| = 0 \quad and \quad \lim_{x \to \infty} \frac{\phi^*(x,t)}{e^{-\kappa x}} = 1, \quad uniformly \ in \ t.$$

where $u^*(t)$ is the only positive entire solution of (1.3).

Assume that the claim holds. Let $U(x,t) = \phi^*(x,t)$ and $V(x,t) = \Psi(x,t;U)$. Then

$$(u(x,t),v(x,t)) = (U(x - C_{\kappa}(t),t),V(x - C_{\kappa}(t),t))$$

where $C_{\kappa}(t) = \int_0^t c_{\kappa}(s) ds$ and $c_{\kappa}(t) = \frac{a(t)+\kappa^2}{\kappa}$, is a transition front solution of (1.5) connecting (0,0) and $(u^*(t), v^*(t))$ satisfying (1.14). In the following, we prove that the claim holds.

We first prove that the mapping Φ has a fixed point. To this end, let $C^b_{\text{unif}}(\mathbb{R} \times \mathbb{R})$ be endowed with the compact-open topology, that is, a sequence of elements of $C^b_{\text{unif}}(\mathbb{R} \times \mathbb{R})$ converges if and only it converges uniformly on every compact subsets of $\mathbb{R} \times \mathbb{R}$. It is clear that $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ is a compact and convex subset of $C^b_{\text{unif}}(\mathbb{R} \times \mathbb{R})$ in the compact-open topology. To show that the mapping Φ has a fixed point, it is then enough to show that Φ is continuous. We show this in the following.

Let $\{\phi_n\}_{n\geq 1}$ and ϕ be elements of $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ such that $\phi_n(x,t) \to \phi(x,t)$ as $n \to \infty$ uniformly on every compact subsets of $\mathbb{R} \times \mathbb{R}$. Since $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ is compact, there is a subsequence $\{n_k\}$ of $\{n\}$ and a function $\Phi(\cdot, \cdot) \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ such that $\Phi(x,t;\phi_{n_k}) \to \Phi(x,t)$ as $n_k \to \infty$ on every compact subset of $\mathbb{R} \times \mathbb{R}$. Since $\phi_{n_k}(x,t) \to \phi(x,t)$ as $n_k \to \infty$ on very compact subset of $\mathbb{R} \times \mathbb{R}$, it follows from Lemma 2.11 (ii) that

$$\psi(x,t;\phi_{n_k}) \to \psi(x,t;\phi) \quad \text{and} \quad \partial_x \psi(x,t;\phi_{n_2}) \to \partial_x \psi(x,t;\phi) \quad \text{as} \quad n_k \to \infty,$$

uniformly on every compact subset of $\mathbb{R} \times \mathbb{R}$, where $\psi(\cdot, \cdot; \phi)$ is given by (2.9). Therefore, it follows from [12, Chapter 3 Theorem 3] that $\Phi(x, t) \in C^{2,1}(\mathbb{R} \times \mathbb{R})$ and satisfies

$$\Phi_t = \Phi_{xx} + (c_{\kappa}(t) - \psi_x(x,t;\phi))\Phi_x + (a(t) - \chi\lambda\psi(x,t;\phi) - (b(t) - \chi\mu)\Phi)\Phi, \ x \in \mathbb{R}, \ t \in \mathbb{R}.$$

Therefore, it follows from Lemma 3.3 that $\Phi(x,t) = \Phi(x,t;\phi)$ for every $(x,t) \in \mathbb{R} \times \mathbb{R}$. Hence the mapping Φ is continuous, and by Schauder's fixed point theorem, there is $\phi^* \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ such that

$$\Phi(x,t;\phi^*) = \phi^*(x,t) \ \forall \ (x,t) \in \mathbb{R} \times \mathbb{R}.$$

Next, we prove (1.16). Suppose that $\phi^* = \Phi(\cdot, \cdot, \phi^*)$. Since

$$\lim_{x \to \infty} \frac{\phi_{\kappa}^+(x)}{\phi_{\kappa}^-(x,t)} = 1,$$

uniformly in $t \in \mathbb{R}$ and $\phi_{\kappa}^{-}(x,t) \leq \phi^{*}(x,t) \leq \phi_{\kappa}^{+}(x)$, for every $x \in \mathbb{R}$ and $t \in \mathbb{R}$, we get that

$$\lim_{x \to \infty} \frac{\phi^*(x,t)}{e^{-\kappa x}} = 1, \quad \text{uniformly in t.}$$

It remains to show that

$$\lim_{x \to -\infty} |\phi^*(x,t) - u^*(t)| = 0, \quad \text{uniformly in } t, \tag{3.30}$$

where $u^*(t)$ is given by (1.3).

Suppose by contradiction that (3.30) does not hold. Then, there exist a sequence of real numbers $\{x_n\}_{n\geq 1}$ with $\lim_{n\to\infty} x_n = -\infty$, a sequence of real numbers $\{t_n\}_{n\geq 1}$, and a positive real number $\delta > 0$ such that

$$\delta \le \inf_{n \ge 1} |\phi^*(x_n, t_n) - u^*(t_n)|.$$
(3.31)

Let $\psi^*(x,t) = \psi(x,t;\phi^*)$. Hence $(u(x,t),v(x,t)) := (\phi^*(x - \int_0^t c_\kappa(s)ds,t),\psi^*(x - \int_0^t c_\kappa(s)ds,t))$ solves (1.5).

For each $n \ge 1, x \in \mathbb{R}$, and $t \in \mathbb{R}$, let

$$u_n(x,t) = u(x+x_n + \int_0^{t_n} c_\kappa(s)ds, t+t_n), \quad u_n^*(t) = u^*(t+t_n), \quad a_n(t) = a(t+t_n), \quad b_n(t) = b(t+t_n),$$

and $v_n = \mu(\lambda I - \Delta)^{-1} u_n$. Note (u_n, v_n) solves

$$\begin{cases} \partial_t u_n = \partial_{xx} u_n - \chi \partial_x (u_n \partial_x v_n) + (a_n(t) - b_n(t) u_n) u_n, & x \in \mathbb{R}, \ t \in \mathbb{R} \\ 0 = \partial_{xx} v_n - \lambda v_n + \mu u_n, & x \in \mathbb{R}, \ t \in \mathbb{R}, \end{cases}$$

and

$$\frac{d}{dt}u_n^* = (a_n(t) - b_n(t)u_n)u_n, \quad t \in \mathbb{R}.$$

Hence, up to a subsequence, using similar arguments as those used in the proof of Lemma 3.2, we may assume that there exist $(\tilde{u}, \tilde{v}) \in C^{2,1}(\mathbb{R} \times \mathbb{R}), \ \tilde{u}^* \in C^1(\mathbb{R}), \ \tilde{a}, \tilde{b} \in C^{\nu_0}_{\text{unif}}(\mathbb{R})$ such that $(u_n, v_n) \to (\tilde{u}, \tilde{v}), \ u_n^* \to \tilde{u}^*, \ a_n \to \tilde{a}, \text{ and } b_n \to \tilde{b} \text{ as } n \to \infty$ locally uniformly. Furthermore, it holds that

$$\begin{cases} \partial_t \tilde{u} = \partial_{xx} \tilde{u} - \chi \partial_x (\tilde{u} \partial_x \tilde{v}) + (\tilde{a}(t) - \tilde{b}(t) \tilde{u}) \tilde{u}, & x \in \mathbb{R}, \ t \in \mathbb{R} \\ 0 = \partial_{xx} \tilde{v} - \lambda \tilde{v} + \mu \tilde{u}, & x \in \mathbb{R}, \ t \in \mathbb{R}, \end{cases}$$
(3.32)

and

$$\frac{d}{dt}\tilde{u}^* = (\tilde{a}(t) - \tilde{b}(t)\tilde{u}^*)\tilde{u}^*, \quad t \in \mathbb{R}.$$
(3.33)

It is clear that $0 < \inf_{t \in \mathbb{R}} \tilde{u}^*(t) \le \sup_{t \in \mathbb{R}} \tilde{u}^*(t) < \infty$. Using the fact that $\inf_{t \in \mathbb{R}} x_{\kappa,\varepsilon,d,A}(t;\delta_0) \ge \inf_{t \in \mathbb{R}} x_{\kappa,\varepsilon,d,A}^-(t) > 0$, it follows from (2.29) that

$$u_n(x,t) = \phi^*(x + x_n - \int_{t_n}^{t_n+t} c_\kappa(s) ds, t + t_n)$$

$$\geq \phi_\kappa^-(x + x_n - \int_{t_n}^{t_n+t} c_\kappa(s) ds, t + t_n)$$

$$\geq \delta_0$$

whenever $x + x_n - \int_{t_n}^{t_n+t} c_{\kappa}(s) ds \leq 0$. Since $x_n \to -\infty$ as $n \to \infty$, and $\|c_{\kappa}\|_{\infty} \leq \frac{a_{\sup} + \kappa^2}{\kappa}$, then

$$0 < \delta_0 \le \tilde{u}(x, t) \le \frac{a_{\sup}}{b_{\inf} - \chi \mu}, \quad \forall x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}.$$

Hence (\tilde{u}, \tilde{v}) is a positive entire solution of (3.32). But, [46, Theorem 1.4] implies that $(\tilde{u}^*, \tilde{v}^*)$, where $\tilde{v}^* = \mu(\lambda I - \Delta)^{-1}\tilde{u}^*$, is the only strictly positive entire solution (3.32), when $0 < \mu\chi \leq \frac{b_{inf}}{2}$. Thus we must have

$$\tilde{u}(x,t) = \tilde{u}^*(t), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R}.$$
(3.34)

But inequality (3.31) gives that

$$0 < \delta \le |\tilde{u}(0,0) - \tilde{u}^*(0)|.$$

This contradicts to (3.34). Therefore, Claim 1 holds and Theorem 1.1(1) is thus proved.

Proof of Theorem 1.1(2). For given $c > c_{\chi}^*$, let $\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{\hat{a}}\})$ $(\hat{a} = \frac{1}{T} \int_0^T a(t) dt)$ be such that $c = \frac{\hat{a} + \kappa^2}{\kappa}$. To prove Theorem 1.1(2), we first change the set $\mathcal{E}_{\kappa,\beta_0,M}$ to

$$\mathcal{E}_{\kappa,\beta_{0},M}^{T} = \{ \phi \in C(\mathbb{R}, C_{\text{unif}}^{b}(\mathbb{R})) \mid 0 \leq \phi(x,t) \leq \phi_{\kappa}^{+}(x), \ \phi(x,t+T) = \phi(x,t) \\ |\phi(x+h,t) - \phi(x,t)| \leq M |h|^{\beta_{0}}, \text{ and} \\ |\phi(x,t+h) - \phi(x,t)| \leq M |h|^{\beta_{0}} \ \forall x,t,h \in \mathbb{R}, \ |h| \leq 1 \},$$
(3.35)

and change $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ to

$$\tilde{\mathcal{E}}^{T}_{\kappa,\beta_{0},M} = \{ \phi \in \mathcal{E}^{1}_{\kappa,\beta_{0},M} \, | \, \phi^{-}_{\kappa}(x,t) \le \phi(x,t) \le \phi^{+}_{\kappa}(x), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R} \}.$$
(3.36)

Note that by the periodicity of a(t) and b(t), A(t) in (2.17) can be chosen to be periodic in t with period T and then $\phi_{\kappa}^{-}(x,t)$ is periodic in t with period T.

Next, note that for any $\phi \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}^T$ and any $t_1 > t_2$,

$$0 \le \Phi(x,t;t_2,\phi_{\kappa}^+,\phi) \le \Phi(x,t;t_1,\phi_{\kappa}^+,\phi) \le \phi_{\kappa}^+(x)$$

for every $x \in \mathbb{R}$ and $t \ge t_1$. Let

$$\Phi(x,t;\phi) = \lim_{t_0 \to \infty} \Phi(x,t;t_0,\phi_{\kappa}^+,\phi) = \lim_{n \to \infty} \Phi(x,t;-nT,\phi_{\kappa}^+,\phi).$$

By the periodicity of a(t) and b(t), we have

$$\Phi(x,t;\phi) = \lim_{n \to \infty} \Phi(x,t;-nT,\phi_{\kappa}^{+},\phi)$$

=
$$\lim_{n \to \infty} \Phi(x,t+nT;0,\phi_{\kappa}^{+},\phi)$$

=
$$\lim_{n \to \infty} \Phi(x,t+(n+1)T;0,\phi_{\kappa}^{+},\phi)$$

=
$$\Phi(x,t+T;\phi).$$

Hence $\Phi(\cdot, \cdot; \phi) \in \tilde{\mathcal{E}}^T_{\kappa, \beta_0, M}$.

Now, by the similar arguments as in Claim 1, we can prove the following claim.

Claim 2. The mapping $\Phi : \tilde{\mathcal{E}}_{\kappa,\beta_0,M}^T \to \tilde{\mathcal{E}}_{\kappa,\beta_0,M}^T$, $\phi \mapsto \Phi(\cdot,\cdot,\phi)$, has a fixed point ϕ^* satisfying $\Phi(\cdot,\cdot,\phi^*) = \phi^*$. Furthermore, any element $\phi^* \in \tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ with the property $\Phi(\cdot,\cdot,\phi^*) = \phi^*$ also satisfies

$$\lim_{x \to -\infty} |\phi^*(x,t) - u^*(t)| = 0 \quad and \quad \lim_{x \to \infty} \frac{\phi^*(x,t)}{e^{-\kappa x}} = 1, \quad uniformly \ in \ t,$$

where $u^*(t)$ is the only positive entire solution of (1.3).

Finally, let $\tilde{U}(x,t) = \phi^*(x,t)$ and $\tilde{V}(x,t) = \Psi(x,t;\tilde{U})$. Then

$$(u(x,t), v(x,t)) = (U(x - C_{\kappa}(t), t), V(x - C_{\kappa}(t), t))$$

where $C_{\kappa}(t) = \int_0^t c_{\kappa}(s) ds$ and $c_{\kappa}(t) = \frac{a(t)+\kappa^2}{\kappa}$, is a transition front solution of (1.5) connecting (0,0) and $(u^*(t), v^*(t))$ satisfying (1.14) and (1.15). Let

$$U(x,t) = \tilde{U}(x+ct - \int_0^t c_k(s)ds, t)$$
 and $V(x,t) = \tilde{V}(x+ct - \int_0^t c_k(s)ds, t).$

Note that the function $\mathbb{R} \ni t \mapsto ct - \int_0^t c_\kappa(s) ds$ is periodic with period T. Then

$$(u(x,t), v(x,t)) = (U(x - ct, t), V(x - ct, t))$$

is a (periodic) transition front solution of (1.5) connecting (0,0) and $(u^*(t), v^*(t))$ satisfying (1.15) and (1.16).

Proof of Theorem 1.1(3). Suppose that $a(t) \equiv a$ and $b(t) \equiv b$ are independent of t. For every $c > c_{\chi}^*$, let $\kappa \in (0, \min\{\kappa_{\chi}, \sqrt{a}\})$ be such that $c = \frac{a+\kappa^2}{\kappa}$. Similarly, to prove Theorem 1.1(3), we change the set $\mathcal{E}_{\kappa,\beta_0,M}$ to

$$\mathcal{E}^{0}_{\kappa,\beta_{0},M} = \{\phi \in C^{b}_{\mathrm{unif}}(\mathbb{R}) \mid 0 \le \phi(x) \le \phi^{+}_{\kappa}(x) \text{ and } |\phi(x+h) - \phi(x)| \le M|h|^{\beta_{0}} \forall x, t, h \in \mathbb{R}, |h| \le 1\}.$$

$$(3.37)$$

and change $\tilde{\mathcal{E}}_{\kappa,\beta_0,M}$ to

$$\tilde{\mathcal{E}}^{0}_{\kappa,\beta_{0},M} = \{ \phi \in \mathcal{E}^{2}_{\kappa,\beta_{0},M} \, | \, \phi^{-}_{\kappa}(x,t) \le \phi(x) \le \phi^{+}_{\kappa}(x), \quad \forall \ x \in \mathbb{R}, \ \forall \ t \in \mathbb{R} \}.$$
(3.38)

Note that A(t) in (2.17) can be chosen to be independent of t and then $\phi_{\kappa}^{-}(x,t)$ is independent of t.

Note that, for any T > 0,

$$\tilde{\mathcal{E}}^0_{\kappa,\beta_0,M} \subset \tilde{\mathcal{E}}^T_{\kappa,\beta_0,M}$$

and $\tilde{\mathcal{E}}^0_{\kappa,\beta_0,M}$ is a closed convex subset of $\tilde{\mathcal{E}}^T_{\kappa,\beta_0,M}$ with respect to the open compact topology.

Note also that, for any $\phi \in \tilde{\mathcal{E}}^0_{\kappa,\beta_0,M}$ and any $t_1 > t_2$, we have

$$0 \le \Phi(x,t;t_2,\phi_{\kappa}^+,\phi) \le \Phi(x,t;t_1,\phi_{\kappa}^+,\phi) \le \phi_{\kappa}^+(x)$$

for every $x \in \mathbb{R}$ and $t \ge t_1$. Let

$$\Phi(x,t;\phi) = \lim_{t_0 \to \infty} \Phi(x,t;t_0,\phi_{\kappa}^+,\phi).$$

We have

$$\Phi(x,t;\phi) = \lim_{t_0 \to -\infty} \Phi(x,t;t_0,\phi_{\kappa}^+,\phi)$$
$$= \lim_{t_0 \to -\infty} \Phi(x,t-t_0;0,\phi_{\kappa}^+,\phi)$$
$$= \lim_{t_0 \to -\infty} \Phi(x,-t_0;0,\phi_{\kappa}^+,\phi)$$
$$= \Phi(x,0;\phi) \quad \forall t \in \mathbb{R}.$$

Hence $\Phi(x,t;\phi) \equiv \Phi(x;\phi)$ and $\Phi(\cdot;\phi) \in \tilde{\mathcal{E}}^{0}_{\kappa,\beta_{0},M}$. Then Φ in **Claim 2** has a fixed point $\phi^{*} \in \tilde{\mathcal{E}}^{0}_{\kappa,\beta_{0},M}$. Let $U(x,t) = \phi^{*}(x)$ and $V(x) = \Psi(x,t;\tilde{U})$. Then

$$(u(x,t), v(x,t)) = (U(x - ct), V(x - ct))$$

is a traveling wave solution of (1.5) connecting (0,0) and $\left(\frac{a}{b}, \frac{\mu}{\lambda}\frac{a}{b}\right)$ satisfying (1.17).

4 Nonexistence of transition front solutions

In this section, we prove Theorem 1.2 on the nonexistence of transition front solutions of (1.5) with least mean speed small then $2\sqrt{\underline{a}}$.

Proof of Theorem 1.2. Suppose that (u(t,x), v(t,x)) = (U(x - C(t), t), V(x - C(t), t)) is a transition front solution of (1.5) connecting $(u^*(t), \frac{\mu}{\lambda}u^*(t))$ and (0,0). It suffices to prove that

$$2\sqrt{\underline{a}} \le \liminf_{t-s \to \infty} \frac{C(t) - C(s)}{t-s}.$$
(4.1)

We prove this in five steps.

Step 1. In this step, we get some pointwise estimate for v(t, x) and $v_x(t, x)$.

Note that

$$0 < m_0 := \inf_{x \le 0, t \in \mathbb{R}} u(t, x + C(t)) = \inf_{x \le 0, t \in \mathbb{R}} U(x, t),$$
(4.2)

and that

$$0 \le u(t,x) \le M := \frac{a_{\sup}}{b_{\inf} - \chi \mu}, \quad \forall \ t, x \in \mathbb{R}.$$

$$(4.3)$$

Note also that for every $R \gg 1$, there is $C_R \gg 1$ and $\varepsilon_R > 0$ such that

$$|\chi v_x(t,\cdot)|_{L^{\infty}(B_{\frac{R}{2}}(x_0))} + |\chi \lambda v(t,\cdot)|_{L^{\infty}(B_{\frac{R}{2}}(x_0))} \le C_R ||u(t,\cdot)||_{L^{\infty}(B_R(x_0))} + \varepsilon_R M, \quad \forall \ x_0 \in \mathbb{R}$$
(4.4)

with $\lim_{R\to\infty} \varepsilon_R = 0$. By the arguments of [14, Lemma 2.2], for every p > 1, $t_0 > 0$, $s_0 \ge 0$ and R > 0, there is $C_{t_0,s_0,R,M,p}$

$$u(t+t_1,x) \le C_{t_0,s_0,R,M,p}[u(t+s+t_1,y)]^{\frac{1}{p}}(M+1), \quad \forall s \in [0,s_0], \ t \ge t_0, \ t_1 \in \mathbb{R}, \ |x-y| \le R.$$
(4.5)

By (4.4) and (4.5) with p > 1, $s_0 = 0$ and $t_0 = 1$, we have

$$\begin{aligned} |\chi v_x(t,x)| + \chi \lambda v(t,x) &= |\chi v_x(t_0 + t - t_0,x)| + \chi \lambda v(t_0 + t - t_0,x) \\ &\leq C_{R,p} \left(u(t_0 + t - t_0,x) \right)^{\frac{1}{p}} + \varepsilon_R M \\ &= C_{R,p} \left(u(t,x) \right)^{\frac{1}{p}} + \varepsilon_R M \quad \forall \ t \in \mathbb{R}, \ x \in \mathbb{R}, \end{aligned}$$
(4.6)

where $C_{R,p} = C_R \cdot C_{1,0,R,M,p} \cdot (M+1) (> 0).$

Step 2. In this step, we construct super-solutions of some equation related to the first equation in (1.1).

By [43, Lemma 2.2], for any $0 < a_0 < \underline{a}$, there is $A_{a_0}(t) \in W^{1,\infty}_{\text{loc}}(\mathbb{R}) \cap L^{\infty}(\mathbb{R})$ such that

$$a(t) + A'_{a_0}(t) \ge a_0, \quad \forall \ t \in \mathbb{R}.$$

Fix $0 < a_0 < \underline{a}$. For every $\gamma \in \mathbb{R}$ and $s \in \mathbb{R}$, let

$$\tilde{u}^{s,\gamma}(t,x) = e^{A_{a_0}(t)}u(t,x-\gamma s + C(s))$$

for $t \geq s$ and $x \in \mathbb{R}$. Then

$$\begin{split} \tilde{u}_t^{s,\gamma} &= A'_{a_0}(t)\tilde{u}^{s,\gamma} + e^{A_{a_0}(t)}u_t^{s,\gamma} \\ &= \tilde{u}^{s,\gamma}_{xx} - \chi v_x(t,x-\gamma s + C(s))\tilde{u}^{s,\gamma}_x \\ &+ \tilde{u}^{s,\gamma}(a(t) + A'_{a_0}(t) - \chi\lambda v(t,x-\gamma s + C(s)) - (b(t) - \chi\mu)u) \\ &\geq \tilde{u}^{s,\gamma}_{xx} - \chi v_x(t,x-\gamma s + C(s))\tilde{u}^{s,\gamma}_x \\ &+ \tilde{u}^{s,\gamma}(a_0 - \chi\lambda v(t,x-\gamma s + C(s)) - (b_{\sup} - \chi\mu)e^{\|A_{a_0}\|_{\infty}}\tilde{u}^{s,\gamma}) \\ &\geq \tilde{u}^{s,\gamma}_{xx} - \chi v_x(t,x-\gamma s + C(s))\tilde{u}^{s,\gamma}_x \\ &+ \tilde{u}^{s,\gamma}(a_0 - \chi\lambda\varepsilon_R M - \chi\lambda C_{R,p}e^{\frac{1}{p}\|A_{a_0}\|_{\infty}}(\tilde{u}^{s,\gamma})^{\frac{1}{p}} - (b_{\sup} - \chi\mu)e^{\|A_{a_0}\|_{\infty}}\tilde{u}^{s,\gamma}). \end{split}$$

Let $b_0 = (b_{\sup} - \chi \mu) e^{\|A_{a_0}\|_{\infty}}$ and

$$\mathcal{A}_{0}(u) = u_{xx} - \chi v_{x}(t, x - \gamma s + C(s))u_{x} + u(a_{0} - \chi\lambda\varepsilon_{R}M - \chi\lambda C_{R,p}e^{\frac{1}{p}\|A_{a_{0}}\|_{\infty}}u^{\frac{1}{p}} - (b_{\sup} - \chi\mu)e^{\|A_{a_{0}}\|_{\infty}}u).$$

Then $\tilde{u}^{s,\gamma}$ is a super-solution of

$$u_t = \mathcal{A}_0 u \tag{4.7}$$

for $t \geq s$ and $x \in \mathbb{R}$.

Step 3. In this step, we construct sub-solutions of some equation related to the first equation in (1.1).

For any $0 < \gamma < 2\sqrt{a_0}$, choose $0 < \varepsilon \ll 1$ such that

$$4(a_0 - 4\varepsilon) - (\gamma + \varepsilon)^2 \ge 2\varepsilon.$$

By [33, Lemma A1, Appendix], there exist is r > 0 and $h \in C^2(\mathbb{R})$ satisfying that

$$h(\rho) = 0 \text{ for } \rho \in (-\infty, 0], \ h(\rho) = 1 \text{ for } \rho \in [r, \infty), \ h'(\rho) > 0 \text{ for } 0 < \rho < r$$

and that

$$-h''(\rho) + (\gamma + \varepsilon)h'(\rho) - (a_0 - 2\varepsilon)h < 0 \quad \text{for } 0 < \rho < r.$$
(4.8)

Choose $R_0 \gg 1$ and $0 < \sigma \ll 1$ such that

$$C_{R_0,p}\sigma^{\frac{1}{p}} + \varepsilon_{R_0}M < \varepsilon,$$

$$\sigma < m_0 e^{-\|A_{a_0}\|_{\infty}},$$

and

$$\chi\lambda\varepsilon_{R_0}M + \chi\lambda C_{R_0,p}e^{\frac{1}{p}\|A_{a_0}\|_{\infty}}\sigma^{\frac{1}{p}} + (b_{\sup} - \chi\mu)e^{\|A_{a_0}\|_{\infty}}\sigma < 2\varepsilon.$$

Let

$$\underline{u}(t,x) = \sigma h(\gamma t - x)$$

and

$$B_s(t,x) = \min\{-\chi v_x(t,x-\gamma s + C(s)),\varepsilon\}.$$

Then

$$\begin{split} \underline{u}_{t} &= \sigma \gamma h'(\gamma t - x) \\ &= \sigma (\gamma + \varepsilon) h'(\gamma t - x) - \varepsilon \sigma h'(\gamma t - x) \\ &= (\gamma + \varepsilon) \sigma h'(\gamma t - x) + \varepsilon \underline{u}_{x} \\ &\leq \sigma h''(\gamma t - x) + (a_{0} - 2\varepsilon) \sigma h(\gamma t - x) + \varepsilon \underline{u}_{x} \quad (by (4.8)) \\ &= \underline{u}_{xx} + \varepsilon \underline{u}_{x} + (a_{0} - 2\varepsilon) \underline{u} \\ &\leq \underline{u}_{xx} + \min\{\varepsilon, -\chi v_{x}(t, x - \gamma s + C(s)) \underline{u}_{x} + (a_{0} - 2\varepsilon) \underline{u} \quad (since \ \underline{u}_{x} \leq 0) \\ &\leq \underline{u}_{xx} + B_{s}(t, x) \underline{u}_{x} + \underline{u} \Big(a_{0} - \chi \lambda \varepsilon_{R} M - \chi \lambda C_{R,p} e^{\frac{1}{p} ||A_{a_{0}}||_{\infty}} \underline{u}^{\frac{1}{p}} - (b_{\sup} - \chi \mu) e^{||A_{a_{0}}||_{\infty}} \underline{u} \Big). \end{split}$$

where $B_s(t, x) = \min\{\varepsilon, -\chi v_x(t, x - \gamma s + C(s))\}$. Let

$$\tilde{\mathcal{A}}_0 u = u_{xx} + B_s(t, x)u_x + u\Big(a_0 - \chi\lambda\varepsilon_R M - \chi\lambda C_{R,p}e^{\frac{1}{p}\|A_{a_0}\|_{\infty}}u^{\frac{1}{p}} - (b_{\sup} - \chi\mu)e^{\|A_{a_0}\|_{\infty}}u\Big).$$

Then $\underline{\mathbf{u}}(t, x)$ is a sub-solution of

$$u_t = \tilde{\mathcal{A}}_0 u \tag{4.9}$$

for $t \geq s$ and $x \in \mathbb{R}$.

Observe that (4.7) and (4.9) are the same at (t, x) such that $B_s(t, x) = -\chi v_x(t, x - \gamma s + C(s))$, and hence (4.7) and (4.9) are the same at (t, x) such that u(t, x) is sufficiently small.

Step 4. In this step, we prove $\underline{u}(t,x) \leq \tilde{u}^{s,\gamma}(t,x)$ for $t \geq s$ and $x \in \mathbb{R}$.

First of all, for any given $t_0 > s$, there are $x_0^- < x_0^+$ such that

$$\widetilde{u}^{s,\gamma}(t,x) > \sigma \ge \underline{\mathbf{u}}(t,x) \quad \text{for} \quad s \le t \le t_0, \ x \le x_0^-$$
(4.10)

and

$$\tilde{u}^{s,\gamma}(t,x) < \sigma, \quad \chi v_x(t,x-\gamma s + C(s)) = -B_s(t,x) \quad \text{for} \quad s \le t \le t_0, \ x \ge x_0^+.$$
(4.11)

Note that

$$\underline{\mathbf{u}}(s,x) < \tilde{u}^{s,\gamma}(s,x) \quad \forall \ x \in \mathbb{R}.$$
(4.12)

Hence, there is $s < \tilde{t}_0 \le t_0$ such that

$$\underline{\mathbf{u}}(t, x_0^+) \le \tilde{u}^{s, \gamma}(t, x_0^+) \quad \text{for} \quad s \le t \le \tilde{t}_0.$$

Then by (4.11) and comparison principle for parabolic equations, we have

$$\underline{\mathbf{u}}(t,x) \le \tilde{u}^{s,\gamma}(t,x) \quad \text{for} \quad s \le t \le \tilde{t}_0, \ x \ge x_0^+.$$

$$(4.13)$$

Next, we prove that

$$\underline{u}(t,x) \le \tilde{u}^{s,\gamma}(t,x) \quad \text{for} \quad s \le t \le t_0, \ x_0^- \le x \le x_0^+.$$

$$(4.14)$$

Assume on the contrary that there is $s < \tilde{t}_1 \le t_0$ and $x_0^- \le \tilde{x}_1 \le x_0^+$ such that

$$\underline{u}(\tilde{t}_1, \tilde{x}_1) > \tilde{u}^{s,\gamma}(\tilde{t}_1, \tilde{x}_1).$$

Let

$$t_1 := \inf\{t \in (s, t_0] \mid \exists x_t \in [x_0^-, x_0^+] \text{ satisfying } \underline{u}(t, x_t) \ge \tilde{u}^{s, \gamma}(t, x_t)\}.$$

Note that $t_1 > s$, otherwise there would exist a sequence $\{(t_n, x_n)\}_{n \ge 1}, t_n \in (s, t_0], x_n \in [x_0^-, x_0^+],$ and $\underline{u}(t_n, x_n) \ge \tilde{u}^{s,\gamma}(t_n, x_n)$ for every n with $t_n \to s$. Without loss of generality, we may suppose that $x_n \to x \in [x_0^-, x_0^+]$. Thus we must have that $\underline{u}(s, x) \ge \tilde{u}^{s,\gamma}(s, x)$, which contradicts to (4.12). Therefore by (4.10), (4.12), and (4.13) we have that $t_1 \le \tilde{t}_1$,

$$\underline{\mathbf{u}}(t,x) < \tilde{u}^{s,\gamma}(t,x) \quad \forall \ s \le t < t_1, \ x \in \mathbb{R},$$

$$(4.15)$$

and

$$\sigma \ge \underline{\mathbf{u}}(t_1, x_1) = \tilde{u}^{s, \gamma}(t_1, x_1) \quad \text{for some } x_0^- \le x_1 \le x_0^+.$$
(4.16)

This implies that there is $0 < \delta \ll 1$ such that $s < t_1 - \delta$ and

$$B_s(t,x) = -\chi v_x(t,x-\gamma s + C(s)) \quad \text{for} \quad t_1 - \delta \le t \le t_1, \ |x-x_1| \le \delta.$$

Note that

$$\underline{\mathbf{u}}(t_1 - \delta, x) < \tilde{u}^{s,\gamma}(t_1 - \delta, x) \quad \forall \ x \in \mathbb{R}$$

and

$$\underline{\mathbf{u}}(t, x_1 \pm \delta) \le \tilde{u}^{s, \gamma}(t, x \pm \delta) \quad \forall \ t_1 - \delta \le t \le t_1.$$

Then by comparison principle for parabolic equations, we have

$$\underline{\mathbf{u}}(t_1, x_1) < \tilde{u}^{s,\gamma}(t_1, x_1),$$

which contradicts to (4.16). Hence (4.14) holds.

Now, by (4.11), (4.14), and comparison principle for parabolic equations, we have

$$\underline{\mathbf{u}}(t,x) \le \tilde{u}^{s,\gamma}(t,x) \quad \forall \ s \le t \le t_0, \ x \in \mathbb{R}.$$

Then, since $t_0 > s$ is arbitrary, we have

$$\underline{\mathbf{u}}(t,x) \le \tilde{u}^{s,\gamma}(t,x) \quad \forall \ t \ge s, \ x \in \mathbb{R}.$$
(4.17)

Step 5. In this step, we prove that (4.1) holds.

Taking $x = \gamma t - \frac{r}{2}$ in (4.17), we obtain that

$$0 < e^{-\|A_{a_0}\|_{\infty}} h(\frac{r}{2}) \le u(t, \gamma(t-s) - \frac{r}{2} + C(s)) = U(t, \gamma(t-s) - \frac{r}{2} + C(s) - C(t)), \forall t > s.$$

This together with the fact that $\lim_{x\to\infty} U(t,x) = 0$ uniformly in $t \in \mathbb{R}$ implies that there is $L \gg 1$ such that

$$\gamma \leq \frac{C(t) - C(s)}{t - s} + \frac{L + r/2}{t - s} \quad \text{for} \quad t > s,$$

and then

$$\gamma \leq \liminf_{t-s \to \infty} \frac{C(t) - C(s)}{t-s}.$$

Since, γ is arbitrarily chosen less that $2\sqrt{a_0}$, we infer that

$$2\sqrt{a_0} \le \liminf_{t-s \to \infty} \frac{C(t) - C(s)}{t-s}, \quad \forall \ 0 < a_0 < \underline{a}.$$

This implies that (4.1) holds. The theorem is thus proved.

References

- S. Ai, W. Huang, and Z.-A. Wang, Reaction, diffusion and chemotaxis in wave propagation, Discrete Contin. Dyn. Syst. Ser. B, 20 (2015), no. 1, 1-21.
- [2] S. Ai and Z.-A. Wang, Traveling bands for the Keller-Segel model with population growth, Math. Biosci. Eng. 12 (2015), no. 4, 717-737.
- [3] D. G. Aronson and H. F. Weinberger, Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation, in "Partail Differential Equations and Related Topics" (J. Goldstein, Ed.), Lecture Notes in Math., Vol. 466, Springer-Verlag, New York, 1975, pp. 5-49.
- [4] D. G. Aronson and H. F. Weinberger, Multidimensional nonlinear diffusions arising in population genetics, Adv. Math., 30 (1978), pp. 33-76.
- [5] H. Berestycki, F. Hamel, and N. Nadirashvili, The speed of propagation for KPP type problems, I - Periodic framework, J. Eur. Math. Soc., 7 (2005), pp. 172-213.
- [6] H. Berestycki, F. Hamel, and L. Roques, Analysis of periodically fragmented environment model: II - Biological invasions and pulsating traveling fronts, J. Math. Pures Appl., 84 (2005), pp. 1101-1146.
- [7] H. Berestycki and F. Hamel, Generalized traveling waves for reaction-diffusion equations. Perspectives in nonlinear partial differential equations, 101-123, *Contemp. Math.*, 446, Amer. Math. Soc., Providence, RI, 2007.
- [8] H. Berestycki and F. Hamel, Generalized transition waves and their properties. Comm. Pure Appl. Math. 65 (2012), no. 5, 592-648.
- [9] M. Bramson, Convergence of Solutions of the Kolmogorov Equations to Traveling Waves, Mem. Amer. Math. Soc., 44 (1983), No. 285.
- [10] R. Fisher, The wave of advance of advantageous genes, Ann. of Eugenics, 7 (1937), pp. 335-369.
- [11] M. Freidlin and J. Gärtner, On the propagation of concentration waves in periodic and ramdom media, Soviet Math. Dokl., 20 (1979), pp. 1282-1286.
- [12] A. Friedman, Partial Differential Equations of Parabolic Type, Prentice–Hall, Inc., Englewood Cliffs, N.J., 1964.
- [13] M. Funaki, M. Mimura and T. Tsujikawa, Travelling front solutions arising in the chemotaxisgrowth model, *Interfaces Free Bound.*, 8 (2006), 223-245.
- [14] F. Hamel and C. Henderson, Propagation in a Fisher-KPP equation with non-local advection, preprint (2017), https://arxiv.org/abs/1709.00923.
- [15] F. Hamel, Qualitative properties of monostable pulsating fronts : exponential decay and monotonicity, J. Math. Pures Appl., (9) 89 (2008), pp. 355-399.

- [16] S. Heinze, G. Papanicolaou, and A. Stevens, A variational principle for propagation speeds in inhomogeneous media, SIAM J. Appl. Math., 62 (2001), pp. 129-148.
- [17] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Lecture Notes in Math., 840, Springer-Verlag, Berlin, 1981.
- [18] D. Horstmann and A. Stevens, A constructive approach to traveling waves in chemotaxis, J. Nonlin. Sci., 14 (2004), 1-25.
- [19] J. H. Huang and W. Shen, Speeds of spread and propagation for KPP models in time almost and space periodic media, SIAM J. Applied Dynamical Systems, 8 (2009), pp. 790-821.
- [20] W. Hudson and B. Zinner, Existence of traveling waves for reaction diffusion equations of Fisher type in periodic media, Boundary value problems for functional-differential equations, 187–199, World Sci. Publ., River Edge, NJ, 1995.
- [21] Y. Kametaka, On the nonlinear diffusion equation of Kolmogorov-Petrovskii- Piskunov type, Osaka J. Math., 13 (1976), pp. 11-66.
- [22] E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an instability, J. Theoret. Biol., 26 (1970), 399-415.
- [23] E. F. Keller and L. A. Segel, A Model for chemotaxis, J. Theoret. Biol., 30 (1971), 225-234.
- [24] A. Kolmogorov, I. Petrowsky, and N. Piscunov, A study of the equation of diffusion with increase in the quantity of matter, and its application to a biological problem, *Bjul. Moskovskogo Gos. Univ.*, 1 (1937), pp. 1-26.
- [25] J. Li, T. Li, and Z.-A. Wang, Stability of traveling waves of the Keller-Segel system with logarithmic sensitivity, Math. Models Methods Appl. Sci. 24 (2014), no. 14, 2819-2849.
- [26] X. Liang, Y. Yi, and X.-Q. Zhao, Spreading speeds and traveling waves for periodic evolution systems, J. Diff. Eq., 231 (2006), no. 1, pp. 57-77.
- [27] X. Liang and X.-Q. Zhao, Asymptotic speeds of spread and traveling waves for monotone semiflows with applications, *Comm. Pure Appl. Math.*, **60** (2007), no. 1, pp. 1-40.
- [28] X. Liang and X.-Q. Zhao, Spreading speeds and traveling waves for abstract monostable evolution systems, J. Funct. Anal., 259 (2010), no. 4, pp. 857-903
- [29] B. P. Marchant, J. Norbury, and J. A. Sherratt, Travelling wave solutions to a haptotaxisdominated model of malignant invasion, *Nonlinearity*, 14 (2001), 1653-1671.
- [30] H. Matano, Traveling waves in spatially random media, RIMS Kokyuroku, 1337 (2003), pp. 1-9.
- [31] G. Nadin, Traveling fronts in space-time periodic media, J. Math. Pures Appl., (9) 92 (2009), pp. 232-262.
- [32] G. Nadin, Critical travelling waves for general heterogeneous one-dimensional reactiondiffusion equations, Ann. Inst. H. Poincar Anal. Non Linaire, 32 (2015), no. 4, pp. 841-873.

- [33] G. Nadin and L. Rossi, Propagation phenomena for time heterogeneous KPP reactiondiffusion equations, J. Math. Pures Appl. (9) 98 (2012), no. 6, pp. 633-653.
- [34] G. Nadin and L. Rossi, Transition waves for Fisher-KPP equations with general timeheterogeneous and space-periodic coefficients, Anal. PDE, 8 (2015), no. 6, pp. 1351-1377.
- [35] G. Nadin and L. Rossi, Generalized transition fronts for one-dimensional almost periodic Fisher-KPP equations, Arch. Ration. Mech. Anal., 223 (2017), pp. 1239-1267.
- [36] G. Nadin, B. Perthame, and L. Ryzhik, Traveling waves for the Keller-Segel system with Fisher birth terms, *Interfaces Free Bound.* 10 (2008), no. 4, 517-538.
- [37] J. Nolen, J.-M. Roquejoffre, L. Ryzhik, and A. Zlatoš, Existence and non-existence of Fisher-KPP transition fronts, Arch. Ration. Mech. Anal., 203 (2012), no. 1, pp. 217-246.
- [38] J. Nolen, M. Rudd, and J. Xin, Existence of KPP fronts in spatially-temporally periodic adevction and variational principle for propagation speeds, *Dynamics of PDE*, 2 (2005), pp. 1-24.
- [39] J. Nolen and J. Xin, Existence of KPP type fronts in space-time periodic shear flows and a study of minimal speeds based on variational principle, *Discrete Contin. Dyn. Syst.*, 13 (2005), no. 5, pp. 1217-1234.
- [40] R. B. Salako and Wenxian Shen, Global existence and asymptotic behavior of classical solutions to a parabolic-elliptic chemotaxis system with logistic source on \mathbb{R}^N , Journal of Differential Equations, **262** (2017), pp. 5635-5690.
- [41] R. B. Salako and Wenxian Shen, Spreading Speeds and Traveling waves of a parabolic-elliptic chemotaxis system with logistic source on ℝ^N, Discrete and Continuous Dynamical Systems Series A, 37 (2017), pp. 6189-6225.
- [42] R. B. Salako and Wenxian Shen, Existence of Traveling wave solution of parabolic-parabolic chemotaxis systems, *Nonlinear Analysis: Real World Applications* Volume 42, August 2018, Pages 93-119.
- [43] R. B. Salako and W. Shen, Long time behavior of random and nonautonomous Fisher-KPP equations. Part I. Stability of equilibria and spreading speeds. (https://arxiv.org/pdf/1806.01354.pdf).
- [44] R. B. Salako and W. Shen, Long time behavior of random and nonautonomous Fisher-KPP equations. Part II. Transition fronts. (https://arxiv.org/pdf/1806.03508.pdf).
- [45] R. B. Salako and Wenxian Shen, Parabolic-elliptic chemotaxis model with space-time dependent logistic sources on R^N. I. Persistence and asymptotic spreading, to appear in Mathematical Models and Methods in Applied Sciences. (https://arxiv.org/pdf/1709.05785.pdf)
- [46] R. B. Salako and Wenxian Shen, Parabolic-elliptic chemotaxis model with space-time dependent logistic sources on \mathbb{R}^N . II. Existence, uniqueness and stability of positive entire solutions, to appear in Journal of mathematical analysis and applications.

- [47] D. H. Sattinger, On the stability of waves of nonlinear parabolic systems, Advances in Math., 22 (1976), pp. 312-355.
- [48] W. Shen, Traveling waves in diffusive random media, J. Dynam. Diff. Eq., 16 (2004), pp. 1011-1060.
- [49] W. Shen, Variational principle for spatial spreading speeds and generalized wave solutions in time almost and space periodic KPP models, *Trans. Amer. Math. Soc.*, **362** (2010), no. 10, pp. 5125-5168
- [50] W. Shen, Existence, uniqueness, and stability of generalized traveling solutions in time dependent monostable equations, J. Dynam. Diff. Eq., 23 (2011), no. 1, pp. 1-44.
- [51] W. Shen, Existence of generalized traveling waves in time recurrent and space periodic monostable equations, J. Appl. Anal. Comput., 1 (2011), no. 1, pp. 69-93.
- [52] T. Tao, B. Zhu and A. Zlatoš, Transition fronts for inhomogeneous monostable reactiondiffusion equations via linearization at zero, *Nonlinearity* 27 (2014), no. 9, pp. 2409-2416.
- [53] K. Uchiyama, The behavior of solutions of some nonlinear diffusion equations for large time, J. Math. Kyoto Univ., 183 (1978), pp. 453-508.
- [54] H. F. Weinberger, Long-time behavior of a class of biology models, SIAM J. Math. Anal., 13 (1982), pp. 353-396.
- [55] H. F. Weinberger, On spreading speeds and traveling waves for growth and migration models in a periodic habitat, J. Math. Biol., 45 (2002), pp. 511-548.
- [56] Z.-A. Wang, Mathematics of traveling waves in chemotaxisreview paper, Discrete Contin. Dyn. Syst. Ser. B 18 (2013), no. 3, 601-641.
- [57] J. Xin, Front propagation in heterogeneous media, SIAM Review, 42 (2000), pp. 161-230.
- [58] A. Zlatoš, Transition fronts in inhomogeneous Fisher-KPP reaction-diffusion equations, J. Math. Pures Appl., (9) 98 (2012), no. 1, pp. 89-102.