

# Spectral Mapping Theorems of Differentiable $C_0$ Semigroups

Abdelaziz Tajmouati<sup>1</sup>, Hamid Boua<sup>2</sup> and Mohammed Karmouni<sup>3</sup>

<sup>(1,2)</sup>Sidi Mohamed Ben Abdellah University

Faculty of Sciences Dhar El Mahraz

Laboratory of Mathematical Analysis and Applications

Fez, Morocco

Email: abdelaziz.tajmouati@usmba.ac.ma hamid12boua@yahoo.com

<sup>(3)</sup>Cadi Ayyad University

Multidisciplinary Faculty

Safi, Morocco

med89karmouni@gmail.com

## Abstract

Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on a Banach space  $X$  with infinitesimal generator  $A$ . In this work, we give conditions for which the spectral mapping theorem  $\sigma_*(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_*(A)\}$  holds, where  $\sigma_*$  can be equal to the essential, Browder and Kato spectrum. Also, we will be interested in the relations between the spectrum of  $A$  and the spectrum of the  $n$ th derivative  $T(t)^{(n)}$  of a differentiable  $C_0$  semigroup  $(T(t))_{t \geq 0}$ .

2010 Mathematics Subject Classification: *47D03, 47A10, 47A11*

Keywords and phrases: *Semigroup, differentiable, essential spectrum.*

## 1 Introduction and Preliminaries

Many equations of mathematical physics can be cast in the abstract form:

$$u'(t) = Au(t), t \geq 0, u(0) = u_0 \quad (1)$$

on a Banach space  $X$ . Here  $A$  is a given linear operator with domain  $D(A)$  and the initial value  $u_0$ . The semigroups can be used to solve a large class of problems commonly known

as the Cauchy problem. The solution of (1) will be given by  $u(t) = T(t)u_0$  for an operator semigroup  $(T(t))_{t \geq 0}$  on  $X$ .

In order to understand the behavior of the solutions in terms of the data concerning  $A$ , one seeks information about the spectrum of  $T(t)$  in terms of the spectrum of  $A$ . Unfortunately the spectral mapping theorem  $e^{t\sigma_*(A)} = \sigma_*(T(t)) \setminus \{0\}$  often fails, sometimes in dramatic ways. However, the inclusion

$$e^{t\sigma_*(A)} \subseteq \sigma_*(T(t)) \setminus \{0\} \quad (2)$$

always holds, where  $\sigma_* \in \{\sigma, \sigma_e, \sigma_b\}$  is the spectrum, essential spectrum or Browder spectrum.

A one-parameter family  $(T(t))_{t \geq 0}$  of bounded operators on a Banach space  $X$  is called a  $C_0$  semigroup of operators or a strongly continuous semigroup of operators if:

1.  $T(0) = I$ .
2.  $T(t+s) = T(t)T(s)$ ,  $\forall t, s \geq 0$ .
3.  $\lim_{t \rightarrow 0} T(t)x = x$ ,  $\forall x \in X$ .

$(T(t))_{t \geq 0}$  has a unique infinitesimal generator  $A$  defined on the domain  $D(A)$  by:

$$Ax = \lim_{t \rightarrow 0} \frac{T(t)x - x}{t}, \forall x \in D(A),$$

$$D(A) = \{x \in X : \lim_{t \rightarrow 0} \frac{T(t)x - x}{t} \text{ exists}\}.$$

Also,  $A$  is a closed operator, see [2, 3].

We introduce the following operator acting on  $X$  and depending on the parameters  $\lambda \in \mathbb{C}$  and  $t \geq 0$ :

$$B_\lambda(t)x = \int_0^t e^{\lambda(t-s)}T(s)x ds, x \in X$$

It is well known that  $B_\lambda(t)$  is a bounded linear operator from  $X$  to  $D(A)$  and we have ([2, 3]):

- a.  $(e^{\lambda t} - T(t))^n x = (\lambda - A)^n B_\lambda^n(t)x$ ,  $\forall x \in X, n \in \mathbb{N}$ ;
- b.  $(e^{\lambda t} - T(t))^n x = B_\lambda^n(t)(\lambda - A)^n x$ ,  $\forall x \in D(A^n), n \in \mathbb{N}$ ;
- c.  $R((e^{\lambda t} - T(t))^n) \subseteq R((\lambda - A)^n)$ ;

d.  $N((\lambda - A)^n) \subseteq N(e^{\lambda t} - T(t))^n$ .

The semigroup  $(T(t))_{t \geq 0}$  is called differentiable for  $t > t_0$  if for every  $x \in X$ ,  $t \mapsto T(t)x$  is differentiable for  $t > t_0$ .  $(T(t))_{t \geq 0}$  is called differentiable if it is differentiable for  $t > 0$ . If  $(T(t))_{t \geq 0}$  is differentiable,  $B_\lambda(t)x$  is differentiable and  $B'_\lambda(t)x = T(t)x + \lambda B_\lambda(t)x$ .

Let  $(T(t))_{t \geq 0}$  be a  $C_0$ -semigroup differentiable on  $X$  with infinitesimal generator  $A$ . We have ([3]):

a.  $(\lambda e^{\lambda t} - AT(t))x = (\lambda - A)B'_\lambda(t)x, \quad \forall x \in X;$

b.  $(\lambda e^{\lambda t} - AT(t))x = B'_\lambda(t)(\lambda - A)x, \quad \forall x \in D(A);$

Let  $\Delta = \{z \in \mathbb{C} : \alpha_1 < \arg z < \alpha_2\}$ ,  $-\frac{\pi}{2} < \alpha_1 < \alpha_2 < \frac{\pi}{2}$ , and for  $z \in \Delta$  let  $T(z)$  be a bounded linear operator. The family  $(T(z))_{z \in \Delta}$  is an analytic semigroup in  $\Delta$  if

i.  $z \mapsto T(z)$  is analytic in  $\Delta$ .

ii.  $T(0) = I$  and  $\lim_{z \rightarrow 0} T(z)x = x$  for every  $x \in X$ .

iii.  $T(z_1 + z_2) = T(z_1)T(z_2)$  for  $z_1, z_2 \in \Delta$ .

A semigroup  $(T(t))_{t \geq 0}$  will be called analytic if it is analytic in some sector  $\Delta$  containing the nonnegative real axis.

One of the classical approaches to finding information about the solution  $T(t)$  is to study the spectrum of the semigroup directly. In several applications, we have the explicit expression of the generator  $A$ . Thus we need information on the spectrum of the semigroup  $T(t)$  in terms of that of the generator  $A$ . In addition, the study of the spectral mapping theorem for the different parts of the spectrum proves to be essential.

The spectral inclusions for various reduced spectra of a differentiable  $C_0$  semigroup were studied by A. Tajmouati et. al [5][6]. But, the equality (spectral mapping theorem) is not examined. So it is natural to ask the question: can we establish equality?

In this work, we will continue in this direction, we will prove, under some conditions, the spectral equality for  $C_0$  semigroup for essential spectrum, Browder spectrum, upper(lower)semi-Fredholm spectrum. Also, we give an affirmative answer to the question we just asked.

## 2 $C_0$ semigroup

Throughout,  $X$  denotes a complex Banach space, let  $A$  be a closed linear operator on  $X$  with domain  $D(A)$ , we denote by  $A^*$ ,  $R(A)$ ,  $N(A)$ ,  $\sigma_{ap}(A)$ ,  $\sigma_s(A)$ ,  $\rho(A)$  and  $\sigma(A)$ , respec-

tively the adjoint, the range, the null space, the approximate point spectrum, the surjective spectrum, the resolvent set and the spectrum of  $A$ .

We start by the following results, they will be needed in the sequel.

**Lemma 1** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Suppose that there exists  $t_0 > 0$  such that  $AT(t_0)$  is bounded. Then  $(T(t)|_F)_{t \geq 0}$  is a uniformly continuous  $C_0$  semigroup on  $X$ , where  $F = N(e^{\lambda s} - T(s))$ ,  $\lambda \in \mathbb{C}$ ,  $s > 0$ .*

**Proof.** It is clear that  $F$  is a closed subspace of  $X$ ,  $A$ -invariant and  $T(t)$ -invariant, furthermore  $D(A|_F) = D(A) \cap F \subseteq F$ . Let  $x \in F$  and  $x_n = n \int_t^{t+\frac{1}{n}} T(s)x ds \in D(A) \cap F$ ,  $x_n \xrightarrow{n \rightarrow \infty} 0$ , so that  $\overline{D(A|_F)} = F$ . Let us show that  $A|_F$  is bounded. Let  $\mu \in \rho(A)$ , then  $R(\mu, A|_F)$  is bounded below. Indeed: If not, there exists  $(x_n)_n \subset F$  such that  $\|x_n\| = 1$  and  $R(\mu, A|_F)x_n \xrightarrow{n \rightarrow \infty} 0$ . Since  $AT(t_0)$  is bounded, then  $(\mu - A|_F)T(t_0)R(\mu, A|_F)x_n \xrightarrow{n \rightarrow \infty} 0$ . So,  $T(t_0)x_n \xrightarrow{n \rightarrow \infty} 0$ .

If  $t \geq t_0$ , then  $T(t)x_n = T(t_0)T(t-t_0)x_n = T(t-t_0)T(t_0)x_n \rightarrow 0$ , then  $e^{\lambda t}x_n = T(t)x_n \rightarrow 0$ , which is impossible.

If  $0 < t < t_0$ , there exists  $p \in \mathbb{N}^*$  such that  $pt > t_0$ , then  $T(pt)x_n = (T(t))^p x_n = T(pt-t_0)T(t_0)x_n \rightarrow 0$ , so  $e^{p\lambda t}x_n = (T(t))^p x_n \rightarrow 0$ , then  $x_n \rightarrow 0$  which contradicts the fact that  $\|x_n\| = 1$ . Hence, there exists  $M \in \mathbb{R}^+$  such that for all  $x \in F$   $\|R(\mu, A)x\| \geq M \|x\|$ .

Whence, for all  $x \in D(A|_F)$

$$\|R(\mu, A)(A - \mu)x\| \geq M \|(A - \mu)x\|.$$

Which gives

$$\|A|_F x\| \leq (|\mu| + M^{-1}) \|x\|$$

Now, let  $x \in F = \overline{D(A|_F)}$ , there exists  $(u_p)_p \subset D(A|_F)$ ,  $u_p \rightarrow x$ . For all  $p, q \in \mathbb{N}$ ,

$$\|Au_p - Au_q\| \leq (|\lambda_0| + C^{-1}) \|u_p - u_q\|$$

Hence is a Cauchy sequence on  $X$ , so converge. Since  $A$  is a closed operator, then  $x \in D(A|_F)$ . Hence the result. ■

A closed linear operator  $A$  is called an upper semi-Fredholm (resp, lower semi Fredholm) if  $\dim N(A) < \infty$  and  $R(A)$  is closed (resp,  $\text{codim}R(A) < \infty$ ).  $A$  is a Fredholm operator if it is a lower and upper semi-Fredholm operator, The essential spectrum of  $T$  is defined by :

$$\sigma_e(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Fredholm operator}\};$$

We denote by  $\sigma_{lf}(T)$  and  $\sigma_{uf}(T)$ , respectively the lower and upper semi-Fredholm spectra which are defined in the same manner.

**Theorem 1** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Suppose that there exists  $t_0 > 0$  such that  $AT(t_0)$  is bounded. Then*

$$\sigma_{uf}(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_{uf}(A)\}$$

If  $X$  is reflexive, we have :

$$\sigma_{lf}(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_{lf}(A)\}$$

$$\sigma_e(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_e(A)\}$$

**Proof.** Let  $\mu = e^{\lambda t} \notin \sigma_{uf}(T(t))$ , then  $\dim N(e^{\lambda t} - T(t)) < \infty$  and  $R(e^{\lambda t} - T(t))$  is closed. According [5, Lemma 2.3],  $R(\lambda - A)$  is closed. Since  $N(\lambda - A) \subseteq N(e^{\lambda t} - T(t))$ , we have  $\dim N(\lambda - A) < \infty$ . Then  $\lambda - A$  is upper semi-Fredholm. Therefore

$$\{e^{\lambda t}, \lambda \in \sigma_{uf}(A)\} \subseteq \sigma_{uf}(T(t)) \setminus \{0\} \quad (1).$$

Now, let  $\mu \in \sigma_{uf}(T(t)) \setminus \{0\}$  and  $F = N(\mu - T(t))$ , then  $F$  is an  $A$ -invariant and  $T(t)$ -invariant closed subspace of  $X$ . From lemma 1,  $A|_F$  is a bounded operator that generates the  $C_0$  semigroup  $(T(t)|_F)_{t \geq 0}$ . By (1), we have

$$\{e^{\lambda t}, \lambda \in \sigma_{uf}(A|_F)\} \subseteq \sigma_{uf}(T(t)|_F) \setminus \{0\} \quad .$$

Since  $A|_F$  is bounded, then  $\sigma_{uf}(A|_F) \neq \emptyset$ . Let  $\lambda_0 \in \sigma_{uf}(A|_F)$ , then  $e^{\lambda_0 t} \in \sigma_{uf}(T(t)|_F) \subseteq \sigma(T(t)|_F) = \{\mu\}$ . Then  $\mu = e^{\lambda_0 t} \in \{e^{\lambda t}, \lambda \in \sigma_{uf}(A|_F)\}$ .

Now, let's show that  $\lambda_0 \in \sigma_{uf}(A)$ . If  $\lambda_0 \notin \sigma_{uf}(A)$ , so  $A - \lambda_0$  is upper semi Fredholm, then  $\dim N(\lambda_0 - A) < \infty$  and  $R(\lambda_0 - A)$  is closed. Since  $N(\lambda_0 - A|_F) \subseteq N(\lambda_0 - A)$ , then  $\dim N(\lambda_0 - A|_F) < \infty$ . Let  $(\lambda_0 - A|_F)x_n \rightarrow y$  with  $(x_n)_n \subseteq F$ . Since  $R(\lambda_0 - A)$  is closed, then there exists  $x \in D(A)$  such that  $y = (\lambda_0 - A)x$ .  $B_{\lambda_0}(t)$  is bounded, so that then  $B_{\lambda_0}(t)(\lambda_0 - A|_F)x_n \rightarrow B_{\lambda_0}(t)(\lambda_0 - A)x$ , then  $(\mu - T(t))x_n \rightarrow (\mu - T(t))x$ . Hence,  $(\mu - T(t))x = 0$ , consequently  $x \in F$  and  $y \in R(\lambda_0 - A|_F)$ , which is absurd. Thus

$$\sigma_{uf}(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_{uf}(A)\}.$$

We now turn to the second statement. Since  $A$  is the generator of the  $C_0$  semigroup  $(T(t))_{t \geq 0}$ , then  $A^*$  is the generator of the  $C_0$  semigroup  $(T^*(t))_{t \geq 0}$ . Then

$$\sigma_{uf}(T^*(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_{uf}(A^*)\}.$$

By duality, we have

$$\sigma_{lf}(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_{lf}(A)\}$$

and consequently

$$\sigma_e(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_e(A)\}. \blacksquare$$

Recall that the ascent of  $A$  is defined as  $\text{asc}(A) = \inf\{n \in \mathbb{N} : N(A^n) = N(A^{n+1})\}$  and the descent of  $A$  is defined as  $\text{des}(A) = \inf\{n \in \mathbb{N} : R(A^n) = R(A^{n+1})\}$ . Where  $\inf \emptyset = \infty$ . We say that a closed operator  $A$  is upper semi-Browder if it is upper semi-Fredholm and has finite ascent. Similarly,  $A$  is lower semi-Browder if it is lower semi-Fredholm and has finite descent.  $A$  is Browder if it is both lower and upper semi-Browder. Equivalently, this means that  $A$  is Fredholm and has finite both ascent and descent.

The Browder spectrum of  $T$  is defined by :

$$\sigma_b(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Browder operator}\}.$$

Also, we denote by  $\sigma_{lb}(T)$  and  $\sigma_{ub}(T)$ , respectively the lower and upper semi-Browder spectra which are defined in the same manner.

The following corollary is an immediate consequence of the previous theorem.

**Corollary 1** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Suppose that there exists  $t_0 > 0$  such that  $AT(t_0)$  is bounded and  $X$  is reflexive. Then*

$$\sigma_{ub}(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_{ub}(A)\}$$

*If  $X$  is reflexive, we have :*

$$\sigma_{lb}(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_{lb}(A)\}$$

$$\sigma_b(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_b(A)\}$$

*Since every differentiable  $C_0$  semigroup matches the condition of the Theorem 1, we have the following result.*

**Corollary 2** *Let  $(T(t))_{t \geq 0}$  be a differentiable  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Then*

$$\sigma_*(T(t)) \setminus \{0\} = \{e^{\lambda t}, \lambda \in \sigma_*(A)\}$$

*with  $\sigma_* \in \{\sigma_{uf}, \sigma_{lf}, \sigma_f, \sigma_{ub}, \sigma_{lb}, \sigma_b\}$ .*

**Remark 1** Note that the inclusion  $\{e^{\lambda t}, \lambda \in \sigma_*(A)\} \subseteq \sigma_*(T(t)) \setminus \{0\}$ , where  $\sigma_* = \sigma_e, \sigma_{uf}, \sigma_{lf}, \sigma_{ub}, \sigma_{lb}, \sigma_b$  holds in the general setting where  $T(t)$  is a  $C_0$  semigroup. This inclusion is strict as shown in the following example.

**Example 1** Consider the translation group on the space  $C_{2\pi}(\mathbb{R})$  of all  $2\pi$  periodic continuous functions on  $\mathbb{R}$  and denote its generator by  $A$  (see [2, Paragraph I.4.15]). From [2, Examples 2.6.iv] we have,  $\sigma(A) = i\mathbb{Z}$ , then  $e^{t\sigma(A)}$  is at most countable, therefore  $e^{t\sigma_*(A)}$  are also. The spectra of the operators  $T(t)$  are always contained in  $\Gamma = \{z \in \mathbb{C} : |z| = 1\}$  and contain the eigenvalues  $e^{ikt}$  for  $k \in \mathbb{Z}$ . Since  $\sigma(T(t))$  is closed, it follows from [2, Theorem IV.3.16] that  $\sigma(T(t)) = \Gamma$  whenever  $t/2\pi \notin \mathbb{Q}$ , then  $\sigma(T(t))$  is not countable, so  $\sigma_*(T(t))$  are also. Therefore the inclusions of the Remark 1 are strict. According Corollary 2, the semigroup  $(T(t))_{t \geq 0}$  is not differentiable.

**Example 2** On  $C([0, 1])$  we consider the operator  $B$  given by

$$Bu := mu'' + qu'$$

where  $m(0) = m(1) = 0$ ,  $\sqrt{m} \in C^1[0, 1]$ ,  $q \in C[0, 1]$ , and  $q/\sqrt{m}$  is bounded in  $]0, 1[$ . The operator  $(B, D(B))$  generates an analytic semigroup  $(T(t))_{t \geq 0}$  of angle  $\frac{\pi}{2}$  in  $C[0, 1]$ , see [3, Theorem 4.20], where  $D(B) := \{u \in C[0, 1] \cap C^2(0, 1) : Bu \in C[0, 1]\}$ . Since  $(T(t))_{t \geq 0}$  is an analytic semigroup,  $(T(t))_{t \geq 0}$  matches the condition of the Theorem 1, then

$$\sigma_b(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_b(B)\};$$

$$\sigma_e(T(t)) \setminus \{0\} = \{e^{\lambda s}, \lambda \in \sigma_e(B)\}.$$

Next, let  $A$  be a closed linear operator,  $A$  is said to have the single valued extension property at  $\lambda_0 \in \mathbb{C}$  (SVEP) if for every open neighborhood  $U \subseteq \mathbb{C}$  of  $\lambda_0$ , the only analytic function  $f : U \rightarrow D(A)$  which satisfies the equation  $(A - zI)f(z) = 0$  for all  $z \in U$  is the function  $f \equiv 0$ .  $A$  is said to have the SVEP if  $A$  has the SVEP for every  $\lambda \in \mathbb{C}$ .

Recall that a closed operator  $A$  is said to be semi-regular if  $R(A)$  is closed and  $N(A) \subseteq R^\infty(A)$ , where  $R^\infty(A) = \bigcap_{n \geq 0} R(A^n)$ . The semi-regular spectrum of  $A$  is defined by:

$$\sigma_k(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not semi-regular}\}.$$

We have the following result.

**Proposition 1** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Then*

$$\{e^{\lambda s}, \lambda \in \sigma_k(A)\} \subseteq \sigma_k(T(t)) \setminus \{0\} \subseteq \{e^{\lambda s}, \lambda \in \sigma_{ap}(A)\}$$

*Furthermore, if  $A^*$  has the SVEP, we have*

$$\{e^{\lambda s}, \lambda \in \sigma_k(A)\} = \sigma_k(T(t)) \setminus \{0\}$$

**Proof.**

According to [1, Theorem 2.2], we have the inclusion  $\{e^{\lambda s}, \lambda \in \sigma_k(A)\} \subseteq \sigma_k(T(t)) \setminus \{0\}$ .

Now, let  $\mu \in \sigma_k(T(t)) \setminus \{0\}$  and  $F = N(\mu - T(t))$ , then  $F$  is an  $A$ -invariant and  $T(t)$ -invariant closed subspace of  $X$ . From lemma 1,  $A|_F$  is a bounded operator that generates the  $C_0$  semigroup  $(T(t)|_F)_{t \geq 0}$ . We have

$$\{e^{\lambda t}, \lambda \in \sigma_k(A|_F)\} \subseteq \sigma_k(T(t)|_F) \setminus \{0\} \quad .$$

Since  $A|_F$  is bounded, then  $\sigma_k(A|_F) \neq \emptyset$ . Let  $\lambda_0 \in \sigma_k(A|_F)$ , then  $e^{\lambda_0 t} \in \sigma_k(T(t)|_F) = \sigma(T(t)|_F) = \{\mu\}$ . Then  $\mu = e^{\lambda_0 t} \in \{e^{\lambda t}, \lambda \in \sigma_k(A|_F)\} \cup \{0\}$ . Now, let us show that  $\lambda_0 \in \sigma_*(A)$ . If  $\lambda_0 \notin \sigma_*(A)$ , so  $\lambda_0 \notin \sigma_*(A|_F)$ , then  $\lambda_0 \notin \sigma_k(A|_F)$ , absurd. ■

The following corollary is an immediate consequence of the previous proposition.

**Corollary 3** *Let  $(T(t))_{t \geq 0}$  be a  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . If  $X$  is reflexive and  $A$  has the SVEP, then*

$$\{e^{\lambda s}, \lambda \in \sigma_k(A)\} = \sigma_k(T(t)) \setminus \{0\}$$

### 3 Differentiable $C_0$ semigroup

**Lemma 2** *Let  $(T(t))_{t \geq 0}$  be a differentiable  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Let  $F = N(\mu - AT(t))$ ,  $\mu \in \sigma(AT(t)) \setminus \{0\}$ . Then  $(T(t)|_F)_{t \geq 0}$  is a uniformly continuous  $C_0$ -semigroup on  $X$  with infinitesimal generator  $A|_F$ .*

**Proof.** For all  $x \in F$ ,

$$\begin{aligned} (\mu - AT(t))T(s)x &= \mu T(s)x - AT(t)T(s)x \\ &= T(s)(\mu - AT(t))x = 0 \end{aligned}$$



For all  $x \in F \cap D(A)$ ,

$$\begin{aligned} (\mu - AT(t))Ax &= \mu Ax - AT(s)Ax \\ &= A(\mu - AT(t))x = 0 \end{aligned}$$

Hence  $F$  is a closed subspace of  $X$ ,  $A$ -invariant and  $T(t)$ -invariant.  $\overline{D(A|_F)} = F$ . Indeed: Let  $x \in F$  and  $x_p = p \int_t^{t+\frac{1}{p}} T(t)x dx$   $p \geq 1$ , then  $x_p \in D(A) \cap F = D(A|_F)$  and  $x_p \rightarrow x$ ,  $p \rightarrow +\infty$ . Then  $x \in \overline{D(A|_F)}$ , so  $\overline{D(A|_F)} = F$ . Let us show that  $A|_F$  is bounded. Let  $\lambda_0 \in \rho(A|_F)$ , then  $R(\lambda_0, A|_F)$  is bounded below, indeed: If not, there exists  $(x_p)_p \subset F$  such that  $\|x_p\| = 1$  and  $R(\lambda_0, A|_F)x_p \rightarrow 0$ , where  $p \rightarrow +\infty$ . Since  $(T(t))_{t \geq 0}$  is differentiable, then  $AT(t)$  is bounded, it is the same for  $(\lambda_0 - A)AT(t)$ .  $(\lambda_0 - A)AT(t)R(\lambda_0, A|_F)x_p = AT(t)x_p \rightarrow 0$ , where  $p \rightarrow +\infty$ . So,  $T(t)x_n \rightarrow 0$ , where  $n \rightarrow +\infty$ , then  $x_p \rightarrow 0$  which contradicts the fact that  $\|x_n\| = 1$ . Hence, there exists  $C > 0$  such that for all  $x \in F$   $\|R(\lambda_0, A|_F)x\| \geq C \|x\|$ . Then, for all  $x \in D(A|_F)$

$$\|Ax\| \leq (|\lambda_0| + C^{-1}) \|x\|$$

Hence the result. ■

**Theorem 2** *Let  $(T(t))_{t \geq 0}$  be a differentiable  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Then for  $t > 0$ , we have*

$$\begin{aligned} \sigma(AT(t)) \cup \{0\} &= \{\lambda e^{\lambda t}, \lambda \in \sigma(A) \cup \{0\}\} \\ \sigma_{ap}(AT(t)) \cup \{0\} &= \{\lambda e^{\lambda t}, \lambda \in \sigma_{ap}(A) \cup \{0\}\} \end{aligned}$$

**Proof.** From [3, Lemma 4.6], we have  $\{\lambda e^{\lambda t}, \lambda \in \sigma(A)\} \subseteq \sigma(AT(t))$ . Let  $\mu \in \sigma(AT(t)) \setminus \{0\}$ . From lemma 2  $(T(t)|_F)_{t \geq 0}$  is a uniformly continuous  $C_0$ -semigroup on  $X$  with infinitesimal generator  $A|_F$  where  $F = N(\mu - AT(t))$ ,  $\mu \in \sigma(A(t)) \setminus \{0\}$ . Then,  $T(t)|_F = e^{tA|_F}$ . Since the function  $f : z \rightarrow ze^{tz}$  is analytic on  $\mathbb{C}$ , from the spectral mapping theorem we have:

$$\sigma((AT(t))|_F) = \sigma(A|_F T(t)|_F) = \sigma(A|_F e^{tA|_F}) = \sigma(f(A|_F)) = f(\sigma(A|_F)).$$

$A|_F$  is bounded, then  $\sigma(A|_F) \neq \emptyset$ . Let  $\lambda_0 \in \sigma(A|_F)$ , then  $\lambda_0 e^{t\lambda_0} = f(\lambda_0) \in \sigma((AT(t))|_F) = \{\mu\}$ . Therefore,  $\mu = \lambda_0 e^{t\lambda_0}$ . Now, let us show that  $\lambda_0 \in \sigma(A)$ . If  $\lambda_0 - A$  is bijective, so  $\lambda_0 - A|_F$  is injective. Let  $y \in F$ ,  $\lambda_0 - A$  is surjective, then there exists  $x \in D(A)$  such that  $y = (\lambda_0 - A)x$ , since  $y \in F$  then  $(\lambda_0 e^{\lambda_0 t} - AT(t))y = (\lambda_0 - A)B'_{\lambda_0}(t)x = 0$ , so  $B'_{\lambda_0}(t)y = 0$

and as a result  $(\lambda_0 e^{\lambda_0 t} - AT(t))y = (\lambda_0 - A)B'_{\lambda_0}(t)x = 0$ , thus  $x \in F$ . Hence,  $\lambda_0 \in \rho(A|_F)$ , absurd. Therefore,

$$\sigma(AT(t)) \cup \{0\} = \{\lambda e^{\lambda t}, \lambda \in \sigma(A) \cup \{0\}\}.$$

Now, let  $\lambda_0 \in \sigma_{ap}(A)$ , then there exists  $(x_p)_p \subset D(A)$  such that  $\|x_p\| = 1$  and  $(\lambda_0 - A)x_p \rightarrow 0$  where  $p \rightarrow +\infty$ . We have  $B'_{\lambda_0}(t)$  is bounded, then  $(\lambda_0 e^{\lambda_0 t} - AT(t))x_p = B'_{\lambda_0}(t)(\lambda_0 - A)x_p \rightarrow 0$  where  $p \rightarrow +\infty$ . Therefore  $\lambda_0 e^{\lambda_0 t} \in \sigma_{ap}(AT(t))$ . So, we have

$$\{\lambda e^{\lambda t}, \lambda \in \sigma_{ap}(A) \cup \{0\}\} \subseteq \sigma_{ap}(AT(t)) \cup \{0\}$$

Let  $\mu \in \sigma_{ap}(AT(t)) \setminus \{0\}$ . Since  $\sigma_{ap}((AT(t))|_F) = \sigma(AT(t)|_F) = f(\sigma_{ap}(A|_F)) = \{\mu\}$ .  $A|_F$  is bounded, then  $\sigma_{ap}(A|_F) \neq \emptyset$ . Let  $\lambda_0 \in \sigma_{ap}(A|_F)$ , then  $\mu = \lambda_0 e^{\lambda_0 t}$ . We have  $\lambda_0 \in \sigma_{ap}(A)$ . Indeed: suppose that  $\lambda_0 \notin \sigma_{ap}(A)$ , then  $\lambda_0 - A$  is injective with closed range, so  $\lambda_0 - A|_F$  is injective. Let  $(\lambda_0 - A|_F)x_n \rightarrow y$ . Since  $R(\lambda_0 - A)$  is closed then there exists  $x \in D(A)$  such that  $y = (\lambda_0 - A|_F)x$ .  $B'_{\lambda_0}(t)$  is bounded, then  $B'_{\lambda_0}(t)(\lambda_0 - A|_F)x_n \rightarrow B'_{\lambda_0}(t)(\lambda_0 - A|_F)x = (\lambda_0 e^{\lambda_0 t} - AT(t))x$ .  $x_n \in F$ , then  $B'_{\lambda_0}(t)(\lambda_0 - A|_F)x_n = 0$ . Then  $x \in F$ . Therefore  $\lambda_0 \in \rho_{ap}(A|_F)$ , absurd. Consequently,  $\sigma_{ap}(AT(t)) \cup \{0\} = \{\lambda e^{\lambda t}, \lambda \in \sigma_{ap}(A) \cup \{0\}\}$ . ■

**Corollary 4** *Let  $(T(t))_{t \geq 0}$  be a differentiable  $C_0$  semigroup on  $X$  with infinitesimal generator  $A$ . Then for  $t > 0$ , we have*

$$\forall n \in \mathbb{N}, \forall t > 0 \quad \sigma_*(T(t)^{(n)}) \cup \{0\} = \{\lambda^n e^{\lambda t}, \lambda \in \sigma_*(A) \cup \{0\}\}$$

where  $\sigma_* = \sigma, \sigma_{ap}$

**Proof.** Let  $n \in \mathbb{N}, t > 0$ . Since  $(T(t))_{t \geq 0}$  is a differentiable  $C_0$  semigroup, then  $T(t)^{(n)}$  is a bounded operator, furthermore

$$T(t)^{(n)} = A^n T(t) = \left( AT \left( \frac{t}{n} \right) \right)^n.$$

From the spectral mapping theorem, we have

$$\begin{aligned} \sigma_*(T(t)^{(n)}) \cup \{0\} &= \sigma_* \left( AT \left( \frac{t}{n} \right) \right)^n \cup \{0\} = \left\{ \mu^n : \mu \in \sigma_* \left( AT \left( \frac{t}{n} \right) \right) \right\} \cup \{0\} \\ &= \left\{ \left( \lambda e^{\lambda \frac{t}{n}} \right)^n, \lambda \in \sigma_*(A) \cup \{0\} \right\} \\ &= \{ \lambda^n e^{\lambda t}, \lambda \in \sigma_*(A) \cup \{0\} \}. \blacksquare \end{aligned}$$

**Example 3** *Let  $X := \ell^p$  and  $M_q$  be the multiplication operator defined by  $M_q(x_n)_{n \in \mathbb{N}} = (q_n x_n)_{n \in \mathbb{N}}$  with  $q = (q_n)_{n \in \mathbb{N}}$  and  $q_n = -n + in^2$ . Then  $\sigma(M_q) = \{q_n : n \in \mathbb{N}\}$ . From [2, Counterexamples II.4.16] the semigroup  $(T_q(t))_{t \geq 0}$  generated by  $M_q$  is differentiable. By Corollary 4, for  $t > 0$ , we have  $\sigma(T_q(t)^{(p)}) \cup \{0\} = \{(-n + in^2)^p e^{t(-n + in^2)} : n \in \mathbb{N}\}$ .*

**Example 4** Consider the heat equation in  $L^p(0, \pi)$ .

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = \frac{\partial^2 u}{\partial x^2}(t, x), (t, x) \in \mathbb{R}^+ \times (0, \pi) \\ u(t, 0) = 0 = u(t, \pi), t \geq 0 \\ u(0, x) = f(x) \quad x \in (0, \pi) \end{cases} .$$

Let  $p > 2$ . On  $X = L^p(0, \pi)$  consider the operator defined by

$$Af(x) = f''(x)$$

with domain  $D(A) = W^{2,p}(0, \pi) \cap W_0^{1,p}(0, \pi)$ ,  $x \in (0, \pi)$  where

$$W_0^{1,p} = \{f \in W^{1,p}(0, \pi) : f(0) = 0 = f(\pi)\}.$$

The operator  $A$  is self-adjoint. The corresponding semigroup  $(T(t))_{t \geq 0}$  is analytic, so is differentiable.

For each  $f \in W^{2,p}(0, \pi) \cap W_0^{1,p}(0, \pi)$  the unique solution of the equation is given by

$$u(t, x) = (T(t)f)(x).$$

The spectrum of  $A$  is  $\sigma(A) = \{-n^2; n \geq 1\}$ . Since  $\text{int}(\sigma(A)) = \emptyset$ , then  $A$  and  $A^*$  have SVEP. So,  $\sigma(A) = \sigma_k(A) = \sigma_{ap}(A) = \sigma_{su}(A)$ .

From Proposition 1, we have  $\sigma_k(T(t)) \cup \{0\} = \{e^{-tn^2}, n \geq 1\} \cup \{0\}$ .

Also, by Corollary 4, we have

$$\sigma(T(t)^{(p)}) \cup \{0\} = \sigma_{ap}(T(t)^{(p)}) \cup \{0\} = \{(-1)^p n^{2p} e^{-tn^2}, n \geq 0\}.$$

## References

- [1] A. ELKOUTRI AND M. A. TAOUDI, *Spectre singulier pour les g en erateurs des semi-groupes*, C. R. Acad. Sci. Paris S er. I Math., 333 (2001), no. 7, 641-644 (French).
- [2] K. J. ENGEL AND R. NAGEL, *One-Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics, vol. 194, Springer-Verlag, New York, 2000.
- [3] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York 1983.
- [4] A. TAJMOUATI AND H. BOUA *Spectral Mapping Theorem for  $C_0$ -Semigroups of Drazin Spectrum*, Bol. Soc. Paran. Mat., to appear.

- [5] A. TAJMOUATI, M. AMOUCH, M.R.F. ALHOMIDI ZAKARIYA *Spectral Inclusion for  $C_0$ -Semigroups Drazin Invertible and Quasi-Fredholm Operators*, Bull. Malays. Math. Sci. Soc., to appear.
- [6] A. TAJMOUATI, M. AMOUCH, M.R.F. ALHOMIDI ZAKARIYA *Spectral Equality for  $C_0$ -Semigroups and Spectral Inclusion of B-Fredholm*, Rend. Circ. Mat. Palermo, 65(3) (2016), 425-434.