

Electromagnetic properties of singly heavy baryons

June-Young Kim^{1,*} and Hyun-Chul Kim^{1,2,3,†}

¹*Department of Physics, Inha University, Incheon 22212, Republic of Korea*

²*School of Physics, Korea Institute for Advanced Study (KIAS), Seoul 02455, Republic of Korea*

³*Advanced Science Research Center, Japan Atomic Energy Agency, Shirakata, Tokai, Ibaraki, 319-1195, Japan*

(Dated: November 7, 2018)

In this presentation, we summarize selectively recent results of the electromagnetic form factors of singly heavy baryons with spin 1/2 and related quantities, which were derived within a framework of the self-consistent chiral quark-soliton model. The results are compared with those from the lattice QCD and their physical implications are discussed.

Keywords: the chiral quark-soliton model, pion mean fields, singly heavy baryons, electromagnetic form factors

arXiv:1811.01936v1 [hep-ph] 5 Nov 2018

* E-mail: junyoung.kim@inha.edu

† E-mail: hchkim@inha.ac.kr

I. INTRODUCTION

The chiral quark-soliton model (χ QSM) was developed as a pion mean-field approach to describe the structures of the nucleon [1, 2]. The model was extended to describe also hyperons [3–5]. Recently, the model was successfully applied to the description of singly heavy baryons [6–10] in the limit of the infinitely heavy quark mass ($m_Q \rightarrow \infty$). In this presentation, we summarize selectively a recent work on the electromagnetic form factors of singly heavy baryons with spin 1/2 [11] and compare the results with those from a lattice QCD [12].

II. RESULTS AND DISCUSSION




FIG. 1. Decomposition of the valence and sea contributions of the electric form factors for the charmed baryons Σ_c^{++} and Ω_c^0 , in the left panel, and the right panel, respectively. The results are compared with those from the lattice QCD.

In Fig. 1, we show the results of the electric form factors of the charmed heavy baryons Σ_c^{++} and Ω_c^0 , comparing them with those from the lattice QCD [12]. The present results seem to fall off faster than those from the lattice QCD. This can be easily understood: When the value of the pion mass employed by the lattice QCD calculation is larger than the experimental one, the nucleon EM form factors decrease less slowly than the experimental data. This tendency can be also found in the magnetic form factors of the heavy baryons as drawn in Fig. 2.




FIG. 2. Decomposition of the valence and sea contributions of the magnetic form factors for the charmed baryons Σ_c^{++} and Ω_c^0 , in the left panel, and the right panel, respectively. The results are compared with those from the lattice QCD.

In Fig. 1, we depict the light-quark contributions of the valence and sea parts to the electric form factors separately. In the present pion mean-field approach and in the limit of $m_Q \rightarrow \infty$, the heavy quark can be considered as a point-like

TABLE I. Dipole magnetic moments of the singly heavy baryons. The second and third columns list the present results without and with the effects of SU(3) symmetry breaking, respectively.

B_c	$\mu_{B_c}^{(m_s=0)}$	$\mu_{B_c}^{(m_s=180)}$	[12, 13]	[10]
Σ_c^{++}	2.147	2.176	2.220(505)	2.15(10)
Σ_c^+	0.537	0.445	-	0.46(3)
Σ_c^0	-1.073	-1.286	-1.073(269)	-1.24(5)
Ξ_c^+	0.537	0.603	0.315(141)	0.60(2)
Ξ_c^0	-1.073	-1.147	-0.599(71)	-1.05(4)
Ω_c^0	-1.073	-1.014	-0.639(88)	-0.85(5)

particle, so that it plays only a part in making the concerned baryon charge complete. Note that the heavy-quark contribution to the magnetic form factors also vanishes in the limit of $m_Q \rightarrow \infty$. So, the light quarks govern all dynamics in a heavy baryon.

In Table I, we list the results of the magnetic moments of the charmed baryons, comparing them with those from the lattice QCD and from a “model-independent” approach of the χ QSM. We want to mention that the nuclear magneton in the model is evaluated not by the experimental value of the nucleon mass but by the model value, which is larger than the experimental one. Keeping this in mind, we find that except for $\mu_{\Sigma^{++}}$ the present results turn out larger than those from the lattice QCD.

The work is supported by Basic Science Research Program through the National Research Foundation (NRF) of Korea funded by the Korean government (Ministry of Education, Science and Technology(MEST)): Grant No. NRF-2018R1A2B2001752.

-
- [1] D. Diakonov, V. Y. Petrov and P. V. Pobylitsa, Nucl. Phys. B **306**, 809 (1988).
[2] D. Diakonov, hep-ph/9802298.
[3] A. Blotz, D. Diakonov, K. Goeke, N. W. Park, V. Petrov and P. V. Pobylitsa, Nucl. Phys. A **555**, 765 (1993).
[4] M. Praszalowicz, T. Watabe and K. Goeke, Nucl. Phys. A **647**, 49 (1999).
[5] C. V. Christov, A. Blotz, H.-Ch. Kim, P. Pobylitsa, T. Watabe, T. Meissner, E. Ruiz Arriola and K. Goeke, Prog. Part. Nucl. Phys. **37**, 91 (1996).
[6] Gh.-S. Yang, H.-Ch. Kim, M. V. Polyakov and M. Praszalowicz, Phys. Rev. D **94**, 071502 (2016).
[7] H.-Ch. Kim, M. V. Polyakov and M. Praszalowicz, Phys. Rev. D **96**, 014009 (2017) Addendum: [Phys. Rev. D **96**, 039902 (2017)].
[8] H.-Ch. Kim, M. V. Polyakov, M. Praszalowicz and G. S. Yang, Phys. Rev. D **96**, 094021 (2017) Erratum: [Phys. Rev. D **97**, 039901 (2018)].
[9] J. Y. Kim, H.-Ch. Kim and G. S. Yang, Phys. Rev. D **98**, 054004 (2018).
[10] G. S. Yang and H.-Ch. Kim, Phys. Lett. B **781**, 601 (2018).
[11] J. Y. Kim and H.-Ch. Kim, Phys. Rev. D **97**, 114009 (2018).
[12] K. U. Can, G. Erkol, B. Isildak, M. Oka and T. T. Takahashi, JHEP **1405**, 125 (2014).
[13] H. Bahtiyar, K. U. Can, G. Erkol, M. Oka and T. T. Takahashi, Phys. Lett. B **772** (2017) 121 doi:10.1016/j.physletb.2017.06.022 [arXiv:1612.05722 [hep-lat]].