

## DIAMETER OF REDUCED SPHERICAL CONVEX BODIES

MAREK LASSAK AND MICHAŁ MUSIELAK

ABSTRACT. The intersection  $L$  of two different non-opposite hemispheres of the unit sphere  $S^2$  is called a lune. By  $\Delta(L)$  we denote the distance of the centers of the semicircles bounding  $L$ . By the thickness  $\Delta(C)$  of a convex body  $C \subset S^2$  we mean the minimal value of  $\Delta(L)$  over all lunes  $L \supset C$ . We call a convex body  $R \subset S^2$  reduced provided  $\Delta(Z) < \Delta(R)$  for every convex body  $Z$  being a proper subset of  $R$ . Our aim is to estimate the diameter of  $R$ , where  $\Delta(R) < \frac{\pi}{2}$ , in terms of its thickness.

## 1. INTRODUCTION

Let  $S^2$  be the unit sphere of the 3-dimensional Euclidean space  $E^3$ . A *great circle* of  $S^2$  is the intersection of  $S^2$  with any two-dimensional linear subspace of  $E^3$ . By a pair of *antipodes* of  $S^2$  we mean any pair of points being the intersection of  $S^2$  with a one-dimensional subspace of  $E^3$ . Observe that if two different points  $a, b \in S^2$  are not antipodes, there is exactly one great circle passing through them. By the *arc*  $ab$  connecting them we understand the shorter part of the great circle through  $a$  and  $b$ . By the *distance*  $|ab|$  of  $a$  and  $b$  we mean the length of the arc  $ab$ . The notion of the *diameter* of any set  $A \subset S^2$  not containing antipodes is taken with respect to this distance and denoted by  $\text{diam}(A)$ .

A subset of  $S^2$  is called *convex* if it does not contain any pair of antipodes of  $S^2$  and if together with every two points it contains the arc connecting them. A closed convex set  $C \subset S^2$  with non-empty interior is called a convex body. Its boundary is denoted by  $\text{bd}(C)$ . If in  $\text{bd}(C)$  there is no arc, we say that the body is *strictly convex*. Convexity on  $S^2$  is considered in very many papers and monographs. For instance in [1], [2], [3], [4], [5] and [14].

---

2010 *Mathematics Subject Classification.* 52A55.

*Key words and phrases.* spherical convex body, spherical geometry, hemisphere, lune, width, constant width, thickness, diameter.

The set of points of  $S^2$  in the distance at most  $\rho$ , where  $0 < \rho \leq \frac{\pi}{2}$ , from a point  $c \in S^2$  is called a *spherical disk*, or shorter a *disk*, of *radius*  $\rho$  and *center*  $c$ . Disks of radius  $\frac{\pi}{2}$  are called *hemispheres*. Two hemispheres whose centers are antipodes are called *opposite hemispheres*. The set of points of a great circle of  $S^2$  which are at distance at most  $\frac{\pi}{2}$  from a fixed point  $p$  of this great circle is called a *semicircle*. We call  $p$  the *center* of this semicircle.

Let  $C \subset S^2$  be a convex body and  $p \in \text{bd}(C)$ . We say that a hemisphere  $K$  *supports*  $C$  at  $p$  provided  $C \subset K$  and  $p$  is in the great circle bounding  $K$ .

Since the intersection of every family of convex sets is also convex, for every set  $A \subset S^2$  contained in an open hemisphere of  $S^2$  there is the smallest convex set  $\text{conv}(A)$  containing  $A$ . We call it *the convex hull of*  $A$ .

If non-opposite hemispheres  $G$  and  $H$  are different, then  $L = G \cap H$  is called a *lune*. The semicircles bounding  $L$  and contained in  $G$  and  $H$ , respectively, are denoted by  $G/H$  and  $H/G$ . The *thickness*  $\Delta(L)$  of  $L$  is defined as the distance of the centers of  $G/H$  and  $H/G$ .

After [8] recall a few notions. For any hemisphere  $K$  supporting a convex body  $C \subset S^2$  we find a hemisphere  $K^*$  supporting  $C$  such that the lune  $K \cap K^*$  is of the minimum thickness (by compactness arguments at least one such a hemisphere  $K^*$  exists). The thickness of the lune  $K \cap K^*$  is called *the width of*  $C$  *determined by*  $K$  and it is denoted by  $\text{width}_K(C)$ . By the *thickness*  $\Delta(C)$  of a convex body  $C \subset S^2$  we understand the minimum of  $\text{width}_K(C)$  over all supporting hemispheres  $K$  of  $C$ . We say that a convex body  $R \subset S^2$  is *reduced* if for every convex body  $Z \subset R$  different from  $R$  we have  $\Delta(Z) < \Delta(R)$ . This definition is analogous to the definition of a reduced body in normed spaces (for a survey of results on reduced bodies see [10]). If for all hemispheres  $K$  supporting  $C$  the numbers  $\text{width}_K(C)$  are equal, we say that  $C$  is *of constant width*. Spherical bodies of constant width are discussed in [12] and applied in [6].

Just bodies of constant width, and in particular disks, are simple examples of reduced bodies on  $S^2$ . Also each of the four parts of a disk dissected by two orthogonal great circles through the center of this disk is a reduced body. It is called a *quarter of a disk*. There is a wide class of reduced odd-gons on  $S^2$  (see [9]). In particular, the regular odd-gons of thickness at most  $\frac{\pi}{2}$  are reduced.

## 2. TWO LEMMAS

**Lemma 2.1.** *Let  $L \subset S^2$  be a lune of thickness at most  $\frac{\pi}{2}$  whose bounding semicircles are  $Q$  and  $Q'$ . For every  $u, v, z$  in  $Q$  such that  $v \in uz$  and for every  $q \in L$  we have  $|qv| \leq \max\{|qu|, |qz|\}$ .*

*Proof.* If  $\Delta(L) = \frac{\pi}{2}$  and  $q$  is the center of  $Q'$ , then the distance between  $q$  and any point of  $Q$  is the same, and thus the assertion is obvious. Consider the opposite case when  $\Delta(L) < \frac{\pi}{2}$ , or  $\Delta(L) = \frac{\pi}{2}$  but  $q$  is not the center of  $Q'$ . Clearly, the closest point  $p \in Q$  to  $q$  is unique. Observe that for  $x \in Q$  the distance  $|qx|$  increases as the distance  $|px|$  increases. This easily implies the assertion of our lemma.  $\square$

In a standard way, an extreme point of a convex body  $C \subset S^2$  is defined as a point for which the set  $C \setminus \{e\}$  is convex (see [8]). The set of extreme points of  $C$  is denoted by  $E(C)$ .

**Lemma 2.2.** *For every convex body  $C \subset S^2$  of diameter at most  $\frac{\pi}{2}$  we have  $\text{diam}(E(C)) = \text{diam}(C)$ .*

*Proof.* Clearly,  $\text{diam}(E(C)) \leq \text{diam}(C)$ .

In order to show the opposite inequality  $\text{diam}(C) \leq \text{diam}(E(C))$ , thanks to  $\text{diam}(C) = \text{diam}(\text{bd}(C))$ , it is sufficient to show that  $|cd| \leq \text{diam}(E(C))$  for every  $c, d \in \text{bd}(C)$ . If  $c, d \in E(C)$ , this is trivial. In the opposite case, at least one of these points does not belong to  $E(C)$ . If, say  $d \notin E(C)$ , then having in mind that  $C$  is a convex body and  $d \in \text{bd}(C)$  we see that there are  $e, f \in E(C)$  different from  $d$  such that  $d \in ef$ . From  $E(C) \subset \text{bd}(C)$ , we see that  $e, f \in \text{bd}(C)$ . Since also  $d \in \text{bd}(C)$ , the arc  $ef$  is a subset of  $\text{bd}(C)$ .

Recall that by Theorem 3 of [8] we have  $\text{width}_K(C) \leq \text{diam}(C)$  for every hemisphere  $K$  supporting  $C$ . In particular, for the hemisphere  $K$  supporting  $C$  at all points of the arc  $ef$ . Thus by the assumption that  $\text{diam}(C) \leq \frac{\pi}{2}$  we obtain  $\text{width}_K(C) \leq \frac{\pi}{2}$  for our particular  $K$ . Hence we may apply Lemma 2.1 taking this  $K/K^*$  in the part of  $Q$  there. We obtain  $|cd| \leq \max\{|ce|, |cf|\}$ .

If  $c \in E(C)$ , from  $e, f \in E(C)$  we conclude that  $|cd| \leq \text{diam}(E(C))$ . If  $c \notin E(C)$ , from  $c \in \text{bd}(C)$  we see that there are  $g, h \in E(C)$  such that  $c \in gh$ . Similarly to the preceding paragraph we show that  $|ec| \leq \max\{|eg|, |eh|\}$

and  $|fc| \leq \max\{|fg|, |fh|\}$ . By these two inequalities and by the preceding paragraph we get  $|cd| \leq \max\{|eg|, |eh|, |fg|, |fh|\} \leq \text{diam}(E(C))$ , which ends the proof.  $\square$

The assumption that  $\text{diam}(C) \leq \frac{\pi}{2}$  is substantial in Lemma 2.2, as it follows from the example of a regular triangle of any diameter greater than  $\frac{\pi}{2}$ . The weaker assumption that  $\Delta(C) \leq \frac{\pi}{2}$  is not sufficient, which we see taking in the part of  $C$  any isosceles triangle  $T$  with  $\Delta(T) \leq \frac{\pi}{2}$  and the arms longer than  $\frac{\pi}{2}$  (so with the base shorter than  $\frac{\pi}{2}$ ). The diameter of  $T$  equals to the distance between the midpoint of the base and the opposite vertex of  $T$ . Hence  $\text{diam}(T)$  is over the length of each of the sides.

### 3. DIAMETER OF REDUCED SPHERICAL BODIES

The following theorem is analogous to the first part of Theorem 9 from [7] and confirms the conjecture from [9], p. 214. By the way, the much weaker estimate  $2 \arctan\left(\sqrt{2} \tan \frac{\Delta(R)}{2}\right)$  results from Theorem 2 in [13].

**Theorem 3.1.** *For every reduced spherical body  $R \subset S^2$  with  $\Delta(R) < \frac{\pi}{2}$  we have  $\text{diam}(R) \leq \arccos(\cos^2 \Delta(R))$ . This value is attained if and only if  $R$  is the quarter of disk of radius  $\Delta(R)$ . If  $\Delta(R) \geq \frac{\pi}{2}$ , then  $\text{diam}(R) = \Delta(R)$ .*

*Proof.* Assume that  $\Delta(R) < \frac{\pi}{2}$ . In order to show the first statement, by Lemma 2.2 it is sufficient to show that the distance between any two different points  $e_1, e_2$  of  $E(R)$  is at most  $\arccos(\cos^2 \Delta(R))$ . Since  $R$  is reduced, according to the statement of Theorem 4 in [8] there exist lunes  $L_j \supset R$ , where  $j \in \{1, 2\}$ , of thickness  $\Delta(R)$  with  $e_j$  as the center of one of the two semicircles bounding  $L_j$  (see Figure). Denote by  $b_j$  the center of the other semicircle bounding  $L_j$ .

If  $e_1 = b_2$  or  $e_2 = b_1$ , then  $|e_1 e_2| = \Delta(R)$ , which by  $\Delta(R) \in (0, \frac{\pi}{2})$  is at most  $\arccos(\cos^2 \Delta(R))$ . Otherwise  $L_1 \cap L_2$  is a non-degenerate spherical quadrangle with points  $e_1, b_2, b_1, e_2$  in its consecutive sides. Therefore, since  $e_1 \neq e_2$ , arcs  $e_1 b_1$  and  $e_2 b_2$  intersect at exactly one point. Denote it by  $g$ . Observe that it may happen  $b_1 = b_2 = g$ .

Let  $F$  be the great circle orthogonal to the great circle containing  $e_1 b_1$  and passing through  $e_2$ . Since  $e_2 \in L_1$ , we see that  $F$  intersects  $e_1 b_1$ . Let  $f$




FIGURE. Illustration to the proof of Theorem 3.1.

be the intersection point of them. Note that we do not exclude the case  $f = b_1$ . From  $|e_2b_2| = \Delta(R)$  we see that  $|ge_2| \leq \Delta(R) < \frac{\pi}{2}$ . Thus from the right spherical triangle  $gfe_2$  we conclude that  $|fe_2| \leq \Delta(R)$ . Moreover, from  $|e_1b_1| = \Delta(R)$  and  $f \in e_1b_1$  we obtain  $|fe_1| \leq \Delta(R)$ . Consequently, from the formula  $\cos k = \cos l_1 \cos l_2$  for the right spherical triangle with hypotenuse  $k$  and legs  $l_1, l_2$  applied to the triangle  $e_1fe_2$  (again see Figure) we obtain  $|e_1e_2| \leq \arccos(\cos^2 \Delta(R))$ .

Observe that thanks to  $\Delta(R) < \frac{\pi}{2}$ , the shown inequality becomes the equality only if  $g = f = b_1 = b_2$ . In this case, by Proposition 3.2 of [12], our body  $R$  is a quarter of disk of radius  $\Delta(R)$ .

Finally, we show the last statement of the theorem. Assume that  $\Delta(R) \geq \frac{\pi}{2}$ . By Theorem 4.3 of [12], the body  $R$  is of constant width  $\Delta(R)$ .

There is an arc  $pq \subset R$  of length equal to  $\text{diam}(R)$ . Clearly,  $p \in \text{bd}(R)$ . Take a lune  $L$  from Theorem 5.2 of [12] such that  $p$  is the center of a semicircle bounding  $L$ . Denote by  $s$  the center of the other semicircle  $S$  bounding  $L$ . By the third part of Lemma 3 in [8], we have  $|px| < |ps|$  for every  $x \in S$  different from  $s$ . Hence  $|px| \leq |ps|$  for every  $x \in S$ . So also  $|pz| \leq |ps|$  for every  $z \in L$ . In particular,  $|pq| \leq |ps|$ . Since  $\text{diam}(R) = |pq|$  and  $\Delta(R) = \Delta(L) = |ps|$ , we obtain  $\text{diam}(R) \leq \Delta(R)$ .

Let us show the opposite inequality.

If  $\text{diam}(R) \leq \frac{\pi}{2}$ , by Theorem 3 of [8] we get  $\Delta(R) = \text{diam}(R)$ . If  $\text{diam}(R) > \frac{\pi}{2}$ , then by Proposition 1 of [8] we get  $\Delta(R) \leq \text{diam}(W)$ . Consequently, always we have  $\Delta(R) \leq \text{diam}(R)$ .

From the inequalities obtained in the two preceding paragraphs we get the equality  $\text{diam}(R) = \Delta(R)$ . Hence the last sentence of the theorem is proved.  $\square$

Proposition 3.5 of [12] implies that *for every reduced spherical body  $R$  with  $\Delta(R) \leq \frac{\pi}{2}$  on  $S^2$  we have  $\text{diam}(R) \leq \frac{\pi}{2}$* . Here is a more precise form of this statement.

**Proposition 3.2.** Let  $R \subset S^2$  be a reduced body. Then  $\text{diam}(R) < \frac{\pi}{2}$  if and only if  $\Delta(R) < \frac{\pi}{2}$ . Moreover,  $\text{diam}(R) = \frac{\pi}{2}$  if and only if  $\Delta(R) = \frac{\pi}{2}$ .

*Proof.* We start with proving the first statement of our proposition. The function  $f(x) = \arccos(\cos^2 x)$  is increasing in the interval  $[0, \frac{\pi}{2}]$  as a composition of the decreasing functions  $\arccos x$  and  $\cos^2 x$ . From  $f(\frac{\pi}{2}) = \frac{\pi}{2}$  we conclude that in the interval  $[0, \frac{\pi}{2})$  the function  $f(x)$  accepts only the values below  $\frac{\pi}{2}$ . Thus by Theorem 3.1, if  $\Delta(R) < \frac{\pi}{2}$ , then  $\text{diam}(R) < \frac{\pi}{2}$ . The opposite implication results from the inequality  $\Delta(C) \leq \text{diam}(C)$  for every spherical convex body  $C$ , which in turn follows from Theorem 3 and Proposition 1 of [8].

Let us show the second part of our proposition.

Assume that  $\text{diam}(R) = \frac{\pi}{2}$ . The inequality  $\Delta(R) < \frac{\pi}{2}$  is impossible, by the first statement of our proposition. Also the inequality  $\Delta(R) > \frac{\pi}{2}$  is impossible, because by  $\Delta(R) \leq \text{diam}(R)$  (see Proposition 1 of [8]), it would imply  $\text{diam}(R) > \frac{\pi}{2}$  in contradiction to the assumption at the beginning of this paragraph. So  $\Delta(R) = \frac{\pi}{2}$ .

Now assume that  $\Delta(R) = \frac{\pi}{2}$ . By the second statement of Theorem 3.1 we get  $\text{diam}(R) = \frac{\pi}{2}$ .  $\square$

## REFERENCES

- [1] BESAU F., SCHUSTER F., Binary operations in spherical convex geometry, *Indiana Univ. Math. J.* **65** (2016), no. 4, 1263–1288.
- [2] DANZER L., GRÜNBAUM B., KLEE V., Helly's theorem and its relatives, *Proc. of Symp. in Pure Math.* vol. VII, Convexity, 1963, pp. 99–180.
- [3] FERREIRA O. P., IUSEM A. N., NÉMETH S. Z., Projections onto convex sets on the sphere, *J. Global Optim.* **57** (2013), 663–676.
- [4] GAO F., HUG D., SCHNEIDER R., Intrinsic volumes and polar sets in spherical space. *Math. Notae*, **41** (2003), 159–176.
- [5] HADWIGER H., Kleine Studie zur kombinatorischen Geometrie der Sphäre, *Nagoya Math. J.* **8** (1955), 45–48.

- [6] HAN H., NISHIMURA T., Self-dual Wulff shapes and spherical convex bodies of constant width  $\pi/2$ , *J. Math. Soc. Japan* **69** (2017), 1475–1484.
- [7] LASSAK M., Reduced convex bodies in the plane, *Israel J. Math.* **70** (1990), 365–379.
- [8] LASSAK M., Width of spherical convex bodies, *Aequationes Math.* **89** (2015), 555–567.
- [9] LASSAK M., Reduced spherical polygons, *Colloq. Math.* **138** (2015), 205–216.
- [10] LASSAK M., MARTINI H., Reduced convex bodies in finite-dimensional normed spaces – a survey, *Results Math.* **66** (2014), No. 3-4, 405–426.
- [11] LASSAK M., MUSIELAK M., Reduced spherical convex bodies, *Bull. Pol. Ac. Math.* **66** (2018), 87–97.
- [12] LASSAK M., MUSIELAK M., Spherical bodies of constant width, *Aequationes Math.* **92** (2018), 627–640.
- [13] MUSIELAK M., Covering a reduced spherical body by a disk, *Ukr. Math. J.*, to appear (see also arXiv:1806.04246).
- [14] VAN BRUMMELEN G., *Heavenly mathematics. The forgotten art of spherical trigonometry*. Princeton University Press (Princeton, 2013).

MAREK LASSAK

INSTYTUT MATEMATYKI I FIZYKI  
UNIVERSITY OF SCIENCE AND TECHNOLOGY  
85-796 BYDGOSZCZ, POLAND  
*E-mail address:* marek.lassak@utp.edu.pl

MICHAŁMUSIELAK

INSTYTUT MATEMATYKI I FIZYKI  
UNIVERSITY OF SCIENCE AND TECHNOLOGY  
85-796 BYDGOSZCZ, POLAND  
*E-mail address:* michal.musielak@utp.edu.pl