

Inflationary perturbation spectrum in extended effective field theory of inflation

Jin Qiao ^{a,*}, Guang-Hua Ding^{a,†}, Qiang Wu^{a,‡}, Tao Zhu^{a,§} and Anzhong Wang^{a,b,¶}

^a *Institute for Theoretical Physics & Cosmology,*

Zhejiang University of Technology, Hangzhou, 310032, China

^b *GCAP-CASPER, Physics Department, Baylor University, Waco, TX 76798-7316, USA*

(Dated: April 30, 2022)

The effective field theory (EFT) of inflation provides a natural framework to study the new physical effects on primordial perturbation spectrum that is original from the higher derivative terms of the theory. Recently a healthy combination of high order operators has been formulated in an extension of the EFT of inflation, which allows one to avoid ghosts and leads to nonlinear dispersion relation for scalar perturbations. This paper devotes to study the new physics effects of these high order operators in primordial perturbations in this extended EFT of inflation by using the uniform asymptotic approximation method. For this purpose, we first construct the approximate and analytical solution of the mode functions for scalar perturbations. Because of the presence of the high order operators in the extended EFT of inflation, the perturbation modes may experience a period of non-adiabatic evolution before it crosses Hubble horizon. We show that it is these high order operators that may lead to the production of excited states and modifications of primordial perturbation spectrum. In particular, after deriving analytically the impact of these new effects on particle production rate and perturbation spectrum, we explain in details for different origins of these new effects.

I. INTRODUCTION

The cosmic inflation has achieved remarkable successes not only in solving several fundamental and conceptual problems (such as flatness, the horizon problem, and exotic relics) of the standard big bang cosmology, but most importantly, in providing a causal mechanism for generating structures in the Universe and the spectrum of cosmic microwave background (CMB) anisotropies [1–3] (see Ref. [4] for an updated review). All these predictions are now matched to cosmological observations with a spectacular precision [5–8]. These observations have provided confirmation of a nearly scale invariant power spectrum of adiabatic inflationary perturbations and support the inflationary paradigm with a scalar degree of freedom and a self-interacting potential that gives rise to a slow-roll phase during which the energy density of the matter field remains nearly constant and the spacetime behaves like a quasi-de Sitter.

While there are a lot of approaches to realize the inflation with scalar degrees of freedom, the EFT provides a general framework for describing the most generic single scalar field theory and the associated fluctuations on a quasi de-Sitter background [9, 10]. In this framework, the scalar field provides a clock that breaks time but preserves spatial diffeomorphism invariance. This allows one to construct the action of the theory around quasi de-Sitter background in terms of only three dimensional spatial diffeomorphism invariants and study the effects of

different terms. This is very similar to the case of Horava-Lifshitz (HL) theory of quantum gravity [11–15], in which the symmetry of the theory is broken from the general covariance down to the foliation-preserving diffeomorphisms. With this property, the action of the HL theory of quantum gravity has to be constructed in terms of three dimensional spatial diffeomorphism invariants. By construction, the HL theory of gravity is power-counting renormalizable and unitary, which is realized by including high-order spatial derivative operators (up to six in (3+1)- dimensional spacetimes) but excluding high-order temporal derivative operators. Such remarkable features have attracted a lot of attention and the observational effects of high-order operators on inflationary perturbation spectra have been extensively studied [16–19] (see [20] for an updated review).

Considering the similarity between two theories based on the analysis of symmetry, it is natural to consider the derivative terms with higher mass dimensions in the framework of EFT of inflation. These new terms provide an efficient way for parametrizing unknown high energy physics effects on low energy scale and producing various interesting inflationary models [21–25], such as inflation models in HL theory mentioned above, DBI inflation [23, 24], or Ghost inflation [25] (for EFT of bouncing universe, see [26, 27]). Recently, the EFT of inflation has been extended by adding higher spatial derivative terms up to the forth-order [28]. With this extension, the usual linear dispersion relation associated with the propagation of inflationary scalar perturbation has been changed to the nonlinear one. The high-order terms in such nonlinear dispersion relation represents the effects of the new high-order operators, which sets a new characteristic energy scale in the theory. Beyond this energy scale, the nonlinear dispersion relation is dominated by effects of the new operators, but reduces to the usual linear one at low energy scale of inflation.

* qiaojin@zjut.edu.cn

† dingguanghua@zjut.edu.cn

‡ wuq@zjut.edu.cn

§ zhut05@zjut.edu.cn

¶ anzhong.wang@baylor.edu

An important question now is whether the new high-order derivative operators in the extended EFT of inflation can leave any important effects in the propagation of inflationary perturbation modes and their primordial perturbation spectrum. An essential step to address this issue is to implement the cosmological perturbations in extended EFT of inflation and calculate the corresponding inflationary observables by evolving perturbation modes starting from high energy regime where higher-order terms dominated until the end of the slow-roll inflation. Such consideration has attracted much attention recently [28–30] and more generally the impact of the nonlinear dispersion relation on primordial perturbation spectrum has been studied extensively (see [31–36] and references therein). The main characteristic of the associated effects is that the cosmological perturbations can experience a period of non-adiabatic evolution in the high energy regime, and after that the perturbations are no longer in the adiabatic Bunch-Davies state, but instead excited states. These excited states in turn lead to particle production during inflation and modify primordial perturbation spectrum.

According to quantum field theory in the curved spacetime and the theory of WKB approximation, particle production can arise from non-adiabatic evolution of the associated field modes [37], which originals from the violation of the adiabatic condition of the WKB approximations. Indeed, this is exactly the case that could occur for cosmological perturbations in the extended EFT of inflation [28–30]. Such non-adiabatic evolution of primordial perturbations also occurs during quantum bounce in loop quantum cosmology, see [38–41] for examples. However, when the adiabatic condition is violated, it is in general impossible to study exact solution and the corresponding spectra for cosmological, and thus one has to use some approximate methods. Recently, we have developed a powerful method, the uniform asymptotic approximation method [33, 35, 42], to calculate precisely the quantum gravitational effects to the primordial perturbation spectra. The robustness of this method has been verified for calculating primordial spectra in k-inflation [43–45], and inflation with nonlinear dispersion relations [33, 35, 36] and quantum gravitational effects of loop quantum cosmology [46–48]. We note here that this method was first applied to inflationary cosmology in the framework of GR in [49–51], and then we have developed it, so it can be applied to more general case, including the ones with nonlinear dispersion relations [33, 36]. The main purpose of the present paper is to use this powerful method to de-

rive the inflationary observables in slow-roll inflation in the framework of the extended EFT of inflation. By analytically solving the equation of motion for scalar perturbation, we concentrate on how the high-order operators in the extended EFT affect the evolution of perturbation modes, with which we calculate the general expressions of perturbation spectrum at the end of slow-roll inflation. The main properties and the new effects of the high order operators have also been discussed in details. These expressions represent a significant improvement over the previous results obtained so far in the literature.

We organize the rest of the paper as follows. In Sec. II, we provide a brief introduction to the extended EFT of inflation and the corresponding equations of motion for both cosmological scalar and tensor perturbations. In Sec. III, we first discuss how the adiabatic condition is violated with the high-order operators introduced in the extended EFT of inflation, and then using the uniform asymptotic approximation to construct the analytical solution for scalar perturbation modes. With the analytical solution we derive the general expression of the corresponding perturbation spectrum. In Sec. IV, we study the main features of both the particle production rate during the inflation and the primordial perturbation spectrum. Our main conclusions and discussions are presented in Sec. V.

II. EXTENDED EFFECTIVE FIELD THEORY OF INFLATION

In this section, we present a brief review of the EFT of inflation with extension by including derivative terms with higher mass dimensions [28]. In general, the effective field theory provides a framework for describing the most generic single scalar field theory on a quasi de-Sitter background [9, 10]. Since the Friedmann-Robterson-Walker (FRW) background metric provides a preferred time foliation, we can in general write the action of a theory around the FRW background in terms of only three dimensional spatial diffeomorphism invariants. This allows us to construct the most general three dimensional spatial diffeomorphism invariant and study the effects of different terms. The basic building blocks of this construction include scalars like g^{00} and pure function of time $c(t)$, and the extrinsic curvature tensor $K^{\mu\nu}$ of the constant time hypersurfaces. Using these blocks, it is shown that the action of the effective field theory of inflation around a flat FRW background reads

$$S_{\text{eft}} = M_{\text{Pl}}^2 \int d^4x \sqrt{-g} \left\{ \frac{R}{2} + \dot{H} g^{00} - (3H^2 + \dot{H}) + \frac{M_2^4}{2M_{\text{Pl}}^2} (g^{00} + 1)^2 + \frac{\bar{M}_1^3}{2M_{\text{Pl}}^2} (g^{00} + 1) \delta K^\mu{}_\mu \right. \\ \left. - \frac{\bar{M}_2^2}{2M_{\text{Pl}}^2} (\delta K^\mu{}_\mu)^2 - \frac{\bar{M}_3^2}{2M_{\text{Pl}}^2} \delta K^\mu{}_\nu \delta K^\nu{}_\mu \right\}, \quad (2.1)$$

where $\delta K_{\mu\nu}$ denotes the perturbation of $K_{\mu\nu}$ about the flat FRW background.

The above action can be extended by introducing higher derivative terms. In this paper, we follow the ex-

tension proposed in [28] which includes operators with higher derivatives up to the fourth-order. The action for these additional terms is given by

$$\Delta S = \int d^4x \sqrt{-g} \left\{ \frac{\bar{M}_4}{2} \nabla g^{00} \nabla^\nu \delta K_{\mu\nu} - \frac{\delta_1}{2} (\nabla_\mu \delta K^{\nu\gamma}) (\nabla^\mu \delta K_{\nu\gamma}) - \frac{\delta_2}{2} (\nabla_\mu \delta K_\nu^\nu)^2 - \frac{\delta_3}{2} (\nabla_\mu \delta K_\nu^\mu) (\nabla_\gamma \delta K^{\gamma\nu}) - \frac{\delta_4}{2} \nabla^\mu \delta K_{\nu\mu} \nabla^\nu \delta K_\sigma^\sigma \right\}, \quad (2.2)$$

where \bar{M}_4 , δ_i ($i = 1, 2, 3, 4$) are free coefficients. The first term (the \bar{M}_4 term) contains three derivatives thus it leads to violation of the symmetry of time reverse. The other terms (δ_i terms) contain fourth-order derivative and lead to sixth-order corrections to the standard linear dispersion relation of scalar perturbations.

We would like to mention that, in writing the above action, one requires the perturbation in inflaton field ϕ to be zero, i.e., $\delta\phi = 0$. This is achieved by considering the following linear transformations under time diffeomorphism,

$$\tilde{t} = t + \xi^0(t, x^i), \quad \delta\tilde{\phi} = \delta\phi + \xi^0(t, x^i) \dot{\phi}_0(t), \quad (2.3)$$

which leads to a particular gauge (unitary gauge) with $\xi^0(t, x^i) = -\delta\phi/\dot{\phi}_0$ where there is no inflaton perturbation. Obviously equations (2.1) and (2.2) of the action that respect the unitary gauge has no time diffeomorphism invariance. In order to restore the time diffeomorphism invariance, one can introduce a Goldstone mode $\pi(t, x^i)$ and requires it transforms as $\pi(t, x^i) \rightarrow \pi(t, x^i) - \xi^0(t, x^i)$. By introducing the Goldstone mode, the perturbation of inflaton field is not required to be zero and it relates to $\pi(t, x^i)$ via $\delta\phi = \dot{\phi}_0 \pi$.

On the other hand, one must be careful when one deals with the higher derivative operators in the theory. The main concern of these terms is they in general produce the time derivative higher than two in the equation of motion, and according to Ostrogradski's theorem, a system with higher time derivatives is usually not-stable. For this reason, as analyzed in [28], one has to impose the condition

$$\delta_1 = 0 = \delta_2, \quad (2.4)$$

in order to avoid the higher time derivatives. In this paper, we will adopt this condition and disregard the δ_1 and δ_2 terms in the action.

A. Tensor perturbations

For tensor perturbations, the perturbed spacetime is set as

$$g_{ij} = a^2(\delta_{ij} + h_{ij}), \quad (2.5)$$

where h_{ij} represents the transverse and traceless tensor perturbations, which satisfies

$$h_i^i = 0 = \partial^i h_{ij}. \quad (2.6)$$

Then expanding the total action $S_{\text{tot}} = S_{\text{eff}} + \Delta S$ up to the second-order gives

$$S_h^2 = \frac{M_{\text{Pl}}^2}{8} \int dt d^3x a^3 (c_t^{-2} \partial_t h_{ij} \partial_t h^{ij} - a^{-2} \partial_k h_{ij} \partial^k h^{ij}), \quad (2.7)$$

where the effective sound speed c_t for tensor perturbations is given by

$$c_t^2 = \left(1 - \frac{\bar{M}_3^2}{M_{\text{Pl}}^2} \right)^{-1}. \quad (2.8)$$

In order to avoid superluminal propagation for tensor modes, one have to require

$$\bar{M}_3 < 0. \quad (2.9)$$

Variation of the action with respect to h_{ij} leads to the equation of motion

$$\frac{d^2 \mu_k^{(t)}(\eta)}{d\eta^2} + \left(c_t^2 k^2 - \frac{a''}{a} \right) \mu_k^{(t)}(\eta) = 0, \quad (2.10)$$

where $d\eta = dt/a$ is the conformal time and $\mu_k^{(t)}(\eta) = a M_{\text{Pl}} h_k / \sqrt{2}$ with h_k denoting the Fourier modes of the two helicities of the tensor perturbations. In the slow-roll inflationary background, if we treat all the slow-rolling quantities as approximately constant, the equation of motion for tensor perturbation can be solved analytically and expressed as a linear combination of Hankel functions,

$$\mu_k^{(t)}(\eta) \simeq \frac{\sqrt{-\pi\eta}}{2} \left[\alpha_k H_{\nu_t}^{(1)}(-c_t k \eta) + \beta_k H_{\nu_t}^{(2)}(-c_t k \eta) \right], \quad (2.11)$$

where $H_{\nu_t}^{(1)}(-c_t k \eta)$ and $H_{\nu_t}^{(2)}(-c_t k \eta)$ denote the first and second kind of Hankel functions respectively, and ν_t is a slow-roll quantity which is defined as

$$\nu_t^2 \equiv \eta^2 \frac{a''}{a} + \frac{1}{4}. \quad (2.12)$$

The coefficients α_k and β_k are two integral constants. In order to fix them, we impose the Bunch-Davis (BD) vacuum state at the initially

$$\mu_k^{(t)}(\eta) = \frac{1}{\sqrt{2k c_t}} e^{-i c_t k \eta}, \quad (2.13)$$

which leads to

$$\alpha_k = 1, \quad \beta_k = 0. \quad (2.14)$$

Then the tensor perturbation spectrum is calculated at the end of inflation, i.e., $\eta \rightarrow 0^-$. By using the asymptotic form of Hankel function when $\eta \rightarrow 0^-$, we obtain

$$\begin{aligned} \mathcal{P}_h &\equiv \frac{k^3}{\pi^2 M_{\text{Pl}}^2} \left| \frac{\mu_k^{(t)}(\eta)}{a(\eta)} \right|^2 \\ &= \frac{k^2}{4c_t \pi^3 a^2(\eta)} \Gamma^2(\nu_t) \left(-\frac{c_t k \eta}{2} \right)^{1-2\nu_t}. \end{aligned} \quad (2.15)$$

B. Scalar perturbations

As mentioned above, the Goldstone mode $\pi(t, x^i)$ can be introduced to restore the time diffeomorphism invariance of action and it also describes the scalar perturbations around the flat FRW background. Thus, in order to study the scalar perturbations, one can transform the action in (2.1) and (2.2) in unitary gauge to π -gauge by evaluating the action explicitly for π . Proceeding this procedure, one obtains

$$S^\pi = S_{\text{eff}}^\pi + \Delta S^\pi = \int d^4x \sqrt{-g} (\mathcal{L}_{\text{eff}} + \mathcal{L}_\Delta), \quad (2.16)$$

where

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= M_{\text{Pl}}^2 \dot{H} (\partial_\mu \pi)^2 + 2M_2^4 \dot{\pi}^2 - \bar{M}_1^3 H \left(3\dot{\pi}^2 - \frac{(\partial_i \pi)^2}{2a^2} \right) \\ &\quad - \frac{\bar{M}_2^2}{2} \left(9H^2 \dot{\pi}^2 - 3H^2 \frac{(\partial_i \pi)^2}{a^2} + \frac{(\partial_i^2 \pi)^2}{a^4} \right) \\ &\quad - \frac{\bar{M}_3^2}{2} \left(3H^2 \dot{\pi}^2 - H^2 \frac{(\partial_i \pi)^2}{a^2} + \frac{(\partial_j^2 \pi)^2}{a^4} \right), \end{aligned} \quad (2.17)$$

and

$$\begin{aligned} \mathcal{L}_{\Delta S} &= \frac{\bar{M}_4}{2} \left(\frac{k^4 H \pi^2}{a^4} + \frac{k^2 H^3 \pi^2}{a^2} - 9H^3 \dot{\pi}^2 \right) \\ &\quad - \frac{1}{2} \delta_3 \left(\frac{k^6 \pi^2}{a^6} + \frac{3H^2 k^4 \pi^2}{a^4} + \frac{H^2 k^2 \dot{\pi}^2}{a^2} - 9H^4 \dot{\pi}^2 \right) \\ &\quad - \frac{1}{2} \delta_4 \left(\frac{k^6 \pi^2}{a^6} + \frac{H^2 k^4 \pi^2}{2a^4} + \frac{9H^4 k^2 \pi^2}{2a^2} + \frac{3H^2 k^2 \dot{\pi}^2}{a^2} \right. \\ &\quad \left. + \frac{27}{2} H^4 \dot{\pi}^2 \right). \end{aligned} \quad (2.18)$$

Variation of the above action S^π with respect to π yields the equation of motion for the Goldstone mode π ,

$$A_1 \ddot{\pi}_k + B_1 \dot{\pi}_k + \left(F_1 \frac{k^2}{a^2} + D_1 \frac{k^4}{a^4} + C_1 \frac{k^6}{a^6} \right) \pi_k = 0, \quad (2.19)$$

where

$$A_1 = -2M_{\text{Pl}}^2 \dot{H} + 4M_2^4 - 6\bar{M}_1^3 H - 9H^2 \bar{M}_2^2 - 3H^2 \bar{M}_3^2 + 2H^4 F_0(k, \tau), \quad (2.20)$$

$$B_1 = -6\bar{M}_1^3 \dot{H} - 18\dot{H} H \bar{M}_2^2 - 6\dot{H} H \bar{M}_3^2 + 3H C_0 + H^5 \left[6F_0(k, \tau) + \frac{2k^2}{a^2 H^2} (\delta_3 + 3\delta_4) \right], \quad (2.21)$$

$$C_1 = \delta_3 + \delta_4, \quad (2.22)$$

$$D_1 = \bar{M}_2^2 + \bar{M}_3^2 + H^2 \frac{\delta_4}{2} + 3H^2 \delta_3 - \bar{M}_4 H, \quad (2.23)$$

$$F_1 = -2M_{\text{Pl}}^2 \dot{H} - \bar{M}_1^3 H - 3H^2 \bar{M}_2^2 - \bar{M}_3^2 H^2 + 3H^4 \left(\delta_3 + \frac{3}{2} \delta_4 \right) - \bar{M}_4 H^3. \quad (2.24)$$

and

$$F_0(k, \tau) \equiv \frac{9}{2} \delta_3 - \frac{27}{4} \delta_4 - \frac{k^2}{2a^2 H^2} (\delta_3 + \delta_4) - \frac{9}{2H} \bar{M}_4. \quad (2.25)$$

In order to simplify the above equation, we follow the same procedure of [52] by requiring $\delta_3 = -3\delta_4$ ¹ and defining $u_k = a\pi_k$, then one obtains

$$u_k'' + \left(\omega_k^2(\eta) - \frac{z''}{z} \right) u_k = 0, \quad (2.26)$$

where

$$\omega_k^2(\eta) = c_s^2 k^2 \left(1 + \frac{D_1}{G_1 c_s^2} \frac{k^2}{a^2} + \frac{C_1}{G_1 c_s^2} \frac{k^4}{a^4} \right), \quad (2.27)$$

and

$$c_s^2 \equiv \frac{F_1}{G_1}, \quad (2.28)$$

$$G_1 \equiv -2M_{\text{Pl}}^2 \dot{H} + 4M_2^4 - 6\bar{M}_1^3 H - 9H^2 \bar{M}_2^2 - 3H^2 \bar{M}_3^2 - \frac{81}{2} H^4 \delta_4 - 9\bar{M}_4 H^3. \quad (2.29)$$

In the de-Sitter limit, we also have

$$\frac{z''}{z} \simeq \frac{2}{\eta^2}, \quad aH \simeq -\eta. \quad (2.30)$$

It is worth noting that, after considering the high derivative terms in the action, the conventional linear dispersion relation becomes nonlinear. To study the nonlinear

¹ The discussion without this condition has been studied in [30] recently.

effects it is convenient to introduce a characteristic energy scale M_* beyond which the nonlinear terms dominate. For this purpose, we set

$$\hat{\alpha}_0 \left(\frac{H}{M_*} \right)^2 = \frac{D_1 H^2}{G_1 c_s^4}, \quad (2.31)$$

$$\hat{\beta}_0 \left(\frac{H}{M_*} \right)^4 = \frac{C_1}{G_1 c_s^6}, \quad (2.32)$$

where $\hat{\alpha}_0$ and $\hat{\beta}_0$ are two dimensionless constants in the de-Sitter limit. Since the inflationary observables are evaluated at the time when the perturbation modes exit the Hubble horizon with energy scales of order the Hubble scale H . Therefore, in order for the high derivative terms in the extended effective theory of inflation to be under control, one would need to require $H \ll M_*$ so that the horizon crossing can occur in the regime where the dispersion relation takes the standard linear form. With this point, it is convenient to write the equation of motion for scalar perturbation as

$$u_k''(\eta) + c_s^2 k^2 \left(1 + \hat{\alpha}_0 \epsilon_*^2 y^2 + \hat{\beta}_0 \epsilon_*^4 y^4 - \frac{2}{y^2} \right) u_k(\eta) = 0, \quad (2.33)$$

where we define $y \equiv -c_s k \eta$ and

$$\epsilon_* \equiv \frac{H}{M_*} \ll 1. \quad (2.34)$$

On the other hand, we also require that all the perturbation modes have to be stable in the ultraviolet limit, this leads to the condition for the healthy ultraviolet behavior

$$\hat{\beta}_0 > 0. \quad (2.35)$$

III. APPROXIMATE SOLUTION IN THE UNIFORM ASYMPTOTIC APPROXIMATION

A. Validity of WKB approximation

In this section, we start to study the evolution of the scalar perturbation during the inflation with the nonlinear dispersion relation given in the above. In general, an important feature of the nonlinear dispersion relation is that it could produce additional excited states for primordial perturbation on the sub-horizon scale during inflation. Before we study the generation of these excited states and their effects on primordial perturbation spectrum in details, we would like first to provide a qualitative analysis by using WKB approximation.

In general, the solution of the mode function $\mu_k^{(s,t)}(\eta)$ of the equation,

$$\mu_k^{(s,t)''}(\eta) + \Omega^2(\eta) \mu_k^{(s,t)}(\eta) = 0, \quad (3.1)$$

can be approximately given in terms of the WKB solutions

$$\mu_k^{(s,t)}(\eta) \simeq \frac{\alpha_k}{\sqrt{2\Omega(\eta)}} e^{-i \int \Omega(\eta) d\eta} + \frac{\beta_k}{\sqrt{2\Omega(\eta)}} e^{i \int \Omega(\eta) d\eta},$$

if the WKB condition

$$\left| \frac{3\Omega'^2}{4\Omega^4} - \frac{\Omega''}{2\Omega^3} \right| \ll 1, \quad (3.2)$$

is satisfied. Here the function

$$\begin{aligned} \Omega^2(\eta) &\equiv \omega_k^2(\eta) - \frac{z''}{z} \\ &= c_s^2 k^2 \left(1 + \hat{\alpha}_0 \epsilon_*^2 y^2 + \hat{\beta}_0 \epsilon_*^4 y^4 - \frac{2}{y^2} \right), \end{aligned} \quad (3.4)$$

and α_k and β_k are the two Bogoliubov coefficients, which can be determined by initial states. Generally an adiabatic state is a good choice, and if the WKB condition is satisfied during the whole process, we have

$$\alpha_k = 1, \quad \beta_k = 0. \quad (3.5)$$

However, in some cases, the WKB condition may be violated or not be fulfilled completely. Then, the non-adiabatic evolution of the mode $\mu_k(\eta)$ will produce excited states (i.e. particle production) during this process and eventually leads to a state with

$$\alpha_k \neq 1, \quad \beta_k \neq 0. \quad (3.6)$$

According to (3.3), there are several facts that can lead to the violation of the WKB condition. One case is $\Omega^2(\eta)$ contains zeros (represented as real turning points of Eq. (3.2) or case (a) and (b) in Fig. 1) or extreme closes to zero (complex conjugated turning points of Eq. (3.2) or case (c) in Fig. 1) in the intervals of interest. It is simple to check that when $\Omega^2(\eta)$ equals zero, the WKB condition in (3.3) becomes divergent. While for the linear dispersion relation, $\Omega^2(\eta)$ in general has only one zero (can be regarded as the time when the mode exit Hubble horizon), $\Omega^2(\eta)$ in Eq. (3.4) may have three zeros because of the presence of the high-order operators in the nonlinear dispersion relation. This fact make the violation of the WKB condition very complicated and heavy mathematics involved. When the extra two zeros are real, the WKB condition is strongly violated, while it is only weakly violated if these two zeros are complex conjugate. Another possible case that could violate the WKB condition is around the second-order pole about $y \rightarrow 0^+$ (i.e. $\eta \rightarrow 0^-$). For the second case, it is easy to check that

$$\left| \frac{3\Omega'^2}{4\Omega^4} - \frac{\Omega''}{2\Omega^3} \right| \rightarrow \frac{3}{8} \sim \mathcal{O}(1), \quad (3.7)$$

so the WKB condition is not satisfied very well. Recently we have developed a powerful method (the uniform asymptotic approximation) for better treatment to equations with $g(y)$ has both the turning points and poles. In the following subsections, we are going to apply this method and study the perturbation modes with high-order operator effects in details.

B. Classification of turning points

For this purpose, let us first write the equation of motion (2.26) for u_k into the standard form in the uniform asymptotic approximation as [33, 53, 54]

$$\frac{d^2 \mu_k}{dy^2} = \{g(y) + q(y)\} \mu_k, \quad (3.8)$$

where

$$g(y) + q(y) = \frac{2}{y^2} - \hat{\beta}_0 \epsilon_*^4 y^4 - \hat{\alpha}_0 \epsilon_*^2 y^2 - 1. \quad (3.9)$$

According to the theory of the second-order ordinary differential equation, the solution of above equation depends on poles and turning points of function $g(y)$ and $q(x)$. Analyzing the corresponding error control function associated with uniform asymptotic approximation around poles or turning points can provide guidance on how to determine the splitting of $g(y)$ and $q(x)$. For the equation of motion given in the above, we can see that $g(y)$ and $q(y)$ in general has two poles: one is located at $y = 0^+$ and the other is at $y = +\infty$. Using the analysis in the uniform asymptotic approximation [33, 53, 54], the splitting of functions $g(y)$ and $q(y)$ can be chosen by

$$\begin{aligned} q(y) &= -\frac{1}{4y^2}, \\ g(y) &= \frac{9}{4y^2} - 1 - \hat{\alpha}_0 \epsilon_*^2 y^2 - \hat{\beta}_0 \epsilon_*^4 y^4. \end{aligned} \quad (3.10)$$

Except the two poles at $y = 0^+$ and $y = +\infty$, $g(y)$ may also have zeros in the range $y \in (0, +\infty)$, which called turning points. By solving the equation $g(y) = 0$, we obtain the corresponding three turning points, which are

$$\begin{aligned} y_0 &= \left\{ \frac{-\hat{\alpha}_0}{3\hat{\beta}_0 \epsilon_*^2} \left[1 - 2\sqrt{1-Y} \cos\left(\frac{\theta}{3}\right) \right] \right\}^{1/2}, \\ y_1 &= \left\{ \frac{-\hat{\alpha}_0}{3\hat{\beta}_0 \epsilon_*^2} \left[1 - 2\sqrt{1-Y} \cos\left(\frac{\theta+2\pi}{3}\right) \right] \right\}^{1/2}, \\ y_2 &= \left\{ \frac{-\hat{\alpha}_0}{3\hat{\beta}_0 \epsilon_*^2} \left[1 - 2\sqrt{1-Y} \cos\left(\frac{\theta+4\pi}{3}\right) \right] \right\}^{1/2}, \end{aligned} \quad (3.11)$$

with $Y \equiv 3\hat{\beta}_0/\hat{\alpha}_0^2$ and

$$\cos \theta \equiv -\left(1 - \frac{3}{2}Y - \frac{27}{8}\hat{\alpha}_0 Y^2 \epsilon_*^2\right)^2. \quad (3.12)$$

Since we have three turning points here, without loss of generality we assume that $0 < y_0 < \text{Re}(y_1) \leq \text{Re}(y_2)$, in which y_0 is assumed to be a single and real turning point but y_1 and y_2 can be both real and single, real and double, or even complex conjugated. In general, the nature of y_1 and y_2 can be determined by the discriminant,

$$\Delta \equiv (\mathcal{Y} - 1)^3 + \left(1 - \frac{3}{2}\mathcal{Y} - \frac{27}{8}\hat{\alpha}_0 Y^2 \epsilon_*^2\right)^2, \quad (3.13)$$




FIG. 1. The function $g(y)$ defined by Eq.(3.10). These cases (a, b, c) correspond, respectively, three different real roots, one single and double roots, and a single and two complex conjugated roots. The case (d) has only a single real root with $\hat{\alpha}_0 < 0$. In all cases, $\hat{\beta}_0 > 0$ is assumed.

When $\Delta < 0$, the three turning points (y_0 , y_1 , and y_2) are all real and different. When $\Delta = 0$, there is one single real turning point (y_0) and one double real turning point ($y_1 = y_2$). When $\Delta > 0$, there is a single real turning point (y_0) and two complex conjugated turning points ($y_1^* = y_2$).

C. Approximate solution in the Uniform Asymptotic approximation

According to the discussion above, there are two poles and three turning points. In the uniform asymptotic approximation, the approximate solution depends on the types of turning points. Thus in the following we are going to discuss the solution around each turning point in details.

We first consider the single turning point y_0 , which lies in the range $(0, \text{Re}(y_1))$. In the uniform asymptotic approximation, the approximate of equation of motion around this single turning point can be expressed in terms of Airy type functions as

$$\mu_k(y) = a_0 \left(\frac{\xi(y)}{g(y)}\right)^{1/4} \text{Ai}(\xi) + b_0 \left(\frac{\xi(y)}{g(y)}\right)^{1/4} \text{Bi}(\xi), \quad (3.14)$$

where $\text{Ai}(\xi)$ and $\text{Bi}(\xi)$ are the Airy functions, a_0 and b_0 are two integration constants, ξ is a monotonous function of y which is given by

$$\xi(y) = \begin{cases} \left(-\frac{3}{2} \int_{y_0}^y \sqrt{g(y')} dy'\right)^{2/3}, & 0 < y \leq y_0, \\ -\left(\frac{3}{2} \int_{y_0}^y \sqrt{-g(y')} dy'\right)^{2/3}, & y_0 < y \leq \text{Re}(y_1), \end{cases} \quad (3.15)$$

The turning points y_1 and y_2 , in the uniform asymptotic approximation, we can treat them together. Around them the approximate solution of $\mu_k(y)$ can be expressed as

$$\begin{aligned} \mu_k(y) = & a_1 \left(\frac{\zeta^2 - \zeta_0^2}{-g(y)} \right)^{\frac{1}{4}} W \left(\frac{1}{2} \zeta_0^2, \sqrt{2}\zeta \right) \\ & + b_1 \left(\frac{\zeta^2 - \zeta_0^2}{-g(y)} \right)^{\frac{1}{4}} W \left(\frac{1}{2} \zeta_0^2, -\sqrt{2}\zeta \right). \end{aligned} \quad (3.16)$$

where $W(\frac{1}{2}\zeta_0^2, \sqrt{2}\zeta)$ and $W(\frac{1}{2}\zeta_0^2, -\sqrt{2}\zeta)$ are the parabolic

$$\begin{cases} \int_{y_1}^y \sqrt{-g(y^*)} dy^* = \frac{1}{2} \zeta \sqrt{\zeta^2 - \zeta_0^2} + \frac{\zeta_0^2}{2} \operatorname{arccosh} \left(-\frac{\zeta}{\zeta_0} \right), & y_0 < y < y_1, \\ \int_{y_1}^y \sqrt{-g(y^*)} dy^* = \frac{1}{2} \zeta \sqrt{\zeta^2 - \zeta_0^2} + \frac{\zeta_0^2}{2} \operatorname{arccosh} \left(-\frac{\zeta}{\zeta_0} \right), & y_1 < y < y_2, \\ \int_{y_2}^y \sqrt{-g(y^*)} dy^* = \frac{1}{2} \zeta \sqrt{\zeta^2 - \zeta_0^2} - \frac{\zeta_0^2}{2} \operatorname{arccosh} \left(\frac{\zeta}{\zeta_0} \right), & y_2 < y, \end{cases} \quad (3.18)$$

When y_1 and y_2 are complex conjugated, we have

$$\int_{\operatorname{Re}(y_1)}^y \sqrt{-g(y^*)} dy^* = \frac{1}{2} \zeta \sqrt{\zeta^2 - \zeta_0^2} - \frac{\zeta_0^2}{2} \ln \left(\frac{\zeta + \sqrt{\zeta^2 - \zeta_0^2}}{|\zeta_0|} \right), \quad (3.19)$$

With the approximate solutions around each of turning points given above, one can turn to the procedure of connecting them together. Before we proceed the procedure for determining all the integration constants (a_1, b_1) and (a_0, b_0), we first need to specify the initial condition of the perturbation modes. As we have mentioned at the end of Sec. II (see Eq. (2.35)), in order to obtain a healthy ultraviolet limit, we have $\hat{\beta}_0 > 0$. This allows us to impose the usual adiabatic Bunch-Davies vacuum state when $y \rightarrow +\infty$ on perturbation modes,

$$\begin{aligned} \lim_{x \rightarrow +\infty} \mu_k(y) &= \frac{1}{\sqrt{2\omega_k}} e^{-i \int \omega_k d\eta} \\ &= \sqrt{\frac{1}{2k}} \left(\frac{1}{-g} \right)^{1/4} \exp \left(-i \int_{y_i}^y \sqrt{-g} dy \right). \end{aligned} \quad (3.20)$$

Connecting the initial condition with the approximate solution (3.16) around turning points y_1 and y_2 in the range of $y \gg \operatorname{Re}(y_2)$, we find

$$a_1 = 2^{-3/4} k^{-1/2} \kappa^{-1} \left(\frac{1}{2} \zeta_0^2 \right), \quad (3.21)$$

$$b_1 = -i 2^{-3/4} k^{-1/2} \kappa \left(\frac{1}{2} \zeta_0^2 \right), \quad (3.22)$$

where $\kappa(\frac{1}{2}\zeta_0^2)$ is given by

$$\kappa \left(\frac{1}{2} \zeta_0^2 \right) \equiv \sqrt{1 + e^{\pi \zeta_0^2}} - e^{\pi \zeta_0^2/2}. \quad (3.23)$$

cylinder functions, a_1 and b_1 are two integration constants, and ζ_0^2 is defined as

$$\zeta_0^2 = \pm \frac{2}{\pi} \left| \int_{y_1}^{y_2} \sqrt{g(y)} dy \right|. \quad (3.17)$$

Here \pm correspond to $y_{1,2}$ are both real and complex conjugated respectively. We observe that ζ_0^2 is sensitive to the nature of the turning points y_1 and y_2 . ζ_0^2 is positive when y_1 and y_2 are real and it is negative if y_1 and y_2 are complex conjugated. The variable ξ is a monotonous increasing function of y . When y_1 and y_2 are both real, ζ is related to y via

Then we turn to match the approximate solution (3.14) around the single turning point y_1 and approximate solution (3.16) around turning points y_1 and y_2 in their overlap region between y_0 and y_1 . This procedure determine the coefficients a_0 and b_0 as

$$\begin{aligned} a_0 &= \sqrt{\frac{\pi}{2k}} \left[\kappa^{-1} \left(\frac{1}{2} \zeta_0^2 \right) \sin \mathfrak{B} - i \kappa \left(\frac{1}{2} \zeta_0^2 \right) \cos \mathfrak{B} \right], \\ b_0 &= \sqrt{\frac{\pi}{2k}} \left[\kappa^{-1} \left(\frac{1}{2} \zeta_0^2 \right) \cos \mathfrak{B} + i \kappa \left(\frac{1}{2} \zeta_0^2 \right) \sin \mathfrak{B} \right], \end{aligned} \quad (3.24)$$

where

$$\mathfrak{B} \equiv \int_{y_0}^{y_1} \sqrt{-g} dy + \phi(\zeta_0^2/2), \quad (3.25)$$

with $\phi(x) = \frac{\pi}{2} - \frac{\pi}{4} \ln x^2 + \frac{1}{2} \operatorname{ph} \Gamma(\frac{1}{2} + ix)$, where the phase $\Gamma(\frac{1}{2} + ix)$ is zero when $x = 0$, and is determined by continuity otherwise.

Having matching the approximate solutions around each turning point and initial condition together, we determine all the integration constants appear in the approximate solutions. Therefore, with these solutions, we are able to track the perturbation modes starting from the high energy regime until the end of the slow-roll inflation. Let us now consider some representative cases. The cases with three different single turning points and one and two complex conjugate roots are plotted, respectively, in the left panel and right panel of Fig. 2. From these figures, one can see clearly that the exact

solutions are well approximated by the analytical ones. Many other cases have also been considered, the results show that two solutions (numerical and analytical) are extremely well approximated.

IV. NON-ADIABATIC EFFECTS ON POWER SPECTRUM OF SCALAR PERTURBATIONS

As we have mentioned in the above section, the presence of the extra two turning points leads to the violation of the adiabatic evolution of the perturbation modes. This fact has also been indicated and discussed in details in Refs. [28, 29, 33, 36]. As shown and pointed out in these works, the non-adiabatic evolution of the perturbation modes lead to highly populated excited states and amplify the standard perturbation spectrum with linear dispersion relation. Since the presence of the extra two turning points is a direct consequence of the high order operators included in the extended EFT of inflation, these adiabatic effects can be regarded as the new high energy effects due to the high-order operators. In this section, by using the analytical solution we derived in the above, we discuss in details how these new effects affect both the generation of excited states and the primordial perturbation spectrum.

A. Generation of excited states and particle production rate

Let us first consider the non-adiabatic effects on the generation of the excited state. For this purpose, when the perturbation modes are inside the Hubble horizon (between y_0 and y_1), one approximately has $g(y) \simeq -\omega_k^2/k^2$ and

$$\int_{y_0}^y \sqrt{-g(y')} dy' \simeq - \int_{\eta_0}^{\eta} \omega_k(\eta') d\eta'. \quad (4.1)$$

Then using the asymptotic form of Airy function and above relation, the approximate solution (3.14) can be casted into the form

$$\begin{aligned} \mu_k(\eta) \simeq & \frac{1}{\sqrt{2\omega_k}} \sqrt{\frac{k}{2\pi}} \frac{1}{i} \left\{ (ia_0 - b_0) e^{i \int_{\eta_0}^{\eta} \omega_k(\eta') d\eta' - i \frac{\pi}{4}} \right. \\ & \left. + (ia_0 + b_0) e^{-i \int_{\eta_0}^{\eta} \omega_k(\eta') d\eta' - i \frac{\pi}{4}} \right\}. \end{aligned} \quad (4.2)$$

From this form one can immediately identify the Bogoliubov coefficients of the excited modes at the subhorizon scale as

$$\begin{aligned} |\beta_k|^2 &= \frac{k}{2\pi} |ia_0 - b_0|^2 \\ &= \frac{1}{4} (\kappa^2 + \kappa^{-2} - 2) \\ &= e^{\pi \xi_0^2}. \end{aligned} \quad (4.3)$$

Here β_k is the Bogoliubov coefficient that measures the particle production rate. From the above expression, we

can see that β_k^2 is determined by the quantity ξ_0^2 . Before proceeding further, several remarks about the nature of ξ_0^2 now are in order. First, the sign of ξ_0^2 is sensitive to the nature of the turning points y_1 and y_2 , for which we classify them into several classes.

- When y_1 and y_2 are both single and real, we have $\xi_0^2 > 0$, which implies that the particle production during the process is exponentially enhanced. As we have shown in Sec. III. B, for this case to happen, one must require the discriminant $\Delta < 0$. This corresponds to a requirement on the parameters in the high order operators.
- When y_1 and y_2 are two real and equal, i.e., $y_1 = y_2$, we have $\xi_0^2 = 0$. Then, we have $\beta_k^2 = 1$.
- When y_1 and y_2 are complex conjugate, i.e., $y_1 = y_2^*$, ξ_0^2 is negative. This implies that the particle production is exponentially suppressed.

Now an important question arises for the case of the exponentially enhanced particle production rate, namely, whether or not the backreaction of the excited modes is small enough to allow inflation to last long enough. According to the analysis in [55, 56], in order to avoid large backreactions, one has to impose the condition

$$|\beta_k|^2 \lesssim 8\pi \frac{H_{inf}^2 M_{Pl}^2}{M_*^4}, \quad (4.4)$$

where H_{inf} is the energy scale of the inflation and the Planck 2015 data yield the constraint $H_{inf}/M_{Pl} \leq 3.5 \times 10^{-5}$ [7]. Thus, if we take $H_{inf}/M_{Pl} \sim 2 \times 10^{-3}$, one can infer that

$$|\beta_k|^2 \lesssim \mathcal{O}(1). \quad (4.5)$$

And the modified factor can be expressed as

$$\mathcal{A} \simeq |\alpha_k + \beta_k|^2. \quad (4.6)$$

It is easy to obtain that

$$\sqrt{1 + |\beta_k|^2} - |\beta_k| \lesssim \sqrt{\mathcal{A}} \lesssim |\beta_k| + \sqrt{1 + |\beta_k|^2}, \quad (4.7)$$

which leads to the constraint on the modified factor \mathcal{A} based on the analysis of the backreaction,

$$3 - 2\sqrt{2} \lesssim \mathcal{A} \lesssim 3 + 2\sqrt{2}. \quad (4.8)$$

B. Scalar perturbation spectrum

With the solutions determined in the above, we are able to calculate the perturbation spectrum for scalar perturbation at the end of the slow-roll inflation in the limit $y \rightarrow 0^+$. In this limit, the scalar perturbation is described by the approximate solution (3.14) around y_0 . This solution also can be represented as a combination of one growing and one decaying modes, in which $\text{Bi}(\xi)$ is the growing mode and $\text{Bi}(\xi)$ represents the decaying




FIG. 2. Comparison of the analytical approximate solutions in the uniform asymptotic approximation to the numerical (exact) solutions in a de Sitter background.

mode. When $y \rightarrow 0^+$, only the growing mode is relevant, then using the asymptotic form of $\text{Bi}(\xi)$ we find

$$u_k \simeq b_0 \left(\frac{1}{\pi^2 g(y)} \right)^{1/4} \exp \left(\int_y^{y_0} \sqrt{g(y)} dy \right), \quad (4.9)$$

where b_0 is given by Eq.(3.24). Then, the scalar power spectrum is given by

$$\begin{aligned} \mathcal{P}_s &\equiv \frac{k^3}{2\pi^2} \left| \frac{u_k(y)}{z} \right|^2 \\ &= \mathcal{A}(k) \left(\frac{k^2 y}{9\pi^2 z^2} \right) \exp \left(2 \int_y^{y_0} \sqrt{g(y)} dy \right), \end{aligned} \quad (4.10)$$

where

$$\begin{aligned} \mathcal{A} &\equiv \frac{2k|b_0|^2}{\pi} \\ &= 1 + 2e^{\pi\zeta_0^2} + 2e^{\pi\zeta_0^2/2} \sqrt{1 + e^{\pi\zeta_0^2}} \cos 2\mathfrak{B}. \end{aligned} \quad (4.11)$$

Obviously the perturbation spectrum can be modified due to the high-order operators included in the extended EFT of inflation by two quantities: the modified factor $\mathfrak{A}(k)$ and the exponential integration of $\sqrt{-g(y)}$ from y_0 to 0. In the following we discuss these two effects in details.

1. Non-adiabatic effects on perturbation spectrum

The modified factor $\mathcal{A}(k)$ measures the contribution due to presence of the two turning points y_1 and y_2 . For the perturbation modes with linear dispersion relation in the EFT of inflation without high-order operators, the equation of motion can in general has only one single turning point. Therefore, the modified factor $\mathcal{A}(k)$ represents a direct effect of the presence of high-order operators included in the extended EFT of inflation, namely, the terms with \bar{M}_4 , δ_3 , and δ_4 in the action (2.2). We

observe that \mathcal{A} depends on the quantity ζ_0^2 , which related to the strength of the violation of adiabatic evolution because of the extra two turning points, as we mentioned above. When ζ_0^2 is positive and large, which corresponds to the case in which both y_1 and y_2 are real (i.e. case (a) in Fig. 1), we have

$$e^{\pi\zeta_0^2} \gg 1. \quad (4.12)$$

Then the modified factor \mathcal{A} now reads

$$\mathcal{A} \simeq 2e^{\pi\zeta_0^2} (1 + \cos 2\mathfrak{B}), \quad (4.13)$$

which indicates the power spectrum is exponentially enhanced for most of parameter space. When $\zeta_0^2 = 0$, which corresponds to the double turning point case that $y_1 = y_2$ (i.e. case (b) in Fig. 1), we have

$$\mathcal{A} = 3 + 2\sqrt{2} \cos 2\mathfrak{B}. \quad (4.14)$$

For the case in which ζ_0^2 is negatively large, which corresponds to the case in which the turning points y_1 and y_2 are complex conjugated, we have

$$e^{\pi\zeta_0^2} \ll 1. \quad (4.15)$$

Thus the modified factor $\mathcal{A}(k)$ is

$$\mathcal{A}(k) = 1 + 2e^{\pi\zeta_0^2/2} \cos 2\mathfrak{B} + \mathcal{O}(e^{\pi\zeta_0^2}), \quad (4.16)$$

which reduces to the usual one with only one single turning point y_0 . In Fig. 3 and 4, we plot the behavior of the modified factor $\mathcal{A}(k)$ with respect to different parameters. All these figures show clearly the modified factor \mathcal{A} enhances when y_1 and y_2 are both real and single, and reduces to one if y_1 and y_2 are complex conjugated. Our analytical results presented here are also in agreement with the numerical results obtained in [28].

One essential question related to the enhanced perturbation spectrum is if these non-adiabatic effects can lead to the violation of the nearly scale invariance of the

scalar spectrum. In fact, as shown in [36], if we assume that both the parameters $\hat{\alpha}_0$ and $\hat{\beta}_0$ are slow-varying, the resulting primordial perturbation spectrum is still nearly scale invariant. Therefore, the presence of high order operators in the theory can mainly affect the overall amplitude of the perturbation spectrum.

2. Impact of the exponential integration of $\sqrt{-g(y)}$

Another effects of high order operators it may also change the exponential integral of $\sqrt{-g(y)}$ over range from y_0 to 0, in comparison to that with linear dispersion relation. In order to see this fact clearly, let us first consider the integral with linear dispersion relation,

$$\mathcal{M}_0 = \exp\left(2 \int_y^{y_0} \sqrt{\frac{9}{4y'^2} - 1} dy'\right), \quad (4.17)$$

Here the turning point $y_0 = \frac{3}{2}$ in de-Sitter background. For the nonlinear dispersion relation, this integral changes into

$$\mathcal{M} = \exp\left(2 \int_y^{y_0} \sqrt{g(y')} dy'\right). \quad (4.18)$$

With the above two definitions, we can cast the perturbation spectrum in (4.10) into the form

$$\mathcal{P}_s = \mathcal{A} \times \frac{\mathcal{M}}{\mathcal{M}_0} \times \mathcal{P}_s^{\text{GR}}, \quad (4.19)$$

where $\mathcal{P}_s^{\text{GR}}$ denotes the standard power-law spectrum in the framework in GR. While the effects of \mathcal{A} , which related to the non-adiabatic effects due to the presence of the two extra turning points, is discussed in the above subsection, here we concentrate on the effects by $\mathcal{M}/\mathcal{M}_0$.

From Eq. (4.18), we can see that the ratio $\mathcal{M}/\mathcal{M}_0$ is more sensitive to the magnitude of the parameter $\epsilon_* = \frac{H}{M_*}$ rather than the parameters $\hat{\alpha}_0$ and $\hat{\beta}_0$. When $\epsilon_* \ll 1$, as we have assumed in Eq. (2.34), we find the turning point y_0 now can be approximated by

$$y_0 = \frac{3}{2} + \mathcal{O}(\epsilon_*). \quad (4.20)$$

During the range between 0 and y_0 , we also have

$$\sqrt{g(y)} = \sqrt{\frac{9}{4y^2} - 1} + \mathcal{O}(\epsilon_*). \quad (4.21)$$

With above two approximated expressions, we have

$$\frac{\mathcal{M}}{\mathcal{M}_0} = 1 + \mathcal{O}(\epsilon_*). \quad (4.22)$$

This indicates that if $\epsilon_* \ll 1$, the exponential integral in (4.18) does not lead to any significant effects on the primordial perturbation spectrum. The only effects come from the modified factor \mathcal{A} .

If we relax the condition $\epsilon_* \ll 1$ to $\epsilon \lesssim \mathcal{O}(1)$, namely, we consider $\hat{\alpha}_0 \epsilon_*^2 \lesssim \mathcal{O}(1)$ and $\hat{\beta}_0 \epsilon_*^4 \lesssim \mathcal{O}(1)$, the above conclusion will significant changes. To see this clearly, in Fig. 5 we plot the $g(y)$ for different values of parameters and compare it with the one of linear dispersion relation. When both $\hat{\alpha}_0 \epsilon_*^2$ and $\hat{\beta}_0 \epsilon_*^4$ are positive, which corresponds to case (d) in Fig. 1, $g(y)$ only has one single turning point and the corresponding modified factor $\mathcal{A} \simeq 1$. However, as shown in the top panel of Fig. 5, this case leads to a shift of y_0 from $\frac{3}{2}$ in the linear case to a smaller value. As a result, the curve of $g(y)$ with the nonlinear dispersion relation is always beneath the one with linear dispersion relation between y_0 and 0. In this case, it is obvious to conclude that $\mathcal{M}/\mathcal{M}_0 < 1$ is significant, which implies the perturbation spectrum is suppressed in comparison to the standard one in GR.

When $\hat{\alpha}_0 \epsilon_*^2$ is negative, thing becomes a little bit complicated. In this case, the shift of turning point y_0 from $\frac{3}{2}$ is a result of competition between $\hat{\alpha}_0 \epsilon_*^2$ and $\hat{\beta}_0 \epsilon_*^4$. The former one can make $y_0 > \frac{3}{2}$ and the later one make it smaller. When the effects of $\hat{\alpha}_0 \epsilon_*^2$ is larger than that of $\hat{\beta}_0 \epsilon_*^4$, y_0 becomes larger than $3/2$, otherwise it becomes smaller. However, from the shift of y_0 itself we still can not conclude if the ratio $\mathcal{M}/\mathcal{M}_0$ is larger than one or not. As shown in the middle and the bottom panels of Fig. 5, even when $y_0 < 3/2$, the curve of $g(y)$ is not always beneath the one with linear dispersion relation during $(0, y_0)$. This makes the analysis of the ratio $\mathcal{M}/\mathcal{M}_0$ analytically in this case very difficult. Anyway, the ratio $\mathcal{M}/\mathcal{M}_0$ can be significant derived from one in this case and it could be either larger than one or smaller than one, depending on the values of the parameters.

In fact, the impact of high-order operators on perturbation spectrum with larger values of $\hat{\alpha}_0 \epsilon_*^2$ and $\hat{\beta}_0 \epsilon_*^4$ has also been studied numerically in [28]. The results we presented in this subsection is in agreement with their numerical analysis. Here we would like to emphasize that our analytical results show clearly the impact of high order operators of the extended EFT of inflation on perturbation spectrum original from two different effects, one measures by the modified factor \mathcal{A} and the other measures by the ratio of the exponential integrals $\mathcal{M}/\mathcal{M}_0$. In addition, we note that, the suppression or enhancement on the perturbation spectrum due to the ratio $\mathcal{M}/\mathcal{M}_0$ is not an effect of the excited state generated by the non-adiabatic evolution of modes around y_1 and y_2 . These effects are mainly original from the derivation of the turning point y_0 from that with the linear dispersion relation.

V. CONCLUSION AND OUTLOOK

In this paper, we have provided a detailed and analytical study of the effects of high order operators on primordial comoving curvature perturbation. These high order operators are naturally included as a combination in the




FIG. 3. The modified factor \mathcal{A} as a function of $\hat{\beta}_0$ with $\hat{\alpha}_0$ and ϵ_* fixed. In both figures, the modified factor changes from the two real turning points region (corresponds to $\mathcal{A} \gg 1$) to the two complex conjugated region (corresponds to $\mathcal{A} \simeq 1$). Left panel: $\hat{\alpha}_0 = 2$ and $\epsilon_* = 0.01$. Right panel: $\hat{\alpha}_0 = 2.5$ and $\epsilon_* = 0.01$.




FIG. 4. The modified factor \mathcal{A} as a function of $\hat{\beta}_0$ with $\hat{\alpha}_0 = 2$ for several values of ϵ_* .

framework of the extended EFT of inflation. We show clearly that the new effects of these high order operators can naturally generate an excited state on the primordial comoving curvature perturbation rather than the usual BD vacuum state at subhorizon scale before the perturbation modes cross Hubble horizon. As we showed in Sec. IV, this excited state can produce enhanced effects

if these high order operators can lead to strong violation of adiabaticity of the perturbation modes (corresponding to two extra real and single turning points y_1 and y_2), but reduce to the BD state when the violation is weak (corresponding to two complex conjugated turning points). With this excited state, we calculated explicitly these non-adiabatic effects on primordial curvature perturbation spectrum and showed that the modification of the spectrum come from two effects, one is from the modified factor $\mathcal{A}(k)$ and the other is from the exponential integral of $\sqrt{g(y)}$ over the range of $(0, y_0)$. In particular, we showed clearly that the modified factor \mathcal{A} measures the contribution due to presence of the two turning points y_1 and y_2 , which represents a direct effects of the inclusion of the high order operators in the theory. For the effects of exponential integral, which either leads to suppression or enhancement on the perturbation spectrum, are mainly original from the derivation of the turning point y_0 from that with respect to the linear dispersion relation.

ACKNOWLEDGEMENTS

This work is supported in part by National Natural Science Foundation of China with the Grants Nos. 11675143 (Q.W. & T.Z.) and 11675145 (A.W.).

-
- [1] A. H. Guth, Inflationary universe: A possible solution to the horizon and flatness problems, *Phys. Rev. D* **23**, 347 (1981).
 - [2] A. A. Starobinsky, A new type of isotropic cosmological models without singularity. *Phys. Lett. B* **91**, 99-102 (1980).
 - [3] K. Sato, First-order phase transition of a vacuum and the expansion of the Universe, *Mon. Not. R. Astron. Soc.* **195**, 467-479 (1981).
 - [4] Daniel Baumann, TASI Lectures on Inflation, [arXiv:0907.5424](https://arxiv.org/abs/0907.5424).
 - [5] E. Komatsu, K. M. Smith, J. Dunkley, C. L. Bennett, et. al., Seven-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation, *ApJS* **192**, 18 (2011).
 - [6] Y. Akrami, F. Arroja, M. Ashdown, J. Aumont, et. al. (Planck Collaboration), Planck 2018 results. X. Constraints on inflation, [arXiv: 1807.06211](https://arxiv.org/abs/1807.06211) [astro-ph].




FIG. 5. Comparison of $g(y)$ with nonlinear and linear dispersion relations in the range of from $(0, y_0)$ for different values of parameters.

[7] P. A. R. Ade, N. Aghanim, M. Arnaud, F. Arroja, et. al. (Planck Collaboration), Planck 2015. XX. Constraints on inflation, [arXiv: 1502.02114 \[astro-ph\]](#).

[8] P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, et. al. (Planck Collaboration), Planck 2013 results. XXII. Constraints on inflation, [Astron. Astrophys. 571, A22 \(2014\)](#).

[9] C. Cheung, A. L. Fitzpatrick, J. Kaplan, L. Senatore, and P. Creminelli, The effective field theory of inflation, [J. High Energy Phys. 03, 014 \(2008\)](#).

[10] S. Weinberg, Effective field theory for inflation, [Phys. Rev. D, 77\(12\):123541, June 2008](#).

[11] P. Hořava. Quantum gravity at a Lifshitz point, [Phys. Rev. D, 79\(8\):084008, April 2009](#).

[12] P. Hořava, and C. M. Melby-Thompson, General covariance in quantum gravity at a Lifshitz point, [Phys. Rev. D, 82\(6\):064027, 2010](#).

[13] T. Zhu, Q. Wu, A. Wang, and F.-W. Shu, U(1) symmetry and elimination of spin-0 gravitons in Horava-Lifshitz gravity without the projectability condition, [Phys. Rev. D, 84, 101502 \(2011\)](#).

[14] T. Zhu, F.-W. Shu, Q. Wu, and A. Wang. General covariant Horava-Lifshitz gravity without projectability condition and its applications to cosmology, [Phys. Rev. D, 85\(4\):044053, February 2012](#).

[15] K. Lin, S. Mukohyama, A. Wang, and T. Zhu. Post-Newtonian approximations in the Hořava-Lifshitz gravity with extra U(1) symmetry, [Phys. Rev. D, 89\(8\):084022, April 2014](#).

[16] Y. Huang, A. Wang, R. Yousefi, and T. Zhu. Primordial non-Gaussianity of gravitational waves in HoVision Res.ava-Lifshitz gravity, [Phys. Rev. D, 88\(2\):023523, July 2013](#).

[17] T. Zhu, Y. Huang, and A. Wang. Inflation in general covariant Hořava-Lifshitz gravity without projectability, [J. High Energy Phys., 2013\(1\):1–23, January 2013](#).

[18] T. Zhu, W. Zhao, Y. Huang, A. Wang, and Q. Wu. Effects of parity violation on non-Gaussianity of primordial gravitational waves in Hořava-Lifshitz gravity, [Phys. Rev. D, 88\(6\):063508, September 2013](#).

[19] A. Wang, Q. Wu, W. Zhao, and T. Zhu. Polarizing primordial gravitational waves by parity violation, [arXiv:1208.5490, August 2012](#).

[20] A. Wang, Hořava gravity at a Lifshitz point: A progress report, [Int. J. Mod. Phys. D 26, 1730014 \(2017\)](#).

[21] X. Tong, Y. Wang, and S. Zhou, On the effective field theory for quasi-single field inflation, [J. Cosmol. Astropart. Phys. 11, 045 \(2017\)](#).

[22] J.-O. Gong, M.-S. Seo, and S. Sypsas, Higher derivatives and power spectrum in effective single field inflation, [J. Cosmol. Astropart. Phys. 03 \(2015\) 009](#).

[23] M. Alishahiha, E. Silverstein, and D. Tong, DBI in the sky: Non-Gaussianity from inflation with a speed limit, [Phys. Rev. D, 70\(12\):123505, December 2004](#).

[24] S. E. Shandera, and S.-H. H. Tye, Observing brane inflation, [J. Cosmol. Astropart. Phys., 2006\(05\):007, May 2006](#).

[25] N. Arkani-Hamed, P. Creminelli, S. Mukohyama, and M. Zaldarriaga, Ghost inflation, [J. Cosmol. Astropart. Phys., 2004\(04\):001, April 2004](#).

[26] Y. Cai, Y. Wan, H.-G. Li, T. Qiu, and Y.-S. Piao, The Effective Field Theory of nonsingular cosmology, [J. High Energy Phys. 01, 90 \(2017\)](#).

[27] Y. Cai and Y.-S. Piao, Higher order derivative coupling to gravity and its cosmological implications, [Phys. Rev. D 96, \(2017\) 124028](#).

[28] A. Ashoorioon, R. Casadio, M. Cicoli, G. Geshnizjani, and H. J. Kim, Extended Effective Field Theory of Inflation, [J. High Energy Phys., 2018\(2\), February 2018](#).

[29] A. Ashoorioon, R. Casadio, G. Geshnizjani, and H. J. Kim, Getting Super-Excited with Modified Dispersion Relations, [J. Cosmol. Astropart. Phys., 2017\(09\):008–008, September 2017](#).

[30] A. Ashoorioon, Non-Unitary Evolution in the General Extended EFT of Inflation & Excited Initial States, [ArXiv180706511 Astro-Ph Physicsgr-Qc Physicshep-Ph Physicshep-Th, July 2018](#).

- [31] A. Ashoorioon, D. Chialva, and U. Danielsson, Effects of nonlinear dispersion relations on non-Gaussianities, *J. Cosmol. Astropart. Phys.*, 2011(06):034–034, June 2011.
- [32] A. Ashoorioon, and G. Shiu. A note on calm excited states of inflation, *J. Cosmol. Astropart. Phys.*, 2011(03):025, 2011.
- [33] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng, Inflationary cosmology with nonlinear dispersion relations, *Phys. Rev. D*, 89(4):043507, February 2014.
- [34] T. Zhu, and A. Wang, Gravitational quantum effects in light of BICEP2 results, *Phys. Rev. D*, 90(2):027304, July 2014.
- [35] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng, Gravitational quantum effects on power spectra and spectral indices with higher-order corrections, *Phys. Rev. D*, 90(6):063503, September 2014.
- [36] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, and Q. Sheng, High-order primordial perturbations with quantum gravitational effects, *Phys. Rev. D*, 93(12):123525, June 2016.
- [37] S. Winitzki, Cosmological particle production and the precision of the WKB approximation, *Phys. Rev. D*, 72(10):104011, November 2005.
- [38] Q. Wu, T. Zhu, and A. Wang, Non-adiabatic Evolution of Primordial Perturbations and non-Gaussianity in Hybrid Approach of Loop Quantum Cosmology, *ArXiv180903172 Astro-Ph Physicsgr-Qc Physicshep-Th*, September 2018.
- [39] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, and Q. Sheng, Primordial non-Gaussianity and power asymmetry with quantum gravitational effects in loop quantum cosmology, *Phys. Rev. D*, 97(4):043501, February 2018.
- [40] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng, Pre-inflationary universe in loop quantum cosmology, *Phys. Rev. D*, 96(8):083520, October 2017.
- [41] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, and Q. Sheng, Universal features of quantum bounce in loop quantum cosmology, *Phys. Lett. B*, 773:196–202, October 2017.
- [42] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng, Constructing analytical solutions of linear perturbations of inflation with modified dispersion relations. *Int. J. Mod. Phys. A*, 29(23):1450142, September 2014.
- [43] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng, Power spectra and spectral indices of k-inflation: High-order corrections. *Phys. Rev. D*, 90(10):103517, November 2014.
- [44] J. Martin, C. Ringeval, and V. Vennin, K-inflationary power spectra at second order, *J. Cosmol. Astropart. Phys.*, 2013(06):021, June 2013.
- [45] C. Ringeval, Dirac-Born-Infeld and k-inflation: The CMB anisotropies from string theory, *J. Phys. Conf. Ser.*, 203:012056, January 2010.
- [46] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, Q. Sheng, and Q. Wu, Scalar and tensor perturbations in loop quantum cosmology: High-order corrections, *J. Cosmol. Astropart. Phys.*, 2015(10):052, 2015.
- [47] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, Q. Sheng, and Q. Wu, Detecting Quantum Gravitational Effects of Loop Quantum Cosmology in the Early Universe? *ApJ*, 807(1):L17, 2015.
- [48] T. Zhu, A. Wang, K. Kirsten, G. Cleaver, Q. Sheng, and Q. Wu. Inflationary spectra with inverse-volume corrections in loop quantum cosmology and their observational constraints from Planck 2015 data. *J. Cosmol. Astropart. Phys.*, 2016(03):046, 2016.
- [49] S. Habib, K. Heitmann, G. Jungman, and C. Molina-París, The Inflationary Perturbation Spectrum, *Phys. Rev. Lett.*, 89(28):281301, December 2002.
- [50] S. Habib, A. Heinen, K. Heitmann, and G. Jungman, Inflationary perturbations and precision cosmology, *Phys. Rev. D*, 71(4):043518, February 2005.
- [51] S. Habib, A. Heinen, K. Heitmann, G. Jungman, and C. Molina-París, Characterizing inflationary perturbations: The uniform approximation, *Phys. Rev. D*, 70(8):083507, October 2004.
- [52] A. Ashoorioon, R. Casadio, M. Cicoli, G. Geshnizjani, and H. J. Kim, Extended effective field theory of inflation, *J. High Energ. Phys.*, 2018(2):172, February 2018.
- [53] F. W. J. Olver, *Asymptotics and Special Functions*, AKP classics. A.K. Peters, Wellesley, Mass, 1997.
- [54] F. W. J. Olver, Second-Order Linear Differential Equations with Two Turning Points, *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 278(1279):137–174, March 1975.
- [55] M. Lemoine, M. Lubo, J. Martin, and J.-P. Uzan, Stress-energy tensor for trans-Planckian cosmology, *Phys. Rev. D*, 65(2):023510, December 2001.
- [56] R. H. Brandenberger and J. Martin, Back-reaction and the trans-Planckian problem of inflation reexamined, *Phys. Rev. D*, 71(2):023504, January 2005.