

Uniqueness for the inverse fixed angle scattering problem

J.A. Barceló* C. Castro* T. Luque† M. C. Vilela*

November 9, 2018

Abstract

We present a uniqueness result in dimensions 2 and 3 for the inverse fixed angle scattering problem associated to the Schrödinger operator $-\Delta + q$, where q is a small real valued potential with compact support in the Sobolev space $W^{\beta,2}$ with $\beta > 0$. This result improves the known result, due to Stefanov [6], in the sense that almost no regularity is required for the potential. The uniqueness result still holds in dimension 4, but for more regular potentials in $W^{\beta,2}$ with $\beta > 2/3$.

The proof is a consequence of the reconstruction method presented in our previous work [1].

1 Introduction

We consider the scattering problem appearing in quantum mechanics of finding u the solution to

$$\begin{cases} (-\Delta + q(x) - k^2)u(x) = 0, & x \in \mathbb{R}^d, \\ u = u_i + u_s, \\ \partial_r u_s - iku_s = o(r^{-(d-1)/2}), & r = |x| \longrightarrow \infty, \end{cases} \quad (1)$$

where $q \equiv q(x)$ is a real valued potential with compact support, $u_i \equiv u_i(x, \theta, k) = e^{ik\theta \cdot x}$ is the incident wave, with wave number $k > 0$ and direction of propagation $\theta \in \mathbb{S}^{d-1}$.

*Departamento de Matemática e Informática aplicadas a las Ingenierías Civil y Naval, Universidad Politécnica de Madrid. E-mails: juanantonio.barcelo@upm.es, carlos.castro@upm.es, maricruz.vilela@upm.es.

†Departamento de Análisis Matemático y Matemática Aplicada, Universidad Complutense de Madrid. E-mail: t.luque@ucm.es.

The solution u of problem (1) is known as the outgoing scattering solution, and u_s , which is the perturbation of the incident wave due to the potential, as the scattered wave.

It is well known that for appropriate potentials q , u_s satisfies

$$u_s(x, \theta, k) = C_d k^{(d-3)/2} \frac{e^{ik|x|}}{|x|^{(d-1)/2}} u_\infty(\omega, \theta, k) + o\left(|x|^{-(d-1)/2}\right), \quad |x| \rightarrow \infty, \quad (2)$$

where $\omega = x/|x|$ and

$$u_\infty(\omega, \theta, k) = \int_{\mathbb{R}^d} e^{-ik\omega \cdot y} q(y) u(y, \theta, k) dy. \quad (3)$$

The function $u_\infty \equiv u_\infty(\omega, \theta, k)$, known as the scattering amplitude or far-field pattern of u , depends on the wave number $k > 0$, the incident angle $\theta \in \mathbb{S}^{d-1}$ and the reflecting angle $\omega \in \mathbb{S}^{d-1}$. The central problem in inverse scattering for the Schrödinger equation is to recover the unknown potential q from the scattering amplitude u_∞ . This is an overdetermined problem, since u_∞ depends on $2d - 1$ variables while q only depends on d .

Therefore, it is natural to recover the potential from partial knowledge of the scattering amplitude. From the applications point of view, the most widely studied cases are the fixed angle scattering data, the fixed energy scattering data and the backscattering. An important question that arise is if this partial knowledge of the scattering amplitude determines the potential. Here we are interested in this problem when considering fixed angle scattering data, i.e. given $\theta_0 \in \mathbb{S}^{d-1}$, we assume known $u_\infty \equiv u_\infty(\omega, \pm\theta_0, k)$ for $\omega \in \mathbb{S}^{d-1}$ and $k > 0$. More precisely, we are interested in the following uniqueness question: Are there two different potentials q_1 and q_2 with the same fixed angle scattering data? Note that in this case both the potential and the scattering data depend on the same number of parameters d .

Mathematical analysis of reconstruction algorithms for fixed angle scattering data was first considered by Prosser [3], who obtained a potential in series form that fits the fixed angle scattering data in dimension three, under some restrictions on the size of the potentials in a Friedrichs type norm. However, the uniqueness result remained open. The first uniqueness result is due to Bayliss, Li and Morawetz [2], that proved uniqueness when the fixed angle scattering amplitude vanishes. This result was extended by Ramm [4] to scattering data coming from compactly supported potentials q that are constant along the lines defined by the direction $\theta = \theta_0$.

As far as we know, the most important result in this subject is due to Stefanov [6], who proves uniqueness for small complex valued potentials in $W^{4,\infty}$ with compact support in \mathbb{R}^3 . In the particular case of real potentials,

he shows that it is enough to consider scattering data for a unique incident angle θ_0 (see Remark 2.1).

Our uniqueness result, given in the following theorem, improves Stefanov's result since much less regularity is required on the potentials and also holds in dimensions two and four. However, we consider a larger set of scattering data since we use the scattering amplitude coming from both θ_0 and $-\theta_0$.

Theorem 1.1. *Let $\beta > 0$ if $d = 2$ or 3 and $\beta > 2/3$ if $d = 4$. For $i = 1, 2$ let q_i be a real valued potential in $W^{\beta,2}(\mathbb{R}^d)$ with compact support and scattering amplitude given by $u_\infty^i(\omega, \theta, k)$. Then, there exists a constant B small enough (see Remark 2.5), depending on β, d and the supports of q_1 and q_2 , such that if*

$$\|q_i\|_{W^{\beta,2}} < B, \quad i = 1, 2, \quad (4)$$

and

$$u_\infty^1(\omega, \theta_0, k) = u_\infty^2(\omega, \theta_0, k), \quad \text{for all } \omega \in \mathbb{S}^{d-1} \text{ and } k > 0, \quad (5)$$

then $q_1 = q_2$.

In general, $W^{\alpha,p}(\mathbb{R}^d)$ denotes the Sobolev space

$$W^{\alpha,p}(\mathbb{R}^d) = \{f \in \mathcal{S}'(\mathbb{R}^d) : \Lambda^\alpha f \in L^p(\mathbb{R}^d)\}, \quad \alpha \in \mathbb{R}, 1 \leq p \leq \infty,$$

where

$$\Lambda^\alpha = (1 - \Delta)^{\alpha/2} = \mathcal{F}^{-1} \langle \xi \rangle^\alpha \mathcal{F},$$

\mathcal{F} denotes the Fourier transform and $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$.

2 Proof of Theorem 1.1

The proof of Theorem 1.1 is a direct consequence of Theorem 1.1. of [1], where authors construct a sequence of approximations using the fixed angle scattering data and prove that it converges to the potential in certain Sobolev space and for certain potentials.

For completeness, we include here the statement of Theorem 1.1. of [1]. We first observe that the scattered wave u_s in (1) for $\theta = \theta_0$ satisfies the Lippmann-Schwinger integral equation given by

$$u_s(x, \theta_0, k) = R_k \left(q e^{ik\theta_0 \cdot (\cdot)} \right) (x) + R_k (q u_s(\cdot, \theta_0, k)) (x), \quad x \in \mathbb{R}^d, \quad (6)$$

where R_k denotes the outgoing resolvent of the Laplace operator, which in terms of the Fourier transform, is defined by

$$\widehat{R_k(f)}(\xi) = \frac{\widehat{f}(\xi)}{-|\xi|^2 + k^2 + i0}. \quad (7)$$

Under the conditions of Theorem 1.1, the existence and uniqueness of u_s is guaranteed by Theorem 2.11 of [1].

Moreover, since $u(x, \theta_0, k) = e^{ik\theta_0 \cdot x} + u_s(x, \theta_0, k)$, from (3) we obtain

$$u_\infty(\omega, \theta_0, k) = \widehat{q}(k(\omega - \theta_0)) + \int_{\mathbb{R}^d} e^{-ik\omega \cdot y} q(y) u_s(y, \theta_0, k) dy. \quad (8)$$

This provides a first approximation to the unknown potential q called the Born approximation. Formally, if we remove the last term in (8), the right hand side is the Fourier transform of q along the so-called Ewald spheres. More precisely, given θ_0 fixed, we have, up to a zero measure set,

$$\mathbb{R}^d = H_{\theta_0} \cup H_{-\theta_0} = \left\{ \xi \in \mathbb{R}^d : \xi \cdot \theta_0 < 0 \right\} \cup \left\{ \xi \in \mathbb{R}^d : \xi \cdot \theta_0 > 0 \right\}.$$

For $\xi \in H_{\pm\theta_0}$ there exist unique $\omega(\xi) \in \mathbb{S}^{d-1}$ and $k(\xi) > 0$ such that

$$\xi = k(\xi)(\omega(\xi) - \theta_0(\xi)), \quad \text{with } \theta_0(\xi) = \begin{cases} \theta_0, & \xi \in H_{\theta_0} \\ -\theta_0, & \xi \in H_{-\theta_0} \end{cases}.$$

Actually, if $\xi \in H_{\theta_0}$ we have that

$$\omega(\xi) = -\frac{2\xi \cdot \theta_0}{|\xi|^2} \xi + \theta_0, \quad \text{and } k(\xi) = -\frac{|\xi|^2}{2\xi \cdot \theta_0}. \quad (9)$$

The Born approximation for fixed angle scattering data θ_0 of a potential q is defined by

$$\widehat{q_{\theta_0}}(\xi) = u_\infty(\omega(\xi), \theta_0(\xi), k(\xi)). \quad (10)$$

Remark 2.1. Definition (10) is equivalent to the one used by Stefanov in [6] when $\xi \in H_{\theta_0}$, but it differs when $\xi \in H_{-\theta_0}$. More precisely, instead of considering scattering data for $-\theta_0$ and $k > 0$, the scattering amplitude in [6] is extended to $k < 0$ by $u_\infty(w, \theta_0, k) = \overline{u_\infty(w, \theta_0, -k)}$. In this way, only scattering data coming from θ_0 are required.

Now we insert iteratively the Lippmann-Schwinger integral equation (6) into (8) to obtain for $\xi = k(\xi)(\omega(\xi) - \theta_0(\xi))$ and $m \geq 1$

$$\widehat{q_{\theta_0}}(\xi) = \widehat{q}(\xi) + \sum_{j=1}^m \widehat{\mathcal{Q}_j(q)}(\xi) + \widehat{q_m^*}(\xi), \quad (11)$$

where

$$\widehat{\mathcal{Q}_j(q)}(\xi) = \sum_{j=1}^m \int_{\mathbb{R}^d} e^{-ik(\xi)\omega(\xi) \cdot y} (qR_{k(\xi)})^j \left(qe^{ik(\xi)\theta_0(\xi) \cdot (\cdot)} \right) (y) dy, \quad (12)$$

and

$$\widehat{q}_m^r(\xi) = \int_{\mathbb{R}^d} e^{-ik(\xi)\omega(\xi)\cdot y} (qR_{k(\xi)})^j (qu_s(\cdot, \theta_0(\xi), k(\xi)))(y) dy.$$

In order to construct our sequence of approximations of q , we fix m and consider the operator

$$\mathcal{T}_m(f) = \phi q_{\theta_0} - \phi \sum_{j=1}^m \mathcal{Q}_j(f)(\xi), \quad (13)$$

where $\phi \in \mathcal{C}^\infty$ is a cut-off function with compact support satisfying

$$\phi(x) = 1, \quad \text{if } |x| < 1 \quad \text{and} \quad \phi(x) = 0, \quad \text{if } |x| > 2R,$$

with $R > 0$ such that the support of q is contained in $B(0, R)$.

We define the sequence $\{q_{m,\ell}\}_{\ell \in \mathbb{N}}$ recursively by

$$\begin{cases} q_{m,1} = 0, \\ q_{m,\ell+1} = \mathcal{T}_m(q_{m,\ell}), \quad \ell \geq 1. \end{cases} \quad (14)$$

Remark 2.2. Observe that $q_{m,2}$ is nothing but ϕq_{θ_0} . On the other hand, from (13), the sequence $\{q_{m,\ell}\}_{\ell \in \mathbb{N}}$ only depends on $u_\infty(\omega, \theta_0, k)$ and R .

The following theorem states that the iterated limit, first in ℓ and then in m , of the sequence $\{q_{m,\ell}\}_{m,\ell \in \mathbb{N}}$ is equal to q , under certain hypotheses.

Theorem 2.3. (Theorem 1.1 of [1]) For $d \geq 2$ and α satisfying

$$0 < \alpha \leq 1, \quad \text{and} \quad \frac{d}{2} - \frac{d}{d-1} < \alpha < \frac{d}{2}, \quad (15)$$

let $q \in W^{\alpha,2}(\mathbb{R}^d)$ be a real function with compact support in $B(0, R)$ and such that

$$\|q\|_{W^{\alpha,2}} < A, \quad (16)$$

for an appropriate constant $A := A(d, \alpha, R) > 0$ small enough. For each $m \in \mathbb{N}$, let $\{q_{m,\ell}\}_{\ell \in \mathbb{N}}$ be the sequence defined by (14). Then, there exists $q_m \in W^{\alpha,2}(\mathbb{R}^d)$ satisfying

$$q_m = \lim_{\ell \rightarrow \infty} q_{m,\ell} \quad \text{in } W^{\alpha,2}(\mathbb{R}^d). \quad (17)$$

Moreover, the sequence $\{q_m\}_{m \in \mathbb{N}}$ satisfies

$$\lim_{m \rightarrow \infty} q_m = q \quad \text{in } W^{\alpha,2}(\mathbb{R}^d). \quad (18)$$

Remark 2.4. The proof of Theorem 2.3 depends on very accurate estimates for R_k , the resolvent of the Laplace operator, which can be found in [5]. The smallness condition (16) is written in terms of the constants appearing in these estimates (see Remark 1.2 of [1]). On the other hand, conditions in (15) imply that $d < 5$ and, therefore, Theorem 2.3 is only valid for $2 \leq d \leq 4$.

Proof of Theorem 1.1. The proof is a straightforward consequence of Theorem 2.3. Precisely, since any q under the conditions of Theorem 1.1 satisfies the assumptions of Theorem 2.3, for $i = 1, 2$ we can construct the approximation sequence $\left\{q_{m,\ell}^i\right\}_{m,\ell \in \mathbb{N}}$ for q_i , which only depends on $u_\infty^i(x, \pm\theta_0, k)$ and the support of q_i . Therefore, by (5), $q_{m,\ell}^1 = q_{m,\ell}^2$. From (17)-(18) we easily obtain $q_1 = q_2$. \square

Remark 2.5. Notice that in the particular cases $\beta < 1$ and $d = 2$ or $\beta \leq 1$ and $d = 3$ or $d = 4$, the constant B given in (4) equals the constant A given in (16), with $\alpha = \beta$. In other case, we can chose any α under the conditions of Theorem 1.1 such that $0 < \alpha < \beta$, and thus

$$\|q_i\|_{W^{\alpha,2}} \leq C(d, \alpha, \beta, R)\|q_i\|_{W^{\beta,2}} < A, \quad i = 1, 2,$$

if we take

$$B = \frac{A}{C(d, \alpha, \beta, R)}.$$

References

- [1] J. A. Barceló, C. Castro, T. Luque, and M. C. Vilela, A new convergent algorithm to approximate potentials from fixed angle scattering data, to appear in: *SIAM J. Appl. Math.* (<https://doi.org/10.1137/18M1172247>), available at [1807.04820](https://arxiv.org/abs/1807.04820).
- [2] A. Bayliss, Y. Li, and C. Morawetz, Scattering by potential using hyperbolic methods, in *Math. Comp.* **52** (1989), 321–328.
- [3] R. T. Prosser, Formal solutions of inverse scattering problems. V, *J. Math. Phys.* **33**, 3493 (1998).
- [4] A. G. Ramm, Multidimensional inverse scattering problems and completeness of the products of solutions to homogeneous PDE, *Z. Angew. Math. Mech.* **69** (1989), 4, pp. T13–T22.
- [5] A. Ruiz, Recovery of the singularities of a potential from fixed angle scattering data, *Commun. Partial Diffe. Equations* **26** (2001), pp. 1721–1738.
- [6] P. Stefanov, Generic uniqueness for two inverse problems in potential scattering, *Commun. Partial Diffe. Equations* **17** (1992), pp. 55–68.