

# Identities And Relations On The Hermite-based Tangent Polynomials

Burak Kurt

Department of Mathematics, Faculty of Educations  
University of Akdeniz  
TR-07058 Antalya, Turkey  
E-mail: [burakkurt@akdeniz.edu.tr](mailto:burakkurt@akdeniz.edu.tr)

## Abstract

In this note, we introduce and investigate the Hermite-based Tangent numbers and polynomials, Hermite-based modified degenerate-Tangent polynomials, poly-Tangent polynomials. We give some identities and relations for these polynomials.

**2010 Mathematics Subject Classification.** 11B75, 11B68, 11B83, 33E30, 33F99.

**Key Words and Phrases.** Bernoulli polynomials and numbers, Genocchi polynomials and numbers, Hermite polynomials, Stirling numbers of the second kind, Tangent polynomials and numbers, Hermite-based Bernoulli polynomials, Hermite-based Tangent polynomials, polylogarithm function, Poly-Bernoulli polynomials, Degenerate Bernoulli and Genocchi polynomials.

## 1. INTRODUCTION AND NOTATION

Many mathematicians have studied in the area of the Bernoulli numbers and polynomials, Euler number and polynomials, Genocchi numbers and polynomials, poly-Bernoulli numbers and polynomials, poly-Euler numbers and polynomials, poly-Genocchi numbers and polynomials, poly-Tangent numbers and polynomials, Hermite polynomials, Hermite-based Bernoulli polynomials, Hermite-based Tangent polynomials, modified degenerate Bernoulli polynomials, modified degenerate Euler polynomials and modified degenerate Genocchi polynomials (see [1]-[20]). In this note we define the Hermite-based tangent polynomials, modified Hermite-based tangent polynomials and poly-tangent polynomials. We obtain some relations and identities for these polynomials. Throughout this paper, we always make use of the following notations:  $\mathbb{N}$  denotes the set of natural numbers and  $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ . We recall that

the classical Stirling numbers of the first kind  $S_1(n, k)$  and second kind  $S_2(n, k)$  are defined by the relations [15]

$$(x)_n = \sum_{k=0}^n S_1(n, k)x^k \quad \text{and} \quad x^n = \sum_{k=0}^n S_2(n, k)(x)_k \quad (1.1)$$

respectively. Here  $(x)_n = x(x-1)\cdots(x-n+1)$  denotes the falling factorial polynomials of order  $n$ . The numbers  $S_2(n, m)$  also admit a representation in terms of a generating function

$$\frac{(e^t - 1)^m}{m!} = \sum_{n=m}^{\infty} S_2(n, m) \frac{t^n}{n!}. \quad (1.2)$$

The Bernoulli polynomials  $B_n^{(r)}(x)$  of order  $\alpha$ , the Euler polynomials  $E_n^{(r)}(x; \lambda)$  of order  $\alpha$  and the Genocchi polynomials  $G_n^{(r)}(x; \lambda)$  of order  $\alpha$  are defined as respectively:

$$\left(\frac{t}{e^t - 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} B_n^{(r)}(x) \frac{t^n}{n!}, \quad |t| < 2\pi, \quad (1.3)$$

$$\left(\frac{2}{e^t + 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}, \quad |t| < \pi \quad (1.4)$$

and

$$\left(\frac{2t}{e^t + 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} G_n^{(r)}(x) \frac{t^n}{n!}, \quad |t| < \pi. \quad (1.5)$$

When  $x = 0$ ,  $B_n^{(r)}(0) = B_n^{(r)}$ ,  $E_n^{(r)}(0) = E_n^{(r)}$  and  $G_n^{(r)}(0) = G_n^{(r)}$  are called Bernoulli numbers of order  $r$ , Euler numbers of order  $r$  and Genocchi numbers of order  $r$ , respectively.

The familiar tangent polynomials  $T_n^{(r)}(x)$  of order  $r$  are defined by the generating functions ([12]-[15], [17])

$$\left(\frac{2}{e^{2t} + 1}\right)^r e^{xt} = \sum_{n=0}^{\infty} T_n^{(r)}(x) \frac{t^n}{n!}, \quad |2t| < \pi. \quad (1.6)$$

When  $x = 0$ ,  $T_n^{(r)}(0) = T_n^{(r)}$  are called the tangent numbers.

2-variable Hermite-Kampéde Fériet polynomials are defined in ([5], [11]) as

$$\sum_{n=0}^{\infty} H_n(x, y) \frac{t^n}{n!} = e^{xt+yt^2}. \quad (1.7)$$

Khan *et al.* in [5] defined and studied on Hermite-based Bernoulli polynomials and Hermite-based Euler polynomials as

$$\sum_{n=0}^{\infty} ({}_H\mathcal{B}_n(x, y)) \frac{t^n}{n!} = \frac{t}{e^t - 1} e^{xt+yt^2}, \quad |t| < 2\pi \quad (1.8)$$

and

$$\sum_{n=0}^{\infty} ({}_H\mathcal{E}_n(x, y)) \frac{t^n}{n!} = \frac{2}{e^t + 1} e^{xt+yt^2}, \quad |t| < \pi, \quad (1.9)$$

respectively.

Carlitz in [1] defined degenerate Bernoulli polynomials which are given by the generating functions to be

$$\frac{t}{(1 + \lambda t)^{1/\lambda} - 1} (1 + \lambda t)^{x/\lambda} = \sum_{n=0}^{\infty} \mathfrak{B}_n(x | \lambda) \frac{t^n}{n!}. \quad (1.10)$$

When  $x = 0$ ,  $\mathfrak{B}_n(\lambda) = \mathfrak{B}_n(0 | \lambda)$  are called the degenerate Bernoulli numbers. From (1.10), we can easily derive the following equation

$$\mathfrak{B}_n(x | \lambda) = \sum_{l=0}^n \binom{n}{l} \mathfrak{B}_{n-l}(\lambda) (x | \lambda)_l, \quad n \geq 0$$

where  $(x | \lambda)_n = x(x - \lambda) \cdots (x - \lambda(n - 1))$ ,  $(x | \lambda)_0 = 1$ .

Dolgy *et. al.* [2] defined the modified degenerate Bernoulli polynomials, which are different from Carlitz's degenerate Bernoulli polynomials as

$$\frac{t}{(1 + \lambda)^{t/\lambda} - 1} (1 + \lambda)^{xt/\lambda} = \sum_{n=0}^{\infty} \mathfrak{B}_{n,\lambda}(x) \frac{t^n}{n!}. \quad (1.11)$$

When  $x = 0$ ,  $\mathfrak{B}_{n,\lambda} = \mathfrak{B}_{n,\lambda}(0)$  are called the modified degenerate Bernoulli numbers. From (1.11), we note that

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \sum_{n=0}^{\infty} \mathfrak{B}_{n,\lambda}(x) \frac{t^n}{n!} &= \lim_{\lambda \rightarrow 0} \frac{t}{(1 + \lambda)^{t/\lambda} - 1} (1 + \lambda)^{xt/\lambda} \\ &= \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}. \end{aligned} \quad (1.12)$$

Thus, by (1.12)

$$\lim_{\lambda \rightarrow 0} \mathfrak{B}_{n,\lambda}(x) = B_n(x).$$

H.-In Known *et. al.* [8] defined the modified degenerate Euler polynomials as

$$\frac{2}{(1 + \lambda)^{t/\lambda} + 1} (1 + \lambda)^{xt/\lambda} = \sum_{n=0}^{\infty} \mathfrak{E}_{n,\lambda}(x) \frac{t^n}{n!} \quad (1.13)$$

and T. Kim *et. al.* in [6] defined the modified degenerate Genocchi polynomials as

$$\frac{2t}{(1 + \lambda)^{t/\lambda} + 1} (1 + \lambda)^{tx/\lambda} = \sum_{n=0}^{\infty} \mathfrak{G}_{n,\lambda}(x) \frac{t^n}{n!}. \quad (1.14)$$

From (1.13) and (1.14), we get

$$\lim_{\lambda \rightarrow 0} \mathfrak{E}_{n,\lambda}(x) = E_n(x), \quad \lim_{\lambda \rightarrow 0} \mathfrak{G}_{n,\lambda}(x) = G_n(x).$$

For  $k \in \mathbb{Z}$ ,  $k > 1$ , then  $k$ -th polylogarithm is defined by Kaneko [4] as

$$L_{i_k}(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^k}. \quad (1.15)$$

Thus function is convergent for  $|z| < 1$ , when  $k = 1$

$$L_{i_1}(z) = -\log(1 - z). \quad (1.16)$$

Kim *et. al.* in [7] defined the poly-Bernoulli polynomials and the poly-Genocchi polynomials as

$$\sum_{n=0}^{\infty} \mathfrak{B}_n^{(k)}(x) \frac{t^n}{n!} = \frac{L_{i_k}(1 - e^{-t})}{1 - e^{-t}} e^{xt} \quad (1.17)$$

and

$$\sum_{n=0}^{\infty} \mathfrak{G}_n^{(k)}(x) \frac{t^n}{n!} = \frac{2L_{i_k}(1 - e^{-t})}{e^t + 1} e^{xt} \quad (1.18)$$

respectively.

For  $k = 1$ , by use (1.16) in (1.17) and (1.18). We get

$$\mathfrak{B}_n^{(1)}(x) = (-1)^{n+1} B_n(x), \quad \mathfrak{G}_n^{(1)}(x) = G_n(x).$$

Hamahata [3] defined poly-Euler polynomials by

$$\sum_{n=0}^{\infty} \mathfrak{E}_n^{(k)}(x) \frac{t^n}{n!} = \frac{2L_{i_k}(1 - e^{-t})}{t(e^t + 1)} e^{xt}.$$

For  $k = 1$ , we get  $\mathfrak{E}_n^{(1)}(x) = E_n(x)$ .

From (1.6), we obtain the following equalities easily

$$T_n^{(r)}(x) = \sum_{k=0}^n \binom{n}{k} T_k^{(r)} x^{n-k},$$

$$T_n^{(r)}(x + y) = \sum_{l=0}^k \binom{k}{l} T_k^{(r)}(x) y^{k-l},$$

$$T_n^{(r_1+r_2)}(x + y) = \sum_{k=0}^n \binom{n}{k} T_k^{(r_1)}(x) T_{n-k}^{(r_2)}(y)$$

and

$$T_n^{(r)}(2(x + 1)) = 2T_n^{(r-1)}(2x).$$

## 2. HERMITE BASED TANGENT POLYNOMIALS

Khan *et. al.* in [5] and Ozarslan [11] introduced and investigated the Hermite-based Bernoulli polynomials and Hermite-based Euler polynomials. They proved some identities and relations for these polynomials.

By this motivation, we define Hermite-based Tangent polynomials of order  $r$  as

$$\sum_{n=0}^{\infty} ({}_H T_n^{(r)}(x, y)) \frac{t^n}{n!} = \left( \frac{2}{e^{2t} + 1} \right)^r e^{xt+yt^2}. \quad (2.1)$$

**Theorem 1.** Let  $r_1, r_2 \in \mathbb{Z}_+$ . We have

$$\begin{aligned} {}_H T_n^{(r)}(x, y) &= \sum_{k=0}^n \binom{n}{k} T_k^{(r)}(0, 0) H_{n-k}(x, y), \\ {}_H T_n^{(r)}(x+u, y+v) &= \sum_{k=0}^n \binom{n}{k} \left( {}_H T_k^{(r)}(x, y) \right) H_{n-k}(u, v) \end{aligned}$$

and

$${}_H T_n^{(r_1+r_2)}(x+u, y+v) = \sum_{k=0}^n \binom{n}{k} \left( {}_H T_k^{(r_1)}(x, y) \right) \left( {}_H T_{n-k}^{(r_2)}(u, v) \right).$$

**Theorem 2.** Let  $r \in \mathbb{Z}_+$ .

$${}_H T_n^{(r)}(2(x+u), 2(y+v)) = \sum_{m=0}^n \binom{n}{m} \left( {}_H T_{n-m}^{(r)}(x, y) \right) \sum_{p=0}^m \binom{m}{p} H_p(x, y) H_{m-p}(x, y).$$

**Theorem 3.** There is the following implicit relation for the Hermite-based Tangent polynomials as

$${}_H T_{n+m}^{(r)}(u, v) = \sum_{p=0}^n \binom{n}{p} \sum_{q=0}^m \binom{m}{q} (v-y)^{p+q} \left( {}_H T_{n+m-p-q}^{(r)}(x, y) \right). \quad (2.2)$$

*Proof.* From (2.1), we replace  $t$  by  $t+u$  and rewrite the generating function as

$$\frac{2e^{y(t+u)^2}}{e^{2t} + 1} = e^{-x(t+u)} \sum_{n=0}^{\infty} T_{n+m}^{(r)}(x, y) \frac{t^n u^m}{n! m!}.$$

Replacing  $x$  by  $v$  in the above equation to the above equation.

We get

$$\sum_{n,m=0}^{\infty} \left( {}_H T_{n+m}^{(r)}(v, y) \right) \frac{t^n u^m}{n! m!} = e^{(t+u)(v-x)} \sum_{n,m=0}^{\infty} \left( {}_H T_{n+m}^{(r)}(x, y) \right) \frac{t^n u^m}{n! m!}$$

which on using formula [19, Srivastava p. 52]

$$\sum_{N=0}^{\infty} f(N) \frac{(x+y)^N}{N!} = \sum_{n,m=0}^{\infty} f(n+m) \frac{x^n y^m}{n! m!}$$

in the right hand side becomes

$$\sum_{p=0}^{\infty} \sum_{q=0}^{\infty} (v-x)^{p+q} \frac{t^p u^q}{p! q!} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \left( {}_H T_{n+m}^{(r)}(x, y) \right) \frac{t^n u^m}{n! m!} = \sum_{n,m=0}^{\infty} \left( {}_H T_{n+m}^{(r)}(v, y) \right) \frac{t^n u^m}{n! m!}.$$

By using Cauchy product and comparing the coefficients of both sides, we have (2.2).  $\square$

**Theorem 4.** There is the following relation between the Hermite-based Tangent polynomials and the Hermite-based Bernoulli polynomials as

$$\left( {}_H \mathfrak{B}_n^{(r)} \left( \frac{x+u}{4}, \frac{y+v}{16} \right) \right) = 2^{r-n-k} \sum_{k=0}^n \binom{n}{k} \left( {}_H T_k^{(r)}(x, y) \right) \left( {}_H \mathfrak{B}_{n-k}^{(r)} \left( \frac{u}{2}, \frac{v}{4} \right) \right). \quad (2.3)$$

*Proof.* From (2.1),

$$\begin{aligned} \sum_{n=0}^{\infty} \left( {}_H\mathfrak{B}_n^{(r)} \left( \frac{x+u}{4}, \frac{y+v}{16} \right) \right) \frac{(4t)^n}{n!} &= \left( \frac{2 \times 4t}{e^{4t} - 1} \right)^{(r)} e^{(x+u)t+(y+v)t^2} \\ &= \left( \frac{2}{e^{2t} + 1} \right)^{(r)} e^{xt+yt^2} 2^r \left( \frac{2t}{e^{2t} - 1} \right)^{(r)} e^{ut+vt^2} \\ &= \sum_{n=0}^{\infty} ({}_HT_n^{(r)}(x, y)) \frac{t^n}{n!} 2^r \sum_{q=0}^{\infty} \left( {}_H\mathfrak{B}_q^{(r)} \left( \frac{u}{2}, \frac{v}{4} \right) \right) \frac{(2t)^q}{q!}. \end{aligned}$$

By using Cauchy product and comparing the coefficients of both sides. We get (2.3).  $\square$

### 3. MODIFIED DEGENERATE HERMITE-BASED TANGENT POLYNOMIALS

Dolgy *et. al.* [2] introduced and investigated the modified degenerate Bernoulli polynomials. Known *et. al.* [8] defined and investigated the modified degenerate Euler polynomials. They proved some properties for these polynomials.

By these motivations, we define 2-variable modified degenerate Hermite polynomials and the modified degenerate Hermite-based Tangent polynomials of order  $r$

$$\sum_{n=0}^{\infty} H_n(x, y : \lambda) \frac{t^n}{n!} = (1 + \lambda)^{\frac{xt+yt^2}{\lambda}} \quad (3.1)$$

and

$$\sum_{n=0}^{\infty} ({}_HT_n^{(r)}(x, y : \lambda)) \frac{t^n}{n!} = \frac{2}{(1 + \lambda)^{\frac{2t}{\lambda}} + 1} (1 + \lambda)^{\frac{xt+yt^2}{\lambda}} \quad (3.2)$$

respectively.

From (3.1) and (3.2), we get

$$\lim_{\lambda \rightarrow 0} H_n(x, y : \lambda) = H_n(x, y), \quad \lim_{\lambda \rightarrow 0} ({}_HT_n^{(r)}(x, y : \lambda)) = ({}_HT_n^{(r)}(x, y)).$$

Similiary, we define the modified Hermite-based Bernoulli polynomials and the modified Hermite-based Euler polynomials as

$$\sum_{n=0}^{\infty} ({}_H\mathfrak{B}_n(x, y : \lambda)) \frac{t^n}{n!} = \frac{t}{(1 + \lambda)^{\frac{t}{\lambda}} - 1} (1 + \lambda)^{\frac{xt+yt^2}{\lambda}} \quad (3.3)$$

and

$$\sum_{n=0}^{\infty} ({}_H\mathfrak{E}_n(x, y : \lambda)) \frac{t^n}{n!} = \frac{2}{(1 + \lambda)^{\frac{t}{\lambda}} + 1} (1 + \lambda)^{\frac{xt+yt^2}{\lambda}} \quad (3.4)$$

respectively.

From (3.2), we obtain the following relations easily

$$({}_HT_n^{(r_1+r_2)}(x+u, y+v : \lambda)) = \sum_{k=0}^n \binom{n}{k} ({}_HT_k^{(r_1)}(x, y : \lambda)) ({}_HT_{n-k}^{(r_2)}(u, v : \lambda)),$$

$$\begin{aligned}({}_H T_n^{(r)}(x, y : \lambda)) &= \sum_{k=0}^n \binom{n}{k} \left( {}_H T_k^{(r)}(0, 0 : \lambda) \right) (H_{n-k}(x, y : \lambda)), \\({}_H T_n^{(r)}(x+2, y : \lambda) + {}_H T_n^{(r)}(x, y : \lambda)) &= 2 ({}_H T_n^{(r-1)}(x, y : \lambda))\end{aligned}$$

for  $r = 1$ ,

$$({}_H T_n(x+2, y : \lambda) + {}_H T_n(x, y : \lambda)) = 2 (H_n(x, y : \lambda))$$

and

$$({}_H T_n^{(r)}(x, y : \lambda)) = \sum_{k=0}^n \binom{n}{k} \left( {}_H T_n^{(r)}\left(\frac{1}{2}, 0 : \lambda\right) \right) \left( H_{n-k}\left(x - \frac{1}{2}, y : \lambda\right) \right).$$

**Theorem 5.** *There is the following relation between the modified degenerate Bernoulli polynomials, the modified degenerate Euler polynomials and the modified degenerate Tangent polynomials as*

$$({}_H \mathfrak{B}_n(x, y : \lambda)) 2^{2n+1} = \sum_{q=0}^n \binom{n}{q} ({}_H T_{n-q}(x, y : \lambda)) \sum_{k=0}^q \binom{q}{k} ({}_H \mathfrak{B}_{q-k}(x, y : \lambda)) ({}_H \mathfrak{E}_n(2x, 14y : \lambda)). \quad (3.5)$$

*Proof.* From (3.3), (3.4) and (3.2), we write as

$$\begin{aligned}\sum_{n=0}^{\infty} ({}_H \mathfrak{B}_n(x, y : \lambda)) \frac{(4t)^n}{n!} &= \left( \frac{4t}{(1+\lambda)^{\frac{4t}{\lambda}} - 1} \right) (1+\lambda)^{\frac{4tx+y(4t)^2}{\lambda}} \\&= \frac{1}{2} \frac{2e^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{2t}{\lambda}} + 1} \frac{2te^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{t}{\lambda}} - 1} \frac{2e^{\frac{2xt+14yt^2}{\lambda}}}{(1+\lambda)^{\frac{t}{\lambda}} + 1} \\&= \frac{1}{2} \sum_{n=0}^{\infty} ({}_H T_n(x, y : \lambda)) \frac{t^n}{n!} \sum_{p=0}^{\infty} ({}_H \mathfrak{B}_p(x, y : \lambda)) \frac{t^p}{p!} \sum_{q=0}^{\infty} ({}_H \mathfrak{E}_q(2x, 14y : \lambda)) \frac{t^q}{q!}.\end{aligned}$$

By using Cauchy product and comparing the coefficient of  $\frac{t^n}{n!}$ , we have (3.5).  $\square$

**Theorem 6.**  $n \in \mathbb{Z}_+$ , we have

$$({}_H T_n(x+2, y : \lambda) + {}_H T_n(x, y : \lambda)) = \frac{1}{n+1} \{ ({}_H \mathfrak{B}_{n+1}(x+1, y : \lambda)) - ({}_H \mathfrak{B}_{n+1}(x, y : \lambda)) \}. \quad (3.6)$$

*Proof.* By using definition (3.2)

$$\begin{aligned}\frac{2t(1+\lambda)^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{2t}{\lambda}} + 1} \left[ (1+\lambda)^{\frac{2t}{\lambda}} + 1 \right] &= \frac{2t(1+\lambda)^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{t}{\lambda}} - 1} \left[ (1+\lambda)^{\frac{t}{\lambda}} - 1 \right] \\ \frac{2t(1+\lambda)^{\frac{(x+2)t+yt^2}{\lambda}}}{(1+\lambda)^{\frac{2t}{\lambda}} + 1} + \frac{2t(1+\lambda)^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{2t}{\lambda}} + 1} &= \frac{2t(1+\lambda)^{\frac{(x+1)t+yt^2}{\lambda}}}{(1+\lambda)^{\frac{t}{\lambda}} - 1} - \frac{2t(1+\lambda)^{\frac{xt+yt^2}{\lambda}}}{(1+\lambda)^{\frac{t}{\lambda}} - 1}\end{aligned}$$

$$t \sum_{n=0}^{\infty} \{({}_H T_n(x+2, y: \lambda)) + ({}_H T_n(x, y: \lambda))\} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \{({}_H \mathfrak{B}_n(x+1, y: \lambda)) - ({}_H \mathfrak{B}_n(x, y: \lambda))\} \frac{t^n}{n!}.$$

From the above equality, we have (3.6).  $\square$

#### 4. POLY-TANGENT POLYNOMIALS

In this section, we define the poly-tangent numbers and polynomials and provide some of their relevant properties.

**Definition 1.** We define the Hermite-based poly-tangent polynomials by

$$\frac{2L_{i_k}(1 - e^{-t})}{t(e^{2t} + 1)} e^{xt+yt^2} = \sum_{n=0}^{\infty} ({}_H \mathcal{T}_n^{(k)}(x, y)) \frac{t^n}{n!}, \quad (4.1)$$

when  $x = 0$ ,  $({}_H \mathcal{T}_n^{(k)}) := ({}_H \mathcal{T}_n^{(k)}(0, 0))$  are called the Hermite-based poly-tangent numbers.

For  $k = 1$  and  $L_{i_k}(z) = -\log(1 - z)$ , from (4.1)

$$\frac{2L_{i_1}(1 - e^{-t})}{t(e^{2t} + 1)} e^{xt+yt^2} = \frac{2e^{xt+yt^2}}{e^{2t} + 1} = \sum_{n=0}^{\infty} ({}_H \mathcal{T}_n(x, y)) \frac{t^n}{n!}. \quad (4.2)$$

By (4.2), we get

$$({}_H \mathcal{T}_n^{(1)}(x, y)) = ({}_H T_n(x, y)).$$

**Theorem 7.**  $n, k \in \mathbb{Z}_+$ , we have

$$({}_H \mathcal{T}_n^{(k)}(x, y)) = \frac{1}{n+1} \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} ({}_H \mathcal{T}_{n+1}(x-j, y)). \quad (4.3)$$

*Proof.*

$$\begin{aligned} \sum_{n=0}^{\infty} ({}_H \mathcal{T}_n^{(k)}(x, y)) \frac{t^n}{n!} &= 2 \sum_{m=0}^{\infty} \frac{(1 - e^{-t})^{m+1}}{(m+1)^k} \frac{e^{xt+yt^2}}{t(e^{2t} + 1)} \\ &= 2 \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} \frac{e^{-tj+xt+yt^2}}{t(e^{2t} + 1)} \\ &= \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} \frac{1}{t} \frac{2}{e^{2t} + 1} e^{t(x-j)+yt^2} \\ &= \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} \sum_{n=0}^{\infty} ({}_H \mathcal{T}_n(x-j, y)) \frac{t^{n-1}}{n!} \\ &= \sum_{m=0}^{\infty} \frac{1}{(m+1)^k} \sum_{j=0}^{m+1} (-1)^j \binom{m+1}{j} \sum_{n=-1}^{\infty} \frac{({}_H \mathcal{T}_{n+1}(x-j, y))}{n+1} \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients both sides, we have (4.3).  $\square$



**Theorem 8.** *There is the following relation between poly-tangent polynomials and the Stirling numbers of the second kind and the Hermite-based Bernoulli polynomials as*

$$({}_H\mathcal{T}_n^{(k)}(x, y)) = \sum_{l=0}^n \frac{\binom{n}{l}}{\binom{l+r}{r}} S_2(l+r, r) \sum_{i=0}^{n-l} \left( {}_H\mathfrak{B}_i^{(r)}(x, y) \right) \left( {}_H\mathcal{T}_{n-l-i}^{(r)} \right). \quad (4.4)$$

*Proof.* From (4.1), we write as

$$\begin{aligned} \sum_{n=0}^{\infty} ({}_H\mathcal{T}_n^{(k)}(x, y)) \frac{t^n}{n!} &= \frac{2L_{i_k}(1-e^{-t})}{t(e^{2t}+1)} e^{xt+yt^2} \\ &= \frac{(e^t-1)^r r!}{r! t^r} \left( \frac{t}{e^t-1} \right)^r e^{xt+yt^2} \frac{2L_{i_k}(1-e^{-t})}{t(e^{2t}+1)} \\ &= \frac{(e^t-1)^r}{r!} \left( \sum_{n=0}^{\infty} ({}_H\mathfrak{B}_n^{(r)}(x, y)) \frac{t^n}{n!} \right) \left( \sum_{q=0}^{\infty} ({}_H\mathcal{T}_q^{(r)}) \frac{t^q}{q!} \right) \frac{r!}{t^r} \\ &= \sum_{n=0}^{\infty} \left( \sum_{l=0}^n \frac{\binom{n}{l}}{\binom{l+r}{r}} S_2(l+r, r) \sum_{i=0}^{n-l} \left( {}_H\mathfrak{B}_i^{(r)}(x, y) \right) \left( {}_H\mathcal{T}_{n-l-i}^{(r)} \right) \right) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients of  $\frac{t^n}{n!}$ , we obtain (4.4).  $\square$

**Theorem 9.** *There is the following relation between the poly-tangent polynomials, the poly-Genocchi numbers and the Hermite-based tangent polynomials*

$$({}_H\mathcal{T}_n^{(k)}(x, y)) = \frac{1}{2} \sum_{p=0}^n \binom{n}{p} G_{n-p}^{(k)} \{ ({}_H\mathcal{T}_n(x+1, y)) + ({}_H\mathcal{T}_n(x, y)) \}. \quad (4.5)$$

*Proof.* From (4.1) and (1.18)

$$\begin{aligned} \sum_{n=0}^{\infty} ({}_H\mathcal{T}_n^{(k)}(x, y)) \frac{t^n}{n!} &= \frac{2L_{i_k}(1-e^{-t})}{t(e^{2t}+1)} e^{xt+yt^2} \\ &= \frac{1}{2} \left( \frac{2L_{i_k}(1-e^{-t})}{e^t+1} \right) \frac{2(e^t+1)e^{xt+yt^2}}{t(e^{2t}+1)} \\ &= \frac{1}{2} \frac{2L_{i_k}(1-e^{-t})}{e^t+1} \left( \frac{2e^{(x+1)t+yt^2}}{t(e^{2t}+1)} + \frac{2e^{xt+yt^2}}{t(e^{2t}+1)} \right) \\ &= \frac{1}{2} \sum_{n=0}^{\infty} G_n^{(k)} \frac{t^n}{n!} \left\{ \sum_{p=0}^{\infty} ({}_H\mathcal{T}_p(x+1, y)) + ({}_H\mathcal{T}_p(x, y)) \frac{t^p}{p!} \right\}. \end{aligned}$$

Comparing the coefficients of both sides, we have (4.5).  $\square$

**Acknowledgement 1.** *The present investigation was supported, by the Scientific Research Project Administration of Akdeniz University.*

## REFERENCES

- [1] Carlitz L., Degenerate Stirling Bernoulli and Eulerian numbers, *Util. Math.*, 15 (1979), 51-88.
- [2] Dolgy D. V., Kim T., Known H.-In and Seo J. J., On the modified degenerate Bernoulli polynomials, *Advanced Studies in Contemp. Math.*, 26 (2016), 203-209.
- [3] Hamahata Y., Poly-Euler polynomials and Arakawa-Kaneko type zeta functions, *Functiones et Approximation Commentari Math.*, 51(1), (2014), 7-22.
- [4] Kaneko M., Poly-Bernoulli numbers, *J. Théor. Nr. Bordx*, 9, (1999), 199-206.
- [5] Khan S., Yasmin G., Khan R., and Hassan N. A., Hermite-based Appell polynomials: Properties and applications, *J. of Math. Anal. and Appl.*, 351, (2009), 756-764.
- [6] Kim T., Kim D. S., and Known H.-In, Some identities relating to degenerate Bernoulli polynomials, *Filomat*, 30.4, 2016, 905-912.
- [7] Kim T., Jang V. S. and Seo J. J., A note on poly-Genocchi numbers and polynomials, *Appl. Math. Sci.*, 8 (96), (2014), 4775-4781.
- [8] Known H.-In, Kim T. and Seo J. J., Modified degenerate Euler polynomials, *Advanced Studies in Contemp. Math.*, 26(1), (2016), 1-9.
- [9] Liu H. and Wang W., Some identities on the Bernoulli, Euler and Genocchi polynomials via power sum and alternate power sums, *Discrete Math.*, 309 (2009), 3346-3363.
- [10] Luo M.-Q., The multiplication formulas for the Apostol-Bernoulli and Apostol-Euler polynomials of higher order, *Integral Trans. Spec. Func.*, 20 (2009), 337-391.
- [11] Ozarslan M. A., Hermite-based unified Apostol-Bernoulli, Euler and Genocchi polynomials, *Advances in Diff. Equa.*, 2013.2013:116.
- [12] Ryoo C. S., A note on the Tangent numbers and polynomials, *Adv. Stud. Theo. Phys.*, 7(9), (2013), 447-454.
- [13] Ryoo C. S., On degenerate Carlitz' s type  $q$ -Tangent numbers and polynomials associated with  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$ , *App. Math. Sci.*, 11(48), (2017), 2367-2375.
- [14] Ryoo C. S., Symmetric identities for degenerate  $(h, q)$ -Tangent polynomials associated with the  $p$ -adic integral on  $\mathbb{Z}_p$ , *Int. J. of Math. Anal.*, 11(8), (2017), 353-362.
- [15] Ryoo C. S. and Agarwal R. P., Some identities involving  $q$ -poly-Tangent numbers and polynomials and distribution of their zeros, *Advances in Diff. Equa.*, (2017), 2017.2013.
- [16] Son J.-W. and Kim M.-S., On poly-Eulerian numbers, *Bull. Korean Math. Soc.*, 36 (1996), 47-61.
- [17] Shin H. and Zeng J., The  $q$ -Tangent and  $q$ -secant numbers via continued fractions, *European J. of Combinatorics*, 31(7), (2010), 1689-1705.
- [18] Srivastava H. M., Some generalization and basic (or  $-q$ ) extension of the Bernoulli, Euler and Genocchi polynomials, *App. Math. and Infor. Sci.*, 5(3), (2011), 390-444.
- [19] Srivastava H. M. and Manocha H. L., A treatise on generating functions, John-Wiley and Soc., (1984), New York-Toronto.
- [20] Young P. T., Degenerate Bernoulli polynomials, generalized factorial sums and their application, *J. of Number Theory*, 128 (2008), 738-758.