

New Tribonacci Recurrence Relations and Addition Formulas

Kunle Adegoke*, Adenike Olatinwo and Winning Oyekanmi

Department of Physics and Engineering Physics,
Obafemi Awolowo University, 220005 Ile-Ife, Nigeria

Abstract

Only one three-term recurrence relation, namely, $W_r = 2W_{r-1} - W_{r-4}$, is known for the generalized Tribonacci numbers, W_r , $r \in \mathbb{Z}$, defined by $W_r = W_{r-1} + W_{r-2} + W_{r-3}$ and $W_{-r} = W_{-r+3} - W_{-r+2} - W_{-r+1}$, where W_0 , W_1 and W_2 are given, arbitrary integers, not all zero. Also, only one four-term addition formula is known for these numbers, which is, $W_{r+s} = T_{s-1}W_{r-1} + (T_{s-1} + T_{s-2})W_r + T_sW_{r+1}$, where $(T_r)_{r \in \mathbb{Z}}$ is the Tribonacci sequence, a special case of the generalized Tribonacci sequence, with $W_0 = T_0 = 0$ and $W_1 = W_2 = T_1 = T_2 = 1$. In this paper we discover three new three-term recurrence relations and two identities from which a plethora of new addition formulas for the generalized Tribonacci numbers may be discovered. We obtain a simple relation connecting the Tribonacci numbers and the Tribonacci-Lucas numbers. Finally, we derive quadratic and cubic recurrence relations for the generalized Tribonacci numbers.

1 Introduction

For $r \geq 3$, we define the generalized Tribonacci numbers W_r by the third order recurrence relation:

$$W_r = W_{r-1} + W_{r-2} + W_{r-3}, \quad (1.1)$$

where W_0 , W_1 and W_2 are arbitrary integers. By writing $W_{r-1} = W_{r-2} + W_{r-3} + W_{r-4}$ and subtracting this from relation (1.1), we see that W_r also obeys the useful three-term recurrence

$$W_r = 2W_{r-1} - W_{r-4}. \quad (1.2)$$

Extension of the definition of the generalized Tribonacci numbers to negative subscripts is provided by writing identity (1.2) as $W_{r+4} = 2W_{r+3} - W_r$; so that

$$W_{-r} = 2W_{-r+3} - W_{-r+4}. \quad (1.3)$$

Well known examples of W_r are the Tribonacci sequence, (T_r) , $r \in \mathbb{Z}$, for which $W = T$, $W_0 = T_0 = 0$, $W_1 = W_2 = T_1 = T_2 = 1$ and the Tribonacci-Lucas sequence, (K_r) , $r \in \mathbb{Z}$,

r	-20	-19	-18	-17	-16	-15	-14	-13	-12
T_r	-56	159	-103	0	56	-47	9	18	-20
K_r	795	-571	47	271	-253	65	83	-105	43
r	-11	-10	-9	-8	-7	-6	-5	-4	-3
T_r	7	5	-8	4	1	-3	2	0	-1
K_r	21	-41	23	3	-15	11	-1	-5	5
r	-2	-1	0	1	2	3	4	5	6
T_r	1	0	0	1	1	2	4	7	13
K_r	-1	-1	3	1	3	7	11	21	39
r	7	8	9	10	11	12	13	14	15
T_r	24	44	81	149	274	504	927	1705	3136
K_r	71	131	241	443	815	1499	2757	5071	9327
r	16	17	18	19	20	21	22	23	24
T_r	5768	10609	19513	35890	66012	121415	223317	410744	755476
K_r	17155	31553	58035	106743	196331	361109	664183	1221623	2246915

Table 1: The first few Tribonacci and Tribonacci-Lucas numbers.

for which $W = K$, $W_0 = K_0 = 3$, $W_1 = K_1 = 1$, $W_2 = K_2 = 3$. Table 1 shows the first few Tribonacci and Tribonacci-Lucas numbers for $-20 \leq r \leq 24$.

The following references contain useful information on the properties of the Tribonacci numbers and related: [1, 2, 3, 4, 5, 6].

Among other interesting results, we found the following three-term recurrence relations which are presumably new:

$$W_{r-16} = -103W_r + 56W_{r+1},$$

$$2W_{r-17} = 9W_r - 103W_{r-4}$$

and

$$W_{r-14} = 9W_{r+2} - 56W_{r-1}.$$

We also found a simple relation linking the Tribonacci numbers and the Tribonacci-Lucas numbers:

$$K_{r-2} = 5T_{r-1} - T_{r+1}.$$

2 Linear recurrence relations and addition formulas

Theorem 1. *The following identity holds for integers a , b , c , d and e :*

$$\begin{aligned} & (T_{a-c}T_{c-b}T_{b-a} + T_{b-c}T_{c-a}T_{a-b})W_{d+e} \\ &= (T_{b-c}T_{c-a}T_{e-b} - T_{b-c}T_{c-b}T_{e-a} + T_{c-b}T_{b-a}T_{e-c})W_{d+a} \\ & \quad + (T_{a-c}T_{c-b}T_{e-a} - T_{a-c}T_{c-a}T_{e-b} + T_{c-a}T_{a-b}T_{e-c})W_{d+b} \\ & \quad + (T_{b-c}T_{a-b}T_{e-a} - T_{a-b}T_{b-a}T_{e-c} + T_{a-c}T_{b-a}T_{e-b})W_{d+c} \end{aligned}$$

*Corresponding author: adegoke00@gmail.com

Proof. We seek to express a generalized Tribonacci number as a linear combination of three Tribonacci numbers. Let

$$W_{d+e} = f_1 T_{e-a} + f_2 T_{e-b} + f_3 T_{e-c}, \quad (2.1)$$

where a, b, c, d and e are arbitrary integers and the coefficients f_1, f_2 and f_3 are to be determined. Setting $e = a, e = b$ and $e = c$, in turn, we obtain three simultaneous equations:

$$W_{d+a} = f_2 T_{a-b} + f_3 T_{a-c}, \quad W_{d+b} = f_1 T_{b-a} + f_3 T_{b-c}, \quad W_{d+c} = f_1 T_{c-a} + f_2 T_{c-b}.$$

The identity of Theorem 1 is established by solving these equations for f_1, f_2 and f_3 and substituting the solutions into identity (2.1). □

Corollary 2. *The following identities hold for integers r and s :*

$$4W_{r+s} = 2T_{s-1}W_{r-4} + (T_{s+4} - 7T_s)W_r + 4T_sW_{r+1}, \quad (2.2)$$

$$4W_{r+s} = 2T_{s-4}W_{r-1} + (4T_{s+1} - 7T_s)W_r + T_sW_{r+4}, \quad (2.3)$$

$$W_{r+s} = T_{s-1}W_{r-1} + (T_{s+1} - T_s)W_r + T_sW_{r+1}, \quad (2.4)$$

$$4W_{r+s} = T_{s-4}W_{r-4} + (T_{s+4} - 11T_s)W_r + T_sW_{r+4}, \quad (2.5)$$

$$W_{r+s} = (T_{s+1} - 2T_s - T_{s-2})W_{r-1} + (T_{s+1} - 2T_s)W_r + T_sW_{r+2}. \quad (2.6)$$

Proof. See Table 2.

Identity	a	b	c	d	e	Identity	a	b	c	d	e
(2.2)	$r-4$	r	$r+1$	0	$r+s$	(2.5)	$r+4$	r	$r-4$	0	$r+s$
(2.3)	$r+4$	r	$r-1$	0	$r+s$	(2.6)	$r+2$	$r-1$	r	0	$r+s$
(2.4)	$r-1$	r	$r+1$	0	$r+s$						

Table 2: Appropriate substitutions in the identity of Theorem 1 to obtain identities (2.2) – (2.6) of Corollary 2. □

Note that identity (2.4) can be written in the familiar form

$$W_{r+s} = T_{s-1}W_{r-1} + (T_{s-1} + T_{s-2})W_r + T_sW_{r+1}. \quad (2.7)$$

Evaluating identity (2.2) at $s = -3, s = -16$ and at $s = -17$, in turn, we find the following three term recurrence relations for the generalized Tribonacci numbers:

$$W_{r-3} = 2W_r - W_{r+1}, \quad (2.8)$$

$$W_{r-16} = -103W_r + 56W_{r+1} \quad (2.9)$$

and

$$2W_{r-17} = 9W_r - 103W_{r-4}. \quad (2.10)$$

Evaluating identity (2.6) at $s = -14$ produces yet another three-term recurrence relation for the generalized Tribonacci numbers:

$$W_{r-14} = 9W_{r+2} - 56W_{r-1}. \quad (2.11)$$

Note that by interchanging r and s in each case and making use of the defining recurrence relation for W and T , the identities (2.2) – (2.6) can also be written

$$4W_{r+s} = 2W_{s-1}T_{r-4} + (W_{s+4} - 7W_s)T_r + 4W_sT_{r+1}, \quad (2.12)$$

$$4W_{r+s} = 2W_{s-4}T_{r-1} + (4W_{s+1} - 7W_s)T_r + W_sT_{r+4}, \quad (2.13)$$

$$W_{r+s} = W_{s-1}T_{r-1} + (W_{s+1} - W_s)T_r + W_sT_{r+1}, \quad (2.14)$$

$$4W_{r+s} = W_{s-4}T_{r-4} + (W_{s+4} - 11W_s)T_r + W_sT_{r+4}, \quad (2.15)$$

$$W_{r+s} = (W_{s+1} - 2W_s - W_{s-2})T_{r-1} + (W_{s+1} - 2W_s)T_r + W_sT_{r+2}. \quad (2.16)$$

Evaluating identity (2.14) at $s = -2$ with $W = K$ gives

$$K_{r-2} = 5T_{r-1} - T_{r+1}, \quad (2.17)$$

a three-term relation connecting the Tribonacci numbers and the Tribonacci-Lucas numbers.

Theorem 3. *The following identity holds for integers a, b, c, d and e :*

$$\begin{aligned} & \{K_{b-a-1}K_{a-b-1} + K_{b-a-1}K_{c-b-1}K_{a-c-1} + K_{c-a-1}K_{b-c-1}K_{a-b-1} \\ & \quad + K_{c-a-1}K_{a-c-1} + K_{c-b-1}K_{b-c-1} - 1\}W_{d+e} \\ & = \{(1 - K_{c-b-1}K_{b-c-1})K_{e-a} + (K_{b-a-1} + K_{c-a-1}K_{b-c-1})K_{e-b} \\ & \quad + (K_{c-a-1} + K_{b-a-1}K_{c-b-1})K_{e-c}\}W_{d+a-1} \\ & + \{(1 - K_{c-a-1}K_{a-c-1})K_{e-b} + (K_{a-b-1} + K_{c-b-1}K_{a-c-1})K_{e-a} \\ & \quad + (K_{c-b-1} + K_{c-a-1}K_{a-b-1})K_{e-c}\}W_{d+b-1} \\ & + \{(1 - K_{b-a-1}K_{a-b-1})K_{e-c} + (K_{b-c-1} + K_{b-a-1}K_{a-c-1})K_{e-b} \\ & \quad + (K_{a-c-1} + K_{b-c-1}K_{a-b-1})K_{e-a}\}W_{d+c-1}. \end{aligned}$$

Proof. We wish to express a generalized Tribonacci number as a linear combination of three Tribonacci-Lucas numbers. Let

$$W_{d+e} = f_1K_{e-a} + f_2K_{e-b} + f_3K_{e-c}, \quad (2.18)$$

where a, b, c, d and e are arbitrary integers and the coefficients f_1, f_2 and f_3 are to be determined. Setting $e = a-1, e = b-1$ and $e = c-1$, in turn, we obtain three simultaneous equations:

$$\begin{aligned} W_{d+a-1} &= -f_1 + f_2K_{a-b-1} + f_3K_{a-c-1}, & W_{d+b-1} &= f_1K_{b-a-1} - f_2 + f_3K_{b-c-1}, \\ W_{d+c-1} &= f_1K_{c-a-1} + f_2K_{c-b-1} - f_3. \end{aligned}$$

The identity of Theorem 3 is obtained by solving these equations for f_1, f_2 and f_3 and substituting the solutions into identity (2.18). □

Corollary 4. *The following identities hold for integers r and s :*

$$\begin{aligned} 44W_{r+s} &= (9K_{s+3} - K_{s+5} + 2K_{s+1})W_{r-6} \\ &+ (9K_{s+1} + K_{s+5} + 2K_{s+3})W_{r-4} \\ &+ (K_{s+3} + 6K_{s+5} - K_{s+1})W_{r-2}, \end{aligned} \quad (2.19)$$

$$\begin{aligned}
88W_{r+s} &= (8K_{s+3} - 30K_{s-1} - 2K_{s-2})W_r \\
&\quad + (K_{s+3} - K_{s-1} + 8K_{s-2})W_{r-4} \\
&\quad + (3K_{s+3} + 19K_{s-1} + 2K_{s-2})W_{r+1},
\end{aligned} \tag{2.20}$$

$$\begin{aligned}
22W_{r+s} &= (5K_{s-2} + K_{s-1} + 2K_s)W_{r-1} \\
&\quad + (K_{s-2} - 2K_{s-1} + 7K_s)W_r \\
&\quad + (2K_{s-2} + 7K_{s-1} + 3K_s)W_{r+1},
\end{aligned} \tag{2.21}$$

$$\begin{aligned}
5060W_{r+s} &= (-264K_{s-3} + K_{s+9} - 1079K_{s-1})W_{r-10} \\
&\quad + (-4279K_{s-3} + 21K_{s+9} - 21394K_{s-1})W_r \\
&\quad + (6325K_{s-1} + 1265K_{s-3})W_{r+2}
\end{aligned} \tag{2.22}$$

and

$$\begin{aligned}
5060W_{r+s} &= (-264K_{s-11} + 1265K_{s+1} - 4279K_{s-1})W_{r-2} \\
&\quad + (-1079K_{s-11} + 6325K_{s+1} - 21394K_{s-1})W_r \\
&\quad + (21K_{s-1} + K_{s-11})W_{r+10}.
\end{aligned} \tag{2.23}$$

Identities (2.19) – (2.23) can also be written as follows.

$$\begin{aligned}
44W_{r+s} &= (9W_{s+3} - W_{s+5} + 2W_{s+1})K_{r-6} \\
&\quad + (9W_{s+1} + W_{s+5} + 2W_{s+3})K_{r-4} \\
&\quad + (W_{s+3} + 6W_{s+5} - W_{s+1})K_{r-2},
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
88W_{r+s} &= (8W_{s+3} - 30W_{s-1} - 2W_{s-2})K_r \\
&\quad + (W_{s+3} - W_{s-1} + 8W_{s-2})K_{r-4} \\
&\quad + (3W_{s+3} + 19W_{s-1} + 2W_{s-2})K_{r+1},
\end{aligned} \tag{2.25}$$

$$\begin{aligned}
22W_{r+s} &= (5W_{s-2} + W_{s-1} + 2W_s)K_{r-1} \\
&\quad + (W_{s-2} - 2W_{s-1} + 7W_s)K_r \\
&\quad + (2W_{s-2} + 7W_{s-1} + 3W_s)K_{r+1},
\end{aligned} \tag{2.26}$$

$$\begin{aligned}
5060W_{r+s} &= (-264W_{s-3} + W_{s+9} - 1079W_{s-1})K_{r-10} \\
&\quad + (-4279W_{s-3} + 21W_{s+9} - 21394W_{s-1})K_r \\
&\quad + (6325W_{s-1} + 1265W_{s-3})K_{r+2}
\end{aligned} \tag{2.27}$$

and

$$\begin{aligned}
5060W_{r+s} &= (-264W_{s-11} + 1265W_{s+1} - 4279W_{s-1})K_{r-2} \\
&\quad + (-1079W_{s-11} + 6325W_{s+1} - 21394W_{s-1})K_r \\
&\quad + (21W_{s-1} + W_{s-11})K_{r+10}
\end{aligned} \tag{2.28}$$

3 Quadratic relations

Our goal in this section is to derive expressions involving only pure squares of generalized Tribonacci numbers. To achieve this we must be able to express the anticipated cross-terms such as $W_{r-1}W_r$ and $W_{r-1}W_{r-4}$ as squares of generalized Tribonacci numbers.

Rearranging identity (1.2) and squaring, we have

$$4W_{r-1}W_r = 4W_{r-1}^2 - W_{r-4}^2 + W_r^2, \tag{3.1}$$

$$4W_{r-1}W_{r-4} = 4W_{r-1}^2 + W_{r-4}^2 - W_r^2 \quad (3.2)$$

and

$$2W_rW_{r-4} = 4W_{r-1}^2 - W_{r-4}^2 - W_r^2. \quad (3.3)$$

Rearranging identity (1.2) and multiplying through by $4W_{r-3}$ to obtain

$$8W_{r-1}W_{r-3} = 4W_rW_{r-3} + 4W_{r-4}W_{r-3}, \quad (3.4)$$

and using identities (3.1) and (3.2) to resolve the right hand side gives

$$8W_{r-1}W_{r-3} = 4W_r^2 + 2W_{r-3}^2 - W_{r+1}^2 + 4W_{r-4}^2 - W_{r-7}^2. \quad (3.5)$$

Multiplying through identity (1.2) by $4W_{r-5}$ to obtain

$$4W_rW_{r-5} = 8W_{r-1}W_{r-5} - 4W_{r-4}W_{r-5}, \quad (3.6)$$

which, with the use of (3.1) and (3.3), translates to

$$4W_rW_{r-5} = 16W_{r-2}^2 - 8W_{r-5}^2 - 4W_{r-1}^2 + W_{r-8}^2 - W_{r-4}^2. \quad (3.7)$$

Rearranging identity (1.2), shifting index r and multiplying through by $2W_r$ gives

$$2W_rW_{r-8} = 4W_rW_{r-5} - 2W_rW_{r-4}, \quad (3.8)$$

from which, using (3.3) and (3.7), we get

$$2W_rW_{r-8} = 16W_{r-2}^2 - 8W_{r-5}^2 - 8W_{r-1}^2 + W_{r-8}^2 + W_r^2. \quad (3.9)$$

Re-arranging identity (2.9) and squaring, we have

$$112W_{r-17}W_r = W_{r-17}^2 + 3136W_r^2 - 10609W_{r-1}^2 \quad (3.10)$$

$$11536W_{r-1}W_r = -W_{r-17}^2 + 3136W_r^2 + 10609W_{r-1}^2, \quad (3.11)$$

and

$$206W_{r-17}W_{r-1} = -W_{r-17}^2 - 10609W_{r-1}^2 + 3136W_r^2. \quad (3.12)$$

Rearranging and squaring identity (2.10), we have

$$36W_{r-17}W_r = 4W_{r-17}^2 + 81W_r^2 - 10609W_{r-4}^2, \quad (3.13)$$

$$1854W_rW_{r-4} = 81W_r^2 + 10609W_{r-4}^2 - 4W_{r-17}^2 \quad (3.14)$$

and

$$412W_{r-17}W_{r-4} = -4W_{r-17}^2 - 10609W_{r-4}^2 + 81W_r^2. \quad (3.15)$$

Finally, squaring and re-arranging identity (2.11) produces

$$18W_{r-17}W_{r-1} = W_{r-17}^2 + 81W_{r-1}^2 - 3136W_{r-4}^2, \quad (3.16)$$

$$1008W_{r-1}W_{r-4} = 81W_{r-1}^2 + 3136W_{r-4}^2 - W_{r-17}^2 \quad (3.17)$$

and

$$112W_{r-17}W_{r-4} = -W_{r-17}^2 - 3136W_{r-4}^2 + 81W_{r-1}^2. \quad (3.18)$$

Theorem 5. *The following identities hold for r an integer:*

$$252W_r^2 - 927W_{r-1}^2 + 2884W_{r-4}^2 - W_{r-17}^2 = 0, \quad (3.19)$$

$$W_r^2 - 2W_{r-1}^2 - 3W_{r-2}^2 - 6W_{r-3}^2 + W_{r-4}^2 + W_{r-6}^2 = 0. \quad (3.20)$$

Proof. Eliminating $W_{r-1}W_r$ between identities (3.1) and (3.11) proves identity (3.19). To prove identity (3.20), write $W_r - W_{r-1} = W_{r-2} + W_{r-3}$, square both sides and use the identity (3.1) to resolve the cross-products W_rW_{r-1} and $W_{r-2}W_{r-3}$. \square

Substituting for W_{r-17}^2 from the identity of Theorem 5 into identities (3.10), (3.12) and (3.18), we have the following simpler versions of these identities:

$$4W_{r-17}W_r = -412W_{r-1}^2 + 103W_{r-4}^2 + 121W_r^2, \quad (3.21)$$

$$W_{r-17}W_{r-1} = -47W_{r-1}^2 - 14W_{r-4}^2 + 14W_r^2 \quad (3.22)$$

and

$$4W_{r-17}W_{r-4} = 36W_{r-1}^2 - 215W_{r-4}^2 - 9W_r^2. \quad (3.23)$$

Next we give how to express the square of a Tribonacci-Lucas number in terms of squares of Tribonacci numbers.

Theorem 6. *The following identity holds for any integer r :*

$$4K_r^2 = 5T_{r+5}^2 - 20T_{r+4}^2 + 4T_{r+3}^2 + 90T_{r+1}^2 - 20T_r^2 + 5T_{r-3}^2. \quad (3.24)$$

Proof. Square identity (2.17) and use identity (3.5) to eliminate the cross term. \square

Theorem 7. *The following identities hold for r and s integers:*

$$\begin{aligned} 16W_{r+s}^2 &= -(T_s + T_{s+4})(-T_{s+4} + 2T_{s-1} + 7T_s)W_r^2 + 4T_{s-1}T_sW_{r-7}^2 \\ &\quad + 2T_{s-1}(-T_{s+4} - 9T_s + 2T_{s-1})W_{r-4}^2 - 2T_s(T_{s+4} - 7T_s + 2T_{s-1})W_{r-3}^2 \\ &\quad + 8T_{s-1}(T_s + T_{s+4})W_{r-1}^2 + 2T_s(T_s + T_{s+4})W_{r+1}^2, \end{aligned} \quad (3.25)$$

$$\begin{aligned} 16W_{r+s}^2 &= 4(2T_s - T_{s+1})(7T_s - T_{s-4} - 4T_{s+1})W_r^2 + 4T_{s-4}(2T_s - T_{s+1})W_{r-4}^2 \\ &\quad - 4T_{s-4}(-T_{s-4} - 4T_{s+1} + 9T_s)W_{r-1}^2 + 16T_{s-4}T_sW_{r+2}^2 \\ &\quad - 4T_s(7T_s + T_{s-4} - 4T_{s+1})W_{r+3}^2 + 4T_s(2T_s - T_{s+1})W_{r+4}^2, \end{aligned} \quad (3.26)$$

$$\begin{aligned} 4W_{r+s}^2 &= -2(-T_{s+1} + T_s)(2T_{s+2} - T_{s-1})W_r^2 - T_{s-1}T_sW_{r-5}^2 \\ &\quad + 2T_{s-1}(-T_{s+1} + T_s)W_{r-4}^2 + 2T_s(-T_{s+1} + T_s)W_{r-3}^2 \\ &\quad + 4T_{s-1}T_sW_{r-2}^2 + 2T_{s-1}(2T_{s-1} + 4T_{s+1} - 3T_s)W_{r-1}^2 \\ &\quad + 2T_s(T_s + T_{s+1})W_{r+1}^2 - T_{s-1}T_sW_{r+3}^2 + 4T_{s-1}T_sW_{r+2}^2. \end{aligned} \quad (3.27)$$

4 Cubic recurrence relations

Theorem 8. *The following identity holds for integer r :*

$$\begin{aligned} W_r^3 - 4W_{r-1}^3 - 9W_{r-2}^3 - 34W_{r-3}^3 + 24W_{r-4}^3 - 2W_{r-5}^3 \\ + 40W_{r-6}^3 - 14W_{r-7}^3 - W_{r-8}^3 - 2W_{r-9}^3 + W_{r-10}^3 = 0. \end{aligned}$$

Proof. Setting $a = r - 8$, $b = r$, $c = r - 10$, $d = 0$ and $e = r - s$ with $s \in \{1, 2, 3, 4, 5, 6, 7, 9\}$ in the identity of Theorem 1, the following linear combinations are formed:

$$\begin{aligned} W_{r-1} &= \frac{37}{68}W_r - \frac{5}{68}W_{r-8} + \frac{1}{17}W_{r-10}, & W_{r-2} &= \frac{5}{17}W_r + \frac{3}{17}W_{r-8} + \frac{1}{17}W_{r-10}, \\ W_{r-3} &= \frac{11}{68}W_r - \frac{7}{68}W_{r-8} - \frac{2}{17}W_{r-10}, & W_{r-4} &= \frac{3}{34}W_r - \frac{5}{34}W_{r-8} + \frac{2}{17}W_{r-10}, \\ W_{r-5} &= \frac{3}{68}W_r + \frac{29}{68}W_{r-8} + \frac{1}{17}W_{r-10}, & W_{r-6} &= \frac{1}{34}W_r - \frac{13}{34}W_{r-8} - \frac{5}{17}W_{r-10}, \\ W_{r-7} &= \frac{1}{68}W_r - \frac{13}{68}W_{r-8} + \frac{6}{17}W_{r-10}, & W_{r-9} &= \frac{1}{68}W_r - \frac{81}{68}W_{r-8} - \frac{11}{17}W_{r-10}. \end{aligned}$$

The above W_{r-i} , $i \in \{1, 2, 3, 4, 5, 6, 7, 9\}$ verify the identity of Theorem 8. \square

Taking the cube in identities (2.8) and (2.9) and solving two simultaneous equations, we find

$$155736W_rW_{r-1}^2 = -W_{r-17}^3 + 199305W_{r-1}^3 - 161504W_{r-4}^3 + 14112W_r^3 \quad (4.1)$$

and

$$77868W_r^2W_{r-1} = -148526W_{r-4}^3 + 27090W_r^3 - W_{r-17}^3 + 95481W_{r-1}^3. \quad (4.2)$$

Theorem 9. *The following identity holds for integer r :*

$$\begin{aligned} 11844W_r^3 + 3W_{r-19}^3 + W_{r-17}^3 + 458556W_{r-6}^3 \\ + 135548W_{r-4}^3 - 442179W_{r-3}^3 - 120204W_{r-2}^3 - 43569W_{r-1}^3 = 0. \end{aligned} \quad (4.3)$$

Proof. Write the defining recurrence relation of the generalized Tribonacci numbers as $W_r - W_{r-1} = W_{r-2} + W_{r-3}$, take the cube of both sides and use identities (4.1) and (4.2) to remove cross terms. \square

References

- [1] K. Adegoke, Weighted Tribonacci sums, *arXiv:1804.06449[math.CA]* (2018).
- [2] J. Feng, More identities on the Tribonacci numbers, *Ars Combinatorial* **C** (2011), 73–78.
- [3] R. Frontczak, Sums of Tribonacci and Tribonacci-Lucas numbers, *International Journal of Mathematical Analysis* **12**:1 (2018), 19–24.
- [4] H. Gabai, Generalized Fibonacci k -sequences, *The Fibonacci Quarterly* **8**:1 (1970), 31–38.
- [5] D. V. Shah, Some Tribonacci identities, *Mathematics Today* **27** (2011), 1–9.
- [6] M. E. Waddill and L. Sacks, Another generalized Fibonacci sequence, *The Fibonacci Quarterly* **5**:3 (1967), 209–222.

2010 *Mathematics Subject Classification*: Primary 11B39; Secondary 11B37.

Keywords: Tribonacci number, Tribonacci-Lucas number, recurrence relation.
