

ULTRA CONTRACTIVITY AND GAUSSIAN BOUNDS FOR EVOLUTION FAMILIES ASSOCIATED WITH NON-AUTONOMOUS FORMS

HAFIDA LAASRI AND DELIO MUGNOLO

ABSTRACT. We develop a variational approach in order to study qualitative properties of non-autonomous parabolic equations. Based on the method of product integrals, we discuss invariance properties and ultracontractivity of evolution families in Hilbert space. Our main results give sufficient conditions for the heat kernel of the evolution family to satisfy Gaussian-type bounds. Along the way, we study examples of non-autonomous equations on graphs, metric graphs, and domains.

1. INTRODUCTION

Non-autonomous evolution equations are partial differential equations in which relevant coefficients of the differential operator and/or in the boundary conditions are time-dependent, thus allowing for underlying models that are variable over time.

In the autonomous case (i.e., evolution equations with time-independent coefficients), well-posedness is known to be equivalent to generation of a semigroup in a suitable Banach space; in comparison, the theory for well-posedness of non-autonomous problems on general Banach spaces is more rudimentary. If the coefficients of a non-autonomous equation are piecewise constant, then one may find a solution by following the orbit of the semigroup governing a given problem as long as the coefficient stays constant; then “freeze” the system; use the final state as an initial condition for a new evolution equation with new (constant) coefficient, and so on: this boils down to consider the composition of a finite numbers of semigroups.

A theory originally developed by J.-L. Lions shows that well-posedness in Hilbert space can be proved under much weaker assumptions, most notably mere measurability of the time dependence, provided the problem has a nice variational structure: this is typically the case if the differential equation is parabolic. By adapting the setting of (time-independent) bounded elliptic forms it is thus possible to show that the equation has a solution that is, in particular, continuous in time, cf. Theorem 2.1 for a precise statement. This motivates the study of *non-autonomous forms*, a topic which has received much attention in the last decade: we mention among others [ADLO14, AD17, ADF17, Ouh15, Fac17]. In all these articles, the focus lies on properties of solutions of partial differential equations, with a focus on maximal regularity issues and hence allowing for inhomogeneous terms.

Our main aim in this paper is to develop a more abstract theory with an operator theoretical flavor. Indeed, Lions’ result paves the way to the possibility of defining an *evolution family* (or

2010 *Mathematics Subject Classification.* 47D06, 47D07, 47A07, 35K90.

Key words and phrases. Evolution families, non-autonomous parabolic problems, kernel estimates.

The first author was supported by the Deutsche Forschungsgemeinschaft (Grant LA 4197/1-1). The second author was partially supported by the Deutsche Forschungsgemeinschaft (Grant 397230547).

evolution system, or *propagator*), i.e., a family of operators $U(\cdot, s)$ mapping each initial data

$$u(s) = x \in H$$

to the orbit of the solution

$$\dot{u}(t) + A(t)u(t) = 0 \quad \text{a.e. on } [s, T].$$

Because the initial condition may well be imposed at instants $s \neq 0$, this actually define a two-parameter family

$$\mathcal{U} := (U(t, s))_{(t,s) \in \bar{\Delta}}$$

of bounded linear operators on H by $U(t, s)x := u(t)$, where

$$\Delta := \{(t, s) \in (0, T)^2 : s < t\}.$$

Some good compendia on such evolution families are [Tan79, Chapt. 7], [Paz83, Chapt. 5], [Fat83, Chapt. 7], [EN00, Section VI.9], or the monograph [CL99].

The tumultuous development of Hilbert space methods, and especially the theory of Dirichlet forms, have been fruitful also in the non-autonomous environment: a theory of non-autonomous Dirichlet forms has been recently introduced in [ADO14]. If $A(t) \equiv A$, the above abstract Cauchy problem is autonomous and its solution is simply given by

$$u(t) = U(t, s)x := e^{-(t-s)A}x ;$$

hence the findings in [ADO14] can be regarded as a strict generalization of the classical theory of Markovian operators and Dirichlet forms represented e.g. in [FOT10]. Our goal is to complement these results, thus setting up a non-autonomous variational program analogous to the autonomous one outlined in classical monographs like [Ouh05, Are06]: to this aim we study further operator theoretical properties, including extrapolation to L^p -spaces, ultracontractivity, or Gaussian-type bounds on integral kernels of evolution families. It should be mentioned that ultracontractivity and kernel estimates have been observed already in [Aro68, Dan00] for specific instances of parabolic non-autonomous equations; in particular, Aronson observed in [Aro67] that the fundamental solution $(t, s; x, y) \mapsto \Gamma(t, s; x, y)$ of a certain class of non-autonomous diffusion equations in (domains of) \mathbb{R}^d satisfies

$$(1.1) \quad \Gamma(t, s; x, y) \leq K G(t - s; x - y)$$

where $(t, x) \mapsto G(t, x)$ is the Gaussian kernel that yields the fundamental solution of the (autonomous) heat equation on \mathbb{R}^d . Analogous Gaussian bounds have ever since been proved for integral kernels of semigroups generated by large classes of second-order elliptic operators, possibly with complex coefficients [Ouh04]; in the non-autonomous case, Aronson's original findings have been extended to operators on domains under Dirichlet or Robin boundary conditions in [Aro68, Dan00].

In this paper we are going to introduce a general approach, based on the so-called Davies' Trick, to prove Gaussian bounds for heat kernels of evolution families that govern non-autonomous parabolic equations. Inspired by some techniques introduced in [Dan00, Ouh04], we show the applicability of our methods by showing that a large class of elliptic operators with complex-valued, bounded measurable (both in time and space) coefficients are associated with evolution families that satisfy Gaussian bounds, thus extending the main results in [Ouh04] to the non-autonomous setting.

Our approach will heavily rely upon the method of product integrals, whose historical evolution is thoroughly discussed in [Sla07], [Fat83, § 7.10] and whose scope has been extended to non-autonomous forms with measurable dependence on time in [EL16, SL15]. We adapt it to our present

setting, thus deriving in Theorem 2.5 a version that we will use over and over again in different contexts throughout this paper.

The present paper is organized as follows. After describing our mathematical framework in Section 2, in Section 3 we present sufficient conditions that enforce qualitative properties based on the lattice structure of L^2 -spaces, including stochasticity or domination. Also, we are able to discuss cases where evolution families on L^2 -spaces extrapolate to further L^p -spaces. This is a key feature of the theory of Dirichlet forms in the autonomous case and shows the flexibility of the Hilbert space approach in the non-autonomous context, too. Even evolution equations on structures that change over time can be studied by means of our theory.

Gaussian-type bounds are shown to depend on ultracontractivity properties of certain operator families related to \mathcal{U} . This approach requires, in turn, suitable common bounds in L^p -norm, uniformly on all compact subsets of Δ . Inspired by similar criteria in the autonomous setting we show that efficient conditions based on Sobolev-type inequalities can enforce such bounds. In Section 4 we develop a theory of ultracontractive evolution families: a technical difficulty we face is related to the failure of self-adjointness of evolution families, a phenomenon that typically occurs even when all operators $A(t)$ are self-adjoint. We take over an idea from [Dan00] and circumvent this problem by studying some non-autonomous form associated with a tightly related backward evolution equation.

It has been known since [Dav87] that ultracontractivity is an important ingredient to prove Gaussian bounds for semigroups. Expanding the scope of Davies' trick, in Section 5 we are going to present different sufficient conditions under which an evolution family satisfies Gaussian bounds.

In particular, our approach allows us to show Gaussian bounds for the evolution family associated with a large class of elliptic operators, thus generalizing the pioneering results in [Aro68, Dan00].

Several applications of increasing complexity are reviewed in Section 6: we discuss well-posedness and qualitative properties of dynamical systems on undirected graphs tightly related to the theory of *dynamic (positive) graphs* discussed in [Sil08] as well as models of Black–Scholes-types equations with time-dependent volatility [Hes93]; we extend the kernel estimates in [Mug07] to more general non-autonomous diffusion equations on possibly infinite networks; and finally, we prove Gaussian bounds for the heat kernel for a large class of elliptic operators with time-dependent, possibly complex coefficients, thus deducing the main results in [Dan00, Ouh04] as special cases.

2. EVOLUTION FAMILIES: NOTATIONS AND PRELIMINARY RESULTS

Throughout this paper H is a separable, complex Hilbert space and V is a further complex Hilbert space that is densely and continuously embedded into H . Let V' denote the antidual of V with respect to the pivot space H ; the duality between V' and V is denoted by $\langle \cdot, \cdot \rangle$. We also denote by $(\cdot | \cdot)_V$ and $\| \cdot \|_V$ the scalar product and the norm on V , respectively; and by $(\cdot | \cdot)$ and $\| \cdot \|$ the corresponding quantities in H .

We fix $T \in]0, \infty[$ and consider a time-dependent family $(a(t))_{t \in [0, T]}$ of mappings such that $a(t; \cdot, \cdot) : V \times V \rightarrow \mathbb{C}$ is for all $t \in [0, T]$ a sesquilinear form and

$$(2.1) \quad [0, T] \ni t \mapsto a(t; u, v) \in \mathbb{C} \text{ is measurable} \quad \text{for all } u, v \in V;$$

and such that furthermore there exist constants $M, \alpha > 0$ and $\omega \geq 0$ such that the boundedness and H -ellipticity estimates

$$(2.2) \quad |a(t; u, v)| \leq M \|u\|_V \|v\|_V \quad \text{for a.e } t \in [0, T] \text{ and } u, v \in V,$$

$$(2.3) \quad \operatorname{Re} a(t; u, u) + \omega \|u\|_H^2 \geq \alpha \|u\|_V^2 \quad \text{for a.e } t \in [0, T] \text{ and } u \in V,$$

hold. In what follows we call such a family $\mathbf{a} := (a(t))_{t \in [0, T]}$ *bounded H -elliptic non-autonomous form*: following [AD17] we denote by $\text{FORM}([0, T]; V, H)$ the class of all such forms.

By the Lax–Milgram theorem, for each $t \in [0, T]$ there exists an *operator associated* with $a(t, \cdot, \cdot)$, i.e., an isomorphism $A(t) : V \rightarrow V'$ such that

$$\langle A(t)u, v \rangle = a(t, u, v) \quad \text{for all } u, v \in V :$$

accordingly we refer to the family $(A(t))_{t \in [0, T]}$ as the *operator family* associated with $\mathbf{a} := (a(t))_{t \in [0, T]}$.

Regarded as an unbounded operator with domain V , $-A(t)$ generates a holomorphic semigroup on V' , and in fact by [Are06, Thm. 7.1.5] on H too, since $a(t)$ is for all t a bounded, H -elliptic sesquilinear form: with an abuse of notation we denote its generator – the part of $-A(t)$ in H – again by $-A(t)$, and the semigroup by

$$\mathcal{T}_t := \{e^{-rA(t)} \mid r \geq 0\}.$$

Hence, for each fixed $t, s \in [0, T]$ the Cauchy problem

$$\begin{aligned} \dot{u}(r) + A(t)u(r) &= 0, & r \in [s, T], \\ u(s) &= x \in H, \end{aligned}$$

is well-posed, its solution being given by $u(r, x) := e^{-(r-s)A(t)}x$. However, we are rather going to focus on the non-autonomous Cauchy problem

$$(2.4) \quad \begin{aligned} \dot{u}(t) + A(t)u(t) &= 0, & t \in [s, T], \\ u(s) &= x \in H. \end{aligned}$$

The Hilbert space setting discussed above is rather benign and we can combine a few known results to observe the following non-autonomous counterpart of the Lumer–Phillips Theorem. Throughout this article we adopt the notation

$$\Delta := \{(t, s) \in (0, T)^2 : s < t\}.$$

The following result has already been proved under slightly stronger assumptions in [Laa18, Prop. 2.1]. We re-formulate it for the sake of self-containedness.

Theorem 2.1. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ and $(A(t))_{t \in [0, T]}$ be the associated operator family. Then for all $s \in [0, T)$ and all $x \in H$ there is a unique $[s, T] \ni t \mapsto u(t) \in MR(V, V')$, where*

$$MR(V, V') := MR(s, T; V, V') := L^2(s, T; V) \cap H^1(s, T; V'),$$

that satisfies

- $\dot{u}(t) + A(t)u(t) = 0$ (as equality of elements of V')¹ for a.e. $t \in (s, T)$ and
- $u(s) = x$ (as equality of elements of H).

Because $MR(V, V') \hookrightarrow C([s, T]; H)$,

$$U(\cdot, s) : (s, T] \times H \ni (t, x) \mapsto u(t) =: U(t, s)x \in H$$

defines a strongly continuous evolution family on H , i.e., the following properties hold:

- (i) $U(s, s) = \text{Id}_H$ for all $s \in [0, T]$,
- (ii) $U(t, s) = U(t, r)U(r, s)$ for all $0 \leq s \leq r \leq t \leq T$,
- (iii) $(t, s) \mapsto U(t, s)x$ is for all $x \in H$ continuous from $\overline{\Delta}$ into H .

¹ Even as equality of elements of H if additionally the dependence $t \mapsto a(t)$ is Hölder continuous of exponent $\beta > \frac{1}{2}$; or else of bounded variation.

In the following we will refer to $\mathcal{U} := (U(t, s))_{(t, s) \in \overline{\Delta}}$ as the *evolution family associated with the non-autonomous form \mathbf{a}* or *with the operator family $(A(t))_{t \in [0, T]}$* .

Proof. Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$. A classical well-posedness theorem by J.-L. Lions [DL92, §§ XVIII.3.2–3] states that for each $s \in [0, T)$ and each $x \in H$ (2.4) admits a unique solution u in the *maximal regularity space* $MR(V, V')$. It is well-known that $MR(V, V')$ is continuously embedded into $C([s, T]; H)$, see e.g. [Sho97, Prop. III.1.2]: this allows us to introduce a family of linear operators by

$$(2.5) \quad U(t, s) : H \ni x \mapsto U(t, s)x := u(t) \in H, \quad (t, s) \in \overline{\Delta},$$

where u is the unique solution of the Cauchy problem (2.4) in $MR(V, V')$. The proof of (i)–(iii) now follows from [ACFP07, Prop. 2.3 and Prop. 2.4] letting $X := V'$ and $D := V$, whence in particular $Tr = H$. \square

Remark 2.2. *If in fact $\mathbf{a} \in \text{FORM}([0, \infty[; V, H)$, then arguing as above we deduce that*

$$\begin{aligned} \dot{u}(t) + A(t)u(t) &= 0, & t \in [s, \infty), \\ u(s) &= x \in H, \end{aligned}$$

has for all $s > 0$ and all $x \in H$ a unique solution $u \in L^2_{Loc}(s, \infty; V) \cap H^1_{Loc}(s, \infty; V')$, hence $u \in C([0, \infty); H)$: this defines a strongly continuous evolution family $(U(t, s))_{0 \leq s \leq t < \infty}$.

Remark 2.3. *It follows from the above mentioned result by Lions that $U(t, s)$ maps H into V for all $s \in [0, T]$ and almost all $t \in [0, T]$ with $(t, s) \in \Delta$. Thus using the evolution family law (ii) we deduce that $U(t, s)$ is a compact operator on H for all $(t, s) \in \Delta$ provided V is compactly embedded in H . Likewise, if the embedding of V in H is of p -Schatten class for some $p \in [1, \infty[$, then $U(t, s)$ is of p -Schatten class – hence by (ii) of trace class – for all $(t, s) \in \Delta$.*

Remark 2.4. *A non-autonomous form is called coercive if (2.3) is satisfied with $\omega = 0$. Now observe that \mathbf{a} satisfies (2.3) if and only if the form \mathbf{a}_ω given by*

$$\mathbf{a}_\omega(t; u, v) := a(t; u, v) + \omega(u | v)$$

is coercive: because $\mathbf{a}_\omega \in \text{FORM}([0, T]; V, H)$ in its own right, it is associated with an evolution family. Moreover, u is a solution of class $MR(V, V')$ of (2.4) if and only if $v := e^{-\omega(\cdot-s)}u$ is a solution of class $MR(V, V')$ of

$$\begin{aligned} \dot{v}(t) + (\omega + A(t))v(t) &= 0, & t \in [s, T], \\ v(s) &= x. \end{aligned}$$

Thus, the evolution family associated with \mathbf{a}_ω is simply obtained by rescaling, i.e.,

$$(2.6) \quad U_\omega(t, s) := e^{-\omega(t-s)}U(t, s), \quad (t, s) \in \overline{\Delta}.$$

The earliest well-posedness results for (2.4) were obtained by Kato based on an approximation method based on the theory of product integrals under strong regularity assumptions on the dependence $t \mapsto a(t)$. Kato's approach has been extended to non-autonomous form of class $\text{FORM}([0, T]; V, H)$ in [LE13, EL16]; this has, in turn, allowed for a new proof of Lions' Theorem 2.1 in [SL15]. Let us sketch now his approach here for the sake of self-containedness, since we are going to use it repeatedly in the next sections.

Let $\Lambda = (\lambda_0, \dots, \lambda_{n+1})$ be a partition of $[0, T]$, i.e., $0 = \lambda_0 < \lambda_1 < \dots < \lambda_{n+1} = T$. Let $(a_k)_{k \in \mathbb{N}}$ be a family of sesquilinear forms defined by

$$(2.7) \quad a_k : V \times V \ni (u, v) \mapsto \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} a(r; u, v) dr \in \mathbb{C}, \quad k = 0, 1, \dots, n.$$

All these forms lie in $\text{FORM}([0, T]; V, H)$ with constants M , α , and ω . The associated operators $A_k \in \mathcal{L}(V, V')$ are given by

$$(2.8) \quad A_k : V \ni u \mapsto \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} A(r)u dr \in V', \quad k = 0, 1, \dots, n.$$

The mapping $A(\cdot) : [0, T] \rightarrow \mathcal{L}(V, V')$ is strongly measurable by Pettis' Theorem [ABHN01, Thm. 1.1.1] since $t \mapsto A(t)u$ is weakly measurable and V' is assumed to be separable. On the other hand, $\|A(t)u\|_{V'} \leq M\|u\|_V$ for all $u \in V$ and a.e. $t \in [0, T]$. Thus $[0, T] \ni t \mapsto A(t)u \in V'$ is Bochner integrable for all $u \in V$. Hence the integrals in (2.7) and (2.8) are well defined.

Next, consider the bounded H -elliptic non-autonomous form $\mathfrak{a}_\Lambda := (a_\Lambda(t))_{t \in [0, T]}$ defined by

$$(2.9) \quad a_\Lambda(t; \cdot, \cdot) : V \times V \ni (u, v) \mapsto \begin{cases} a_k(u, v) & \text{if } t \in [\lambda_k, \lambda_{k+1}[\\ a_n(u, v) & \text{if } t = T. \end{cases}$$

Its associated time dependent operator family $\mathcal{A}_\Lambda := (A_\Lambda(t))_{t \in [0, T]} \subset \mathcal{L}(V, V')$ is given by

$$(2.10) \quad A_\Lambda(t) := \begin{cases} A_k & \text{if } t \in [\lambda_k, \lambda_{k+1}[\\ A_n & \text{if } t = T. \end{cases}$$

For each $k = 0, 1, \dots, n$ we denote by $\mathcal{T}_k := \{e^{-rA_k} \mid r \geq 0\}$ the C_0 -semigroup generated by $-A_k$. For each interval $[a, b] \subset [0, T]$ such that

$$(2.11) \quad \lambda_{m-1} \leq a < \lambda_m < \dots < \lambda_{l-1} \leq b < \lambda_l$$

we define the operator families $\mathcal{U}_\Lambda := (U_\Lambda(t, s))_{(t, s) \in \Delta} \subset \mathcal{L}(V')$ by

$$(2.12) \quad U_\Lambda(b, a) = e^{-(b-\lambda_{l-1})A_{l-1}} e^{-(\lambda_{l-1}-\lambda_{l-2})A_{l-2}} \dots e^{-(\lambda_{m-1}-\lambda_{m-2})A_{m-2}} e^{-(\lambda_m-a)A_{m-1}},$$

and for $\lambda_{l-1} \leq a \leq b < \lambda_l$ by

$$(2.13) \quad U_\Lambda(b, a) = e^{-(b-a)A_l}.$$

Remark that \mathcal{U}_Λ defines an evolution family on H (as well as on V' and V), since all semigroups \mathcal{T}_k consist of bounded linear operators on H . Additionally, one sees that the conditions (2.1)–(2.3) are satisfied by the forms \mathfrak{a}_Λ , too: hence for all $x \in H$ also the function $u_\Lambda(\cdot) := U_\Lambda(\cdot, s)x$ is the unique solution of class $MR(s, T; V, V')$ of the problem

$$(2.14) \quad \begin{aligned} \dot{u}_\Lambda(t) + A_\Lambda(t)u_\Lambda(t) &= 0, & t \in [s, T], \\ u_\Lambda(s) &= x. \end{aligned}$$

More precisely, let $x \in H$. On the interval $[\lambda_0, \lambda_1[$, the function u_Λ coincides with the solution of the autonomous problem

$$\begin{aligned} \dot{u}_0(t) + A_0u_0(t) &= 0, & t \in [\lambda_0, \lambda_1[, \\ u_0(\lambda_0) &= x, \end{aligned}$$

which belongs to $MR(\lambda_0, \lambda_1; V, V') \hookrightarrow C([\lambda_0, \lambda_1]; H)$.

Next, for each $k = 1, 2, \dots, n$, the restriction $u_{\Lambda|_{[\lambda_k, \lambda_{k+1}[}}$ coincides with the solution of the autonomous problem

$$\begin{aligned} \dot{u}_k(t) + A_k u_k(t) &= 0, & t \in [\lambda_k, \lambda_{k+1}[, \\ u_k(\lambda_k) &= u_{k-1}(\lambda_k), \end{aligned}$$

which belongs to $MR(\lambda_k, \lambda_{k+1}; V, V')$. We conclude that $U_\Lambda(\cdot, s) \in MR(s, T; V, V')$ and is the unique solution of (2.14).

A similar approximation scheme was introduced in [EL16] in the more general context of inhomogeneous non-autonomous problems; several convergence results could be deduced there, depending on conditions satisfied by the non-autonomous form. In the language of evolution families, we can paraphrase the main result in [EL16] and state the following.

Theorem 2.5. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ and let \mathcal{U} and \mathcal{U}_Λ be the evolution families associated with \mathbf{a} and \mathbf{a}_Λ , respectively. Then*

$$\lim_{|\Lambda| \rightarrow 0} U_\Lambda(t, s) = U(t, s) \quad \text{for all } (t, s) \in \Delta$$

in the strong operator topology of $\mathcal{L}(H)$.

Proof. By Remark 2.4 we can assume that $\omega = 0$ in (2.3). For each $u \in L^2(s, T; V)$, $A_\Lambda(\cdot)u(\cdot)$ converges to $A(\cdot)u(\cdot)$ on $L^2(s, T; V')$ as $|\Lambda| \rightarrow 0$ by [LE13, Lemma 2.3 and Lemma 3.1]. Let $x \in V$. Let $u, u_\Lambda \in MR(V, V')$ solve (2.4) and (2.14), respectively. Set $w_\Lambda := u_\Lambda - u$ and $g_\Lambda := (A - A_\Lambda)u$. Then $w_\Lambda \in MR(V, V')$ and satisfies

$$\begin{aligned} \dot{w}_\Lambda(t) + A_\Lambda(t)w_\Lambda(t) &= g_\Lambda(t), & t \in [s, T], \\ w_\Lambda(s) &= 0. \end{aligned}$$

If $y \in MR(V, V')$ then $\|y(\cdot)\|_H^2 \in W^{1,1}(s, T; V')$ and

$$(2.15) \quad \frac{d}{dt} \|y(\cdot)\|_H^2 = 2 \operatorname{Re} \langle \dot{y}(\cdot), y(\cdot) \rangle$$

by [Sho97, Prop. III.1.2]: accordingly,

$$\begin{aligned} \frac{d}{dt} \|w_\Lambda(t)\|_H^2 &= 2 \operatorname{Re} \langle g_\Lambda(t) - A_\Lambda(t)w_\Lambda(t), w_\Lambda(t) \rangle \\ &= -2 \operatorname{Re} a_\Lambda(t, w_\Lambda(t), w_\Lambda(t)) + 2 \operatorname{Re} \langle g_\Lambda(t), w_\Lambda(t) \rangle \end{aligned}$$

for a.e. $t \in [s, T]$. Integrating this equality between s and t and using (2.3) we obtain

$$\alpha \int_s^t \|w_\Lambda(s)\|_V^2 ds + \|w_\Lambda(s)\|_V^2 \leq \int_s^t \|g_\Lambda(s)\|_V' \|w_\Lambda(s)\|_V ds.$$

This estimate and Young's inequality yield the estimate

$$\alpha \|w_\Lambda\|_{L^2(s, T; V)}^2 \leq \frac{1}{\alpha} \|g_\Lambda\|_{L^2(s, T; V')}^2.$$

The term of the right-hand side of this inequality converges to 0 as $|\Lambda| \rightarrow 0$. It follows that $u_\Lambda \rightarrow u$ in $L^2(s, T; V)$. Again by [LE13, Lemma 2.3 and Lemma 3.1], $A_\Lambda u_\Lambda \rightarrow Au$ in $L^2(s, T; V')$. Letting $|\Lambda| \rightarrow 0$ in

$$\dot{w}_\Lambda = \dot{u}_\Lambda - \dot{u} = f - A_\Lambda u_\Lambda - \dot{u}$$

and recalling the continuous embedding of $MR(V, V')$ into $C([s, T]; H)$ implies the claim. \square

While all strongly continuous semigroups are exponentially bounded, this is not the case for general evolution families, cf. [EN00, § VI.9]. Evolution families associated with non-autonomous forms are rather special, though. Let us show how to apply the product integral method to deduce two known results about long-time behavior of evolution families: the assertion about quasi-contractivity is [Laa18, Prop. 2.1], whereas *strong* stability was proved by similar means in a special case in [ADKF14, Thm. 5.4].

Proposition 2.6. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$. Then the associated evolution family \mathcal{U} is quasi-contractive, i.e.,*

$$\|U(t, s)\|_{\mathcal{L}(H)} \leq e^{\hat{\omega}(t-s)} \quad \text{for all } (t, s) \in \overline{\Delta},$$

where $\hat{\omega}$ is the mean value over $[s, t]$ of some $\omega \in L^1(0, T)$ such that

$$(2.16) \quad \operatorname{Re} a(t; u, u) + \omega(t) \|u\|_H^2 \geq 0 \quad \text{for a.e. } t \in [0, T] \text{ and } u \in V.$$

If in particular $\mathbf{a} \in \text{FORM}([0, \infty[; V, H)$ and $\int_{t_0}^t \omega(s) ds < 0$ for some t_0 and all $t > t_0$, then \mathcal{U} is uniformly exponentially stable.

In view of Proposition 2.6.(1) and Remark 2.4, we will often assume without loss of generality that $\omega \equiv 0$, and thus that an evolution family is contractive up to a scalar perturbation of the family $(A(t))_{t \in [0, T]}$.

Proof. Let $[a, b] \subset [0, T]$, Λ be a partition of $[a, b]$ as in (2.11) and consider the discretized evolution family \mathcal{U}_Λ . Let $\omega_k \in \mathbb{R}$ be defined by

$$(2.17) \quad \omega_k := \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} \omega(r) dr, \quad k = 0, 1, \dots, n.$$

Then by definition of a_k in (2.7), (2.16) implies

$$\operatorname{Re} a_k(u, u) + \omega_k \|u\|_H^2 \geq 0 \quad \text{for all } u \in V \text{ and } k = 0, 1, \dots, n,$$

hence the associated semigroup satisfies

$$(2.18) \quad \|e^{-rA_k}\|_{\mathcal{L}(H)} \leq e^{r\omega_k} \quad \text{for all } k = 0, 1, \dots, n, r \geq 0.$$

Now we obtain from (2.12)

$$(2.19) \quad \|U_\Lambda(a, b)\|_{\mathcal{L}(H)} \leq \prod_{k=0}^n e^{\int_{\lambda_{k-1}}^{\lambda_k} \omega(r) dr} = e^{\int_a^b \omega(r) dr}.$$

The claim now follows from Theorem 2.5 and Fatou's Lemma. \square

We conclude this section by formulating a perturbation lemma that will prove useful will discussing concrete examples.

Lemma 2.7. *Let $\alpha \in (0, 1)$ and let H_α be some normed space such that $V \hookrightarrow H_\alpha \hookrightarrow H$ and such that additionally*

$$\|f\|_{H_\alpha} \leq M_\alpha \|f\|_V^\alpha \|f\|_H^{1-\alpha} \quad \text{for all } f \in V.$$

Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ and \mathbf{b}, \mathbf{c} be family of sesquilinear mappings such that $b(t; \cdot, \cdot) : V \times H_\alpha \rightarrow \mathbb{C}$ and $c(t; \cdot, \cdot) : H_\alpha \times V \rightarrow \mathbb{C}$ are for all $t \in [0, T]$ bounded with bound independent of t . Then $\mathbf{a} + \mathbf{b} + \mathbf{c} \in \text{FORM}([0, T]; V, H)$.

This extends [Mug14, Lemma 6.22] to the case of non-autonomous forms and can be proved likewise.

3. INVARIANCE PROPERTIES

Let us discuss invariance of a given subset C of H under \mathcal{U} , i.e., whether $u(s) \in C$ implies that the solution $u(t)$ of (2.4) lies in C for any $(t, s) \in \Delta$. The following criterion is known: it combines [SL15, Thm. 4.1] with an extension to non-accretive forms [MVV05, Thm. 2.1] of Ouhabaz' classical invariance criterion [Ouh96, Thm. 2.1].

Proposition 3.1. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$. Let C be a closed convex subset of H and denote by P the projector of H onto C . Consider the following assertions:*

- (i) C is invariant under the semigroup \mathcal{T}_t associated with $a(t)$ for a.e. $t \in [0, T]$;
- (ii) C is invariant under the evolution family \mathcal{U} .
- (iii) $Pu \in V$ and $\text{Re } a(t; Pu, u - Pu) \geq 0$ for all $u \in V$ and a.e. $t \geq 0$;

Then (i) is equivalent to (iii) and both imply (ii).

The implication (iii) \Rightarrow (ii) has been proved in [ADO14, Thm. 2.2] in the more general case of inhomogeneous equations. Special instances of the same assertion have been obtained in [Tho03, § 3.5.5]. The implication (i) \Rightarrow (ii) allows us to deduce invariance properties for \mathcal{U} even if P is not explicitly known, e.g., when \mathcal{T}_t is known to preserve convexity for a.e. t [BB12].

Remark 3.2. *We have remarked that the mapping $[0, T] \ni t \mapsto \mathcal{A}(t) \in \mathcal{L}(V, V')$ is strongly measurable under our standing assumptions. It is unclear whether the same assumptions also imply strong measurability of its resolvents, apart from the somewhat trivial case where the operators commute and are therefore simultaneously diagonalizable.*

If however strong measurability of $[0, T] \ni t \mapsto R_t(\lambda) \in \mathcal{L}(V')$ for each λ is imposed, then in view of the representation of the holomorphic semigroups \mathcal{T}_t as inverse Laplace transforms of $R_t(\cdot)$ [ABHN01, Thm. 3.7.11] we can deduce that $[0, T] \ni t \mapsto T_t(r) \in \mathcal{L}(V')$ is strongly measurable, too, for each r . Given a partition Λ of $[0, T]$ we can define the ‘‘averaged operator family’’ $\mathcal{T}_\Lambda := (T_\Lambda(t))_{t \in [0, T]}$

$$(3.1) \quad T_\Lambda(t)x := e^{-rA_k}x := \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} e^{-rA(s)}x \, ds \quad \text{if } t \in (\lambda_k, \lambda_{k+1}), \quad x \in V',$$

and show that, again, their product integrals converge towards the evolution family \mathcal{U} as the partition becomes finer. In this case, if a closed (but not necessarily convex!) subset C of H is invariant under all semigroups \mathcal{T}_t , then we can apply the strategy in Remark 3.2 and introduce \mathcal{U}_Λ based on the semigroups in (3.1); accordingly, C is also invariant under $U_\Lambda(b, a)$ for all $(b, a) \in \Delta$ and all partitions Λ , hence by Theorem 2.5 also under \mathcal{U} .

It is known that under (rather strong) conditions on the dependence on t of the resolvent operators of $A(t)$, \mathcal{U} can be showed to be immediately differentiable and even holomorphic (i.e., continuously differentiable, resp. holomorphic from Δ to $\mathcal{L}(H)$), and to map H into $D(A(t))$, see [Fat83, Thm. 7.2.5 and Thm. 7.4.1].

For our purposes, a particularly interesting instance of closed convex sets are order intervals in Hilbert lattices: we hence assume in the following H to be a Hilbert lattice. It is known that each separable Hilbert lattice is isometrically lattice isomorphic to a Lebesgue space $L^2(X)$ for some σ -finite measure space (X, Σ, μ) , see e.g. [MN91, Cor. 2.7.5]. Accordingly, we can consider the set $H_{\mathbb{R}} := L^2(X; \mathbb{R})$ of real-valued functions. Let $a, b \in \mathbb{R} \cup \{\pm\infty\}$: we introduce the (bounded or unbounded) *order intervals*

$$[a, b]_H := \{f \in H_{\mathbb{R}} : a \leq f(x) \leq b \text{ for a.e. } x \in X\} :$$

or, more generally,

$$[\phi, \psi]_H := \{f \in H_{\mathbb{R}} : \phi(x) \leq f(x) \leq \psi \text{ for a.e. } x \in X\},$$

where $\phi, \psi : X \rightarrow [-\infty, \infty]$ are measurable functions: they are closed convex subsets of H . Many qualitative properties of solutions to evolution equations can be described by means of invariance of order intervals under the flow that governs the associated Cauchy problems.

Definition 3.3. *Let (X, Σ, μ) be a σ -finite measure space. An evolution family \mathcal{U} on the Hilbert lattice $L^2(X)$ is called*

- (a) *real if $U(t, s)H_{\mathbb{R}} \subset H_{\mathbb{R}}$ for all $(t, s) \in \Delta$;*
- (b) *positive if it is real and $U(t, s)[0, \infty[_H \subset [0, \infty[_H$ for all $(t, s) \in \Delta$;*
- (c) *positivity improving if \mathcal{U} is positive and additionally $U(t, s)f(x) > 0$ for all $(t, s) \in \Delta$ and a.e. $x \in X$ provided $f \in [0, \infty[_H$ and $f \neq 0$.*
- (d) *L^∞ -contractive if $U(t, s)[-1, 1]_H \subset [-1, 1]_H$ for all $(t, s) \in \Delta$;*
- (e) *completely contractive if it is both L^1 -contractive and L^∞ -contractive;*
- (f) *completely quasi-contractive if there is some constant $\tilde{\omega}$ such that the rescaled evolution family $\mathcal{U}_{\tilde{\omega}}$ defined by*

$$(3.2) \quad U_{\tilde{\omega}}(t, s) := e^{-\tilde{\omega}(t-s)}U(t, s), \quad (t, s) \in \Delta,$$

is completely contractive;

- (g) *sub-Markovian if it is positive and L^∞ -contractive and Markovian if additionally $\|U(t, s)\|_{\mathcal{L}(L^\infty)} = 1$;*
- (h) *sub-stochastic if it is positive and L^1 -contractive and stochastic if additionally and $\|U(t, s)f\|_{L^1(X)} = \|f\|_{L^1(X)}$ for all $0 \leq f \in L^2(X) \cap L^1(X)$ and all $(t, s) \in \Delta$.*

Remark 3.4. (i) *Let \mathcal{U} be a completely quasi-contractive evolution family on $L^2(X)$. Then by Riesz–Thorin the rescaled evolution family $\mathcal{U}_{\tilde{\omega}}$ is L^p -contractive for all $p \in [1, \infty]$. Hence each $U(t, s)$ can be extended from $L^p(X) \cap L^2(X)$ to a quasi-contractive operator $U_p(t, s)$ on $L^p(X)$ for all $p \in [1, \infty]$. The extrapolated family $\mathcal{U}_p := \{U_p(t, s) \mid (t, s) \in \Delta\}$ is consistent, i.e., for all $p \in [1, \infty]$*

$$U_p(t, s)f = U_2(t, s)f \quad \text{for all } (t, s) \in \Delta \text{ and all } f \in L^p(X) \cap L^2(X).$$

It is clear that the extended family \mathcal{U}_p satisfies conditions (i)–(ii) in Theorem 2.1. Moreover, by the interpolation inequality (Hölder inequality) we obtain that \mathcal{U}_p satisfies (iii) in Theorem 2.1 for all $p \in]1, \infty[$. Using a similar argument as in [Voi92, Prop. 4] we conclude that \mathcal{U}_p is a strongly continuous evolution family on $L^p(X)$ for all $p \in [1, \infty[$.

- (ii) *We also deduce from Remark 2.3 and [Are06, Prop. 4.4.3] that if V is compactly embedded in H , then $U_p(t, s)$ is a compact operators for all $p \in]1, \infty[$ and a.e. $(t, s) \in \Delta$.*

For future reference let us note explicitly the following consequence of Proposition 3.1.

Proposition 3.5. *The evolution family \mathcal{U} associated with $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ is*

- (1) *positive provided $(\text{Re } v)^+ \in V$, $\text{Re } a(t; \text{Re } v, \text{Im } v) \in \mathbb{R}$, and $\text{Re } a(t; (\text{Re } v)^+, (\text{Re } v)^-) \leq 0$ for all $v \in V$ and a.e. $t \in [0, T]$.*
- (2) *L^∞ -contractive, provided $(1 \wedge |v|) \text{sgn } v \in V$ and $\text{Re } a(t; (1 \wedge |v|) \text{sgn } v, (|v| - 1)^+ \text{sgn } v) \geq 0$ for all $v \in V$ and a.e. $t \in [0, T]$.*

Likewise, owing to [Ein08, Prop. 4.4.1] we obtain the following.

Proposition 3.6. *Let $\psi : X \rightarrow [0, \infty)$ and $\phi : X \rightarrow (-\infty, 0]$ be measurable. The evolution family \mathcal{U} associated with $\mathfrak{a} \in \text{FORM}([0, T]; V, H)$ leaves the order interval*

- (i) $] -\infty, \psi]_H$ invariant provided $\min\{\text{Re } v, \psi\} \in V$, $\text{Re } a(t; \min\{\text{Re } v, \psi\}, (\text{Re } v - \psi)^+) \geq 0$ for all $v \in V$ and a.e. $t \in [0, T]$.
- (ii) $[\phi, \psi]_H$ invariant provided $\max\{\phi, \min\{\text{Re } v, \psi\}\} \in V$,

$$\text{Re } a(t; \max\{\phi, \min\{\text{Re } v, \psi\}\}, (\text{Re } v - \psi)^+ - (\text{Re } v - \phi)^+) \geq 0$$

for all $v \in V$ and a.e. $t \in [0, T]$.

Let us state a further consequence of Proposition 3.1 concerning irreducibility of evolution families on $L^2(X)$ on a given σ -finite measure space (X, Σ, μ) . We denote by $\mathbf{1}_\Xi$ the characteristic function of any given $\Xi \in \Sigma$.

Proposition 3.7. *Let $\mathfrak{a} \in \text{FORM}([0, T]; V, H)$ and assume that $(\text{Re } v)^+ \in V$, $\text{Re } a(t; \text{Re } v, \text{Im } v) \in \mathbb{R}$, and $\text{Re } a(t; (\text{Re } v)^+, (\text{Re } v)^-) \leq 0$ for all $v \in V$ and a.e. $t \in [0, T]$. If additionally*

$$\mathbf{1}_\Xi V \subset V \quad \text{implies} \quad \mu(\Xi) = 0 \quad \text{or} \quad \mu(X \setminus \Xi) = 0,$$

for all $\Xi \in \Sigma$, then \mathcal{U} is positivity improving.

Proof. Let Λ be a partition of $[a, b]$ as in (2.11) and consider the discretized C_0 -semigroups $\mathcal{T}_k, k = 0, 1, \dots, n$. It follows from [Are06, Thm. 10.1.5 and Thm. 10.1.2] that \mathcal{T}_k is positivity improving for all $k = 0, 1, \dots, n$. The assertion now follows from Theorem 2.5. \square

This enables us to provide sufficient conditions for the evolution family to converge towards a rank-one projector, thus extending to the non-autonomous setting one of the main results of the classical Perron–Frobenius theory for semigroups.

Corollary 3.8. *Under the assumptions of Proposition 3.7, let additionally $\mathfrak{a} \in \text{FORM}([0, \infty[; V, H)$ and V be compactly embedded in H . Assume that for all $t \in [0, \infty[$*

- the spectral bound of $A(t)$ is 0,
- for a.e. $t \in]0, \infty[$ $\text{Ker}(A(t))$ is spanned by the same strictly positive vector u , and
- for a.e. $t \in]0, \infty[$ $\text{Ker}(A^*(t))$ is spanned by the same strictly positive vector ϕ , with $(\varphi|u) = 1$.

Then

$$\lim_{t \rightarrow \infty} U(t, s) = P := \phi \otimes u \quad \text{strongly in } H \text{ for all } s \geq 0.$$

Proof. Under our assumptions the semigroups \mathcal{T}_t are for a.e. $t \in]0, \infty[$ holomorphic, positivity improving, and compact: hence by classical Perron–Frobenius theory there is a spectral gap of size $\tilde{s}(A(t)) > 0$ between their common dominant eigenvalue 0 and the bottom of the strictly positive part of the spectrum of their generators $A(t)$: accordingly, the part $\tilde{A}(t)$ in the Hilbert space $\text{Ker}(P)$ is associated with a form $\tilde{\mathfrak{a}} \in \text{FORM}(0, \infty; V \cap \text{Ker}(P), \text{Ker}(P))$ that satisfies

$$\tilde{\mathfrak{a}}(t; u, u) \geq \varepsilon(t) \|u\|^2 \quad \text{for a.e. } t \in]0, \infty[\text{ and all } u \in V \cap \text{Ker}(P)$$

and any $\varepsilon \in L^1(0, \infty)$ s.t.

$$0 < \varepsilon(t) \leq \tilde{s}(A(t)) := \inf \{ \text{Re } \lambda > 0 : \lambda \in \sigma(A(t)) \} \quad \text{for a.e. } t \in]0, \infty[.$$

Given a compact interval $[a, b]$ and a partition $\Lambda = (\lambda_0, \dots, \lambda_{n+1})$ of $[a, b]$, we can hence define in the usual way the averaged forms $\tilde{\mathfrak{a}}_k$, which satisfy

$$\text{Re } \tilde{\mathfrak{a}}_k(u, u) \geq \varepsilon_k \|u\|^2, \quad u \in V \cap \text{Ker}(P),$$

for

$$\varepsilon_k := \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} \varepsilon(r) dr.$$

Therefore, the associated semigroup $(e^{-r\tilde{A}_k})_{r \geq 0}$ is uniformly exponentially stable: more precisely

$$\|e^{-r\tilde{A}_k}\| = \|e^{-rA_k}(I - P)\| \leq e^{-\varepsilon_k r} \quad \text{for all } r \geq 0.$$

Observe now that

$$\begin{aligned} U_\Lambda(b, a) - P &= U_\Lambda(b, a)P - P + U_\Lambda(b, a)(I - P) \\ &= e^{-(\lambda_{n+1} - \lambda_n)A_{n+1}} \dots e^{-(\lambda_1 - \lambda_0)A_1} P - P + U_\Lambda(b, a)(I - P)^n \\ &= U_\Lambda(b, a)(I - P)^n. \end{aligned}$$

Accordingly,

$$\begin{aligned} \|U_\Lambda(b, a) - P\| &\leq \|e^{-(\lambda_{n+1} - \lambda_n)A_{n+1}}(I - P)\| \dots \|e^{-(\lambda_1 - \lambda_0)A_1}(I - P)\| \\ &\leq e^{-(\lambda_{n+1} - \lambda_n)\varepsilon_n} \dots e^{-(\lambda_1 - \lambda_0)\varepsilon_1} = e^{\int_a^b \varepsilon(r) dr} \end{aligned}$$

and for all $x \in H$ by Theorem 2.5

$$\|U(b, a)x - Px\| = \lim_{|\Lambda| \rightarrow 0} \|U_\Lambda(b, a)x - Px\| \leq e^{-\int_a^b \varepsilon(r) dr} \|x\|.$$

Now the claim follows, since $\varepsilon(r) > 0$ for a.e. r . □

In the following sections we will often need to discuss complete contractivity. In order to find sufficient conditions therefor, observe that \mathcal{U} is L^1 -contractive if and only if $U(t, s)^*$ is L^∞ -contractive for all $(t, s) \in \Delta$. How to prove L^∞ -contractivity of all $U(t, s)^*$? Consider the non-autonomous adjoint form $\mathfrak{a}^* : [0, T] \times V \times V \rightarrow \mathbb{C}$ of \mathfrak{a} defined by $\mathfrak{a}^*(t; u, v) := \overline{a(t; v, u)}$ for all $t \in [0, T]$ and $u, v \in V$. While $\mathfrak{a}^* \in \text{FORM}([0, T]; V, H)$, too, and hence \mathfrak{a}^* is associated with an evolution family $(U_*(t, s))_{(t, s) \in \overline{\Delta}}$, one has in general $U_*(t, s) \neq U(t, s)^*$. However, it was observed in [Dan00, Thm. 2.6] that the *returned adjoint form* $\overleftarrow{\mathfrak{a}}^* : [0, T] \times V \times V \rightarrow \mathbb{C}$ of \mathfrak{a} defined by

$$\overleftarrow{\mathfrak{a}}^*(t; u, v) := \mathfrak{a}^*(T - t; v, u), \quad t \in [0, T], \quad u, v \in V,$$

which clearly belongs to $\text{FORM}([0, T]; V, H)$, too, is associated with an evolution family $\overleftarrow{\mathcal{U}}^*$ that satisfies

$$(3.3) \quad [\overleftarrow{\mathcal{U}}^*(t, s)]^* f = U(T - s, T - t)f \quad \text{for all } f \in H \text{ and } (t, s) \in \Delta.$$

In particular, \mathcal{U} is L^1 -contractive if and only if $\overleftarrow{\mathcal{U}}^*$ is L^∞ -contractive; \mathcal{U} is completely contractive if so is $\overleftarrow{\mathcal{U}}^*$; and by Proposition 3.5 we conclude the following.

Proposition 3.9. *The evolution family \mathcal{U} associated with $\mathfrak{a} \in \text{FORM}([0, T]; V, H)$ is*

- (1) *L^1 -contractive provided $(1 \wedge |u|) \text{sgn } u \in V$ and $\text{Re } a(t; (|v| - 1)^+ \text{sgn } v, (1 \wedge |v|) \text{sgn } v) \geq 0$ for all $v \in V$ and a.e. $t \in [0, T]$;*
- (2) *completely contractive provided $(1 \wedge |u|) \text{sgn } u \in V$ and $\text{Re } a(t; (|v| - 1)^+ \text{sgn } v, (1 \wedge |v|) \text{sgn } v) \geq 0$, $\text{Re } a(t; (1 \wedge |v|) \text{sgn } v, (|v| - 1)^+ \text{sgn } v) \geq 0$ for all $v \in V$ and a.e. $t \in [0, T]$.*

Recall that a C_0 -semigroup $\mathcal{S} = (S(r))_{r \geq 0}$ on $L^2(X)$ is said to be L^p -quasi-contractive for some $p \in [1, \infty]$ if there exists a constant $\hat{\omega}_p \in \mathbb{R}$ such that

$$(3.4) \quad \|S(r)f\|_{L^p(X)} \leq e^{r\hat{\omega}_p} \|f\|_{L^p(X)} \quad \text{for all } r \geq 0 \text{ and } f \in L^2(X) \cap L^p(X).$$

In this case, \mathcal{S} extends by continuity to a bounded linear operator on $L^p(X)$ and, by the Riesz–Thorin Theorem, on $L^q(X)$ for all q between 2 and p .

We can now give a sufficient condition for L^p -quasi-contractivity of the evolution family \mathcal{U} that governs the Cauchy problem (2.4).

Theorem 3.10. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ and let $p \in]1, \infty[$ be given. Assume that there exists a function $\hat{\omega}_p \in L^\infty(0, T)$ such that $(\mathcal{T}_t)_{t \in [0, T]}$ satisfies*

$$(3.5) \quad \|e^{-rA(t)}f\|_{L^p(X)} \leq e^{r\hat{\omega}_p(t)} \|f\|_{L^p(X)} \quad \text{for all } f \in L^2(X) \cap L^p(X), r \geq 0, \text{ and a.e. } t \in [0, T].$$

Then the evolution family \mathcal{U} associated with \mathbf{a} extrapolates to a consistent evolution family on $L^p(X)$ and

$$(3.6) \quad \|U(t, s)f\|_{L^p(X)} \leq e^{\int_s^t \hat{\omega}_p(r) dr} \|f\|_{L^p(X)} \quad \text{for all } f \in L^2(X) \cap L^p(X) \text{ and } (t, s) \in \Delta.$$

Proof. The case where $\hat{\omega}_p \equiv 0$ follows directly from Proposition 3.1; the general case is slightly more delicate. First, applying [Nit12, Thm. 4.4] to the non-autonomous form $a(t, \cdot, \cdot) + \hat{\omega}_p(t)(\cdot | \cdot)_{L^2}$, we see that the assumption (3.5) is equivalent to the following condition: $P_{B^p}V \subset V$ and for a.e. $t \in [0, T]$

$$(3.7) \quad \text{Re } a(t, u, |u|^{p-2}u) + \hat{\omega}_p(t) \|u\|_{L^p}^p \geq 0 \quad \text{for all } u \in V \text{ s.t. } |u|^{p-2}u \in V.$$

(Here B^p denotes the L^p -unit ball and P_{B^p} is the projector of $L^2(X)$ onto B^p .)

Let now $[s, t] \subset [0, T]$ and let $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_n)$ be a partition of $[s, t]$ and let $\mathbf{a}_k : V \times V \rightarrow \mathbb{C}$, $k = 0, 1, \dots, n$, be the family of bounded H -elliptic forms given by (2.7) and $(e^{-rA_k})_{r \geq 0}$ be the associated C_0 -semigroup. Furthermore, define the finite real sequence $\hat{\omega}_{k,p}$, $k = 0, 1, \dots, n$, as follows

$$(3.8) \quad \hat{\omega}_{k,p} := \frac{1}{\lambda_{k+1} - \lambda_k} \int_{\lambda_k}^{\lambda_{k+1}} \hat{\omega}_p(r) dr \quad k = 0, 1, \dots, n.$$

Then (3.7) implies that for all $k = 0, 1, \dots, n$

$$(3.9) \quad \text{Re } a_k(u, |u|^{p-2}u) + \hat{\omega}_{k,p} \|u\|_{L^p}^p \geq 0, \quad \text{for all } u \in V \text{ s.t. } |u|^{p-2}u \in V.$$

Again applying [Nit12, Thm. 4.4] to the form $a_k + \hat{\omega}_{k,p}(\cdot | \cdot)_{L^2}$ we obtain

$$(3.10) \quad \|e^{-rA_k}\|_{L^p(X)} \leq e^{r\hat{\omega}_{k,p}} \quad \text{for all } s \geq 0 \text{ and } k = 0, 1, \dots, n.$$

Thus, using (2.12)-(2.13) we find

$$(3.11) \quad \|U_\Lambda(t, s)\|_{L^p(X)} \leq e^{\int_s^t \hat{\omega}_p(r) dr} \quad \text{for all } (t, s) \in \Delta \text{ and each partition } \Lambda \text{ of } [s, t].$$

Thus the desired estimate (3.6) follows from Theorem 2.5 and Fatou's Lemma. \square

Let us now discuss stochasticity, another feature that cannot be easily interpreted as an invariance property.

Proposition 3.11. *The evolution family \mathcal{U} associated with $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ is stochastic provided $(\text{Re } u)^+ \in V$, $\text{Re } a(t, \text{Re } u, \text{Im } u) \in \mathbb{R}$, $\text{Re } a(t; (\text{Re } u)^+, (\text{Re } u)^-) \leq 0$, $\mathbf{1} \in V$, and $a(t; \text{Re } u, \mathbf{1}) = 0$ for all $v \in V$ and a.e. $t \in [0, T]$.*

Proof. We will again use the approximation techniques described in Section 2. Let $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_n)$ be a partition of $[0, T]$, $\mathbf{a}_k : V \times V \rightarrow \mathbb{C}$ be given by (2.7) and \mathcal{T}_k be the C_0 -semigroup associated with \mathbf{a}_k in H for $k = 0, 1, \dots, n$: under our assumptions it is easy to check that \mathcal{T}_k is positive, $\mathbf{1} \in V$, and $\mathbf{a}_k(\operatorname{Re} v, \mathbf{1}) = 0$ for all $v \in V$ and $k = 0, 1, \dots, n$ which is equivalent to the fact the all semigroups \mathcal{T}_k are stochastic. Now the claim follows from Theorem 2.5. \square

Our last result in this section is devoted to the issue of domination of evolution families.

Proposition 3.12. *Let $\mathbf{a} \in \operatorname{FORM}([0, T]; V, H)$ and denote as usual by \mathcal{U} the associated evolution family. Let furthermore W be a separable Hilbert space that is densely and continuously embedded in H and $\mathbf{b} \in \operatorname{FORM}([0, T]; W, H)$: we denote by \mathcal{V} the associated evolution family. Assume that*

- $\operatorname{Re} u \in V$ and $(\operatorname{Re} u)^+ \in V$ for all $u \in V$;
- V is a generalized ideal of W , i.e.,
 - $u \in W$ implies $|u| \in W$ and
 - $u_1 \in V$ and $u_2 \in W$ are such that $|u_2| \leq |u_1|$, then $u_2 \operatorname{sgn} u_1 \in V$;
- $\operatorname{Re} \mathbf{a}(t; \operatorname{Re} u, \operatorname{Im} u) \in \mathbb{R}$ for all $u \in V$;
- $\operatorname{Re} \mathbf{a}(t; (\operatorname{Re} u)^+, (\operatorname{Re} u)^-) \leq 0$ for all $u \in V$;
- $\operatorname{Re} \mathbf{a}(t; u, v) \geq \mathbf{b}(t; |u|, |v|)$ for all $u, v \in V$ s.t. $u\bar{v} \geq 0$.

Then \mathcal{U} is dominated by \mathcal{V} , i.e.,

$$(3.12) \quad |U(t, s)f| \leq V(t, s)|f| \quad \text{for all } (t, s) \in \Delta \text{ and } f \in H.$$

Proof. Let Λ be a partition of $[0, T]$. Define the piecewise constant $\mathbf{b}_\Lambda \in \operatorname{FORM}([0, T]; W, H)$ via formulae which are analogous to (2.7) and (2.10) and let \mathcal{V}_Λ be the associated evolution family.

By [MVV05, Thm. 4.1], domination holds if and only if V is a generalized ideal of W and $\operatorname{Re} \mathbf{a}(t; u, v) \geq \mathbf{b}(t; |u|, |v|)$ for all $u, v \in V$ such that $u\bar{v} \geq 0$; in this case, the latter property holds also for the averaged forms \mathbf{b}_Λ and hence, again by [MVV05, Thm. 4.1], also the associated semigroups dominate \mathcal{T}_t . It follows from (2.12) that \mathcal{U}_Λ is dominated by \mathcal{V}_Λ : letting $|\Lambda| \rightarrow 0$ and in view of Theorem 2.5 we conclude that \mathcal{U} is dominated by \mathcal{V} . \square

Example 3.13. *To illustrate our results obtained so far, let us briefly dip into the topic of diffusion equations on dynamic graphs that appear in different applications, like flocking models [VZ12].*

Consider a (finite or infinite) simple graph \mathbf{G} with vertex set \mathbf{V} and edge set \mathbf{E} , with V vertices and E edges (i.e., $V = |\mathbf{V}|$ and $E = |\mathbf{E}|$). Fix an orientation of \mathbf{G} and introduce the $V \times E$ (signed) incidence matrix $\mathcal{I} = (\iota_{\mathbf{v}\mathbf{e}})$ of \mathbf{G} by

$$\iota_{\mathbf{v}\mathbf{e}} := \begin{cases} -1 & \text{if } \mathbf{v} \text{ is initial endpoint of } \mathbf{e}, \\ +1 & \text{if } \mathbf{v} \text{ is terminal endpoint of } \mathbf{e}, \\ 0 & \text{otherwise.} \end{cases}$$

Let $m \in \ell^\infty(\mathbf{E})$ be a family of edge weights and consider the (weighted) Laplacian $\mathcal{L} := \mathcal{I}M\mathcal{I}^T$ on $\ell^2(\mathbf{V})$, where $0 \leq M := \operatorname{diag}(m(\mathbf{e}))_{\mathbf{e} \in \mathbf{E}}$. (\mathcal{L} can be shown to be independent of the orientation.)

We assume that \mathbf{G} is uniformly locally finite, i.e., there is $M < \infty$ such that $\sum_{\mathbf{e} \in \mathbf{E}} |\iota_{\mathbf{v}\mathbf{e}}| \leq M$ for all $\mathbf{v} \in \mathbf{V}$: in this case \mathcal{I} is a bounded linear operator from $\ell^2(\mathbf{E})$ to $\ell^2(\mathbf{V})$ [Mug14, Lemma. 4.3], hence \mathcal{L} is a positive semi-definite, bounded self-adjoint operator on $\ell^2(\mathbf{V})$: we can thus take $V = H = V' = \ell^2(\mathbf{V})$. It is well-known that the semigroup generated by $-\mathcal{L}$ is sub-Markovian, see e.g. [Mug14, § 6.4.1]; if the graph is finite, then it is Markovian and stochastic, too.

Let us now regard \mathbf{G} as a reference graph (one may e.g. think of a complete graph, or else of a lattice graph \mathbb{Z}^d) and consider a family $(\mathbf{G}(t))_{t \in [0, T]}$ of modifications of \mathbf{G} – in other words, a graph-valued dynamical system, or dynamic graph [Šil08]. We describe the dependence of $\mathbf{G}(t)$ on t by introducing a measurable function $[0, T] \ni t \mapsto m(t) \in \ell^\infty(\mathbf{E})$: this allows e.g. for sudden switching of edges is allowed (as in the case of adjacency driven by a Poisson process). In particular, we consider the non-autonomous form \mathbf{a} defined by

$$a(t; u, v) := (\text{diag}(m(t, \mathbf{e})) \mathcal{I}^T u | \mathcal{I}^T v)_{\ell^2(\mathbf{E})}, \quad t \in [0, T], \quad u, v \in \ell^2(\mathbf{V}).$$

It is easy to see that $\mathbf{a} \in \text{FORM}([0, T]; \ell^2(\mathbf{V}), \ell^2(\mathbf{V}))$ and the associated operators are the Laplacians $(\mathcal{L}_{\mathbf{G}(t)})_{t \in [0, T]}$. (We are not assuming boundedness from below on m : this is made unnecessary by the boundedness of the operator $\mathcal{L}_{\mathbf{G}(t)}$ for all t ; in fact, even negative weights and hence signed graphs are allowed.) We deduce by Proposition 3.1 that the non-autonomous Cauchy problem

$$\begin{aligned} \dot{u}(t, \mathbf{v}) + \mathcal{L}_{\mathbf{G}(t)} u(t, \mathbf{v}) &= 0, \quad t \geq s, \quad \mathbf{v} \in \mathbf{V}, \\ u(s, \mathbf{v}) &= x_{\mathbf{v}}, \quad \mathbf{v} \in \mathbf{V} \end{aligned}$$

is governed by an evolution family on $\ell^2(\mathbf{V})$; in fact, for all $x \in \ell^2(\mathbf{V})$ the above equation enjoys backward well-posedness, too, and the unique solution u is of class $H^1(\mathbb{R}; \ell^2(\mathbf{V}))$: the corresponding evolution family $(U(t, s))_{(t, s) \in \mathbb{R}^2}$ can be defined via product integrals. As observed in [AD17, Example 7.3] $U(t, s)$ is sub-Markovian for all t, s ; in particular, \mathcal{U} extrapolates to a consistent family of contractive evolution families on $\ell^p(\mathbf{V})$ for all $p \in [1, \infty]$. The evolution family is also positivity improving, and additionally stochastic if \mathbf{G} is finite. Furthermore, [Laa18, Thm. 2.6] yields that the evolution family is immediately norm-continuous if $[0, T] \ni t \mapsto m(t) \in \ell^\infty(\mathbf{E})$ is Hölder continuous with exponent $\alpha > 1/2$; by [Fat83, Thm. 7.4.1] it is even holomorphic if additionally $[0, T] \ni t \mapsto m(t) \in \ell^\infty(\mathbf{E})$ extends to a holomorphic function on an open convex neighborhood in \mathbb{C} of $[0, T]$.

To conclude, let us study Laplacians on subgraphs \mathbf{G}_t induced by subsets \mathbf{V}_t of \mathbf{V} as in [Chu97, Chapt. 8] in the unweighted case ($m(t, \mathbf{e}) \in \{0, 1\}$). Even in the autonomous case, Laplacian on (non-trivial) subgraphs of \mathbf{G} generate semigroup that neither are dominated by, nor dominate $(e^{-r\mathcal{L}_{\mathbf{G}}})_{r \geq 0}$: this can e.g. be seen by applying [Ouh05, Cor. 2.22]. Things change, however, if Dirichlet boundary conditions are imposed, e.g., if \mathcal{L} on \mathbf{G} is restricted to

$$D_t := \{f \in \ell^2(\mathbf{V}) : f|_{\mathbf{V} \setminus \mathbf{V}_t} \equiv 0\}, \quad t \in [0, T].$$

Because D_t is for all t a generalized ideal of $V = \ell^2(\mathbf{V})$, the associated Laplacian $\mathcal{L}|_{D_t}$ generates for all t a semigroup $(e^{-r\mathcal{L}|_{D_t}})_{r \geq 0}$ each of which is – again by [Ouh05, Cor. 2.22] – indeed dominated by $(e^{-r\mathcal{L}})_{r \geq 0}$. Therefore, by Proposition 3.1 and Proposition 3.12 the evolution family $(U(t, s))_{(t, s) \in \Delta}$ satisfies

$$(3.13) \quad |U(t, s)f| \leq e^{-(t-s)\mathcal{L}}|f| \quad \text{for all } (t, s) \in \Delta \text{ and } f \in H.$$

We have seen in the introduction that if $A(t) \equiv A$, then the evolution family that governs the non-autonomous problem is nothing but

$$U(t, s) = e^{(t-s)A}, \quad (t, s) \in \Delta,$$

hence \mathcal{U} satisfies Gaussian bounds if and only if so does $(e^{rA})_{r \geq 0}$. We are already in the position to show a less trivial instance of Gaussian-type estimate.

Gaussian-type kernel bounds on $(e^{-t\mathcal{L}_G})_{t \geq 0}$ have been proved in [Del99] for certain classes of \mathbf{G} . Thus, if $(\mathbf{G}_t)_{t \in [0, T]}$ is a family of subgraphs of a reference graph \mathbf{G} with measurable $t \mapsto m(t, \mathbf{e})$ for all $\mathbf{e} \in \mathbf{E}$, and if \mathcal{U} is the evolution family associated with the corresponding Laplacians $-\mathcal{L}_{|D_t}$, then (3.13) yields a Gaussian-type kernel estimate. If we e.g. take \mathbf{G} to be \mathbb{Z} , then

$$0 \leq \Gamma(t, s; n_1, n_2) \leq G(t - s; n_1, n_2), \quad (t, s) \in \Delta, \quad n_1, n_2 \in \mathbb{Z},$$

where

$$G(r; n_1, n_2) := \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos((n_1 - n_2)q) e^{-2r(1 - \cos q)} dq$$

is the heat kernel on \mathbb{Z} explicitly computed e.g. in [Dav07, Exa. 12.3.3].

4. ULTRA CONTRACTIVITY

In this and the next section we are going to restrict to the case of $H = L^2(X)$, where X an σ -finite measurable space. Recall that a C_0 -semigroup \mathcal{S} on $L^2(X)$ is said to be *ultracontractive* if there exist constants $c_0, n > 0$, and $\tilde{\omega} \in \mathbb{R}$ such that

$$(4.1) \quad \|\mathcal{S}(r)f\|_{L^\infty(X)} \leq c_0 r^{-\frac{n}{2}} e^{r\tilde{\omega}} \|f\|_{L^1(X)} \quad \text{for all } r \geq 0 \text{ and all } f \in L^2(X) \cap L^1(X).$$

In this section we are going to develop a theory of ultracontractive evolution families.

Definition 4.1. *We call an evolution family \mathcal{U} on $L^2(X)$ ultracontractive if there exist constants $c_0, n > 0$ and $\tilde{\omega} \in \mathbb{R}$ such that*

$$(4.2) \quad \|U(t, s)f\|_{L^\infty(X)} \leq c_0 (t - s)^{-\frac{n}{2}} e^{(t-s)\tilde{\omega}} \|f\|_{L^1(X)} \quad \text{for all } (t, s) \in \Delta \text{ and } f \in L^2(X) \cap L^1(X).$$

By a direct consequence of the Kantorovitch–Vulikh Theorem, see e.g. [AB94, Theorem 1.3], any ultracontractive evolution family \mathcal{U} is given by an integral kernel: more precisely, there exists a family $(\Gamma(t, s))_{(t, s) \in \Delta} \subset L^\infty(\Omega \times \Omega)$ such that

$$U(t, s)f(x) = \int_{\Omega} \Gamma(t, s; x, y) f(y) dy \quad \text{for all } (t, s) \in \Delta \text{ and } f \in L^2(X) \cap L^1(X) \text{ and a.e. } x \in \Omega,$$

with

$$\|U(t, s)\|_{\mathcal{L}(L^1(X), L^\infty(X))} = \|\Gamma(t, s)\|_{L^\infty(\Omega \times \Omega)}, \quad \text{for all } (t, s) \in \Delta.$$

We observe in passing that if \mathcal{U} is ultracontractive and Ω has finite measure, then $\Gamma(t, s) \in L^2(\Omega \times \Omega)$ for all $(t, s) \in \Delta$, hence \mathcal{U} consists of Hilbert–Schmidt operators; in fact even of trace class operators, by the operator equation that defines evolution families ((ii) in Theorem 2.1). Different sufficient conditions for the trace class property of \mathcal{U} have been presented in [Laa18, § 3].

It is well-known that ultracontractivity of semigroups can be deduced from the Nash or Gagliardo–Nirenberg inequalities for the domain of the associated form, see [Ouh05, Chapt. 6]. We are going to extend this result to the non-autonomous setting.

Definition 4.2. *Let V be a subspace of $L^2(X)$. The space V is said to satisfy*

(i) *a Nash inequality if there exist constants $C_N, \mu > 0$ such that*

$$(4.3) \quad \|u\|_{L^2(X)}^{2+\frac{4}{\mu}} \leq C_N \|u\|_V^2 \|u\|_{L^1(X)}^{\frac{4}{\mu}} \quad \text{for all } u \in L^1(X) \cap V;$$

(ii) *a Gagliardo–Nirenberg inequality if there exist constants $C_G, N > 0$ such that*

$$(4.4) \quad \|u\|_{L^q(X)} \leq C_G \|u\|_{L^2(X)}^{1-N\frac{q-2}{2q}} \|u\|_V^{N\frac{q-2}{2q}} \quad \text{for all } u \in V$$

holds for all $q \in]2, \infty[$ such that $N^{\frac{q-2}{2q}} \leq 1$.

Sobolev spaces $H^1(I)$ on intervals $I \subset \mathbb{R}$ satisfy e.g. the Nash inequality, see e.g. [Maz85, § 1.4.8]. More generally, the same is true for each closed subspace V of $H^1(X)$ which has the $L^1 - H^1$ -extension property [AtE97, Lemma 2.7], where Ω is an arbitrary open set of \mathbb{R}^d . Several geometric conditions on $\Omega \subset \mathbb{R}^d$ under which a Sobolev space $V = H^k(\Omega)$ satisfies a Gagliardo–Nirenberg inequality are known, see e.g. [AF03, Chapter 5].

Here and in the following, we are adopting the usual notations introduced in (2.1)–(2.3).

Theorem 4.3. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ such that the associated evolution family \mathcal{U} is completely quasi-contractive with constant $\tilde{\omega} \in \mathbb{R}$. If V satisfies a Nash inequality (4.3) for some constants $\mu, C_N > 0$, then \mathcal{U} is ultracontractive and*

$$(4.5) \quad \|U(t, s)\|_{\mathcal{L}(L^1(\Omega), L^\infty(\Omega))} \leq \left(\frac{\mu C_N}{4\alpha}\right)^{\frac{\mu}{2}} (t-s)^{-\frac{\mu}{2}} e^{\max\{\omega, \tilde{\omega}\}(t-s)} \quad \text{for all } (t, s) \in \Delta.$$

Definition 4.4. *An evolution family \mathcal{U} on $L^2(X)$ is called linearly quasi-contractive if for some constants α_1, α_2 independent of p*

$$(4.6) \quad \|U(t, s)f\|_{L^p(X)} \leq e^{(t-s)(\alpha_1 + p\alpha_2)} \|f\|_{L^p(X)}$$

for all $(t, s) \in \Delta$, $f \in L^2(X) \cap L^p(X)$, and $p \in [2, \infty[$.

Theorem 4.5. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$. Assume that \mathcal{U} and $\overleftarrow{\mathcal{U}}^*$ are both linearly quasi-contractive with constants $\alpha_1, \alpha_2, \alpha_1^*, \alpha_2^*$. If V satisfies a Gagliardo–Nirenberg inequality for some $C_G, N > 0$, then \mathcal{U} is ultracontractive and we have*

$$(4.7) \quad \|U(t, s)\|_{\mathcal{L}(L^1(X), L^\infty(X))} \leq C_G^{\frac{N}{2}} \alpha^{-\frac{N^2}{4(N+2)}} e^{\tilde{\omega}(t-s)} (t-s)^{-\frac{N^2}{2(N+2)}} \quad \text{for all } (t, s) \in \Delta,$$

where

$$\tilde{\omega} := \omega + \alpha_1 + \alpha_1^* + \frac{2(N+2)}{N} [\alpha_2 + \alpha_2^*].$$

Proof of Theorem 4.3. Upon rescaling $U(t, s)$ by $e^{-\max\{\omega, \tilde{\omega}\}(t-s)}$ we can without loss of generality assume both \mathbf{a} to be coercive and the evolution family \mathcal{U} to be completely contractive. The first part of the proof is similar to that of [AtE97, Prop. 3.8]. Let $f \in L^1(X) \cap V$ and let $s \in [0, T]$ be fixed. Using (2.15), (2.3) and since $t \mapsto U(t, s)f \in MR(s, T; V, V')$ we obtain that for all $f \in V \cap L^1(X)$ and a.e. $(t, s) \in \Delta$

$$\begin{aligned} \frac{\partial}{\partial t} \|U(t, s)f\|_{L^2(X)}^2 &= 2 \operatorname{Re} \left\langle \frac{\partial}{\partial t} U(t, s)f, U(t, s)f \right\rangle \\ &= -2 \operatorname{Re} \langle A(t)U(t, s)f, U(t, s)f \rangle \\ &= -2a(t; U(t, s)f, U(t, s)f) \\ &\leq -2\alpha \|U(t, s)f\|_V^2 \\ &\leq -\frac{2\alpha}{C_N} \|U(t, s)f\|_{L^2(X)}^{2+\frac{4}{\mu}} \|U(t, s)f\|_{L^1(X)}^{-\frac{4}{\mu}}. \end{aligned}$$

It follows that

$$\frac{\partial}{\partial t} \left(\|U(t, s)f\|_{L^2(X)}^2 \right)^{-\frac{2}{\mu}} = -\frac{2}{\mu} \|U(t, s)f\|_{L^2(X)}^{-2-\frac{4}{\mu}} \frac{\partial}{\partial t} \|U(t, s)f\|_{L^2(X)}^2 \geq \frac{4\alpha}{\mu C_N} \|f\|_{L^1(X)}^{-\frac{4}{\mu}}.$$

Integrating this inequality between s and t we find

$$(4.8) \quad \|U(t, s)\|_{\mathcal{L}(L^1(X), L^2(X))} \leq \left(\frac{\mu C_N}{4\alpha}\right)^{\frac{\mu}{4}} (t-s)^{-\frac{\mu}{4}} \quad \text{for all } (t, s) \in \Delta.$$

In order to obtain the $L^2 - L^\infty$ -bound and thus prove the claimed ultracontractivity we will use the returned adjoint form $\overleftarrow{\mathbf{a}}^*$ introduced in Section 3. In fact, arguing as in the first part of the proof we find that the evolution family \overleftarrow{U}^* associated with $\overleftarrow{\mathbf{a}}^*$ satisfies (4.8) with the same bound. Then using the identity (3.3) we conclude that

$$(4.9) \quad \|U(t, s)\|_{\mathcal{L}(L^2(X), \mathcal{L}^\infty(X))} \leq \left(\frac{\mu C_N}{4\alpha}\right)^{\frac{\mu}{4}} (t-s)^{-\frac{\mu}{4}} \quad \text{for all } (t, s) \in \Delta.$$

Finally, the evolution law satisfied by \mathcal{U} completes the proof. \square

Linear L^p -quasi-contractivity turns out to be a key notion when it comes to checking ultracontractivity. In the proof of Theorem 4.5 we will need the following lemma.

Lemma 4.6. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ and let \mathcal{U} and $\overleftarrow{\mathcal{U}}^*$ be the evolution families associated with \mathbf{a} and $\overleftarrow{\mathbf{a}}^*$, respectively. Assume that \mathcal{U} and $\overleftarrow{\mathcal{U}}^*$ are both linearly quasi-contractive (with constants $\alpha_1, \alpha_2, \alpha_1^*, \alpha_2^*$). In addition we assume that there exist constants $\kappa_1, \kappa > 0$ such that*

$$(4.10) \quad \|U(t, s)\|_{\mathcal{L}(L^2(X), L^{N_2}(X))} \leq \kappa (t-s)^{-\frac{\kappa_1}{2}} e^{(t-s)[\alpha_1 + N_2\alpha_2]} \quad \text{for all } (t, s) \in \Delta$$

and

$$(4.11) \quad \|\overleftarrow{U}^*(t, s)\|_{\mathcal{L}(L^2(X), L^{N_2}(X))} \leq \kappa (t-s)^{-\frac{\kappa_1}{2}} e^{(t-s)[\alpha_1^* + N_2\alpha_2^*]} \quad \text{for all } (t, s) \in \Delta$$

where $N_2 := \frac{2N}{N-2}$ for some integer $N \geq 3$. Then \mathcal{U} is ultracontractive: more precisely,

$$(4.12) \quad \|U(t, s)\|_{\mathcal{L}(L^1(X), L^\infty(X))} \leq \kappa^{\frac{N}{2}} c (t-s)^{-\frac{N\kappa_1}{2}} e^{\tilde{\omega}(t-s)} \quad \text{for all } (t, s) \in \Delta \text{ with } t > s$$

for some positive constants $c, \mu > 0$ that depend only on N, κ_1 , where

$$\tilde{\omega} := [\alpha_1 + \alpha_1^*] + \mu[\alpha_2 + \alpha_2^*].$$

Proof. For the proof we follow similar argument as in [Ouh04, Thm. 5.2] and [Cou93] where ultracontractivity for semigroups are treated.

Step 1. We will first prove that

$$(4.13) \quad \|U(t, s)\|_{\mathcal{L}(L^2(X), L^\infty(X))} \leq \kappa^{\frac{N}{2}} C (t-s)^{-\frac{\kappa_1 N}{4}} e^{\alpha_1(t-s)} e^{\mu\alpha_2(t-s)},$$

where the positive constants C and μ depend only on N and κ_1 . For some $r > 2$ that will be fixed later we can combine (4.10) with the linear L^r -quasi-contractivity of \mathcal{U} and obtain by a version of Riesz–Thorin interpolation theorem [Dav07, Thm. 2.2.14] that for any $\theta \in [0, 1]$

$$\|U(t, s)\|_{\mathcal{L}(L^{p_1}(X), L^{q_1}(X))} \leq \kappa^\theta (t-s)^{-\frac{\kappa_1\theta}{2}} e^{(1-\theta)(t-s)[\alpha_1 + r\alpha_2]} e^{\theta(t-s)[\alpha_1 + \bar{2}\alpha_2]}$$

where $\frac{1}{p_1} := \frac{1-\theta}{r} + \frac{\theta}{2}$, $\frac{1}{q_1} := \frac{1-\theta}{r} + \frac{\theta}{N_2}$. Let now $p \in]2, \infty[$. Choosing $\theta := \frac{1}{p}$ and $r = 2(p-1)$ in the above equation we obtain that

$$(4.14) \quad \|U(t, s)\|_{\mathcal{L}(L^p(X), L^{N_p}(X))} \leq \kappa^{\frac{1}{p}} (t-s)^{-\frac{\kappa_1}{2p}} e^{(1-\frac{1}{p})(t-s)[\alpha_1 + 2(p-1)\alpha_2]} e^{\frac{1}{p}(t-s)[\alpha_1 + \bar{2}\alpha_2]}$$

holds for all $p \in]2, \infty[$ where $N_p := p\frac{N}{N-1}$.

Next, set $R = \frac{N}{N-1}$, $p_k = 2R^k$ and $t_k = \frac{N+1}{2N}(2R)^{-k} = \frac{N+1}{Np_k}2^{-k}$ for all $k \in \mathbb{N}$. Moreover, let $s_0 = s$ and $s_{k+1} = s_k + t_k(t-s)$ for each integer $k > 0$. Then we have $\sum_k t_k = 1$, $\sum_k \frac{1}{p_k} = \frac{N}{2}$. Furthermore, $s_{k+1} < s_k$ for all $k \in \mathbb{N}$ and $t = \lim_{k \rightarrow \infty} s_k$. Thus, applying (4.14) for $p = p_k$, using (4.6) and the evolution law satisfied by \mathcal{U} we deduce that

$$\begin{aligned}
\|U(t, s)\|_{\mathcal{L}(L^2(X), L^\infty(X))} &\leq \prod_{k \geq 0} \|U(s_{k+1}, s_k)\|_{\mathcal{L}(L^{p_k}(X), L^{Np_k}(X))} \\
&\leq \prod_{k \geq 0} \kappa^{\frac{1}{p_k}} t_k^{-\frac{\kappa_1}{2p_k}} (t-s)^{-\frac{\kappa_1}{2p_k}} e^{(1-\frac{1}{p_k})(s_{k+1}-s_k)[\alpha_1+2(p_k-1)\alpha_2]+\frac{1}{p_k}(s_{k+1}-s_k)[\alpha_1+\bar{2}\alpha_2]} \\
&= \kappa^{\frac{N}{2}} (t-s)^{-\frac{\kappa_1 N}{4}} \prod_{k \geq 0} t_k^{-\frac{\kappa_1}{2p_k}} e^{(1-\frac{1}{p_k})(s_{k+1}-s_k)[\alpha_1+2(p_k-1)\alpha_2]+\frac{1}{p_k}(s_{k+1}-s_k)[\alpha_1+\bar{2}\alpha_2]} \\
&\leq \kappa^{\frac{N}{2}} (t-s)^{-\frac{N\kappa_1}{4}} \prod_{k \geq 0} t_k^{-\frac{\kappa_1}{2p_k}} \prod_{k \geq 0} e^{(t-s)t_k\alpha_1} \prod_{k \geq 0} e^{(t-s)t_k\alpha_2[p_k-1-\frac{p_k-1}{p_k}+\frac{\bar{2}}{p_k}]} \\
&= \kappa^{\frac{N}{2}} (t-s)^{-\frac{\kappa_1 N}{4}} e^{(t-s)\alpha_1} \prod_{k \geq 0} t_k^{-\frac{\kappa_1}{2p_k}} e^{\alpha_2(t-s)\sum_{k \geq 0} t_k \left[\frac{(p_k-1)^2 + \frac{2N}{N-2}}{p_k} \right]} \\
&\leq \kappa^{\frac{N}{2}} (t-s)^{-\frac{\kappa_1 N}{4}} e^{\alpha_1(t-s)} \prod_{k \geq 0} t_k^{-\frac{\kappa_1}{2p_k}} e^{\alpha_2(t-s)\sum_{k \geq 0} t_k \left[\frac{(p_k-1)^2 + \frac{2N}{N-2}}{p_k} \right]} \\
&\leq \kappa^{\frac{N}{2}} C (t-s)^{-\frac{\kappa_1 N}{4}} e^{\alpha_1(t-s)} e^{\mu\alpha_2(t-s)},
\end{aligned}$$

where the positive constants C and μ depend only on N and κ_1 .

Step 2. It remains to estimate $U(t, s)$ in $\mathcal{L}(L^1(X), L^2(X))$. To this end, we will follow an idea in [Dan00, Corollary 5.3] and use the returned adjoint form $\overleftarrow{\mathfrak{a}}^*$. Indeed, by assumption $\overleftarrow{\mathfrak{a}}^*$ is linearly contractive. Thus one can just repeat the argument in Step 1 and obtain

$$(4.15) \quad \|\overleftarrow{U}^*(t, s)\|_{\mathcal{L}(L^2(X), L^\infty(X))} \leq \tilde{C} (t-s)^{-\frac{\kappa_1 N}{4}} e^{\alpha_1^*(t-s)} e^{\tilde{\mu}\alpha_2^*(t-s)}$$

for each $(t, s) \in \Delta$ and some constants $\tilde{C}, \tilde{\mu}$ that depend only on N, κ_1 . This yields, in turn, an estimate of $U(t, s)$ from $L^2(X)$ to $L^1(X)$, thanks to (4.15). Finally, using again the evolution law satisfied by U we conclude that \mathcal{U} is ultracontractive and (4.12) holds. \square

Proof of Theorem 4.5. Upon rescaling the evolution family by $e^{-\omega(t-s)}$ we can without loss of generality assume \mathcal{U} to be contractive. Let $q > 2$, $f \in L^q(X) \cap L^2(X)$ and set

$$\widehat{U}(t, s) := e^{-(t-s)[\alpha_1+q\alpha_2]} U(t, s) \quad \text{for each } (t, s) \in \Delta.$$

Because of (4.6) we have that $t \mapsto \|\widehat{U}(t, s)f\|_{L^q(X)}$ is decreasing on $[s, T]$ for each $s \in [0, T]$. Let now $(t, s) \in \Delta$: the contractivity of \mathcal{U} together with the Gagliardo–Nirenberg inequality and (2.15)

imply that for all $q > 2$ and all

$$\begin{aligned}
(t-s)\|\widehat{U}(t,s)f\|_{L^q(X)}^{\frac{4q}{N(q-2)}} &\leq \int_s^t \|\widehat{U}(r,s)f\|_{L^q(X)}^{\frac{4q}{N(q-2)}} dr \\
&\leq C_G^{\frac{4q}{N(q-2)}} \int_s^t \|\widehat{U}(r,s)f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \|\widehat{U}(r,s)f\|_V^2 dr \\
&\leq C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \int_s^t \left[\operatorname{Re} \mathbf{a}(r; \widehat{U}(r,s)f) + [\alpha_1 + q\alpha_2] \|\widehat{U}(r,s)f\|_{L^2(X)} \right] dr \\
&= -C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \int_s^t \operatorname{Re} \left\langle \frac{\partial}{\partial r} U(r,s)f, U(r,s)f \right\rangle dr \\
&= -C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} 2^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \int_s^t \frac{\partial}{\partial r} \|U(r,s)f\|_{L^2(X)}^2 dr \\
&= -C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} 2^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \left(\|U(t,s)f\|_{L^2(X)}^2 - \|f\|_{L^2(X)}^2 \right) \\
&\leq C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}-2} \|f\|_{L^2(X)}^2 \\
&= C_G^{\frac{4q}{N(q-2)}} \alpha^{-1} \|f\|_{L^2(X)}^{\frac{4q}{N(q-2)}}.
\end{aligned}$$

Here we have used that $t \mapsto \widehat{U}(t,s)f$ solves (2.4) with $A(t)$ replaced by $A(t) + [\alpha_1 + q\alpha_2]$. It follows that for all $(t,s) \in \Delta$ and all $q > 2$

$$(4.16) \quad \|U(t,s)\|_{\mathcal{L}(L^2(X),L^q(X))} \leq C_G \alpha^{-N \frac{q-2}{4q}} (t-s)^{-N \frac{q-2}{4q}} e^{(t-s)[\alpha_1 + q\alpha_2]}$$

and likewise

$$(4.17) \quad \|\overleftarrow{U}^*(t,s)\|_{\mathcal{L}(L^2(X),L^q(X))} \leq C_G \alpha^{-N \frac{q-2}{4q}} (t-s)^{-N \frac{q-2}{4q}} e^{(t-s)[\alpha_1^* + q\alpha_2^]}$$

Choosing now $q = \frac{2(N+2)}{N+2-2} = \frac{2(N+2)}{N}$ in (4.16)-(4.17) we obtain that (4.11)-(4.10) are fulfilled with $\kappa = C_G \alpha^{-\frac{N}{2(N+2)}}$ and $\kappa_1 = \frac{N}{N+2}$. Thus we conclude by Lemma 4.6 that \mathcal{U} is ultracontractive and (4.7) holds. \square

Remark 4.7. *Theorem 4.5 holds in particular for $N \frac{q-2}{2q} = 1$: in this case the Gagliardo–Nirenberg inequality becomes*

$$\|u\|_{L^q(X)} \leq C_G \|u\|_V \quad \text{for all } u \in V,$$

i.e., (4.4) reduces to the elementary assumption that V is continuously embedded in some $L^q(X)$: a classical Sobolev inequality. More precisely, if there exists $N > 2$ such that

$$(4.18) \quad V \subset L^q(X) \text{ for } \frac{1}{q} = \frac{1}{2} - \frac{1}{N},$$

then \mathcal{U} is ultracontractive and (4.7) holds.

5. GAUSSIAN BOUNDS

The existence of integral kernels of the evolution family, established in the previous section, paves the way to the discussion of kernel estimates. We have already seen a simple class of evolution

families satisfying Gaussian-type estimates in Example 3.13. In the following we are going to study this issue more systematically.

Definition 5.1. *Let \mathcal{U} be an evolution family on $L^2(\mathbb{R}^d)$ with an integral kernel Γ . Then \mathcal{U} is said to satisfy Gaussian bounds if there exist $b, c > 0$, $n > 0$, and $\omega \in \mathbb{R}$ such that*

$$(5.1) \quad |\Gamma(t, s; x, y)| \leq ce^{\omega(t-s)}(t-s)^{-\frac{n}{2}} \exp\left(-b\frac{|x-y|^2}{t-s}\right)$$

for all $(t, s) \in \Delta$ and a.e. $x, y \in \mathbb{R}^d$.

We regard $L^2(\Omega)$ as a closed subspace of $L^2(\mathbb{R}^d)$, extending operators on $L^2(\Omega)$ to $L^2(\mathbb{R}^d)$ by 0. In this way we can naturally define Gaussian bounds for operators on $L^2(\Omega)$.

Gaussian bounds for evolution equations can be characterized by ultracontractivity. This characterisation is well-known for autonomous closed forms. It is based on the so-called *Davies' trick*, first appeared in [Dav87], see also [AtE97, Thm. 3.3] and [Are06, Thm. 13.1.4] for more general versions. Davies' trick is essentially an algorithm centered around an auxiliary result, whose non-autonomous counterpart is Theorem 5.2 below.

To begin with we introduce a suitable space

$$(5.2) \quad W := \left\{ \psi \in C^\infty(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d) \mid \|D_j \psi\|_\infty \leq 1, \|D_i D_j \psi\|_\infty \leq 1, i, j = 1, \dots, n \right\}$$

of smooth functions. By [Rob91, p. 200–202], the function $d : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}_+$ defined by

$$d(x, y) := \sup\{|\psi(x) - \psi(y)| \mid \psi \in W\}$$

is a metric equivalent to the Euclidean one: there exists $\beta > 0$ such that

$$(5.3) \quad \beta|x-y| \leq d(x, y) \leq \beta^{-1}|x-y| \quad \text{for all } x, y \in \mathbb{R}^d.$$

Let \mathcal{U} be an evolution family on $L^2(\Omega)$ and, as usual, extend it if needed to $L^2(\mathbb{R}^d)$. For a fixed $\psi \in W$ we define perturbed evolution families \mathcal{U}_ρ on $L^2(\mathbb{R}^d)$ by

$$U_\rho(t, s) := U_\rho^\psi(t, s) := M_\rho U(t, s) M_\rho^{-1}, \quad \rho \in \mathbb{R},$$

where M_ρ is the isomorphism on $L^2(\mathbb{R}^d)$ defined by

$$(M_\rho g)(x) := e^{-\rho\psi(x)} g(x), \quad g \in L^2(\mathbb{R}^d), \quad x \in \mathbb{R}^d.$$

Gaussian bounds for \mathcal{U} can now be derived from uniform ultracontractivity of the perturbed evolution families \mathcal{U}_ρ with respect to ρ and ψ . The proof of this fact is very similar to that of the autonomous case studied in [AtE97, Prop. 3.3]: our result contains [Dan00, Thm. 6.1] as a special case.

Theorem 5.2. *Let \mathcal{U} be an evolution family on $L^2(\Omega)$. Then the following are equivalent:*

(i) *There exist $c > 0$, $n > 0$, and $\tilde{\omega} \in \mathbb{R}$ such that*

$$(5.4) \quad \|U_\rho(t, s)\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} \leq c(t-s)^{-\frac{n}{2}} e^{\tilde{\omega}(1+\rho^2)(t-s)}$$

for all $\rho \in \mathbb{R}$, $\psi \in W$, and $(t, s) \in \Delta$.

(ii) *\mathcal{U} satisfies Gaussian bounds.*

In this case \mathcal{U} satisfies (5.1) with $b = \frac{\beta^2}{4\tilde{\omega}}$, $c, \tilde{\omega}$ are as in (5.4) and β is the constant in (5.3).

Proof. (i) \Rightarrow (ii) Assume that (i) holds, hence \mathcal{U} is given by an integral kernel $(\Gamma(t, s))_{(t, s) \in \Delta} \subset L^\infty(\Omega \times \Omega)$. Thus for each $\psi \in W$ the perturbed evolution family \mathcal{U}_ρ is also given by a kernel Γ_ρ with

$$\Gamma_\rho(t, s; x, y) = e^{\rho(\psi(y) - \psi(x))} \Gamma(t, s; x, y), \quad (t, s) \in \Delta, \quad x, y \in \Omega,$$

whose L^∞ -norm agrees with the $\mathcal{L}(L^1, L^\infty)$ -norm of \mathcal{U}_ρ . Using (5.4) we obtain that for all $(t, s) \in \Delta$, $\rho \in \mathbb{R}$, and $\psi \in W$

$$|\Gamma(t, s; x, y)| \leq c(t-s)^{-\frac{n}{2}} e^{\tilde{\omega}(1+\rho^2)(t-s)} e^{\rho(\psi(y) - \psi(x))}, \quad (t, s) \in \Delta, \quad x, y \in \Omega.$$

Replacing ρ by $-\rho$ in this inequality yields

$$|\Gamma(t, s; x, y)| \leq c(t-s)^{-\frac{n}{2}} e^{\tilde{\omega}(1+\rho^2)(t-s)} e^{-\rho|\psi(y) - \psi(x)|}.$$

By [AtE97, Lemma 3.4] it follows that

$$|\Gamma(t, s; x, y)| \leq c(t-s)^{-\frac{n}{2}} e^{\tilde{\omega}(1+\rho^2)(t-s)} e^{-\rho d(x, y)},$$

for a.e. $x, y \in \Omega$, all $\rho \geq 0$, and all $(t, s) \in \Delta$. Let now $x, y \in \Omega$ and $(t, s) \in \Delta$ be fixed. Then the function

$$\rho \mapsto c(t-s)^{-\frac{n}{2}} e^{\tilde{\omega}(1+\rho^2)(t-s)} e^{-\rho d(x, y)}$$

attains its minimum at $\rho_0 := (2w(t-s))^{-1} d(x, y)$. Applying again [AtE97, Lemma 3.4] and using (5.3) we deduce that \mathcal{U} satisfies (5.1) with $b = \frac{\beta^2}{4\tilde{\omega}}$ and thus (ii) holds.

(ii) \Rightarrow (i)

One finds

$$\begin{aligned} \|U_\rho(t, s)\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} &\leq \|\Gamma_\rho(t, s; \cdot, \cdot)\|_{L^\infty(\Omega \times \Omega)} \\ &\leq \text{ess sup}_{x, y \in \Omega} |\Gamma(t, s; x, y)| e^{|\rho| |\psi(x) - \psi(y)|} \\ &\leq c(t-s)^{-\frac{n}{2}} e^{\omega(t-s)} e^{-b \frac{|x-y|^2}{t-s}} e^{\beta^{-1} |\rho| |x-y|} \\ &\leq c(t-s)^{-\frac{n}{2}} e^{(\omega + \hat{\omega} \rho^2)(t-s)} \end{aligned}$$

where β is as in (5.3) and $\hat{\omega} := \frac{1}{4b\beta^2}$, i.e., \mathcal{U}_ρ is ultracontractive. \square

The form domain V of $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ is said to be W -invariant if $M_\rho V \subset V$ for all $\rho \in \mathbb{R}$ and $\psi \in W$. In this case the family of mappings \mathbf{a}^ρ given by

$$(5.5) \quad \mathbf{a}^\rho(t; u, v) := \mathbf{a}(t; M_\rho u, M_\rho^{-1} v), \quad \rho \in \mathbb{R}, \quad t \in [0, T], \quad u, v \in V,$$

is well-defined and in fact $\mathbf{a}^\rho \in \text{FORM}([0, T]; V, H)$. For each $\rho \in \mathbb{R}$, $\psi \in W$, and $t \in [0, T]$ consider the operator family

$$\begin{aligned} A^\rho(t) &:= M_\rho^{-1} A(t) M_\rho, \\ D(A^\rho(t)) &:= \left\{ u \in L^2(\Omega) \mid M_\rho u \in D(A(t)) \right\}. \end{aligned}$$

Now, $-A^\rho(t)$ is for all $\rho \in \mathbb{R}$ and all $t \in [0, T]$ the generator of the semigroup \mathcal{T}_t^ρ given by

$$T_t^\rho(r) := M_\rho^{-1} e^{-rA(t)} M_\rho, \quad r \geq 0.$$

Lemma 5.3. *Assume that V is W -invariant and $\mathbf{a}^\rho \in \text{FORM}([0, T]; V, H)$ for each $\rho \in \mathbb{R}$ such that there exist $M_\rho, \alpha_\rho > 0$ and $\omega_\rho \in \mathbb{R}$ with*

$$(5.6) \quad \begin{aligned} |a^\rho(t; u, v)| &\leq M_\rho \|u\|_V \|v\|_V \\ \text{Re } a^\rho(t; u, u) + \omega_\rho \|u\|_H^2 &\geq \alpha_\rho \|u\|_V^2 \end{aligned} \quad \text{for all } t \in [0, T], u, v \in V.$$

Then $(A^\rho(t))_{t \in [0, T]}$ and \mathcal{U}_ρ are the operator family and the evolution family on H associated with \mathbf{a}^ρ , respectively.

Proof. Let $\rho \in \mathbb{R}, \psi \in W$ and $t \in [0, T]$ be fixed and denote by $C^\rho(t)$ the operator associated with $a^\rho(t; \cdot, \cdot)$ on H . Then for each $f, u \in L^2(\Omega)$ we have, $u \in D(C^\rho(t))$ and $C^\rho(t)u = f$ if and only if $u \in V$ and

$$a(t; M_\rho u, M_\rho^{-1} v) = (f | v)_H \quad \text{for all } v \in V,$$

or equivalently if and only if $u \in V$ and

$$a(t; M_\rho u, w) = (M_\rho f | w)_H \quad \text{for all } w \in V.$$

This is equivalent to $M_\rho u \in D(A(t))$ and $M_\rho^{-1} A(t) M_\rho u = f$. We conclude that $C^\rho(t) = M_\rho^{-1} A(t) M_\rho$. The last assertion is easy to prove: in fact, \mathcal{U}_ρ solves the Cauchy problem associated with $(A(t))_{t \in [0, T]}$. \square

After all these preparatory results we are finally in the position to present our main theorems: given $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ we introduce two sets of assumptions, which impose a Sobolev-like embedding on V and a contractivity condition on the perturbed semigroups \mathcal{T}_t^ρ , and show that each of them imply Gaussian bounds for the evolution family associated with \mathbf{a} .

Theorem 5.4. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$. Assume that V is W -invariant and that (5.6) holds for a uniform choice of α and for ω_ρ such that*

$$(5.7) \quad \omega_\rho \leq \omega(1 + \rho)$$

for some constant $\omega > 0$ that is independent of ρ . Assume V satisfies a Nash inequality and the semigroups $(e^{-\omega_\rho r} \mathcal{T}_t^\rho(r))_{r \geq 0}$ are completely contractive for a.e. $t \in [0, T]$ and all $\rho \in \mathbb{R}$. Then the evolution family \mathcal{U} associated with \mathbf{a} satisfies Gaussian bounds.

Proof. By Propositions 3.5 and 3.9 the evolution family $(e^{-\omega_\rho(t-s)} U_\rho(t, s))_{(t, s) \in \Delta}$ is completely contractive. Thus by Theorem 4.3

$$(5.8) \quad \|e^{-\omega_\rho(t-s)} U_\rho(t, s)\|_{\mathcal{L}(L^1(\Omega), L^\infty(\Omega))} \leq \left(\frac{\mu C}{4\alpha}\right)^{\frac{\mu}{2}} (t-s)^{-\frac{\mu}{2}} e^{\omega_\rho(t-s)} \quad \text{for all } (t, s) \in \Delta.$$

Now, using (5.7) we obtain that \mathcal{U}_ρ satisfies (5.4): the claim follows from Theorem 5.2. \square

Theorem 5.5. *Let $\mathbf{a} \in \text{FORM}([0, T]; V, H)$ with associated evolution family \mathcal{U} . Assume that V is W -invariant and that (5.6) holds for a uniform choice of α and all t, u, v . Assume that V satisfies a Gagliardo–Nirenberg inequality and both \mathcal{U}_ρ and $\overleftarrow{\mathcal{U}}_\rho^*$ are linearly quasi-contractive for all $\rho \in \mathbb{R}$. Then \mathcal{U}_ρ is ultracontractive for all $\rho \in \mathbb{R}$ with*

$$(5.9) \quad \|U_\rho(t, s)\|_{\mathcal{L}(L^1(\Omega), L^\infty(\Omega))} \leq C \frac{N}{G} \alpha^{-\frac{N^2}{4(N+2)}} e^{(t-s)\tilde{\omega}_\rho} (t-s)^{-\frac{N^2}{2(N+2)}} \quad \text{for all } (t, s) \in \Delta,$$

where

$$\tilde{\omega}_\rho := \left(\omega_\rho + \alpha_{\rho,1} + \alpha_{\rho,1}^* + \frac{2(N+2)}{N} [\alpha_{\rho,2} + \alpha_{\rho,2}^*] \right)$$

and $\alpha_{\rho,i}, \alpha_{\rho,i}^*$ are the constants that appear in the linear quasi-contractivity estimate. Thus, if additionally $\omega_\rho, \alpha_{\rho,i}, \alpha_{\rho,i}^*, i = 1, 2$, can be chosen in such a way that

$$(5.10) \quad \tilde{\omega}_\rho \leq \omega_0(1 + \rho^2)$$

for some constant $\omega_0 > 0$ independent of ρ , then \mathcal{U} satisfies Gaussian bounds.

Proof. The assertion can be proved similarly to Theorem 5.4, based in this case on Theorem 5.2 and Theorem 4.5. \square

6. APPLICATIONS

6.1. Time-dependent pageranks. Let us study a model similar to that of Example 3.13: it is based on an idea originally presented in [Chu07] and thoroughly developed ever since, cf. the survey [Gle15], where the connectivity of \mathbf{G} describes the internal links of a server network – possibly the whole World Wide Web.

We thus consider an orientation of a *finite* complete graph (i.e., a graph such that either $(\mathbf{v}, \mathbf{w}) \in \mathbf{E}$ or $(\mathbf{w}, \mathbf{v}) \in \mathbf{E}$ for any $\mathbf{v}, \mathbf{w} \in \mathbf{V}$ with $\mathbf{v} \neq \mathbf{w}$).

As in 3.13, we assign a weight m to each edge: if e.g. $m(t, \mathbf{e}) \in \{0, 1\}$ for all $\mathbf{e} \in \mathbf{E}$ and all $t \in [0, T]$, then we are effectively shutting off/switching on certain links in the considered network. We then consider the matrix

$$(6.1) \quad A := \mathcal{I}\mathcal{M}(\mathcal{I}^-)^T(\mathcal{D}^{\text{out}})^{-1}$$

where \mathcal{I} is again the incidence matrix of \mathbf{G} (see Example 3.13), $\mathcal{I}^- := (\iota_{\mathbf{ve}}^-)$ is its negative part, $\mathcal{M} = \text{diag}(m(\mathbf{e}))_{\mathbf{e} \in \mathbf{E}}$, and $\mathcal{D}^{\text{out}} := \text{diag}(\text{deg}^{\text{out}}(\mathbf{v}))_{\mathbf{v} \in \mathbf{V}}$, where $\text{deg}^{\text{out}}(\mathbf{v}) := \sum_{\mathbf{e} \in \mathbf{E}} |\iota_{\mathbf{ve}}^- m(\mathbf{e})|$.

Then, A defines a so-called *heat kernel pagerank* $e^{-rA}x$ of \mathbf{G} with parameters r and x : here r is a positive time and x a probability distribution on \mathbf{V} , i.e., $x \in \mathbb{R}^{\mathbf{V}}$, $x_{\mathbf{v}} \geq 0$ for all $\mathbf{v} \in \mathbf{V}$ and $\|x\|_1 = 1$. The rationale behind this definition is that A is a column stochastic matrix, hence $(e^{-rA})_{r \geq 0}$ is a stochastic semigroup and $e^{-rA}f$ is thus again a probability distribution for all $r \geq 0$, which can be used to measure the relevance of a certain node within a network in a way similar to Google's classical PageRank, cf. [Mug14, § 2.1.7.3].

We can now consider a measurable function $t \mapsto (m(t, \mathbf{e}))_{\mathbf{e} \in \mathbf{E}}$ and accordingly a time-dependent matrix family $(A(t))_{t \in [0, T]}$ as in (6.1) these matrices will in general not be symmetric, but in view of finiteness of \mathbf{V} they are certainly associated with a form $\mathbf{a} \in \text{FORM}([0, T]; \ell^2(\mathbf{V}), \ell^2(\mathbf{V}))$. Accordingly, in view of Proposition 3.11 the associated evolution family $(\mathcal{U}(t, s))_{(t, s) \in \Delta}$ consists of stochastic operators and hence $\mathcal{U}(t, s)f$ is a probability distribution on \mathbf{V} for all $(t, s) \in \Delta$ and all probability distributions $f \in \mathbb{R}^{\mathbf{V}}$.

There is a known correspondence between linear transport differential equations on networks and flows on their underlying graphs [Dor05]: accordingly, our results also extend to the space-continuous case. In this way, the well-posedness result in [Bay12, § 6] – which relies on the assumption that the dependence of the graph on time is absolutely continuous – can be strengthened: we omit the details.

6.2. Black–Scholes equation with time-dependent volatility. The Cauchy problem consisting of the backward parabolic equation

$$(6.2) \quad u_t(t, x) + \frac{1}{2}x^2\sigma^2 u_{xx}(t, x) + rxu_x(t, x) - ru(t, x) = 0, \quad t \in [0, \tau], \quad x \in]0, \infty[,$$

along with the final value assignment

$$u(\tau, x) = h(x) \quad x \in]0, \infty[$$

was derived in [BS73] and is currently considered among the main mathematical tool in the pricing theory of European options: the positive constants σ, r describe volatility and interest rate of the system, respectively, whereas τ is the maturity time of an option.

An effective variational approach to the relevant operator appearing in the Black–Scholes equation has been discussed in [Ein08]: it is based on studying the sesquilinear form

$$(6.3) \quad \begin{aligned} a(u, v) := & \frac{\sigma^2}{2} \int_0^\infty x^2 u'(x) \overline{v'(x)} \, dx \\ & + (\sigma^2 - r) \int_0^\infty x u'(x) \overline{v(x)} \, dx + r \int_0^\infty u(x) \overline{v(x)} \, dx, \quad u, v \in V, \end{aligned}$$

defined on the form domain

$$V := \{u \in W_{loc}^{1,1}]0, \infty[\cap L^2]0, \infty[: \text{id} \cdot u' \in L^2]0, \infty[\},$$

which is a Hilbert space with respect to the inner product

$$(u|v)_V := \int_0^\infty x^2 u'(x) \overline{v'(x)} \, dx + \int_0^\infty u(x) \overline{v(x)} \, dx.$$

Then it was proved in [Ein08, § 7.2] that V is dense in $L^2]0, \infty[$ and that furthermore

$$(6.4) \quad \begin{aligned} |a(u, v)| & \leq \left(\frac{\sigma^2}{2} + |\sigma^2 - r| + |r| \right) \|u\|_V \|v\|_V \\ \text{Re } a(u, u) + \left(\sigma^2 - \frac{3r}{2} \right) \|u\|_{L^2}^2 & = \frac{\sigma^2}{2} \|u\|_V^2 \end{aligned} \quad \text{for all } u \in V,$$

i.e., a is bounded and elliptic; and $\text{Re } a(u, u) \geq 0$ if $3r \geq \sigma^2$.

The original Black–Scholes-theory assumes σ to be time-independent, but is rather unrealistic and has been questioned ever since: we mention the celebrated Heston model [Hes93], which leads to a non-autonomous PDE similar to (6.2), based on the assumption that the volatility evolves following a certain Brownian-like motion. This justifies the study of

$$u_t(t, x) + \frac{1}{2} x^2 \sigma^2(t) u_{xx}(t, x) + r x u_x(t, x) - r u(t, x) = 0, \quad t \in [0, \tau], \quad x \in]0, \infty[,$$

with measurable dependence $t \mapsto \sigma(t)$. The computations in (6.4) show that if $0 < \sigma_0 \leq \sigma(t) \leq \Sigma$ for a.e. $t \in [0, \tau]$, then $\mathbf{a} \in \text{FORM}([0, T]; V; H)$, where \mathbf{a} is defined by

$$\begin{aligned} a(t; u, v) := & \frac{\sigma^2(t)}{2} \int_0^\infty x^2 u'(x) \overline{v'(x)} \, dx \\ & + (\sigma^2(t) - r) \int_0^\infty x u'(x) \overline{v(x)} \, dx + r \int_0^\infty u(x) \overline{v(x)} \, dx, \quad u, v \in V. \end{aligned}$$

Furthermore, the semigroup associated with a is quasi-contractive and sub-Markovian: we deduce from Proposition 3.1 that such non-autonomous Black–Scholes equation is governed by a sub-Markovian evolution family \mathcal{U} that extrapolates to all $L^p]0, \infty[$ spaces, $p \in [2, \infty]$. In view of Proposition 3.6 we can also apply [Ein08, Thm. 7.2.5] and deduce that \mathcal{U} leaves invariant the

order interval $] - \infty, \text{id}]_H$. By [Ein08, Rem. 7.2.4] $\min\{\log, 0\} \in V$, hence $V \not\rightarrow L^\infty]0, \infty[$; however, it is unclear whether V satisfies a Nash or Gagliardo–Nirenberg inequality, which would imply ultracontractivity of the evolution family.

(A manifold of financial models exist that display a similar mathematical structure, albeit their meaning is different: the popular Cox–Ingersoll–Ross along with several other so-called *short-rate models* surveyed in [CKLS92] involve time-dependent σ and/or r and can be discussed with only minor variations to our treatment above.)

6.3. Second-order elliptic operators on networks. With the purpose of introducing a differential operator on a network-like structure, we consider once again a graph $G = (V, E)$ (like in Example 3.13) and identify each edge $e \in E$ with an interval $[0, 1]$. In other words, we are considering a collection of copies of $[0, 1]$ and gluing them in a graph-like fashion: we thus obtain what are often called *metric graphs* or *networks* in the literature [BK13, Mug14]. (For the sake of simplicity we are going to assume such a metric graph to be connected.) The history of non-autonomous diffusion equations on networks goes back at least to pioneering investigations by von Below, Lumer, and Schnaubelt: well-posedness results could be proved in [Bel88, LS99], further results on long-time asymptotics have been deduced in [ADKF14].

We are going to apply in this context the theory developed in the previous sections: when introducing operators on metric graphs we avoid to go into full details and refer the interested reader to [Mug14, Chapter 6]. To fix the ideas, consider a possibly infinite, but uniformly locally finite (see Example 3.13) graph $G = (V, E)$ and upon rescaling identify each edge e with an interval $(0, 1)$. On each interval e we consider the operator family

$$A_e(t) : u_e \mapsto -\frac{d}{dx} \left(c_e(t) \frac{d}{dx} u_e \right) - p_e(t) u_e, \quad t \in [0, T] :$$

we assume the coefficients

$$[0, T] \ni t \mapsto c_e(t) \in L^\infty(0, 1; \ell^\infty(E)) \quad \text{and} \quad [0, T] \ni t \mapsto p_e(t) \in L^1(0, 1; \ell^\infty(E))$$

to be measurable: this defines in a natural way an operator A with domain $D(A) := \widetilde{H}^2(\mathcal{G}) := H^2(0, 1; \ell^2(E))$ on the Hilbert space

$$H := L^2(\mathcal{G}) := L^2(0, 1; \ell^2(E)) .$$

We will additionally assume that the operator family is uniformly elliptic, i.e.,

$$(6.5) \quad c_e(t, x) \geq \gamma \quad \text{for all } e \in E \text{ and a.e. } x \in (0, 1), t \in [0, T],$$

for some $\gamma > 0$. In order to reflect the topology of the graph, transmission conditions in the vertices are required: the most common conditions are usually referred to as *continuity/Kirchhoff* and amount to asking that

- u is continuous, i.e., the boundary values of u_e and u_f agree whenever evaluated at endpoints of the intervals e, f that are glued together in the network \mathcal{G} (continuity);
- u satisfies a Kirchhoff-type rule, i.e., at any vertex the sum over all neighboring edges of the normal derivatives evaluated at the vertex vanishes.

However, many more boundary conditions are conceivable: indeed, a parametrization of a large class of boundary conditions that fits very well the setting of sesquilinear forms has been discussed in [Mug14, § 6.5.1], based on the finite case treated in [Kuc04, Thm. 5].

Fix a closed subspace Y of the Hilbert space $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$, let $(\Sigma(t))_{t \in [0, T]}$ be a family of bounded linear operators on Y , and consider the non-autonomous form \mathbf{a} defined by

$$a(t; u, v) = \int_0^1 \left[(c_{\mathbf{e}}(t, x) u'_{\mathbf{e}}(x) | v'_{\mathbf{e}}(x))_{\ell^2(\mathbf{E})} + (p_{\mathbf{e}}(t, x) u_{\mathbf{e}}(x) | v_{\mathbf{e}}(x))_{\ell^2(\mathbf{E})} \right] dx + (\Sigma(t) \underline{u} | \underline{v})_Y$$

with time-independent form domain

$$H_Y^1(\mathcal{G}) := \left\{ u \in \bigoplus_{\mathbf{e} \in \mathbf{E}} H^1(0, 1; \ell^2(\mathbf{E})) : \underline{u} \in Y \right\}$$

where

$$\underline{u} := \begin{pmatrix} (u_{\mathbf{e}}(0))_{\mathbf{e} \in \mathbf{E}} \\ (u_{\mathbf{e}}(1))_{\mathbf{e} \in \mathbf{E}} \end{pmatrix}.$$

Then the conditions in the vertices satisfied by functions in the domain of each operator $A(t)$ associated with \mathbf{a} can be written in a compact form as

$$(6.6) \quad \underline{u} \in Y \quad \text{and} \quad \underline{\underline{u}} + \Sigma(t) \underline{u} \in Y^\perp.$$

where

$$\underline{\underline{u}} := \begin{pmatrix} (-c_{\mathbf{e}}(0) u'_{\mathbf{e}}(0))_{\mathbf{e} \in \mathbf{E}} \\ (c_{\mathbf{e}}(1) u'_{\mathbf{e}}(1))_{\mathbf{e} \in \mathbf{E}} \end{pmatrix}.$$

We finally assume that for some $P, S > 0$

$$\|p(t)\|_{L^1} \leq P \quad \text{and} \quad \|\Sigma(t)\|_{\mathcal{L}(Y)} \leq S \quad \text{for all } t \in [0, T].$$

Then taking into account Lemma 2.7 it is easy to see that $\mathbf{a} \in \text{FORM}([0, T]; V, H)$.

Our abstract results in the previous sections hence yield the following.

Proposition 6.1. *Under the above assumptions on the coefficients $c_{\mathbf{e}}, p_{\mathbf{e}}$, the space Y , and the operators Σ , the form \mathbf{a} is associated with a strongly continuous evolution family \mathcal{U} on H . If $p_{\mathbf{e}}(t) \geq 0$ and $\Sigma(t)$ is accretive for a.e. $t \in [0, T]$, then \mathcal{U} is contractive.*

If all these coefficients are defined on the whole interval $[0, \infty[$, then \mathcal{U} extends to an evolution family on $\{(t, s) : 0 \leq s \leq t \leq \infty\}$.

We denote by P_Y the orthogonal projector of $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$ onto Y . Furthermore, for $a, b \in \mathbb{R} \cup \{\pm\infty\}$ we denote as in Section 3 by $[a, b]_H$, $[a, b]_{\ell^2 \times \ell^2}$, and $[a, b]_Y$ the corresponding order intervals in H , $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$, and Y , respectively (the latter inherits the lattice structure of $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$).

Due to the standing assumption that \mathbf{G} is uniformly locally finite, P_Y is a block operator matrix whose blocks are of the form $\frac{1}{n} J_n$ (J_n denoting the $n \times n$ all-1-matrix, n the degree of the corresponding vertex). Because P_Y leaves invariant the order interval $[-1, 1]_{\ell^2 \times \ell^2}$, we deduce that if merely $(e^{-r\Sigma(t)})_{r \geq 0}$ is L^p -contractive for some $p \in]1, \infty[$ and a.e. $t \in [0, T]$, then so is $(e^{-rA(t)})_{r \geq 0}$ and a.e. $t \in [0, T]$ and, by Proposition 3.1, also the evolution family \mathcal{U} .

Owing to Proposition 3.6 we can formulate the following generalization of [Mug14, Thm. 6.85] (see also [CM09, Prop. 5.1]).

Proposition 6.2. *Let $a, b \in \mathbb{R}$ with $a \leq 0 \leq b$ and $p_{\mathbf{e}}(t, x) \geq 0$ for all $\mathbf{e} \in \mathbf{E}$ and a.e. $t \in (0, T)$ and $x \in (0, 1)$. Then the evolution family \mathcal{U} associated with \mathbf{a} leaves invariant $[a, b]_H$ if*

- P_Y leaves invariant the order interval $[a, b]_Y$; and
- $e^{-r\Sigma(t)}$ leaves invariant $[a, b]_{\ell^2 \times \ell^2}$ for all $r \geq 0$ and a.e. $t \in [0, T]$.

Three special cases are particularly noteworthy.

Corollary 6.3. (1) If $e^{-r\Sigma(t)}$ (for all $r \geq 0$ and a.e. $t \in [0, T]$) and P_Y are real and positive, then \mathcal{U} is positive. If additionally $p_e \equiv 0$ and $\mathbf{1} \in Y$, then \mathcal{U} is stochastic.

(2) Let $p_e(t, x) \geq 0$ for all $e \in \mathbf{E}$ and a.e. $t \in (0, T)$ and $x \in (0, 1)$. If $e^{-r\Sigma(t)}$ (for all $r \geq 0$ and a.e. $t \in [0, T]$) and P_Y are ℓ^∞ -contractive, then \mathcal{U} is $L^\infty(0, 1; \ell^2(\mathbf{E}))$ -contractive.

(3) Under the assumptions of (2), let additionally $e^{-r\Sigma(t)^*}$ be ℓ^∞ -contractive for all $r \geq 0$ and a.e. $t \in [0, T]$. Then \mathcal{U} is completely contractive; accordingly, it extrapolates to a strongly continuous, contractive evolution family on all spaces $L^p(0, 1; \ell^2(\mathbf{E}))$, $p \in [1, \infty]$.

Example 6.4. The continuity/Kirchhoff vertex conditions are special cases of the general conditions in (6.6). Indeed, denote by c_V the vector in $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$ that consists of vertex-wise constants, i.e., entries of c_V agree whenever they correspond to endpoints of edges the same vertex $v \in \mathbf{V}$ is incident with. Let by Y the subspace of $\ell^2(\mathbf{E}) \times \ell^2(\mathbf{E})$ spanned by c_V and take $\Sigma = 0$: then (6.6) agrees with continuity–Kirchhoff conditions: we denote by

$$H^1(\mathcal{G})$$

the Sobolev space $H_Y^1(\mathcal{G})$ with respect to this distinguished space Y . Under stronger assumptions on p_e, c_e , a well-posedness result comparable to Proposition 6.1 has been obtained in [ADKF14, Thm. 3.3].

Because $H^1(\mathcal{G}) \hookrightarrow C(\mathcal{G})$, we furthermore deduce that $U(t, s)$ maps $C(\mathcal{G}) \cap L^2(\mathcal{G})$ into $C(\mathcal{G})$ for all $s \in [0, T]$ and a.e. $t \in (s, T]$.

Under the same assumption that $\Sigma \equiv 0$, we can consider two different cases:

- if $\int_{t_0}^t \text{ess inf } p_e(t, x) > 0$ for a.e. $x \in (0, 1)$ and all $e \in \mathbf{E}$, then \mathcal{U} is by Proposition 2.6 uniformly exponentially stable; if $p_e \equiv 0$ and the network is finite (i.e., $|\mathbf{E}| < \infty$), then by Corollary 3.8 and [Mug14, Proposition 6.70]

$$\lim_{t \rightarrow \infty} U(t, s)f = \frac{1}{\sqrt{|\mathbf{E}|}} \int_{\mathcal{G}} f dx \cdot \mathbf{1} \quad \text{for all } f \in H.$$

A similar result has been obtained for general inhomogeneous diffusion equations on finite networks in [ADKF14, § 5.1].

Proposition 6.5. Let $p_e(t, x) \geq 0$ for all $e \in \mathbf{E}$ and a.e. $t \in [0, T]$, $x \in (0, 1)$. Let furthermore $P_Y = (\pi_{ij}^Y)$ and $\Sigma(t) = (\sigma(t)_{ij})$ satisfy

- $\sum_j |\pi_{ij}| \leq 1$ for all i ;
- $\text{Re } \sigma(t)_{ii} \geq \sum_{j \neq i} |\sigma(t)_{ij}|$ for all i and a.e. $t \in [0, T]$;
- $\text{Re } \sigma(t)_{ii} \geq \sum_{j \neq i} |\sigma(t)_{ji}|$ for all i and a.e. $t \in [0, T]$.

Then \mathcal{U} is completely contractive. If additionally Y is a generalized ideal of $\langle c_V \rangle$, then \mathcal{U} is ultracontractive.

Proof. It follows from [Mug07, Lemma 6.1] and Corollary 6.3 that \mathcal{U} is completely contractive. It has been shown in [Prö, Chapt. 3] that $H^1(\mathcal{G})$ satisfies a Nash inequality whenever \mathcal{G} is a connected, locally finite metric graph with edge lengths uniformly bounded away from 0: accordingly, the non-autonomous form with domain $H^1(\mathcal{G})$ is associated with an ultracontractive \mathcal{U} , owing to Theorem 4.3.

In order to complete the proof, observe that by [CM09, Thm. 6.2] the semigroup associated with $a(t) \equiv a$ with domain $H_Y^1(\mathcal{G})$ is dominated by the semigroup associated with the same form with

domain $H^1(\mathcal{G})$, provided Y is a generalized ideal of $\langle c_V \rangle$ (in fact by [Nag86, Thm. C.II-5.5] the latter is the modulus semigroup of the former one). It is a direct consequence of Proposition 3.1 that the same holds for the associated evolution families, hence the former heat kernel inherits ultracontractivity from the latter one. \square

The following result seems to be new even in the autonomous case: in [Mug07] Gaussian bounds for heat kernels on *finite* networks have been proved only in the special case of $Y = \langle c_V \rangle$, see also [Mug14, Chapt. 7] for an abstract approach based on the theory of Dirichlet forms.

Corollary 6.6. *Under the assumptions of Proposition 6.5, let Y be $\langle c_V \rangle$ -invariant (i.e., the entrywise product ψc_V lies in Y for all $\psi \in Y$), where the vector c_V is defined as in Example 6.4. If furthermore $\Sigma(t)$ is diagonal for a.e. $t \in [0, T]$, then \mathcal{U} satisfies Gaussian bounds.*

We stress that our assumption on P_Y and Σ are only enforcing complete contractivity, but the evolution family need not be positive. An example is given by the non-autonomous parabolic equation on a loop with boundary conditions defined by $\Sigma \equiv 0$ and $Y = \langle \begin{pmatrix} 1 \\ -1 \end{pmatrix} \rangle$, which is $\langle c_V \rangle$ -invariant (here $c_V = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$): this equation is governed by a completely contractive and (in view of the Nash inequality for $H^1(\mathcal{G})$) ultracontractive evolution family \mathcal{U} , which therefore enjoys Gaussian bounds. However, \mathcal{U} is not positive, since neither is P_Y .

Proof of Corollary 6.6. We apply Davies' Trick in a slightly different version. Indeed, we adapt the usual setting to our network environment by introducing the space

$$W_{\mathcal{G}} := \{ \psi \in H^1(\mathcal{G}) \cap C^\infty(0, 1; \ell^2(\mathbf{E})) \mid \|\psi'\|_\infty \leq 1, \|\psi''\|_\infty \leq 1 \} :$$

then one can check that

$$d(x, y) := \sup\{ |\psi(x) - \psi(y)| \mid \psi \in W_{\mathcal{G}} \}, \quad x, y \in \mathcal{G},$$

defines a metric on \mathcal{G} that is equivalent to the canonical one [Mug14, § 3.2]. (Recall that $H^1(\mathcal{G})$ denotes the space $H_{\tilde{Y}}^1(\mathcal{G})$, where $\tilde{Y} = \langle c_V \rangle$ is the space spanned by c_V : each function in $H^1(\mathcal{G})$ is by definition continuous on the metric space \mathcal{G} . Observe that $\langle c_V \rangle$ is in fact an algebra with respect to the entry-wise product.) By definition, $H_{\tilde{Y}}^1(\mathcal{G})$ is $W_{\mathcal{G}}$ -invariant if and only if $\underline{u} \in Y$ implies $e^{\rho\psi}\underline{u} \in Y$, i.e., if and only if Y is $\langle c_V \rangle$ -invariant.

If $\Sigma(t) \equiv 0$, then the assertion has been proved in [Mug07, Thm. 4.7] by showing that the relevant form (let us denote it by \mathfrak{a}_0 to stress the absence of boundary terms) induces perturbed forms \mathfrak{a}_0^ρ that are associated with completely contractive perturbed evolution families (with the form domain being unchanged and still satisfying a Nash inequality). In the general case of $a(t; u, v) = a_0(t; u, v) + (\Sigma(t)\underline{u}|\underline{v})_Y$, we find that $a^\rho(t; u, v) = a_0^\rho(t; u, v) + (\Sigma(t)\underline{u}|\underline{v})_Y$. These forms are associated with completely contractive evolution families, hence the claim follows. \square

6.4. Second-order elliptic operators with complex coefficients on open domains. Let $\Omega \subset \mathbb{R}^d$ be an open set and consider on the complex Hilbert space $L^2(\Omega)$ the non-autonomous form $\mathfrak{a}_V : [0, T] \times V \times V$ defined by

$$(6.7) \quad \mathfrak{a}_V(t; u, v) := \sum_{k,j=1}^d \int_{\Omega} a_{kj}(t; x) D_k u \overline{D_j v} \, dx + \sum_{k=1}^d \int_{\Omega} [b_k(t; x) D_k u \bar{v} + c_k(t; x) u \overline{D_k v}] \, dx + \int_{\Omega} a_0(t; x) u \bar{v} \, dx$$

for all $u, v \in V$ where V is a closed subspace of $H^1(\Omega)$ that contains $H_0^1(\Omega)$. We assume that the coefficients $a_{k,j}, b_k, c_k, a_0$ lie in $L^\infty([0, T] \times \Omega; \mathbb{C})$. Moreover, we assume that the principal part is uniformly elliptic, i.e., there exist a constant $\nu > 0$ such that

$$(6.8) \quad \operatorname{Re} \sum_{k,j=1}^d a_{kj}(t; x) \xi_k \bar{\xi}_j \geq \nu |\xi|^2 \quad \text{for a.e. } t \in [0, T], \text{ a.e. } x \in \Omega, \text{ and all } \xi \in \mathbb{C}^n.$$

Then \mathfrak{a}_V defined in (6.7) belongs to $\text{FORM}([0, T]; V, L^2(\Omega))$ [Ouh05, Section 4.1]. In fact, we have

$$\begin{aligned} |a_V(t; u, v)| &\leq M \|u\|_V \|v\|_V \\ \operatorname{Re} \mathfrak{a}_V(t; u, u) + \omega \|u\|_{L^2(\Omega)}^2 &\geq \frac{\nu}{2} \|u\|_V^2, \end{aligned} \quad \text{for all } u, v \in V \text{ and a.e. } t \in [0, T],$$

where $M > 0$ is a constant depending only on $\|a_{kj}\|_\infty, \|b_k\|_\infty, \|c_k\|_\infty$, and $\|a_0\|_\infty$, and we can choose $\omega = \sum_{k=1}^d \frac{1}{2} (\|\operatorname{Re}(b_k + c_k)\|_\infty + \|\operatorname{Im}(b_k - c_k)\|_\infty^2) + \|(\operatorname{Re} a_0)^-\|_\infty$.

Here $(\operatorname{Re} a_0)^- = \max\{0, -\operatorname{Re} a_0\}$. We can then associate an operator $\mathcal{A}_V(t) \in \mathcal{L}(V, V')$ with the form \mathfrak{a}_V which is formally given by

$$(6.9) \quad \mathcal{A}_V(t) = - \sum_{k,j=1}^d D_j a_{kj} D_k u + \sum_{k=1}^d b_k D_k u - \sum_{k=1}^d D_k (c_k u) + c_0 u.$$

Let $A_V(t)$ be the operator associated with $a(t; \cdot, \cdot)$ on $L^2(\Omega)$. Thus $A_V(t)$ is the realization of $\mathcal{A}_V(t)$ in $L^2(\Omega)$ with various boundary condition which are determined by the form domain V . For example $A_V(t)$ is the realization of $\mathcal{A}_V(t)$ with

- (a) Dirichlet boundary condition if $V = H_0^1(\Omega)$.
- (b) Neumann boundary condition if $V = H^1(\Omega)$.
- (c) Mixed boundary condition if

$$V = \overline{\{u|_\Omega : u \in C_c^\infty(\mathbb{R}^d \setminus \Gamma)\}}^{H^1(\Omega)}$$

where Γ is a closed subset of the boundary of Ω .

In particular, \mathfrak{a}_V is associated with an evolution family \mathcal{U}_V that governs the non-autonomous problem driven by the operator family $(A_V(t))_{t \in [0, T]}$. Each of these evolution families is positive, it dominates the evolution family $\mathcal{U}_{H_0^1}$ and is dominated by \mathcal{U}_{H^1} . Following [Ouh04] we introduce the following notations:

$$\begin{aligned} f_k(t, x) &:= \sum_{j=1}^d D_j (\operatorname{Im} a_{kj}(t, x)), \\ m(t, x) &:= \frac{1}{4\nu} \sum_{k=1}^d \left[f_k(t, x) + \operatorname{Im}(c_k(t, x) - b_k(t, x)) \right], \\ \Re_V(t; u, v) &:= \int_\Omega \sum_{k,j=1}^d \operatorname{Re}(a_{kj}) D_k u \overline{D_j v} dx + \sum_{k,j=1}^d \int_\Omega \left[\operatorname{Re}(b_k) D_k u \overline{v} dx + \operatorname{Re}(c_k) u \overline{D_k v} \right] + \int_\Omega \operatorname{Re}(a_0) u \overline{v} dx. \end{aligned}$$

Lemma 6.7. *Let \mathfrak{a}_V be given by (6.7) and denote by \mathcal{U}_V the associated evolution family on $L^2(\Omega)$. Assume that $(|v| \wedge 1) \operatorname{sgn} v \in V$ for all $v \in V$. Moreover, we assume that $a_{kj}(\cdot, \cdot)$ are real-valued*

functions for all $k, j = 1, 2, \dots, d$. Then the evolution family \mathcal{U}_V is L^p -quasi-contractive for all $p \in (1, \infty[$ and we have

$$(6.10) \quad \|\mathcal{U}_V(t, s)f\|_{L^p(\Omega)} \leq e^{(t-s)\tilde{\omega}_p} \|f\|_{L^p(\Omega)} \quad \text{for all } (t, s) \in \Delta,$$

where

$$(6.11) \quad \tilde{\omega}_p := \begin{cases} \|(\operatorname{Re} a_0)^-\|_\infty + \frac{1}{\nu} \left(\frac{1}{p} + \frac{1}{2} \right) \sum_{k=1}^d \|b_k - c_k\|_\infty^2 + \frac{p}{\nu} \sum_{k=1}^d \|\operatorname{Re} c_k\|_\infty^2 & \text{if } p \in [2, \infty[, \\ \|(\operatorname{Re} a_0)^-\|_\infty + \frac{1}{\nu} \left(\frac{1}{2} + \frac{p-1}{p} \right) \sum_{k=1}^d \|b_k - c_k\|_\infty^2 + \frac{p}{\nu(p-1)} \sum_{k=1}^d \|\operatorname{Re} b_k\|_\infty^2 & \text{if } p \in]1, 2]. \end{cases}$$

Proof. The assertion follows from Theorem 3.10 and [Ouh04, Thm. 4.3]. \square

We can also discuss the case where $a_{k,j}$ are complex-valued functions.

Lemma 6.8. *Let \mathfrak{a}_V be given by (6.7) such that $(|v| \wedge 1) \operatorname{sgn} v \in V$ for all $v \in V$ and denote by \mathcal{U}_V the associated evolution family on $L^2(\Omega)$. Assume that $f_k \in L^\infty([0, T] \times \Omega)$, $\operatorname{Im}(a_{k,j}(t, \cdot) + a_{j,k}(t, \cdot)) = 0$ for all $k, j = 1, 2, \dots, d$ and a.e. $t \in [0, T]$. If either of the conditions*

- (i) $V = H_0^1(\Omega)$;
- (ii) $V \neq H_0^1(\Omega)$, $(\operatorname{Re} u)^+ \in V$ for all $u \in V$ and there exists two constants $c_1, c_2 > 0$ such that

$$\int_\Omega m(t; x) |u|^2 dx \geq c_1 \int_\Omega |u|^2 dx + c_2 \operatorname{Re} \mathfrak{R}_V(t; u, u) \quad u \in V \text{ and } t \in [0, T];$$

are satisfied, then \mathcal{U}_V is L^p -quasi-contractive and (6.10) holds (up to replacing $\operatorname{Re} a_0(t, \cdot)$ by $\operatorname{Re} a_0(t, \cdot) - m$ in the expression of $\tilde{\omega}_p$).

Proof. The assertion follows again from [Ouh04, Thm. 4.4] and Theorem 3.10. \square

Remark 6.9. *If \mathfrak{a}_V fulfills the assumptions of Lemma 6.7 or those of Lemma 6.8, then we see that the evolution family \mathcal{U}_V is linearly quasi-contractive where (4.6) is satisfied with*

$$(6.12) \quad \alpha_1 = \|(\operatorname{Re} a_0 - m)^-\|_\infty + \frac{1}{\nu} \sum_{k=1}^d \|b_k - c_k\|_\infty^2$$

and

$$(6.13) \quad \alpha_2 = \frac{1}{\nu} \sum_{k=1}^d \|\operatorname{Re} c_k\|_\infty^2.$$

Likewise, the evolution family $\overleftarrow{\mathcal{U}}_V^*$ associated with $\overleftarrow{\mathfrak{a}}_V^*$ is linearly quasi-contractive and (4.6) is satisfied with $\alpha_1^* = \alpha_1$ and

$$(6.14) \quad \alpha_2^* = \frac{1}{\nu} \sum_{k=1}^d \|\operatorname{Re} b_k\|_\infty^2.$$

In view of Remark (6.9), the following corollary follows directly from Lemma 6.7 and Lemma 6.8.

Corollary 6.10. *Let \mathfrak{a} be given by (6.7) and denote by \mathcal{U}_V the associated evolution family on $L^2(\Omega)$. Suppose that V satisfies a Gagliardo Nirenberg inequality and that the assumptions of Lemma 6.7 (respectively that of Lemma 6.8) holds. Then \mathcal{U}_V is ultracontractive and satisfies 4.7 with $\mu = \nu$ and $\alpha_1 = \alpha_1^*, \alpha_2, \alpha_2^*$ are defined by (6.12)-(6.14).*

Now we are going to prove that the evolution family \mathcal{U}_V governed by the time dependent elliptic operator (6.9) satisfies Gaussian bounds. We known from Theorem 5.2 that U_V satisfies Gaussian bounds if and only if there exist a constants $c > 0, n > 0$ and $\omega \in \mathbb{R}$ such that

$$\|M_\rho U_V(t, s) M_\rho^{-1}\|_{\mathcal{L}(L^1(\mathbb{R}^d), L^\infty(\mathbb{R}^d))} \leq c(t-s)^{-\frac{n}{2}} e^{\omega(1+\rho^2)(t-s)}$$

for all $\rho \in \mathbb{R}, \psi \in W$ and $0 \leq s < t \leq T$. Let \mathbf{a} given by (6.7). Then the non-autonomous form $\mathbf{a}^\rho(t, u, v) := a(t; M_\rho u, M_\rho^{-1} v)$ is given by

$$(6.15) \quad \mathbf{a}_\rho(t; u, v) := \sum_{k,j=1}^d \int_{\Omega} a_{kj}(t; x) D_k u \overline{D_j v} \, dx + \sum_{k=1}^d \int_{\Omega} [b_{k,\rho}(t; x) D_k u \bar{v} + c_{k,\rho}(t; x) u \overline{D_k v}] \, dx + \int_{\Omega} a_{0,\rho}(t; x) u \bar{v} \, dx$$

where

$$\begin{aligned} b_{k,\rho} &= b_k - \rho \sum_{j=1}^d a_{kj} D_j \psi \\ c_{k,\rho} &= c_k + \rho \sum_{i=1}^d a_{ik} D_i \psi, \\ a_{0,\rho} &= a_0 - \rho^2 \sum_{i,k=1}^d a_{ik} D_i \psi D_k \psi + \rho \sum_{k=1}^d b_k D_k \psi - \rho \sum_{k=1}^d c_k D_k \psi. \end{aligned}$$

In the following we define for each $\rho \in \mathbb{R}, \psi \in W$ the constants $\alpha_{i,\rho}, \alpha_{i,\rho}^*, i = 1, 2$, via formulas which are analogous to (6.12), (6.13) and (6.14) where $\operatorname{Re} a_0$ is replaced by $\operatorname{Re} a_0 - m$ if $a_{k,j}$ are complex-valued functions. Further, we set

$$(6.16) \quad c_0 := \max\{\|a_{k,j}\|_\infty, \|b_k\|_\infty, \|c_k\|_\infty, \|c_0\|_\infty, k, j = 1, 2, \dots, d\}$$

and

$$(6.17) \quad \omega := 4c_0 d^2 + 4c_0 d^3 \nu^{-1}.$$

Lemma 6.11. (a) For all $\rho \in \mathbb{R}, \psi \in W$

$$(6.18) \quad \operatorname{Re} a^\rho(t; u, u) + \omega(1 + \rho^2) \|u\|^2 \geq \frac{\nu}{2} \|u\|_V^2 \quad \text{for a.e. } t \in [0, T], \text{ and all } u \in V$$

(b) Assume that $f_k \in L^\infty([0, T] \times \Omega)$, $\operatorname{Im} [a_{k,j}(t, \cdot) + a_{j,k}(t, \cdot)] = 0$ for all $k, j = 1, 2, \dots, d$ and $t \in [0, T]$. Then for all $\rho \in \mathbb{R}, \psi \in W$ we have

$$(6.19) \quad \alpha_{1,\rho} = \alpha_{1,\rho}^* \leq 2\alpha_1 + \rho^2(1 + 2d^2 c_0 + 4d^3 c_0^2 \nu^{-1}) + c_0 d^2$$

$$(6.20) \quad \alpha_{2,\rho} \leq 2\alpha_2 + 2d^3 \rho^2 c_0^2 \nu^{-1}$$

$$(6.21) \quad \alpha_{2,\rho}^* \leq 2\alpha_2^* + 2d^3 \rho^2 c_0^2 \nu^{-1}.$$

Proof. (a) We first show (6.18). Let $k = 1, \dots, d$ and $u \in V$

$$\begin{aligned}
& \left| \operatorname{Re} \left[b_{k,\rho}(t; x) D_k u \bar{u} + c_{k,\rho}(t; x) u \overline{D_k u} \right] \right| \\
& \leq \left| b_k(t; x) D_k u \bar{u} + c_k(t; x) u \overline{D_k u} \right| + |\rho| \left| \sum_{j=1}^d a_{kj}(t; x) D_j \psi D_k u \bar{u} - \sum_{i=1}^d a_{ik}(t; x) D_i \psi u \overline{D_k u} \right| \\
& \leq 2c_0 |D_k u| |u| + 2d|\rho|c_0 |D_k u| |u| = 2c_0(1 + d|\rho|) |D_k u| |u| \\
& \leq 2c_0(1 + d|\rho|) \varepsilon |D_k u|^2 + \frac{1}{2\varepsilon} |u|^2 c_0(1 + d|\rho|) \quad (\varepsilon > 0).
\end{aligned}$$

Choosing $\varepsilon = \frac{\nu}{4c_0(1+d|\rho|)}$ in the above inequality yields

$$\begin{aligned}
& \left| \operatorname{Re} \left[b_{k,\rho}(t; x) D_k u \bar{u} + c_{k,\rho}(t; x) u \overline{D_k u} \right] \right| \leq \frac{\nu}{2} |D_k u|^2 + 2c_0^2(1 + d|\rho|)^2 \nu^{-1} |u|^2 \\
& \leq \frac{\nu}{2} |D_k u|^2 + 4c_0^2 d^2 (1 + \rho^2) \nu^{-1} |u|^2.
\end{aligned}$$

If follows

$$(6.22) \quad \left| \operatorname{Re} \sum_{k=1}^d \left[b_{k,\rho}(t; x) D_k u \bar{v} + c_{k,\rho}(t; x) u \overline{D_k v} \right] \right| \leq \frac{\nu}{2} \sum_{k=1}^d |D_k u|^2 + 4c_0^2 d^3 (1 + \rho^2) \nu^{-1} |u|^2$$

Likewise,

$$(6.23) \quad \left| \operatorname{Re} a_{0,\rho}(t; x) u \bar{u} \right| \leq 4c_0 d^2 (1 + \rho^2) |u|^2$$

Combining (6.8), (6.22) and (6.23) yields (6.18).

(b) Using again that $|D_j \psi| < 1, i = 1, 2, \dots, d$ we obtain

$$(6.24) \quad \|\operatorname{Re} c_{k,\rho}\|_\infty^2 \leq 2\|\operatorname{Re} c_k\|_\infty^2 + 2\rho^2 \left(\sum_{i=1}^d \|a_{ik}\|_\infty \right)^2 \leq 2\|\operatorname{Re} c_k\|_\infty^2 + 2d^2 \rho^2 c_0^2,$$

and

$$(6.25) \quad \|\operatorname{Re} b_{k,\rho}\|_\infty^2 \leq 2\|\operatorname{Re} b_k\|_\infty^2 + 2d^2 \rho^2 c_0^2,$$

This proves (6.20) and (6.21). Next, using again that $|D_j \psi| < 1, i = 1, 2, \dots, d$ we obtain

$$\begin{aligned}
(6.26) \quad \frac{1}{\nu} \sum_k \|b_{k,\rho} - c_{k,\rho}\|_\infty^2 &= \frac{1}{\nu} \sum_k \left[\|b_k - a_k - \rho \sum_{j=1}^d a_{kj} D_j \psi - \rho \sum_{i=1}^d a_{ik} D_i \psi\|_\infty^2 \right] \\
&\leq \frac{1}{\nu} \sum_k \left[2\|b_k - a_k\|_\infty^2 + 4d^2 \rho^2 c_0^2 \right] \\
&\leq \frac{2}{\nu} \sum_k \|b_k - a_k\|_\infty^2 + \frac{1}{\nu} 4d^3 \rho^2 c_0^2
\end{aligned}$$

Since $\text{Im}(a_{k,j} + a_{j,k}) = 0$ for all $k, j = 1, 2, \dots, d$ we have

$$\begin{aligned} m_\rho &= \frac{1}{4\nu} \sum_{k=1}^d \left[f_k + \text{Im}(c_{k,\rho} - b_{k,\rho}) \right] \\ &= \frac{1}{4\nu} \sum_{k=1}^d \left[f_k + \text{Im}(c_k - b_k) + \rho \text{Im} \left(\sum_{j=1}^d a_{kj} D_j \psi + \sum_{i=1}^d a_{ik} D_i \psi \right) \right] \\ &= m. \end{aligned}$$

Thus,

$$\begin{aligned} \|(\text{Re } a_{0,\rho} - m_\rho)^-\|_\infty &\leq \|(\text{Re } a_0 - m)^-\|_\infty + \|\rho^2 \sum_{i,k=1}^d a_{ik} D_i \psi D_k \psi + \rho \sum_{k=1}^d b_k D_k \psi - \rho \sum_{k=1}^d c_k D_k \psi\|_\infty \\ &= \|(\text{Re } a_0 - m)^-\|_\infty + \rho^2 d^2 c_0 + \rho^2 + c_0 d^2. \end{aligned}$$

This equality together with (6.26) prove (6.19). \square

Combining Theorem 5.5 with Lemma 6.11 and Corollary 6.10 we can finally prove that Gaussian bounds are satisfied by those evolution families associated with families of uniform elliptic operators of the form (6.9).

Theorem 6.12. *Let V be W -invariant and satisfy (4.4). If the assumptions of Lemma 6.7 or those of Lemma 6.8 are satisfied, then \mathcal{U}_V satisfies Gaussian bounds. More precisely we have*

$$(6.27) \quad (\mathcal{U}_V(t, s)f)(x) = \int_{\mathbb{R}^d} \Gamma_V(t, s, x, y) f(y) dy$$

where

$$(6.28) \quad |\Gamma_V(t, s, x, y)| \leq c_0 e^{\omega_0(t-s)} (t-s)^{-\frac{n}{2}} \exp\left(-\frac{\beta_0}{4} \frac{|x-y|^2}{t-s}\right)$$

for a.e. $x \in \mathbb{R}^d$, all $(t, s) \in \Delta$, and all $f \in L^2(\mathbb{R}^d)$, where c, ω_0, n and β_0 are positive constants that depend only on C_G, N, d, ν, c_0 and on the constant β defined in (5.3).

REFERENCES

- [AB94] W. Arendt and A.V. Bukhvalov. Integral representations of resolvents and semigroups. *Forum Math.*, 6:111–135, 1994.
- [ABHN01] W. Arendt, C.J.K. Batty, M. Hieber, and F. Neubrander. *Vector-Valued Laplace Transforms and Cauchy Problems*, volume 96 of *Monographs in Mathematics*. Birkhäuser, Basel, 2001.
- [ACFP07] W. Arendt, R. Chill, S. Fornaro, and C. Poupaud. L^p -maximal regularity for non-autonomous evolution equations. *J. Differ. Equ.*, 237:1–26, 2007.
- [AD17] W. Arendt and D. Dier. Reaction-diffusion systems governed by non-autonomous forms. *J. Differ. Equ.*, 264:6362–6379, 2017.
- [ADF17] W. Arendt, D. Dier, and S. Fackler. J. L. Lions' problem on maximal regularity. *Arch. Math.*, 109:59–72, 2017.
- [ADKF14] W. Arendt, D. Dier, and M. Kramar Fijavž. Diffusion in networks with time-dependent transmission conditions. *Appl. Math. Optim.*, 69:315–336, 2014.
- [ADLO14] W. Arendt, D. Dier, H. Laasri, and E.M. Ouhabaz. Maximal regularity for evolution equations governed by non-autonomous forms, 2014.

- [ADO14] W. Arendt, D. Dier, and E.M. Ouhabaz. Invariance of convex sets for non-autonomous evolution equations governed by forms. *J. London Math. Soc.*, 89:903–916, 2014.
- [AF03] R.A. Adams and J.J.F. Fournier. *Sobolev Spaces*. Elsevier, Amsterdam, 2003.
- [Are06] W. Arendt. Heat Kernels – Manuscript of the 9th Internet Seminar, 2006. (freely available at http://www.uni-ulm.de/fileadmin/website_uni_ulm/mawi.inst.020/arendt/downloads/internetseminar.pdf).
- [Aro67] D.G. Aronson. Bounds for the fundamental solution of a parabolic equation. *Bull. Amer. Math. Soc.*, 73:890–896, 1967.
- [Aro68] D.G. Aronson. Non-negative solutions of linear parabolic equations. *Ann. Sc. Norm. Super. Pisa, Cl. Sci.*, 22:607–694, 1968.
- [AtE97] W. Arendt and T. ter Elst. Gaussian estimates for second order elliptic operators with boundary conditions. *J. Operator Th.*, 38:87–130, 1997.
- [Bay12] F. Bayazit. Positive evolution families solving nonautonomous difference equations. *Positivity*, 16:653–684, 2012.
- [BB12] A. Bátkai and A. Bobrowski. On shape preserving semigroups. *Arch. Math.*, 98:37–48, 2012.
- [Bel88] J. von Below. Classical solvability of linear parabolic equations on networks. *J. Differ. Equ.*, 72:316–337, 1988.
- [BK13] G. Berkolaiko and P. Kuchment. *Introduction to Quantum Graphs*, volume 186 of *Math. Surveys and Monographs*. Amer. Math. Soc., Providence, RI, 2013.
- [BS73] F. Black and M. Scholes. The pricing of options and corporate liabilities. *J. Polit. Econ.*, 81:637–654, 1973.
- [Chu97] F.R.K. Chung. *Spectral Graph Theory*, volume 92 of *Reg. Conf. Series Math.* Amer. Math. Soc., Providence, RI, 1997.
- [Chu07] F. Chung. The heat kernel as the pagerank of a graph. *Proc. Natl. Acad. Sci. USA*, 104:19735–19740, 2007.
- [CKLS92] K.C. Chan, G.A. Karolyi, F.A. Longstaff, and A.B. Sanders. An empirical comparison of alternative models of the short-term interest rate. *J. Finance*, 47:1209–1227, 1992.
- [CL99] C. Cicone and Y. Latushkin. *Evolution semigroups in dynamical systems and differential equations.*, volume 70 of *Math. Surveys and Monographs*. Amer. Math. Soc., Providence, RI, 1999.
- [CM09] S. Cardanobile and D. Mugnolo. Parabolic systems with coupled boundary conditions. *J. Differ. Equ.*, 247:1229–1248, 2009.
- [Cou93] T. Coulhon. Iteration de Moser et estimation gaussienne du noyau de la chaleur. *J. Operator Th.*, 29:157–165, 1993.
- [Dan00] D. Daners. Heat kernel estimates for operators with boundary conditions. *Math. Nachr.*, 217:13–42, 2000.
- [Dav87] E.B. Davies. Explicit constants for Gaussian upper bounds on heat kernels. *Am. J. Math.*, 109:319–333, 1987.
- [Dav07] E.B. Davies. *Linear Operators And Their Spectra*. Cambridge Univ. Press, Cambridge, 2007.
- [Del99] T. Delmotte. Parabolic Harnack inequality and estimates of Markov chains on graphs. *Rev. Mat. Iberoam.*, 15:181–232, 1999.
- [DL92] R. Dautray and J.-L. Lions. *Mathematical Analysis and Numerical Methods for Science and Technology, Vol. 5*. Springer-Verlag, Berlin, 1992.
- [Dor05] B. Dorn. Semigroups for flows on infinite networks. Master’s thesis, Eberhard-Karls-Universität, Tübingen, 2005.
- [Ein08] M. Einemann. *Semigroup Methods in Finance*. PhD thesis, Universität Ulm, 2008.
- [EL16] O. El-Mennaoui and H. Laasri. On evolution equations governed by non-autonomous forms. *Arch. Math.*, 107:43–57, 2016.
- [EN00] K.-J. Engel and R. Nagel. *One-Parameter Semigroups for Linear Evolution Equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [Fac17] S. Fackler. J.-L. Lions’ problem concerning maximal regularity of equations governed by non-autonomous forms. *Ann. Inst. Henri Poincaré., Ann. Non Linéaire*, 34:699–709, 2017.
- [Fat83] H.O. Fattorini. *The Cauchy problem*, volume 18 of *Encyclopedia of Mathematics and its Applications*. Cambridge Univ. Press, Cambridge, 1983.
- [FOT10] M. Fukushima, Y. Oshima, and M. Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *Studies in Math.* de Gruyter, Berlin, 2010.
- [Gle15] D.F. Gleich. PageRank beyond the Web. *SIAM Review*, 57:321–363, 2015.

- [Hes93] S.L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *Rev. Financ. Studies*, 6:327–343, 1993.
- [Kuc04] P. Kuchment. Quantum graphs I: Some basic structures. *Waves Random Media*, 14:107–128, 2004.
- [Laa18] H. Laasri. Regularity properties for evolution families governed by non-autonomous forms. *Arch. Math.*, 111:187–201, 2018.
- [LE13] H. Laasri and O. El-Mennaoui. Stability for non-autonomous linear evolution equations with L^p -maximal regularity. *Czech. Math. J.*, 63:887–908, 2013.
- [LS99] G. Lumer and R. Schnaubelt. Local operator methods and time dependent parabolic equations on non-cylindrical domains. In *Evolution equations, Feshbach resonances, singular Hodge theory*, pages 58–130. Wiley-VCH, Berlin, 1999.
- [Maz85] V.G. Maz’ya. *Sobolev Spaces*. Springer-Verlag, Berlin, 1985.
- [MN91] P. Meyer-Nieberg. *Banach Lattices*. Universitext. Springer-Verlag, Berlin, 1991.
- [Mug07] D. Mugnolo. Gaussian estimates for a heat equation on a network. *Networks Het. Media*, 2:55–79, 2007.
- [Mug14] D. Mugnolo. *Semigroup Methods for Evolution Equations on Networks*. Underst. Compl. Syst. Springer-Verlag, Berlin, 2014.
- [MVV05] A. Manavi, H. Vogt, and J. Voigt. Domination of semigroups associated with sectorial forms. *J. Oper. Theory*, 54:9–25, 2005.
- [Nag86] R. Nagel, editor. *One-Parameter Semigroups of Positive Operators*, volume 1184 of *Lect. Notes Math.* Springer-Verlag, Berlin, 1986.
- [Nit12] R. Nittka. Projections onto convex sets and L^p -quasi-contractivity of semigroups. *Arch. Math.*, 98:341–353, 2012.
- [Ouh96] E.M. Ouhabaz. Invariance of closed convex sets and domination criteria for semigroups. *Potential Analysis*, 5:611–625, 1996.
- [Ouh04] E.M. Ouhabaz. Gaussian upper bounds for heat kernels of second-order elliptic operators with complex coefficients on arbitrary domains. *J. Operator. Theory.*, 51:335–360, 2004.
- [Ouh05] E.M. Ouhabaz. *Analysis of Heat Equations on Domains*, volume 30 of *Lond. Math. Soc. Monograph Series*. Princeton Univ. Press, Princeton, NJ, 2005.
- [Ouh15] E.M. Ouhabaz. Maximal regularity for non-autonomous evolution equations governed by forms having less regularity. *Arch. Math.*, 105:79–91, 2015.
- [Paz83] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*, volume 44 of *Appl. Math. Sci.* Springer-Verlag, New York, 1983.
- [Prö] R. Pröpper. *Heat kernels on metric measure spaces*. PhD thesis, Universität Ulm, (in preparation).
- [Rob91] D.W. Robinson. *Elliptic operators and Lie groups*. Oxford Univ. Press, Oxford, 1991.
- [Sho97] R.E. Showalter. *Monotone Operator in Banach Space and Partial Differential Equations*, volume 49 of *Math. Surveys and Monographs*. Amer. Math. Soc., Providence, RI, 1997.
- [Šil08] D.D. Šiljak. Dynamic graphs. *Nonlinear Analysis: Hybrid Systems*, 2:544–567, 2008.
- [SL15] A. Sani and H. Laasri. Evolution equations governed by Lipschitz continuous non-autonomous forms. *Czech. Math. J.*, 65:475–491, 2015.
- [Sla07] A. Slavík. *Product Integration, its History and Applications*. PhD thesis, Matfyzpress, Praha, 2007.
- [Tan79] H. Tanabe. *Equations of Evolution*, volume 6 of *Monogr. Studies Math.* Pitman, 1979.
- [Tho03] S. Thomaschewski. *Form methods for autonomous and non-autonomous Cauchy problems*. PhD thesis, Universität Ulm, 2003.
- [Voi92] J. Voigt. One-parameter semigroups acting simultaneously on different L_p -spaces. *Bull. Soc. Royale Sci. Liège*, 61:465–470, 1992.
- [VZ12] T. Vicsek and A. Zafeiris. Collective motion. *Phys. Reports*, 517:71–140, 2012.

HAFIDA LAASRI, ARBEITSGRUPPE FUNKTIONALANALYSIS, FAKULTÄT MATHEMATIK UND INFORMATIK, UNIVERSITÄT WUPPERTAL, 42119 WUPPERTAL, GERMANY
E-mail address: laasri@uni-wuppertal.de

DELIO MUGNOLO, LEHRGEBIET ANALYSIS, FAKULTÄT MATHEMATIK UND INFORMATIK, FERNUNIVERSITÄT IN HAGEN, 58084 HAGEN, GERMANY
E-mail address: delio.mugnolo@fernuni-hagen.de