On the Gradient of Harmonic Functions¹

Pisheng Ding

ABSTRACT. For a harmonic function u on a domain in \mathbb{R}^n , this note shows that $\|\nabla u\|$ is essentially determined by the geometry of level hypersurfaces of u. Specifically, the factor by which $\|\nabla u\|$ changes along a gradient flow is completely determined by the mean curvature of the level hypersurfaces intersecting the flow.

1 Introduction

Let u be a harmonic function on an open connected subset Ω of \mathbb{R}^n . Suppose, for simplicity, that u has no critical points in Ω . Starting at a point $p_0 \in \Omega$, follow the gradient flow to reach another point $p \in \Omega$. By how much has $\|\nabla u\|$ changed along the flow? This note seeks to generalize the answer indicated in [1] for the case n = 2; see Remark 2 in §3.

For $a \in u(\Omega)$, let S_a denote the level-*a* hypersurface $\{p \mid u(p) = a\}$. Orient S_a by prescribing the normal field $\mathbf{N} = \nabla u / \|\nabla u\|$. Define $H : \Omega \to \mathbb{R}$ by letting H(p) be the mean curvature of $S_{u(p)}$ at p.

Theorem: Let p be a point on the gradient flow originating from p_0 ; let p_0p be the arc on the flow between p_0 and p. Then,

$$\left\|\nabla u\left(p\right)\right\| = \left\|\nabla u\left(p_{0}\right)\right\| \exp\left(\left(n-1\right)\int_{\widehat{p_{0}p}}H\,ds\right),$$

where s denotes arc length along p_0p .

It turns out that the case n = 3 completely embodies the general case. For simplicity, we treat this case.

2 Preliminaries

We begin by considering the mean curvature H of level surfaces of a C^2 function f on a domain in \mathbb{R}^3 .

Suppose that 0 is a regular value of f and let S denote the level-0 set $\{p \mid f(p) = 0\}$. Let $\mathbf{N} = \nabla f / \|\nabla f\|$. For $p \in S$, let $T_p S$ denote the tangent plane of S at p. For each unit vector $\mathbf{v} \in T_p S$, let $\gamma_{\mathbf{v}}$ be the unit-speed parametrization of the normal section of S with $\gamma_{\mathbf{v}}(0) = p$ and $\gamma'_{\mathbf{v}}(0) = \mathbf{v}$, then the (signed) curvature $\kappa_p(\mathbf{v})$ of $\gamma_{\mathbf{v}}$ at p is defined by the equation $\gamma''_{\mathbf{v}}(0) = \kappa_p(\mathbf{v})\mathbf{N}(p)$. The mean curvature H(p) of S at p is the mean of $\kappa_p(\mathbf{v})$ with \mathbf{v} ranging over the unit circle in T_pS , i.e.,

$$H(p) := \frac{1}{2\pi} \int_{\mathbf{v} \in T_p S; \, \|\mathbf{v}\| = 1} \kappa_p(\mathbf{v}) d\sigma \,,$$

where $d\sigma$ is the arc length element of the circle $\|\mathbf{v}\| = 1$.

¹MSC2010: Primary 31B05; Secondary 53A07.

Key words: harmonic function, gradient, mean curvature, level hypersurface.

As $\gamma_{\mathbf{v}}(t) \in S$, the two vectors $\nabla f(\gamma_{\mathbf{v}}(t))$ and $\gamma'_{\mathbf{v}}(t)$ are orthogonal; hence

$$0 = \frac{d}{dt} \langle \nabla f(\gamma_{\mathbf{v}}(t)), \gamma_{\mathbf{v}}'(t) \rangle$$
$$= \left\langle \frac{d}{dt} \nabla f(\gamma_{\mathbf{v}}(t)), \gamma_{\mathbf{v}}'(t) \right\rangle + \left\langle \nabla f(\gamma_{\mathbf{v}}(t)), \gamma_{\mathbf{v}}''(t) \right\rangle$$

Note that

$$\langle \nabla f(\gamma_{\mathbf{v}}(0)), \gamma_{\mathbf{v}}''(0) \rangle = \kappa_p(\mathbf{v}) \| \nabla f(p) \|,$$

whereas

$$\left\langle \left. \frac{d}{dt} \right|_{t=0} \nabla f(\gamma_{\mathbf{v}}(t)), \ \gamma_{\mathbf{v}}'(0) \right\rangle = Q(p)(\mathbf{v}, \mathbf{v}),$$

where $Q(p): T_p \mathbb{R}^3 \times T_p \mathbb{R}^3 \to \mathbb{R}$ is the Hessian quadratic form, implying that

$$\kappa_p(\mathbf{v}) = -\frac{1}{\|\nabla f(p)\|} Q(p)(\mathbf{v}, \mathbf{v}).$$

To average $\kappa_p(\mathbf{v})$ over the unit circle, it suffices to average $Q(p)(\mathbf{v}, \mathbf{v})$. To that end, let $(\boldsymbol{\xi}, \boldsymbol{\eta})$ be an orthonormal basis for $T_p S$; then $(\boldsymbol{\xi}, \boldsymbol{\eta}, \mathbf{N})$ is an orthonormal basis for $T_p \mathbb{R}^3$. It is simple to verify that

$$\frac{1}{2\pi} \int_{\mathbf{v}\in T_pS; \, \|\mathbf{v}\|=1} Q(p)(\mathbf{v},\mathbf{v}) d\sigma = \frac{1}{2} \left(Q(p)\left(\boldsymbol{\xi},\boldsymbol{\xi}\right) + Q(p)\left(\boldsymbol{\eta},\boldsymbol{\eta}\right) \right).$$

Note that

$$Q(p) (\boldsymbol{\xi}, \boldsymbol{\xi}) + Q(p) (\boldsymbol{\eta}, \boldsymbol{\eta}) + Q(p) (\mathbf{N}, \mathbf{N}) = \operatorname{Tr} Q(p) = \Delta f(p).$$

Thus, we have

$$H(p) = \frac{Q(p) \left(\mathbf{N}, \mathbf{N}\right) - \Delta f(p)}{2 \left\|\nabla f(p)\right\|} .$$
(1)

3 Proof of Theorem

Let u be a harmonic function on an open connected subset Ω of \mathbb{R}^3 without critical points. Let $\mathbf{N} = \nabla u / ||\nabla u||$. Let $s \mapsto \varphi(s)$ be the unit-speed gradient flow originating from p_0 ; φ is such that

$$\varphi'(s) = \mathbf{N}(\varphi(s)) \text{ and } \varphi(0) = p_0.$$

Consider $g(t) := u(\varphi(t))$. Then,

$$\begin{aligned} g'(s) &= \langle \nabla u(\varphi(s)), \ \varphi'(s) \rangle = \| \nabla u(\varphi(s)) \| \,, \\ g''(s) &= \left\langle \frac{d}{ds} \nabla u(\varphi(s), \ \varphi'(s) \right\rangle + \left\langle \nabla u(\varphi(s)), \ \varphi''(s) \right\rangle. \end{aligned}$$

As $\|\varphi'\| \equiv 1$, the vectors φ' and φ'' , and hence ∇u and φ'' , are always orthogonal. Thus,

$$g''(s) = Q(\varphi(s))(\mathbf{N}, \mathbf{N}).$$

By (1) in §2,

$$H(\varphi(s)) = \frac{Q(\varphi(s))(\mathbf{N}, \mathbf{N})}{2 \|\nabla u(\varphi(s))\|} = \frac{1}{2} \frac{g''(s)}{g'(s)} = \frac{1}{2} \frac{d}{ds} \log g'(s)$$
$$= \frac{1}{2} \frac{d}{ds} \log \|\nabla u(\varphi(s))\|.$$
(2)

Integrating both sides yields the result.

Remarks:

1. If u is a harmonic function of two variables, the above analysis can be easily adapted to show that

$$\left\|\nabla u\left(p\right)\right\| = \left\|\nabla u\left(p_{0}\right)\right\| \exp\left(\int_{\widehat{p_{0}p}} \kappa \, ds\right),$$

where κ is the curvature (signed, according to choice made of **N**) of the level curves of u. This formula was suggested in [1] by a purely complexanalytic argument, as u is locally the real part of a holomorphic function. However, this argument does not apply when n > 2.

2. For n > 3, it suffices to notes that a level hypersurface of an *n*-variable C^2 function f has at p mean curvature

$$H(p) = \frac{Q(p) \left(\mathbf{N}, \mathbf{N}\right) - \Delta f(p)}{(n-1) \left\|\nabla f(p)\right\|} ,$$

where the factor 1/(n-1), the counterpart of the factor 1/2 in (1), stems from averaging a quadratic form on \mathbb{R}^n over a unit (n-2)-sphere. Omitting the obvious details, we conclude, as a generalization of (2), that, for a harmonic function u of n variables,

$$H = \frac{1}{(n-1)} D_{\mathbf{N}} \log \|\nabla u\| \quad .$$

from which the general case of the Theorem follows.

References

 R. P. Jerrard and L. A. Rubel, On the Curvature of the Level Lines of a Harmonic Function, *Proceedings of the American Mathematical Society*, Vol. 14, No. 1 (1963), pp.29-32.

Illinois State University, Normal, Illinois pding@ilstu.edu