

ISOMORPHISMS UP TO BOUNDED TORSION BETWEEN RELATIVE K_0 -GROUPS AND CHOW GROUPS WITH MODULUS

RYOMEI IWASA AND WATARU KAI

CONTENTS

0.	Introduction	1
1.	A presentation of Chow group with modulus	2
2.	A presentation of relative K_0 -group	4
3.	Adams decomposition	7
4.	Proof of Theorem 0.1	10
5.	Applications	14
	References	15

0. INTRODUCTION

The purpose of this note is to establish isomorphisms up to bounded torsion between relative K_0 -groups and Chow groups with modulus as defined in [BS17].

Theorem 0.1. *Let X be a separated regular noetherian scheme of dimension d and D an effective Cartier divisor on X . Assume that D has an affine open neighborhood in X . Then there exists a finite descending filtration F^* on $K_0(X, D)$ and, for each integer p , there exists a surjective group morphism*

$$\text{cyc}: \text{CH}^p(X|D) \twoheadrightarrow F^p K_0(X, D) / F^{p+1} K_0(X, D)$$

such that its kernel is $(p-1)!$ -torsion for some positive integer N depending only on p . Furthermore, the filtration F^ coincides with the gamma filtration on $K_0(X, D)$ up to $(d-1)!$ -torsion for some positive integer M depending only on d .*

The case $D = \emptyset$ is a classical theorem of Soulé [So85], which owes its origin to Grothendieck's Riemann-Roch type formula [SGA6]. The filtration F^* and the morphism cyc have been constructed in [Iw19] in a slightly weaker generality. The assumption that D has an affine open neighborhood is essential, see Example 4.8.

Let $\mathcal{S}(X|D)$ be the set of all closed subsets in X not meeting D and $\mathcal{S}(X|D, 1)$ the set of all closed subsets in $X \times \square^1$ satisfying the modulus condition along D . It follows easily from the definition of $\text{CH}^*(X|D)$ that there is an exact sequence

$$\text{colim}_{Y \in \mathcal{S}(X|D, 1)} \mathcal{Z}_Y^*(X \times \square^1) \longrightarrow \text{colim}_{Y \in \mathcal{S}(X|D)} \text{CH}_Y^*(X) \longrightarrow \text{CH}^*(X|D) \longrightarrow 0,$$

where $\mathcal{Z}_Y^*(-)$ is the group of cycles with supports in Y and $\text{CH}_Y^*(-)$ is the Chow group with supports in Y . The real content of this note is to establish an analogous exact sequence for K -groups. In the second section, as Theorem 2.2, we establish an exact sequence

$$\text{colim}_{Y \in \mathcal{S}(X|D, 1)} K_0^Y(X \times \square^1) \longrightarrow \text{colim}_{Y \in \mathcal{S}(X|D)} K_0^Y(X) \longrightarrow K_0(X, D) \longrightarrow 0.$$

Then, from the classical rational isomorphisms between K_0 -groups and Chow groups, we get a rational isomorphism between $K_0(X, D)$ and $\text{CH}^*(X|D)$. The estimate on torsion is obtained by using Adams decomposition.

Convention. All rings are noetherian and all schemes are separated noetherian. For a point v of a scheme X , we denote by $\kappa(v)$ the residue field of v .

1. A PRESENTATION OF CHOW GROUP WITH MODULUS

Chow groups with supports.

Notation 1.1. Let X be a scheme and p an integer.

- (1) We write $X^{(p)}$ for the set of all points of codimension p in X , i.e., points $v \in X$ whose closures in X have codimension p . We understand $X^{(p)} = \emptyset$ if $p < 0$.
- (2) For a closed subset Y of X , we define $\mathcal{Z}_Y^p(X)$ to be the free abelian group with the generators $[V]$, one for each $v \in X^{(p)} \cap Y$, with V being the closure of v in X . We write $\mathcal{Z}^p(X) = \mathcal{Z}_X^p(X)$.
- (3) For a closed subscheme D of pure codimension p in X , we write

$$[D] := \sum_{x_i \in D^{(0)}} \text{length}(\mathcal{O}_{D, x_i}) [D_i] \in \mathcal{Z}_D^p(X),$$

where D_i is the closure of x_i in X .

Construction 1.2. Let X be a scheme and p an integer. Let $w \in X^{(p-1)}$ and write W for its closure in X . For each $v \in X^{(p)} \cap W$, there exists a unique group morphism $\nu_v: \kappa(w)^\times \rightarrow \mathbb{Z}$ which sends $a \in \mathcal{O}_{W, v} \setminus \{0\}$ to the length of $\mathcal{O}_{W, v}/(a)$. For $f \in \kappa(w)^\times$, we define

$$\text{div}(f) := \sum_{v \in X^{(p)} \cap W} \nu_v(f) [V] \in \mathcal{Z}_W^p(X).$$

Definition 1.3. Let X be a scheme and p an integer. For a closed subset Y of X , we define

$$\text{CH}_Y^p(X) := \text{coker} \left(\bigoplus_{w \in X^{(p-1)} \cap Y} \kappa(w)^\times \xrightarrow{\text{div}} \mathcal{Z}_Y^p(X) \right).$$

We write $\text{CH}^p(X) = \text{CH}_X^p(X)$.

Definition 1.4. Let X be a topological space with irreducible components $\{X_i\}_{i \in I}$. We say that X is *unicodimensional* if $\text{codim}_X(V) = \text{codim}_{X_i}(V)$ for any $i \in I$ and any irreducible closed subset V of X_i . A scheme is uncodimensional if the underlying topological space is uncodimensional.

Lemma 1.5. *Let X be a uncodimensional catenary scheme, D a closed subscheme of pure codimension r in X and p an integer. Then D is uncodimensional and $D^{(p-r)} \subset X^{(p)}$. Furthermore, the inclusion $\iota: D \hookrightarrow X$ induces a group morphism*

$$\iota_*: \text{CH}_{D \cap Y}^{p-r}(D) \rightarrow \text{CH}_Y^p(X)$$

for any closed subset Y of X .

Proof. Let $\{X_i\}_{i \in I}$ (resp. $\{D_j\}_{j \in J}$) be the set of irreducible components of X (resp. D). Take $j \in J$ and $v \in D^{(p-r)} \cap D_j$. Then

$$\text{codim}_{D_j}(v) + \text{codim}_{X_i}(D_j) = \text{codim}_{X_i}(v) = \text{codim}_X(v)$$

for any $i \in I$ with $X_i \supset D_j$. Since $\text{codim}_{X_i}(D_j) = r$ regardless of the choices of i, j , we see that D is uncodimensional and that $\text{codim}_X(v) = \text{codim}_D(v) + r = p$. Hence, $D^{(p-r)} \subset X^{(p)}$. The last statement is immediate from this. \square

Lemma 1.6. *Let X be a uncodimensional catenary scheme, Y a closed subset of X , D an effective Cartier divisor on X and p an integer. Let $v \in X^{(p)} \cap Y$ whose closure V in X is not contained in D . Then $[V \times_X D] \in \mathcal{Z}_{D \cap Y}^p(D)$.*

Proof. It suffices to show that $\text{codim}_D(V \times_X D) = p$. First of all, note that $V \times_X D$ is an effective Cartier divisor on V , and thus it is of pure codimension 1 in V . It follows that $\text{codim}_X(V \times_X D) = p + 1$. Since X is catenary, we conclude that $\text{codim}_D(V \times_X D) = \text{codim}_X(V \times_X D) - \text{codim}_X(D) = p$. \square

Construction 1.7. Let X be a unidimensional catenary scheme, Y a closed subset of X , D an effective Cartier divisor on X and p an integer. We define a group morphism

$$\iota^*: \mathcal{Z}_Y^p(X) \rightarrow \text{CH}_{Y \cap D}^p(D)$$

as follows, where ι refers to the inclusion $D \hookrightarrow X$. For an integral closed subscheme V of codimension p in X whose support is in Y ,

$$\iota^*([V]) := \begin{cases} [V \times_X D] & \text{if } V \not\subseteq D \\ j_*[\mathcal{O}_X(D)|_V] & \text{if } V \subseteq D \end{cases} \in \text{CH}_{Y \cap D}^p(D),$$

where the first equation is well-defined by Lemma 1.6 and, for the second, j refers to the inclusion $V \hookrightarrow D$ and $j_*: \text{CH}^1(V) \rightarrow \text{CH}_{Y \cap D}^p(D)$ is the push-forward ensured by Lemma 1.5.

Remark 1.8. It is the classical fact that the morphism ι^* in Construction 1.7 factors through $\text{CH}_Y^p(X)$ if X is an algebraic scheme, cf., [Fu98, Chapter 2]. It would be true more generally, but we do not need such a result for our purpose.

Chow groups with modulus.

Notation 1.9. We set $\overline{\square}^1 := \text{Proj}(\mathbb{Z}[T_0, T_1])$ and let t be the rational coordinate T_0/T_1 . We write $\square^1 := \overline{\square}^1 \setminus (t = \{\infty\})$. For an integer q and a scheme X , we denote by $\iota_{X,q}$ (or simply by ι_q) the inclusion $X \hookrightarrow X \times \square^1$ defined by $t = q$.

Definition 1.10 (Binda-Saito). Let X be a scheme and D an effective Cartier divisor on X . Let W be a closed subset of $X \times \square^1$. Let \overline{W}^N be the normalization of the closure \overline{W} (with the reduced scheme-structure) of W in $X \times \overline{\square}^1$ and denote by ϕ_W the canonical morphism $\overline{W}^N \rightarrow \overline{W}$. We say that W satisfies the modulus condition along D if the following inequality of Cartier divisors on \overline{W}^N holds

$$\phi_W^*(D \times \overline{\square}^1) \leq \phi_W^*(X \times \{\infty\}).$$

Lemma 1.11. *Let X be a scheme and D an effective Cartier divisor on X . Let W be a closed subset of $X \times \square^1$ satisfying the modulus condition along D . Then any closed subset of W satisfies the modulus condition along D .*

Proof. The proof for [KP12, Proposition 2.4] works, noting that every integral morphism of schemes is closed (the theorem of Cohen-Seidenberg, [EGA2, (6.1.10)] or [AM69, Theorem 5.10]).

Here we give an alternative argument which might be useful later. First, since the modulus condition is a local condition, we may assume X is affine $X = \text{Spec}(A)$ and D is principal $D = (f)$. We give $\overline{W} \subset X \times \overline{\square}^1$ the reduced scheme structure and consider its restriction to the open subset $X \times (\overline{\square}^1 \setminus \{0\}) = \text{Spec}(A[1/t])$. Let $A[t^{-1}]/J$ be the coordinate ring of $\overline{W} \cap \text{Spec}(A[1/t])$. The modulus condition for W is equivalent to the condition that the element $1/tf$ in the ring of total quotients of $A[1/t]/J$ is integral over this ring, i.e., that there is a relation in $A[1/t]/J$ of the form

$$\frac{1}{t^n} + g_1 \frac{1}{t^{n-1}} + \cdots + g_n = 0$$

with $g_i \in f^i A[1/t]/J$. Now let $Y \subset W$ be any closed subset. Then we have the image of the above relation to the coordinate ring of $\overline{Y} \cap \text{Spec}(A[1/t])$. This implies the modulus condition for Y . \square

Notation 1.12. Let X be scheme, D an effective Cartier divisor on X and p an integer.

- (1) $\mathcal{S}(X|D)$ is the set of all closed subsets of X not meeting D .
- (2) $\mathcal{S}(X|D, 1)$ is the set of all closed subsets of $X \times \square^1$ satisfying the modulus condition along D .
- (3) $\mathcal{Z}^p(X|D)$ is the free abelian group with generators $[V]$, one for each $v \in X^{(p)}$ whose closure V does not meet D .
- (4) $\mathcal{Z}^p(X|D, 1)$ is the free abelian group with generators $[W]$, one for each $w \in (X \times \square^1)^{(p)}$ whose closure W is dominant over \square^1 and satisfies the modulus condition along D .

Remark 1.13. We remark that

$$\mathcal{Z}^p(X|D) = \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \mathcal{Z}_Y^p(X) \quad \text{and} \quad \mathcal{Z}^p(X|D, 1) \subset \operatorname{colim}_{Y \in \mathcal{S}(X|D, 1)} \mathcal{Z}_Y^p(X \times \square^1).$$

In the latter formula, the difference consists of cycles not dominant over \square^1 .

Definition 1.14. Let X be a unidimensional catenary scheme, D an effective Cartier divisor on X and p an integer. We define

$$\operatorname{CH}^p(X|D) := \operatorname{coker}\left(\mathcal{Z}^p(X|D, 1) \xrightarrow{\iota_0^* - \iota_1^*} \mathcal{Z}^p(X|D)\right),$$

where the morphisms ι_0^* and ι_1^* are well-defined by Lemma 1.6 and Remark 1.13.

Lemma 1.15. *Let X be a unidimensional catenary scheme, D an effective Cartier divisor on X , Y a closed subset of X not meeting D and p an integer. Then the canonical morphism*

$$\mathcal{Z}_Y^p(X) \rightarrow \operatorname{CH}^p(X|D)$$

factors through $\operatorname{CH}_Y^p(X)$.

Proof. Suppose given $w \in X^{(p-1)} \cap Y$ and $f \in \kappa(w)^\times$. We have to show that $\operatorname{div}(f) = 0$ in $\operatorname{CH}^p(X|D)$. Consider the Cartier divisor E on $W \times \square^1$ defined by $f + (1-f)t$. Then E gives an element $[E]$ in $\mathcal{Z}^p(X|D, 1)$ and $\iota_0^*[E] - \iota_1^*[E] = \operatorname{div}(f)$ in $\mathcal{Z}^p(X|D)$. This prove the lemma. \square

Proposition 1.16. *Let X be a unidimensional catenary scheme, D an effective Cartier divisor on X and p an integer. Then the sequence*

$$\operatorname{colim}_{Y \in \mathcal{S}(X|D, 1)} \mathcal{Z}_Y^p(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{CH}_Y^p(X) \xrightarrow{\epsilon} \operatorname{CH}^p(X|D) \longrightarrow 0$$

is exact. Here, ι_0^, ι_1^* are the morphisms defined in Construction 1.7, and ϵ is the canonical morphism as in Lemma 1.15.*

Proof. We only have to show that the composite $\epsilon \circ (\iota_0^* - \iota_1^*)$ is zero. Let $Y \in \mathcal{S}(X|D, 1)$ and $v \in (X \times \square^1)^{(p)} \cap Y$. Then the closure V of v satisfies the modulus condition along D by Lemma 1.11. If $v \notin X \times \{0, 1\}$, then it is immediate from the definition of $\operatorname{CH}^p(X|D)$ that $(\epsilon \circ (\iota_0^* - \iota_1^*))([V]) = 0$. If $v \in X \times \{0, 1\}$, say $v \in X \times \{0\}$, then $\iota_1^*[V] = 0$ and $\iota_0^*[V] = j_*[\mathcal{O}_{X \times \square^1}(X)|_V] = 0$ since X is a principal divisor in $X \times \square^1$. This completes the proof. \square

2. A PRESENTATION OF RELATIVE K_0 -GROUP

For a scheme X , we denote by $K(X)$ Thomason-Trobaugh's K -theory spectrum [TT90, 3.1]. The spectrum $K(X)$ is contravariant functorial in X .

Definition 2.1. Let X be a scheme.

- (1) Let D be a closed subscheme of X . We define $K(X, D)$ to be the homotopy fiber of the canonical morphism $K(X) \rightarrow K(D)$. For an integer n , we write $K_n(X, D) = \pi_n K(X, D)$.
- (2) Let Y be a closed subset of X . We define $K^Y(X)$ to be the homotopy fiber of the canonical morphism $K(X) \rightarrow K(X \setminus Y)$. For an integer n , we write $K_n^Y(X) = \pi_n K^Y(X)$.

The goal of this section is to prove the following theorem.

Theorem 2.2. *Let X be a regular scheme and D an effective Cartier divisor on X . Assume that D admits an affine open neighborhood in X . Then the sequence*

$$\operatorname{colim}_{Y \in \mathcal{S}(X|D, 1)} K_0^Y(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \operatorname{colim}_{Y \in \mathcal{S}(X|D)} K_0^Y(X) \xrightarrow{\epsilon} K_0(X, D) \longrightarrow 0$$

is exact. Here, ϵ denotes the obvious morphism.

The surjectivity of ϵ has been observed in [Iw19].

Lemma 2.3. *Let X be a scheme and D a closed subscheme of X . Assume that D has an affine open neighborhood in X . Then the canonical morphism*

$$\epsilon: \operatorname{colim}_{Y \in \mathcal{S}(X|D)} K_0^Y(X) \rightarrow K_0(X, D)$$

is surjective.

Proof. Let U be an affine open neighborhood of D in X . By the localization theorem [TT90, 7.4], the sequence

$$K_0^{X \setminus U}(X) \longrightarrow K_0(X, D) \longrightarrow K_0(U, D)$$

is exact. Hence, we may replace X by U and reduce to the case X is affine. Then the result follows from [Iw19, Lemma 3.4]. \square

The rigidity.

Lemma 2.4. *Let X be a regular scheme and D an effective Cartier divisor. Let $Y \in \mathcal{S}(X|D, 1)$ and denote its closure in $X \times \overline{\square}^1$ by \overline{Y} . Assume that D admits an affine open neighborhood in X . Then the two morphisms*

$$\iota_0^*, \iota_1^*: K_0^{\overline{Y}}(X \times \overline{\square}^1) \rightarrow K_0(X, D)$$

coincide.

Proof. First of all, let us fix notation for morphisms of schemes:

$$\begin{array}{ccccc} X & \begin{array}{c} \xrightarrow{\iota_0} \\ \xrightarrow{\iota_1} \end{array} & X \times \overline{\square}^1 & \xrightarrow{q} & X & \xleftarrow{i} & D \\ & & \downarrow p & & & & \\ & & \overline{\square}^1 & & & & \end{array}$$

where p, q are the canonical projections and i is the canonical inclusion.

According to [Iw19, Theorem 3.1], $K_0(X, D)$ is generated by triples (P, α, Q) where P, Q are perfect complexes of X and α is a quasi-isomorphism $Li^*P \xrightarrow{\sim} Li^*Q$. The morphism

$$\iota_a^*: K_0^{\overline{Y}}(X \times \overline{\square}^1) \rightarrow K_0(X, D) \quad a \in \{0, 1\}$$

sends $[P]$ to $[(L\iota_a^*P, 0, 0)]$, where P is a perfect complex of $X \times \overline{\square}^1$ whose support lies in \overline{Y} . Since X is regular, $K_0^{\overline{Y}}(X \times \overline{\square}^1)$ is generated by coherent $\mathcal{O}_{\overline{Y}}$ -modules. Hence, it suffices to show that, for any coherent $\mathcal{O}_{\overline{Y}}$ -modules \mathcal{F} ,

$$[(L\iota_0^*\mathcal{F}, 0, 0)] = [(L\iota_1^*\mathcal{F}, 0, 0)]$$

in $K_0(X, D)$. In the sequel, we denote by \mathcal{F} a coherent $\mathcal{O}_{\overline{Y}}$ -module.

First calculation in $K_0(X, D)$. Recall that we have fixed a rational coordinate t of $\overline{\square}^1$. We denote by $\mathcal{O}(-1)$ the invertible sheaf on $\overline{\square}^1$ generated by t . We write j_0 for the canonical inclusion $\mathcal{O}(-1) \rightarrow \mathcal{O}_{\overline{\square}^1}$ sending t to t , and write j_1 for the inclusion $\mathcal{O}(-1) \rightarrow \mathcal{O}_{\overline{\square}^1}$ sending t to $t - 1$. Then we have an exact triangle

$$Lp^*\mathcal{O}(-1) \otimes_{X \times \overline{\square}^1}^L \mathcal{F} \xrightarrow{p^*j_a} \mathcal{F} \longrightarrow \iota_{a*}L\iota_a^*\mathcal{F} \xrightarrow{+}$$

of perfect complexes of $X \times \overline{\square}^1$ for $a \in \{0, 1\}$. Consequently,

$$[(L\iota_a^*\mathcal{F}, 0, 0)] = -Rq_*[(Lp^*\mathcal{O}(-1) \otimes_{X \times \overline{\square}^1}^L \mathcal{F}, p^*j_a, \mathcal{F})]$$

in $K_0(X, D)$. We set $\theta := (t-1)/t$. Then we have a commutative diagram (the vertical arrow is defined after restricting to $\overline{\square^1} \setminus \{0\}$)

$$\begin{array}{ccc} & & \mathcal{O}_{\overline{\square^1}} \\ & \nearrow^{j_0} & \vdots \\ \mathcal{O}(-1) & & \text{multiplication by } \theta \\ & \searrow_{j_1} & \vdots \\ & & \mathcal{O}_{\overline{\square^1}} \end{array}$$

Since Y satisfies the modulus condition, the multiplication by θ on $\mathcal{O}_{\overline{Y}}$ -modules makes sense in a neighborhood of $\overline{Y} \cap (D \times \overline{\square^1})$. It follows from the above diagram that

$$[(Lp^* \mathcal{O}(-1) \otimes_{X \times \overline{\square^1}}^L \mathcal{F}, p^* j_0, \mathcal{F})] + [(\mathcal{F}, \theta, \mathcal{F})] = [(Lp^* \mathcal{O}(-1) \otimes_{X \times \overline{\square^1}}^L \mathcal{F}, p^* j_1, \mathcal{F})].$$

Hence, it remains to show that

$$Rq_*[(\mathcal{F}, \theta, \mathcal{F})] = 0$$

in $K_0(X, D)$.

Adic filtration on \mathcal{F} . Let $\text{Fil}^* \mathcal{F}$ be the adic filtration on \mathcal{F} with respect to the ideal defining $D \times \{\infty\}$ in $X \times \overline{\square^1}$. Since $(1/t) \text{Fil}^l \mathcal{F} \subset \text{Fil}^{l+1} \mathcal{F}$ in a neighborhood of $\overline{Y} \cap (D \times \overline{\square^1})$, we see that

$$[(\text{Fil}^l \mathcal{F} / \text{Fil}^{l+1} \mathcal{F}, \theta, \text{Fil}^l \mathcal{F} / \text{Fil}^{l+1} \mathcal{F})] = 0$$

for all $l \geq 0$. Hence, we are reduced to showing that $Rq_*[(\text{Fil}^l \mathcal{F}, \theta, \text{Fil}^l \mathcal{F})] = 0$ for some $l \geq 0$. We show that θ acts on $Li^* Rq_* \text{Fil}^l \mathcal{F}$ as the identity for sufficiently large l .

Take an affine open neighborhood U of D in X such that $\overline{Y}_U := \overline{Y} \times_X U$ misses $X \times \{0, 1\}$ and that the restriction $q|_{\overline{Y}_U} : \overline{Y}_U \rightarrow U$ is finite. We set some notation:

$$A = \mathcal{O}(U) \quad A/I = \mathcal{O}(D) \quad A[1/t]/J = \mathcal{O}(\overline{Y}_U) \quad M = \mathcal{F}(\overline{Y}_U)$$

The filtration $\text{Fil}^* \mathcal{F}$ on \mathcal{F} descends to a filtration $\text{Fil}^* M$ on M , which is identified with the $(I, 1/t)$ -adic filtration. Observe that $Li^* Rq_* \text{Fil}^l \mathcal{F} = (Li^* \text{Fil}^l M)^\sim$.

We claim that there exists $n \geq 0$ such that $\text{Fil}^{l+1} M = I \text{Fil}^l M$ and $H_k(Li^* \text{Fil}^l M) = 0$ for all $l \geq n$ and $k > 0$. If we admit the claim, then

$$Li^* \text{Fil}^l M = \text{Fil}^l M / I \text{Fil}^l M = \text{Fil}^l M / \text{Fil}^{l+1} M$$

on which we know that θ acts as the identity. The claim is a local question on $\text{Spec } A$, and thus we may assume that I is principal, $I = (f)$ with $0 \neq f \in A$. By the modulus condition, we have a relation in $A[1/t]/J$ of the form

$$\frac{1}{t^n} + g_1 \frac{1}{t^{n-1}} + \cdots + g_{n-1} \frac{1}{t} + g_n = 0$$

for some $n \geq 0$ and $g_k \in f^k A[1/t]/J$ with $1 \leq k \leq n$. Repeated application of this relation gives

$$\text{Fil}^l M = f^l M + \frac{f^{l-1}}{t} M + \cdots + \frac{f^{l-(n-1)}}{t^{n-1}} M$$

for $l \geq n$. In particular, $\text{Fil}^{l+1} M = f \text{Fil}^l M$. Furthermore, since the f -power torsion of M has a bounded exponent, $\text{Fil}^l M = f^{l-n} \text{Fil}^n M$ has no f -power torsion for $l \gg n$. This proves the claim. \square

Corollary 2.5. *Under the situation in Theorem 2.2, $\epsilon \circ (\iota_0^* - \iota_1^*) = 0$.*

Proof. Since X is regular, the restriction morphism

$$K_0^{\overline{Y}}(X \times \overline{\square^1}) \rightarrow K_0^Y(X \times \square^1)$$

is surjective, where $Y \in \mathcal{S}(X|D, 1)$ and \overline{Y} is the closure of Y in $X \times \overline{\square^1}$. Hence, the result follows from Lemma 2.4. \square

End of the proof.

Lemma 2.6. *Let X be a scheme and D an effective Cartier divisor on X admitting an affine open neighborhood in X . Suppose we are given $Z \in \mathcal{S}(X|D)$ and $\alpha \in K_1(X \setminus Z)$ whose restriction to $K_1(D)$ is zero. Then there exist $W \in \mathcal{S}(X|D, 1)$ and $\beta \in K_1(X \times \square^1 \setminus W)$ such that*

$$\beta|_{t=0} - \beta|_{t=1} = \alpha \quad \text{in} \quad \operatorname{colim}_{Y \in \mathcal{S}(X|D)} K_1(X \setminus Y).$$

Proof. By enlarging Z if necessarily, we may assume that $X \setminus Z$ is affine. Set $U = X \setminus Z$. Take a representative of α in $\operatorname{GL}_n(U)$, which we also denote by α . By our assumption, the restriction of α to matrices over D is in the group $E_m(D)$ of elementary matrices for some $m \geq n$. Take a lift $\alpha' \in E_m(U)$ of $\alpha|_D$. Then $\alpha = \alpha' + \epsilon$ in $\operatorname{GL}_m(U)$, where ϵ is a matrix whose entries are all in the ideal I defining D . We define an $(m \times m)$ -matrix over $U \times \square^1$ by

$$\alpha(t) := \alpha' + (1-t)\epsilon.$$

Then the determinant $\det(\alpha(t))$ is an admissible polynomial for D in the sense [BS17, §4], and thus its zero locus $W = V(\det(\alpha(t)))$ satisfies the modulus condition along D . Lastly, by definition, $\alpha(t)$ gives an element β in $K_1(X \times \square^1 \setminus W)$ and it satisfies the desired formula. \square

Proof of Theorem 2.2. By Lemma 2.3 and Corollary 2.5, it remains to show the exactness at the middle term. Suppose we are given $Z \in \mathcal{S}(X|D)$ and $\alpha \in K_0^Z(X)$ which dies in $K_0(X, D)$ along the obvious morphism. Watch the commutative diagram with exact rows

$$\begin{array}{ccccc} K_1(X \setminus Z, D) & \longrightarrow & K_1(X \setminus Z) & \longrightarrow & K_1(D) \\ & & \downarrow & & \downarrow \\ K_1(X \setminus Z, D) & \longrightarrow & K_0^Z(X) & \longrightarrow & K_0(X, D). \end{array}$$

It follows that there exists a lift $\alpha' \in K_1(X \setminus Z)$ of α along the boundary morphism whose restriction to $K_1(D)$ is zero. Hence, by Lemma 2.6, we find an element β' in the group at the left upper corner of the commutative diagram

$$\begin{array}{ccccc} \operatorname{colim}_{Y \in \mathcal{S}(X|D, 1)} K_1(X \times \square^1 \setminus Y) & \xrightarrow{\iota_0^* - \iota_1^*} & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} K_1(X \setminus Y) & & \\ \downarrow & & \downarrow & & \\ \operatorname{colim}_{Y \in \mathcal{S}(X|D, 1)} K_0^Y(X \times \square^1) & \xrightarrow{\iota_0^* - \iota_1^*} & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} K_0^Y(X) & \xrightarrow{\epsilon} & K_0(X, D) \end{array}$$

such that $\iota_0^* \beta' - \iota_1^* \beta' = \alpha'$ in the upper middle group. This proves the exactness of the lower sequence. \square

3. ADAMS DECOMPOSITION

Notation 3.1. For $i \geq 0$, we set

$$w_i = \begin{cases} 1 & \text{if } i \text{ is zero} \\ 2 & \text{if } i \text{ is odd} \\ \text{denominator of } |B_{i/2}|/2i & \text{if } i \text{ is positive even} \end{cases}$$

where B_n denotes the n -th Bernoulli number.

Lemma 3.2.

- (i) *If a prime p divides w_i , then $(p-1)$ divides i . The converse is true if i is even.*
- (ii) *For $i \geq 1$ and for N large enough than i ,*

$$w_i = \operatorname{gcd}_{k \geq 2} k^N (k^i - 1).$$

Proof. See [MS74, Appendix B]. \square

Definition 3.3. Let N be a positive integer. We say that a morphism $f: A \rightarrow B$ of abelian groups is an N -monomorphism (resp. N -epimorphism) if the kernel (resp. cokernel) of f is killed by N . We say that f is an N -isomorphism if it is an N -monomorphism and an N -epimorphism.

Proposition 3.4. Let A be an abelian group equipped with endomorphisms ψ^k for $k > 0$ which commute with each other. Suppose that there is a finite filtration

$$A = F^i \supset F^{i+1} \supset \dots \supset F^j \supset F^{j+1} = 0$$

with $0 \leq i \leq j$ consisting of subgroups preserved by ψ^k such that ψ^k acts on F^p/F^{p+1} by the multiplication by k^p . For $p \geq 0$, let $A^{(p)}$ be the subgroup of A consisting of elements $x \in A$ such that $\psi^k x = k^p x$ for all $k > 0$. Then:

(i) For $i \leq p \leq j$, $(\prod_{q=i}^{p-1} w_{p-q})A^{(p)}$ is in F^p and the induced morphism

$$\prod_{q=i}^{p-1} w_{p-q}: A^{(p)} \rightarrow F^p/F^{p+1}$$

is a $(\prod_{q=i}^j w_{|p-q|})$ -isomorphism.

(ii) The canonical morphism

$$\bigoplus_{p=i}^j A^{(p)} \rightarrow A$$

is a $(\prod_{i \leq q, p \leq j} w_{|p-q|})$ -isomorphism.

Proof. We follow [So85, 2.8]. Take integers A_{pqk} ($p \neq q$) so that

$$w_{|p-q|} = \sum A_{pqk}(k^p - k^q).$$

We fix $i \leq p \leq j$. The morphism

$$\Phi_p := \prod_{q=i}^{p-1} \left(\sum_{k>1} A_{pqk}(\psi^k - k^q) \right)$$

sends A to F^p , and on $A^{(p)}$ it is the multiplication by $\prod_{q=i}^{p-1} w_{p-q}$. On the other hand, the morphism

$$\Psi_p := \prod_{q=p+1}^j \left(\sum_{k>1} A_{pqk}(\psi^k - k^q) \right)$$

sends F^p to $A^{(p)}$, and modulo F^{p+1} it is the multiplication by $\prod_{q=p+1}^j w_{q-p}$. Moreover, this morphism kills F^{p+1} . To summarize, $(\prod_{q=i}^{p-1} w_{p-q})A^{(p)}$ is in F^p and the induced morphism

$$\prod_{q=i}^{p-1} w_{p-q}: A^{(p)} \rightarrow F^p/F^{p+1}$$

is a $(\prod_{q=i}^j w_{|p-q|})$ -isomorphism.

Next, we prove (ii). Let us consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{p=i+1}^q A^{(p)} \cap F^{i+1} & \longrightarrow & \bigoplus_{p=i}^q A^{(p)} & \longrightarrow & \left(\bigoplus_{p=i+1}^q A^{(p)} / A^{(p)} \cap F^{i+1} \right) \oplus A^{(i)} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F^{i+1} & \longrightarrow & A & \longrightarrow & A/F^{i+1} \longrightarrow 0 \end{array}$$

with exact rows. The left vertical arrow is a $(\prod_{i+1 \leq p, q \leq j} w_{|p-q|})$ -isomorphism by induction. Since Φ_{i+1} sends A to F^{i+1} and it is the multiplication by w_{p-i} on $A^{(p)}$, $\bigoplus_{p=i+1}^q A^{(p)} / A^{(p)} \cap F^{i+1}$ is killed by

$\prod_{p=i+1}^j w_{p-i}$. Combining it with (i), we see that the right vertical arrow is a $(\prod_{p=i}^j w_{p-i})^2$ -isomorphism. Consequently, the middle vertical arrow is a $(\prod_{i \leq p, q \leq j} w_{|p-q|})$ -isomorphism. \square

Example I: γ -filtration. We refer to [AT69] for the definition of (non-unital) special λ -rings, γ -filtrations and Adams operations.

Lemma 3.5. *Let I be a non-unital special λ -ring. Assume that the γ -filtration on I is finite. Then the γ -filtration on I together with the Adams operations satisfies the condition of Proposition 3.4.*

Proof. This follows from [loc. cit., Proposition 5.3]. \square

Definition 3.6. Let X be a scheme. We define

$$\tilde{K}(X) := \mathbb{H}_{\text{Zar}}(X, B\text{GL}^+)$$

the global sections over X of a Zariski-fibrant replacement of (a functorial model of) $B\text{GL}^+$. Let Y be a closed subset of X and D a closed subscheme of X . We define $\tilde{K}^Y(X, D)$ to be the iterated homotopy fiber of the square

$$\begin{array}{ccc} \tilde{K}(X) & \longrightarrow & \tilde{K}(D) \\ \downarrow & & \downarrow \\ \tilde{K}(X \setminus Y) & \longrightarrow & \tilde{K}(D \setminus Y). \end{array}$$

For a non-negative integer n , we write $\tilde{K}_n^Y(X, D) := \pi_n \tilde{K}^Y(X, D)$. When $D = \emptyset$ (resp. $Y = X$), we write $\tilde{K}^Y(X)$ (resp. $\tilde{K}(X, D)$) for $\tilde{K}^Y(X, D)$.

Lemma 3.7. *Let X be a scheme, Y a closed subset of X and D a closed subscheme of X . Then $\tilde{K}_n^Y(X, D)$ is naturally a special λ -ring for each $n \geq 0$. The first grading of the γ -filtration is*

$$F_\gamma^1 \tilde{K}_n^Y(X, D) / F_\gamma^2 \tilde{K}_n^Y(X, D) \simeq \begin{cases} \mathbb{H}_{\text{Zar}, Y}^1((X, D), \mathcal{O}^\times) & n = 0 \\ \mathbb{H}_{\text{Zar}, Y}^0((X, D), \mathcal{O}^\times) & n = 1 \\ 0 & n > 1 \end{cases}$$

where $\mathbb{H}_{\text{Zar}, Y}^*(X, D, \mathcal{O}^\times)$ denotes the homology of the iterated homotopy fiber of the square

$$\begin{array}{ccc} \mathbb{H}_{\text{Zar}}(X, \mathcal{O}^\times) & \longrightarrow & \mathbb{H}_{\text{Zar}}(D, \mathcal{O}^\times) \\ \downarrow & & \downarrow \\ \mathbb{H}_{\text{Zar}}(X \setminus Y, \mathcal{O}^\times) & \longrightarrow & \mathbb{H}_{\text{Zar}}(D \setminus Y, \mathcal{O}^\times). \end{array}$$

Proof. Refer to [Le97, Corollary 5.6] for the fact $\tilde{K}_n^Y(X, D)$ is a special λ -ring. The first grading is calculated by the determinant $\det: B\text{GL}^+ \rightarrow B\mathcal{O}^\times$. Indeed, we have $\gamma^1 + \gamma^2 + \dots = 0$ in $SK_*^Y(X, D) = \mathbb{H}_{\text{Zar}, Y}^{-*}((X, D), B\text{SL}^+)$ as in [So85, p524]¹, and thus $F_\gamma^2 SK_*^Y(X, D) = SK_*^Y(X, D)$. \square

Example II: coniveau filtration.

Definition 3.8. Let X be a scheme and Y a closed subset of X . For each $p \geq 0$, we define

$$F^p K_0^Y(X) := \text{colim}_Z \text{image}(K_0^Z(X) \rightarrow K_0^Y(X))$$

where Z runs over all closed subset of Y whose codimension in X is greater or equal to p . We call the filtration the *coniveau filtration*. We write $\text{Gr}^p K_0^Y(X)$ for the p -th grading of the coniveau filtration.

¹In [So85, Théorème 4], it is asserted that $K_0(X) = H^0(X, \mathbb{Z}) \oplus \text{Pic}(X) \oplus F_\gamma^2 K_0(X)$, but it is not true. The proof shows $\text{Gr}_\gamma^1 K_0(X) \simeq \text{Pic}(X)$, but $\text{Pic}(X)$ may not split.

Lemma 3.9. *Let X be a regular scheme of dimension d and Y a closed subset of X of codimension p . Then the coniveau filtration*

$$K_0^Y(X) = F^p \supset F^{p+1} \supset \dots \supset F^d \supset F^{d+1} = 0$$

together with the Adams operations satisfies the condition of Proposition 3.4.

Proof. First, we prove the case Y is a closed point of X . Let j be the inclusion $Y \hookrightarrow X$ and j_* the pushforward $K_0(Y) \rightarrow K_0^Y(X)$. Then, by [So85, Théorème 3], we have $\psi^k \circ j_* = k^d(j_* \circ \psi^k)$. Since ψ^k is the identity on $K_0(Y) \simeq \mathbb{Z}$ and j_* is an isomorphism, we conclude that ψ^k acts by the multiplication by k^d on $K_0^Y(X)$. This proves the case Y is a closed point.

We prove the remaining case by descending induction on p . We have seen the case $p = d$. Let $p < d$. If $q > p$, then the canonical morphism

$$\operatorname{colim}_Z \operatorname{Gr}^q K_0^Z(X) \rightarrow \operatorname{Gr}^q K_0^Y(X),$$

where Z runs over all closed subsets of Y whose codimension in X is greater or equal to $p + 1$, is surjective. By induction, the Adams operation ψ^k acts by the multiplication by k^q on the left term, and so on the right. It remains to show that the Adams operation ψ^k acts by the multiplication by k^p on $\operatorname{Gr}^p K_0^Y(X) = K_0^Y(X)/F^{p+1}K_0^Y(X)$. This follows from the exact sequence [GS87, Lemma 5.2]

$$0 \longrightarrow F^{p+1}K_0^Y(X) \longrightarrow K_0^Y(X) \longrightarrow \bigoplus_{y \in Y \cap X^{(p)}} K_0^y(X_y) \longrightarrow 0.$$

Note that the Adams operations act on the sequence and we have seen that ψ^k acts on the right term by the multiplication by k^p . This completes the proof. \square

Example III: relative coniveau filtration.

Definition 3.10. Let X be a scheme and D a closed subscheme of X . For each $p \geq 0$, we define

$$F^p K_0(X, D) := \operatorname{colim}_Z \operatorname{image}(K_0^Z(X) \rightarrow K_0(X, D))$$

where Z runs over all closed subset in X not meeting D of codimension greater or equal to p . We call the filtration the *relative coniveau filtration*. We write $\operatorname{Gr}^p K_0(X, D)$ for the p -th grading of the relative coniveau filtration.

Lemma 3.11. *Let X be a scheme of dimension d with an ample family of line bundles and D a closed subscheme of X . Assume that $X \setminus D$ is regular and that D has an affine open neighborhood in X . Then the relative coniveau filtration*

$$K_0(X, D) = F^0 \supset F^1 \supset \dots \supset F^d \supset F^{d+1} = 0$$

together with the Adams operations satisfies the condition of Proposition 3.4.

Proof. By Lemma 2.3, $F^0 K_0(X, D) = K_0(X, D)$. By definition, the canonical morphism

$$\operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^p K_0^Y(X) \rightarrow \operatorname{Gr}^p K_0(X, D)$$

is surjective. The morphism is compatible with the Adams operations, and thus the result follows from Lemma 3.9. \square

4. PROOF OF THEOREM 0.1

Cycle class morphisms with supports.

Definition 4.1. Let X be a regular scheme, Y a closed subset of X and p an integer. The *cycle class morphism* is a group morphism

$$\operatorname{cyc}: \mathcal{Z}_Y^p(X) \rightarrow K_0^Y(X)$$

defined by sending the closure V of a point $v \in X^{(p)} \cap Y$ to \mathcal{O}_V .

Theorem 4.2 (Gillet-Soulé). *Let X be a regular scheme, Y a closed subset of X and p an integer. Then the cycle class morphism induces a surjective group morphism*

$$\text{cyc}: \text{CH}_Y^p(X) \rightarrow F^p K_0^Y(X)/F^{p+1} K_0^Y(X)$$

and its kernel is $(\prod_{i=1}^{p-2} \omega_i)$ -torsion.

Proof. This is essentially a consequence of results in [So85] as observed in [GS87, Theorem 8.2]. Since our formulation claims a little bit stronger than the original one, we give a sketch of the proof here.

Consider the Gersten-Quillen spectral sequence

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)} \cap Y} K_{-p-q}(k(x)) \Rightarrow K_{-p-q}^Y(X).$$

The Adams operations act on this spectral sequence, and by the Riemann-Roch type formula [So85, Théorème 3] ψ^k acts by the multiplication by k^{p+i} on $E_r^{p,-p-i}$ for $i = 0, 1$ and $r \geq 1$. It follows that the differential $d_r^p: E_r^{p-r, -p+r-1} \rightarrow E_r^{p,-p}$ is killed by ω_{r-1} for $p \geq r > 1$. In fact, $d_r^p = 0$ for $r > 1$, because

$$E_2^{0,-1} \simeq \bigoplus_{x \in X^{(0)} \cap Y} \mathcal{O}(\overline{\{x\}})^\times \simeq E_\infty^{0,-1}.$$

Consequently, the kernel of the canonical surjection $E_2^{p,-p} \rightarrow E_\infty^{p,-p}$ is killed by $\prod_{i=1}^{p-2} \omega_i$. On the other hand, we have $E_2^{p,-p} \simeq \text{CH}_Y^p(X)$ and $E_\infty^{p,-p} \simeq \text{Gr}^p K_0^Y(X)$. Hence, we get the result. \square

Lemma 4.3. *Let X be a regular scheme, Y a closed subset of X , D a principal effective Cartier divisor on X and p an integer. We denote the inclusion $D \hookrightarrow X$ by ι . Then the diagram*

$$\begin{array}{ccc} K_0^Y(X) & \xrightarrow{\iota^*} & K_0^{Y \cap D}(D)/F^{p+1} K_0^{Y \cap D}(D) \\ \text{cyc} \uparrow & & \uparrow \text{cyc} \\ \mathcal{Z}_Y^p(X) & \xrightarrow{\iota^*} & \text{CH}_{Y \cap D}^p(D) \end{array}$$

commutes. Here, the bottom horizontal map ι^* is the one defined in Construction 1.7.

Proof. Suppose that we are given $v \in X^{(p)} \cap Y$ and denote its closure in X by V . If $V \not\subseteq D$, then the commutativity is clear, i.e., $(\iota^* \circ \text{cyc})[V] = (\text{cyc} \circ \iota^*)[V]$. Suppose that $V \subseteq D$. Then $\iota^*[V] = 0$ in $\text{CH}_{Y \cap D}^p(D)$. Also, $\iota^*(\mathcal{O}_V) = [\mathcal{O}_V \xrightarrow{f} \mathcal{O}_V] = 0$ in $K_0^{Y \cap D}(D)$, where f denotes the defining equation of D . This proves the lemma. \square

Corollary 4.4. *Under the situation in Lemma 4.3, the restriction morphism $K_0^Y(X) \rightarrow K_0^{Y \cap D}(D)$ preserves the coniveau filtration.*

On codimension one.

Lemma 4.5. *Let X be a regular scheme and Y a closed subset of X not containing any irreducible components of X . Then there are natural isomorphisms*

$$\begin{aligned} K_0^Y(X) &\simeq \text{CH}_Y^1(X) \oplus F^2 K_0^Y(X), \\ \text{CH}_Y^1(X) &\simeq H_{\text{Zar}, Y}^1(X, \mathcal{O}^\times) \quad \text{and} \quad F^2 K_0^Y(X) \simeq F_\gamma^2 K_0^Y(X). \end{aligned}$$

Proof. Since $K_0^Y(X) = F^1 K_0^Y(X)$, we get an isomorphism

$$\text{cyc}: \text{CH}_Y^1(X) \xrightarrow{\sim} K_0^Y(X)/F^2 K_0^Y(X)$$

by Theorem 4.2. Since $\text{CH}_Y^1(X)$ is the free abelian group generated by the irreducible components of Y of codimension one in X , the above isomorphism factors through $K_0^Y(X)$. This proves the first isomorphism. The second isomorphism follows from the quasi-isomorphism $\mathcal{Z}^1(X, \bullet) \simeq \mathcal{O}^\times[1]$ ([BI86, Theorem 6.1]) and Lemma 3.7. The last isomorphism follows from the first two isomorphisms. \square

Corollary 4.6. *Let X be a regular scheme and D an effective Cartier divisor on X . Assume that D has an affine open neighborhood in X . Then there are exact sequences*

$$\begin{aligned} \operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} F^2 K_0^Y(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \operatorname{colim}_{Y \in \mathcal{S}(X|D)} F^2 K_0^Y(X) \xrightarrow{\epsilon} F^2 K_0(X, D) \longrightarrow 0 \\ \operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \operatorname{Gr}^1 K_0^Y(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^1 K_0^Y(X) \xrightarrow{\epsilon} \operatorname{Gr}^1 K_0(X, D) \longrightarrow 0 \end{aligned}$$

and the same for the γ -filtration.

Proof. The morphisms are well-defined by Corollary 4.4. We may assume that X is connected. If D is empty, then the result is clear. If D is not empty, then it follows from Theorem 2.2 and Lemma 4.5. \square

Theorem 4.7. *Let X be a regular scheme and D an effective Cartier divisor on X . Assume that D has an affine open neighborhood in X . Then there are natural isomorphisms*

$$\operatorname{CH}^1(X|D) \simeq \operatorname{Gr}^1 K_0(X, D) \simeq \operatorname{Gr}_\gamma^1 K_0(X, D) \simeq \operatorname{Pic}(X, D).$$

Proof. The first isomorphism follows from the commutative diagram

$$\begin{array}{ccccc} \operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \operatorname{Gr}^1 K_0^Y(X \times \square^1) & \xrightarrow{\iota_0^* - \iota_1^*} & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^1 K_0^Y(X) & \xrightarrow{\epsilon} & \operatorname{Gr}^1 K_0(X, D) \longrightarrow 0 \\ \uparrow \text{cyc} & & \uparrow \text{cyc} \simeq & & \uparrow \text{dotted} \\ \operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \mathcal{Z}_Y^1(X \times \square^1) & \xrightarrow{\iota_0^* - \iota_1^*} & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{CH}_Y^1(X) & \xrightarrow{\epsilon} & \operatorname{CH}^1(X|D) \longrightarrow 0. \end{array}$$

This is indeed commutative by Lemma 4.3, and the rows are exact by Proposition 1.16 and Corollary 4.6. The middle vertical arrow is an isomorphism by Theorem 4.2.

The second isomorphism follows from Lemma 4.5 and Corollary 4.6. The last isomorphism has been observed in Lemma 3.7. \square

Example 4.8. Let k be a field, $X = \mathbb{P}_k^1 \times_k \mathbb{P}^1$ and $D = \mathbb{P}_k^1$ which we regard as a Cartier divisor on X by the diagonal embedding. Then $\operatorname{CH}^1(X|D) = 0$, but $\operatorname{Gr}_\gamma^1 K_0(X, D) = \operatorname{Pic}(X, D) = \mathbb{Z}$.

On higher codimension.

Lemma 4.9. *Let X be a regular scheme, D an effective Cartier divisor on X and p an integer. Assume that D has an affine open neighborhood in X . Suppose we are given*

$$\alpha \in \ker \left(\operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^p K_0^Y(X) \xrightarrow{\epsilon} \operatorname{Gr}^p K_0(X, D) \right).$$

Then there exists $\beta \in \operatorname{Gr}^p K_0^W(X \times \square^1)$ for some $W \in \mathcal{S}(X|D, 1)$ such that

$$\iota_0^* \beta - \iota_1^* \beta = \left(\prod_{i=2}^{p-1} w_{p-i} \right) \left(\prod_{2 \leq i, j \leq p} w_{|i-j|} \right)^2 \alpha.$$

Proof. We may assume that D is non-empty. The case $p \leq 1$ is true by Theorem 2.2 and Lemma 4.6. Let $p > 1$. We consider the diagram

$$\begin{array}{ccc}
\operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \bigoplus_{i=2}^p (F^2 K_0^Y(X \times \square^1)/F^{p+1})^{(i)} & \longrightarrow & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \bigoplus_{i=2}^p (F^2 K_0^Y(X)/F^{p+1})^{(i)} \\
\downarrow & & \downarrow \\
\operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} F^2 K_0^Y(X \times \square^1)/F^{p+1} & \longrightarrow & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} F^2 K_0^Y(X)/F^{p+1} \\
\uparrow & & \uparrow \\
\operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \operatorname{Gr}^p K_0^Y(X \times \square^1) & \longrightarrow & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^p K_0^Y(X)
\end{array}$$

Suppose given α as in the statement. According to Corollary 4.6, there exists $\tilde{\beta} \in F^2 K_0^W(X \times \square^1)/F^{p+1}$ for some $W \in \mathcal{S}(X|D, 1)$ such that

$$\iota_0^* \beta - \iota_1^* \beta = \alpha \quad \text{in} \quad \operatorname{colim}_{Y \in \mathcal{S}(X|D)} F^2 K_0^Y(X)/F^{p+1}.$$

By Lemma 3.4, $(\prod_{2 \leq i, j \leq p} w_{|i-j|}) \tilde{\beta}$ lifts to

$$\beta^{(2)} + \beta^{(3)} + \dots + \beta^{(p)} \in \bigoplus_{i=2}^p (F^2 K_0^W(X \times \square^1)/F^{p+1})^{(i)}.$$

By Lemma 3.4 again, we see that

$$\left(\prod_{2 \leq i, j \leq p} w_{|i-j|} \right) (\iota_0^* \beta^{(p)} - \iota_1^* \beta^{(p)}) = \left(\prod_{2 \leq i, j \leq p} w_{|i-j|} \right)^2 \alpha.$$

Since $(\prod_{i=2}^{p-1} w_{p-i}) \beta^{(p)}$ is in $\operatorname{Gr}^p K_0^W(X \times \square^1)$, we are done. \square

Proof of Theorem 0.1. Let X be a regular scheme, D an effective Cartier divisor on X and p an integer. Let us consider the commutative diagram

$$\begin{array}{ccccccc}
\operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \operatorname{Gr}^p K_0^Y(X \times \square^1) & \longrightarrow & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{Gr}^p K_0^Y(X) & \longrightarrow & \operatorname{Gr}^p K_0(X, D) & \longrightarrow & 0 \\
\operatorname{cyc}_1 \uparrow & & \operatorname{cyc}_0 \uparrow & & \operatorname{cyc}_{\text{rel}} \uparrow & & \\
\operatorname{colim}_{Y \in \mathcal{S}(X|D,1)} \operatorname{CH}_Y^p(X \times \square^1) & \longrightarrow & \operatorname{colim}_{Y \in \mathcal{S}(X|D)} \operatorname{CH}_Y^p(X) & \longrightarrow & \operatorname{CH}^p(X|D) & \longrightarrow & 0.
\end{array}$$

Since the bottom row is exact (Proposition 1.16) and the composite of the first two morphisms in the upper row is zero (Theorem 2.2), a morphism $\operatorname{cyc}_{\text{rel}}$ is induced and it is surjective.

Suppose we are given $\alpha \in \operatorname{CH}^p(X|D)$ such that $\operatorname{cyc}_{\text{rel}}(\alpha) = 0$. By Lemma 4.9 and by a simple diagram chase, there exists $\beta \in \ker(\operatorname{cyc}_0)$ which lifts

$$\left(\prod_{i=2}^{p-1} w_{p-i} \right) \left(\prod_{2 \leq i, j \leq p} w_{|i-j|} \right)^2 \alpha.$$

By Theorem 4.2, $(\prod_{i=1}^{p-2} w_i) \beta = 0$.

The last statement (comparison between F^* and the gamma filtration) follows from Proposition 3.4 and Theorem 4.7. \square

5. APPLICATIONS

Multiplicative structure on Chow groups with modulus. As an application of Theorem 0.1, we prove that there is a natural multiplicative structure on $\mathrm{CH}^*(X|D)$ up to torsion. We formulate it keeping track of the changes of D . Note that the Chow groups with modulus yield a contravariant functor

$$\mathrm{CH}^*(X|-) : \mathrm{Div}^+(X)^{\mathrm{op}} \rightarrow \mathrm{GrAb}$$

from the category of effective Cartier divisors on X to the category of graded abelian groups.

Notation 5.1. Let \mathcal{A} be an additive category and l a positive integer. We define a category $\mathcal{A}_{\mathbb{Z}[1/l]}$ having the same objects as \mathcal{A} and $\mathrm{Hom}_{\mathcal{A}[1/l]}(-, -) = \mathrm{Hom}_{\mathcal{A}}(-, -) \otimes_{\mathbb{Z}} \mathbb{Z}[1/l]$. For an object M in \mathcal{A} , we denote its image in $\mathcal{A}_{\mathbb{Z}[1/l]}$ by $M_{\mathbb{Z}[1/l]}$.

Theorem 5.2. *Let X be a regular scheme of dimension d . Let $\mathrm{Div}_{\mathrm{aff}}^+(X)$ be the full subcategory of $\mathrm{Div}^+(X)$ consisting of divisors admitting affine open neighborhoods in X . Then*

$$\mathrm{CH}^*(X|-)_{\mathbb{Z}[1/(d-1)!]} \in \mathrm{Fun}(\mathrm{Div}_{\mathrm{aff}}^+(X)^{\mathrm{op}}, \mathrm{GrAb})_{\mathbb{Z}[1/(d-1)!]}$$

has a natural commutative monoid structure.

Proof. By Theorem 0.1, it suffices to show that $\mathrm{Gr}_{\gamma}^* K_0(X, -)$ has a commutative monoid structure, but it is obvious from its definition. \square

Chow groups with topological modulus. Here, we show that if D is K_1 -regular then the Chow group with modulus $\mathrm{CH}^*(X|D)$ becomes much simpler (at least up to torsion). Compare the following definition with Notation 1.12 and Definition 1.14.

Definition 5.3. Let X be a unidimensional catenary scheme, D an effective Cartier divisor and p an integer.

- (1) $\mathcal{S}(X|D_{\mathrm{top}}) := \mathcal{S}(X|D)$ and $\mathcal{Z}^p(X|D_{\mathrm{top}}) := \mathcal{Z}^p(X|D)$.
- (2) $\mathcal{S}(X|D_{\mathrm{top}}, 1)$ is the set of all closed subsets of $X \times \square^1$ not meeting $D \times \square^1$.
- (3) $\mathcal{Z}^p(X|D_{\mathrm{top}}, 1)$ is the free abelian group with generators $[W]$, one for each $w \in (X \times \square^1)^{(p)}$ whose closure W is dominant over \square^1 and not meeting $D \times \square^1$.
- (4) We define

$$\mathrm{CH}^p(X|D_{\mathrm{top}}) := \mathrm{coker}(\mathcal{Z}^p(X|D_{\mathrm{top}}, 1) \xrightarrow{\iota_0^* - \iota_1^*} \mathcal{Z}^p(X|D_{\mathrm{top}})).$$

Remark 5.4. The groups $\mathrm{CH}^p(X|D_{\mathrm{top}})$ and its higher variant have been studied in [Mi17, IK] by the name of *naïve Chow groups with modulus* and *Chow groups with topological modulus* respectively.

Lemma 5.5. *Let X be a unidimensional catenary scheme, D an effective Cartier divisor on X and p an integer. Then the sequence*

$$\mathrm{colim}_{Y \in \mathcal{S}(X|D_{\mathrm{top}}, 1)} \mathcal{Z}_Y^p(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \mathrm{colim}_{Y \in \mathcal{S}(X|D_{\mathrm{top}})} \mathrm{CH}_Y^p(X) \xrightarrow{\epsilon} \mathrm{CH}^p(X|D_{\mathrm{top}}) \longrightarrow 0$$

is exact.

Proof. Same as Proposition 1.16. \square

The following is a variant of Theorem 2.2, which is a special case of [IK, Lemma xx].

Lemma 5.6. *Let X be a scheme and D an effective Cartier divisor on X admitting an affine open neighborhood in X . Assume that X is K_0 -regular and that D is K_1 -regular. Then the sequence*

$$\mathrm{colim}_{Y \in \mathcal{S}(X|D_{\mathrm{top}}, 1)} K_0^Y(X \times \square^1) \xrightarrow{\iota_0^* - \iota_1^*} \mathrm{colim}_{Y \in \mathcal{S}(X|D_{\mathrm{top}})} K_0^Y(X) \xrightarrow{\epsilon} K_0(X, D) \longrightarrow 0$$

is exact.

Proof. By the assumption, $K_0(X, D) \simeq K_0(X \times \square^1, D \times \square^1)$, from which it follows that the composite $\epsilon \circ (\iota_0^* - \iota_1^*)$ is zero. The surjectivity of ϵ follows from Lemma 2.3, and the exactness at the middle term follows from Lemma 2.6. \square

Proposition 5.7. *Let X be a regular scheme and D an effective Cartier divisor on X . Assume that D is K_1 -regular and admits an affine open neighborhood in X . Then, for each integer p , there exists a surjective group morphism*

$$\mathrm{CH}^p(X|D_{\mathrm{top}}) \twoheadrightarrow F^p K_0(X, D)/F^{p+1} K_0(X, D)$$

such that its kernel is $(p-1)!^N$ -torsion for some positive integer N depending only on p .

Proof. This follows from Lemma 5.5 and Lemma 5.6 as in §4. \square

Theorem 5.8. *Let X be a regular scheme and D an effective Cartier divisor on X . Assume that D is K_1 -regular and admits an affine open neighborhood in X . Then, for each integer p , the canonical morphism*

$$\mathrm{CH}^p(X|D) \rightarrow \mathrm{CH}^p(X|D_{\mathrm{top}})$$

is a $(p-1)!^N$ -isomorphism for some positive integer N depending only in p .

Proof. This follows from Theorem 0.1 and Proposition 5.7. \square

REFERENCES

- [AM69] M. Atiyah, I. MacDonal, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [AT69] M. Atiyah, D. Tall, *Group representations, λ -rings and the J -homomorphism*, *Topology* 8 (1969), 253–297.
- [BS17] F. Binda, S. Saito, *Relative cycles with moduli and regulator maps*, *J. Inst. Math. Jussieu* (2017), 1–61.
- [B186] S. Bloch, *Algebraic cycles and higher K -theory*, *Adv. in Math.* 61 (1986), no. 3, 267–304.
- [GS87] H. Gillet, C. Soulé, *Intersection theory using Adams operations* *Invent. Math.* 90 (1987), no. 2, 243–277.
- [Fu98] W. Fulton, *Intersection theory*, Second edition, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, 2. Springer-Verlag, 1998.
- [Iw19] R. Iwasa, *Relative K_0 and relative cycle class map*, *J. Pure Appl. Algebra* 223 (2019), 48–71.
- [IK] R. Iwasa, A. Krishna, *Relative homotopy K -theory and algebraic cycles with modulus*, in preparation.
- [Kr80] C. Kratzer, *λ -structure en K -théorie algébrique*, *Comment. Math. Helv.* 55 (1980), no. 2, 233–254.
- [KP12] A. Krishna, J. Park, *Moving lemma for additive higher Chow groups*, *Algebra Number Theory* 6 (2012), no. 2, 293–326.
- [Le97] M. Levine, *Lambda-operations, K -theory and motivic cohomology*, *Algebraic K -theory* (Toronto, ON, 1996), 131–184, *Fields Inst. Commun.*, 16, Amer. Math. Soc., Providence, RI, 1997.
- [MS74] J. Milnor, J. Stasheff, *Characteristic classes*, *Annals of Mathematics Studies*, No. 76., Princeton University Press; University of Tokyo Press, 1974.
- [Mi17] H. Miyazaki, *Cube invariance of higher Chow groups with modulus*, arXiv:1604.06155, (2017).
- [So85] C. Soulé, *Opérations en K -théorie algébrique*, *Canad. J. Math.* 37 (1985), no. 3, 488–550.
- [TT90] R. W. Thomason, T. Trobaugh, *Higher algebraic K -theory of schemes and of derived categories*, *The Grothendieck Festschrift*, Vol. III, 247–435, *Progr. Math.* 88, Birkhäuser Boston, 1990.
- [EGA2] J. Dieudonné, A. Grothendieck, *Éléments de géométrie algébrique II*, *Publ. math. I.H.É.S.* 8 (1961), 5–222.
- [SGA6] P. Berthelot, A. Grothendieck, L. Illusie, *Théorie des intersections et théorème de Riemann-Roch*, *Séminaire de Géométrie Algébrique de Bois-Marie 1966-1967 (SGA 6)*, *Lecture Notes in Math.*, Vol. 225, Springer-Verlag, 1971.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5, DK-2100 COPENHAGEN Ø.
E-mail address: ryomei@math.ku.dk

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, AZA-AOBA 6-3, SENDAI 980-8578, JAPAN.
E-mail address: kaiw@tohoku.ac.jp