Critical points of solutions to a kind of linear elliptic equations in multiply connected domains^{*}

Haiyun Deng¹[†], Hairong Liu², Xiaoping Yang³

¹School of Science, Nanjing University of Science and Technology, Nanjing, Jiangsu, 210094, China; ²School of Science, Nanjing Forestry University, Nanjing, Jiangsu, 210037, China;

³Department of Mathematics, Nanjing University, Nanjing, Jiangsu, 210093, China

Abstract: In this paper, we mainly study the critical points and critical zero points of solutions u to a kind of linear elliptic equations with nonhomogeneous Dirichlet boundary conditions in a multiply connected domain Ω in \mathbb{R}^2 . Based on the fine analysis about the distributions of connected components of the super-level sets $\{x \in \Omega : u(x) > t\}$ and sub-level sets $\{x \in \Omega : u(x) < t\}$ for some t, we obtain the geometric structure of interior critical point sets of u. Precisely, let Ω be a multiply connected domain with the interior boundary γ_I and the external boundary γ_E , where $u|_{\gamma_I} = \psi_1(x), u|_{\gamma_E} = \psi_2(x)$. When $\psi_1(x)$ and $\psi_2(x)$ have N_1 and N_2 local maximal points on γ_I and γ_E respectively, we deduce that $\sum_{i=1}^k m_i \leq N_1 + N_2$, where m_1, \dots, m_k are the respective multiplicities of interior critical points x_1, \dots, x_k of u. In addition, when $\min_{\gamma_E} \psi_2(x) \geq \max_{\gamma_I} \psi_1(x)$ and u has only N_1 and N_2 equal local maxima relative to $\overline{\Omega}$ on γ_I and γ_E respectively, we develop a new method to show that one of the following three results holds $\sum_{i=1}^k m_i = N_1 + N_2$ or $\sum_{i=1}^k m_i + 1 = N_1 + N_2$ or $\sum_{i=1}^k m_i + 2 = N_1 + N_2$. Moreover, we investigate the geometric structure of interior critical zero points of u. We obtain that the sum of multiplicities of the interior critical zero points of u is less than or equal to the half of the number of its isolated zero points on the boundaries.

Key Words: critical point, critical zero point, multiplicity, level sets.

2010 Mathematics Subject Classification. 35J25; 35B38.

1 Introduction and main results

In this paper we mainly investigate the interior critical points and critical zero points of solutions to the following elliptic equations

$$Lu := \sum_{i,j=1}^{2} a_{ij}(x) u_{x_i x_j}(x) + \sum_{i=1}^{2} b_i(x) u_{x_i}(x) = 0 \quad \text{in } \Omega,$$
(1.1)

where Ω is a bounded, smooth and multiply connected domain in \mathbb{R}^2 , a_{ij} , b_i are smooth and L is uniformly elliptic in Ω .

Critical points of solutions to elliptic equations is a significant research topic. There are many known results about the critical points. In 1987 Alessandrini [2] investigated the geometric structure of the critical point sets of solutions to linear elliptic equations with nonhomogeneous Dirichlet boundary condition in a planar bounded simply connected domain. In 1994, Sakaguchi [28] considered the critical points of solutions to an obstacle problem in a planar bounded smooth simply

^{*}The work is supported by The work is supported by National Natural Science Foundation of China (No.11401310) and Postgraduate Research & Practice Innovation Program of Jiangsu Province (KYCX17_0321). The first author is fully supported by China Scholarship Council(CSC) for visiting Rutgers University (201806840122).

[†]E-mail: haiyundengmath1989@163.com(H. Deng), hrliu@njfu.edu.cn(H. Liu), xpyang@nju.edu.cn(X. Yang)

connected domain. He showed that if the number of critical points of the obstacle is finite and the obstacle has only N local (global) maximum points, then $\sum_{i=1}^{k} m_i + 1 \leq N$ ($\sum_{i=1}^{k} m_i + 1 = N$), where m_1, m_2, \dots, m_k are the respective multiplicities of the critical points x_1, x_2, \dots, x_k of u in the noncoincidence set. In 2012 Arango and Gómez [6] studied the critical points of the solutions to a quasilinear elliptic equation with Dirichlet boundary condition in planar strictly convex and nonconvex domains respectively. In 2018, Deng, Liu and Tian [13] investigated the geometric structure of interior critical point sets of solutions u to a quasilinear elliptic equation with nonhomogeneous Dirichlet boundary condition in a simply connected or multiply connected domain Ω in \mathbb{R}^2 , and proved that $\sum_{i=1}^{k} m_i + 1 = N$ or $\sum_{i=1}^{k} m_i = N$, where m_1, \dots, m_k are the respective multiplicities of interior critical points x_1, \dots, x_k of u and N is the number of global maximal points of u on $\partial\Omega$. For the related research work of critical points, see [3, 4, 5, 8, 10, 12, 16, 17, 21, 27].

The geometric structure of critical point sets of solutions to elliptic equations in higher dimensional spaces has been studied by many people. Under the assumption of the existence of a semistable solution of Poisson equation $-\Delta u = f(u)$, Cabré and Chanillo [7] showed that the solution u has exactly one nondegenerate critical point in bounded smooth convex domains of $\mathbb{R}^n (n \ge 2)$. In 1998, Han, Hardt and Lin [18] concerned with the geometric measure of singular sets of weak solutions of second-order elliptic equations. They gave an estimate on the measure of singular sets in terms of the frequency of solutions. In 2015, Cheeger, Naber and Valtorta [9] introduced some techniques for giving measure estimates of the critical sets, including the Minkowski type estimates on the effective critical set $C_r(u)$. In 2017, Naber and Valtorta [26] introduced some techniques for estimating the critical sets and singular sets which avoid the need of any ε -regularity lemmas. In 2017 Alberti, Bal and Di Cristo [1] studied the existence of critical points for solutions to secondorder elliptic equations of the form $\nabla \cdot \sigma(x) \nabla u = 0$ in a bounded domain with prescribed boundary conditions in $\mathbb{R}^n (n \ge 3)$. See [11, 14, 15, 19, 22, 23, 24, 25, 29, 30] for some related work.

The goal of this paper is further to study the interior critical points and critical zero points of solutions to a kind of linear elliptic equations with nonhomogeneous Dirichlet boundary conditions in a multiply connected domain. Our main results are as follows.

Theorem 1.1. Let Ω be a bounded, smooth and multiply connected domain with one interior boundary γ_I and the external boundary γ_E in \mathbb{R}^2 and $\psi_1(x), \psi_2(x) \in C^1(\overline{\Omega})$. Suppose that $\psi_1(x)$ and $\psi_2(x)$ have N_1 local maximal points and N_2 local maximal points on γ_I and γ_E respectively. Let u be a non-constant solution of the following boundary value problem

$$\begin{cases} \sum_{i,j=1}^{2} a_{ij}(x) u_{x_i x_j}(x) + \sum_{i=1}^{2} b_i(x) u_{x_i}(x) = 0 & \text{in } \Omega, \\ u|_{\gamma_I} = \psi_1(x), u|_{\gamma_E} = \psi_2(x). \end{cases}$$
(1.2)

Then u has finite interior critical points, denoting by x_1, x_2, \dots, x_k , and the following inequality holds

$$\sum_{i=1}^{k} m_i \le N_1 + N_2, \tag{1.3}$$

where m_1, m_2, \cdots, m_k are the multiplicities of critical points x_1, x_2, \cdots, x_k respectively.

Theorem 1.2. Suppose that the domain Ω satisfies the hypothesis of Theorem 1.1 and $\min_{\gamma_E} \psi_2(x) \ge \max_{\gamma_I} \psi_1(x)$. Let u be a non-constant solution of (1.2). In addition, suppose that u has only N_1 equal local maxima and N_1 equal local minima relative to $\overline{\Omega}$ on γ_I and that u has only N_2 equal local maxima

and N_2 equal local minima relative to $\overline{\Omega}$ on γ_E , i.e., the values of all local maximal (minimum) points on the corresponding boundary are equal. Then u has finite interior critical points, and one of the following three results holds

$$\sum_{i=1}^{k} m_i = N_1 + N_2, \tag{1.4}$$

or

$$\sum_{i=1}^{k} m_i + 1 = N_1 + N_2, \tag{1.5}$$

or

$$\sum_{i=1}^{k} m_i + 2 = N_1 + N_2, \tag{1.6}$$

where m_i is as in Theorem 1.1.

Theorem 1.3. Let Ω be a bounded, smooth and multiply connected domain with the interior boundary γ_I and the external boundary γ_E in \mathbb{R}^2 . Suppose that $\psi(x) \in C^1(\overline{\Omega})$ is sign-changing, H is a given constant and that ψ has \widetilde{N} zero points on γ_E . Let u be a non-constant solution of the following boundary value problem

$$\begin{cases} \sum_{i,j=1}^{2} a_{ij}(x) u_{x_i x_j}(x) + \sum_{i=1}^{2} b_i(x) u_{x_i}(x) = 0 & \text{in } \Omega, \\ u|_{\gamma_I} = H, \quad u|_{\gamma_E} = \psi(x). \end{cases}$$
(1.7)

Then u has finite interior critical zero points, denoting by x_1, \dots, x_l , and if $H \neq 0$, we have

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}}{2},\tag{1.8}$$

else H = 0, we have

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}}{2} - 1, \tag{1.9}$$

where critical zero point x means that $u(x) = |\nabla u(x)| = 0$ and m_i is the multiplicity of corresponding critical zero point x_i .

Theorem 1.4. Suppose that the domain Ω satisfies the hypothesis of Theorem 1.1. Let u be a nonconstant solution of (1.2). In addition, suppose that $\psi_1(x)$ and $\psi_2(x)$ are sign-changing and have \widetilde{N}_1 and \widetilde{N}_2 zero points on γ_I and γ_E respectively. Then u has finite interior critical zero points and

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}_1 + \tilde{N}_2}{2},\tag{1.10}$$

where m_i is as in Theorem 1.3.

Remark 1.5. According to the assumptions of Theorem 1.2 and the strong maximum principle, we know that the local minima of u on γ_I and the local maxima of u on γ_E are the global minima and global maxima of u on $\overline{\Omega}$ respectively.

Remark 1.6. If the assumption $\min_{\gamma_E} \psi_2(x) \ge \max_{\gamma_I} \psi_1(x)$ is replaced by $\max_{\gamma_E} \psi_2(x) \le \min_{\gamma_I} \psi_1(x)$ in Theorem 1.2, the conclusions are still valid. In addition, if the elliptic operator L in (1.1) is replaced by $Lu := \sum_{i,j=1}^{2} a_{ij}(\nabla u) u_{x_i x_j}(x) = 0$, where a_{ij} is smooth and L is uniformly elliptic in Ω , then the conclusions of Theorem 1.1, Theorem 1.2, Theorem 1.3 and Theorem 1.4 still hold. **Remark 1.7.** If the elliptic operator L in (1.1) is replaced by $Lu := \sum_{i,j=1}^{2} a_{ij}(x)u_{x_ix_j}(x) + \sum_{i=1}^{2} b_i(x)u_{x_i}(x) + c(x)u(x) = 0$, where c(x) is smooth, $c(x) \leq 0$ in Ω and a_{ij}, b_i is as in (1.1). By the results of [18], we know that the interior critical zero points are isolated. Then the conclusions of Theorem 1.3 and Theorem 1.4 still hold.

Throughout this paper, we set $z_1 := \min_{\gamma_I} \psi_1(x)$, $Z_1 := \max_{\gamma_I} \psi_1(x)$, $z_2 := \min_{\gamma_E} \psi_2(x)$, $Z_2 := \max_{\gamma_E} \psi_2(x)$. For the sake of clarity, we now explain the key ideas which are used to prove the main results. Based on the fine analysis about the distributions of connected components of some level sets, we prove (1.3) by induction and the strong maximum principle. On the other hand, we develop a new method to prove (1.4), (1.5) and (1.6), which is different from the methods in [2, 28]. Concerning the case of simply connected domains. In [2], the author proved the result by induction on the number N of global maximal points on $\partial \Omega$. In [28], the author showed the result by using induction and a differential homeomorphism method.

Actually, when $z_2 \ge Z_1$ and u has only N_1 (N_2) equal local maxima and N_1 (N_2) equal local minima relative to $\overline{\Omega}$ on γ_I (γ_E), we will prove that all critical values are equal to one of the two values, i.e., $u(x_1) = \cdots = u(x_l) = t_1$ and $u(x_{l+1}) = \cdots = u(x_k) = t_0$ for some $t_1 \neq t_0$. We obtain (1.4), (1.5) or (1.6) by showing that there are the following three "just right"s similar to those in [13]:

(i) the first "just right" is that all critical values are equal to one of the two values;

(ii) the second "just right" is that critical points together with the corresponding level lines of $\{x \in \Omega : u(x) = t_1 \ (t_0) \text{ for some } t_1 \in (z_2, Z_2) \ (t_0 \in (z_1, Z_1))\}$, which meet $\gamma_E \ (\gamma_I)$, clustering round these points exactly form one connected set;

(iii) the third "just right" is that every connected component of $\{x \in \Omega : u(x) > t_1 > z_2\}$ $(\{x \in \Omega : u(x) < t_0 < Z_1\})$ has exactly one global maximal (minimum) point on $\gamma_E(\gamma_I)$.

The rest of this paper is organized as follows. In Section 2, we give the fine analysis about the distributions of connected components of super-level sets $\{x \in \Omega : u(x) > t\}$ and sub-level sets $\{x \in \Omega : u(x) < t\}$ for some t. Among others, we will prove Theorems 1.1 and 1.2 in Sections 3 and 4 respectively, and Theorems 1.3 and 1.4 in Section 5.

2 Preliminaries

According to the assumption of Theorem 1.1, we know that one of the following three occurs: (I) the global maximal points are only on γ_E , (II) the global maximal points are only on γ_I , (III) the global maximal points are on γ_E and γ_I at the same time.

For (I), we have one of the following three cases

$$z_1 < Z_1 \le z_2 < Z_2$$
, $z_1 < z_2 < Z_1 < Z_2$ and $z_2 \le z_1 < Z_1 < Z_2$

One can easily observe that the third case is similar to the second one from the proofs of Lemmas in this section. Moreover during the course of proofs we also know that (II) is obviously similar to (I) and (III) is simpler than (I). In fact (I) has three cases and (III) has only two cases $z_1 \leq z_2 < Z_1 = Z_2$ and $z_2 < z_1 < Z_1 = Z_2$. Furthermore, there is a case $z_1 < z_2 < Z_1 < Z_2$ in (I), which is the most complex case that we discuss. Thus, from later on we only describe the lemmas and prove them for Case (I). Our main idea is to sort the size of the four values Z_2, z_2, Z_1, z_1 , and then study the distribution of the super-level (sub-level) sets of the corresponding interval.

In order to prove Theorem 1.1 for the cases of $z_1 < Z_1 \leq z_2 < Z_2$ and $z_1 < z_2 < Z_1 < Z_2$, we need the following lemmas.

Lemma 2.1. Suppose that x_0 is an interior critical point of u in Ω and that m is the multiplicity of x_0 . Then there exist m + 1 distinct connected components of $\{x \in \Omega : u(x) > u(x_0)\}$ and $\{x \in \Omega : u(x) < u(x_0)\}$ clustering around the point x_0 respectively.

Proof. By the results of Hartman and Wintner [20], in a neighborhood of x_0 , the level line $\{x \in \Omega : u(x) = u(x_0)\}$ consists of m + 1 simple arcs intersecting at x_0 . Then the conclusion can be easily obtained.

Lemma 2.2. (i) Suppose that $z_1 < Z_1 \leq z_2 < Z_2$ and u is a non-constant solution of (1.2), then we have:

(1) For any $t \in (z_2, Z_2)$, any connected component ω of $\{x \in \Omega : u(x) > t\}$ has to meet the external boundary γ_E .

(2) For any $t \in (z_1, Z_1)$, any connected component ω of $\{x \in \Omega : u(x) < t\}$ has to meet the interior boundary γ_I .

(ii) Suppose that $z_1 < z_2 < Z_1 < Z_2$ and u is a non-constant solution of (1.2), then we have:

(3) For any $t \in [Z_1, Z_2)$, any connected component ω of $\{x \in \Omega : u(x) > t\}$ has to meet the external boundary γ_E .

(4) For any $t \in (z_2, Z_1)$, any connected component ω of $\{x \in \Omega : u(x) > t\}$ or $\{x \in \Omega : u(x) < t\}$ may meet γ_I or γ_E .

(5) For any $t \in (z_1, z_2]$, any connected component ω of $\{x \in \Omega : u(x) < t\}$ has to meet the interior boundary γ_I .

Proof. (1) For any $t \in (z_2, Z_2)$, suppose that A is a connected component of $\{x \in \Omega : u(x) > t\}$. According to the assumption of $u|_{\gamma_I} = \psi_1(x)$ and $z_2 \geq Z_1$, we know that A can not contain the interior boundary γ_I . Then the strong maximum principle shows that the connected component A has to meet the external boundary γ_E . The proofs of (2), (3) and (5) are same as the proof of (1) and the results of (4) naturally hold.

Lemma 2.3. Let u be a non-constant solution to (1.2). Then u has finite interior critical points in Ω .

Proof. We set up the usual contradiction argument. Since the theorem of Hartman and Wintner [20] shows that the interior critical points of u are isolated, so we suppose that u has infinite interior critical points in Ω , denoting by x_1, x_2, \cdots . The results of Lemma 2.1 and Lemma 2.2 show that there exist infinite connected components of $\{x \in \Omega : u(x) > u(x_i)\}$ and $\{x \in \Omega : u(x) < u(x_i)\}$ $(i = 1, 2, \cdots)$, which meet the boundary γ_E or γ_I . The strong maximum principle implies that there totally exist infinite local maximal points and minimum points on γ_E and γ_I , this contradicts with the assumption of Theorem 1.1.

Lemma 2.4. Suppose that $z_1 < Z_1 \leq z_2 < Z_2$ and u is a non-constant solution to (1.2). Then there does not exist any interior critical point x such that u(x) = t for any $t \in [Z_1, z_2]$.

Proof. We divide the proof into two cases.

(i) Case 1: When $Z_1 = z_2$, without loss of generality, we suppose by contradiction that there exists an interior critical point x_0 such that $u(x_0) = Z_1 = z_2$ and that the multiplicity of x_0 is one. By Lemma 2.1 and Lemma 2.2, one of the distributions for x_0 and the corresponding level lines of $\{x \in \Omega : u(x) = z_2\}$ is show in Fig. 1.



Fig. 1. The distributions for x_0 and the corresponding level lines of $\{x \in \Omega : u(x) = z_2\}$.

Any one of the distributions of Fig. 1 contradicts with $Z_1 = z_2 < Z_2$ and the continuity of u.

(ii) Case 2: When $Z_1 < z_2$, without loss of generality, we suppose by contradiction that there exists an interior critical point x_0 such that $u(x_0) = t_0$ for $t_0 = Z_1$ or $t_0 \in (Z_1, z_2)$ or $t_0 = z_2$ and that the multiplicity of x_0 is one. The proof is same as Case 1.

Lemma 2.5. Let $z_1 < Z_1 \leq z_2 < Z_2$ and x_1, x_2, \dots, x_k be the interior critical points of u in Ω . Suppose that $z_1 < u(x_{l+1}) = \dots = u(x_k) = t_0 < Z_1 \leq z_2 < u(x_1) = \dots = u(x_l) = t_1 < Z_2$ and that x_1, \dots, x_l and x_{l+1}, \dots, x_k together with the corresponding level lines of $\{x \in \Omega : u(x) = t_1\}$ and $\{x \in \Omega : u(x) = t_0\}$ clustering round these points form q_1, q_0 connected sets respectively, where $q_1, q_0 \geq 1$ and m_1, m_2, \dots, m_k are the multiplicities of critical points x_1, x_2, \dots, x_k respectively.

Case 1: Suppose that there exists a non-simply connected component ω of $\{x \in \Omega : u(x) < t_1\}$ and the external boundary γ of ω is a simply closed curve between γ_I and γ_E such that u has at least one critical point on γ , then

$$\sharp \Big\{ \text{the simply connected components } \widetilde{\omega} \text{ of the sub-level set } \{x \in \Omega : u(x) < t_1 \} \\ \text{such that } \widetilde{\omega} \text{ meet the external boundary } \gamma_E \Big\} = \sum_{i=1}^l m_i + q_1 - 1.$$

$$(2.1)$$

Case 2: Suppose that there exists a non-simply connected component ω of $\{x \in \Omega : u(x) < t_1\}$ such that ω meets γ_E . In addition, we set M_1 and M_2 as the number of the connected components of the super-level set $\{x \in \Omega : u(x) > t_1\}$ and the sub-level set $\{x \in \Omega : u(x) < t_1\}$, respectively. Then

$$M_1 \ge \sum_{i=1}^{l} m_i + 1, \ M_2 \ge \sum_{i=1}^{l} m_i + 1, \ and \ M_1 + M_2 = 2 \sum_{i=1}^{l} m_i + q_1 + 1.$$
 (2.2)

Case 3: Suppose that there exists a simply closed curve γ of $\{x \in \Omega : u(x) = t_0\}$ between γ_I and γ_E such that u has at least one critical point on γ , then

 $\sharp \Big\{ \text{the simply connected components } \omega \text{ of the super-level set } \{x \in \Omega : u(x) > t_0\} \\ \text{such that } \omega \text{ meet the interior boundary } \gamma_I \Big\} = \sum_{i=l+1}^k m_i + q_0 - 1.$ (2.3)

Case 4: For $t_0 \in (z_1, Z_1)$, if Case 3 does not occur, we set M_1 and M_2 as the number of the connected components of $\{x \in \Omega : t_0 < u(x) < z_2\}$ and the sub-level set $\{x \in \Omega : u(x) < t_0\}$, respectively. Then

$$\widetilde{M}_1 \ge \sum_{i=l+1}^k m_i + 1, \ \widetilde{M}_2 \ge \sum_{i=l+1}^k m_i + 1, \ and \ \widetilde{M}_1 + \widetilde{M}_2 = 2 \sum_{i=l+1}^k m_i + q_0 + 1.$$
 (2.4)

Proof. (i) Case 1: We divide the proof into two steps.

Step 1: When $q_1 = 1$, by induction. When l = 1, the result holds by Lemma 2.1 and Lemma 2.2. For $1 \leq l \leq n$, we assume that the connected set, which consists of l critical points and the connected components clustering round these points, contains exactly $\sum_{i=1}^{l} m_i$ components ω of the sub-level set $\{x \in \Omega : u(x) < t_1\}$ such that ω meet γ_E . Let l = n + 1. Let A be the set which consists of the points x_1, x_2, \dots, x_{n+1} together with the respective components clustering round these points. We may assume that the points x_1, x_2, \dots, x_n together with the respective components clustering round these points form a connected set, denotes by B. By Lemma 2.2, we know that A cannot surround a component of $\{x \in \Omega : u(x) > t_1\}$. Up to renumbering, therefore there is only one component of $\{x \in \Omega : u(x) < t_1\}$ whose boundary α contains both x_n and x_{n+1} . The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$ is shown in Fig. 2.



Fig. 2. The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$.

Noting that both A and B are connected, by using Lemma 2.1 and the inductive assumption to B, we know that A contains exactly

$$\sum_{i=1}^{n} m_i + (m_{n+1} + 1) - 1 = \sum_{i=1}^{n+1} m_i$$

connected components $\widetilde{\omega}$ of the sub-level set $\{x \in \Omega : u(x) < t_1\}$ such that $\widetilde{\omega}$ meet γ_E . This completes the proof of step 1.

Step 2: When $q_1 \ge 2$. We easily know that the number of connected sets of the level lines $\{x \in \Omega : u(x) = t_1\}$ together with x_1, \dots, x_l increases one leading the number of connected components of $\{x \in \Omega : u(x) < t_1\}$ increases one. If all the critical points x_1, x_2, \dots, x_l together with the level lines $\{x \in \Omega : u(x) = t_1\}$ clustering round these points form q_1 connected sets. By the results of step 1, then we have

This completes the proof of Case 1.

(ii) Case 2: We divide the proof of Case 2 into two steps.

Step 1: When $q_1 = 1$, by induction. When l = 1, the result holds by Lemma 2.1 and Lemma 2.2. For $1 \leq l \leq n$, we assume that the connected set, which consists of l critical points and the connected components clustering round these points, contains exactly $\sum_{i=1}^{l} m_i + 1$ components of the super-level set $\{x \in \Omega : u(x) > t_1\}$. Let l = n + 1. Let A be the set which consists of the points x_1, x_2, \dots, x_{n+1} together with the respective components clustering round these points. We may assume that the points x_1, x_2, \dots, x_n together with the respective components clustering round

these points form a connected set, denotes by *B*. By Lemma 2.2, we know that *A* cannot surround a component of $\{x \in \Omega : u(x) < t_1\}$. Up to renumbering, therefore there is only one component of $\{x \in \Omega : u(x) > t_1\}$ whose boundary γ contains both x_n and x_{n+1} . The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$ is shown in Fig. 3.



Fig. 3. The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$.

Noting that both A and B are connected, by using Lemma 2.1 and the inductive assumption to B, we know that A contains exactly

$$\left(\sum_{i=1}^{n} m_i + 1\right) + (m_{n+1} + 1) - 1 = \sum_{i=1}^{n+1} m_i + 1$$

connected components of the super-level set $\{x \in \Omega : u(x) > t_1\}$.

Step 2: When $q_1 \ge 2$. We know that the number of connected sets of the level lines $\{x \in \Omega : u(x) = t_1\}$ together with x_1, \dots, x_l increases one leading the number of connected components of $\{x \in \Omega : u(x) > t_1\}$ or $\{x \in \Omega : u(x) < t_1\}$ increases one. If the critical points x_1, x_2, \dots, x_l together with the level lines of $\{x \in \Omega : u(x) = t_1\}$ clustering round these points form q_1 connected sets. By the results of step 1, then we have

$$M_1 \ge \sum_{i=1}^{l} m_i + 1, \quad M_2 \ge \sum_{i=1}^{l} m_i + 1,$$

and

$$M_1 + M_2 = 2(\sum_{i=1}^{l} m_i + 1) + (q_1 - 1) = 2\sum_{i=1}^{l} m_i + q_1 + 1.$$

This completes the proof of Case 2.

(iii) Case 3: We divide the proof of Case 3 into two steps. For convenience, up to renumbering, we denote x_{l+1}, \dots, x_k and m_{l+1}, \dots, m_k by y_1, \dots, y_{k-l} and $\widetilde{m}_1, \dots, \widetilde{m}_{k-l}$ respectively.

Step 1: When $q_0 = 1$, by induction. When k - l = 1, the result holds by Lemma 2.1 and Lemma 2.2. For $1 \le k - l \le n$, we assume that the connected set, which consists of k - l critical points and the connected components clustering round these points, contains exactly $\sum_{i=1}^{k-l} \tilde{m}_i$ components ω of the super-level set $\{x \in \Omega : u(x) > t_0\}$ such that ω meet the interior boundary γ_I . Let k - l = n + 1. Let A be the set which consists of the points y_1, \dots, y_{n+1} together with the respective components clustering round these points. We may assume that the points y_1, \dots, y_n together with the respective components clustering round these points form a connected set, denotes by B. By Lemma 2.2 we know that A cannot surround a component of $\{x \in \Omega : u(x) > t_0\}$ whose boundary α contains both y_n and y_{n+1} . The distribution for the level lines of $\{x \in \Omega : u(x) = t_0\}$ is shown in Fig. 4.



Fig. 4. The distribution for the level lines of $\{x \in \Omega : u(x) = t_0\}$.

Noting that both A and B are connected, by using Lemma 2.1 and the inductive assumption to B, we know that A contains exactly

$$\sum_{i=1}^{n} \widetilde{m}_i + (\widetilde{m}_{n+1} + 1) - 1 = \sum_{i=1}^{n+1} \widetilde{m}_i = \sum_{i=l+1}^{k} m_i$$

connected components ω of the super-level set $\{x \in \Omega : u(x) > t_0\}$ such that ω meet γ_I . This completes the proof of step 1.

Step 2: When $q_0 \ge 2$. Since the number of connected sets of the level lines $\{x \in \Omega : u(x) = t_0\}$ together with x_{l+1}, \dots, x_k increases one leading the number of connected components of $\{x \in \Omega : u(x) > t_0\}$ increases one. If all the critical points x_{l+1}, \dots, x_k together with the level lines $\{x \in \Omega : u(x) = t_0\}$ clustering round these points form q_0 connected sets. By the results of step 1, then we have

 $\sharp \Big\{ \text{the simply connected components } \omega \text{ of the super-level set } \{x \in \Omega : u(x) > t_0 \} \\ \text{such that } \omega \text{ meet the interior boundary } \gamma_I \Big\} = \sum_{i=l+1}^k m_i + (q_0 - 1).$

This completes the proof of Case 3.

(iv) Case 4: Case 2 implies Case 4.

Remark 2.6. Note that if $z_1 < u(x_{l+1}) = \cdots = u(x_k) = t_0 < Z_1 \le z_2 < u(x_1) = \cdots = u(x_l) = t_1 < Z_2$. Suppose that there exists a non-simply connected component ω of $\{x \in \Omega : u(x) < t_1\}$ for critical value t_1 , where the external boundary γ_1 of ω is a simply closed curve in Ω as in Case 1 of Lemma 2.5 or that there exists a simply closed curve γ_0 of $\{x \in \Omega : u(x) = t_0\}$ between γ_I and γ_E as in Case 3 of Lemma 2.5, then u has at least one critical point on γ_1 and γ_0 respectively. For the sake of clarity, we give the proof of the first situation. In fact, suppose by contradiction that u has no critical point on γ_1 . Without loss of generality, we may assume that u has only two local maximal points q_1, q_2 on γ_E and one critical point x_1 in $\Omega \setminus \overline{\omega}$ such that $u(x_1) = t_1$ and the multiplicity of x_1 is one, we denote the non-simply connected component of $\{x \in \Omega : u(x) > t_1\}$ by A. The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$ is shown in Fig. 5.



Fig. 5. The distribution for the level lines of $\{x \in \Omega : u(x) = t_1\}$.

By using the method of step 3 of the proof of Theorem 1.2 in [13], this would imply that either: $u(x) = u(q_2)$ in interior points of A, or: there exist two level lines intersecting in A, i.e., there exist critical points in A. This is a contradiction.

Lemma 2.7. Let $z_1 < z_2 < Z_1 < Z_2$ and x_1, x_2, \dots, x_k be the interior critical points of u in Ω and m_1, m_2, \dots, m_k be the multiplicities of critical points x_1, x_2, \dots, x_k respectively. Situation 1: Suppose that $z_1 < u(x_{l+1}) = \dots = u(x_k) = t_0 \le z_2 < Z_1 \le u(x_1) = \dots = u(x_l) = t_1 < Z_2$ or $z_2 < u(x_{l+1}) = \dots = u(x_k) = t_0 < Z_1 \le u(x_1) = \dots = u(x_l) = t_1 < Z_2$ or $z_1 < u(x_{l+1}) = \dots = u(x_k) = t_1 < Z_1$ and that x_1, \dots, x_l and x_{l+1}, \dots, x_k together with the corresponding level lines of $\{x \in \Omega : u(x) = t_1\}$ and $\{x \in \Omega : u(x) = t_0\}$ clustering round these points meet γ_E, γ_I and form q_1, q_0 connected sets respectively, where $q_1, q_0 \ge 1$.

Case 1: If there exists a simply closed curve γ_1 of $\{x \in \Omega : u(x) = t_1\}$ between γ_I and γ_E such that u has at least one critical point on γ_1 and there exists a simply closed curve γ_0 of $\{x \in \Omega : u(x) = t_0\}$ between γ_I and γ_1 such that u has at least one critical point on γ_0 or there exists totally one simply closed curve γ of $\{x \in \Omega : u(x) = t\}$ for two critical values between γ_I and γ_E such that u has at least one critical point on γ . Then

and

$$\sharp \Big\{ \text{the simply connected components } \omega \text{ of the super-level set } \{x \in \Omega : u(x) > t_0 \} \\ \text{such that } \omega \text{ meet the interior boundary } \gamma_I \Big\} \ge \sum_{i=l+1}^k m_i + q_0 - 1.$$

$$(2.6)$$

Case 2: If there does not exist a simply closed curve γ of $\{x \in \Omega : u(x) = t\}$ for any critical value t between γ_I and γ_E , then

and

Situation 2: Suppose that $z_2 < u(x_1) = u(x_2) = \cdots = u(x_k) = t < Z_1$ and that x_1, \cdots, x_k together with the corresponding level lines of $\{x \in \Omega : u(x) = t\}$ clustering round these points form q connected sets, where $q \ge 1$. In addition, we set M_1 and M_2 as the number of the connected components of $\{x \in \Omega : u(x) > t\}$ and $\{x \in \Omega : u(x) < t\}$, respectively.

Case 3: If there exists a simply closed curve γ of $\{x \in \Omega : u(x) = t\}$ between γ_I and γ_E such that u has at least two critical points on γ , then

$$M_1 \ge \sum_{i=1}^k m_i, \ M_2 \ge \sum_{i=1}^k m_i, \ and \ M_1 + M_2 = 2\sum_{i=1}^k m_i + q - 1,$$
 (2.9)

Case 4: If there does not exist a simply closed curve γ of $\{x \in \Omega : u(x) = t\}$ for critical value t between γ_I and γ_E , then

$$M_1 \ge \sum_{i=1}^k m_i + 1, \ M_2 \ge \sum_{i=1}^k m_i + 1, \ and \ M_1 + M_2 = 2\sum_{i=1}^k m_i + q + 1.$$
 (2.10)

Proof. According to the fine analysis about the distributions of connected components of the superlevel sets $\{x \in \Omega : u(x) > t\}$ and sub-level sets $\{x \in \Omega : u(x) < t\}$ for some t, we can easily know that the Case 1, 2, 3, 4 of Lemma 2.5 implies Case 1, 2, 3, 4 respectively.

3 Proof of Theorem 1.1

In this section, we only give out the proofs of these two cases $z_1 < Z_1 \leq z_2 < Z_2$ and $z_1 < z_2 < Z_1 < Z_2$. It can be easily observed from the proofs that other cases can be dealt with in a similar way.

3.1 Proof for the case of $z_1 < Z_1 \le z_2 < Z_2$

In this subsection, we suppose that x_1, x_2, \dots, x_k are the interior critical points of u in Ω and $z_1 < u(x_{l+1}), \dots, u(x_k) < Z_1 \leq z_2 < u(x_1), \dots, u(x_l) < Z_2$. From the preparations in Section 2, we are now ready to present the proof of Theorem 1.1 for the case of $z_1 < Z_1 \leq z_2 < Z_2$.

Proof of Theorem 1.1 for the case of $z_1 < Z_1 \leq z_2 < Z_2$. (i) Case A_1 : If $z_1 < u(x_{l+1}) = \cdots = u(x_k) = t_0 < Z_1 \leq z_2 < u(x_1) = \cdots = u(x_l) = t_1 < Z_2$, by the results of Lemma 2.5, we know that

and

 $\sharp \{$ the simply connected components ω of the super-level set $\{x \in \Omega : u(x) > t_0\}$

such that ω meet the interior boundary $\gamma_I \Big\} \ge \sum_{i=l+1}^k m_i$.

By the strong maximum principle, we have that u exists at least $\sum_{i=1}^{l} m_i$ and $\sum_{i=l+1}^{k} m_i$ local minimal points and maximal points on γ_E and γ_I respectively. Hence, we have

$$\sum_{i=1}^{l} m_i \le N_2, \quad \sum_{i=l+1}^{k} m_i \le N_1,$$

then

$$\sum_{i=1}^{k} m_i \le N_1 + N_2$$

(ii) Case A_2 : If the values at critical points x_1, \dots, x_l are not totally equal or the values at critical points x_{l+1}, \dots, x_k are not totally equal. Next we need divide the proof of Case A_2 into two situations.

(1) Situation 1: If the values at critical points x_1, \dots, x_l are not totally equal, without loss of generality, we may assume that

$$z_{2} < u(x_{1}) = \dots = u(x_{j_{1}}) < u(x_{j_{1}+1}) = \dots = u(x_{j_{2}}) < \dots < \dots < u(x_{j_{n-1}+1}) = \dots = u(x_{j_{n}}),$$
(3.1)

where $x_1, \dots, x_{j_1}, \dots, x_{j_2}, \dots, x_{j_n}$ are different critical points in Ω , $j_n = l$ and $n \ge 2$. Next we put

 $E_j := \left\{ \omega : \text{open set } \omega \text{ is a connected component of } \{x \in \Omega : u(x) > u(x_j)\} \right\} \ (j = j_1, j_2, \cdots, j_n),$

and

 $G_{j_n} := \Big\{ \omega : \text{open set } \omega \text{ is a connected component of } \{x \in \Omega : u(x) > u(x_j)\} (j = j_1, j_2, \cdots, j_n) \\ \text{ such that there does not exist interior critical point in } \omega \Big\}.$

By the definition, we know that G_{j_n} consists of disjoint connected components. We set

 $|G_{j_n}| := \sharp \{ \omega : \omega \text{ is a connected component of } G_{j_n} \}.$

To illustrate G_{j_n} , let us consider an illustration for G_{j_2} . Assume that u has only three critical points $x_{j_1}, x_{j_1+1}, x_{j_2}$ such that $z_2 < u(x_{j_1}) < u(x_{j_1+1}) = u(x_{j_2})$ in Ω and the respective multiplicity $m_{j_1} = 1, m_{j_1+1} = 1, m_{j_2} = 2$. The distributions of elements of G_{j_2} is shown in Fig. 6.



Fig. 6. The distributions of elements of G_{j_2} .

By the definition of G_{j_n} , we know $|G_{j_2}| = 6$.

Now let us show that $|G_{j_s}| \ge \sum_{i=1}^{j_s} m_i + 1$ by induction on the number s. When s = 1, the result holds by Lemma 2.2. Suppose that $|G_{j_s}| \ge \sum_{i=1}^{j_s} m_i + 1$ for $1 \le s \le n-1$. Let s = n. Then, by (3.1) and the definition of E_j , we have

$$\{x_{j_{n-1}+1}, \cdots, x_{j_n}\} \subset \bigcup_{\omega \in E_{j_{n-1}}} \omega,$$

where ω is a connected component of $\{x \in \Omega : u(x) > u(x_{j_{n-1}})\}$.

Let us assume that $\{x_{j_{n-1}+1}, \dots, x_{j_n}\}$ are contained in exactly \hat{q} components $\omega_1, \dots, \omega_{\hat{q}}$. Then $x_{j_{n-1}+1}, \dots, x_{j_n}$ together with the corresponding level lines of $\{x \in \Omega : u(x) = u(x_{j_n})\}$ clustering round these points at least form \hat{q} connected sets. By the Case 2 of Lemma 2.5, we have

$$M := \sharp \left\{ \text{the connected components of } \{x \in \Omega : u(x) > u(x_{j_n}) \} \text{ in all } \omega_j \ (j = 1, \cdots, \widehat{q}) \right\}$$
$$\geq \sum_{i=j_{n-1}+1}^{j_n} m_i + \widehat{q}.$$

By using the definition of $|G_{j_n}|$ and the inductive assumption to $1 \le s \le n-1$, since $\{x_{j_{n-1}+1}, \dots, x_{j_n}\}$ are contained in exactly \hat{q} components $\omega_1, \dots, \omega_{\hat{q}}$, so when we calculate the number of $|G_{j_n}|$, the number of $|G_{j_{n-1}}|$ will be reduced by \hat{q} . Then we have

$$|G_{j_n}| = |G_{j_{n-1}}| + M - \widehat{q} \ge |G_{j_{n-1}}| + \left(\sum_{i=j_{n-1}+1}^{j_n} m_i + \widehat{q}\right) - \widehat{q} \ge \sum_{i=1}^{j_n} m_i + 1.$$

By the strong maximum principle and Lemma 2.2, we have that u has at least $\sum_{i=1}^{l} m_i + 1$ local maximum points on γ_E . Therefore, we obtain

$$\sum_{i=1}^{l} m_i + 1 \le N_2.$$

(2) Situation 2: If the values at critical points x_{l+1}, \dots, x_k are not totally equal. For convenience, we denote x_{l+1}, \dots, x_k and m_{l+1}, \dots, m_k by y_1, \dots, y_{k-l} and $\tilde{m}_1, \dots, \tilde{m}_{k-l}$ respectively. The proof is same as the case 2 of the proof of Theorem 1.1 in [13]. Without loss of generality, we may assume that

$$u(y_1) = \dots = u(y_{j_1}) < u(y_{j_1+1}) = \dots = u(y_{j_2}) < \dots < \dots < u(y_{j_{n-1}+1}) = \dots = u(y_{j_n}) < Z_1,$$
(3.2)

where $y_1, \dots, y_{j_1}, \dots, y_{j_2}, \dots, y_{j_n}$ are different critical points in Ω , $j_n = k - l$ and $n \ge 2$. Next we put

 $E_j := \left\{ \omega : \text{open set } \omega \text{ is a connected component of } \{x \in \Omega : u(x) > u(y_j)\} \right\} \ (j = j_1, j_2, \cdots, j_n),$

and

$$F_{j_n} := \Big\{ \omega : \text{open set } \omega \text{ is a connected component of } \{ x \in \Omega : u(x) < u(y_{j_1}) \} \text{ or } \omega \text{ is a connected component of } \{ x \in \Omega : u(y_{j_i}) < u(x) < u(y_{j_{i+1}}) \text{ for some } 1 \le i \le n-1 \} \Big\}.$$

By the definition, we know that F_{j_n} consists of disjoint connected components. We set

 $|F_{j_n}| := \sharp \{ \omega : \omega \text{ is a connected component of } F_{j_n} \}.$

To illustrate F_{j_n} , let us consider an illustration for F_{j_2} . Assume that u has only two critical points y_{j_1}, y_{j_2} such that $u(y_{j_1}) < u(y_{j_2}) < Z_1$ in Ω and the respective multiplicity $\tilde{m}_{j_1} = 1, \tilde{m}_{j_2} = 1$. The distributions of elements of F_{j_2} is shown in Fig. 7.



Fig. 7. The distributions of elements of F_{j_2} .

By the definition of F_{j_n} , we know $|F_{j_2}| = 4$.

Now let us show that $|F_{j_s}| \ge \sum_{i=1}^{j_s} \widetilde{m}_i + 1$ by induction on the number s. When s = 1, the result holds by Lemma 2.2. Suppose that $|F_{j_s}| \ge \sum_{i=1}^{j_s} \widetilde{m}_i + 1$ for $1 \le s \le n-1$. Let s = n. Then, by (3.2) and the definition of E_j , we have

$$\{y_{j_{n-1}+1}, \cdots, y_{j_n}\} \subset \bigcup_{\omega \in E_{j_{n-1}}} \omega,$$

where ω is a connected component of $\{x \in \Omega : u(x) > u(y_{j_{n-1}})\}$.

Let us assume that $\{y_{j_{n-1}+1}, \dots, y_{j_n}\}$ are contained in exactly \tilde{q} components $\omega_1, \dots, \omega_{\tilde{q}}$. Then $\{y_{j_{n-1}+1}, \dots, y_{j_n}\}$ together with the corresponding level lines of $\{x \in \Omega : u(x) = u(y_{j_n})\}$ clustering round these points at least form \tilde{q} connected sets. By Lemma 2.2, we have

$$\widetilde{M} := \sharp \Big\{ \text{the connected components of } \{ x \in \Omega : u(y_{j_{n-1}}) < u(x) < u(y_{j_n}) \} \text{ in all } \omega_j \ (j = 1, \cdots, \widetilde{q}) \Big\} \\ \ge \sum_{i=j_{n-1}+1}^{j_n} \widetilde{m}_i + \widetilde{q}.$$

By using the definition of $|F_{j_n}|$ and the inductive assumption to $1 \le s \le n-1$, then we have

$$\left|F_{j_n}\right| = \left|F_{j_{n-1}}\right| + \widetilde{M} \ge \left|F_{j_{n-1}}\right| + \left(\sum_{i=j_{n-1}+1}^{j_n} \widetilde{m}_i + \widetilde{q}\right) - \widetilde{q} \ge \sum_{i=1}^{j_n} \widetilde{m}_i + 1.$$

By the strong maximum principle, we have that u has at least $\sum_{i=1}^{k-l} \widetilde{m}_i + 1$ local minimal points on γ_I . Therefore, we obtain

$$\sum_{i=1}^{k-l} \widetilde{m}_i + 1 = \sum_{i=l+1}^k m_i + 1 \le N_1.$$

According to Case A_1 , Case A_2 and the assumption of

$$z_1 < u(x_{l+1}), \cdots, u(x_k) < Z_1 \le z_2 < u(x_1), \cdots, u(x_l) < Z_2,$$

then we have

$$\sum_{i=l+1}^{k} m_i \le N_1, \quad \sum_{i=1}^{l} m_i \le N_2,$$

that is

$$\sum_{i=1}^{k} m_i \le N_1 + N_2. \tag{3.3}$$

This completes the proof of Theorem 1.1 for the case of $z_1 < Z_1 \leq z_2 < Z_2$.

3.2 Proof for the case of $z_1 < z_2 < Z_1 < Z_2$

In this subsection, we suppose that x_1, x_2, \dots, x_k are the interior critical points of u in Ω and $z_1 < z_2 < Z_1 < Z_2$. From the preparations in Section 2, we are now ready to present the proof of Theorem 1.1 for the case of $z_1 < z_2 < Z_1 < Z_2$.

Proof of Theorem 1.1 for the case of $z_1 < z_2 < Z_1 < Z_2$. (i) Case B_1 : If $z_1 < u(x_{l+1}) = \cdots = u(x_k) = t_0 \le z_2 < Z_1 \le u(x_1) = \cdots = u(x_l) = t_1 < Z_2$ or $z_2 < u(x_{l+1}) = \cdots = u(x_k) = t_0 < Z_1 \le u(x_1) = \cdots = u(x_l) = t_1 < Z_2$ or $z_1 < u(x_{l+1}) = \cdots = u(x_k) = t_0 \le z_2 < u(x_1) = \cdots = u(x_l) = t_1 < Z_2$

 $u(x_l) = t_1 < Z_1$, by the Case 1 and Case 2 of Lemma 2.7 and the strong maximum principle, we know that

$$\sum_{i=1}^{l} m_i \le N_2, \qquad \sum_{i=l+1}^{k} m_i \le N_1,$$

then

$$\sum_{i=1}^k m_i \le N_1 + N_2.$$

(ii) Case B_2 : If $z_2 < u(x_1) = u(x_2) = \cdots = u(x_k) = t < Z_1$. By the Case 3 and Case 4 of Lemma 2.7 and the strong maximum principle, we have

$$\sum_{i=1}^{k} m_i \le N_1 + N_2$$

(iii) Case B_3 : For the other cases, the idea of proof is essentially same as the Case A_2 in section 3.1. Here we omit the proof.

4 Proof of Theorem 1.2

In this section, we assume that $z_2 \ge Z_1$. We investigate the geometric structure of interior critical point sets of a solution u in a planar bounded smooth multiply connected domain Ω with one interior boundary γ_I and the external boundary γ_E , where u has only N_1 (N_2) equal local maxima and N_1 (N_2) equal local minima relative to $\overline{\Omega}$ on γ_I (γ_E), i.e., the values of all local maximal (minimum) points on the corresponding boundary are equal. We develop a new method to prove that one of the following three results holds $\sum_{i=1}^k m_i = N_1 + N_2$ or $\sum_{i=1}^k m_i + 1 = N_1 + N_2$ or $\sum_{i=1}^k m_i + 2 = N_1 + N_2$, where $N_1, N_2 \ge 2$.

Proof of Theorem 1.2. We need divide the proof into three cases.

Firstly, we should show that u has at least one interior critical point in Ω . Suppose by contradiction that $|\nabla u| > 0$ in Ω . The strong maximum principle implies that u has no interior maximum point and minimal point in Ω , then we have

$$z_1 < u(x) < Z_2$$
 for any $x \in \Omega$.

According to the assumption of Theorem 1.2 and Remark 1.5, let q_1, \dots, q_{N_2} and p_1, \dots, p_{N_2} be the local maximal points and local minimal points relative to $\overline{\Omega}$ on γ_E respectively and $u(q_1) = \dots = u(q_{N_2}), u(p_1) = \dots = u(p_{N_2}).$

Without loss of generality, we may assume that there only exist two different local maximal points q_1, q_2 and minimum points p_1, p_2 on boundary γ_E . Note that u is monotonically decreasing on the connected components of γ_E from one maximal point to the near minimum point. Therefore, $\{x \in \Omega : u(x) = Z_2 - \epsilon\}$ exactly exists two level lines in Ω for any ϵ such that $0 < \epsilon < Z_2 - z_2$. Here, we need consider the following two situations:

Situation 1: If $Z_1 = z_2$, this is impossible, because this would imply that either: $u(x) = \min_{\gamma_E} \psi_2(x) = z_2$ in interior points of Ω , i.e., there exist two level lines of $\{x \in \Omega; u(x) = z_2\}$ connecting γ_I from p_1, p_2 respectively, this contradicts with the continuity of u, or: there exist two level lines of $\{x \in \Omega; u(x) = t_0 \in (z_2, Z_2)\}$ intersecting in Ω , i.e., there exist critical points in Ω (see Fig. 8).



Fig. 8. The distributions of some level lines $\{x \in A : u(x) = t \in (z_2, Z_2)\}$.

Situation 2: If $Z_1 < z_2$, this is also impossible. Because this would imply that there exist two level lines of $\{x \in \Omega; u(x) = t_0 \in (z_2, Z_2)\}$ intersecting in Ω , i.e., there exist critical points in Ω . Then u has at least one interior critical point in Ω .

(i) Case C_1 : Suppose that there exists a non-simply connected component ω of $\{x \in \Omega : u(x) < t\}$ for some $t \in (z_2, Z_2)$ such that ω meets γ_E and that the Case 3 of Lemma 2.5 does not occur for $t \in (z_1, Z_1)$. Next we need divide the proof of Case C_1 into four steps.

Step 1, the first "just right": we show that all critical values are equal to one of the two values. By Lemma 2.3, we suppose that the interior critical points of u are x_1, x_2, \dots, x_k and $z_1 < u(x_{l+1}), \dots, u(x_k) < Z_1 \leq z_2 < u(x_1), \dots, u(x_l) < Z_2$. Next we show that $z_1 < u(x_{l+1}) = \dots = u(x_k) < Z_1 \leq z_2 < u(x_1) = \dots = u(x_l) < Z_2$. Here, we only give the proof of $u(x_1) = \dots = u(x_l)$, the proof of $u(x_{l+1}) = \dots = u(x_k)$ is similar. By the assumption of Theorem 1.2, let q_1, \dots, q_{N_2} and p_1, \dots, p_{N_2} be the equal local maximal points and minimum points on γ_E , respectively. We set up the usual contradiction argument. Without loss of generality, we suppose that $z_2 < u(x_1) < u(x_2) < Z_2$ and that m_1, m_2 are the respective multiplicities of x_1, x_2 . Then, by Lemma 2.2, we know that any connected component of $\{x \in \Omega : u(x) > u(x_1)\}$ has to meet the boundary γ_E and $u(p_1) < u(x_1)$, where p_1 is the minimal point of some one connected component of $\{x \in \Omega : u(x) < u(x_2)\}$ meeting γ_E such that $u(x_1) < u(p_2) < u(x_2)$, where p_2 is the minimal point of connected component C of $\{x \in \Omega : u(x_1) < u(x_2) < u(x_2)\}$ meeting γ_E such that $u(x_1) < u(p_2) < u(x_2)$, where p_2 is the minimal point of connected component C on γ_E (see Fig. 9). Then $u(p_1) \neq u(p_2)$, which contradicts with the assumption of Theorem 1.2. This completes the proof of step 1.



Fig. 9. The distributions of the connected components.

Step 2, the second "just right": we show that critical points together with the corresponding level lines of $\{x \in \Omega : u(x) = t \text{ for some } t \in (z_2, Z_2)\}$ ($\{x \in \Omega : u(x) = t \text{ for some } t \in (z_1, Z_1)\}$), which meet $\gamma_E(\gamma_I)$, clustering round these points exactly form one connected set. Without loss of generality, we suppose by contradiction that x_1, x_2, \dots, x_l together with the level lines of $\{x \in \Omega : u(x) = t > z_2\}$ clustering round these points form two connected sets. Therefore, there exists a connected components of $\{x \in \Omega : u(x) < t\}$, which meets two parts γ_1, γ_2 of γ_E , denoting by A (see Fig. 10). Note that u is monotonically decreasing on the connected components of boundary γ_E from one maximal point to the near minimum point. Therefore, $\{x \in A : u(x) = t - \epsilon\}$ exactly exists two level lines in A for some ϵ such that $0 < \epsilon < t - z_2$. Here, we need consider the following two situations:

Situation 1: If $Z_1 = z_2$, this is impossible, because this would imply that either: $u(x) = \min_{\gamma_E} \psi_2(x) = z_2$ in interior points of A, i.e., there exist two level lines of $\{x \in A; u(x) = z_2\}$ connecting γ_I from p_1, p_2 respectively, this contradicts with the continuity of u, or: there exist two level lines of $\{x \in A; u(x) = t_0 \in (z_2, t)\}$ intersecting in A, i.e., there exist critical points in A (see Fig. 10).



Fig. 10. The distributions of some level lines $\{x \in A : u(x) = t - \epsilon, 0 < \epsilon < t - z_2\}$.

Situation 2: If $Z_1 < z_2$, this is also impossible. Because this would imply that there exist two level lines of $\{x \in A; u(x) = t_0 \in (z_2, t)\}$ intersecting in A, i.e., there exist critical points in A. This completes the proof of step 2.

Step 3, the third "just right": we show that every connected component of $\{x \in \Omega : u(x) > t > z_2\}$ ($\{x \in \Omega : u(x) < t \text{ for } t \in (z_1, Z_1)\}$) has exactly one global maximal (minimum) point on boundary γ_E (γ_I). In fact, we assume that some connected component B of $\{x \in \Omega : u(x) > t > z_2\}$ exists two global maximal points on boundary γ_E . According to u has N_2 equal local maxima and N_2 equal local minima on γ_E , then there must exist a minimum point \tilde{p} between the two maximal points on γ_E such that $u(\tilde{p}) = z_2$. Since $u(x) > t > z_2$ in B, by the continuity of solution u, this contradicts with the definition of connected component B. This completes the proof of step 3.

Step 4, By the results of step 2 and the results of Case 2 and Case 4 in Lemma 2.5, we have

and

On the other hand, by the results of step 3 and the strong maximum principle, therefore we obtain

$$\sum_{i=1}^{l} m_i + 1 = N_2 \text{ and } \sum_{i=l+1}^{k} m_i + 1 = N_1.$$

That is

$$\sum_{i=1}^{k} m_i + 2 = N_1 + N_2.$$
(4.1)

(ii) Case C_2 : Suppose that there exists a non-simply connected component ω of $\{x \in \Omega : u(x) < t\}$ for some $t \in (z_2, Z_2)$ and the external boundary γ of ω is a simply closed curve between γ_I and γ_E such that u has at least one critical point on γ . The idea of proof is essentially same as the proof of Case C_1 . Next we need divide the proof of Case C_2 into four steps.

Step 1, the first "just right": According to Lemma 2.3, we assume that $z_2 < u(x_1), \dots, u(x_l) < Z_2$. We show that $u(x_1) = \dots = u(x_l) = t$, i.e., we exclude the situation 1 of case A_2 in Theorem 1.1. The proof is same as the step 1 of the proof of Case C_1 .

Step 2, the second "just right": we show that x_1, x_2, \dots, x_l together with the corresponding level lines of $\{x \in \Omega : u(x) = t\}$ clustering round these points exactly form one connected set. The proof is same as the step 2 of the proof of Case C_1 .

Step 3, the third "just right": we show that every simply connected component ω of $\{x \in \Omega : u(x) < t\}$ has exactly one minimum point on γ_E , where ω meets the external boundary γ_E . In fact, we assume that some simply connected component ω of $\{x \in \Omega : u(x) < t\}$ exists two minimum points on boundary γ_E . According to u has only N_2 local maximal points and N_2 local minimum points on γ_E , then there must exist a maximal point \overline{q} between the two minimum points on γ_E such that $u(\overline{q}) = Z_2$. Since $u(x) < t < Z_2$ in ω , by the continuity of solution u, this contradicts with the definition of connected component ω . This completes the proof of step 3.

Step 4, By the results of step 2 and the results of step 1 of Case 1 in Lemma 2.5, we have

On the other hand, using the results of step 3 and the strong maximum principle, therefore we obtain

$$\sum_{i=1}^{l} m_i = N_2$$

(iii) Case C_3 : Suppose that there exists a simply closed curve γ of $\{x \in \Omega : u(x) = t\}$ for $t \in (z_1, Z_1)$ between γ_I and γ_E such that u has at least one critical point on γ and that $z_1 < u(x_{l+1}), \dots, u(x_k) < Z_1$. The proof is same as the proof of Case C_2 . Then we have

$$\sum_{i=l+1}^{k} m_i = N_1.$$

According to the above discussion, if Case C_2 and Case C_3 both occur, then we have

$$\sum_{i=1}^{k} m_i = N_1 + N_2. \tag{4.2}$$

In addition, if Case 1 and Case 4 of Lemma 2.5 occur, or Case 2 and Case 3 of Lemma 2.5 occur, then we have

$$\sum_{i=1}^{k} m_i + 1 = N_1 + N_2. \tag{4.3}$$

This completes the proof of Theorem 1.2.

In addition, according to Theorem 1.2 and Lemma 2.7, we can easily have the following results.

Corollary 4.1. Suppose that domain Ω satisfies the hypothesis of Theorem 1.1, $z_1 < z_2 < Z_1 < Z_2$ and that u is a non-constant solution of (1.2). In addition, suppose that u has only N_1 equal local maxima and N_1 equal local minima relative to $\overline{\Omega}$ on γ_I and that u has only N_2 equal local maxima and N_2 equal local minima relative to $\overline{\Omega}$ on γ_E . Then u has finite interior critical points, and one of the following three holds

$$\sum_{i=1}^{k} m_i = N_1 + N_2, \tag{4.4}$$

or

$$\sum_{i=1}^{k} m_i + 1 = N_1 + N_2, \tag{4.5}$$

or

$$\sum_{i=1}^{k} m_i + 2 = N_1 + N_2, \tag{4.6}$$

where m_i is as in Theorem 1.1.

Remark 4.2. Notice that in Theorem 1.2 and Corollary 4.1, the assumption of all the local maximal and local minimum points of u on γ_I and γ_E are the local maximal and local minimum points relative to $\overline{\Omega}$ is necessary. If there exist some local maximal and local minimum points only relative to γ_I and γ_E , there will be two counterexamples.

Counterexample 1. Suppose that the planar multiply connected domain Ω with the following two boundaries:

$$\gamma_I := \left\{ r(\theta) = R_1 + \sin(N_1\theta) \right\}, \quad \gamma_E := \left\{ r(\theta) = R_2 + \sin(N_2\theta) \right\},$$

where we describe the curves γ_I and γ_E by using polar coordinates (r, θ) , R_1 , $R_2 > 1$ and N_1 , N_2 are integers bigger than 1. We assume that $R_2 > R_1 + 2$, then γ_I and γ_E are disjoint. Let

$$u(x_1, x_2) := \log \sqrt{x_1^2 + x_2^2},$$

which is obviously a harmonic function in Ω . Note that $z_2 := \min_{\gamma_E} u = \log \sqrt{R_2 - 1} > Z_1 := \max_{\gamma_I} u = \log \sqrt{R_1 + 1}$, which is a condition demanded in Theorem 1.2. Moreover, $u|_{\gamma_I}$ and $u|_{\gamma_E}$ has only N_1 equal maxima and N_2 equal minima respectively, which are not the local maxima and minima relative to $\overline{\Omega}$. However, u has no critical points in Ω .

Counterexample 2. Suppose that the planar multiply connected domain Ω with the following two boundaries:

$$\gamma_I := \Big\{ r(\theta) = R_1 + \sin(N\theta) \Big\}, \quad \gamma_E := \Big\{ r(\theta) = R_2 + \sin(N\theta) \Big\},$$

where $R_2 > R_1 > 1$ such that $R_2 - R_1 < 2$ and N is an integer bigger than 1. Then γ_I and γ_E are disjoint. Let

$$u(x_1, x_2) := \log \sqrt{x_1^2 + x_2^2},$$

Note that $z_1 < z_2 < Z_1 < Z_2$, which is a condition demanded in Corollary 4.1. Moreover, $u|_{\gamma_I}$ and $u|_{\gamma_E}$ has only N equal maxima and N equal minima respectively, which are not the local maxima and minima relative to $\overline{\Omega}$. However, u has no critical points in Ω .

5 The geometric structure of interior critical zero points

In this section, we will study the geometric structure of interior critical zero points of solutions u to linear elliptic equations with nonhomogeneous Dirichlet boundary conditions in a multiply connected domain Ω .

5.1 Proof of Theorem 1.3

In this subsection, we will investigate the geometric structure of interior critical zero points of a solution u in a planar, bounded, smooth and multiply connected domain Ω with $u|_{\gamma_I} = H, u|_{\gamma_E} = \psi(x)$ and ψ is sign-changing and has \widetilde{N} zero points on γ_E .

Proof of Theorem 1.3. We need divide the proof into two cases.

Firstly, we should show that u has finite critical zero points in Ω , denoting by x_1, \dots, x_l , and we set m_i as the multiplicity of corresponding critical zero point x_i $(i = 1, \dots, l)$. Suppose by contradiction that u has infinite critical zero points in Ω . According to the results of Lemma 3.1 and Lemma 4.1 in [13], we know that every non-closed zero level line and γ_E have at least one intersection point. In addition, the theorem of Hartman and Wintner [20] shows that the interior critical points of u are isolated. Then there exist infinite non-closed zero level lines across these critical zero points and there are infinite zero points on γ_E , this contradicts with the assumption of Theorem 1.3.

(i) Case D_1 : If all the critical zero points x_1, \dots, x_l together with the corresponding zero level lines of $\{x \in \Omega : u(x) = 0\}$ clustering round these points form one connected set C. Next we need divide the proof of Case D_1 into two situations.

(1) Situation 1: If there exists a simply closed curve γ of $\{x \in \Omega : u(x) = 0\}$ between γ_I and γ_E such that u has at least one critical point on γ , that is $H \neq 0$. Then there are just $\sum_{i=1}^{l} m_i$ non-closed zero level lines across these critical zero points. In addition, when $H \neq 0$, we know that every non-closed zero level line and γ_E must have two intersection points. Then we have

$$2(\sum_{i=1}^{l} m_i) \le \widetilde{N},$$

$$\sum_{i=1}^{l} m_i \le \frac{\widetilde{N}}{2}.$$
(5.1)

that is

(2) Situation 2: Suppose that there does not exist a simply closed curve γ of $\{x \in \Omega : u(x) = 0\}$ between γ_I and γ_E such that u has at least one critical point on γ . Then there are just $(\sum_{i=1}^l m_i + 1)$ non-closed zero level lines across these critical zero points.

When $H \neq 0$, we know that every non-closed zero level line and γ_E must have two intersection points. Then we have

$$2(\sum_{i=1}^{l} m_i + 1) \le \tilde{N},$$

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}}{2} - 1.$$
 (5.2)

that is

When H = 0, by strong maximum principle, we know that above connected set C and γ_I have at most one intersection point. If above connected set C and γ_I do not have intersection points, then every non-closed zero level line and γ_E must have two intersection points and (5.2) holds. If above connected set C and γ_I have one intersection point, by the results of Lemma 4.1 in [13], then there must be a independent zero level line connecting γ_I and γ_E , and (5.2) still hold.

(ii) Case D_2 : If all the critical zero points x_1, \dots, x_l together with the corresponding zero level lines of $\{x \in \Omega : u(x) = 0\}$ clustering round these points form $q \ (q \ge 2)$ connected sets. Then there are just $(\sum_{i=1}^{l} m_i + q)$ non-closed zero level lines across these critical zero points. Next we need divide the proof of Case D_2 into two situations.

(3) Situation 3: When $H \neq 0$, we know that each non-closed zero level line and γ_E must have two intersection points. Then we have

$$2(\sum_{i=1}^{l} m_i + q) \le \widetilde{N},$$

that is

$$\sum_{i=1}^{l} m_i \le \frac{\widetilde{N}}{2} - q$$

(4) Situation 4: When H = 0, according to the strong maximum principle, we know that each connected set and γ_I have at most one intersection point. So we know that the number of intersection point of q connected sets and γ_E is at least $2(\sum_{i=1}^l m_i + q) - q$. Then we have

$$2(\sum_{i=1}^{l} m_i + q) - q \le \widetilde{N},$$

that is

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}-q}{2}$$

 $\sum_{i=1}^{l} m_i \le \frac{\widetilde{N}}{2},$

 $\sum_{i=1}^{l} m_i \le \frac{\tilde{N}}{2} - 1.$

Therefore, we have: If $H \neq 0$, we have

else
$$H = 0$$
, we have

5.2 Proof of Theorem 1.4

In this subsection, we will investigate the geometric structure of interior critical zero points of a solution u in a planar, bounded, smooth and multiply connected domain Ω with $u|_{\gamma_I} = \psi_1(x), u|_{\gamma_E} = \psi_2(x), \psi_1(x)$ and $\psi_2(x)$ are sign-changing and has \widetilde{N}_1 zero points and \widetilde{N}_2 zero points on γ_I and γ_E respectively.

Proof of Theorem 1.4. The proof of finiteness of critical zero points is same as Theorem 1.3, denoting by x_1, \dots, x_l , and we set m_i as the multiplicity of corresponding critical zero point x_i . Next we need divide the proof into two cases.

(i) Case E_1 : If all the critical zero points x_1, \dots, x_l together with the corresponding zero level lines of $\{x \in \Omega : u(x) = 0\}$ clustering round these points form one connected set and there exists a simply closed curve γ of $\{x \in \Omega : u(x) = 0\}$ between γ_I and γ_E such that u has at least two critical points on γ . Then there are just $\sum_{i=1}^{l} m_i$ non-closed zero level lines across these critical zero points. In addition, by the results of Lemma 2.2, we know that every non-closed zero level line and boundaries must have two intersection points. Then we have

$$2(\sum_{i=1}^{l} m_i) \le \widetilde{N}_1 + \widetilde{N}_2,$$
$$\sum_{i=1}^{l} m_i \le \frac{\widetilde{N}_1 + \widetilde{N}_2}{2}.$$

that is

(ii) Case E_2 : If Case E_1 does not occur. Then there are at least $(\sum_{i=1}^{l} m_i + 1)$ non-closed zero level lines across these critical zero points. According to the analysis of above Case E_1 , then we have

$$2(\sum_{i=1}^{l} m_i + 1) \le \widetilde{N}_1 + \widetilde{N}_2,$$

that is

$$\sum_{i=1}^{l} m_i \le \frac{\widetilde{N}_1 + \widetilde{N}_2}{2} - 1.$$

According to Theorem 1.3 and Theorem 1.4, we can easily have the following result.

Remark 5.1. Let Ω be a bounded, smooth and simply connected domain in \mathbb{R}^2 . Suppose that $\psi(x) \in C^1(\overline{\Omega})$ is sign-changing and that ψ has \widetilde{N} zero points on $\partial\Omega$. Let u be a non-constant solution of the following boundary value problem

$$\begin{cases} \sum_{i,j=1}^{2} a_{ij}(x) u_{x_i x_j}(x) + \sum_{i=1}^{2} b_i(x) u_{x_i}(x) = 0 & \text{in } \Omega, \\ u = \psi(x) & \text{on } \partial\Omega. \end{cases}$$
(5.3)

Then u has finite interior critical zero points, denoting by x_1, \dots, x_l , and

$$\sum_{i=1}^{l} m_i \le \frac{\tilde{N}}{2} - 1, \tag{5.4}$$

where m_i is as in Theorem 1.3.

References

- G.S. Alberti, G. Bal, M. Di Cristo, Critical points for elliptic equations with prescribed boundary conditions, Arch. Ration. Mech. Anal. 226 (2017), no. 1, 117-141.
- [2] G. Alessandrini, Critical points of solutions of elliptic equations in two variables, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 14 (1987) 229-256.
- [3] G. Alessandrini, R. Magnanini, The index of isolated critical points and solutions of elliptic equations in the plane, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 19 (4) (1992) 567-589.
- [4] G. Alessandrini, D. Lupo, E. Rosset, Local behavior and geometric properties of solutions to degenerate quasilinear elliptic equations in the plane, Appl. Anal. 50 (1993), no. 3-4, 191-215.
- [5] G. Alessandrini, R. Magnanini, Elliptic equations in divergence form, geometric critical points of solutions, and Stekloff eigenfunctions, SIAM J. Math. Anal. 25 (1994) 1259-1268.
- [6] J. Arango, A. Gómez, Critical points of solutions to quasilinear elliptic problems, Nonlinear Anal. 75 (2012) 4375-4381.

- [7] X. Cabré, S. Chanillo, Stable solutions of semilinear elliptic problems in convex domains, Selecta Math. (N.S.) 4 (1998) 1-10.
- [8] S. Cecchini, R. Magnanini, Critical points of solutions of degenerate elliptic equations in the plane, Calc. Var. Partial Differential Equations 39 (2010) 121-138.
- [9] J. Cheeger, A. Naber, D. Valtorta, Critical sets of elliptic equations, Comm. Pure Appl. Math. 68 (2015), no. 2, 173-209.
- [10] J.T. Chen, W.H. Huang, Convexity of capillary surfaces in the outer space, Invent. Math. 67 (1982) 253-259.
- [11] H.Y. Deng, H.R. Liu, L. Tian, Critical points of solutions for mean curvature equation in strictly convex and nonconvex domains, arXiv preprint arXiv:1712.08431 (2017).
- [12] H.Y. Deng, H.R. Liu, L. Tian, Uniqueness of critical points of solutions to the mean curvature equation with Neumann and Robin boundary conditions, arXiv preprint arXiv:1712.08454 (2017).
- [13] H.Y. Deng, H.R. Liu, L. Tian, Critical points of solutions to a quasilinear elliptic equation with nonhomogeneous Dirichlet boundary conditions, J. Differential Equations 265 (2018), no. 9, 4133-4157.
- [14] H. Donnelly, C. Fefferman, Nodal sets of eigenfunctions on Riemannian manifolds, Invent. Math. 93 (1988), 161-183.
- [15] H. Donnelly, C. Fefferman, Nodal sets for eigenfunctions of the Laplacian on surfaces, J. Amer. Math. Soc. 31 (1990), 333-353.
- [16] A. Enciso, D. Peralta-Salas, Critical points and level sets in exterior boundary problems, Indiana Univ. Math. J. 58 (2009) 1947-1968.
- [17] A. Enciso, D. Peralta-Salas, Critical points of Green's functions on complete manifolds, J. Differential Geom. 92 (2012) 1-29.
- [18] Q. Han, R. Hardt, F.-H. Lin, Geometric measure of singular sets of elliptic equations, Comm. Pure Appl. Math. 51 (1998), no. 11-12, 1425-1443.
- [19] R. Hardt, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, N. Nadirashvili, Critical sets of solutions to elliptic equations, J. Differential Geom. 51 (1999), no. 2, 359-373.
- [20] P. Hartman, A. Wintner, On the local behavior of solutions of non-parabolic partial differential equations, Amer. J. Math. 75, (1953) 449-476.
- [21] D. Kraus, Critical sets of bounded analytic functions, zero sets of Bergman spaces and nonpositive curvature, Proc. Lond. Math. Soc. 106 (3) (2013) 931-956.
- [22] F.H. Lin, Nodal sets of solutions of elliptic and parabolic equations, Comm. Pure Appl. Math. 44 (1991), no. 3, 287-308.
- [23] A. Logunov, Nodal sets of Laplace eigenfunctions: polynomial upper estimates of the Hausdorff measure, Ann. of Math. (2) 187 (2018), no. 1, 221-239.
- [24] A. Logunov, Nodal sets of Laplace eigenfunctions: proof of Nadirashvili's conjecture and of the lower bound in Yau's conjecture, Ann. of Math. (2) 187 (2018), no. 1, 241-262.
- [25] A. Logunov, E. Malinnikova, Nodal sets of Laplace eigenfunctions: estimates of the Hausdorff measure in dimension two and three, Operator Theory: Advances and Applications, (2018) Vol. 261, 333-344.
- [26] A. Naber, D. Valtorta, Volume estimates on the critical sets of solutions to elliptic PDEs, Comm. Pure Appl. Math. 70 (2017), no. 10, 1835-1897.
- [27] S. Sakaguchi, Uniqueness of critical point of the solution to some semilinear elliptic boundary value problem in \mathbb{R}^2 , Trans. Amer. Math. Soc. 319 (1990) 179-190.
- [28] S. Sakaguchi, Critical points of solutions to the obstacle problem in the plane, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 21 (4) (1994) 157-173.
- [29] L. Tian, X.P. Yang, Measure estimates of nodal sets of bi-harmonic functions, J. Differential Equations 256 (2014), no. 2, 558-576.
- [30] L. Tian, X.P. Yang, Measure upper bounds for nodal sets of eigenfunctions of the bi-harmonic operator, arXiv preprint arXiv: 1709.00153 (2017).