

Chiral photons from chiral gravitational waves

Keisuke Inomata^{1,2} and Marc Kamionkowski³

¹*ICRR, University of Tokyo, Kashiwa, 277-8582, Japan*

²*Kavli IPMU (WPI), UTIAS, University of Tokyo, Kashiwa, 277-8583, Japan*

³*Department of Physics and Astronomy, Johns Hopkins University,
3400 N. Charles Street, Baltimore, MD 21218, U.S.A.*

(Dated: November 14, 2018)

We show that a parity-breaking *uniform* (averaged over all directions on the sky) circular polarization of amplitude $V_{00} \simeq 2.6 \times 10^{-17} \Delta\chi(r/0.06)$ can be induced by chiral gravitational-wave (GW) background with tensor-to-scalar ratio r and chirality parameter $\Delta\chi$ (which is ± 1 for a maximally chiral background). We also show, however, that a uniform circular polarization can arise from a realization of a *non*-chiral GW background that spontaneously breaks parity. The magnitude of this polarization is drawn from a distribution of root-variance $\sqrt{\langle V_{00}^2 \rangle} \simeq 1.5 \times 10^{-18} (r/0.06)^{1/2}$ implying that the chirality parameter must be $\Delta\chi \gtrsim 0.12(r/0.06)^{-1/2}$ to establish that the GW background is chiral. Although these values are too small to be detected by any experiment in the foreseeable future, the calculation is a proof of principle that cosmological parity breaking in the form of a chiral gravitational-wave background can be imprinted in the chirality of the photons in the cosmic microwave background. It also illustrates how a seemingly parity-breaking cosmological signal can arise from parity-conserving physics.

The cosmic microwave background (CMB) is linearly polarized as a consequence of the anisotropic Thomson scattering of CMB photons that arises at linear order in the amplitude of cosmological perturbations [1]. This Thomson scattering does not, however, induce circular polarization. Recent work has shown that circular polarization can be generated, from primordial perturbations through post-recombination photon-photon scattering [2–6], at second order in the density-perturbation amplitude. Still, the *mean* value of the circular polarization, when averaged over the sky, is predicted to be zero. Since a nonzero circular polarization implies a particular handedness, the absence of a uniform circular polarization can be seen as a consequence of the absence of parity breaking in primordial density perturbations.

Here we show that a uniform circular polarization can arise if parity is broken in the form of a chiral primordial gravitational-wave (GW) background. Chiral GWs may arise if, for example, there is a Chern-Simons coupling of the inflaton to gravity [7], from gravity at a Lifshitz point [8], graviton self-couplings [9, 10], gaugeflation and chromo-natural inflation [11–14], Holst gravity [15], and in models that connect leptogenesis to primordial gravitational waves [16]. The circular polarization arises through interactions of linearly polarized CMB photons with CMB anisotropies along the line of sight to the surface of last scatter. The correlation between the photon anisotropy induced by the gravitational wave and the primordial linear polarization also induced by the gravitational wave leads, if the GW background is chiral, to a uniform circular polarization.

We also show, however, that a uniform circular polarization can arise as a statistical fluctuation—a cosmic variance—if the stochastic GW background is *not* chiral. Even if the expected value is zero, there will be some variation, in any given realization of the GW background, in the amplitudes of the right- and left-handed

gravitational waves. The theory predicts that the uniform circular polarization V_{00} (where the 00 indices indicate the $l = 0, m = 0$ spherical-harmonic coefficient of the circular-polarization pattern) is selected from a distribution centered on $V_{00} = 0$ but with nonzero variance. As we show, though, such a nonzero value for the uniform circular polarization cannot arise from primordial density perturbations, and so a nonzero uniform value would indicate a GW background, even if not necessarily chiral.

This work builds upon a detailed analysis presented in Ref. [17] of the circular polarization induced by GWs. That work builds upon a re-calculation [18], obtained with the TAM formalism [19, 20], of circular polarization induced by photon-photon scattering, in Ref. [6], which itself extends several earlier analyses [2–5] of this effect.

As discussed in that earlier work, circular polarization is induced through Faraday conversion as a linearly polarized light ray propagates through a medium with an anisotropic index-of-refraction tensor. The circular polarization $V(\hat{n})$ in direction \hat{n} is

$$V(\hat{n}) = \epsilon_{ac} P^{ab}(\hat{n}) \Phi_b^c(\hat{n}). \quad (1)$$

Here,

$$P_{ab}(\hat{n}) = \frac{1}{\sqrt{2}} \begin{pmatrix} Q(\hat{n}) & U(\hat{n}) \\ U(\hat{n}) & -Q(\hat{n}) \end{pmatrix}, \quad (2)$$

is the polarization tensor, whose components are the Stokes parameters $Q(\hat{n})$ and $U(\hat{n})$, and

$$\Phi_{ab}(\hat{n}) = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_Q(\hat{n}) & \phi_U(\hat{n}) \\ \phi_U(\hat{n}) & -\phi_Q(\hat{n}) \end{pmatrix}, \quad (3)$$

is a phase-shift tensor that describes the phase shifts induced by the index-of-refraction tensor. These are ob-

tained as line-of-sight integrals,

$$\phi_{Q,U}(\hat{\mathbf{n}}) = \frac{2}{c} \int_0^{\chi_{\text{LSS}}} \frac{d\chi}{1+z} \omega(\chi) n_{Q,U}(\hat{\mathbf{n}}\chi), \quad (4)$$

where z is redshift and χ_{LSS} the comoving distance to the last-scattering surface. Here, n_Q and n_U are components, in a plane transverse to the line of sight, of the index-of-refraction tensor,

$$n_{ab} = \delta_{ab} + \frac{1}{2}(\chi_{e,ab} + \chi_{m,ab}) = \begin{pmatrix} n_I + n_Q & n_U + in_V \\ n_U - in_V & n_I - n_Q \end{pmatrix}. \quad (5)$$

Finally, ϵ_{ab} is the antisymmetric tensor on the 2-sphere.

The polarization and phase-shift tensors can both be expanded,

$$P_{ab}(\hat{\mathbf{n}}) = \sum_{lm} \left[P_{lm}^E Y_{(lm)ab}^{TE}(\hat{\mathbf{n}}) + P_{lm}^B Y_{(lm)ab}^{TB}(\hat{\mathbf{n}}) \right], \quad (6)$$

$$\Phi_{ab}(\hat{\mathbf{n}}) = \sum_{lm} \left[\Phi_{lm}^E Y_{(lm)ab}^{TE}(\hat{\mathbf{n}}) + \Phi_{lm}^B Y_{(lm)ab}^{TB}(\hat{\mathbf{n}}) \right], \quad (7)$$

in terms of tensor spherical harmonics [21–23] $Y_{(lm)ab}^{TE}(\hat{\mathbf{n}})$ and $Y_{(lm)ab}^{TB}(\hat{\mathbf{n}})$.

The expansion coefficients in Eqs. (6) and (7) are random variables chosen from a distribution of zero mean and variances $C_l^{XX} = \langle |X_{lm}|^2 \rangle$ for $X = P^E, P^B, \Phi^E, \Phi^B$. In the absence of parity breaking, there are, moreover, cross-correlations $C_l^{P^E\Phi^E}$ and $C_l^{P^B\Phi^B}$. If there is no parity breaking, then there is no cross-correlation between the E modes and the B modes—the two have opposite parity and any cross-correlation would imply a preferred handedness. Expressions for all the power spectra C_l^{XY} are given as integrals over the primordial gravitational-wave power spectra in Ref. [17].

The uniform circular polarization is obtained by setting $l = m = 0$ in Eq. (60) in Ref. [17], and taking $G_{l_1 m_1 l_2 m_2}^{00} = \delta_{l_1 l_2} \delta_{-m_1 m_2} / \sqrt{4\pi}$:

$$V_{00} = \frac{1}{\sqrt{4\pi}} \sum_{lm} (P_{lm}^E \Phi_{lm}^{B*} - P_{lm}^B \Phi_{lm}^{E*}). \quad (8)$$

Taking the expectation value, over all realizations of metric perturbations, we find,

$$\langle V_{00} \rangle = \frac{1}{\sqrt{4\pi}} \sum_{lm} (C_l^{P^E\Phi^B} - C_l^{P^B\Phi^E}). \quad (9)$$

Through calculations that parallel those in Ref. [17] and lead to the power spectrum $C_l^{P^E\Phi^B}$ induced by photon-photon scattering, we infer power spectra,

$$\begin{aligned} C_l^{P^E\Phi^B} &= 4\pi A \int \frac{dk}{k} 2 \left(\frac{k^3}{2\pi^2} P^{(TE,TB)}(k) \right) \\ &\times \int_{\eta_{\text{ss}}}^{\eta_0} d\eta (1+z)^4 (\bar{a}_{2,2}^E(k, \eta)) \beta_l^{(2)}(k(\eta_0 - \eta)) \\ &\times \int_0^{\eta_0} d\eta g(\eta) \left(-\sqrt{6} \mathcal{P}^{(2)}(k, \eta) \right) \epsilon_l^{(2)}(k(\eta_0 - \eta)), \end{aligned} \quad (10)$$

where all the quantities are as defined in Ref. [17]. Here, $\mathcal{P}^{(TE,TB)}(k)$ is defined as the power spectrum for the cross-correlation between the amplitudes $h_{(lm)}^{k,TE}$ and $h_{(lm)}^{k,TB}$,

$$\langle h_{(lm)}^{k,TE} \left(h_{(lm)}^{k',TB} \right)^* \rangle = i \frac{(2\pi)^3}{k^2} \delta(k - k') P^{(TE,TB)}, \quad (11)$$

where $h_{(lm)}^{k,TE}$ and $h_{(lm)}^{k,TB}$ are the TE and TB modes in the TAM decomposition for the transverse-traceless metric perturbation. The i in Eq. (11) is canceled out by the i in front of $\beta_l^{(2)}$ in Eq. (41) in Ref. [17]. $C_l^{P^B\Phi^E}$ are also given by the same equation except for $\epsilon_l^{(2)} \leftrightarrow \beta_l^{(2)}$. In the absence of parity breaking, this power spectrum is zero, but if parity is broken in the form of a disparity between the amplitudes of the right- and left-circularly polarized gravitational waves, it will be nonzero. This can be seen by writing the amplitudes,

$$h_{(lm)}^{k,\pm} = \frac{1}{\sqrt{2}} \left(h_{(lm)}^{k,TE} \mp i h_{(lm)}^{k,TB} \right), \quad (12)$$

for the \pm (R and L) helicity-basis TAM waves (cf., Eq. (14) in Ref. [17]). Following Ref. [24], we define the chirality parameter $\Delta\chi$ through $P_{\pm}(k) = (1 \mp \Delta\chi) P_T(k)$, where $P_T(k)$ is the primordial GW power spectrum, and P_+ and P_- correspond to P_R and P_L in Ref [24], respectively. From this and Eq. (12), it follows that

$$P^{(TE,TB)}(k) = \Delta\chi P_T(k). \quad (13)$$

As a result, we find by numerical evaluation the uniform circular polarization to be

$$\begin{aligned} \langle V_{00} \rangle &= \sum_l \frac{2l+1}{\sqrt{4\pi}} (C_l^{P^E\Phi^B} - C_l^{P^B\Phi^E}) \\ &= 2.6 \times 10^{-17} \Delta\chi \left(\frac{r}{0.06} \right), \end{aligned} \quad (14)$$

for a gravitational-wave background with tensor-to-scalar ratio r and chirality parameter $\Delta\chi$.

So far, we have calculated the *expectation value* for V_{00} . However, the *observed* value of the circular polarization arises as the result of a single realization of a random field. The prediction is thus that V_{00} is selected from a random distribution, with some nonzero variance $\langle (\Delta V_{00})^2 \rangle = \langle V_{00}^2 \rangle - \langle V_{00} \rangle^2$, with an expectation value $\langle V_{00} \rangle$. Note that, in the absence of the parity breaking, $\langle (\Delta V_{00})^2 \rangle = \langle V_{00}^2 \rangle$ is satisfied. From Eq. (8), we find,

$$\begin{aligned} \langle (\Delta V_{00})^2 \rangle &= \sum_l \frac{(2l+1)}{4\pi} \left(C_l^{P^E P^E} C_l^{\Phi^B \Phi^B} + C_l^{P^B P^B} C_l^{\Phi^E \Phi^E} \right. \\ &\quad \left. - 2 C_l^{P^E \Phi^E} C_l^{P^B \Phi^B} \right. \\ &\quad \left. + \left(C_l^{P^E \Phi^B} \right)^2 + \left(C_l^{P^B \Phi^E} \right)^2 - 2 C_l^{P^E P^B} C_l^{\Phi^E \Phi^B} \right). \end{aligned} \quad (15)$$

There are contributions—those in the first and second lines—that arise even in the absence of parity breaking (while those that arise from parity-breaking physics are listed in the third line). Thus, the observed Universe can have a uniform circular polarization as a consequence of a parity-breaking realization of scalar and tensor metric perturbations, even in the absence of parity breaking in the underlying physics (in analogy to spontaneous symmetry breaking in particle theory). Such a possibility does not arise if there are only density perturbations (and thus only E-mode polarization and index-of-refraction tensor); a uniform circular polarization requires tensor perturbations (and thus B modes). Moreover, given current measurements of CMB polarization power spectra, and an upper limit $r \lesssim 0.06$ [25] to the tensor-to-scalar ratio, the power spectra for P^E and Φ^E are dominated by the contributions from scalar perturbations. As a result, we find numerically,

$$\langle (\Delta V_{00})^2 \rangle^{1/2} = 1.5 \times 10^{-18} \left(\frac{r}{0.06} \right)^{1/2}. \quad (16)$$

Comparing with Eq. (14), we see that the chirality parameter must be $\Delta\chi \gtrsim 0.12(r/0.06)^{-1/2}$ if detection of a uniform circular polarization can be attributed, at the 2σ level, to a chiral GW background. Although de-

tection of a nonzero uniform circular polarization with $V_{00} \lesssim 3 \times 10^{-18} (r/0.06)^{1/2}$ would not necessarily indicate a chiral GW background, it would still indicate the presence of tensor (or perhaps vector) perturbations.

In practice, these values of the circular polarization are too small to be detected in the foreseeable future. Still, the result is a proof of principle that gravitational-wave chirality can be imprinted in the chirality of the cosmic microwave background. Measurement of the circular polarization would, in the event of detection of a non-zero EB correlation in the CMB polarization, help distinguish a chiral-GW explanation [26] for such an effect [7, 24, 27] from cosmic birefringence. It would also complement probes of the GW chirality at nanoHertz frequencies [28] and at LIGO/LISA frequencies [29, 30].

We thank Daniel Green for useful discussion. K.I. is supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, Advanced Leading Graduate Course for Photon Science, and JSPS Research Fellowship for Young Scientists, and thanks Johns Hopkins University for hospitality. This work was supported at Johns Hopkins by NASA Grant No. NNX17AK38G, NSF Grant No. 1818899, and the Simons Foundation.

-
- [1] M. J. Rees, “Polarization and spectrum of the primeval radiation in an anisotropic Universe,” *Astrophys. J.* **153**, L1 (1968).
 - [2] I. Motie and S. S. Xue, “Euler-Heisenberg lagrangian and photon circular polarization,” *EPL* **100**, no. 1, 17006 (2012) [arXiv:1104.3555 [hep-ph]].
 - [3] R. F. Sawyer, “Photon-photon interactions can be a source of CMB circular polarization,” arXiv:1408.5434 [astro-ph.CO].
 - [4] D. Ejlli, “Magneto-optic effects of the cosmic microwave background,” *Nucl. Phys. B* **935**, 83 (2018) [arXiv:1607.02094 [astro-ph.CO]].
 - [5] M. Sadegh, R. Mohammadi and I. Motie, “Generation of circular polarization in CMB radiation via nonlinear photon-photon interaction,” *Phys. Rev. D* **97**, no. 2, 023023 (2018) [arXiv:1711.06997 [astro-ph.CO]].
 - [6] P. Montero-Camacho and C. M. Hirata, “Exploring circular polarization in the CMB due to conventional sources of cosmic birefringence,” *JCAP* **1808**, no. 08, 040 (2018) [arXiv:1803.04505 [astro-ph.CO]].
 - [7] A. Lue, L. M. Wang and M. Kamionkowski, “Cosmological signature of new parity violating interactions,” *Phys. Rev. Lett.* **83**, 1506 (1999) [astro-ph/9812088].
 - [8] T. Takahashi and J. Soda, “Chiral primordial gravitational waves from a Lifshitz point,” *Phys. Rev. Lett.* **102**, 231301 (2009) [arXiv:0904.0554 [hep-th]].
 - [9] J. M. Maldacena and G. L. Pimentel, “On graviton non-Gaussianities during inflation,” *JHEP* **1109**, 045 (2011) [arXiv:1104.2846 [hep-th]].
 - [10] M. M. Anber and L. Sorbo, “Non-Gaussianities and chiral gravitational waves in natural steep inflation,” *Phys. Rev. D* **85**, 123537 (2012) [arXiv:1203.5849 [astro-ph.CO]].
 - [11] A. Maleknejad, M. M. Sheikh-Jabbari and J. Soda, “Gauge fields and inflation,” *Phys. Rept.* **528**, 161 (2013) [arXiv:1212.2921 [hep-th]].
 - [12] P. Adshead, E. Martinec and M. Wyman, “Gauge fields and inflation: Chiral gravitational waves, fluctuations, and the Lyth bound,” *Phys. Rev. D* **88**, no. 2, 021302 (2013) [arXiv:1301.2598 [hep-th]].
 - [13] I. Obata and J. Soda, “Chiral primordial gravitational waves from dilaton induced delayed chromonatural inflation,” *Phys. Rev. D* **93**, no. 12, 123502 (2016) Addendum: [*Phys. Rev. D* **95**, no. 10, 109903 (2017)] [arXiv:1602.06024 [hep-th]].
 - [14] J. Bielefeld and R. R. Caldwell, “Chiral imprint of a cosmic gauge field on primordial gravitational waves,” *Phys. Rev. D* **91**, no. 12, 123501 (2015) [arXiv:1412.6104 [astro-ph.CO]].
 - [15] C. R. Contaldi, J. Magueijo and L. Smolin, “Anomalous CMB polarization and gravitational chirality,” *Phys. Rev. Lett.* **101**, 141101 (2008) [arXiv:0806.3082 [astro-ph]].
 - [16] S. H. S. Alexander, M. E. Peskin and M. M. Sheikh-Jabbari, “Leptogenesis from gravity waves in models of inflation,” *Phys. Rev. Lett.* **96**, 081301 (2006) [hep-th/0403069].
 - [17] K. Inomata and M. Kamionkowski, “Circular polarization of the cosmic microwave background from vector and tensor perturbations,” arXiv:1811.nnnnn [astro-ph.CO].
 - [18] M. Kamionkowski, “Circular polarization in a spherical basis,” *Phys. Rev. D* **97**, no. 12, 123529 (2018)

- [arXiv:1804.06412 [astro-ph.CO]].
- [19] L. Dai, M. Kamionkowski and D. Jeong, “Total angular momentum waves for scalar, vector, and tensor fields,” *Phys. Rev. D* **86**, 125013 (2012) [arXiv:1209.0761 [astro-ph.CO]].
- [20] L. Dai, D. Jeong and M. Kamionkowski, “Wigner-Eckart theorem in cosmology: Bispectra for total-angular-momentum waves,” *Phys. Rev. D* **87**, no. 4, 043504 (2013) [arXiv:1211.6110 [astro-ph.CO]].
- [21] M. Kamionkowski, A. Kosowsky and A. Stebbins, “Statistics of cosmic microwave background polarization,” *Phys. Rev. D* **55**, 7368 (1997) [astro-ph/9611125].
- [22] M. Zaldarriaga and U. Seljak, “An all sky analysis of polarization in the microwave background,” *Phys. Rev. D* **55**, 1830 (1997) [astro-ph/9609170].
- [23] M. Kamionkowski and E. D. Kovetz, “The quest for B modes from inflationary gravitational waves,” *Ann. Rev. Astron. Astrophys.* **54**, 227 (2016) [arXiv:1510.06042 [astro-ph.CO]].
- [24] V. Gluscevic and M. Kamionkowski, “Testing parity-violating mechanisms with cosmic microwave background experiments,” *Phys. Rev. D* **81**, 123529 (2010) [arXiv:1002.1308 [astro-ph.CO]].
- [25] P. A. R. Ade *et al.* [BICEP2 and Keck Array Collaborations], “BICEP2 / Keck Array X: Constraints on primordial gravitational waves using Planck, WMAP, and new BICEP2/Keck observations through the 2015 season,” [arXiv:1810.05216 [astro-ph.CO]].
- [26] S. Saito, K. Ichiki and A. Taruya, *JCAP* **0709**, 002 (2007) doi:10.1088/1475-7516/2007/09/002 [arXiv:0705.3701 [astro-ph]].
- [27] A. Ferté and J. Grain, “Detecting chiral gravity with the pure pseudospectrum reconstruction of the cosmic microwave background polarized anisotropies,” *Phys. Rev. D* **89**, no. 10, 103516 (2014) [arXiv:1404.6660 [astro-ph.CO]].
- [28] W. Qin, K. K. Boddy, M. Kamionkowski and L. Dai, “Pulsar-timing arrays, astrometry, and gravitational waves,” arXiv:1810.02369 [astro-ph.CO].
- [29] N. Seto and A. Taruya, “Measuring a parity violation signature in the early Universe via ground-based laser interferometers,” *Phys. Rev. Lett.* **99**, 121101 (2007) [arXiv:0707.0535 [astro-ph]].
- [30] T. L. Smith and R. R. Caldwell, “Sensitivity to a frequency-dependent circular polarization in an isotropic stochastic gravitational wave background,” *Phys. Rev. D* **95**, no. 4, 044036 (2017) [arXiv:1609.05901 [gr-qc]].