# B-type Landau-Ginzburg models with one-dimensional target

Calin Iuliu Lazaroiu<sup>1</sup> and Mehdi Tavakol<sup>2</sup>

 $^1$  Center for Geometry and Physics, Institute for Basic Science, Pohang 37673, Korea $^2$  School of mathematics and statistics, University of Melbourne, VIC 3010, Australia

E-mail: calin@ibs.re.kr, mehdi.tavakol@unimelb.edu.au

**Abstract.** We summarize the main results of our investigation of B-type topological Landau-Ginzburg models whose target is an arbitrary open Riemann surface. Such a Riemann surface need not be affine algebraic and in particular it may have infinite genus or an infinite number of Freudenthal ends. Under mild conditions on the Landau-Ginzburg superpotential, we give a complete description of the triangulated structure of the category of topological D-branes in such models as well as counting formulas for the number of topological D-branes considered up to relevant equivalence relations.

#### 1. Introduction

It is well-known [1, 2] that classical B-type topological Landau-Ginzburg (LG) models with Dbranes can be associated to any pair (X, W), where X is a non-compact Kähler manifold and  $W: X \to \mathbb{C}$  is a non-constant holomorphic function defined on X. When the canonical line bundle of X is holomorphically trivial and the critical set of W is compact, it is expected that the quantization of such models produces a non-anomalous two-dimensional topological field theory (2d TFT) which obeys the general axioms introduced in [3] and hence is characterized entirely by an open-closed *TFT datum*, an algebraic structure subject to certain axioms which encode the sewing constraints of the TFT. As shown in [2], path integral arguments lead to a proposal for the open-closed TFT datum of the quantum B-type LG model defined by such a pair (X, W), a proposal which was clarified and analyzed mathematically in [4, 5] (see [6] for a brief summary). When X is Stein and Calabi-Yau, this proposal simplifies as explained in [7] and summarized in [8].

As already pointed out in [4], an interesting particular case arises when X is complex onedimensional, i.e. an open Riemann surface (a non-compact Riemann surface without boundary, which we assume to be connected). Since any open Riemann surface is Stein and holomorphically parallelizable (hence Calabi-Yau), this fits into the class of models considered in [7], whose TFT datum admits the simplified description discussed in loc. cit. and summarized in [8].

In this contribution, we outline our investigation of B-type LG models with open Riemann surface target [9], which in turn relies on algebraic results derived for a large class of rings in references [10, 11]. We make free use of certain concepts and terminology which were introduced in [3] and [1, 2, 4, 7] and which are summarized in [6] and [8].

The open Riemann surfaces considered in this contribution are fully general. In particular, they can have infinite genus and an infinite number of Freudenthal ends<sup>1</sup>. As such, the topological LG models discussed herein are the most general non-anomalous B-type LG models with non-singular complex one-dimensional target.

## 2. Open Riemann surfaces

In this section, we recall some well-known facts regarding open Riemann surfaces.

**Definition 2.1.** A Riemann surface (in the sense of Weyl-Radó) is a Hausdorff, borderless and connected complex manifold  $\Sigma$  such that dim<sub> $\mathbb{C}</sub> \Sigma = 1$ .</sub>

**Theorem 2.2** (Radó). The following mutually-equivalent statements hold for any Riemann surface  $\Sigma$ :

- (a)  $\Sigma$  is paracompact.
- (b)  $\Sigma$  has a countable basis.
- (c)  $\Sigma$  is countable at infinity.

Definition 2.3. A Riemann surface is called open if it is non-compact.

## 2.1. The topological type of open Riemann surfaces

Unlike the compact case, the topological classification of open Riemann surfaces is quite involved, since such a surface can have infinite genus as well as an infinity of Freudenthal ends (a.k.a. "ideal points"). The ends of Riemann surfaces were studied in the classical work of Kerékjártó [12] and Stoilow [13] (see [14] for a summary in modern mathematical language).

# Definition 2.4.

• The transfinite genus of  $\Sigma$  is the cardinal number:

$$g(\Sigma) = \frac{1}{2} \mathrm{rk}_{\mathbb{Z}} \mathrm{H}^1(\Sigma, \mathbb{Z})$$
 .

• The ideal boundary  $\partial_{\infty} \Sigma$  of  $\Sigma$  is the set of Freudenthal ends of  $\Sigma$ , endowed with its natural topology.

**Proposition 2.5** (Kerékjártó). The following statements hold for any open Riemann surface  $\Sigma$ :

1.  $g(\Sigma)$  is finite or countable.

2.  $\partial_{\infty}\Sigma$  is a finite or countable compact Hausdorff topological space which is totally disconnected.

# Remarks.

- 1.  $\partial_{\infty}\Sigma$  can be a Cantor space.
- 2. Adjoining  $\partial_{\infty}\Sigma$  to  $\Sigma$  produces the so-called *Kerékjártó-Stoilow compactification*  $\hat{\Sigma}$  of  $\Sigma$  (see [12, 13, 14]).

There exists a natural disjoint union decomposition  $\partial_{\infty}\Sigma = \partial_{\infty}^{1}\Sigma \sqcup \partial_{\infty}^{2}\Sigma$  where the ends belonging to  $\partial_{\infty}^{1}\Sigma$  and  $\partial_{\infty}^{2}\Sigma$  are called of *first* and *second* kind.

<sup>&</sup>lt;sup>1</sup> An open Riemann surface coincides with the analyticization of a complex affine curve iff it has finite genus and a finite number of ends, in which case it can be obtained by removing a finite set of points from a compact Riemann surface.

#### 2.2. The complex geometry of open Riemann surfaces

The following statements summarize classical results due to Kerékjártó and Stoilow:

**Theorem 2.6** (Kerékjártó). Let  $\Sigma$  and  $\Sigma'$  be two open Riemann surfaces. Then the following statements are equivalent:

(a)  $\Sigma$  and  $\Sigma'$  are unoriented-homeomorphic.

(b)  $g(\Sigma) = g(\Sigma')$  and there exists a homeomorphism  $\psi: \partial_{\infty}\Sigma \to \partial_{\infty}\Sigma'$  such that  $\psi(\partial_{\infty}^{1}\Sigma) = \partial_{\infty}^{1}\Sigma'$ .

**Theorem 2.7** (Richards-Stoilow). The following statements hold:

- (i) The unoriented homeomorphism type of an open Riemann surface  $\Sigma$  is entirely determined by the triplet  $(g(\Sigma), \partial_{\infty}^{1}\Sigma, \partial_{\infty}\Sigma)$ .
- (ii) Consider any triplet  $(g, F^1, F)$  with  $g \in \mathbb{Z}_{\geq 0} \sqcup \{\aleph_0\}$ , F a non-empty compact, Hausdorff and totally disconnected countable topological space and  $F^1 \subset F$  a (possibly empty) subset of F, endowed with the topology induced from F. Then there exists an open Riemann surface  $\Sigma$  with  $g(\Sigma) = g$ ,  $\partial_{\infty}\Sigma \simeq_{\text{homeo}} F$  and  $\partial_{\infty}^1\Sigma \simeq_{\text{homeo}} F^1$ .

The following results are crucial in the context of B-type LG models:

Theorem 2.8 (Behnke-Stein). Every open Riemann surface is Stein.

**Theorem 2.9** (Grauert-Röhrl). Any holomorphic vector bundle on an open Riemann surface  $\Sigma$  is holomorphically trivial. In particular, the analytic Picard group  $\operatorname{Pic}(\Sigma)$  vanishes and the canonical line bundle  $K_{\Sigma}$  is holomorphically trivial.

It follows that any open Riemann surface is Stein (and hence Kählerian) and holomorphically Calabi-Yau. Hence any open Riemann surface can be used as the target space of a B-type LG model whose TFT datum admits the simplified description discussed in [7] and summarized in [8]. Since the holomorphic tangent bundle  $T\Sigma$  is trivial by Grauert-Röhrl theorem, any open Riemann surface is holomorphically parallelizable.

**Proposition 2.10** (various authors). Every non-compact Riemann surface  $\Sigma$  admits a holomorphic embedding in  $\mathbb{C}^3$  and a holomorphic immersion in  $\mathbb{C}^2$ .

**Remark.** It is not known if any open Riemann surface admits a proper holomorphic embedding in  $\mathbb{C}^2$ .

2.3. The ring  $\mathcal{O}(\Sigma)$ 

Let  $\Sigma$  be an open Riemann surface. This subsection recalls certain results regarding the ring  $\emptyset(\Sigma)$  of complex-valued holomorphic functions defined on  $\Sigma$ .

**Theorem 2.11** (Bers). Let  $\Sigma_1$  and  $\Sigma_2$  be two connected non-compact Riemann surfaces. Then  $\Sigma_1$  and  $\Sigma_2$  are biholomorphic iff their rings of holomorphic functions  $\emptyset(\Sigma_1)$  and  $\emptyset(\Sigma_2)$  are isomorphic as unital  $\mathbb{C}$ -algebras.

**Theorem 2.12** (Iss'sa). Let  $\Sigma_1$  and  $\Sigma_2$  be two connected non-compact Riemann surfaces. Then  $\Sigma_1$  and  $\Sigma_2$  are biholomorphic iff their fields of meromorphic functions  $\mathcal{M}(\Sigma_1)$  and  $\mathcal{M}(\Sigma_2)$  are isomorphic as unital  $\mathbb{C}$ -algebras.

**Proposition 2.13** (Henriksen-Alling). The following statements hold:

- 1. The cardinal Krull dimensions of all open Riemann surfaces are equal to each other (denote this cardinal number by  $\mathbf{k}$ ).
- 2. We have  $\mathbf{k} \geq 2^{\aleph_1}$ .

**Definition 2.14.** An elementary divisor domain (EDD) is an integral domain R such that any matrix with coefficients from R admits a Smith normal form.

**Theorem 2.15** (Helmer-Henriksen-Alling). For any open Riemann surface  $\Sigma$ , the ring  $\emptyset(\Sigma)$  is an elementary divisor domain.

## 2.4. Special uniformizers

Let  $\Sigma$  be an open Riemann surface and  $\operatorname{ord}_x : \mathcal{M}(\Sigma) \to \mathbb{Z}$  be the map which assigns to a meromorphic function  $f \in \mathcal{M}(\Sigma)$  its order at the point  $x \in \Sigma$ .

**Definition 2.16.** A special local uniformizer for  $\Sigma$  at the point  $x \in \Sigma$  is a meromorphic function  $t_x \in \mathcal{M}(\Sigma)$  such that  $\operatorname{ord}_x(t_x) = 1$  and  $\operatorname{ord}_y(t_x) = 0$  for all  $y \in \Sigma \setminus \{x\}$ . Notice that  $t_x \in \mathcal{O}(\Sigma)$ .

**Proposition 2.17.** The special local uniformizers of  $\Sigma$  coincide with the prime elements of  $\emptyset(\Sigma)$ . In particular, a special local uniformizer at x is determined up to multiplication by a unit of  $\emptyset(\Sigma)$  (i.e. a nowhere-vanishing element of  $\emptyset(\Sigma)$ ).

# 2.5. Critically-finite elements of $\mathcal{O}(\Sigma)$

Most of our results for B-type Landau-Ginzburg models with open Riemann surface target depend on the assumption that the Landau-Ginzburg superpotential is *critically finite* in the sense discussed below. The notion of critically finite element was introduced in [10, 11], which studied categories of finite rank matrix factorizations for Bézout and elementary divisor domains.

**Definition 2.18.** A divisor  $r \in \mathcal{O}(\Sigma)$  of  $f \in \mathcal{O}(\Sigma)$  is called critical if  $r^2 | f$ .

**Definition 2.19.** A non-zero non-unit  $f \in \mathcal{O}(\Sigma)$  is called:

- non-critical, if f has no critical divisors;
- critically-finite, if f has a factorization of the form:

$$f = f_0 f_c$$
 with  $f_c = p_1^{n_1} \dots p_N^{n_N}$ 

where  $N \ge 1$ ,  $n_i \ge 2$ ,  $p_1, \ldots, p_N \in \mathcal{O}(\Sigma)$  are critical prime divisors of f with  $(p_i) \ne (p_j)$ for  $i \ne j$  and  $f_0 \in \mathcal{O}(\Sigma)$  is non-critical and coprime with  $f_c$ .

**Lemma 2.20.** Let  $f \in \mathcal{O}(\Sigma)$  be a non-zero element and D(f) be the analytic divisor of f. Then the following statements hold:

- 1. f is a unit of  $\mathcal{O}(\Sigma)$  iff D(f) = 0.
- 2. f is a prime element of  $\emptyset(\Sigma)$  (i.e. a special local uniformizer of  $\Sigma$ ) iff D(f) = x for some  $x \in \Sigma$ .
- 3. f is non-critical iff D(f) is multiplicity-free at any point in its support, i.e. iff it has the form:

$$D(f) = \sum_{x \in \mathcal{Z}(f)} x \quad .$$

where  $\mathcal{Z}(f)$  denotes the zero set of f (which is at most countable).

- 4. If  $g \in \mathcal{O}(\Sigma)$  is another holomorphic function, then f|g iff  $D(f) \leq D(g)$ .
- 5. f is critically-finite iff f = gh, where  $g \in \mathcal{O}(\Sigma)$  is non-critical and  $h \in \mathcal{O}(\Sigma)$  satisfies:

$$D(h) = \sum_{x \in \mathcal{Z}(h)} n_x \cdot x \quad ,$$

with  $\mathcal{Z}(h)$  a non-empty finite set and  $n_x > 1$  for every  $x \in \mathcal{Z}(h)$ .

#### 3. B-type Landau-Ginzburg models with one-dimensional target

**Definition 3.1.** A Landau-Ginzburg (LG) pair is a pair (X, W) such that:

- 1. X is a non-compact complex and Kählerian connected manifold of dimension d > 0, which is holomorphically Calabi-Yau in the sense that its holomorphic canonical line bundle  $K_X$  is holomorphically trivial.
- 2.  $W \in \mathcal{O}(X)$  is a non-constant holomorphic complex-valued function defined on X, which is called the Landau-Ginzburg superpotential.

Given an LG pair (X, W), let  $Z_W \stackrel{\text{def.}}{=} \{x \in X | (\partial W)(x) = 0\}$  be the critical set of W. Since any Stein manifold is Kählerian, a particular case of LG pair is obtained by taking X to be Stein. This situation was studied in [7] (see [8] for a summary).

**Definition 3.2.** A one-dimensional LG pair is an LG pair (X, W) such that  $\dim_{\mathbb{C}} X = 1$ .

Since any open Riemann surface is Stein and holomorphically Calabi-Yau (see Subsection 2.2), we have:

**Proposition 3.3.** An LG pair is one-dimensional iff X coincides with an open Riemann surface  $\Sigma$ .

Since any compact analytic subset of a Stein manifold is finite, the critical set of  $W \in \mathcal{O}(\Sigma)$ is compact iff it is finite. Let  $(\Sigma, W)$  be a one-dimensional LG pair with compact (and hence finite) critical set  $Z_W$ . For any  $p \in Z_W$ , let:

$$\mathbf{M}(\hat{W}_p) \stackrel{\text{def.}}{=} \frac{\mathcal{O}_{\Sigma,p}}{\langle \hat{W}'_p \rangle}$$

denote the **analytic** Milnor algebra of the analytic function germ  $\hat{W}_p$  of W at p and  $\nu_p \stackrel{\text{def.}}{=} \dim_{\mathbb{C}} \mathcal{M}(\hat{W}_p)$  denote the analytic Milnor number of W at p. Here  $\langle W'_p \rangle$  is the ideal of  $\mathcal{O}(\Sigma)$  generated by  $\frac{\mathrm{d}W(p)}{\mathrm{d}z}$ , where z is any local holomorphic coordinate of  $\Sigma$  centered at p (this ideal is independent of the choice of such a local coordinate). Let  $t_p \in \mathcal{M}(\Sigma)$  be a special local uniformizer of  $\Sigma$  at p.

**Proposition 3.4.** [9] We have  $M(\hat{W}_p) \simeq_{\emptyset(\Sigma)} \oplus_{i=1}^N \mathcal{O}_{\Sigma,p} / \langle \hat{t}_p^{\nu_p} \rangle$ .

**Proposition 3.5.** [9] The bulk state space of the open-closed TFT defined by  $(\Sigma, W)$  can be identified with the direct sum of analytic Milnor algebras:

$$\mathcal{H} \simeq_{\mathcal{O}(X)} \oplus_{p \in Z_W} \mathcal{M}(\hat{W}_p)$$

In particular, we have:

$$\dim_{\mathbb{C}} \mathcal{H} = \sum_{p \in Z_W} \nu_p$$

#### 3.1. The topological D-brane category

For any unital commutative ring R, let MF(R, W) denote the category of finite rank matrix factorizations of W over R and HMF(R, W) denote its total cohomology category (which is  $\mathbb{Z}_2$ -graded and R-linear). Let  $hmf(R, W) \stackrel{\text{def.}}{=} HMF^{\hat{0}}(R, W)$  denote the sub-category obtained from HMF(R, W) by keeping only the even morphisms. It is well-known that hmf(R, W) has a natural structure of triangulated category with involutive shift functor. **Proposition 3.6.** [9] Let  $(\Sigma, W)$  be a one-dimensional LG pair. Then the D-brane category  $\mathcal{T}$  of the associated 2d TFT (i.e. of the quantum LG model with target (X, W)) can be identified with:

$$\mathcal{T} \simeq_{\mathcal{O}(\Sigma)} \mathrm{HMF}(\mathcal{O}(\Sigma), W)$$
.

**Definition 3.7.** A matrix factorization (M, D) of W over  $\emptyset(\Sigma)$  is called elementary if  $\operatorname{rk} M^{\hat{0}} = \operatorname{rk} M^{\hat{1}} = 1$ .

In the limit  $W \to 0$ , the B-type LG model parameterized by (X, W) reduces to a B-type nonlinear sigma model with non-compact target X. Intuitively, elementary factorizations correspond to 'elementary D-branes', i.e. those topological D-branes which are obtained by 'topological tachyon condensation' in a pair formed of a single brane and a single anti-brane of this nonlinear sigma model, where tachyon condensation is driven by turning on the Landau-Ginzburg superpotential W. See [15, 16, 17, 18] for the origin of this interpretation in open topological string field theory.

**Definition 3.8.** The  $\mathbb{Z}_2$ -graded cocycle category  $\text{ZMF}(\emptyset(\Sigma), W)$  has the same objects as  $\text{MF}(\Sigma, W)$  but its morphisms are the closed morphisms of  $\text{MF}(\emptyset(\Sigma), W)$ .

Let  $\operatorname{zmf}(\emptyset(\Sigma), W)$  be the even subcategory of  $\operatorname{ZMF}(\emptyset(\Sigma), W)$ . We say that two matrix factorizations of W are *strongly isomorphic* if they are isomorphic in  $\operatorname{zmf}(\emptyset(\Sigma), W)$ . Notice that an elementary factorization of W is indecomposable in the additive category  $\operatorname{zmf}(\emptyset(\Sigma), W)$ , but it need not be indecomposable in  $\mathcal{T}^{\hat{0}} = \operatorname{hmf}(\emptyset(\Sigma), W)$ .

### 3.2. Critically-finite superpotentials

The following result characterizes critically-finite superpotentials:

**Proposition 3.9.** [9] Let  $(\Sigma, W)$  be a one-dimensional LG pair. Then the following statements are equivalent:

- (a) W is a critically-finite element of  $\mathcal{O}(\Sigma)$ .
- (b) The intersection  $\mathcal{Z}(W) \cap Z_W$  is finite.
- (c) The divisor of W has the form:

$$D(W) = D_0 + \sum_{i=1}^N n_i x_i$$
,

where  $D_0$  is either the trivial divisor or an effective divisor whose multiplicity at every point of its support is one, the symbols  $x_1, \ldots, x_N$  (with  $N \ge 1$ ) denote a finite collection of points of  $\Sigma$  which do not belong to the support of  $D_0$  and  $n_i \ge 2$  for all  $i = 1, \ldots, N$ .

(d) We have  $W = W_0 W_c$ , where  $W_0$  has no zeros or only simple zeros,  $W_c$  has a finite number of zeros, each of which has multiplicity at least two and  $W_0$ ,  $W_c$  have no common zero.

In particular, any holomorphic function W with finite critical set is a critically-finite element of  $\mathcal{O}(\Sigma)$ .

**Remark.** Any critically-finite superpotential  $W \in \mathcal{O}(\Sigma)$  can be written as:

$$W = W_0 W_c$$
, with  $W_c = \prod_{i=1}^N t_{x_i}^{n_i}$ , (1)

where:

- (i)  $W_0 \in \mathcal{O}(\Sigma)$  has a finite or countable number of zeros, all of which are simple.
- (ii)  $x_1, \ldots, x_N$  (with  $N \ge 1$ ) are distinct points of  $\Sigma$  and  $t_{x_i}$  are special uniformizers at these points.
- (iii) None of the points  $x_i$  is a zero of  $W_0$ .

In this case, we have  $D_0 = D(W_0)$  and we set  $D_c \stackrel{\text{def.}}{=} \sum_{i=1}^N n_i x_i$ .

3.3. Primary factorizations and the Krull-Schmidt property of  $hmf(\mathcal{O}(\Sigma), W)$ 

Recall that an additive category  $\mathcal{A}$  is called *Krull-Schmidt* if any object of  $\mathcal{A}$  decomposes as a finite sum of indecomposable objects with quasi-local endomorphism rings. We say that a matrix factorization of W is *trivial* if it is a zero object in hmf( $\Sigma, W$ ).

**Proposition 3.10.** [9] Any elementary factorization of W over  $\emptyset(\Sigma)$  is strongly isomorphic to an elementary factorization of the form  $e_v \stackrel{\text{def.}}{=} (\emptyset(\Sigma)^{1|1}, D_v)$ , where  $v \in \emptyset(\Sigma)$  is a divisor of Wand  $D_v \stackrel{\text{def.}}{=} \begin{bmatrix} 0 & v \\ u & 0 \end{bmatrix}$ , with  $u \stackrel{\text{def.}}{=} W/v \in \emptyset(\Sigma)$ .

**Definition 3.11.** An element  $f \in \mathcal{O}(\Sigma)$  is called primary if it is a power of a prime element of  $\mathcal{O}(\Sigma)$ . An elementary factorization  $e_v$  of W is called primary if v is a primary divisor of W.

If  $e_v$  is a primary factorization of W over  $\mathcal{O}(\Sigma)$ , then we have  $v = at_x^k$  for some  $a \in \mathcal{O}(\Sigma)^{\times}$ , some  $k \in \mathbb{Z}_{>0}$  and some  $x \in \Sigma$ . In this case, W has a zero of order at least k at the point  $x \in \Sigma$ .

**Theorem 3.12.** [9] Suppose that W is critically-finite. Then the additive category  $\operatorname{hmf}(\emptyset(\Sigma), W)$  is a Krull-Schmidt category whose non-zero indecomposables are the nontrivial primary matrix factorizations of W.

Hence any finite rank matrix factorization (in particular, any non-primary elementary factorization) of W decomposes into primary factorizations in the category hmf( $\emptyset(\Sigma), W$ ). The primary factorizations of W play the role of 'simple' (a.k.a 'irreducible') topological D-branes of the B-type LG model.

## 3.4. The triangulated structure of $hmf(\mathcal{O}(\Sigma), W)$

Let R be a unital commutative ring. Let  $\text{mod}_R$  be the Abelian category of finitely-generated R-modules.

#### Definition 3.13.

(i) The projectively stable category  $\underline{\mathrm{mod}}_R$  has the same objects as  $\mathrm{mod}_R$  and modules of morphisms given by:

 $\underline{\operatorname{Hom}}_{R}(M,N) \stackrel{\text{def.}}{=} \operatorname{Hom}_{R}(M,N) / \mathcal{P}_{R}(M,N) \quad \forall M,N \in \operatorname{Ob}(\operatorname{mod}_{R}) ,$ 

where  $\mathcal{P}_R(M, N) \subset \operatorname{Hom}_R(M, N)$  consists of those morphisms of  $\operatorname{mod}_R$  which factor through a projective module of finite rank.

(ii) The injectively stable category  $\overline{\text{mod}}_R$  has the same objects as  $\text{mod}_R$  and modules of morphisms given by:

 $\overline{\operatorname{Hom}}_R(M,N) \stackrel{\text{def.}}{=} \operatorname{Hom}_R(M,N) / \mathcal{I}_R(M,N) \quad \forall M,N \in \operatorname{Ob}(\operatorname{mod}_R) ,$ 

where  $\mathcal{I}_R(M, N) \subset \operatorname{Hom}_R(M, N)$  consists of those morphisms of  $\operatorname{mod}_R$  which factor through an injective module of finite rank.

**Remark.** When R is a self-injective ring, the categories  $\underline{mod}_R$  and  $mod_R$  are equivalent to each other and the category  $\underline{mod}_R$  is naturally triangulated since in this case  $mod_R$  is a Frobenius category (i.e. an exact category whose projective and injective objects coincide).

The following result gives a complete description of the triangulated structure of the category  $\mathcal{T}^{\hat{0}} = \operatorname{hmf}(\mathcal{O}(\Sigma), W)$  for the case when W is critically finite:

**Theorem 3.14.** [9] Let  $W \in \mathcal{O}(\Sigma)$  be a critically-finite superpotential of the form (1). Then the following statements hold:

- 1. The ring  $A_i \stackrel{\text{def.}}{=} \emptyset(\Sigma)/\langle t_{x_i}^{n_i} \rangle$  is Artinian and Frobenius (hence also self-injective) for all  $i = 1, \ldots, N$ .
- 2. For each i = 1, ..., N, there exist equivalences of triangulated categories:

$$D_{\text{Sing}}(A_i) \simeq \underline{\text{mod}}_{A_i}$$

where  $\underline{\mathrm{mod}}_{A_i}$  denotes the projectively stable category of finitely-generated  $A_i$ -modules.

3. The triangulated category  $\underline{\text{mod}}_{A_i}$  is Krull-Schmidt with non-zero indecomposable objects given by the  $A_i$ -modules:

$$V_i^{(k)} \stackrel{\text{def.}}{=} \mathcal{O}(\Sigma)/\langle t_{x_i}^k \rangle \simeq \langle t_{x_i}^{n_i-k} \rangle/\langle t_{x_i}^{n_i} \rangle \text{ where } k = 1, \dots, n_i - 1$$
.

This category admits Auslander-Reiten triangles, having the Auslander-Reiten quiver shown in Figure 1. Moreover, it is classically generated by the residue field  $V_i^{(1)} = \emptyset(\Sigma)/\langle t_{x_i} \rangle$  of  $A_i$ . The triangulated extracting  $D_{i}$  (A)  $\Sigma$  model is 1. Calabi Yau for all i with involution shift


4. The triangulated category  $D_{Sing}(A_i) \simeq \underline{mod}_{A_i}$  is 1-Calabi-Yau for all *i*, with involutive shift functor:

$$\Omega(V_i^{(k)}) = V_i^{(n_i - k)} \quad \forall k = 1, \dots, n_i - 1$$

5. There exists an equivalences of triangulated categories:

$$\operatorname{hmf}(\emptyset(\Sigma), W) \simeq \vee_{i=1}^{N} \operatorname{D}_{\operatorname{Sing}}(A_{i}) \simeq \vee_{i=1}^{N} \underline{\operatorname{mod}}_{A_{i}}$$

6. The triangulated category  $hmf(\emptyset(\Sigma), W)$  is 1-Calabi-Yau, Krull-Schmidt and admits Auslander-Reiten triangles. Its Auslander-Reiten quiver is disconnected, with connected components given by the Auslander-Reiten quivers of the categories  $\underline{mod}_{A_i}$ .



**Figure 1.** Auslander-Reiten quiver for  $\underline{\text{mod}}_{A_i}$  when  $n_i = 5$ . The Auslander-Reiten translation fixes all vertices and the multiplicities of all arrows are trivial.

## 3.5. Counting indecomposables in $hmf(\mathcal{O}(\Sigma), W)$

The following result gives the count of simple topological D-branes of the LG model:

**Proposition 3.15.** [9] Suppose that W is critically finite of the form given in eq. (1). Then the number of isomorphism classes of indecomposable non-zero objects of the category  $\operatorname{hmf}(\emptyset(\Sigma), W)$  equals  $\sum_{i=1}^{N} (n_i - 1) = -N + \sum_{i=1}^{N} n_i$ .

The degrees  $n_i$  of the prime factors  $t_{x_i}$  arising in the decomposition of  $W_c$  define a  $\mathbb{Z}_2$ -grading on the set  $I_N \stackrel{\text{def.}}{=} \{1, \ldots, N\}$  whose components are given by:

$$I_N^{\hat{0}} \stackrel{\text{def.}}{=} \{i \in I_N \mid n_i \text{ is even}\}, \ I_N^{\hat{1}} \stackrel{\text{def.}}{=} \{i \in I_N \mid n_i \text{ is odd}\}.$$

Let:

$$N^{\hat{0}} \stackrel{\text{def.}}{=} |I_N^{\hat{0}}| \text{ and } N^{\hat{1}} \stackrel{\text{def.}}{=} |I_N^{\hat{1}}|$$

denote the cardinalities of these subsets of I, which satisfy  $N^{\hat{0}} + N^{\hat{1}} = N$ . Any non-empty subset  $K \subset I_N$  is endowed with the  $\mathbb{Z}_2$ -grading induced from  $I_N$ , which has components:

$$K^{\hat{0}} \stackrel{\text{def.}}{=} K \cap I_N^{\hat{0}} , \quad K^{\hat{1}} \stackrel{\text{def.}}{=} K \cap I_N^{\hat{1}}$$

3.6. Counting elementary factorizations in  $hmf(\mathcal{O}(\Sigma), W)$  and  $HMF(\mathcal{O}(\Sigma), W)$ 

The following result allows one to count elementary topological D-branes as well as elementary brane-antibrane pairs:

**Theorem 3.16.** [9] Suppose that W is a critically-finite element of  $\emptyset(\Sigma)$  with the decomposition given in eq. (1). Then:

1. The number of isomorphism classes of elementary factorizations in the category  $\operatorname{hmf}(\mathcal{O}(\Sigma), W)$  is given by:

$$\widehat{\mathcal{N}}_W = \sum_{k=0}^{N^1} \sum_{\substack{K \subseteq I_N, \\ |K^{\widehat{1}}| = k}} 2^{N^{\widehat{0}} + k} \prod_{i \in K} \left\lfloor \frac{n_i - 1}{2} \right\rfloor$$

2. The number of isomorphism classes of elementary matrix factorizations in the category  $\text{HMF}(\mathcal{O}(\Sigma), W)$  is given by:

$$\mathcal{N}_W = 2^{r^0} + \sum_{k=0}^{N^1} 2^{N^0 + k - 1} \sum_{\substack{K \subseteq I_N \\ |K^1| = k}} \prod_{i \in K} \left\lfloor \frac{n_i - 1}{2} \right\rfloor$$

#### Acknowledgments

This work was supported by the research grants IBS-R003-S1 ("Constructive string field theory of open-closed topological B-type strings") and IBS-R003-D1. The second author was also supported by the Max Planck Institute for Mathematics, Bonn and by the Australian Research Council grant DP180103891.

#### References

- [1] Lazaroiu, C. I., On the boundary coupling of topological Landau-Ginzburg models, JHEP 0505 (2005) 037.
- [2] Herbst, M., Lazaroiu, C. I., Localization and traces in open-closed topological Landau-Ginzburg models, *JHEP* 0505 (2005) 044.
- [3] Lazaroiu, C. I., On the structure of open-closed topological field theory in two dimensions, Nucl. Phys. B 603 (2001) 497–530.
- [4] Babalic, E. M., Doryn, D., Lazaroiu, C. I., Tavakol, M., Differential models for B-type open-closed topological Landau-Ginzburg theories, Commun. Math. Phys. 361 (2018) 1169–1234 (arXiv:1610.09103v3 [math.DG]).

- [5] Doryn, D., Lazaroiu, C. I., Non-degeneracy of cohomological traces for general Landau-Ginzburg models, preprint, arXiv:1802.06261 [math.AG].
- [6] Babalic, E. M., Doryn, D., Lazaroiu, C. I., Tavakol, M., A differential model for B-type Landau-Ginzburg theories, *preprint*, arXiv:1709.00684 [math.DG].
- [7] Babalic, E. M., Doryn, D., Lazaroiu, C. I., Tavakol, M., On B-type open-closed Landau-Ginzburg theories defined on Calabi-Yau Stein manifolds, *Commun. Math. Phys.* **362** (2018) 129–165 (arXiv:1610.09813v3 [math.DG]).
- [8] Babalic, E. M., Doryn, D., Lazaroiu, C. I., Tavakol, M., B-type Landau-Ginzburg models on Stein manifolds, preprint, conference proceedings for "Group 32", Prague, July 9-13, 2018.
- [9] Lazaroiu, C. I., Tavakol, M., B-type topological Landau-Ginzburg models over general non-compact Riemann surfaces, preprint available at https://cgp.ibs.re.kr/archive/preprints.
- [10] Doryn, D., Lazaroiu, C. I., Tavakol, M., Elementary matrix factorizations over Bézout domains, preprint, arXiv:1801.02369 [math.AC].
- [11] Doryn, D., Lazaroiu, C. I., Tavakol, M., Matrix factorizations over elementary divisor domains, preprint, arXiv:1802.07635 [math.AC].
- [12] Kerékjártó, B., Vorlesungen über Topologie I, Springer, Berlin, 1923.
- [13] Stoilow, S., Leçons sur les principes topologiques de la théorie de fonctions analytiques, 2nd ed., Gauthier-Villars, Paris, 1956.
- [14] Richards, I., On the Classification of Non-Compact Surfaces, Transactions of the AMS 106 (1963) 2, 259–269.
- [15] Lazaroiu, C. I., Generalized complexes and string field theory, JHEP 0106 (2001) 052.
- [16] Lazaroiu, C. I., Unitarity, D-brane dynamics and D-brane categories, JHEP 0112 (2001) 031.
- [17] Lazaroiu, C. I., D-brane categories, Int. J. Mod. Phys. A18 (2003) 5299–5335.
- [18] Herbst, M., Lazaroiu, C. I., Lerche, W., D-brane effective action and tachyon condensation in topological minimal models, JHEP 0503 (2005) 078.