A PROBABILISTIC TAKENS THEOREM

KRZYSZTOF BARAŃSKI¹, YONATAN GUTMAN², AND ADAM ŚPIEWAK¹

ABSTRACT. Let $X \subset \mathbb{R}^N$ be a Borel set, μ a Borel probability measure on X and $T: X \to X$ a locally Lipschitz and injective map. Fix $k \in \mathbb{N}$ strictly greater than the (Hausdorff) dimension of X and assume that the set of p-periodic points of T has dimension smaller than p for $p = 1, \ldots, k - 1$. We prove that for a typical polynomial perturbation \tilde{h} of a given locally Lipschitz function $h: X \to \mathbb{R}$, the k-delay coordinate map $x \mapsto (\tilde{h}(x), \tilde{h}(Tx), \ldots, \tilde{h}(T^{k-1}x))$ is injective on a set of full μ -measure. This is a probabilistic version of the Takens delay embedding theorem as proven by Sauer, Yorke and Casdagli. We also provide a nondynamical probabilistic embedding theorem of similar type, which strengthens a previous result by Alberti, Bölcskei, De Lellis, Koliander and Riegler. In both cases, the key improvements compared to the non-probabilistic counterparts are the reduction of the number of required coordinates from 2 dim X to dim X and using Hausdorff dimension instead of the box-counting one. We present examples showing how the use of the Hausdorff dimension improves the previously obtained results.

1. INTRODUCTION

Consider an experimentalist observing a physical system modeled by a discrete time dynamical system (X, T), where $T: X \to X$ is the evolution rule and the phase space X is a subset of the Euclidean space \mathbb{R}^N . It often happens that, for a given point $x \in X$, instead of an actual sequence of k states $x, Tx, \ldots, T^{k-1}x$, the observer's access is limited to the values of k measurements $h(x), h(Tx), \ldots, h(T^{k-1}x)$, for a real-valued observable $h: X \to \mathbb{R}$. Therefore, it is natural to ask, to what extent the original system can be reconstructed from such sequences of measurements and what is the minimal number k, referred to as the number of delay-coordinates, required for a reliable reconstruction. These questions have emerged in the physical literature (see e.g. [PCFS80]) and inspired a number of mathematical results, known as Takens-type delay embedding theorems, stating that the reconstruction of (X,T)is possible for certain observables h, as long as the measurements $h(x), h(Tx), \ldots, h(T^{k-1}x)$ are known for all $x \in X$ and large enough k.

¹Institute of Mathematics, University of Warsaw, ul. Banacha 2, 02-097 Warszawa, Poland

²Institute of Mathematics, Polish Academy of Sciences, ul. Sniadeckich 8, 00-656 Warszawa, Poland

E-mail addresses: baranski@mimuw.edu.pl, y.gutman@impan.pl, a.spiewak@mimuw.edu.pl.

²⁰⁰⁰ Mathematics Subject Classification. 37C45 (Dimension theory of dynamical systems), 28A78 (Hausdorff and packing measures), 28A80 (Fractals).

Key words and phrases. Takens delay embedding theorem, probabilistic embedding, Hausdorff dimension, box-counting dimension.

The possibility of performing measurements at every point of the phase space is clearly unrealistic. However, such an assumption enables one to obtain theoretical results which justify the validity of actual procedures used by experimentalists (see e.g. [HGLS05, KY90, SGM90, SM90]). Note that one cannot expect a reliable reconstruction of the system based on the measurements of a given observable h, as it may fail to distinguish the states of the system (e.g. if h is a constant function). It is therefore necessary (and rather realistic) to assume that the experimentalists are able to *perturb* the given observable. The first result obtained in this area is the celebrated Takens delay embedding theorem for smooth systems on manifolds [Tak81, Theorem 1]. Due to its strong connections with actual reconstruction procedures used in the natural sciences, Takens theorem has been met with great interest among mathematical physicists (see e.g. [HBS15, SYC91, Vos03]). Let us recall its extension due to Sauer, Yorke and Casdagli [SYC91]. In this setting, the number k of the delaycoordinates should be two times larger than the upper box-counting dimension of the phase space X (denoted by $\overline{\dim}_B X$; see Section 2 for the definition), and the perturbation is a polynomial of degree 2k. The formulation of the result given here follows [Rob11].

Theorem 1.1 ([Rob11, Theorem 14.5]). Let $X \subset \mathbb{R}^N$ be a compact set and let $T: X \to X$ be Lipschitz and injective. Let $k \in \mathbb{N}$ be such that $k > 2\overline{\dim}_B X$ and assume $2\overline{\dim}_B (\{x \in X : T^p x = x\}) < p$ for $p = 1, \ldots, k - 1$. Let $h: \mathbb{R}^N \to \mathbb{R}$ be a Lipschitz function and $h_1, \ldots, h_m: \mathbb{R}^N \to \mathbb{R}$ a basis of the space of polynomials of degree at most 2k. For $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$ denote by $h_\alpha: \mathbb{R}^N \to \mathbb{R}$ the map

$$h_{\alpha}(x) = h(x) + \sum_{j=1}^{m} \alpha_j h_j(x).$$

Then for Lebesgue almost every $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$, the transformation

$$\phi_{\alpha}^T \colon X \to \mathbb{R}^k, \qquad \phi_{\alpha}^T(x) = (h_{\alpha}(x), h_{\alpha}(Tx), \dots, h_{\alpha}(T^{k-1}x))$$

is injective on X.

The map ϕ_{α}^{T} is called the *delay-coordinate map*. Note that Theorem 1.1 applies to any compact set $X \subset \mathbb{R}^{N}$, not necessarily a manifold. This is a useful feature, as it allows one to consider sets with a complicated geometrical structure, such as fractal sets arising as attractors in chaotic dynamical systems, see e.g. [ER85]. Moreover, the upper box-counting dimension of X can be smaller than the dimension of any smooth manifold containing X, so Theorem 1.1 may require fewer delay-coordinates than its smooth counterpart in [Tak81].

As it was noted above, usually an experimentalist may perform only a finite number of observations $h(x_j), \ldots, h(T^{k-1}x_j)$ for some points $x_j \in X$, $j = 1, \ldots, l$. We believe it is realistic to assume there is an (explicit or implicit) random process determining which initial states x_j are accessible to the experimentalist. In this paper we are interested in the question of reconstruction of the system in presence of such process. Mathematically speaking, this corresponds to fixing a probability measure μ on X and asking whether the delay-coordinate map ϕ_{α}^T is injective almost surely with respect to μ . Since in this setting we are allowed to neglect sets of probability zero, it is reasonable to ask whether the minimal number of delay-coordinates sufficient for the reconstruction of the system can be smaller than 2 dim X.

Our main result states that this is indeed the case, and the number of delay-coordinates can be reduced by half for *any* (Borel) probability measure.

The problem of determining the minimal number of delay-coordinates required for reconstruction has been already considered in the physical literature. In [PCFS80], the authors analyzed an algorithm which may by interpreted as an attempt to determine this number in a probabilistic setting. Our work provides rigorous results in this direction. The following theorem is a simplified version of our result.

Theorem 1.2 (Probabilistic Takens delay embedding theorem). Let $X \subset \mathbb{R}^N$ be a Borel set, μ a Borel probability measure on X and $T: X \to X$ an injective, locally Lipschitz map. Take $k \in \mathbb{N}$ such that $k > \dim X$ and assume that for $p = 1, \ldots, k - 1$ we have $\dim(\{x \in X : T^p x = x\}) < p$ or $\mu(\{x \in X : T^p x = x\}) = 0$. Let $h: X \to \mathbb{R}$ be a locally Lipschitz function and $h_1, \ldots, h_m: \mathbb{R}^N \to \mathbb{R}$ a basis of the space of real polynomials of Nvariables of degree at most 2k - 1. For $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$ denote by $h_\alpha: \mathbb{R}^N \to \mathbb{R}$ the map

$$h_{\alpha}(x) = h(x) + \sum_{j=1}^{m} \alpha_j h_j(x).$$

Then for Lebesgue almost every $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$, there exists a Borel set $X_\alpha \subset X$ of full μ -measure, such that the delay-coordinate map

$$\phi_{\alpha}^T \colon X \to \mathbb{R}^k, \qquad \phi_{\alpha}^T(x) = (h_{\alpha}(x), h_{\alpha}(Tx), \dots, h_{\alpha}(T^{k-1}x))$$

is injective on X_{α} .

In the above theorem, the dimension dim can be chosen to be any of $\dim_H, \underline{\dim}_B, \overline{\dim}_B$ (Hausdorff, lower and upper box-counting dimension; for the definitions see Section 2). Recall that for any Borel set X one has

(1.1)
$$\dim_H X \le \underline{\dim}_B X \le \overline{\dim}_B X$$

(see e.g. [Fal14, Proposition 3.4]). Since the inequalities in (1.1) may be strict, using the Hausdorff dimension instead of the box-counting one(s) may reduce the required number of delay-coordinates. In particular, there are compact sets $X \subset \mathbb{R}^N$ with $\dim_H X = 0$ and $\overline{\dim}_B X = N$, hence Theorem 1.2 can reduce significantly the number of required delay-coordinates compared to Theorem 1.1 (in a probabilistic setting).

Notice that in Theorem 1.2 we do not assume that the measure μ is *T*-invariant. However, the invariance of μ provides some additional benefits, as shown in the following remark.

Remark 1.3 (Invariant measure case). Suppose that the measure μ in Theorem 1.2 is additionally *T*-invariant, i.e. $\mu(Y) = \mu(T^{-1}(Y))$ for every Borel set $Y \subset X$. Then the set X_{α} can be chosen to satisfy $T(X_{\alpha}) = X_{\alpha}$. Moreover, if μ is *T*-invariant and ergodic (i.e. $T^{-1}(Y) = Y$ can occur only for sets Y of 0 or full μ -measure), then the assumption on the periodic points of T in Theorem 1.2 can be omitted.

Note that in the case when the measure μ is *T*-invariant, Theorem 1.2 and Remark 1.3 show that for a suitable choice of X_{α} , the map ϕ_{α}^{T} is injective on the invariant set X_{α} , which

implies that the dynamical system (\hat{X}, \hat{T}) for $\hat{X} = \phi_{\alpha}^{T}(X_{\alpha}), \hat{T} = \phi_{\alpha}^{T} \circ T \circ (\phi_{\alpha}^{T})^{-1}$, is a model of the system (X, T) embedded in \mathbb{R}^{k} .

An extended versions of Theorem 1.2 and Remark 1.3 are presented and proved in Section 4 as Theorem 4.3 and Remark 4.4, respectively. Theorem 4.3 shows that the assumption $k > \dim X$ can be slightly weakened, and in addition to locally Lipschitz functions h, one can consider locally β -Hölder functions for suitable $\beta \in (0, 1]$. Moreover, one can replace the probabilistic measure μ by any Borel σ -finite measure on X. For details, see Section 4.

Notice that to eliminate the assumption on the periodic points of T in Theorem 1.2, one can also consider systems with 'few' or no periodic points. For instance, as proved in [Yor69], a flow on a subset of Euclidean space given by an autonomous differential equation $\dot{x} = F(x)$, where F is Lipschitz with a constant L, has no periodic orbits of period smaller than $\frac{2\pi}{L}$. It follows that if T is a t-time map for such a flow with $t < \frac{2\pi}{L \dim X}$, then it has no periodic points of periods smaller than $\dim X$ and therefore the assumption on periodic points in Theorem 1.2 can be omitted (compare also [Gut16, Remark 1.2]). The same holds if the number of periodic points of a given period is finite, which by the Kupka–Smale theorem is a generic condition in the space of C^r -diffeomorphisms ($r \ge 1$) of a compact manifold equipped with the uniform C^r -topology¹.

As has been mentioned already, Takens theorems are used in order to justify actual (approximate) delay map procedures based on real experimental data (see e.g. [MGNS18, HGLS05, SGM90, SM90]). Note, however, that in the cited papers the dimension of the phase space X is deduced a *posteriori* from the properties of the *time series* (orbits of the delay coordinate map for a given observable). It would be very interesting to know whether in the literature it has been observed for some experimental data originating from a space X with known dimension that it is sufficient to have $k \approx \dim X$ (instead of $k \approx 2 \dim X$) delay-coordinates (in other words, time series of length k) in the framework of such procedures.

In this paper we focus our attention to the case when the space X is a subset of a finitedimensional Euclidean space. Takens-type delay embedding theorems have also been extended to finite-dimensional subsets of Banach spaces (see e.g. [Rob05]). It is a natural question, whether our probabilistic embedding theorems can also be transferred into the infinite-dimensional setup. This problem will be considered in a subsequent work.

Takens-type delay embedding theorems can be seen as dynamical versions of *embedding* theorems which specify when a finite-dimensional set can be embedded into a Euclidean space. Indeed, under the assumptions of Theorem 1.1, the delay-coordinate map ϕ_{α}^{T} is an embedding of X into \mathbb{R}^{k} for typical α . Embedding theorems in various categories have been extensively studied in a number of papers (see Section 3 for a more detailed discussion). Recently, Alberti, Bölcskei, De Lellis, Koliander and Riegler [ABD+19] proved a probabilistic embedding theorem involving the modified lower box-counting dimension of the set (see Theorem 3.6). We are able to improve this result by considering the Hausdorff dimension.

¹According to the Kupka–Smale theorem ([PdM82, Chapter 3, Theorem 3.6]) it is generic that the periodic points are hyperbolic and thus periodic points of a given period are isolated by the Hartman–Grobman theorem ([PdM82, Chapter 2, Theorem 4.1]).

Below we present a simplified version of our theorem, which can be seen as a non-dynamical counterpart of Theorem 1.2.

Theorem 1.4 (Probabilistic embedding theorem). Let $X \subset \mathbb{R}^N$ be a Borel set and let μ be a Borel probability measure on X. Take $k \in \mathbb{N}$ such that the k-th Hausdorff measure of X is zero (it suffices to take $k > \dim_H X$) and let $\phi: X \to \mathbb{R}^k$ be a locally Lipschitz map. Then for Lebesgue almost every linear transformation $L: \mathbb{R}^N \to \mathbb{R}^k$ there exists a Borel set $X_L \subset X$ of full μ -measure, such that $\phi_L = \phi + L$ is injective on X_L .

The extended version of the theorem is formulated and proved in Section 3 as Theorem 3.1. In particular, we obtain the following geometric corollary (see Section 3 for details).

Corollary 1.5 (Probabilistic injective projection theorem). Let $X \subset \mathbb{R}^N$ be a Borel set and let μ be a Borel probability measure on X. Then for every $k > \dim_H X$ and almost every k-dimensional linear subspace $S \subset \mathbb{R}^N$, the orthogonal projection of X into S is injective on a full μ -measure subset of X.

Notice that by the Marstrand–Mattila projection theorem (see [Mar54, Mat75]), if $X \subset \mathbb{R}^N$ is Borel and $k \geq \dim_H X$, then for almost all k-dimensional linear subspaces $S \subset \mathbb{R}^N$, the image of X under the orthogonal projection into S has Hausdorff dimension equal to $\dim_H X$. Note also that Sauer and Yorke proved in [SY97] that the dimension² of a bounded Borel subset X of \mathbb{R}^N is preserved under typical smooth maps and typical delay-coordinate maps into \mathbb{R}^k as long as $k \geq \dim X$.

In this paper we also provide several examples. Example 3.5 shows that in general the condition $k > \dim_H X$ in Theorem 1.4 cannot be replaced by $k \ge \dim_H X$. Example 4.6 shows that linear perturbations of the observable are not sufficient for Takens theorem. Section 5 contains a pair of examples. The first one is based on Kan's example from the Appendix to [SYC91], showing that condition $k > 2 \dim_H X$ is not sufficient for existence of a linear transformation into \mathbb{R}^k which is injective on X. As in the probabilistic setting one can work with the Hausdorff dimension, we consider a set $X \subset \mathbb{R}^2$ similar to the one provided by Kan, which cannot be embedded linearly into \mathbb{R} , but when endowed with a natural probability measure, almost every linear transformation $L: \mathbb{R}^2 \to \mathbb{R}$ is injective on a set of full measure. The second example provides a probability measure with $\dim_H \mu < \underline{\dim}_{MB} \mu$, showing that Theorem 1.4 strengthens a previous result from [ABD+19].

Organization of the paper. The paper is organized as follows. In Section 2 we introduce notation, definitions and preliminary results. Section 3 contains the formulation and proof of the extended version of the probabilistic embedding theorem (Theorem 3.1), while Section 4 is devoted to the proof of the extended version of the probabilistic Takens delay embedding theorem (Theorem 4.3). In Section 5 we present examples showing how the use of the Hausdorff dimension improves the previously obtained results.

Acknowledgements. We are grateful to Erwin Riegler for helpful discussions and to the anonymous referees for helpful comments. Y. G. and A. Ś. were partially supported by the National Science Centre (Poland) grant 2016/22/E/ST1/00448.

 $^{^2\}mathrm{Any}$ one of the dimensions mentioned above and denoted by dim.

2. Preliminaries

Consider the Euclidean space \mathbb{R}^N for $N \in \mathbb{N}$, with the standard inner product $\langle \cdot, \cdot \rangle$ and the norm $\|\cdot\|$. The open δ -ball around a point $x \in \mathbb{R}^N$ is denoted by $B_N(x, \delta)$. By |X| we denote the diameter of a set $X \subset \mathbb{R}^N$. We say that function $\phi: X \to \mathbb{R}^k$, $X \subset \mathbb{R}^N$ is *locally* β -Hölder for $\beta > 0$ if for every $x \in X$ there exists an open set $U \subset \mathbb{R}^N$ containing x such that ϕ is β -Hölder on $U \cap X$, i.e. there exists C > 0 such that

$$\|\phi(x) - \phi(y)\| \le C \|x - y\|^{\beta}$$

for every $x, y \in U \cap X$. We say that ϕ is *locally Lipschitz* if it is locally 1-Hölder.

For $k \leq N$ we write $\operatorname{Gr}(k, N)$ for the (k, N)-Grassmannian, i.e. the space of all kdimensional linear subspaces of \mathbb{R}^N , equipped with the standard rotation-invariant (Haar) measure, see [Mat95, Section 3.9] (and [FR02] for another construction of a rotation-invariant measure on the Grassmannian). By η_N we denote the normalized Lebesgue measure on the unit ball $B_N(0, 1)$, i.e.

$$\eta_N = \frac{1}{\operatorname{Leb}(B_N(0,1))} \operatorname{Leb}|_{B_N(0,1)},$$

where Leb is the Lebesgue measure on \mathbb{R}^N .

For s > 0, the s-dimensional (outer) Hausdorff measure of a set $X \subset \mathbb{R}^N$ is defined as

$$\mathcal{H}^{s}(X) = \lim_{\delta \to 0} \inf \Big\{ \sum_{i=1}^{\infty} |U_i|^s : X \subset \bigcup_{i=1}^{\infty} U_i, \ |U_i| \le \delta \Big\}.$$

The Hausdorff dimension of X is given as

$$\dim_H X = \inf\{s > 0 : \mathcal{H}^s(X) = 0\} = \sup\{s > 0 : \mathcal{H}^s(X) = \infty\}.$$

For a bounded set $X \subset \mathbb{R}^N$ and $\delta > 0$, let $N(X, \delta)$ denote the minimal number of balls of diameter at most δ required to cover X. The *lower* and *upper box-counting* (*Minkowski*) *dimensions* of X are defined as

$$\underline{\dim}_B X = \liminf_{\delta \to 0} \frac{\log N(X, \delta)}{-\log \delta} \quad \text{and} \quad \overline{\dim}_B X = \limsup_{\delta \to 0} \frac{\log N(X, \delta)}{-\log \delta}$$

The lower (resp. upper) box-counting dimension of an unbounded set is defined as the supremum of the lower (resp. upper) box-counting dimensions of its bounded subsets.

The lower and upper modified box-counting dimensions of $X \subset \mathbb{R}^N$ are defined as

$$\underline{\dim}_{MB} X = \inf \Big\{ \sup_{i \in \mathbb{N}} \underline{\dim}_B K_i : X \subset \bigcup_{i=1}^{\infty} K_i, \ K_i \text{ compact} \Big\},\$$
$$\overline{\dim}_{MB} X = \inf \Big\{ \sup_{i \in \mathbb{N}} \overline{\dim}_B K_i : X \subset \bigcup_{i=1}^{\infty} K_i, \ K_i \text{ compact} \Big\}.$$

With this notation, the following inequalities hold:

(2.1)
$$\dim_{H} X \leq \underline{\dim}_{MB} X \leq \overline{\dim}_{MB} X \leq \overline{\dim}_{B} X, \\ \dim_{H} X \leq \underline{\dim}_{MB} X \leq \underline{\dim}_{B} X \leq \overline{\dim}_{B} X.$$

We define dimension of a finite Borel measure μ in \mathbb{R}^N as

 $\dim \mu = \inf \{\dim X : X \subset \mathbb{R}^N \text{ is a Borel set of full } \mu\text{-measure} \}.$

Here dim may denote any one of the dimensions defined above. Recall that for a measure μ on a set X and a measurable $Y \subset X$ we say that Y is of *full* μ -measure, if $\mu(X \setminus Y) = 0$. For more information on dimension theory in Euclidean space see [Fal14, Mat95, Rob11].

For $N, k \in \mathbb{N}$ let $\operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ be the space of all linear transformations $L: \mathbb{R}^N \to \mathbb{R}^k$. Such transformations are given by

(2.2)
$$Lx = (\langle l_1, x \rangle, \dots, \langle l_k, x \rangle),$$

where $l_1, \ldots, l_k \in \mathbb{R}^N$. Thus, the space $\operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ can be identified with $(\mathbb{R}^N)^k$, and the Lebesgue measure on $\operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ is understood as $\bigotimes_{j=1}^k$ Leb, where Leb is the Lebesgue measure in \mathbb{R}^N . Within the space $\operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ we consider the space E_k^N consisting of all linear transformations $L: \mathbb{R}^N \to \mathbb{R}^k$ of the form (2.2), for which $l_1, \ldots, l_k \in B_N(0, 1)$. Note that by the Cauchy-Schwarz inequality,

$$\|Lx\| \le \sqrt{N} \|x\|$$

for every $L \in E_k^N$ and $x \in \mathbb{R}^N$.

By $\eta_{N,k}$ we denote the normalized Lebesgue measure on E_k^N , i.e. the probability measure on E_k^N given by

$$\eta_{N,k} = \bigotimes_{j=1}^{k} \frac{1}{\text{Leb}(B_N(0,1))} \text{Leb}|_{B_N(0,1)}.$$

The following geometrical inequality, used in [HK99] (see also [Rob11, Lemma 4.1]) is the key ingredient of the proof of Theorem 3.1.

Lemma 2.1. Let $L: \mathbb{R}^N \to \mathbb{R}^k$ be a linear transformation. Then for every $x \in \mathbb{R}^N \setminus \{0\}$, $z \in \mathbb{R}^k$ and $\varepsilon > 0$,

$$\eta_{N,k}(\{L \in E_k^N : \|Lx + z\| \le \varepsilon\}) \le CN^{k/2} \frac{\varepsilon^k}{\|x\|^k}$$

where C > 0 is an absolute constant.

For $L \in \text{Lin}(\mathbb{R}^m; \mathbb{R}^k)$, where $m, k \in \mathbb{N}$, denote by $\sigma_p(L)$, $p \in \{1, \ldots, k\}$, the *p*-th largest singular value of the matrix L, i.e. the *p*-th largest square root of an eigenvalue of the matrix L^*L . In the proof of Theorem 4.3, instead of Lemma 2.1 we will use the following lemma, proved as [SYC91, Lemma 4.2] (see also [Rob11, Lemma 14.3]).

Lemma 2.2. Let $L: \mathbb{R}^m \to \mathbb{R}^k$ be a linear transformation. Assume that $\sigma_p(L) > 0$ for some $p \in \{1, \ldots, k\}$. Then for every $z \in \mathbb{R}^k$ and $\rho, \varepsilon > 0$,

$$\frac{\operatorname{Leb}(\{\alpha \in B_m(0,\rho) : \|L\alpha + z\| \le \varepsilon\})}{\operatorname{Leb}(B_m(0,\rho))} \le C_{m,k} \left(\frac{\varepsilon}{\sigma_p(L)\rho}\right)^p,$$

where $C_{m,k} > 0$ is a constant depending only on m, k and Leb is the Lebesgue measure on \mathbb{R}^m .

To verify the measurability of the sets occurring in subsequent proofs, we will use the two following elementary lemmas. A measure μ on a set X is called σ -finite if there exists a countable collection of measurable sets A_n , $n \in \mathbb{N}$ such that $\mu(A_n) < \infty$ for each $n \in \mathbb{N}$ and $\bigcup_{n=1}^{\infty} A_n = X$. Recall that a σ -compact set is a countable union of compact sets.

Lemma 2.3. Let $X \subset \mathbb{R}^N$ be a Borel set and let μ be a Borel σ -finite measure on X. Then there exists a σ -compact set $K \subset X$ of full μ -measure.

Proof. Follows directly from the fact that a σ -finite Borel measure in a Euclidean space is regular (see e.g. [Bil99, Theorem 1.1]).

Lemma 2.4. Let \mathcal{X}, \mathcal{Z} be metric spaces. Then the following hold.

- (a) If $K \subset \mathcal{X} \times \mathcal{Z}$ is σ -compact, then so is $\pi_{\mathcal{X}}(K)$, where $\pi_{\mathcal{X}} \colon \mathcal{X} \times \mathcal{Z} \to \mathcal{X}$ is the projection given by $\pi_{\mathcal{X}}(x, z) = x$. In particular, $\pi_{\mathcal{X}}(K)$ is Borel.
- (b) If \mathcal{X} is σ -compact, $F \colon \mathcal{X} \to \mathcal{Z}$ is continuous and $K \subset \mathcal{Z}$ is σ -compact, then $F^{-1}(K)$ is σ -compact, hence Borel.
- (c) If \mathcal{X}, \mathcal{Z} are σ -compact, $F: \mathcal{X} \times \mathcal{Z} \to \mathbb{R}^k$, $k \in \mathbb{N}$, is continuous and $K \subset \mathcal{X}$ is σ -compact, then the set

$$\{(x,z) \in \mathcal{X} \times \mathcal{Z} : F(x,z) = F(y,z) \text{ for some } y \in K \setminus \{x\}\}$$

is σ -compact and hence Borel.

Proof. The statement (a) follows from the fact that $\pi_{\mathcal{X}}$ is continuous, and a continuous image of a compact set is also compact. To show (b), it is enough to notice that $F^{-1}(K)$ is a countable union of closed subsets of a σ -compact space. To check (c), let $\pi_{\mathcal{X}\times\mathcal{Z}} \colon \mathcal{X}\times K\times\mathcal{Z} \to \mathcal{X}\times\mathcal{Z}$ be the projection $\pi_{\mathcal{X}\times\mathcal{Z}}(x, y, z) = (x, z)$. Then

$$\begin{aligned} &\{(x,z) \in \mathcal{X} \times \mathcal{Z} : F(x,z) = F(y,z) \text{ for some } y \in K \setminus \{x\}\} \\ &= \pi_{\mathcal{X} \times \mathcal{Z}} \big(\{(x,y,z) \in \mathcal{X} \times K \times \mathcal{Z} : F(x,z) = F(y,z), \ d(x,y) \neq 0\}\big) \\ &= \bigcup_{n=1}^{\infty} \pi_{\mathcal{X} \times \mathcal{Z}} \big(\{(x,y,z) \in \mathcal{X} \times K \times \mathcal{Z} : F(x,z) = F(y,z), \ d(x,y) \geq \frac{1}{n}\}\big), \end{aligned}$$

where d is the metric in \mathcal{X} . Since d is continuous, we can use (a) and (b) to end the proof. \Box

3. PROBABILISTIC EMBEDDING THEOREM

In this section we prove an extended version of the probabilistic embedding theorem, formulated below. Obviously, Theorem 1.4 follows from Theorem 3.1

Theorem 3.1 (Probabilistic embedding theorem – extended version). Let $X \subset \mathbb{R}^N$ be a Borel set and μ be a Borel σ -finite measure on X. Take $k \in \mathbb{N}$ and $\beta \in (0, 1]$ such that $\mathcal{H}^{\beta k}(X) = 0$ and let $\phi: X \to \mathbb{R}^k$ be a locally β -Hölder map. Then for Lebesgue almost every linear transformation $L: \mathbb{R}^N \to \mathbb{R}^k$ there exists a Borel set $X_L \subset X$ of full μ -measure, such that the map $\phi_L = \phi + L$ is injective on X_L .

Remark 3.2. It is straightforward to notice that if $\dim_H X = 0$, then ϕ can be taken to be an arbitrary Hölder map.

Proof of Theorem 3.1. Note first that it sufficient to prove that the set X_L exists for $\eta_{N,k}$ almost every $L \in E_k^N$. Indeed, if this is shown, then for a given locally β -Hölder map $\phi: X \to \mathbb{R}^k$ we can take sets $\mathcal{L}_j \subset E_k^N$, $j \in \mathbb{N}$, such that $\eta_{N,k}(\mathcal{L}_j) = 1$ and for every $\tilde{L} \in \mathcal{L}_j$ the map $(\phi/j)_{\tilde{L}} = \phi/j + \tilde{L}$ is injective on a Borel set $X_{\tilde{L}}^{(j)} \subset X$ of full μ -measure. Then the set $\mathcal{L} = \bigcup_{j \in \mathbb{N}} \{j\tilde{L} : \tilde{L} \in \mathcal{L}_j\} \subset \operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ has full Lebesgue measure and for every $L \in \mathcal{L}$ there exists j such that $L/j \in \mathcal{L}_j$, so $(\phi/j)_{L/j} = (\phi + L)/j$ (and hence ϕ_L) is injective on $X_L = \bigcap_{j \in \mathbb{N}} X_{L/j}^{(j)}$, which has full μ -measure.

By Lemma 2.3, we can assume that X is σ -compact. Take $k \in \mathbb{N}, \beta \in (0, 1]$ with $\mathcal{H}^{\beta k}(X) = 0$ and a locally β -Hölder map $\phi \colon X \to \mathbb{R}^k$. Set

$$A = \{ (x, L) \in X \times E_k^N : \phi_L(x) = \phi_L(y) \text{ for some } y \in X \setminus \{x\} \}.$$

By Lemma 2.4, A is Borel. For $x \in X$ and $L \in E_k^N$, denote by A_x and A^L , respectively, the sections

$$A_x = \{ L \in E_k^N : (x, L) \in A \}, \quad A^L = \{ x \in X : (x, L) \in A \}.$$

The sets A_x and A^L are Borel as sections of a Borel set. Observe first that in order to prove the theorem it is enough to show $\eta_{N,k}(A_x) = 0$ for every $x \in X$, since then by Fubini's theorem ([Rud87, Thm. 8.8]), $(\eta_{N,k} \otimes \mu)(A) = 0$ and, consequently, $\mu(A^L) = 0$ for $\eta_{N,k}$ -almost every $L \in E_k^N$. Since ϕ_L is injective on $X \setminus A^L$, the assertion of the theorem is true.

Take a point $x \in X$. Since ϕ is locally β -Hölder and X is separable, there exists a countable covering of X by open sets $U_j \subset \mathbb{R}^N$, $j \in \mathbb{N}$, such that

(3.1)
$$\|\phi(y) - \phi(y')\| \le C_j \|y - y'\|^{\beta} \quad \text{for every} \quad y, y' \in U_j \cap X$$

for some $C_i > 0$. Let

$$K_n = \left\{ y \in X : \frac{1}{n} \le ||x - y|| \right\}.$$

To show $\eta_{N,k}(A_x) = 0$, it suffices to prove $\eta_{N,k}(A_{x,j,n}) = 0$ for every $j, n \in \mathbb{N}$, where

$$A_{x,j,n} = \{ L \in E_k^N : \phi_L(x) = \phi_L(y) \text{ for some } y \in U_j \cap K_n \}$$

Note that by Lemma 2.4, the set $A_{x,j,n}$ is Borel.

Take $j, n \in \mathbb{N}$ and fix a small $\varepsilon > 0$. Since $\mathcal{H}^{\beta k}(U_j \cap K_n) \leq \mathcal{H}^{\beta k}(X) = 0$, there exists a collection of balls $B_N(y_i, \varepsilon_i), i \in \mathbb{N}$, for some $y_i \in U_j \cap K_n$ and $\varepsilon_i > 0$, such that

(3.2)
$$U_j \cap K_n \subset \bigcup_{i \in \mathbb{N}} B_N(y_i, \varepsilon_i) \text{ and } \sum_{i=1}^{\infty} \varepsilon_i^{\beta k} \leq \varepsilon$$

Take $L \in A_{x,j,n}$ and $y \in U_j \cap K_n$ such that $\phi_L(x) = \phi_L(y)$. Then $y \in B_N(y_i, \varepsilon_i)$ for some $i \in \mathbb{N}$ and

$$\begin{aligned} \|L(y_{i} - x) + \phi(y_{i}) - \phi(x)\| &= \|\phi_{L}(y_{i}) - \phi_{L}(x)\| \\ &= \|\phi_{L}(y_{i}) - \phi_{L}(y)\| \\ &\leq \|\phi(y_{i}) - \phi(y)\| + \|L(y_{i} - y)\| \\ &\leq C_{j}\|y_{i} - y\|^{\beta} + \sqrt{N}\|y_{i} - y\| \\ &\leq M_{j}\varepsilon_{i}^{\beta} \end{aligned}$$

for some $M_j > 0$, by (2.3) and (3.1). This shows that

$$A_{x,j,n} \subset \bigcup_{i \in \mathbb{N}} \{ L \in E_k^N : \|L(y_i - x) + \phi(y_i) - \phi(x)\| \le M_j \varepsilon_i^\beta \}.$$

By Lemma 2.1, (3.2) and the fact $y_i \in K_n$, we have

$$\eta_{N,k}(A_{x,j,n}) \leq \sum_{i=1}^{\infty} \eta_{N,k}(\{L \in E_k^N : \|L(y_i - x) + \phi(y_i) - \phi(x)\| \leq M_j \varepsilon_i^\beta\})$$
$$\leq \frac{CN^{k/2}M_j^k}{1/n^k} \sum_{i=1}^{\infty} \varepsilon_i^{\beta k} \leq CN^{k/2}M_j^k n^k \varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, we obtain $\eta_{N,k}(A_{x,j,n}) = 0$, which ends the proof.

Remark 3.3. Note that the assumption $\mathcal{H}^{\beta k}(X) = 0$ is fulfilled if $\dim_H X < \beta k$, so Theorem 3.1 is indeed a Hausdorff dimension embedding theorem. Moreover, it may happen that $\mathcal{H}^{\beta k}(X) = 0$ and $\dim_H X = \beta k$.

As a simple consequence of Theorem 3.1, we obtain the following corollary, formulated in a slightly simplified version in Section 1 as Corollary 1.5.

Corollary 3.4 (Probabilistic injective projection theorem – extended version). Let $X \subset \mathbb{R}^N$ be a Borel set and let μ be a Borel σ -finite measure on X. Then for every $k \in \mathbb{N}, k \leq N$ such that $\mathcal{H}^k(X) = 0$ and almost every k-dimensional linear subspace $S \subset \mathbb{R}^N$ (with respect to the standard measure on the Grassmannian Gr(k, N)), the orthogonal projection of X into S is injective on a full μ -measure subset of X (depending on S).

Proof of Corollary 3.4. Apply Theorem 3.1 for the map $\phi \equiv 0$. Then we know that a linear map $L \in \operatorname{Lin}(\mathbb{R}^N; \mathbb{R}^k)$ of the form (2.2) is injective on a set $X_L \subset X$ of full μ -measure for Lebesgue almost every $(l_1, \ldots, l_k) \in (\mathbb{R}^N)^k$. We can assume that l_1, \ldots, l_k are linearly independent for all such L, which also implies that the same holds for Ll_1, \ldots, Ll_k . Setting

 $S_L = \operatorname{Span}(l_1, \ldots, l_k)$

and taking $V_L \in \text{Lin}(\mathbb{R}^k; \mathbb{R}^N)$ defined by $V_L(Ll_j) = l_j$ for $j = 1, \ldots, k$, we have

$$V_L \circ L = \prod_{S_L}$$

where Π_{S_L} is the orthogonal projection from \mathbb{R}^N onto S_L and V_L is injective. It follows that Π_{S_L} is injective on X_L for almost every (l_1, \ldots, l_k) , so Π_S is injective on a full μ -measure subset of X for almost every k-dimensional linear subspace $S \subset \mathbb{R}^N$.

Let us note that in general, the requirement $\mathcal{H}^{\beta k}(X) = 0$ in Theorem 3.1 cannot be replaced by the weaker condition $\dim_H(X) \leq \beta k$.

Example 3.5. Let $k = \beta = 1$, $X = \mathbb{S}^1 \subset \mathbb{R}^2$ be the unit circle and let μ be the normalized Lebesgue measure on \mathbb{S}^1 . We shall prove that there is no Lipschitz transformation $\phi \colon \mathbb{S}^1 \to \mathbb{R}$ which is injective on a set of full μ -measure. Let ϕ be such a transformation. Then $\phi(\mathbb{S}^1) = [a, b]$ for some compact interval. As ϕ is injective on a set of full measure, the interval [a, b] is non-degenerate, i.e. a < b. Fix points $x, y \in \mathbb{S}^1$ with $\phi(x) = a, \phi(y) = b$. As $x \neq y$, there are

exactly two open arcs $I, J \subset \mathbb{S}^1$ of positive measure joining x and y such that $\overline{I} \cap \overline{J} = \{x, y\}$ and $\overline{I} \cup \overline{J} = \mathbb{S}^1$. Clearly $\phi(\overline{I}) = \phi(\overline{J}) = [a, b]$. Let $A \subset \mathbb{S}^1$ be a Borel set such that ϕ is injective on A and $\mu(A)=1$. As Lipschitz maps transform sets of zero Lebesgue measure to sets of zero Lebesgue measure, we conclude that $\phi(I \cap A)$ and $\phi(J \cap A)$ are disjoint Lebesgue measurable subsets of [a, b] with Lebesgue measure equal to b - a. This contradiction shows that no Lipschitz transformation $\phi \colon \mathbb{S}^1 \to \mathbb{R}$ is injective on a full measure set.

Theorem 3.1 strengthens the following embedding theorem, proved recently by Alberti, Bölcskei, De Lellis, Koliander and Riegler in [ABD⁺19].

Theorem 3.6 ([ABD⁺19, Theorem II.1]). Let μ be a Borel probability measure in \mathbb{R}^N and let $k \in \mathbb{N}$ be such that $k > \underline{\dim}_{MB} \mu$. Then for Lebesgue almost every linear transformation $L: \mathbb{R}^N \to \mathbb{R}^k$ there exists a Borel set $X_L \subset \mathbb{R}^N$ such that $\mu(X_L) = 1$ and L is injective on X_L .

In fact, in [ABD⁺19] the authors introduced the notion of $\underline{\dim}_{MB}\mu$, denoting it by $K(\mu)$ and calling it the *description complexity* of the measure. In particular, Theorem 3.6 holds for measures μ supported on a Borel set $X \subset \mathbb{R}^N$ with $\underline{\dim}_B X < k$. By (2.1), we have $\underline{\dim}_H \mu \leq \underline{\dim}_{MB}\mu$, and in Section 5 we present an example (Theorem 5.5) showing that the inequality may be strict. Therefore, Theorem 3.1 actually strengthens Theorem 3.6.

Non-probabilistic embedding theorems were first obtained in topological and smooth categories. The well-known Menger-Nöbeling embedding theorem (see e.g. [HW41, Theorem V.2]) states that for a compact metric space X with Lebesgue covering dimension at most k, a generic continuous transformation $\phi: X \to \mathbb{R}^{2k+1}$ is injective (and hence defines a homeomorphism between X and $\phi(X)$). Genericity means here that the set of injective transformations $\phi: X \to \mathbb{R}^{2k+1}$ is a dense G_{δ} subset of $C(X; \mathbb{R}^{2k+1})$ endowed with the supremum metric. The dimension 2k + 1 is known to be optimal. The corresponding result in the category of smooth manifolds is the Whitney embedding theorem (see [Whi36]). It states that for a given k-dimensional C^r -manifold M, a generic C^r -transformation from M to \mathbb{R}^{2k+1} is a C^r -embedding (i.e. an injective immersion of class C^r).

Let us now compare Theorem 3.1 to non-probabilistic embedding theorems involving the box-counting dimension. One of the first results in this area was a theorem by Mañé [Mn81, Lemma 1.1]. We present its formulation following [SYC91, Theorem 4.6] and [Rob11, Theorem 6.2] (originally, Mañé proved that topologically generic linear transformation is injective on X).

Theorem 3.7. Let $X \subset \mathbb{R}^N$ be a compact set. Let $k \in N$ be such that $k > 2\overline{\dim}_B X$ (it suffices to take $k > \dim_H(X - X)$). Then Lebesgue almost every linear transformation $L: \mathbb{R}^N \to \mathbb{R}^k$ is injective on X.

Remark 3.8. As noticed by Mañé and communicated in [ER85, p. 627], his original statement in [Mn81] is incorrect. Namely, he assumed $k > 2 \dim_H X + 1$ instead of $k > \dim_H (X - X)$. However, this is known to be insufficient for the existence of a linear embedding of X into \mathbb{R}^k . In fact, in [SYC91, Appendix A], Kan presented an example of a set $X \subset \mathbb{R}^m$ with $\dim_H X = 0$, such that any linear transformation $L: \mathbb{R}^m \to \mathbb{R}^{m-1}$ fails to be injective on X. It turns out that the assumption $k > 2 \dim_H X$ is insufficient, while $k > 2 \dim_B X$ is sufficient. This stems from the fact that the proof of Theorem 3.7 actually requires the property $k > \dim_H (X - X)$, and the upper box-counting dimension satisfies

(3.3)
$$\overline{\dim}_B (A \times B) \le \overline{\dim}_B (A) + \overline{\dim}_B (B),$$

for $A, B \subset \mathbb{R}^N$, hence

$$\dim_{H}(X - X) \le \dim_{H}(X \times X) \le \overline{\dim}_{B}(X \times X) \le 2\overline{\dim}_{B}X$$

(note that this calculation shows that $k > 2\overline{\dim}_B X$ is a stronger assumption than $k > \dim_H(X-X)$). On the other hand, (3.3) does not hold for the Hausdorff dimension (nor for the lower box-counting dimension), and $\dim_H X$ does not control $\dim_H(X-X)$. The fact that in Theorem 3.1 we can work with the Hausdorff dimension comes from the application of Fubini's theorem, which enables us to consider covers of the set X itself, instead of X-X. In Section 5 we analyze Kan's example from the point of view of Theorem 3.1.

Theorem 3.7 is also true for subsets of an arbitrary Banach space \mathfrak{B} for a prevalent set of linear transformations $L: \mathfrak{B} \to \mathbb{R}^k$ (see [Rob11, Chapter 6] for details).

Note that the linear embedding from Theorem 3.1 need not preserve the dimension of X. Indeed, the Hausdorff and box-counting dimensions are invariants for bi-Lipschitz transformations, yet inverse of a linear map on a compact set does not have to be Lipschitz. Therefore, we only know that dim $\phi_L(X) \leq \dim X$ (see [Rob11, Proposition 2.8.iv and Lemma 3.3.iv]) and the inequality can be strict. For example, let $\phi \equiv 0$ and $X = \{(x, f(x)) : x \in [0, 1]\}$ be a graph of a (Hölder continuous) function $f: [0, 1] \to \mathbb{R}$ with dim_H X > 1, e.g. the Weierstrass non-differentiable function. Then the linear projection $L: \mathbb{R}^2 \to \mathbb{R}$ given by L(x, y) = x satisfies $1 = \dim L(X) < \dim_H X$. The following theorem shows that in the non-probabilistic setting, one can obtain β -Hölder continuity of the inverse map for small enough $\beta \in (0, 1)$ (see [BAEFN93, EFNT94, HK99] and [Rob11, Chapter 4]).

Theorem 3.9. Let $X \subset \mathbb{R}^N$ be a compact set. Let $k \in \mathbb{N}$ be such that $k > 2\overline{\dim}_B X$ and let β be such that $0 < \beta < 1 - 2\overline{\dim}_B X/k$. Then Lebesgue almost every linear transformation $L \colon \mathbb{R}^N \to \mathbb{R}^k$ is injective on X with β -Hölder continuous inverse.

However, this is not true in the case of Theorem 3.1.

Remark 3.10. In general, we cannot claim that the injective map $\phi_L|_{X_L}$ from Theorem 3.1 has a Hölder continuous inverse. Indeed, it is well-known that for $n \in \mathbb{N}$ there are examples of compact sets $X \subset \mathbb{R}^N$ of Hausdorff and topological dimension equal to n, which do not embed topologically into \mathbb{R}^k for $k \leq 2n$ (showing the optimality of the bounds in the Menger– Nöbeling embedding theorem, see [HW41, Example V.3]). Consider a probability measure μ on X with $\operatorname{supp} \mu = X$, where supp denotes the topological support of the measure (the intersection of all closed sets of full measure). It is known that such measure exists for any compact set. If the map $\phi_L|_{X_L}$ from Theorem 3.1 for k = n+1 had a Hölder continuous inverse $f = \phi_L^{-1}$, then we could extend f from $\phi_L(X_L)$ to \mathbb{R}^{n+1} preserving the Hölder continuity ([Ban51, Theorem IV.7.5], see also [Min70]). Then $Y = \{x \in X : f \circ \phi_L(x) = x\}$ would be a closed subset of X with $\mu(Y) = 1$, hence Y = X, so ϕ_L would be homeomorphism between X and $\phi_L(X) \subset \mathbb{R}^{n+1}$, which would give a contradiction.

4. PROBABILISTIC TAKENS DELAY EMBEDDING THEOREM

In this section we present the proof of the extended probabilistic Takens delay embedding theorem. It turns out that linear perturbations are insufficient for Takens-type theorems, see Example 4.6. As observed in [SYC91], it is enough to take perturbations over the space of polynomials of degree 2k. This can be easily extended to more general families of functions.

Definition 4.1. Let X be a subset of \mathbb{R}^N . A family of transformations $h_1, \ldots, h_m \colon X \to \mathbb{R}$ is called a *k*-interpolating family on set X, if for every collection of distinct points $x_1, \ldots, x_k \in X$ and every $\xi = (\xi_1, \ldots, \xi_k) \in \mathbb{R}^k$ there exists $(\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$ such that $\alpha_1 h_1(x_i) + \cdots + \alpha_m h_m(x_i) = \xi_i$ for each $i = 1, \ldots, k$. In other words, the matrix

$$\begin{bmatrix} h_1(x_1) & \dots & h_m(x_1) \\ \vdots & \ddots & \vdots \\ h_1(x_k) & \dots & h_m(x_k) \end{bmatrix}$$

has full row rank as a transformation from \mathbb{R}^m to \mathbb{R}^k . Note that the same is true for any collection of l distinct points with $l \leq k$.

Remark 4.2. It is known that any linear basis h_1, \ldots, h_m of the space of real polynomials of N variables of degree at most k-1 is a k-interpolating family (see e.g. [GS00, Section 1.2, eq. (1.9)]).

For a transformation $T: X \to X$ and $p \in \mathbb{N}$ denote by $\operatorname{Per}_p(T)$ the set of periodic points of minimal period p, i.e.

$$\operatorname{Per}_{p}(T) = \{x \in X : T^{p}x = x \text{ and } T^{j}x \neq x \text{ for } j = 1, \dots, p-1\}.$$

Let μ and ν be measures on a measurable space $(\mathcal{X}, \mathcal{F})$. The measure μ is called *singular* with respect to ν , if there exists a measurable set $Y \subset \mathcal{X}$ such that $\mu(\mathcal{X} \setminus Y) = \nu(Y) = 0$. In this case we write $\mu \perp \nu$. By $\mu|_A$ we denote the restriction of μ to a set $A \in \mathcal{F}$.

Theorem 4.3 (Probabilistic Takens delay embedding theorem – extended version). Let $X \subset \mathbb{R}^N$ be a Borel set, μ be a Borel σ -finite measure on X and $T: X \to X$ an injective, locally Lipschitz map. Take $k \in \mathbb{N}$ and $\beta \in (0,1]$ such that $\mathcal{H}^{\beta k}(X) = 0$ and assume $\mu|_{\operatorname{Per}_p(T)} \perp \mathcal{H}^{\beta p}$ for every $p = 1, \ldots, k-1$. Let $h: X \to \mathbb{R}$ be a locally β -Hölder function and $h_1, \ldots, h_m: X \to \mathbb{R}$ a 2k-interpolating family on X consisting of locally β -Hölder functions. For $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$ denote by $h_{\alpha}: X \to \mathbb{R}$ the transformation

$$h_{\alpha}(x) = h(x) + \sum_{j=1}^{m} \alpha_j h_j(x)$$

Then for Lebesgue almost every $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{R}^m$, there exists a Borel set $X_\alpha \subset X$ of full μ -measure, such that the delay-coordinate map

$$\phi_{\alpha}^T \colon X \to \mathbb{R}^k, \qquad \phi_{\alpha}^T(x) = (h_{\alpha}(x), h_{\alpha}(Tx), \dots, h_{\alpha}(T^{k-1}x))$$

is injective on X_{α} .

Notice that Theorem 1.2 follows from Theorem 4.3 by Remark 4.2.

Remark 4.4 (Invariant measure case – extended version). Under the assumptions of Theorem 4.3, the following hold.

- (a) If the measure μ is *T*-invariant, then the set X_{α} can be chosen to satisfy $T(X_{\alpha}) \subset X_{\alpha}$.
- (b) If the measure μ is finite and T-invariant, then the set X_{α} can be chosen to satisfy $T(X_{\alpha}) = X_{\alpha}.$
- (c) If the measure μ is T-invariant and ergodic, then the assumption on the periodic points of T in Theorem 4.3 can be omitted.

Under the notation of Theorem 4.3, we first show a preliminary lemma. For $x \in X$ define its full orbit Orb(x) as

$$Orb(x) = \{T^n x : n \ge 0\} \cup \{y \in X : T^n y = x \text{ for some } n \in \mathbb{N}\}.$$

Note that since T is injective, all full orbits are at most countable, and any two full orbits $\operatorname{Orb}(x)$ and $\operatorname{Orb}(y)$ are either equal or disjoint. For $x, y \in X$ let $D_{x,y}$ be the $k \times m$ matrix defined by

$$D_{x,y} = \begin{bmatrix} h_1(x) - h_1(y) & \dots & h_m(x) - h_m(y) \\ h_1(Tx) - h_1(Ty) & \dots & h_m(Tx) - h_m(Ty) \\ \vdots & \ddots & \vdots \\ h_1(T^{k-1}x) - h_1(T^{k-1}y) & \dots & h_m(T^{k-1}x) - h_m(T^{k-1}y) \end{bmatrix}.$$

Lemma 4.5. For $x, y \in X$, the following statements hold.

- (i) If $y \neq x$, then rank $D_{x,y} \geq 1$.
- (ii) If $y \notin \operatorname{Orb}(x)$ and $y \in \operatorname{Per}_p(T)$ for some $p \in \{1, \ldots, k-1\}$, then $\operatorname{rank} D_{x,y} \ge p$. (iii) If $y \notin \operatorname{Orb}(x)$ and $y \notin \bigcup_{p=1}^{k-1} \operatorname{Per}_p(T)$, then $\operatorname{rank} D_{x,y} = k$.

Proof. For (i), it suffices to observe that the first row of $D_{x,y}$ is non-zero as long as $x \neq y$ and therefore rank $(D_{x,y}) \geq 1$. Indeed, otherwise we would have $h_j(x) = h_j(y)$ for $j = 1, \ldots, m$ which contradicts the fact that h_1, \ldots, h_m is an interpolating family.

Assume now $y \notin \operatorname{Orb}(x)$, which implies $\operatorname{Orb}(y) \cap \operatorname{Orb}(x) = \emptyset$. Let q (resp. r) be a maximal number from $\{1, \ldots, k\}$ such that the points $x, Tx, \ldots, T^{q-1}x$ (resp. $y, Ty, \ldots, T^{r-1}y$) are distinct. Notice that if $y \in \operatorname{Per}_p(T)$ for some $p \in \{1, \ldots, k-1\}$, then r = p, and if $y \notin p$ \bigcup Per_p(T), then r = k. Thus, the assertions (ii)–(iii) of the lemma can be written simply as one condition

(4.1)
$$\operatorname{rank} D_{x,y} \ge r.$$

To show that (4.1) holds, denote the points $x, Tx, \ldots, T^{q-1}x, y, Ty, \ldots, T^{r-1}y$, preserving the order, by z_1, \ldots, z_l , for l = q + r. By the definition of q, r, we have $1 \le l \le 2k$ and the points z_1, \ldots, z_l are distinct. Thus, the matrix $D_{x,y}$ can be written as the product

$$D_{x,y} = \underbrace{J_{x,y}V_{x,y}}_{14}$$

where

$$V_{x,y} = \begin{bmatrix} h_1(z_1) & \dots & h_m(z_1) \\ \vdots & \ddots & \vdots \\ h_1(z_l) & \dots & h_m(z_l) \end{bmatrix}$$

and $J_{x,y}$ is a $k \times l$ matrix with entries in $\{-1, 0, 1\}$ and block structure of the form

$$J_{x,y} = \begin{bmatrix} * & -\operatorname{Id}_{r \times r} \\ \hline * & * \end{bmatrix},$$

where $\operatorname{Id}_{r \times r}$ is the $r \times r$ identity matrix. It follows that rank $J_{x,y} \geq r$. Moreover, since z_1, \ldots, z_l are distinct and h_1, \ldots, h_m is a 2k-interpolating family, the matrix $V_{x,y}$ is of full rank, hence rank $D_{x,y} = \operatorname{rank} J_{x,y} \geq r$, which ends the proof.

Proof of Theorem 4.3. We proceed similarly as in the proof of Theorem 3.1, using Lemma 2.2 instead of Lemma 2.1, together with the suitable rank estimates coming from Lemma 4.5. In the same way as in the proof of Theorem 3.1, we show that it is enough to check that the suitable set X_{α} exists for η_m -almost every $\alpha \in B_m(0, 1)$.

Applying Lemma 2.3 to the sets $\operatorname{Per}_p(T)$, $p = 1, \ldots, k - 1$ and (possibly zero) measures $\mu|_{\operatorname{Per}_p(T)}$, we find (possibly empty) disjoint σ -compact sets $X_1, \ldots, X_{k-1} \subset X$ such that

$$X_p \subset \operatorname{Per}_p(T), \quad \mu(\operatorname{Per}_p(T) \setminus X_p) = 0, \quad \mathcal{H}^{\beta p}(X_p) = 0 \quad \text{for } p = 1, \dots, k-1.$$

Similarly, there exists a σ -compact set $X_k \subset X \setminus \bigcup_{p=1}^{k-1} \operatorname{Per}_p(T)$ such that

$$\mu\left(\left(X \setminus \bigcup_{p=1}^{k-1} \operatorname{Per}_p(T)\right) \setminus X_k\right) = 0 \quad \text{and} \quad \mathcal{H}^{\beta k}(X_k) = 0.$$

Note that X_k contains both aperiodic and periodic points (with period at least k). Let

$$\tilde{X} = \bigcup_{p=1}^{k} X_p.$$

Then $\tilde{X} \subset X$ is a σ -compact set of full μ -measure. Define

$$A = \{ (x, \alpha) \in \tilde{X} \times B_m(0, 1) : \phi_\alpha^T(x) = \phi_\alpha^T(y) \text{ for some } y \in \tilde{X} \setminus \{x\} \}.$$

The set A is Borel by Lemma 2.4. For $x \in \tilde{X}$ and $\alpha \in B_m(0,1)$, denote, respectively, by A_x and A^{α} , the Borel sections

$$A_x = \{ \alpha \in B_m(0,1) : (x,\alpha) \in A \}, \quad A^\alpha = \{ x \in \tilde{X} : (x,\alpha) \in A \}.$$

Observe that to show the injectivity of ϕ_{α}^{T} on a set of full μ -measure, it is enough to prove $\eta_{m}(A_{x}) = 0$ for every $x \in \tilde{X}$, since then by Fubini's theorem ([Rud87, Thm. 8.8]), ($\eta_{m} \otimes \mu$)(A) = 0 and, consequently, $\mu(A^{\alpha}) = 0$ for η_{m} -almost every $\alpha \in B_{m}(0, 1)$. As ϕ_{α}^{T} is injective on $\tilde{X} \setminus A^{\alpha}$ and \tilde{X} has full μ -measure, the proof of the claim is finished.

Fix $x \in \tilde{X}$. To show $\eta_m(A_x) = 0$, note that for $y \in \tilde{X}$,

(4.2)
$$\phi_{\alpha}^{T}(x) - \phi_{\alpha}^{T}(y) = D_{x,y}\alpha + w_{x,y}$$

for

$$w_{x,y} = \begin{bmatrix} h(x) - h(y) \\ h(Tx) - h(Ty) \\ \vdots \\ h(T^{k-1}x) - h(T^{k-1}y) \end{bmatrix}.$$

Write A_x as

$$A_x = A_x^{\text{orb}} \cup \bigcup_{p=1}^k A_x^p,$$

where

$$A_x^{\text{orb}} = \{ \alpha \in B_m(0,1) : \phi_\alpha^T(x) = \phi_\alpha^T(y) \text{ for some } y \in \tilde{X} \cap \operatorname{Orb}(x) \setminus \{x\} \},\$$
$$A_x^p = \{ \alpha \in B_m(0,1) : \phi_\alpha^T(x) = \phi_\alpha^T(y) \text{ for some } y \in X_p \setminus \{x\} \}, \quad p = 1, \dots, k.$$

The set A_x^{orb} is Borel as a countable union of closed sets of the form

(4.3)
$$\{\alpha \in B_m(0,1) : \phi_\alpha^T(x) = \phi_\alpha^T(y)\}, \quad y \in \tilde{X} \cap \operatorname{Orb}(x) \setminus \{x\},\$$

while each set A_x^p is Borel as a section of the set

$$\{(x,\alpha) \in \tilde{X} \times B_m(0,1) : \phi_\alpha^T(x) = \phi_\alpha^T(y) \text{ for some } y \in X_p \setminus \{x\}\},\$$

which is Borel by Lemma 2.4. To end the proof, it is enough to show that the sets A_x^{orb} and $A_x^p, p = 1, \ldots, k$, have η_m measure zero.

To prove $\eta_m(A_x^{\text{orb}}) = 0$ it suffices to check that the sets of the form (4.3) have η_m measure zero. By (4.2), we have

$$\{\alpha \in B_m(0,1) : \phi_{\alpha}^T(x) = \phi_{\alpha}^T(y)\} = \{\alpha \in B_m(0,1) : D_{x,y}\alpha = -w_{x,y}\}$$

and Lemma 4.5 gives rank $D_{x,y} \ge 1$ whenever $y \ne x$, so each set of the form (4.3) is contained in an affine subspace of \mathbb{R}^m of codimension at least 1. Consequently, it has η_m measure zero.

Since T is locally Lipschitz, h, h_1, \ldots, h_m are locally β -Hölder and X is separable, there exists a countable covering \mathcal{V} of X by open sets in \mathbb{R}^N , such that for every $V \in \mathcal{V}$, the map T is Lipschitz on V and h, h_1, \ldots, h_m are β -Hölder on V. Let \mathcal{U} be the collection of all sets of the form $U = V_0 \cap T^{-1}(V_1) \cap \ldots \cap T^{-(k-1)}(V_{k-1})$, where $V_0, \ldots, V_{k-1} \in \mathcal{V}$. Then \mathcal{U} is a countable covering of X by open sets, and we can write $\mathcal{U} = \{U_j\}_{j \in \mathbb{N}}$. By definition, for every $j \in \mathbb{N}$ there exists $C_j > 0$ such that

$$||T^{s+1}(y) - T^{s+1}(y')|| \le C_j ||T^s(y) - T^s(y')||,$$

$$||h(T^s(y)) - h(T^s(y'))|| \le C_j ||T^s(y) - T^s(y')||^{\beta}$$

$$||h_r(T^s(y)) - h_r(T^s(y'))|| \le C_j ||T^s(y) - T^s(y')||^{\beta}$$

for every $y, y' \in U_i \cap X$, $s \in \{0, \dots, k-1\}$, $r \in \{1, \dots, m\}$. By induction, it follows that $||T^{s}(y) - T^{s}(y')|| \le C_{1}^{s}||y - y'||$

(4.4)
$$\|h(T^{s}(y)) - h(T^{s}(y'))\| \leq C_{j}^{\beta s+1} \|y - y'\|^{\beta}, \\\|h_{r}(T^{s}(y)) - h_{r}(T^{s}(y'))\| \leq C_{j}^{\beta s+1} \|y - y'\|^{\beta}$$

for $y, y' \in U_j \cap X$, $s \in \{0, \dots, k-1\}$, $r \in \{1, \dots, m\}$.

To prove $\eta_m(A_x^p) = 0$ for p = 1, ..., k, we follow the strategy used in [SYC91] (see also [Rob11]). Fix $n \in \mathbb{N}$ and for $j \in \mathbb{N}$ define

$$X_x^{p,n} = \left\{ y \in X_p : \sigma_p(D_{x,y}) \ge \frac{1}{n} \right\},\$$

$$A_x^{p,j,n} = \left\{ \alpha \in B_m(0,1) : \phi_\alpha^T(x) = \phi_\alpha^T(y) \text{ for some } y \in U_j \cap X_x^{p,n} \setminus \{x\} \right\}$$

where $\sigma_p(D_{x,y})$ is the *p*-th largest singular value. Note that singular values of given order depend continuously on the coefficients of the matrix, see e.g. [GVL13, Corollary 8.6.2]. Hence, the set $X_x^{p,n}$ is σ -compact as a closed subset of X_p and by Lemma 2.4, the set $A_x^{p,j,n}$ is Borel.

By Lemma 4.5, for every $y \in X_p \setminus \operatorname{Orb}(x)$ we have rank $D_{x,y} \ge p$. This implies $\sigma_p(D_{x,y}) > 0$ (see e.g. [Rob11, Lemma 14.2]). Hence,

$$A_x^p \setminus A_x^{\operatorname{orb}} = \bigcup_{j=1}^\infty \bigcup_{n=1}^\infty A_x^{p,j,n} \setminus A_x^{\operatorname{orb}}.$$

Consequently, it is enough to prove $\eta_m(A_x^{p,j,n} \setminus A_x^{\text{orb}}) = 0$ for every $n \in \mathbb{N}$.

Fix $\varepsilon > 0$. Since $\mathcal{H}^{\beta p}(U_j \cap X_x^{p,n} \setminus \operatorname{Orb}(x)) \leq \mathcal{H}^{\beta p}(X_p) = 0$, there exists a collection of balls $B_N(y_i, \varepsilon_i)$, for $y_i \in U_j \cap X_x^{p,n} \setminus \operatorname{Orb}(x)$ and $0 < \varepsilon_i < \varepsilon$, $i \in \mathbb{N}$, such that

(4.5)
$$U_j \cap X_x^{p,n} \setminus \operatorname{Orb}(x) \subset \bigcup_{i \in \mathbb{N}} B_N(y_i, \varepsilon_i) \quad \text{and} \quad \sum_{i=1}^{\infty} \varepsilon_i^{\beta p} \leq \varepsilon.$$

Take $\alpha \in A_x^{p,j,n} \setminus A_x^{\text{orb}}$ and let $y \in U_j \cap X_x^{p,n} \setminus \operatorname{Orb}(x)$ be such that $\phi_{\alpha}^T(x) = \phi_{\alpha}^T(y)$. Then for y_i with $y \in B(y_i, \varepsilon_i)$ we have

(4.6)
$$\|D_{x,y_{i}}\alpha + w_{x,y_{i}}\| = \|\phi_{\alpha}^{T}(x) - \phi_{\alpha}^{T}(y_{i})\| = \|\phi_{\alpha}^{T}(y) - \phi_{\alpha}^{T}(y_{i})\| \\ \leq \sqrt{\sum_{s=0}^{k-1} \left(\|h(T^{s}y) - h(T^{s}y_{i})\| + \sum_{r=1}^{m} \alpha_{r} \|h_{r}(T^{s}y) - h_{r}(T^{s}y_{i})\| \right)^{2}} \\ \leq M_{j} \|y - y'\|^{\beta} \leq M_{j} \varepsilon_{i}^{\beta}$$

for

$$M_j = (1 + \sqrt{m}) \sqrt{\sum_{s=0}^{k-1} C_j^{2(\beta s+1)}},$$

by (4.4) and the fact $\alpha \in B_m(0,1)$. By (4.6),

$$A_x^{p,j,n} \setminus A_x^{\text{orb}} \subset \bigcup_{i \in \mathbb{N}} \{ \alpha \in B_m(0,1) : \|D_{x,y_i}\alpha + w_{x,y_i}\| \le M_j \varepsilon_i^\beta \}.$$

Since for every $i \in \mathbb{N}$ we have $\sigma_p(D_{x,y_i}) \geq 1/n$, we can apply Lemma 2.2 and (4.5) to obtain

$$\eta_m(A_x^{p,j,n} \setminus A_x^{\text{orb}}) \le \sum_{i=1}^{\infty} C_{m,k} \frac{M_j^p \varepsilon_i^{\beta p}}{1/n^p} \le C_{m,k} M_j^p n^p \varepsilon_i$$

Since $\varepsilon > 0$ was arbitrary, we conclude that $\eta_m(A_x^{p,j,n} \setminus A_x^{\text{orb}}) = 0$, so in fact $\eta_m(A_x^{p,j,n}) = 0$. This ends the proof of Theorem 4.3. *Proof of Remark* 4.4. Suppose that the measure μ is *T*-invariant. Then it is easy to check that the set

$$\tilde{X}_{\alpha} = \bigcap_{n=0}^{\infty} T^{-n}(X_{\alpha})$$

is a Borel subset of X_{α} of full μ -measure satisfying $T(\tilde{X}_{\alpha}) \subset \tilde{X}_{\alpha}$. Hence, to show (a), it suffices to replace the set X_{α} by \tilde{X}_{α} .

In the case when μ is additionally finite, we first remark that the measure μ is also forward invariant, i.e. $\mu(T(Y)) = \mu(Y)$ for Borel sets $Y \subset X$. Note that if Y is Borel, then so is T(Y) as the image of a Borel set under a continuous and injective mapping (see e.g. [Kec95, Theorem 15.1]). Using this together with the invariance of μ and the injectivity of T, we check that

$$\tilde{X}_{\alpha} = \bigcap_{n \in \mathbb{Z}} T^{-n}(X_{\alpha})$$

is a Borel subset of X_{α} of full μ -measure satisfying $T(\tilde{X}_{\alpha}) = \tilde{X}_{\alpha}$. This gives (b). Notice that the finiteness of μ is indeed necessary, as for $X = \mathbb{N}$, T(x) = x + 1 and μ the counting measure, there does not exist a set $Y \subset X$ of full μ -measure satisfying T(Y) = Y.

To show (c), suppose that μ is *T*-invariant and ergodic. Obviously, we can assume that the μ -measure of the set of all periodic points of *T* is positive (including $+\infty$). Then there exists $p \in \mathbb{N}$ such that the measure of the set *P* of all *p*-periodic points of *T* is positive (including $+\infty$).

Suppose first that μ restricted to P is non-atomic. Then there exists a Borel set $Y \subset P$ with $0 < \mu(Y) < \mu(X)/p$. Let $Z = Y \cup T^{-1}(Y) \cup \ldots \cup T^{-(p-1)}(Y)$. Then $0 < \mu(Z) < \mu(X)$ and, by the injectivity of T, we have $T^{-1}(Z) = Z$, which contradicts the ergodicity of μ .

Suppose now that μ has an atom in P. Since μ is a Borel σ -finite measure in a Euclidean space, this is equivalent to the fact that $\mu(\{x\}) > 0$ for some $x \in P$. Let \mathcal{O} be the periodic orbit of x. Again by the injectivity of T, we have $T^{-1}(\mathcal{O}) = \mathcal{O}$, so by the ergodicity of μ , the set \mathcal{O} has full μ -measure. This means that μ is supported on a set of Hausdorff dimension 0, which obviously gives (c).

The original Takens delay embedding theorem states that for given finite dimensional C^2 manifold M and generic pair of C^2 -diffeomorphism $T: M \to M$ and C^2 -function $h: M \to \mathbb{R}$, the corresponding delay-coordinate map $\phi: M \to \mathbb{R}^k$, $\phi(x) = (h(x), h(Tx), \dots, h(T^{k-1}x))$ is a C^2 -embedding (an injective immersion) as long as $k > 2 \dim M$. It was followed by the boxcounting dimension version of Sauer, Yorke and Casdagli (Theorem 1.1) and subsequently by the infinite-dimensional result of [Rob05] (see also [Rob11, Section 14.3]). Refer to [NV18] for a version of Takens' theorem with a fixed observable and perturbation performed on the dynamics. Takens' theorem involving Lebesgue covering dimension on compact metric spaces and a continuous observable was proved in [Gut16] (see [GQS18] for a detailed proof). See also [Sta99, Cab00] for Takens theorem for deterministically driven smooth systems and [SBDH97, SBDH03] for stochastically driven smooth systems.

Example 4.6. It turns out that linear perturbations are not sufficient for Theorems 1.1 and 4.3, i.e. it may happen that $\phi_L = (\phi(x) + Lx, \dots, \phi(T^{k-1}x) + LT^{k-1}x)$ is not (almost

surely) injective for a generic linear map $L: \mathbb{R}^N \to \mathbb{R}$. As an example, let $X = B_2(0, 1)$, fix $a \in (0, 1)$ and define $T: X \to X$ as

$$T(x) = ax.$$

Then T is a Lipschitz injective transformation on the unit disc $X \subset \mathbb{R}^2$ with zero being the unique periodic point. Fix $\phi \equiv 0$. We claim that there is no linear observable $L: \mathbb{R}^2 \to \mathbb{R}$ which makes the delay map injective, i.e. for every $k \in \mathbb{N}$ and every $v \in \mathbb{R}^2$ the transformation $x \mapsto \phi_v^T(x) = (\langle x, v \rangle, \langle Tx, v \rangle, \dots, \langle T^{k-1}x, v \rangle) \in \mathbb{R}^k$ is not injective on X. This follows from the fact that for each 1-dimensional linear subspace $W \subset \mathbb{R}^2$ the set $W \cap X$ is T-invariant, hence $\phi_v^T = 0$ on an infinite set $\operatorname{Ker}(\langle \cdot, v \rangle) \cap X$. We have seen that ϕ_v^T is not injective for any $v \in \mathbb{R}^2$. No we will see that it also not almost surely injective for μ being the Lebesgue measure on X. Note that for $v \in \mathbb{R}^2$ and $c \in \mathbb{R}$, the segment $W_c = \{z \in X : \langle z, v \rangle = c\}$ satisfies $T(W_c) \subset W_{ac}$, hence all points on W_c will have the same observation vector $(\langle x, v \rangle, \langle Tx, v \rangle, \dots, \langle T^{k-1}x, v \rangle) = (c, ac, a^2c, \dots, a^{k-1}c)$. Therefore, a set $X_v \subset X$ on which ϕ_v^T is injective can only have one point on each of the parallel segments W_c contained in X. However, such a set X_v cannot be of full Lebesgue measure. Note that the above example can be easily modified to make T a homeomorphism.

5. EXAMPLES

In this section we present two examples which illustrate the usage of Theorem 3.1. Let us begin with fixing some notation. For $x \in [0, 2)$ we will write

$$x = x_0 . x_1 x_2 \dots,$$

where $x_0.x_1x_2...$ is the binary expansion of x, i.e.

$$x = \sum_{j=0}^{\infty} \frac{x_j}{2^j}, \quad x_0, x_1, x_2, \ldots \in \{0, 1\}.$$

For a dyadic rational we agree to choose its eventually terminating expansion, i.e. the one with $x_j = 0$ for j large enough. Let $\pi \colon \{0, 1\}^{\mathbb{N}} \to [0, 1]$ be the coding map

$$\pi(x_1, x_2, \ldots) = \sum_{j=1}^{\infty} \frac{x_j}{2^j}.$$

5.1. A modified Kan example. In the Appendix to [SYC91], Kan presented an example of a compact set $K \subset \mathbb{R}^N$ with $\dim_H K = 0$ and such that every linear transformation $L: \mathbb{R}^N \to \mathbb{R}^{N-1}$ fails to be injective on K (see also Remark 3.8). It follows from Theorem 3.1, that whenever we are given a Borel σ -finite measure μ on such a set, then almost every linear transformation $L: \mathbb{R}^N \to \mathbb{R}$ is injective on a set of full μ -measure. To illustrate this, we construct a σ -compact set $X \subset \mathbb{R}^2$ with $\dim_H X = 0$, which is a slight modification of Kan's example, equipped with a natural Borel σ -finite measure μ , such that no linear transformation $L: \mathbb{R}^2 \to \mathbb{R}$ is injective on X, while for almost every L we explicitly show a set $X_L \subset X$ of full μ -measure, such that L is injective on X_L . Following [SYC91, Appendix], we begin with constructing compact sets $A, B \subset [0, 1]$ such that

(5.1)
$$\dim_H A = \underline{\dim}_B A = \dim_H B = \underline{\dim}_B B = 0 \quad (\text{hence } \dim_H (A \cup B) = 0),$$

and

(5.2)
$$\overline{\dim}_B A = \overline{\dim}_B B = 1, \quad \underline{\dim}_B (A \cup B) = \overline{\dim}_B (A \cup B) = 1.$$

To this aim, let M_k , $k \ge 0$, be an increasing sequence of positive integers such that $M_0 = 1$ and $M_k \nearrow \infty$ with $\lim_{k\to\infty} \frac{M_{k+1}}{M_k} = \infty$. Define

$$\widetilde{A} = \{ (x_1, x_2, \ldots) \in \{0, 1\}^{\mathbb{N}} : \text{ for every even } k, x_j = 0 \text{ for all } j \in [M_k, M_{k+1}) \\ \text{ or } x_j = 1 \text{ for all } j \in [M_k, M_{k+1}) \}, \\ \widetilde{B} = \{ (x_1, x_2, \ldots) \in \{0, 1\}^{\mathbb{N}} : \text{ for every odd } k, x_j = 0 \text{ for all } j \in [M_k, M_{k+1}) \\ \text{ or } x_j = 1 \text{ for all } j \in [M_k, M_{k+1}) \}, \end{cases}$$

and set

$$A = \pi(\widetilde{A}), \quad B = \pi(\widetilde{B})$$

It is a straightforward calculation to check that A and B satisfy (5.1) and (5.2) (see [SYC91, Appendix], [Fal14, Example 7.8] or [Rob11, Section 6.1]). Define $X \subset \mathbb{R}^2$ as

$$X = \left(\{0\} \times \bigcup_{n \in \mathbb{Z}} (A+n)\right) \cup \left(\{1\} \times \bigcup_{n \in \mathbb{Z}} (B+n)\right).$$

By (5.1), we have $\dim_H X = 0$. The following two propositions describe the embedding properties of the set X.

Proposition 5.1. No linear transformation $L: \mathbb{R}^2 \to \mathbb{R}$ is injective on X.

Proof. The map L has the form $L(x, y) = \alpha x + \beta y$ for $\alpha, \beta \in \mathbb{R}$. Obviously, we can assume $\beta \neq 0$. Note that the points

$$u = (0, a + n), \quad v = (1, b + m), \quad \text{for } a \in A, \ b \in B, \ n, m \in \mathbb{Z}$$

are in X and

(5.3)
$$L(u) = L(v)$$
 if and only if $b - a = z$,

where

$$z = -\frac{\alpha}{\beta} + n - m.$$

For given α and β , choose $n, m \in \mathbb{Z}$ such that $z \in [0, 1)$. Consider the binary expansion $z = 0.z_1 z_2 \dots$ and define

$$a = 0.a_1 a_2 \ldots \in A, \quad b = 0.b_1 b_2 \ldots \in B$$

setting

(5.4)
$$a_{j} = 0, \qquad b_{j} = z_{j} \qquad \text{for } j \in [M_{k}, M_{k+1}), \text{ if } k \text{ is even}, \\ a_{j} = 1 - z_{j}, \qquad b_{j} = 1 \qquad \text{for } j \in [M_{k}, M_{k+1}), \text{ if } k \text{ is odd}$$

(if all b_j are equal to 1, we set b = 1). Then z = b - a and (5.3) implies that L is not injective on X.

Let us now define a natural Borel σ -finite measure μ on X, starting from a pair of probability measures ν_1, ν_2 on \widetilde{A} and \widetilde{B} , respectively. Let

$$\nu_1 = \bigotimes_{k=0}^{\infty} \mathbf{p}_k, \quad \nu_2 = \bigotimes_{k=0}^{\infty} \mathbf{q}_k,$$

where \mathbf{p}_k and \mathbf{q}_k are probability measures on $\{0,1\}^{M_{k+1}-M_k}$ given as

$$\mathbf{p}_{k} = \begin{cases} \frac{1}{2}\delta_{(0,\dots,0)} + \frac{1}{2}\delta_{(1,\dots,1)} & \text{if } k \text{ is even} \\ \left(\frac{1}{2}\delta_{0} + \frac{1}{2}\delta_{1}\right)^{\otimes(M_{k+1} - M_{k})} & \text{if } k \text{ is odd} \end{cases}, \quad \mathbf{q}_{k} = \begin{cases} \left(\frac{1}{2}\delta_{0} + \frac{1}{2}\delta_{1}\right)^{\otimes(M_{k+1} - M_{k})} & \text{if } k \text{ is even} \\ \frac{1}{2}\delta_{(0,\dots,0)} + \frac{1}{2}\delta_{(1,\dots,1)} & \text{if } k \text{ is odd} \end{cases}$$

and the symbol δ_a denotes the Dirac measure at a. Then supp $\nu_1 = \widetilde{A}$, supp $\nu_2 = \widetilde{B}$, hence defining

$$\mu_1 = \pi_*(\nu_1), \quad \mu_2 = \pi_*(\nu_2),$$

we obtain probability measures on A, B, respectively, with $\operatorname{supp} \mu_1 = A$, $\operatorname{supp} \mu_2 = B$. Finally, let

$$\mu = \sum_{n \in \mathbb{Z}} \delta_0 \otimes (\tau_n)_* \mu_1 + \sum_{n \in \mathbb{Z}} \delta_1 \otimes (\tau_n)_* \mu_2,$$

where $\tau_n \colon \mathbb{R} \to \mathbb{R}, \ \tau_n(x) = x + n, \ n \in \mathbb{Z}$. Clearly, μ is a Borel σ -finite measure with $\operatorname{supp} \mu = X$.

For $a \in A$, $b \in B$ let

$$A_a = \{x \in A \setminus \{1\} : x + a = z_0.z_1z_2\dots \text{ such that the sequence } (z_0, z_1, \dots) \\ \text{ is constant on } [M_k, M_{k+1} - 1) \cap \mathbb{N} \text{ for every odd } k\}, \\ B_b = \{x \in B \setminus \{1\} : x + b = z_0.z_1z_2\dots \text{ such that the sequence } (z_0, z_1, \dots) \\ \text{ is constant on } [M_k, M_{k+1} - 1) \cap \mathbb{N} \text{ for every even } k\}.$$

Lemma 5.2. For every $a \in A$, $b \in B$, we have $\mu_1(A_a) = \mu_2(B_b) = 0$.

Proof. Fix $b = b_0.b_1b_2... \in B$. We will show $\mu_2(B_b) = 0$ (the fact $\mu_1(A_a) = 0$ can be proved analogously). The proof proceeds by showing that for each even k, the vector $(x_{M_k}, \ldots, x_{M_{k+1}-2})$, where $x = x_0.x_1x_2... \in B_b$, can assume at most four values. This will imply $\mu_2(B_b) \leq 8 \cdot 2^{-(M_{k+1}-M_k)}$ for each even k and, consequently, $\mu_2(B_b) = 0$. To show the assertion, fix an even k and let

$$\xi = \sum_{j=M_{k+1}-1}^{\infty} \frac{x_j + b_j}{2^j}$$

Note that $\xi < 2^{-(M_{k+1}-3)}$ (as $\xi < 2$ and we exclude expansions with digits eventually equal to 1). Hence, $\xi = \xi_0.\xi_1\xi_2...$ with $\xi_j = 0$ for $j \le M_{k+1}-3$. Note that, since b is fixed, the values of $\xi_{M_{k+1}-2} \in \{0,1\}$ and $(x_{M_k}+b_{M_k},\ldots,x_{M_{k+1}-2}+b_{M_{k+1}-2}) \in \{(0,\ldots,0),(1,\ldots,1)\}$ determine uniquely the value of $(x_{M_k},\ldots,x_{M_{k+1}-2})$. Therefore, $(x_{M_k},\ldots,x_{M_{k+1}-2})$ can assume at most four values.

Now for Lebesgue almost every linear transformation $L: \mathbb{R}^2 \to \mathbb{R}$ we will construct a set $X_L \subset X$ of full μ -measure, such that L is injective on X_L . As previously, write $L(x, y) = \alpha x + \beta y$ for $\alpha, \beta \in \mathbb{R}$. Neglecting a set of zero Lebesgue measure, we can assume $\beta \neq 0$. Let $l \in \mathbb{Z}$ be such that

(5.5)
$$z = -\frac{\alpha}{\beta} + l \text{ belongs to } [0,1).$$

Similarly as in (5.4), we can write

(5.6)
$$z = a' - b', \quad z - 1 = a'' - b'' \text{ for some } a', a'' \in A, \ b', b'' \in B.$$

Let

$$X_L = \left(\{0\} \times \bigcup_{n \in \mathbb{Z}} (A+n)\right) \cup \left(\{1\} \times \bigcup_{n \in \mathbb{Z}} \left((B \setminus (B_{b'} \cup B_{b''} \cup \{1\})) + n \right) \right).$$

Then $X_L \subset X$ and Lemma 5.2 implies that X_L has full μ -measure.

Proposition 5.3. For every $\alpha \in \mathbb{R}$, $\beta \in \mathbb{R} \setminus \{0\}$, the linear transformation $L \colon \mathbb{R}^2 \to \mathbb{R}$, $L(x, y) = \alpha x + \beta y$, is injective on X_L .

For the proof of the proposition we will need the following simple lemma. The proof is left to the reader.

Lemma 5.4. Let $x = x_0.x_1x_2... \in [0, 1]$, $y = y_0.y_1y_2... \in [0, 1]$, $M, N \in \mathbb{N}$, M < N - 1, be such that x + y < 2 and sequences $(x_M, ..., x_N)$ and $(y_M, ..., y_N)$ are constant. Then $x + y = z_0.z_1z_2...$, where the sequence $(z_M, ..., z_{N-1})$ is constant.

Proof of Proposition 5.3. Assume, on the contrary, that there exist points $u, v \in X_L$ such that L(u) = L(v). As $\beta \neq 0$, we cannot have $u, v \in \{0\} \times \mathbb{R}$ or $u, v \in \{1\} \times \mathbb{R}$. Hence, we can assume $u \in \{0\} \times \mathbb{R}, v \in \{1\} \times \mathbb{R}$. Then, following the previous notation, we have u = (0, a + n), v = (1, b + m) for $a \in A, b \in B \setminus (B_{b'} \cup B_{b''} \cup \{1\}), n, m \in \mathbb{Z}$. Note that $b - a \in [-1, 1)$, so by (5.3), we have

$$b-a=z$$
 or $b-a=z-1$,

for z from (5.5), and (5.6) implies

$$b - a = a' - b'$$
 or $b - a = a'' - b''$.

Hence,

$$a + a' = b + b'$$
 or $a + a'' = b + b''$

This is a contradiction, as Lemma 5.4 implies that the binary expansion sequences of a + a'and a + a'' are constant on $[M_k, M_{k+1} - 1) \cap \mathbb{N}$ for every even k, while by the condition $b \in B \setminus (B_{b'} \cup B_{b''} \cup \{1\})$, the binary expansion sequences of b + b' and b + b'' are not constant on $[M_k, M_{k+1} - 1) \cap \mathbb{N}$ for some even k. 5.2. Measure with $\dim_H \mu < \underline{\dim}_{MB} \mu$. To show that Theorem 3.1 is an actual strengthening of Theorem 3.6, we present an example of a measure μ , for which $\dim_H \mu < \underline{\dim}_{MB} \mu$. More precisely, we show the following.

Theorem 5.5. There exists a Borel probability measure μ on $[0, 1]^2$, such that $\dim_H \mu = 1$ and $\underline{\dim}_{MB} \mu = 2$.

To begin the construction of μ , fix an increasing sequence of positive integers N_k , $k \in \mathbb{N}$, such that $N_k \nearrow \infty$ with $\frac{S_k}{S_{k+1}} \le \frac{1}{k+1}$, where $S_k = \sum_{j=1}^k N_j$. Consider the probability distributions $\mathbf{p}_0, \mathbf{p}_1$ on $\{0, 1\}$ given by

$$\mathbf{p}_0(\{0\}) = 0, \ \mathbf{p}_0(\{1\}) = 1, \qquad \mathbf{p}_1(\{0\}) = \mathbf{p}_1(\{1\}) = \frac{1}{2}$$

For $y = 0.y_1y_2... \in [0, 1]$ (in this subsection we assume that the binary expansion of 1 is 0.111...), define the probability measure ν_y on $\{0, 1\}^{\mathbb{N}}$ as the infinite product

$$\nu_y = \bigotimes_{j=1}^{\infty} \bigotimes_{i=1}^{N_j} \mathbf{p}_{y_j}.$$

Further, let μ_y be the Borel probability measure on [0, 1] given by

$$\mu_y = \pi_* \nu_y.$$

Finally, let μ be the Borel probability measure on $[0, 1]^2$ defined as

$$\mu = \int_{[0,1]} \mu_y d\operatorname{Leb}(y), \quad \text{i.e. } \mu(A) = \int_{[0,1]} \mu_y(A^y) d\operatorname{Leb}(y) \text{ for a Borel set } A \subset [0,1]^2,$$

where $A^y = \{x \in [0, 1] : (x, y) \in A\}$. It is easy to see that μ is well-defined, as the function $y \mapsto \mu_y(A^y)$ is measurable for every Borel set $A \subset [0, 1]^2$.

The proof of Theorem 5.5 is based on the analysis of the local dimension of μ , defined in terms of dyadic squares (rather then balls). For $n \in \mathbb{N}$ and $x_1, \ldots, x_n \in \{0, 1\}$ let $[x_1, \ldots, x_n]$ denote the dyadic interval corresponding to the sequence (x_1, \ldots, x_n) , i.e.

$$[x_1, \dots, x_n] = \begin{cases} \left[\sum_{j=1}^n \frac{x_j}{2^j}, \sum_{j=1}^n \frac{x_j}{2^j} + \frac{1}{2^n}\right) & \text{if } \sum_{j=1}^n \frac{x_j}{2^j} + \frac{1}{2^n} < 1\\ \left[1 - \frac{1}{2^n}, 1\right] & \text{otherwise.} \end{cases}$$

Under this notation, for $n \in \mathbb{N}$ and $(x, y) \in [0, 1]^2$ let $D_n(x, y)$ be the dyadic square of sidelength 2^{-n} containing (x, y), i.e.

$$D_n(x,y) = [x_1, \dots, x_n] \times [y_1, \dots, y_n],$$
 where $x = 0.x_1x_2...$ and $y = 0.y_1y_2...$

Recall that the box-dimensions can be defined equivalently in terms of dyadic squares. Precisely, let $N'(X, 2^{-n})$ be the number of dyadic squares $D_n(x, y)$ of sidelength 2^{-n} intersecting X. Then (see e.g. [Fal14, Section 2.1])

(5.7)
$$\underline{\dim}_B(X) = \liminf_{n \to \infty} \frac{\log N'(X, 2^{-n})}{n \log 2} \text{ and } \overline{\dim}_B(X) = \limsup_{n \to \infty} \frac{\log N'(X, 2^{-n})}{n \log 2}.$$

For a Borel finite measure μ on $[0,1]^2$ and $(x,y) \in [0,1]^2$ define the *lower* and *upper local* dimension of μ at (x,y) as

$$\underline{d}(\mu, (x, y)) = \liminf_{n \to \infty} \frac{-\log \mu(D_n(x, y))}{n \log 2}, \quad \overline{d}(\mu, (x, y)) = \limsup_{n \to \infty} \frac{-\log \mu(D_n(x, y))}{n \log 2}.$$

It is well-known (see e.g. [Hoc14, Propositions 3.10 and 3.20]) that

(5.8)
$$\dim_{H} \mu = \underset{(x,y)\sim\mu}{\operatorname{ess sup}} \underline{d}(\mu, (x, y)).$$

The following lemma gives estimates on the measure of dyadic squares at suitable scales.

Lemma 5.6. Let $x = 0.x_1x_2..., \in [0,1], y = 0.y_1y_2... \in [0,1], n \in \mathbb{N}$ and $D = D_n(x,y) = [x_1,...,x_n] \times [y_1,...,y_n]$. Let $k \in \mathbb{N}$ be such that $S_k < n \leq S_{k+1}$. Then the following hold. (a) If $y_k = y_{k+1} = 1$, then $\mu(D) \leq 2^{-(2-\frac{1}{k})n}$. (b) If $n = S_{k+1}$ and $y_{k+1} = 0$, then either $\mu(D) = 0$ or $\mu(D) \geq 2^{-(1+\frac{1}{k+1})n}$.

Proof. Note that for $y' = 0.y'_1y'_2... \in [0,1]$ such that $(y'_1,...,y'_n) = (y_1,...,y_n)$ we have

(5.9)
$$\mu_{y'}(D^{y'}) = \mu_{y'}([x_1, \dots, x_n]) = \mathbf{p}_{y'_1}(\{x_1\}) \cdots \mathbf{p}_{y'_1}(\{x_{S_1}\})\mathbf{p}_{y'_2}(\{x_{S_1+1}\}) \cdots \mathbf{p}_{y'_2}(\{x_{S_2}\}) \\ \cdots \mathbf{p}_{y'_{k+1}}(\{x_{S_k+1}\}) \cdots \mathbf{p}_{y'_{k+1}}(\{x_n\}).$$

Moreover, as k < n, the value of $\mu_{y'}(D^{y'})$ depends only on (y_1, \ldots, y_n) and (x_1, \ldots, x_n) . Using (5.9), we can prove both assertions of the lemma, as follows. Ad (a).

If $y_k = y_{k+1} = 1$, then for $j \in \{S_{k-1} + 1, \dots, n\}$ we have $\mathbf{p}_{y_l}(x_j) = \frac{1}{2}$, where $l \in \{k, k+1\}$ is such that $S_{l-1} < j \leq S_l$. Therefore, in the product (5.9) there is at least $n - S_{k-1}$ terms equal to $\frac{1}{2}$. Consequently,

$$\mu_{y'}(D^{y'}) \le 2^{-(n-S_{k-1})} = 2^{-(1-\frac{S_{k-1}}{n})n} \le 2^{-(1-\frac{S_{k-1}}{S_k})n} \le 2^{-(1-\frac{1}{k})n},$$

hence

$$\mu(D) = \int_{[y_1,\dots,y_n]} \mu_{y'}(D^{y'}) d\operatorname{Leb}(y') \le \operatorname{Leb}([y_1,\dots,y_n]) 2^{-n(1-\frac{1}{k})} = 2^{-n(2-\frac{1}{k})}.$$

Ad (b).

Assume that $\mu(D) \neq 0$. Then all the terms in (5.9) have to be non-zero, so every term is equal to either $\frac{1}{2}$ or 1. Moreover, as $y_{k+1} = 0$ and $n = S_{k+1}$, we have

$$\mathbf{p}_{y_{k+1}}(\{x_{S_k+1}\})\cdots\mathbf{p}_{y_{k+1}}(\{x_n\})=1$$

and, consequently,

$$\mu(D) = 2^{-n} \mathbf{p}_{y_1}(\{x_1\}) \cdots \mathbf{p}_{y_1}(\{x_{S_1}\}) \mathbf{p}_{y_2}(\{x_{S_1+1}\}) \cdots \mathbf{p}_{y_2}(\{x_{S_2}\})$$
$$\cdots \mathbf{p}_{y_k}(\{x_{S_{k-1}+1}\}) \cdots \mathbf{p}_{y_k}(\{x_{S_k}\}) \ge 2^{-n-S_k} = 2^{-(1+\frac{S_k}{S_{k+1}})n} \ge 2^{-(1+\frac{1}{k+1})n}.$$

Now we are ready to give the proof of Theorem 5.5.

Proof of Theorem 5.5. We begin by proving $\dim_H \mu = 1$. Note that $\dim_H \mu \ge 1$, as μ projects under $[0,1]^2 \ni (x,y) \mapsto y \in [0,1]$ to the Lebesgue measure, so it is sufficient to show dim_H $\mu \leq$ 1. By (5.8), it is enough to prove that $\underline{d}(\mu, (x, y)) \leq 1$ for μ -almost every $(x, y) \in [0, 1]$. Note that for Lebesgue almost every $y = 0.y_1y_2... \in [0,1]$, the sequence $(y_1, y_2,...)$ contains infinitely many zeros. Hence, it is sufficient to show $\underline{d}(\mu, (x, y)) \leq 1$ for μ_y -almost every $x \in [0, 1]$, assuming that $y \in [0, 1]$ has this property. Moreover, for μ_y -almost every $x \in [0, 1]$, we have $\mu(D_n(x, y)) > 0$ for all $n \in \mathbb{N}$ (see (5.9)). For such x, by Lemma 5.6(b), we have

$$\underline{d}(\mu, (x, y)) \le \liminf_{k \to \infty} \frac{-\log \mu(D_{S_{n_k}}(x, y))}{S_{n_k} \log 2} \le \lim_{k \to \infty} \frac{(1 + \frac{1}{n_k})S_{n_k}}{S_{n_k}} = 1$$

Therefore, $\dim_H \mu \leq 1$, so in fact $\dim_H \mu = 1$.

Let us prove now $\underline{\dim}_{MB} \mu = 2$. Since μ is supported on $[0, 1]^2$, it suffices to show $\underline{\dim}_{MB} \mu \geq$ 2. Let $A \subset [0,1]^2$ be a Borel set with $\mu(A) > 0$. We show $\underline{\dim}_B A \ge 2$. Note that there exists c > 0 such that the set

(5.10)
$$B = \{ y \in [0,1] : \mu_y(A^y) \ge c \}$$

satisfies $\operatorname{Leb}(B) > 0$. Fix $\varepsilon \in (0, \frac{1}{4})$. By the Lebesgue density theorem (see e.g. [Hoc14, Corollary 3.16]), there exists a dyadic interval $I \subset [0, 1]$ such that

(5.11)
$$\frac{\operatorname{Leb}(B \cap I)}{|I|} \ge 1 - \varepsilon,$$

where $|I| = 2^{-N}$ is the length of I. Fix $k \ge N+2$ and $n \in \{S_k + 1, \dots, S_{k+1}\}$. Consider the collection \mathcal{C}_n of dyadic intervals of length 2^{-n} defined as

$$C_n = \{ [y_1, \dots, y_n] : y_k = y_{k+1} = 1 \text{ and } [y_1, \dots, y_n] \cap B \cap I \neq \emptyset \}.$$

By (5.11), we have

(5.12)
$$\operatorname{Leb}\left(B \cap \bigcup \mathcal{C}_n\right) \ge \left(\frac{1}{4} - \varepsilon\right) 2^{-N}$$

Let

$$A_n = A \cap \left([0, 1] \times \left(B \cap \bigcup \mathcal{C}_n \right) \right).$$

Then $A_n \subset A$ and (5.10) together with (5.12) imply

(5.13)
$$\mu(A_n) = \int_{B \cap \bigcup \mathcal{C}_n} \mu_y(A^y) d\operatorname{Leb}(y) \ge c \left(\frac{1}{4} - \varepsilon\right) 2^{-N}.$$

Note that the above lower bound does not depend on k and n. Let $N'(A_n, 2^{-n})$ be the number of dyadic squares of sidelength 2^{-n} intersecting A_n . If $D = I_1 \times I_2$ is a dyadic square of sidelength 2^{-n} intersecting A_n , then $I_2 \in \mathcal{C}_n$, hence by Lemma 5.6(a) we have

$$\mu(D) \le 2^{-(2-\frac{1}{k})n}.$$

As any two dyadic squares of the same sidelength are either equal or disjoint, (5.13) gives

$$N'(A, 2^{-n}) \ge N'(A_n, 2^{-n}) \ge c \left(\frac{1}{4} - \varepsilon\right) 2^{-N + (2 - \frac{1}{k})n}$$

Since k and n can be taken arbitrary large, invoking (5.7) gives $\underline{\dim}_B A \ge 2$. Hence, $\underline{\dim}_{MB} \mu \ge 2$, so in fact $\underline{\dim}_{MB} \mu = 2$.

Remark 5.7. Note that as

$$\operatorname{ess\,sup}_{z\sim\mu} \underline{d}(\mu, z) = \dim_{H} \mu \leq \underline{\dim}_{MB} \mu \leq \overline{\dim}_{MB} \mu = \dim_{P} \mu = \operatorname{ess\,sup}_{z\sim\mu} \overline{d}(\mu, z)$$

(dim_P denotes the packing dimension, see e.g. [Fal14, Proposition 3.9] and [Fal97, Proposition 10.3]), the equality dim_H $\mu = \underline{\dim}_{MB} \mu$ holds for all *exact dimensional* measures μ , i.e. the measures μ with $\underline{d}(\mu, z) = \overline{d}(\mu, z) = \text{const}$ for μ -almost every z.

References

- [ABD⁺19] G. Alberti, H. Bölcskei, C. De Lellis, G. Koliander, and E. Riegler. Lossless analog compression. IEEE Transactions on Information Theory, 65(11):7480–7513, 2019.
- [BAEFN93] Asher Ben-Artzi, Alp Eden, Ciprian Foias, and Basil Nicolaenko. Hölder continuity for the inverse of Mañé's projection. J. Math. Anal. Appl., 178(1):22–29, 1993.
- [Ban51] Stefan Banach. Wstęp do teorii funkcji rzeczywistych (Polish) [Introduction to the theory of real functions]. Monografie Matematyczne. Tom XVII. Polskie Towarzystwo Matematyczne, Warszawa-Wrocław, 1951.
- [Bil99] Patrick Billingsley. Convergence of probability measures. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999. A Wiley-Interscience Publication.
- [Cab00] Victoria Caballero. On an embedding theorem. Acta Math. Hungar., 88(4):269–278, 2000.
- [EFNT94] Alp Eden, Ciprian Foias, Basil Nicolaenko, and Roger M. Temam. Exponential attractors for dissipative evolution equations, volume 37 of RAM: Research in Applied Mathematics. Masson, Paris; John Wiley & Sons, Ltd., Chichester, 1994.
- [ER85] Jean-Pierre Eckmann and David Ruelle. Ergodic theory of chaos and strange attractors. Rev. Modern Phys., 57(3, part 1):617–656, 1985.
- [Fal97] Kenneth Falconer. Techniques in fractal geometry. John Wiley & Sons, Ltd., Chichester, 1997.
- [Fal14] Kenneth Falconer. Fractal geometry. John Wiley & Sons, Ltd., Chichester, third edition, 2014. Mathematical foundations and applications.
- [FR02] Peter K. Friz and James C. Robinson. Constructing an elementary measure on a space of projections. J. Math. Anal. Appl., 267(2):714–725, 2002.
- [GQS18] Yonatan Gutman, Yixiao Qiao, and Gábor Szabó. The embedding problem in topological dynamics and Takens' theorem. *Nonlinearity*, 31(2):597–620, 2018.
- [GS00] Mariano Gasca and Thomas Sauer. Polynomial interpolation in several variables. Adv. Comput. Math., 12(4):377–410, 2000. Multivariate polynomial interpolation.
- [Gut16] Yonatan Gutman. Taken's embedding theorem with a continuous observable. In *Ergodic theory*, pages 134–141. De Gruyter, Berlin, 2016.
- [GVL13] Gene H. Golub and Charles F. Van Loan. *Matrix computations*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, fourth edition, 2013.
- [HBS15] Franz Hamilton, Tyrus Berry, and Timothy Sauer. Predicting chaotic time series with a partial model. Phys. Rev. E, 92:010902, Jul 2015.
- [HGLS05] Chih-Hao Hsieh, Sarah M. Glaser, Andrew J. Lucas, and George Sugihara. Distinguishing random environmental fluctuations from ecological catastrophes for the North Pacific Ocean. Nature, 435(7040):336–340, 2005.
- [HK99] Brian R. Hunt and Vadim Yu. Kaloshin. Regularity of embeddings of infinite-dimensional fractal sets into finite-dimensional spaces. *Nonlinearity*, 12(5):1263–1275, 1999.

- [Hoc14] Michael Hochman. Lectures on dynamics, fractal geometry, and metric number theory. J. Mod. Dyn., 8(3-4):437–497, 2014.
- [HW41] Witold Hurewicz and Henry Wallman. Dimension Theory. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941.
- [Kec95] Alexander S. Kechris. Classical descriptive set theory, volume 156 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995.
- [KY90] Eric J. Kostelich and James A. Yorke. Noise reduction: finding the simplest dynamical system consistent with the data. *Phys. D*, 41(2):183–196, 1990.
- [Mar54] John M. Marstrand. Some fundamental geometrical properties of plane sets of fractional dimensions. Proc. London Math. Soc. (3), 4:257–302, 1954.
- [Mat75] Pertti Mattila. Hausdorff dimension, orthogonal projections and intersections with planes. Ann. Acad. Sci. Fenn. Ser. A I Math., 1(2):227–244, 1975.
- [Mat95] Pertti Mattila. Geometry of sets and measures in Euclidean spaces, volume 44 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [MGNS18] Stephan B. Munch, Alfredo Giron-Nava, and George Sugihara. Nonlinear dynamics and noise in fisheries recruitment: A global meta-analysis. *Fish and Fisheries*, 19(6):964–973, 2018.
- [Min70] George J. Minty. On the extension of Lipschitz, Lipschitz-Hölder continuous, and monotone functions. *Bulletin of the American Mathematical Society*, 76(2):334–339, 1970.
- [Mn81] Ricardo Mañé. On the dimension of the compact invariant sets of certain nonlinear maps. In Dynamical systems and turbulence, Warwick 1980 (Coventry, 1979/1980), volume 898 of Lecture Notes in Math., pages 230–242. Springer, Berlin-New York, 1981.
- [NV18] Raymundo Navarrete and Divakar Viswanath. Prevalence of delay embeddings with a fixed observation function. *Preprint. https://arxiv.org/abs/1806.07529*, 2018.
- [PCFS80] Norman H. Packard, James P. Crutchfield, J. Doyne Farmer, and Robert S. Shaw. Geometry from a time series. *Phys. Rev. Lett.*, 45:712–716, Sep 1980.
- [PdM82] Jacob Palis, Jr. and Welington de Melo. Geometric theory of dynamical systems. Springer-Verlag, New York-Berlin, 1982. An introduction, Translated from the Portuguese by A. K. Manning.
- [Rob05] James C. Robinson. A topological delay embedding theorem for infinite-dimensional dynamical systems. Nonlinearity, 18(5):2135–2143, 2005.
- [Rob11] James C. Robinson. Dimensions, embeddings, and attractors, volume 186 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2011.
- [Rud87] Walter Rudin. Real and complex analysis. McGraw-Hill Book Co., New York, third edition, 1987.
- [SBDH97] Jaroslav Stark, David S. Broomhead, Michael Evan Davies, and Jeremy P. Huke. Takens embedding theorems for forced and stochastic systems. In *Proceedings of the Second World Congress* of Nonlinear Analysts, Part 8 (Athens, 1996), volume 30, pages 5303–5314, 1997.
- [SBDH03] Jaroslav Stark, David S. Broomhead, Michael Evan Davies, and Jeremy P. Huke. Delay embeddings for forced systems. II. Stochastic forcing. J. Nonlinear Sci., 13(6):519–577, 2003.
- [SGM90] George Sugihara, Bryan Grenfell, and Robert May. Distinguishing error from chaos in ecological time-series. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 330(1257):235–251, 1990.
- [SM90] George Sugihara and Robert May. Nonlinear forecasting as a way of distinguishing chaos from measurement error in time series. *Nature*, 344(6268):734–741, 1990.
- [Sta99] Jaroslav Stark. Delay embeddings for forced systems. I. Deterministic forcing. J. Nonlinear Sci., 9(3):255–332, 1999.
- [SY97] Timothy D. Sauer and James A. Yorke. Are the dimensions of a set and its image equal under typical smooth functions? *Ergodic Theory Dynam. Systems*, 17(4):941–956, 1997.
- [SYC91] Timothy D. Sauer, James A. Yorke, and Martin Casdagli. Embedology. J. Statist. Phys., 65(3-4):579–616, 1991.

[Tak81]	Floris Takens. Detecting strange attractors in turbulence. In Dynamical systems and turbulence,
	Warwick 1980, volume 898 of Lecture Notes in Math., pages 366-381. Springer, Berlin-New
	York, 1981.

- [Vos03] Henning U. Voss. Synchronization of reconstructed dynamical systems. *Chaos*, 13(1):327–334, 2003.
- [Whi36] Hassler Whitney. Differentiable manifolds. Ann. of Math. (2), 37(3):645–680, 1936.
- [Yor69] James A. Yorke. Periods of periodic solutions and the Lipschitz constant. Proc. Amer. Math. Soc., 22:509–512, 1969.