

On conjugates and the asymptotic distortion of 1-dimensional C^{1+bv} diffeomorphisms

Andrés Navas

Abstract: We show that a C^{1+bv} circle diffeomorphism with absolutely continuous derivative and irrational rotation number can be conjugated into diffeomorphisms that are C^{1+bv} arbitrary close of the corresponding rotation. This improves a theorem of M. Herman, who established the same result but starting with a C^2 diffeomorphism. We prove that the same holds for Abelian groups of diffeomorphisms acting freely on the circle, a result which is new even in the C^∞ context. Related results and examples concerning the asymptotic distortion of diffeomorphisms are presented.

Keywords: Herman's theorem, invariant distribution, conjugacy.

Mathematical Subject Classification: 37E05, 37E10, 37E45.

In memory of Jean-Christophe Yoccoz

Since the seminal work of Poincaré and Denjoy, circle diffeomorphisms became a central object in dynamics. One of the most celebrated results of the theory is the linearization theorem of M. Herman [6], which establishes the differentiability of the conjugating map to the corresponding rotation in the case of an irrational rotation number satisfying a Diophantine condition. The key idea of Herman's proof consists in conjugating the diffeomorphism into one that is arbitrarily close to the rotation in a good topology, and finally applying local linearization results coming from standard KAM techniques (see also [5]).

Herman's theorem was later extended via different approaches by many people, including Yoccoz [18], Katznelson and Orstein [10], and Khanin and Teplinsky [9]. However, the problem of approximation by conjugation raised by Herman in Chapter VII of his seminal work [6] is interesting by itself. In this regard, just a couple of results that apply to diffeomorphisms of arbitrary irrational rotation number are known. On the one hand, it was shown by the author in [14] that every C^1 circle diffeomorphism f of irrational rotation number $\rho = \rho(f)$ has a sequence of conjugates that converges in the C^1 topology to the rotation R_ρ ; actually, there is a continuous path of C^1 conjugates that starts at f and finishes at R_ρ . On the other hand, Herman proved in [6] that if f is of class C^2 , then there exists a sequence of conjugates of f that converges to R_ρ in the C^{1+bv} sense; that is, they converge to f in the uniform topology, their derivatives uniformly converge to 1, and the total variation of the logarithm of these derivatives converge to 0. The first goal of this paper is to relax the hypothesis of regularity of f for this result.

Theorem A. *Given a C^{1+bv} (orientation-preserving) circle diffeomorphism with absolutely continuous derivative and having an irrational rotation number, there exists a continuous path of C^{1+bv} diffeomorphisms g_t , $t \in [0, 1[$, that starts at $g_0 = id$ such that the conjugates $g_t f g_t^{-1}$ converge to the rotation of angle $\rho = \rho(f)$ in the C^{1+bv} sense as t goes to 1.*

The method of proof of this result is somewhat similar to that of [14]; namely, we look for a reduction of the problem of approximation by conjugates into a problem of finding approximate solutions to a certain cohomological equation. In the C^1 case, this is relatively easy, since the associated equation concerns the logarithmic derivative, and it is hence linear. In the present setting, however, the equation that arises deals with the affine derivative, and it is hence twisted by the action of the (classical) derivative. A key role is then played by a (slight variation and simplification of a) result established by the author and Triestino in [15], which establishes the existence of approximate solutions to these twisted cohomological equations in the most general C^{1+bv} setting. (The main result of [15] is stated in terms of invariant 1-distributions, but it is equivalent to the previous formulation via a standard Hahn-Banach type argument.)

It is worth comparing this method of proof with that of the weaker statement of Herman for C^2 diffeomorphisms. On the one hand, Herman shows his theorem of approximation by conjugates via quite intricate estimates for certain ergodic averages which, as a byproduct, yield the ergodicity of the diffeomorphism with respect to the Lebesgue measure. (He needs the C^2 –or, at least, a C^{1+Lip} – hypothesis for making this procedure working.) On the other hand, as we already mentioned, our approach morally uses the uniqueness of the invariant 1-distribution, which essentially follows from the uniqueness of the 1-conformal measure established by Douady and Yoccoz in [4] and which, in its turn, strongly relies on Katok’s geometric proof of the ergodicity of C^{1+bv} circle diffeomorphisms with irrational rotation number (see [13]). In summary, in both approaches, a key role is played by the ergodicity of the diffeomorphism, which is essentially a one-dimensional phenomenon. (Compare [2].)

Although our approach still applies in the C^2 setting, a key technical point (concerning the “size” of the conjugating map) at the end of the proof fails, so that the question of whether conjugates of C^2 circle diffeomorphisms of arbitrary irrational rotation number can approximate the corresponding rotation in the C^2 topology remains open. (It should be pointed out that A. Avila and R. Krikorian have recently announced a result in this direction but for C^∞ conjugates of C^∞ diffeomorphisms.) On the positive side, since our approach uses the affine derivative as a cocycle in L^1 , it is suitable for a generalization to Abelian group actions, which have deserved much attention over the last years (see for instance [3, 7, 11]). The result in this context, which is new in the C^∞ setting (and even in the real-analytic context), reads as follows.

Theorem B. *Let Γ be a finitely-generated, Abelian group of C^{1+bv} (orientation-preserving) circle diffeomorphisms with absolute continuous derivative. If the action of Γ is free, then there exists a path g_t , $t \in [0, 1[$, of C^{1+bv} diffeomorphisms starting at $g_0 = id$ such that, for each $f \in \Gamma$, the conjugates $g_t f g_t^{-1}$ converge to the rotation $R_{\rho(f)}$ in the C^{1+bv} sense as t goes to 1.*

We close this work with a discussion on the notion of asymptotic distortion of diffeomorphisms of one-dimensional manifolds, which corresponds to the linear growth of the total variation of the logarithm of the derivative:

$$\text{dist}_\infty(f) := \lim_{n \rightarrow \infty} \frac{\text{var}(\log(Df^n))}{n}.$$

We show that the vanishing of this number is equivalent to that the diffeomorphism f can be conjugated into maps that are arbitrarily close to isometries in the C^{1+bv} sense. Some results and examples are exhibited: it is proved that, in $\text{PSL}(2, \mathbb{R})$, only the hyperbolic elements have positive asymptotic distortion, though there exist C^∞ diffeomorphisms of the interval with positive asymptotic distortion but without hyperbolic fixed points. Actually, this is the typical behavior of diffeomorphisms with no hyperbolic fixed points.

Acknowledgments. I'm strongly indebted to Adolfo Guillot, with whom formula (5) arose in a discussion seminar (more than ten years ago). I'm also indebted to Jairo Bochi and Alejandro Kocsard for quite stimulating discussions. The preparation of this text was funded by the CONICYT Project MathAmsud 17-03 "GDAR: geometry, dynamics, and Anosov representations" as well as the projects FONDECYT 1160541 and FORDECYT 265667.

On C^{1+bv} conjugates of circle diffeomorphisms with irrational rotation number

All maps considered in this article will be assumed to be orientation preserving. Before passing to the proof of Theorem A, we will give a variation of one of the main results of [15] (which deals with continuous functions) to an L^1 setting, so that it becomes more suitable for our needs. As the reader will notice, this version is much simpler in that it does not require Douady-Yoccoz' result of uniqueness of the 1-conformal measure, but just Katok-Herman's theorem of ergodicity (with respect to the Lebesgue measure) of C^{1+bv} circle diffeomorphisms with irrational rotation number [13]. To strength this point, and also for the reader's convenience, we provide a detailed proof. (It is not hard to see that an alternative but more direct proof can be given using the stronger result of [15].)

Proposition 1. *Let f be a C^{1+bv} circle diffeomorphism of irrational rotation number. If φ is an L^1 function defined on the circle that has zero mean, then there exists a sequence of L^1 functions ψ_n of zero mean such that the sequence*

$$\psi_n - \psi_n \circ f \cdot Df$$

converges to φ in the L^1 sense as n goes to infinity.

Proof. Assume first that φ cannot be approximated in L^1 by functions of the form

$$\psi - \psi \circ f \cdot Df,$$

where no condition on the mean of $\psi \in L^1$ is imposed. Then, by the Hahn-Banach theorem, there exists a linear functional $L: L^1(S^1) \rightarrow \mathbb{R}$ that vanishes on the set of functions of the form $\psi - \psi \circ f \cdot Df$ and such that $L(\varphi) = 1$. This functional comes from integration against an L^∞ function ξ :

$$L(\phi) = \int_{S^1} \xi \cdot \phi.$$

Hence, the equality $L(\psi - \psi \circ f \cdot Df) = 0$ becomes

$$\int_{S^1} \xi \cdot \psi = \int_{S^1} \xi \cdot \psi \circ f \cdot Df = \int_{S^1} \xi \circ f^{-1} \cdot \psi,$$

where the last equality comes from change of variable. Since this holds for every $\psi \in L^1$, this implies that a.e. we have the equality $\xi = \xi \circ f^{-1}$. By the Katok-Herman's ergodicity theorem, the function ξ is a.e. constant, say equal to c . However, this implies that

$$L(\varphi) = c \int_{S^1} \varphi = 0,$$

which is a contradiction.

Let hence $\tilde{\psi}_n$ be a sequence of L^1 functions such that the sequence

$$\tilde{\psi}_n - \tilde{\psi}_n \circ f \cdot Df$$

converges to φ in L^1 . For each n let

$$c_n := \int \tilde{\psi}_n.$$

Then the function $\tilde{\psi}_n - c_n$ has zero mean, and satisfies

$$(\tilde{\psi}_n - c_n) - (\tilde{\psi}_n - c_n) \circ f \cdot Df + c_n(1 - Df) = \tilde{\psi}_n - \tilde{\psi}_n \circ f \cdot Df \longrightarrow \varphi.$$

To deal with the unpleasant expression $c_n(1 - Df)$, let

$$\eta_k := \frac{1}{q_k} [1 + Df + Df^2 + \dots + Df^{q_k-1}] - 1,$$

where (q_k) is the sequence of denominators in the rational approximation of the rotation number of f . Notice that, for all $k \geq 1$,

$$\int_{S^1} \eta_k = 0.$$

Moreover, an easy computation yields

$$1 - Df = (\eta_k - \eta_k \circ f \cdot Df) - \frac{1}{q_k} [Df^{q_k} - 1].$$

By the Denjoy's inequality [13], the last expression on the right uniformly converges to zero, hence

$$\eta_k - \eta_k \circ f \cdot Df$$

uniformly converges to $1 - Df$. The desired sequence ψ_n is then obtained by letting

$$\psi_n = \tilde{\psi}_n - c_n + c_n \eta_{k_n}$$

for a well-chosen sequence k_n . □

Proof of Theorem A. For a diffeomorphism f , belonging to the class C^{1+bv} with an absolutely continuous derivative is equivalent to being C^1 and having a second derivative a.e. defined so that the function D^2f/Df is integrable. The total variation of the logarithm of the derivative of f then equals

$$\text{var}(\log(Df); S^1) = \int_{S^1} \left| \frac{D^2f}{Df} \right| = \left\| \frac{D^2f}{Df} \right\|_{L^1}.$$

Notice that $D \log(Df) = D^2f/Df$. Therefore, if f is a C^{1+bv} diffeomorphism with an absolutely continuous derivative, then

$$\int_{S^1} \frac{D^2f}{Df} = 0.$$

By Proposition 1 above, there exists a sequence of L^1 functions ψ_n , all of which have zero mean, such that the sequence

$$\psi_n - (\psi_n \circ f) \cdot Df$$

converges to D^2f/Df in the L^1 topology. Let h_n be defined by

$$h_n(x) := \frac{\int_0^x \exp\left(\int_0^s \psi_n(r) dr\right) ds}{\int_0^1 \exp\left(\int_0^s \psi_n(r) dr\right) ds}.$$

Since ψ_n has zero mean, this map h_n is a C^1 circle diffeomorphism. (The condition on the mean translates into that $Dh_n(0) = Dh_n(1)$.) Besides, in the L^1 sense, we have the equality

$$\frac{D^2h_n}{Dh_n} = D(\log(Dh_n)) = \psi_n.$$

Since the cocycle relation

$$\frac{D^2(g_1g_2)}{D(g_1g_2)} = \frac{D^2g_1}{Dg_1} \circ g_2 \cdot Dg_2 + \frac{D^2g_2}{Dg_2} \tag{1}$$

implies

$$\frac{D^2(h_n f h_n^{-1})}{D(h_n f h_n^{-1})} = \frac{D^2h_n}{Dh_n} \circ (f h_n^{-1}) \cdot D(f h_n^{-1}) + \frac{D^2f}{Df} \circ h_n^{-1} \cdot Dh_n^{-1} - \frac{D^2h_n}{Dh_n} \circ h_n^{-1} \cdot Dh_n^{-1},$$

we have

$$\frac{D^2(h_n f h_n^{-1})}{D(h_n f h_n^{-1})} = \left[\psi_n \circ f \cdot Df + \frac{D^2 f}{Df} - \psi_n \right] \circ h_n^{-1} \cdot Dh_n^{-1}.$$

Besides, given $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that, for all $n \geq N$, we have

$$\left\| \psi_n \circ f \cdot Df + \frac{D^2 f}{Df} - \psi_n \right\|_{L^1} \leq \varepsilon.$$

As a consequence, by change of variable,

$$\left\| \left[\psi_n \circ f \cdot Df + \frac{D^2 f}{Df} - \psi_n \right] \circ h_n^{-1} \cdot Dh_n^{-1} \right\|_{L^1} \leq \varepsilon.$$

Therefore,

$$\text{var}(\log(D(h_n f h_n^{-1}))) = \left\| \frac{D^2(h_n f h_n^{-1})}{D(h_n f h_n^{-1})} \right\|_{L^1} \leq \varepsilon,$$

which implies that $h_n f h_n^{-1}$ converges to the rotation of angle $\rho(f)$ in the C^{1+bv} sense.

Finally, to build the desired path of conjugating maps g_t , just interpolate the functions ψ_n above. More precisely, let $u : [0, 1[\rightarrow [0, \infty[$ be a homeomorphism. For $t \in [0, 1[$, let n be the integer such that $n \leq u(t) < n + 1$, and define

$$\psi^t := (n + 1 - u(t)) \psi_n + (u(t) - n) \psi_{n+1},$$

where $\psi_0 \equiv 0$. Finally, define the diffeomorphism g_t by letting

$$g_t(x) := \frac{\int_0^x \exp\left(\int_0^s \psi^t(r) dr\right) ds}{\int_0^1 \exp\left(\int_0^s \psi^t(r) dr\right) ds}.$$

We leave to the reader the task of checking that the sequence of conjugates $g_t f g_t^{-1}$ satisfies the desired properties. \square

In order to treat the case of Abelian groups of diffeomorphisms, it is worth coming back to the proof of the proposition above and provide explicit approximating coboundaries.

Lemma 1. *If f is a C^{1+bv} circle diffeomorphism with absolutely continuous derivative and irrational rotation number, then the sequence*

$$\frac{1}{n} \left\| \frac{D^2 f^n}{Df^n} \right\|_{L^1} = \frac{\text{var}(\log(Df^n); S^1)}{n}$$

converges to 0 as n goes to infinite.

Proof. Denote by $U : L^1 \rightarrow L^1$ the linear isometry given by $U(\varphi) = \varphi \circ f \cdot Df$. Then the cocycle relation (1) yields

$$\frac{D^2 f^n}{Df^n} = \sum_{i=0}^{n-1} U^i \left(\frac{D^2 f}{Df} \right). \quad (2)$$

By Proposition 1, given $\varepsilon > 0$, there exists an L^1 function ψ such that

$$\left\| \frac{D^2 f}{Df} - (\psi - \psi \circ f \cdot Df) \right\|_{L^1} \leq \varepsilon.$$

Since U is an isometry, this implies that, for all $i \geq 0$,

$$\left\| U^i \left(\frac{D^2 f}{Df} \right) - (\psi \circ f^i \cdot Df^i - \psi \circ f^{i+1} \cdot Df^{i+1}) \right\|_{L^1} = \left\| U^i \left(\frac{D^2 f}{Df} - (\psi - \psi \circ f \cdot Df) \right) \right\|_{L^1} \leq \varepsilon.$$

Using the triangular inequality together with (2), this gives

$$\left\| \frac{D^2 f^n}{Df^n} - (\psi - \psi \circ f^n \cdot Df^n) \right\|_{L^1} \leq \sum_{i=0}^{n-1} \left\| U^i \left(\frac{D^2 f}{Df} \right) - (\psi \circ f^i \cdot Df^i - \psi \circ f^{i+1} \cdot Df^{i+1}) \right\|_{L^1} \leq n\varepsilon.$$

Therefore,

$$\frac{1}{n} \left\| \frac{D^2 f^n}{Df^n} \right\|_{L^1} \leq \varepsilon + \frac{1}{n} \left(\|\psi\|_{L^1} + \|(\psi \circ f^n) \cdot Df^n\|_{L^1} \right) = \varepsilon + \frac{2\|\psi\|_{L^1}}{n}.$$

Since this holds for every $\varepsilon > 0$, we deduce that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \left\| \frac{D^2 f^n}{Df^n} \right\|_{L^1} = 0$$

Finally, the cocycle relation (1) easily implies that the sequence $\|D^2 f^n / Df^n\|_{L^1}$ is subadditive, which shows the desired convergence. \square

Remark 1. In the proof above, the crucial property for $D^2 f / Df$ is that it has zero mean. It is hence not hard to check along the very same lines that for every $\varphi \in L^1$ having zero mean, one has the convergence

$$\frac{1}{n} \left\| \varphi + \varphi(f) \cdot Df + \varphi(f^2) \cdot Df^2 + \dots + \varphi(f^{n-1}) \cdot Df^{n-1} \right\|_{L^1} \longrightarrow 0.$$

Remark 2. It would be worth knowing whether the previous result still holds without the hypothesis of absolute continuity for the derivative. For piecewise affine homeomorphisms this is the case whenever the map belongs to an S^1 (faithful) action by piecewise affine homeomorphisms, as it follows from Minakawa's classification theorem [12]. Indeed, if the action is not conjugate to a rotation by a piecewise-affine homeomorphism, then it is conjugate by such a map together with a homeomorphism of the form $x \mapsto \frac{\sigma^x - 1}{\sigma - 1}$, which is piecewise real-analytic with a derivative of finite total variation. (In particular, the sequence $\text{var}(\log(Df^n); S^1)$ is bounded.)

Following the lines of [6, Chapter IV], it is not hard to prove that, for a C^{1+bv} circle diffeomorphism (independently of the rotation number), the sequence $\text{var}(\log(Df^n); S^1)$ is

bounded if and only if f is conjugated to the corresponding isometry (the identity, in the case of the interval) by a C^{1+bv} diffeomorphism. The next question is inspired by [17].

Question 1. In case of unboundedness, does there exist a constraint for the growth of the sequence $\text{var}(\log(Df^n); S^1)$ that is stronger than sublinearity?

The next lemma is stated in a quite general and conceptual framework for potential uses in different contexts. For our purposes, it will be used for the isometry $U: \varphi \mapsto (\varphi \circ f) \cdot Df$ on L^1 introduced above and the associated cocycle

$$c(f) = \frac{D^2 f}{Df}.$$

Lemma 2. *Let U be a linear isometric action of a finitely-generated Abelian group Γ on a Banach space \mathbb{B} , and let $c: \Gamma \rightarrow \mathbb{B}$ be a cocycle, that is, a function that satisfies the relation*

$$c(g_1 g_2) = c(g_2) + U(g_2)(c(g_1)).$$

Assume that, for every element g of infinite order in Γ , one has

$$\frac{\|c(g^n)\|_{\mathbb{B}}}{n} \rightarrow 0$$

as $n \rightarrow \infty$. Then there exists a sequence of functions $\psi_n \in \mathbb{B}$ such that, for all $f \in \Gamma$, the sequence $\psi_n - U(f)(\psi_n)$ converges to $c(f)$ as n goes to infinite.

Proof. Let us first assume that Γ has no torsion, say $\Gamma \sim \mathbb{Z}^k$. Let $\{g_1, \dots, g_k\}$ be a (minimal) system of generators of Γ . For each $n \in \mathbb{N}$, let us denote by $B(n)$ be the *positive square ball* of radius n , that is,

$$B(n) = \{g_1^{n_1} g_2^{n_2} \cdots g_k^{n_k} : 0 \leq n_i \leq n\}.$$

For each $n \geq 1$, define

$$\psi_n := \frac{1}{|B(n-1)|} \sum_{g \in B(n-1)} c(g). \tag{3}$$

Then, for each $f \in \Gamma$,

$$\begin{aligned} U(f)(\psi_n) &= \frac{1}{|B(n-1)|} \sum_{g \in B(n-1)} U(f)(c(g)) \\ &= \frac{1}{|B(n-1)|} \sum_{g \in B(n-1)} [c(gf) - c(f)] \\ &= -c(f) + \frac{1}{|B(n-1)|} \sum_{g \in B(n-1)} c(gf) \\ &= -c(f) + \frac{1}{|B(n-1)|} \sum_{g \in B(n-1)} c(fg). \end{aligned}$$

Therefore, for each $i \in \{1, 2, \dots, k\}$,

$$\left\| c(g_i) - (\psi_n - U(g_i)(\psi_n)) \right\|_{\mathbb{B}} \leq \frac{1}{B(n-1)} \left\| \sum_{g \in B(n-1)} [c(g_i g) - c(g)] \right\|_{\mathbb{B}},$$

and the last expression equals

$$\frac{1}{B(n-1)} \left\| \sum_{\substack{0 \leq n_j \leq n-1 \\ j \neq i}} [c(g_i^{n_1} g_1^{n_1} \cdots g_{i-1}^{n_{i-1}} g_{i+1}^{n_{i+1}} \cdots g_k^{n_k}) - c(g_1^{n_1} \cdots g_{i-1}^{n_{i-1}} g_{i+1}^{n_{i+1}} \cdots g_k^{n_k})] \right\|_{\mathbb{B}}.$$

By the cocycle relation, this reduces to

$$\frac{1}{B(n-1)} \left\| \sum_{\substack{0 \leq n_j \leq n-1 \\ j \neq i}} U(g_1^{n_1} \cdots g_{i-1}^{n_{i-1}} g_{i+1}^{n_{i+1}} \cdots g_k^{n_k})(c(g_i^{n_j})) \right\|_{\mathbb{B}},$$

which, by the triangular inequality, is smaller than or equal to

$$\frac{1}{B(n-1)} \sum_{\substack{0 \leq n_j \leq n-1 \\ j \neq i}} \|U(g_1^{n_1} \cdots g_{i-1}^{n_{i-1}} g_{i+1}^{n_{i+1}} \cdots g_k^{n_k})(c(g_i^{n_j}))\|_{\mathbb{B}} = \frac{1}{B(n-1)} \sum_{\substack{0 \leq n_j \leq n-1 \\ j \neq i}} \|c(g_i^{n_j})\|_{\mathbb{B}}.$$

A simple counting argument yields the following upper bound for the last expression:

$$\frac{1}{n^k} n^{k-1} \|c(g_i^{n_j})\|_{\mathbb{B}} = \frac{\|c(g_i^{n_j})\|_{\mathbb{B}}}{n},$$

According to the hypothesis, this goes to zero as $n \rightarrow \infty$. Therefore,

$$\|c(g_i) - (\psi_n - U(g_i)(\psi_n))\|_{\mathbb{B}}$$

converges to zero as $n \rightarrow \infty$. To show that this holds changing g_i by any $f \in \Gamma$, just write f as a product of generators, apply the previous conclusion to each of these generators, and use the triangular inequality.

Finally, assume that Γ has torsion, say $\Gamma \sim \mathbb{Z}^k \times T$, where T is a finite group. Let ψ_n^* be a sequence for which $\|c(f) - (\psi_n^* - U(f)(\psi_n^*))\|_{\mathbb{B}}$ converges to zero for every $f \in \mathbb{Z}^k$. Replace ψ_n^* by

$$\psi_n := \frac{1}{|T|} \sum_{g \in T} [U(g)(\psi_n^*) + c(g)]. \quad (4)$$

For each $f \in \Gamma$, we have

$$\begin{aligned} U(f)\psi_n &= \frac{1}{|T|} \sum_{g \in T} [U(f)U(g)(\psi_n^*) + U(f)c(g)] \\ &= \frac{1}{|T|} \sum_{g \in T} [U(f)U(g)(\psi_n^*) + c(gf) - c(f)] \\ &= -c(f) + \frac{1}{|T|} \sum_{g \in T} [U(gf)(\psi_n^*) + c(gf)]. \end{aligned}$$

If f belongs to T , then the last expression reduces to $-c(f) + \psi_n$, thus showing that we actually have an equality:

$$c(f) = \psi_n - U(f)(\psi_n).$$

Assume otherwise that f belongs to \mathbb{Z}^k . Then

$$U(f)\psi_n = -c(f) + \frac{1}{|T|} \sum_{g \in T} [U(gf)(\psi_n^*) + c(gf)] = -c(f) + \psi_n + \frac{1}{|T|} \sum_{g \in T} [U(gf)(\psi_n^*) + c(gf)] - \psi_n,$$

hence the value of

$$\|c(f) - (\psi_n - U(f)\psi_n)\|_{\mathbb{B}}$$

is bounded from above by

$$\frac{1}{|T|} \left\| \sum_{g \in T} [U(gf)(\psi_n^*) + c(gf)] - [U(g)(\psi_n^*) + c(g)] \right\|_{\mathbb{B}} = \frac{1}{|T|} \left\| \sum_{g \in T} U(g)[U(f)(\psi_n^*) - \psi_n^*] + [c(fg) - c(g)] \right\|_{\mathbb{B}},$$

that is, by

$$\begin{aligned} \frac{1}{|T|} \left\| \sum_{g \in T} U(g)[U(f)(\psi_n^*) - \psi_n^*] + U(g)(c(f)) \right\|_{\mathbb{B}} &= \frac{1}{|T|} \left\| \sum_{g \in T} U(g)[U(f)(\psi_n^*) - \psi_n^* + c(f)] \right\|_{\mathbb{B}} \\ &= \left\| U(f)(\psi_n^*) - \psi_n^* + c(f) \right\|_{\mathbb{B}}, \end{aligned}$$

which, by construction, converges to zero as $n \rightarrow \infty$. Finally, to deal with an arbitrary $f \in \Gamma$, just write $f = h_1 h_2$, with $h_1 \in \mathbb{Z}^k$ and $h_2 \in T$, and use the cocycle relation and the triangular inequality to show the desired convergence for the function ψ_n . \square

Remark 3. The proof above mimics one of the key arguments of [14]. (Compare [1].) Actually, some of the ideas therein allow extending it to nilpotent group actions.

Proof of Theorem B. Let us again consider the isometries $U : \varphi \mapsto (\varphi \circ f) \cdot Df$ defined on the space L_0^1 made of the L^1 functions with zero mean, as well as the associated cocycle

$$c(f) = \frac{D^2 f}{Df} \in L_0^1.$$

Lemma 1 establishes the hypothesis of Lemma 2, which yields a sequence of functions $\psi_n \in L_0^1$ such that, for all $f \in \Gamma$, the value of

$$\|c(f) - (\psi_n - \psi_n \circ f \cdot Df)\|_{L^1}$$

converges to zero as $n \rightarrow \infty$. Therefore, as in the proof of Theorem A, this function ψ_n can be seen as the affine derivative of a circle diffeomorphism h_n (that is, $D^2 h_n / Dh_n = \psi_n$) such that, for all $f \in \Gamma$, the conjugates $h_n f h_n^{-1}$ converge to the rotation $R_{\rho(f)}$ in the C^{1+bv} sense. To produce the path of approximating conjugates, it suffices to linearly interpolate the functions ψ_n , as it was done at the end of the proof of Theorem A. We leave the verification of the details as a task to the reader. \square

Remark 4. Following the lines of the proofs above, the conjugating maps become explicit, even for the case of single maps (*i.e.* for actions of $\mathbb{Z} = \langle f \rangle$). Indeed, in this setting, definition (3) becomes

$$\psi_n = \frac{1}{n} \sum_{k=0}^{n-1} \frac{D^2 f^k}{Df^k},$$

and the associated conjugating maps h_n are given by

$$h_n(x) = \frac{\int_0^x \exp\left(\int_0^s \psi_n(r) dr\right) ds}{\int_0^1 \exp\left(\int_0^s \psi_n(r) dr\right) ds}.$$

A simple computation then shows that

$$Dh_n = \frac{1}{\int_0^1 \left(\prod_{k=0}^{n-1} \frac{Df^k(s)}{Df^k(0)}\right)^{1/n} ds} \left(\prod_{k=0}^{n-1} \frac{Df^k(x)}{Df^k(0)}\right)^{1/n}.$$

Notice that these conjugating maps are quite different from those used by Herman, which are given by

$$h_n^*(x) := \frac{1}{n} \sum_{k=0}^{n-1} (f^k - k\rho(f)),$$

where, on the right, f denotes a lift to the real-line of the original map and $\rho(f)$ is its corresponding translation number. It is worth mentioning that certain analogs of Herman's conjugating maps h_n^* still work for Abelian (and even nilpotent) group actions, yet providing explicit estimates for the derivatives of the conjugate action in the smooth case seems out of reach. (Herman managed to control derivatives for \mathbb{Z} -actions.)

On the asymptotic distortion of diffeomorphisms

Given a C^{1+bv} diffeomorphism of a compact one-dimensional manifold M , let us define its asymptotic distortion as

$$\text{dist}_\infty(f) := \lim_{n \rightarrow \infty} \frac{\text{var}(\log(Df^n); M)}{n}.$$

Notice that the existence of the limit follows from subadditivity:

$$\text{var}(\log(Df^{m+n}); M) \leq \text{var}(\log(Df^m); M) + \text{var}(\log(Df^n); M).$$

Besides, the subadditivity relation

$$\text{var}(\log(D(fg)); M) \leq \text{var}(\log(Df); M) + \text{var}(\log(Dg); M)$$

easily implies that dist_∞ is invariant under conjugacy.

It should become evident from the previous paragraph that the vanishing of the asymptotic distortion is closely related to approximating (in the C^{1+bv} sense) an isometry by a sequence of conjugates. We explicitly record this fact in the statement below. Notice that, in one of the implications, no assumption on absolute continuity for the derivative is made.

Proposition 3. Let f be a C^{1+bv} diffeomorphism of a compact one-dimensional manifold M . If there exists a sequence h_n of C^{1+bv} diffeomorphisms such that the conjugates $h_n f h_n^{-1}$ converge to an isometry in the C^{1+bv} sense, then the asymptotic distortion of f vanishes. Conversely, if f has zero asymptotic distortion and its derivative is absolutely continuous, then there exists a sequence of conjugates of f converging to an isometry in the C^{1+bv} sense.

Proof. If $h_n f h_n^{-1}$ converges to an isometry then, for each $\varepsilon > 0$, there exists an integer N such that, for all $n \geq N$,

$$\text{var}(\log(D(h_n f h_n^{-1})); M) < \varepsilon.$$

This yields, for every integer k ,

$$\text{var}(\log(D(h_n f^k h_n^{-1})); M) = \text{var}(\log(D((h_n f h_n^{-1})^k)); M) \leq k \text{var}(\log(D(h_n f h_n^{-1})); M) < k \varepsilon.$$

Therefore,

$$\text{var}(\log(Df^k); M) \leq \text{var}(\log(D(h_n f^k h_n^{-1})); M) + 2 \text{var}(\log(Dh_n); M) < k \varepsilon + 2 \text{var}(\log(Dh_n); M),$$

hence

$$\frac{\text{var}(\log(Df^k); M)}{k} < \varepsilon + \frac{2 \text{var}(\log(Dh_n); M)}{k}.$$

Letting $k \rightarrow \infty$, we obtain

$$\lim_{k \rightarrow \infty} \frac{\text{var}(\log(Df^k); M)}{k} \leq \varepsilon.$$

Since this holds for all $\varepsilon > 0$, the asymptotic distortion of f vanishes.

The converse statement is proved following the lines of the proof of Theorem B. Namely, if f is a C^{1+bv} diffeomorphism with absolute continuous derivative, the vanishing of the asymptotic distortion is equivalent to the vanishing of the linear growth of the L^1 -cocycle $D^2 f^n / Df^n$ in the space of functions of zero mean. (Notice that, for the case of the interval, the vanishing of the asymptotic distortion implies that Df equals 1 at the endpoints, hence $D^2 f / Df$ has zero mean.) By Lemma 2, this implies the existence of a sequence of zero-mean functions ψ_n such that the sequence

$$\psi_n - \psi_n \circ f \cdot Df$$

converges to $D^2 f / Df$ as $n \rightarrow \infty$. Writing the function ψ_n as $D^2 h_n / Dh_n$ for a C^{1+bv} diffeomorphism h_n provides the desired sequence of conjugating maps. This has been explicitly done for the case of the circle, and the very same formula provided for that case works in the framework of diffeomorphisms of $[0, 1]$. \square

We have showed that the asymptotic distortion of a C^{1+bv} circle diffeomorphism with absolute continuous derivative and irrational rotation number is equal to zero. In case of

rational rotation number, this is no longer true, since hyperbolic periodic points obviously force positive asymptotic distortion. Actually, as we next see, hyperbolicity is the only obstruction to zero asymptotic distortion for maps in $\mathrm{PSL}(2, \mathbb{R})$, yet it is not the only one in general.

Proposition 4. *A Möbius transformation has positive asymptotic distortion if and only if it is hyperbolic.*

Proof. There are two ways to prove this. First, since hyperbolic Möbius transformations have hyperbolic fixed points and, hence, positive asymptotic distortion, we need to check that both elliptic and parabolic Möbius transformations have zero asymptotic distortion. This can be showed using Proposition 3. Namely, elliptic Möbius transformations are conjugate to Euclidean rotations, and parabolic Möbius maps admit sequences of conjugates that converge to the identity, as it readily follows by letting $\lambda \rightarrow 0$ in the identity

$$\begin{bmatrix} \lambda & 0 \\ 0 & 1/\lambda \end{bmatrix} \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \lambda & 0 \\ 0 & 1/\lambda \end{bmatrix}^{-1} = \begin{bmatrix} 1 & \lambda^2 t \\ 0 & 1 \end{bmatrix}.$$

The starting observation for the second proof is the following: consider a Möbius transformation $f \in \mathrm{PSL}(2, \mathbb{R})$ both as an isometry of the Poincaré disk (endowed with the hyperbolic distance dist_h) and as a circle diffeomorphism. Then the following equality holds:

$$\mathrm{var}(\log(Df); S^1) = 4 \mathrm{dist}_h(f(0), 0). \quad (5)$$

Assuming this equality, we conclude that the rate of $\mathrm{var}(\log(Df^n); S^1)$ is linear if and only if the same happens for $\mathrm{dist}_h(f^n(0), 0)$, and it is very well known that the behavior of the last sequence classifies Möbius maps: this sequence is bounded, unbounded but sublinear, or linear, in the elliptic, parabolic, or hyperbolic case, respectively.

We are hence left to prove (5), which is just a computation. Namely, if we write f in complex form as

$$f(e^{i\theta}) = e^{i\alpha} \cdot \frac{e^{i\theta} - a}{1 - \bar{a}e^{i\theta}}, \quad a = re^{i\xi}, \quad |r| < 1,$$

then, as a circle diffeomorphism, $df/d\theta$ is the norm of the derivative of this expression, which equals

$$\left\| \frac{e^{i\alpha} [ie^{i\theta}(1 - re^{i(\theta-\xi)}) + (e^{i\theta} - re^{i\alpha})rie^{i(\theta-\alpha)}]}{(1 - re^{i(\theta-\xi)})^2} \right\| = \frac{1 - r^2}{1 - 2r \cos(\theta - \xi) + r^2}.$$

The last expression attains its maximum at $\theta = \xi + \pi/2$ and its minimum at $\theta = \xi - \pi/2$, and it is monotone (as a function of θ) on both intervals with these endpoints. Using this, one easily concludes that

$$\mathrm{var}(\log(Df); S^1) = 2 \left[\log \left(\frac{1 - r^2}{(1 + r)^2} \right) - \log \left(\frac{1 - r^2}{(1 - r)^2} \right) \right] = 4 \log \left(\frac{1 - r}{1 + r} \right).$$

Finally, it is well known that the last expression corresponds to $4 \mathrm{dist}_h(0, -e^{i\alpha}a)$, which closes the proof since $f(0) = -e^{i\alpha}a$. \square

Despite Proposition 4 above, the naive conjecture that, for general diffeomorphisms, positive asymptotic distortion implies the existence of hyperbolic periodic points is false. Indeed, below we provide an example of a C^∞ diffeomorphism of $[0, 1]$ having no fixed point in the interior and with positive asymptotic distortion yet the endpoints are parabolic fixed points.

Example 1. Start with a parabolic Möbius transformation, and denote \hat{f} its restriction to a fixed interval having no fixed point inside. After conjugacy by an affine map, we may assume that this interval is $[0, 1]$ and (passing to the inverse if necessary) that $\hat{f}(x) > x$ for all $x \in (0, 1)$. By Proposition 4 above, the asymptotic distortion of \hat{f} vanishes.

Fix $a < b < f(a)$ in $(0, 1)$, and let f be a C^∞ diffeomorphism with no fixed point in the interior that coincides with \hat{f} on a neighborhood of $[0, a] \cup [f(a), 1]$ and such that $f(b) = \hat{f}(b)$ but $Df(b) \neq D\hat{f}(b)$. (Notice that f cannot be achieved as a conjugate of \hat{f} .) We claim that f has nonzero asymptotic distortion. Indeed, since the asymptotic distortion of \hat{f} is zero, given $\varepsilon > 0$ there exists N such that, for all $n \geq N$,

$$\sum_{i=1}^n |\log(D\hat{f}^n(\hat{f}^{-n+i}(b))) - \log(D\hat{f}^n(\hat{f}^{-n+i}(a)))| \leq \text{var}(\log(D\hat{f}^n); [0, 1]) \leq n\varepsilon.$$

Fix $\varepsilon < |\log(D\hat{f}(b)) - \log(Df(b))|/2$. The preceding inequality may be rewritten as

$$\sum_{i=1}^n \left| \sum_{j=0}^{n-1} \log(D\hat{f}(\hat{f}^{-n+i+j}(b))) - \log(D\hat{f}(\hat{f}^{-n+i+j}(a))) \right| \leq n\varepsilon.$$

In the left-hand side expression, we may replace \hat{f} by f at each occurrence except for that, for each index i , the value of Df is different from that of $D\hat{f}$ at the point $f^{-n+i+j}(b) = \hat{f}^{-n+i+j}(b)$ for $j = n - i$. This yields the inequality

$$\sum_{i=1}^n \left| [\log(D\hat{f}(b)) - \log(Df(b))] + \sum_{j=0}^{n-1} \log(Df(f^{-n+i+j}(b))) - \log(Df(f^{-n+i+j}(a))) \right| \leq n\varepsilon.$$

By the triangle inequality $|y| \geq |x| - |x + y|$, this implies that

$$\sum_{i=1}^n \left| \sum_{j=0}^{n-1} \log(Df(f^{-n+i+j}(b))) - \log(Df(f^{-n+i+j}(a))) \right| \geq n |\log(D\hat{f}(b)) - \log(Df(b))| - n\varepsilon,$$

Since the right-hand side expression above may be rewritten as

$$\sum_{i=1}^n |\log(Df^n(f^{-n+i}(b))) - \log(Df^n(f^{-n+i}(a)))|,$$

the choice of ε implies that

$$\text{var}(\log(Df^n); [0, 1]) \geq \frac{n |\log(D\hat{f}(b)) - \log(Df(b))|}{2}.$$

Letting n go to infinite, this shows that

$$dist_\infty(f) \geq \frac{|\log(D\hat{f}(b)) - \log(Df(b))|}{2}.$$

Starting with arbitrary nontrivial diffeomorphisms of zero asymptotic distortion, the technique above easily shows that the set of diffeomorphisms with positive asymptotic distortion is dense in the (closed) set of diffeomorphisms having only parabolic fixed points. Since the former set coincides with

$$\bigcup_m \bigcap_n \left\{ f : \frac{\text{var}(\log(Df^n); \mathbb{S}^1)}{n} \geq \frac{1}{m} \right\},$$

this shows that positive asymptotic distortion is a generic property in the latter set. However, the construction reveals even more than this: zero asymptotic distortion seems to force a very particular structure for the diffeomorphism. Despite this, we were not able to settle the question below:

Question 2. Does there exist a real-analytic diffeomorphism of the interval with no hyperbolic fixed point and having positive asymptotic distortion?

Many other natural questions concerning the value of the asymptotic distortion for specific types of diffeomorphisms arise.

Question 3. Can the asymptotic distortion of a nontrivial piecewise-projective C^1 diffeomorphism of the interval vanish?

Notice that the asymptotic distortion of every nontrivial piecewise-affine homeomorphism of the interval is positive, due to the presence of semi-hyperbolic fixed points.

Question 4. What is the range of values of the asymptotic distortion for the C^∞ diffeomorphisms of the interval that arise in the Ghys-Sergiescu's smooth realizations of Thompson's group F (see [8])? Is it possible in this framework to produce (nontrivial) diffeomorphisms with the same asymptotical distortion which are not smoothly conjugate?

Finally, another natural question (inspired by [16]) deals with the behavior of the sequence of distortions along iterates. (Compare Question 1.)

Question 5. Assume that the asymptotic distortion of a diffeomorphism f of $[0, 1]$ with no hyperbolic fixed point vanishes. Can the sequence $\text{var}(\log(Df^n); [0, 1])$ grow faster than logarithmically?

References

- [1] BOCHI, J. & NAVAS, A. A geometric path from zero Lyapunov exponents to rotation cocycles. *Erg. Theory and Dyn. Systems.* **35** (2015), 374-402.
- [2] DEROIN, B., KLEPTSYN, V. & NAVAS, A. On the ergodic theory of free group actions by real-analytic circle diffeomorphisms. *Invent. Math.* **212** (2018), 731-779.
- [3] DEROIN, B., KLEPTSYN, V. & NAVAS, A. Sur la dynamique unidimensionnelle en régularité intermédiaire. *Acta Math.* **199** (2007), 199-262.
- [4] DOUADY, R. & YOCCOZ, J.-C. Nombre de rotation des difféomorphismes du cercle et mesures automorphes. *Regul. Chaotic Dyn.* **4** (1999), 19-38.
- [5] HERMAN, M. Simple proofs of local conjugacy theorems for diffeomorphisms of the circle with almost every rotation number. *Bol. Soc. Brasil. Mat.* **16** (1985), 45-83.
- [6] HERMAN, M. Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Publ. Math. de l'IHÉS* **49** (1979), 5-234.
- [7] FAYAD, B. & KHANIN, K. Smooth linearization of commuting circle diffeomorphisms. *Ann. Math.* **170** (2009), 961-980.
- [8] GHYS, É. & SERGIESCU, V. Sur un groupe remarquable de difféomorphismes du cercle. *Comment. Math. Helv.* **62** (1987), 185-239.
- [9] KHANIN, K. & TEPLINSKY, A. Herman's theory revisited. *Invent. Math.* **178** (2009), 333-344.
- [10] KATZNELSON, Y. & ORNSTEIN, D. The differentiability of the conjugation of certain diffeomorphisms of the circle. *Ergodic Theory Dynam. Systems* **9** (1989), 643-680.
- [11] KLEPTSYN, V. & NAVAS, A. A Denjoy type theorem for commuting circle diffeomorphisms with derivatives having different Hölder regularity classes. *Moscow Math. J.* **8** (2009), 477-492.
- [12] MINAKAWA, H. Classification of exotic circles of $PL_+(S^1)$. *Hokkaido Math. Journal* **25** (1997), 685-697.
- [13] NAVAS, A. *Groups of circle diffeomorphisms*. Chicago Lectures in Mathematics (2011).
- [14] NAVAS, A. Sur les rapprochements par conjugaison en dimension 1 et classe C^1 . *Compositio Math.* **150** (2014), 1183-1195.
- [15] NAVAS, A. & TRIESTINO, M. On the invariant distributions of C^2 circle diffeomorphisms of irrational rotation number. *Math. Z.* **274** (2013), 315-321.
- [16] POLTEROVICH, L. & SODIN, M.. A growth gap for diffeomorphisms of the interval. *Journal d'Analyse Mathématique* **92** (2004), 191-209.
- [17] WATANABE, N. Growth sequences for circle diffeomorphisms. *Geometric and Functional Analysis GAFA* **17** (2007), 320-331.
- [18] YOCCOZ, J.-C. Conjugaison différentiable des difféomorphismes du cercle dont le nombre de rotation vérifie une condition diophantienne. *Ann. Sci. École Norm. Sup.* **17** (1984), 333-359.

Andrés Navas

Dpto de Matemática y C.C., Fac. de Ciencia, Univ. de Santiago de Chile
 Alameda 3363, Estación Central, Santiago, Chile
 Email address: andres.navas@usach.cl

Unidad Cuernavaca Instituto de Matemáticas
 Universidad Nacional Autónoma de México
 Campus Morelos