

**LIPSCHITZ STABILITY IN INVERSE SOURCE AND INVERSE
COEFFICIENT PROBLEMS FOR A FIRST- AND HALF-ORDER
TIME-FRACTIONAL DIFFUSION EQUATION**

ATSUSHI KAWAMOTO AND MANABU MACHIDA

ABSTRACT. We consider inverse problems for the first and half order time fractional equation. We establish the stability estimates of Lipschitz type in inverse source and inverse coefficient problems by means of the Carleman estimates.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with the boundary $\partial\Omega$ of C^2 class. We set $Q = \Omega \times (0, T)$, where $T > 0$. We use notations $\partial_t = \frac{\partial}{\partial t}$, $\partial_i = \frac{\partial}{\partial x_i}$ ($i = 1, 2, \dots, n$). We also use the multi index $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ with $\alpha_j \in \mathbb{N} \cup \{0\}$ ($j = 1, 2, \dots, n$), $\partial_x^\alpha = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_n^{\alpha_n}$, $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$. Let $\nu = \nu(x)$ be the outwards unit normal vector to $\partial\Omega$ at x and let $\partial_\nu = \nu \cdot \nabla$. In general, the β th order Caputo-type fractional derivative is defined by

$$\partial_t^\beta u(x, t) := \frac{1}{\Gamma(n - \beta)} \int_0^t \frac{1}{(t - \tau)^{\beta+1-n}} \frac{\partial^n u(x, \tau)}{\partial \tau^n} d\tau, \quad (x, t) \in Q,$$

for $n - 1 < \beta < n$, $n \in \mathbb{N}$ (See e.g., [8, 31]). Here, Γ is the gamma function. We consider the following first and half order time-fractional diffusion equation.

$$\begin{aligned} (1) \quad & (\rho_1 \partial_t + \rho_2 \partial_t^{\frac{1}{2}} - L)u(x, t) = g(x, t), & (x, t) \in Q, \\ (2) \quad & u(x, t) = h_1(x, t), & (x, t) \in \partial\Omega \times (0, T), \\ (3) \quad & u(x, 0) = h_2(x), & x \in \Omega, \end{aligned}$$

where $\rho_1 > 0$, $\rho_2 \neq 0$ are constants, and L is a symmetric uniformly elliptic operator given by

$$Lu(x, t) := \sum_{i,j=1}^n \partial_i(a_{ij}(x)\partial_j u(x, t)) - \sum_{j=1}^n b_j(x)\partial_j u(x, t) - c(x)u(x, t), \quad (x, t) \in Q.$$

We assume that $a_{ij} \in C^3(\bar{\Omega})$, $a_{ij} = a_{ji}$ ($i, j = 1, 2, \dots, n$), $b_j \in C^2(\bar{\Omega})$ ($j = 1, 2, \dots, n$), $c \in C^2(\bar{\Omega})$, and moreover there exists a constant $\mu > 0$ such that

$$\frac{1}{\mu} |\xi|^2 \leq \sum_{i,j=1}^n a_{ij}(x)\xi_i \xi_j \leq \mu |\xi|^2, \quad \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n, \quad x \in \bar{\Omega}.$$

In fluid dynamics, (1) appears in the Basset problem [4] when the motion of a particle in a nonuniform flow is considered [6, 21, 30]. The first and half order time-fractional equation (1) also appears in porous media. Starting with the microscopic diffusion in a heterogeneous medium which has two length scales: the microscopic length scale of a typical porous block and the relative fracture width, a diffusion

equation with the first- and half-order time derivatives is obtained at the large scale limit by the homogenization process [1, 2].

The first- and half-order equation (1) is one of parabolic equations with multiple time-fractional terms, i.e., the time-derivative part in the equation is given by $\sum_{j=0}^{\ell} p_j \partial_t^{\alpha_j}$, where $0 < \alpha_{\ell} < \cdots < \alpha_1 < \alpha_0 \leq 1$ and coefficients p_j generally depend on x . Initial-boundary-value problems for multi-term time-fractional diffusion equations were considered in [29]. In the case that all time-derivatives are non-integer order and the time-derivative part is given by $\sum_{j=1}^{\ell} p_j \partial_t^{\alpha_j}$, the well-posedness was investigated [23] and moreover the uniqueness in inverse boundary-value problems was proven [24]. An exact solution was obtained in the special case of a two-term time-fractional diffusion equation [5]. The uniqueness for two kinds of inverse problems of identifying fractional orders in diffusion equations with multiple time-fractional derivatives was proved [25]. The uniqueness in determining the spatial component of the source term from interior observation was established [14]. The maximum principle and uniqueness was considered in [27] for the determination of the temporal component of the source term from a single point observation. Also the unique continuation was considered for multi-term time-fractional diffusion equations [26].

In [16], the Hölder stability is proven for the inverse source problem of (1) (See also [22]). In this paper, we further prove the Lipschitz stability not only for the inverse source problem but also for the inverse coefficient problem for (1).

The methodology of our stability analysis is based on the technique of the Carleman estimate [9], which was pioneered by Bukhgeim and Klivanov [7] when they proved the global uniqueness in inverse problems. See also [17, 18], recent reviews [19, 35], and textbooks [13, 20]. The Carleman estimate is a weighted L^2 inequality for a solution of a partial differential equation. In the case of parabolic equations with one first-order time derivative, the global Lipschitz stability was proven by using this method of Carleman estimates [12]. In this paper we make use of the Carleman estimate for parabolic equations. The Carleman estimates have been used for differential equations with a single term time-fractional derivative [15, 34, 36].

This paper is organized as follows. In §2, inverse source problems are considered. In §3, inverse coefficient problems are considered. In §4, the Carleman estimate necessary for our paper is established. Finally, proofs of the main theorems are given in §5.

2. INVERSE SOURCE PROBLEMS

We consider the inverse problems of determining the time-independent source factor of (1) from spatial data and two types of observations. One is the boundary observation and the other is the interior observation.

Let $t_0 \in (0, T)$ be an arbitrarily fixed time. Let γ be an arbitrarily fixed open connected sub-boundary of $\partial\Omega$ and let ω be an arbitrary fixed sub-domain of Ω such that $\omega \Subset \Omega$. We set $\Sigma = \gamma \times (0, T)$ and $Q_{\omega} = \omega \times (0, T)$.

Moreover we choose $\delta > 0$ such that

$$0 < t_0 - \delta < t_0 < t_0 + \delta < T,$$

and we set $Q_\delta = \Omega \times (t_0 - \delta, t_0 + \delta)$, $\Sigma_\delta = \gamma \times (t_0 - \delta, t_0 + \delta)$, $Q_{\omega, \delta} = \omega \times (t_0 - \delta, t_0 + \delta)$.

We assume that

$$(4) \quad \begin{cases} R \in C([0, T]; C(\overline{\Omega})) \cap C^2((0, T); C^2(\overline{\Omega})) \cap C^3((0, T); C(\overline{\Omega})), \\ \partial_t^{\frac{1}{2}} R \in C^2((0, T); C(\overline{\Omega})) \text{ and } |R(x, t_0)| > 0, \quad x \in \overline{\Omega}. \end{cases}$$

Furthermore we define

$$\mathcal{U} = L^2(0, T; H^4(\Omega)) \cap H^1(0, T; H^2(\Omega)) \cap H^2(0, T; L^2(\Omega)).$$

Let us assume that $g(x, t)$ in (1) has the form

$$(5) \quad g(x, t) = f(x)R(x, t),$$

and set $h_1 = 0$ in Q , $h_2 = 0$ in Ω .

We consider

$$(6) \quad (\rho_1 \partial_t + \rho_2 \partial_t^{\frac{1}{2}} - L)u(x, t) = f(x)R(x, t), \quad (x, t) \in Q,$$

$$(7) \quad u(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T),$$

$$(8) \quad u(x, 0) = 0, \quad x \in \Omega,$$

and we investigate the two kinds of inverse problems depending on the way of observations.

In the inverse source problem via boundary observation, we determine $f(x)$, $x \in \Omega$ by spatial data $u(x, t_0)$, $x \in \Omega$ and boundary data on Σ . In the inverse source problem via interior observation, we determine $f(x)$, $x \in \Omega$ by spatial data $u(x, t_0)$, $x \in \Omega$ and interior data in Q_ω . The main theorems Theorem 2.1 and Theorem 2.2 are stated as follows.

Theorem 2.1. *Let us assume that $u, \partial_t u, \partial_t^2 u \in \mathcal{U}$ and u satisfies (6)–(8). We suppose that $f \in H^2(\Omega)$ with $f = 0$ on $\partial\Omega$ and $\nabla f = 0$ on γ and R satisfies (4). Then there exist constants $C > 0$ such that*

$$(9) \quad \|f\|_{H^2(\Omega)} \leq C \|u(\cdot, t_0)\|_{H^4(\Omega)} + CB,$$

where

$$B = \|\nabla \partial_t^3 u\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^{\frac{5}{2}} u\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^2 u\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^{\frac{3}{2}} u\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t u\|_{L^2(\Sigma_\delta)}.$$

Theorem 2.2. *Let us assume that $u, \partial_t u, \partial_t^2 u \in \mathcal{U}$ and u satisfies (6)–(8). We suppose that $f \in H^2(\Omega)$ with $f = 0$ on $\partial\Omega$ and $f = 0$ in ω and R satisfies (4). Then there exist constants $C > 0$ such that*

$$(10) \quad \|f\|_{H^2(\Omega)} \leq C \|u(\cdot, t_0)\|_{H^4(\Omega)} + CI,$$

where

$$I = \|\partial_t^3 u\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^{\frac{5}{2}} u\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^2 u\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^{\frac{3}{2}} u\|_{L^2(Q_{\omega, \delta})} + \|\partial_t u\|_{L^2(Q_{\omega, \delta})}.$$

Remark 1. There is another approach to obtain the Lipschitz stability in inverse source problems by final observation data. In [33], Sakamoto and Yamamoto considered the perturbation of the single term time-fractional diffusion equations with a parameter as the diffusion coefficient and they obtained the stability estimate by means of the analytic perturbation theory under the appropriate assumptions on the parameter. In our case, however, we may not adopt their methodology directly since we consider the diffusion coefficient without the perturbation.

3. INVERSE COEFFICIENT PROBLEMS

3.1. Determination of the zeroth-order coefficient. Let us consider the inverse problem of determining the zeroth-order coefficient. In (1), we consider two coefficients $c_k(x)$, $x \in \Omega$ ($k = 1, 2$), where $c_k \in C^2(\overline{\Omega})$, $c_k(x) \geq 0$, $x \in \Omega$ ($k = 1, 2$). Let $u_k(x, t)$ be the corresponding solutions. We write L as

$$(11) \quad Lu_k(x, t) = Au_k(x, t) - c_k(x)u_k(x, t),$$

where A is defined as

$$Au(x, t) = \sum_{i,j=1}^n \partial_i(a_{ij}(x)\partial_j u(x, t)) - \sum_{j=1}^n b_j(x)\partial_j u(x, t), \quad (x, t) \in Q.$$

By subtraction we obtain

$$\begin{cases} \left(\rho_1 \partial_t + \rho_2 \partial_t^{\frac{1}{2}} - A + c_1(x) \right) u(x, t) = f(x)R(x, t), & (x, t) \in Q, \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = 0, & x \in \Omega, \end{cases}$$

where

$$u(x, t) = u_1(x, t) - u_2(x, t), \quad f(x) = c_1(x) - c_2(x), \quad R(x, t) = -u_2(x, t)$$

for $(x, t) \in Q$. Thus we arrive at the following two theorems as a direct consequence of the inverse source problems. In both cases the Lipschitz stability is obtained. Theorem 3.1 is proved using the inverse source problem via boundary observation stated in Theorem 2.1. Theorem 3.2 is proved using the inverse source problem via interior observation stated in Theorem 2.2.

Theorem 3.1 (boundary observation). *Let $u_k, \partial_t u_k, \partial_t^2 u_k \in \mathcal{U}$ ($k = 1, 2$) and u_1, u_2 satisfy (1)–(3) with (11). We suppose that $c_1, c_2 \in C^2(\overline{\Omega})$ with $c_1 = c_2$ on $\partial\Omega$ and $\nabla c_1 = \nabla c_2$ on γ , and $R = -u_2$ satisfies (4). Then there exist constants $C > 0$ such that*

$$(12) \quad \|c_1 - c_2\|_{H^2(\Omega)} \leq C \|u_1(\cdot, t_0) - u_2(\cdot, t_0)\|_{H^4(\Omega)} + CB,$$

where

$$\begin{aligned} B = & \|\nabla \partial_t^3(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^{\frac{5}{2}}(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^2(u_1 - u_2)\|_{L^2(\Sigma_\delta)} \\ & + \|\nabla \partial_t^{\frac{3}{2}}(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t(u_1 - u_2)\|_{L^2(\Sigma_\delta)}. \end{aligned}$$

Theorem 3.2 (interior observation). *Let $u_k, \partial_t u_k, \partial_t^2 u_k \in \mathcal{U}$ ($k = 1, 2$) and u_1, u_2 satisfy (1)–(3) with (11). We suppose that $c_1, c_2 \in C^2(\overline{\Omega})$ with $c_1 = c_2$ in $\partial\Omega \cup \omega$ and $R = -u_2$ satisfies (4). Then there exist constants $C > 0$ such that*

$$(13) \quad \|c_1 - c_2\|_{H^2(\Omega)} \leq C \|u_1(\cdot, t_0) - u_2(\cdot, t_0)\|_{H^4(\Omega)} + CI,$$

where

$$I = \|\partial_t^3(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^{\frac{5}{2}}(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^2(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} \\ + \|\partial_t^{\frac{3}{2}}(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})}.$$

Remark 2. In the case of diffusion in porous media, the condition $|u_2(x, t_0)| = |R(x, t_0)| > 0$ for $x \in \overline{\Omega}$ means that the concentration of the target particles is nonzero at the macroscopic scale.

3.2. Determination of the diffusion coefficient. We consider diffusion coefficients a_k ($k = 1, 2$) and corresponding solutions u_k . Let us express L as

$$(14) \quad Lu(x, t) = \mathcal{A}_k u(x, t),$$

where \mathcal{A}_k is defined as

$$\mathcal{A}_k u(x, t) = \operatorname{div}(a_k(x) \nabla u(x, t)) - \mathbf{b}(x) \cdot \nabla u(x, t) - c(x)u(x, t), \quad (x, t) \in Q,$$

for $k = 1, 2$. We suppose that $a_k \in C^4(\overline{\Omega})$ ($k = 1, 2$), $\mathbf{b} = (b_1, b_2, \dots, b_n) \in \{C^3(\overline{\Omega})\}^n$ and $c \in C^3(\overline{\Omega})$. Moreover we assume that there exists a constant $m > 0$ such that $a_k(x) \geq m$, $x \in \Omega$ ($k = 1, 2$). We investigate the inverse problems of determining the diffusion coefficients a_k ($k = 1, 2$) by boundary observations and interior observations.

Set

$$u(x, t) = u_1(x, t) - u_2(x, t), \quad a(x) = a_1(x) - a_2(x), \quad r(x, t) = u_2(x, t)$$

for $(x, t) \in Q$. Then by subtracting the equations for $k = 2$ from ones for $k = 1$, we obtain

$$(15) \quad \begin{cases} (\rho_1 \partial_t + \rho_2 \partial_t^{\frac{1}{2}} - \mathcal{A}_1) u(x, t) = \operatorname{div}(a(x) \nabla r(x, t)), & (x, t) \in Q, \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, T), \\ u(x, 0) = 0, & x \in \Omega, \end{cases}$$

We assume that

$$(16) \quad \begin{cases} r \in C([0, T]; C^3(\overline{\Omega})) \cap C((0, T); C^5(\overline{\Omega})) \\ \quad \cap C^2((0, T); C^4(\overline{\Omega})) \cap C^3((0, T); C^2(\overline{\Omega})), \\ \partial_t^{\frac{1}{2}} r \in C((0, T); C^3(\overline{\Omega})) \cap C^2((0, T); C^2(\overline{\Omega})). \end{cases}$$

Let us introduce weight functions for the Carleman estimates introduced in §4. According to observation types we consider in this paper, we prepare two kinds of distance functions d_1 and d_2 . We choose $d_1 \in C^2(\overline{\Omega})$ such that

$$d_1(x) > 0, \quad x \in \Omega, \quad |\nabla d_1(x)| > \sigma_1, \quad x \in \overline{\Omega},$$

$$\sum_{i,j=1}^n a_{ij}(x) \partial_i d_1 \nu_j \leq 0, \quad x \in \partial\Omega \setminus \gamma,$$

where $\sigma_1 > 0$ is a constant. Let ω_0 be an arbitrarily fixed sub-domain of Ω such that $\omega_0 \Subset \omega$. We take $d_2 \in C^2(\overline{\Omega})$ such that

$$d_2(x) > 0, \quad x \in \Omega, \quad |\nabla d_2(x)| > \sigma_2, \quad x \in \overline{\Omega \setminus \omega_0}, \quad d_2(x) = 0, \quad x \in \partial\Omega,$$

where $\sigma_2 > 0$ is a constant. The existence of the distance functions d_1 and d_2 is proved in [10, 11, 12]. Then we introduce weight functions φ_k, ψ_k ($k = 1, 2$) as

$$\varphi_k(x, t) = \frac{e^{\lambda d_k(x)}}{\ell(t)}, \quad \psi_k(x, t) = \frac{e^{\lambda d_k(x)} - e^{2\lambda \|d_k\|_{C(\overline{\Omega})}}}{\ell(t)}, \quad (x, t) \in Q,$$

where $\ell(t) = t(T - t)$. Moreover we assume that there exists a constant $m_1 > 0$ such that

$$(17) \quad |\nabla r(x, t_0) \cdot \nabla d_1(x)| \geq m_1, \quad x \in \overline{\Omega},$$

or that there exists a constant $m_2 > 0$ such that

$$(18) \quad |\nabla r(x, t_0) \cdot \nabla d_2(x)| \geq m_2, \quad x \in \overline{\Omega \setminus \omega}.$$

Let D' be an arbitrary sub-domain such that $\omega \Subset D' \Subset \Omega$. Set $D = \Omega \setminus D'$. Henceforth we suppose that $a \equiv 0$ in D .

Now we are ready to state our main results.

Theorem 3.3 (boundary observation). *Let $u_k, \partial_t u_k, \partial_t^2 u_k, \nabla u_k \in \mathcal{U}$ ($k = 1, 2$) and u_1, u_2 satisfy (1)–(3) with (14). We suppose that $a_1, a_2 \in C^4(\overline{\Omega})$ with $a_1 = a_2$ in D and $r = u_2$ satisfies (16) and (17). Then there exist constants $C > 0$ such that*

$$(19) \quad \|a_1 - a_2\|_{H^3(\Omega)} \leq C \|u_1(\cdot, t_0) - u_2(\cdot, t_0)\|_{H^5(\Omega)} + CB,$$

where

$$B = \|\nabla \partial_t^3(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^{\frac{5}{2}}(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t^2(u_1 - u_2)\|_{L^2(\Sigma_\delta)} \\ + \|\nabla \partial_t^{\frac{3}{2}}(u_1 - u_2)\|_{L^2(\Sigma_\delta)} + \|\nabla \partial_t(u_1 - u_2)\|_{L^2(\Sigma_\delta)}.$$

Theorem 3.4 (interior observation). *Let $u_k, \partial_t u_k, \partial_t^2 u_k, \nabla u_k \in \mathcal{U}$ ($k = 1, 2$) and u_1, u_2 satisfy (1)–(3) with (14). We suppose that $a_1, a_2 \in C^4(\overline{\Omega})$ with $a_1 = a_2$ in $D \cup \omega$ and $r = u_2$ satisfies (16) and (18). Then there exist constants $C > 0$ such that*

$$(20) \quad \|a_1 - a_2\|_{H^3(\Omega)} \leq C \|u_1(\cdot, t_0) - u_2(\cdot, t_0)\|_{H^5(\Omega)} + CI,$$

where

$$I = \|\partial_t^3(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^{\frac{5}{2}}(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t^2(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} \\ + \|\partial_t^{\frac{3}{2}}(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})} + \|\partial_t(u_1 - u_2)\|_{L^2(Q_{\omega, \delta})}.$$

Remark 3. In one dimensional case in space, we may relax some assumptions on u_k ($k = 1, 2$). It depends on the assumptions of the Carleman estimate for the third order partial differential equations (Lemma 4.8 and Lemma 4.9). See also [32].

4. CARLEMAN ESTIMATE

In this section, we establish the Carleman estimates for (1). We transform (1) into an integer-order partial differential equation. The calculation is similar to [34]. Let us begin with the following Lemma.

Lemma 4.1 (Lemma 3.1 in [16]). *If $u \in C([0, T]; H^4(\Omega)) \cap C^1((0, T); H^2(\Omega)) \cap C^2((0, T); L^2(\Omega))$ satisfies (1) through (3), then u satisfies*

$$(21) \quad \rho_2^2 \partial_t u(x, t) - (\rho_1 \partial_t - L)^2 u(x, t) = G(x, t), \quad (x, t) \in Q$$

where

$$(22) \quad G(x, t) = \left[\rho_2 \partial_t^{\frac{1}{2}} - (\rho_1 \partial_t - L) \right] g(x, t) + \frac{\rho_2 g(x, 0)}{\sqrt{\pi t}}, \quad (x, t) \in Q.$$

Although $\partial_t^{\frac{1}{2}} \partial_t^{\frac{1}{2}} \neq \partial_t$ in general, we may obtain the above lemma by applying $\rho_2 \partial_t^{\frac{1}{2}} - (\rho_1 \partial_t - L)$ to the both hand side of (1) and using $u(x, 0) = 0$, $x \in \Omega$.

Now we are ready to state our Carleman estimates.

Theorem 4.2 (Carleman estimate for (1) with boundary data). *Let $p \geq 0$. Suppose that $g(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ and $\nabla g(x, t) = 0$, $(x, t) \in \Sigma$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$(23) \quad \begin{aligned} & \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t^2 u|^2 + \sum_{i,j=1}^n |\partial_t \partial_i \partial_j u|^2 \right) + (s\varphi_1)^{p+1} |\nabla \partial_t u|^2 \right. \\ & \quad \left. + (s\varphi_1)^{p+2} |\nabla(\rho_1 \partial_t - L)u|^2 + (s\varphi_1)^{p+3} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p+5} |\nabla u|^2 + (s\varphi_1)^{p+7} |u|^2 \right] e^{2s\psi_1} dx dt \\ & \leq C \int_Q (s\varphi_1)^{p+1} |[\rho_2^2 \partial_t - (\rho_1 \partial_t - L)^2] u|^2 e^{2s\psi_1} dx dt \\ & \quad + C \int_{\Sigma} \left[(s\varphi_1)^{p+1} |\nabla \partial_t u|^2 + (s\varphi_1)^{p+2} |\nabla \partial_t^{\frac{1}{2}} u|^2 + (s\varphi_1)^{p+5} |\nabla u|^2 \right] e^{2s\psi_1} dS dt, \end{aligned}$$

for all $s > s_0$ and all $u \in \mathcal{U}$ satisfying (1) with $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ and $u(x, 0) = 0$, $x \in \Omega$.

Theorem 4.3 (Carleman estimate for (1) with interior data). *Let $p \geq 0$. Suppose that $g(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ and $g(x, t) = 0$, $(x, t) \in Q_\omega$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$(24) \quad \int_Q \left[(s\varphi_2)^{p-1} \left(|\partial_t^2 u|^2 + \sum_{i,j=1}^n |\partial_t \partial_i \partial_j u|^2 \right) + (s\varphi_2)^{p+1} |\nabla \partial_t u|^2 \right.$$

$$\begin{aligned}
& + (s\varphi_2)^{p+2} |\nabla(\rho_1 \partial_t - L)u|^2 + (s\varphi_2)^{p+3} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \\
& + (s\varphi_2)^{p+5} |\nabla u|^2 + (s\varphi_2)^{p+7} |u|^2 \Big] e^{2s\psi_2} dxdt \\
& \leq C \int_Q (s\varphi_2)^{p+1} |[\rho_2^2 \partial_t - (\rho_1 \partial_t - L)^2] u|^2 e^{2s\psi_2} dxdt \\
& + C \int_{Q_\omega} \left[(s\varphi_2)^{p+3} |\partial_t u|^2 + (s\varphi_2)^{p+4} |\partial_t^{\frac{1}{2}} u|^2 + (s\varphi_2)^{p+7} |u|^2 \right] e^{2s\psi_2} dxdt,
\end{aligned}$$

for all $s > s_0$ and all $u \in \mathcal{U}$ satisfying (1) with $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ and $u(x, 0) = 0$, $x \in \Omega$.

To prove Theorems 4.2 and 4.3, we start with the global Carleman estimates for parabolic equations (see e.g., [11, 35]) stated in Lemmas 4.4 and 4.5 below.

Lemma 4.4. *Let $p \geq 0$. There exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\begin{aligned}
& \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t v|^2 + \sum_{i,j=1}^n |\partial_i \partial_j v|^2 \right) + (s\varphi_1)^{p+1} |\nabla v|^2 + (s\varphi_1)^{p+3} |v|^2 \right] e^{2s\psi_1} dxdt \\
& \leq C \int_Q (s\varphi_1)^p |(\rho_1 \partial_t - L)v|^2 e^{2s\psi_1} dxdt + C \int_\Sigma (s\varphi_1)^{p+1} |\nabla v|^2 e^{2s\psi_1} dSdt,
\end{aligned}$$

for all $s > s_0$ and all $v \in L^2(0, T; H^2(\Omega)) \cap H^1(0, T; L^2(\Omega))$ satisfying $v(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$.

Lemma 4.5. *Let $p \geq 0$. There exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\begin{aligned}
& \int_Q \left[(s\varphi_2)^{p-1} \left(|\partial_t v|^2 + \sum_{i,j=1}^n |\partial_i \partial_j v|^2 \right) + (s\varphi_2)^{p+1} |\nabla v|^2 + (s\varphi_2)^{p+3} |v|^2 \right] e^{2s\psi_2} dxdt \\
& \leq C \int_Q (s\varphi_2)^p |(\rho_1 \partial_t - L)v|^2 e^{2s\psi_2} dxdt + C \int_{Q_\omega} (s\varphi_2)^{p+3} |v|^2 e^{2s\psi_2} dxdt,
\end{aligned}$$

for all $s > s_0$ and all $v \in L^2(0, T; H^2(\Omega)) \cap H^1(0, T; L^2(\Omega))$ satisfying $v(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$.

Proof of Theorem 4.2. Throughout the proof, we assume that $s > 1$ is large enough to satisfy $s\varphi > 1$ in Q .

Equation (21) yields

$$(25) \quad \rho_1 \partial_t w(x, t) - Lw(x, t) = \rho_2^2 \partial_t u(x, t) - G(x, t), \quad (x, t) \in Q,$$

where

$$(26) \quad w(x, t) = \rho_1 \partial_t u(x, t) - Lu(x, t), \quad (x, t) \in Q.$$

Since $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ and $g(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$, we have by (1),

$$w(x, t) = \rho_1 \partial_t u(x, t) - Lu(x, t) = g(x, t) - \rho_2 \partial_t^{\frac{1}{2}} u(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T).$$

Applying the Lemma 4.4 to (25), we obtain

$$(27) \quad \begin{aligned} & \int_Q [(s\varphi_1)^{p_1-1} |\partial_t w|^2 + (s\varphi_1)^{p_1+1} |\nabla w|^2 + (s\varphi_1)^{p_1+3} |w|^2] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p_1} |\partial_t u|^2 e^{2s\psi_1} dxdt + C \int_Q (s\varphi_1)^{p_1} |G|^2 e^{2s\psi_1} dxdt \\ & \quad + C \int_\Sigma (s\varphi_1)^{p_1+1} |\nabla w|^2 e^{2s\psi_1} dSdt, \end{aligned}$$

for $p_1 \geq 0$. Next by applying Lemma 4.4 to (26), we obtain

$$(28) \quad \begin{aligned} & \int_Q \left[(s\varphi_1)^{p_2-1} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p_2+1} |\nabla u|^2 + (s\varphi_1)^{p_2+3} |u|^2 \right] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p_2} |w|^2 e^{2s\psi_1} dxdt + C \int_\Sigma (s\varphi_1)^{p_2+1} |\nabla u|^2 e^{2s\psi_1} dSdt, \end{aligned}$$

for $p_2 \geq 0$.

Putting $p_2 = p_1 + 1$ and substituting the estimate of $|\partial_t u|^2$ in (28) into the right-hand side of (27), we obtain

$$\begin{aligned} & \int_Q [(s\varphi_1)^{p_1-1} |\partial_t w|^2 + (s\varphi_1)^{p_1+1} |\nabla w|^2 + (s\varphi_1)^{p_1+3} |w|^2] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p_1+1} |w|^2 e^{2s\psi_1} dxdt + C \int_Q (s\varphi_1)^{p_1} |G|^2 e^{2s\psi_1} dxdt + CB_{1,p_1}, \end{aligned}$$

where

$$B_{1,p_1} = \int_\Sigma [(s\varphi_1)^{p_1+1} |\nabla w|^2 + (s\varphi_1)^{p_1+2} |\nabla u|^2] e^{2s\psi_1} dSdt.$$

Taking sufficiently large $s > 0$, we can absorb the first term on the right-hand side of the above inequality into the left-hand side and we have

$$(29) \quad \begin{aligned} & \int_Q [(s\varphi_1)^{p_1-1} |\partial_t w|^2 + (s\varphi_1)^{p_1+1} |\nabla w|^2 + (s\varphi_1)^{p_1+3} |w|^2] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p_1} |G|^2 e^{2s\psi_1} dxdt + CB_{1,p_1}. \end{aligned}$$

By (28) with $p_2 = p_1 + 3$ and (29), we obtain

$$(30) \quad \begin{aligned} & \int_Q \left[(s\varphi_1)^{p_1+1} |\nabla(\rho_1 \partial_t - L)u|^2 + (s\varphi_1)^{p_1+2} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p_1+4} |\nabla u|^2 + (s\varphi_1)^{p_1+6} |u|^2 \right] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p_1} |G|^2 e^{2s\psi_1} dxdt + CB_{2,p_1}, \end{aligned}$$

where

$$B_{2,p_1} = \int_\Sigma [(s\varphi_1)^{p_1+1} |\nabla w|^2 + (s\varphi_1)^{p_1+4} |\nabla u|^2] e^{2s\psi_1} dSdt.$$

Let us choose $p_1 = p + 1$ in (29). Then from (26) and (29), we have

$$\int_Q (s\varphi_1)^p |\partial_t(\rho_1 \partial_t u - Lu)|^2 e^{2s\psi_1} dxdt \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dxdt + CB_{1,p+1}.$$

Setting $u_0 = \partial_t u$, we obtain

$$(31) \quad \int_Q (s\varphi_1)^p |\rho_1 \partial_t u_0 - Lu_0|^2 e^{2s\psi_1} dxdt \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dxdt + CB_{1,p+1}.$$

If we use Lemma 4.4 with $v = u_0$ and applying (31), we obtain

$$\begin{aligned} & \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t u_0|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u_0|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p+1} |\nabla u_0|^2 + (s\varphi_1)^{p+3} |u_0|^2 \right] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dxdt + C \int_{\Sigma} (s\varphi_1)^{p+1} |\nabla u_0|^2 e^{2s\psi_1} dSdt + CB_{1,p+1}. \end{aligned}$$

Recalling $u_0 = \partial_t u$, we have

$$\begin{aligned} & \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t^2 u|^2 + \sum_{i,j=1}^n |\partial_t \partial_i \partial_j u|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p+1} |\nabla \partial_t u|^2 + (s\varphi_1)^{p+3} |\partial_t u|^2 \right] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dxdt + CB_{3,p}, \end{aligned}$$

where

$$B_{3,p} = \int_{\Sigma} [(s\varphi_1)^{p+1} |\nabla \partial_t u|^2 + (s\varphi_1)^{p+2} |\nabla w|^2 + (s\varphi_1)^{p+3} |\nabla u|^2] e^{2s\psi_1} dSdt.$$

Hence using (30), we obtain

$$\begin{aligned} & \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t^2 u|^2 + \sum_{i,j=1}^n |\partial_t \partial_i \partial_j u|^2 \right) + (s\varphi_1)^{p+1} |\nabla \partial_t u|^2 \right. \\ & \quad \left. + (s\varphi_1)^{p+2} |\nabla(\rho_1 \partial_t - L)u|^2 + (s\varphi_1)^{p+3} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \right. \\ & \quad \left. + (s\varphi_1)^{p+5} |\nabla u|^2 + (s\varphi_1)^{p+7} |u|^2 \right] e^{2s\psi_1} dxdt \\ & \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dxdt + CB_{4,p}, \end{aligned}$$

where

$$B_{4,p} = \int_{\Sigma} [(s\varphi_1)^{p+1} |\nabla \partial_t u|^2 + (s\varphi_1)^{p+2} |\nabla w|^2 + (s\varphi_1)^{p+5} |\nabla u|^2] e^{2s\psi_1} dSdt.$$

Finally, we consider the boundary term B_4 . Since $\nabla g = 0$ on Σ is assumed, $\nabla w = \nabla g - \rho_2 \nabla \partial_t^{\frac{1}{2}} u = -\rho_2 \nabla \partial_t^{\frac{1}{2}} u$ on Σ . Hence we have

$$\begin{aligned} & \int_Q \left[(s\varphi_1)^{p-1} \left(|\partial_t^2 u|^2 + \sum_{i,j=1}^n |\partial_t \partial_i \partial_j u|^2 \right) + (s\varphi_1)^{p+1} |\nabla \partial_t u|^2 \right. \\ & \quad + (s\varphi_1)^{p+2} |\nabla(\rho_1 \partial_t - L)u|^2 + (s\varphi_1)^{p+3} \left(|\partial_t u|^2 + \sum_{i,j=1}^n |\partial_i \partial_j u|^2 \right) \\ & \quad \left. + (s\varphi_1)^{p+5} |\nabla u|^2 + (s\varphi_1)^{p+7} |u|^2 \right] e^{2s\psi_1} dx dt \\ & \leq C \int_Q (s\varphi_1)^{p+1} |G|^2 e^{2s\psi_1} dx dt \\ & \quad + C \int_{\Sigma} \left[(s\varphi_1)^{p+1} |\nabla \partial_t u|^2 + (s\varphi_1)^{p+2} |\nabla \partial_t^{\frac{1}{2}} u|^2 + (s\varphi_1)^{p+5} |\nabla u|^2 \right] e^{2s\psi_1} dS dt. \end{aligned}$$

Thus we obtain (23). \square

Proof of Theorem 4.3. By using Lemma 4.5 instead of Lemma 4.4, we can prove Theorem 4.3 in the same way as Theorem 4.2. \square

Furthermore we need Carleman estimates for elliptic equations in the proof of the stability estimates in inverse source problems which we will develop in §5.

Let us assume that $\tilde{a}_{ij} \in C^1(\bar{\Omega})$, $\tilde{a}_{ij} = \tilde{a}_{ji}$ ($i, j = 1, \dots, n$), $\tilde{b}_j \in C(\bar{\Omega})$ ($j = 1, \dots, n$), $\tilde{c} \in C(\bar{\Omega})$, and that there exists a constant $\tilde{\mu} > 0$ such that

$$\frac{1}{\tilde{\mu}} |\xi|^2 \leq \sum_{i,j=1}^n \tilde{a}_{ij}(x) \xi_i \xi_j \leq \tilde{\mu} |\xi|^2, \quad \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n, \quad x \in \bar{\Omega}.$$

We consider the following symmetric uniformly elliptic operator.

$$\tilde{L}\tilde{v}(x) := \sum_{i,j=1}^n \partial_i(\tilde{a}_{ij}(x)\partial_j\tilde{v}(x)) - \sum_{j=1}^n \tilde{b}_j(x)\partial_j\tilde{v}(x) - \tilde{c}(x)\tilde{v}(x), \quad x \in \Omega.$$

Set $\tilde{\varphi}_k(x) := \varphi_k(x, t_0)$, $x \in \Omega$ and $\tilde{\psi}_k(x) := \psi_k(x, t_0)$, $x \in \Omega$ for $k = 1, 2$. Then we have the following Lemmas.

Lemma 4.6. *Let $p \geq 0$. There exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\begin{aligned} & \int_{\Omega} \left[(s\tilde{\varphi}_1)^{p-1} \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_1)^{p+1} |\nabla \tilde{v}|^2 + (s\tilde{\varphi}_1)^{p+3} |\tilde{v}|^2 \right] e^{2s\tilde{\psi}_1} dx \\ & \leq C \int_{\Omega} (s\tilde{\varphi}_1)^p |\tilde{L}\tilde{v}|^2 e^{2s\tilde{\psi}_1} dx + C \int_{\gamma} (s\tilde{\varphi}_1)^{p+1} |\nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dS, \end{aligned}$$

for all $s > s_0$ and all $\tilde{v} \in H^2(\Omega)$ satisfying $\tilde{v}(x) = 0$, $x \in \partial\Omega$.

Lemma 4.7. *Let $p \geq 0$. There exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\int_{\Omega} \left[(s\tilde{\varphi}_2)^{p-1} \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_2)^{p+1} |\nabla \tilde{v}|^2 + (s\tilde{\varphi}_2)^{p+3} |\tilde{v}|^2 \right] e^{2s\tilde{\psi}_2} dx$$

$$\leq C \int_{\Omega} (s\tilde{\varphi}_2)^p |\tilde{L}\tilde{v}|^2 e^{2s\tilde{\psi}_2} dx + C \int_{\omega} (s\tilde{\varphi}_2)^{p+3} |\tilde{v}|^2 e^{2s\tilde{\psi}_2} dx,$$

for all $s > s_0$ and all $\tilde{v} \in H^2(\Omega)$ satisfying $\tilde{v}(x) = 0$, $x \in \partial\Omega$.

These lemmas can be shown in the same manner as the parabolic case by means of integration by parts. Hence we omit the proofs of these lemmas here.

We conclude this section by introducing Carleman estimates for the third order partial differential equations which we use in the proof of the stability estimates in inverse problems of determining the diffusion coefficients.

Let $\mathbf{p} = (p_1, \dots, p_2) \in \{C^1(\bar{\Omega})\}^n$.

Lemma 4.8. *We assume that there exists $m_1 > 0$ such that $|\mathbf{p}(x) \cdot \nabla d_1(x)| \geq m_1$, $x \in \bar{\Omega}$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\begin{aligned} & \int_{\Omega} \left[s\tilde{\varphi}_1 \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k \tilde{v}|^2 + (s\tilde{\varphi}_1)^2 |\nabla \Delta \tilde{v}|^2 \right. \\ & \quad \left. + (s\tilde{\varphi}_1)^3 \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_1)^5 (|\nabla \tilde{v}|^2 + |\tilde{v}|^2) \right] e^{2s\tilde{\psi}_1} dx \\ & \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \Delta \tilde{v})|^2 + |\mathbf{p} \cdot \nabla \Delta \tilde{v}|^2) e^{2s\tilde{\psi}_1} dx, \end{aligned}$$

for all $s > s_0$ and all $\tilde{v} \in H^4(\Omega)$ satisfying $|\tilde{v}(x)| = |\nabla \tilde{v}(x)| = |\Delta \tilde{v}(x)| = |\nabla \Delta \tilde{v}(x)| = 0$, $x \in \partial\Omega$ and $|\nabla \partial_k \tilde{v}(x)| = 0$, $x \in \gamma$ ($k = 1, 2, \dots, n$).

Lemma 4.9. *We assume that there exists $m_2 > 0$ such that $|\mathbf{p}(x) \cdot \nabla d_2(x)| \geq m_2$, $x \in \bar{\Omega} \setminus \omega$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\begin{aligned} & \int_{\Omega} \left[s\tilde{\varphi}_2 \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k \tilde{v}|^2 + (s\tilde{\varphi}_2)^2 |\nabla \Delta \tilde{v}|^2 \right. \\ & \quad \left. + (s\tilde{\varphi}_2)^3 \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_2)^5 (|\nabla \tilde{v}|^2 + |\tilde{v}|^2) \right] e^{2s\tilde{\psi}_2} dx \\ & \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \Delta \tilde{v})|^2 + |\mathbf{p} \cdot \nabla \Delta \tilde{v}|^2) e^{2s\tilde{\psi}_2} dx, \end{aligned}$$

for all $s > s_0$ and all $\tilde{v} \in H^4(\Omega)$ satisfying $|\tilde{v}(x)| = |\nabla \tilde{v}(x)| = |\Delta \tilde{v}(x)| = |\nabla \Delta \tilde{v}(x)| = 0$, $x \in \partial\Omega$ and $\tilde{v}(x) = 0$, $x \in \omega$.

To establish the Carleman estimate for $\mathbf{p} \cdot \nabla \Delta \tilde{v}$, we start by proving the first order partial differential equations $\mathbf{p} \cdot \nabla \tilde{v}$.

Lemma 4.10. *We assume that there exists $m_1 > 0$ such that $|\mathbf{p}(x) \cdot \nabla d_1(x)| \geq m_1$, $x \in \bar{\Omega}$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\int_{\Omega} (s\tilde{\varphi}_1)^2 (|\nabla \tilde{v}|^2 + |\tilde{v}|^2) e^{2s\tilde{\psi}_1} dx \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \tilde{v})|^2 + |\mathbf{p} \cdot \nabla \tilde{v}|^2) e^{2s\tilde{\psi}_1} dx,$$

for all $s > s_0$ and all $\tilde{v} \in H^2(\Omega)$ satisfying $|\tilde{v}(x)| = |\nabla \tilde{v}(x)| = 0$, $x \in \partial\Omega$.

Lemma 4.11. *We assume that there exists $m_2 > 0$ such that $|\mathbf{p}(x) \cdot \nabla d_2(x)| \geq m_2$, $x \in \overline{\Omega} \setminus \overline{\omega}$. Then there exists $\lambda_0 > 0$ such that for any $\lambda > \lambda_0$, we can choose $s_0(\lambda) > 0$ for which there exists $C = C(s_0, \lambda) > 0$ such that*

$$\int_{\Omega} (s\tilde{\varphi}_2)^2 (|\nabla \tilde{v}|^2 + |\tilde{v}|^2) e^{2s\tilde{\psi}_2} dx \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \tilde{v})|^2 + |\mathbf{p} \cdot \nabla \tilde{v}|^2) e^{2s\tilde{\psi}_2} dx,$$

for all $s > s_0$ and all $\tilde{v} \in H^2(\Omega)$ satisfying $|\tilde{v}(x)| = |\nabla \tilde{v}(x)| = 0$, $x \in \partial\Omega$ and $\tilde{v}(x) = 0$, $x \in \omega$.

Proof of Lemma 4.10. Setting $\tilde{w} = \tilde{v}e^{s\tilde{\psi}_1}$ in Ω , we have

$$e^{s\tilde{\psi}_1}(\mathbf{p} \cdot \nabla \tilde{v}) = \mathbf{p} \cdot \nabla \tilde{w} - s\lambda\tilde{\varphi}_1(\mathbf{p} \cdot \nabla d_1)\tilde{w} \quad \text{in } \Omega.$$

Taking the weighted L^2 norm, by integrating by parts we obtain

$$\begin{aligned} & \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx \\ &= \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{w}|^2 dx + \int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 (\mathbf{p} \cdot \nabla d_1)^2 |\tilde{w}|^2 dx \\ & \quad - 2 \int_{\Omega} s\lambda\tilde{\varphi}_1(\mathbf{p} \cdot \nabla d_1) \sum_{j=1}^n p_j \tilde{w} \partial_j \tilde{w} dx \\ & \geq \int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 (\mathbf{p} \cdot \nabla d_1)^2 |\tilde{w}|^2 dx - \int_{\Omega} s\lambda\tilde{\varphi}_1(\mathbf{p} \cdot \nabla d_1) \sum_{j=1}^n p_j \partial_j (\tilde{w})^2 dx \\ &= \int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 (\mathbf{p} \cdot \nabla d_1)^2 |\tilde{w}|^2 dx + \int_{\Omega} s\lambda^2 \tilde{\varphi}_1 (\mathbf{p} \cdot \nabla d_1)^2 |\tilde{w}|^2 dx \\ & \quad + \int_{\Omega} s\lambda\tilde{\varphi}_1 [(\mathbf{p} \cdot \nabla d_1)(\operatorname{div} \mathbf{p}) + \mathbf{p} \cdot \nabla(\mathbf{p} \cdot \nabla d_1)] |\tilde{w}|^2 dx \end{aligned}$$

Hence we have

$$\int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\tilde{w}|^2 dx \leq C \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx + C \int_{\Omega} (s\lambda^2 \tilde{\varphi}_1 + s\lambda\tilde{\varphi}_1) |\tilde{w}|^2 dx.$$

Taking sufficiently large $s > 0$, we may absorb the second term on the right-hand side of the above inequality into the left-hand side and we see that

$$\int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\tilde{w}|^2 dx \leq C \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx,$$

that is,

$$(32) \quad \int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\tilde{v}|^2 e^{2s\tilde{\psi}_1} dx \leq C \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx.$$

Set $\tilde{v}_k = \partial_k \tilde{v}$ in Ω for $k = 1, 2, \dots, n$. We consider

$$\mathbf{p} \cdot \nabla \tilde{v}_k = \partial_k(\mathbf{p} \cdot \nabla \tilde{v}) - (\partial_k \mathbf{p}) \cdot \nabla \tilde{v}$$

Applying the estimate (32) to the above equation, we may obtain

$$\begin{aligned} \int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\tilde{v}_k|^2 e^{2s\tilde{\psi}_1} dx &\leq C \int_{\Omega} |\mathbf{p} \cdot \nabla \tilde{v}_k|^2 e^{2s\tilde{\psi}_1} dx \\ &\leq C \int_{\Omega} |\partial_k(\mathbf{p} \cdot \nabla \tilde{v})|^2 e^{2s\tilde{\psi}_1} dx + C \int_{\Omega} |(\partial_k \mathbf{p}) \cdot \nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx. \end{aligned}$$

Hence we have

$$\int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx \leq C \int_{\Omega} |\nabla(\mathbf{p} \cdot \nabla \tilde{v})|^2 e^{2s\tilde{\psi}_1} dx + C \int_{\Omega} |\nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx.$$

Choosing sufficiently large $s > 0$, we can absorb the second term on the right-hand side of the above inequality into the left-hand side and we may get

$$\int_{\Omega} s^2 \lambda^2 \tilde{\varphi}_1^2 |\nabla \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx \leq C \int_{\Omega} |\nabla(\mathbf{p} \cdot \nabla \tilde{v})|^2 e^{2s\tilde{\psi}_1} dx.$$

Combining this with (32), we obtain the Carleman estimate of Lemma 4.10. Thus we conclude the Lemma 4.10. \square

Proof of Lemma 4.11. By an argument similar to that used in the proof of Lemma 4.10, we may obtain Lemma 4.11. \square

Remark 4. In one spatial dimension, the assumption $|\nabla \tilde{v}| = 0$ on $\partial\Omega$ is not necessary in Lemma 4.10 and Lemma 4.11. In this case, we have the following Carleman estimate by integration by parts.

$$\int_{\Omega} (s\tilde{\varphi}_k)^2 (|\partial_1 \tilde{v}|^2 + |\tilde{v}|^2) e^{2s\tilde{\psi}_k} dx \leq C \int_{\Omega} |p_1 \partial_1 \tilde{v}|^2 e^{2s\tilde{\psi}_k} dx,$$

for $k = 1, 2$.

Now we are ready to prove Lemma 4.8 and Lemma 4.9.

Proof of Lemma 4.8. Set $\tilde{y} = \Delta \tilde{v}$ in Ω . By the assumptions $|\Delta \tilde{v}(x)| = |\nabla \Delta \tilde{v}(x)| = 0$, $x \in \partial\Omega$, we see that $|\tilde{y}(x)| = |\nabla \tilde{y}(x)| = 0$, $x \in \partial\Omega$. By Lemma 4.10, we obtain

$$\int_{\Omega} (s\tilde{\varphi}_1)^2 (|\nabla \tilde{y}|^2 + |\tilde{y}|^2) e^{2s\tilde{\psi}_1} dx \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \tilde{y})|^2 + |\mathbf{p} \cdot \nabla \tilde{y}|^2) e^{2s\tilde{\psi}_1} dx,$$

that is,

$$(33) \quad \begin{aligned} & \int_{\Omega} (s\tilde{\varphi}_1)^2 (|\nabla \Delta \tilde{v}|^2 + |\Delta \tilde{v}|^2) e^{2s\tilde{\psi}_1} dx \\ & \leq C \int_{\Omega} (|\nabla(\mathbf{p} \cdot \nabla \Delta \tilde{v})|^2 + |\mathbf{p} \cdot \nabla \Delta \tilde{v}|^2) e^{2s\tilde{\psi}_1} dx. \end{aligned}$$

Next we use the Carleman estimate for elliptic equations to estimate the left-hand side of the above inequality. By Lemma 4.6 with $p = 2$, we have

$$(34) \quad \begin{aligned} & \int_{\Omega} \left[s\tilde{\varphi}_1 \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_1)^3 |\nabla \tilde{v}|^2 + (s\tilde{\varphi}_1)^5 |\tilde{v}|^2 \right] e^{2s\tilde{\psi}_1} dx \\ & \leq C \int_{\Omega} (s\tilde{\varphi}_1)^2 |\Delta \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx. \end{aligned}$$

Setting $\tilde{v}_k = \partial_k \tilde{v}$ in Ω for $k = 1, 2, \dots, n$ and using Lemma 4.6 again, we see that

$$\begin{aligned} & \int_{\Omega} \left[s\tilde{\varphi}_1 \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}_k|^2 + (s\tilde{\varphi}_1)^3 |\nabla \tilde{v}_k|^2 + (s\tilde{\varphi}_1)^5 |\tilde{v}_k|^2 \right] e^{2s\tilde{\psi}_1} dx \\ & \leq C \int_{\Omega} (s\tilde{\varphi}_1)^2 |\Delta \tilde{v}_k|^2 e^{2s\tilde{\psi}_1} dx, \end{aligned}$$

that is,

$$(35) \quad \int_{\Omega} \left[s\tilde{\varphi}_1 \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k \tilde{v}|^2 + (s\tilde{\varphi}_1)^3 \sum_{i,j=1}^n |\partial_i \partial_j \tilde{v}|^2 + (s\tilde{\varphi}_1)^5 |\nabla \tilde{v}|^2 \right] e^{2s\tilde{\psi}_1} dx \\ \leq C \int_{\Omega} (s\tilde{\varphi}_1)^2 |\nabla \Delta \tilde{v}|^2 e^{2s\tilde{\psi}_1} dx.$$

Summing up the inequalities (33)–(35), we may obtain the Carleman estimate of Lemma 4.8 \square

Proof of Lemma 4.9. Using Lemma 4.11 and Lemma 4.7 instead of Lemma 4.10 and Lemma 4.6, we may prove Lemma 4.9 in the same way as Lemma 4.8. \square

5. PROOF OF STABILITY ESTIMATES

Hereafter we let C denote a generic constant which is independent of s, x, t and let $C(s)$ denote a generic constant which is independent of x, t but depends on s .

5.1. Stability for the zeroth-order coefficient.

Proof of Theorem 2.1. By using Lemma 4.1 for (6), we obtain

$$(36) \quad \rho_2^2 \partial_t u(x, t) - (\rho_1 \partial_t - L)^2 u(x, t) = F(x, t), \quad (x, t) \in Q,$$

where we introduced $F(x, t)$ as

(37)

$$F(x, t) = \left[\rho_2 \partial_t^{\frac{1}{2}} - (\rho_1 \partial_t - L) \right] (f(x)R(x, t)) + \rho_2 f(x) \frac{R(x, 0)}{\sqrt{\pi t}} \\ = R(x, t) \sum_{i,j=1}^n \partial_i (a_{ij}(x) \partial_j f(x)) \\ + \sum_{j=1}^n \left(2 \sum_{i=1}^n a_{ij}(x) \partial_i R(x, t) - b_j(x) R(x, t) \right) \partial_j f(x) \\ + \left[\rho_2 \partial_t^{\frac{1}{2}} R(x, t) - \rho_1 \partial_t R(x, t) + \sum_{i,j=1}^n \partial_i (a_{ij}(x) \partial_j R(x, t)) \right. \\ \left. - \sum_{j=1}^n b_j(x) \partial_j R(x, t) - c(x) R(x, t) + \frac{\rho_2 R(x, 0)}{\sqrt{\pi t}} \right] f(x), \quad (x, t) \in Q.$$

Let us set $y = \partial_t u$, $z = \partial_t^2 u$ in Q . By differentiating (36) with respect to t , we have

$$(38) \quad \rho_2^2 \partial_t y(x, t) - (\rho_1 \partial_t - L)^2 y(x, t) = \partial_t F(x, t), \quad (x, t) \in Q,$$

$$(39) \quad \rho_2^2 \partial_t z(x, t) - (\rho_1 \partial_t - L)^2 z(x, t) = \partial_t^2 F(x, t), \quad (x, t) \in Q.$$

Since $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$, we see that

$$y(x, t) = z(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T).$$

To use the Carleman estimate in Q_δ , we introduce the weight functions. Set

$$\varphi_{\delta,1}(x, t) = \frac{e^{\lambda d_1(x)}}{\ell_\delta(t)}, \quad \psi_{\delta,1}(x, t) = \frac{e^{\lambda d_1(x)} - e^{2\lambda \|d_1\|_C(\overline{\Omega})}}{\ell_\delta(t)}, \quad (x, t) \in Q_\delta,$$

where $\ell_\delta(t) = (t - t_0 + \delta)(t_0 + \delta - t)$. Fixing $\lambda > 0$ and applying Theorem 4.2 ($p = 0$) to (38) and (39) in Q_δ , we have

$$(40) \quad \int_{Q_\delta} \left[(s\varphi_{\delta,1})^3 \left(|\partial_t y|^2 + |\partial_t z|^2 + \sum_{i,j=1}^n |\partial_i \partial_j y|^2 + \sum_{i,j=1}^n |\partial_i \partial_j z|^2 \right) \right. \\ \left. + (s\varphi_{\delta,1})^5 (|\nabla y|^2 + |\nabla z|^2) + (s\varphi_{\delta,1})^7 (|y|^2 + |z|^2) \right] e^{2s\psi_{\delta,1}} dxdt \\ \leq C \int_{Q_\delta} s\varphi_{\delta,1} (|\partial_t F|^2 + |\partial_t^2 F|^2) e^{2s\psi_{\delta,1}} dxdt + C\tilde{B},$$

where

$$\tilde{B} = \int_{\Sigma_\delta} \left[s\varphi_{\delta,1} (|\nabla \partial_t y|^2 + |\nabla \partial_t z|^2) \right. \\ \left. + (s\varphi_{\delta,1})^2 (|\nabla \partial_t^{\frac{1}{2}} y|^2 + |\nabla \partial_t^{\frac{1}{2}} z|^2) + (s\varphi_{\delta,1})^5 (|\nabla y|^2 + |\nabla z|^2) \right] e^{2s\psi_{\delta,1}} dSdt.$$

We can estimate \tilde{B} by B^2 . We note that $\partial_t^{\frac{1}{2}} \partial_t^m = \partial_t^{m+\frac{1}{2}}$, $m \in \mathbb{N}$. Since there exist constants $C_k(s) > 0$ such that $\varphi_{\delta,1}^k e^{2s\psi_{\delta,1}} \leq C_k(s)$ on Σ_δ for $k = 0, 1, 2, \dots$, we have

$$\tilde{B} \leq C \int_{\Sigma_\delta} \left[s\varphi_{\delta,1} |\nabla \partial_t^3 u|^2 + (s\varphi_{\delta,1})^2 (|\nabla \partial_t^{\frac{1}{2}} \partial_t u|^2 + |\nabla \partial_t^{\frac{1}{2}} \partial_t^2 u|^2) \right. \\ \left. + (s\varphi_{\delta,1})^5 (|\nabla \partial_t u|^2 + |\nabla \partial_t^2 u|^2) \right] e^{2s\psi_{\delta,1}} dSdt \\ \leq C(s)B^2.$$

Note that

$$\int_{Q_\delta} s\varphi_{\delta,1} (|\partial_t F|^2 + |\partial_t^2 F|^2) e^{2s\psi_{\delta,1}} dxdt \leq C \int_{Q_\delta} s\varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dxdt.$$

This together with (40) gives

$$(41) \quad \int_{Q_\delta} \left[(s\varphi_{\delta,1})^3 \left(|\partial_t y|^2 + |\partial_t z|^2 + \sum_{i,j=1}^n |\partial_i \partial_j y|^2 + \sum_{i,j=1}^n |\partial_i \partial_j z|^2 \right) \right. \\ \left. + (s\varphi_{\delta,1})^5 (|\nabla y|^2 + |\nabla z|^2) + (s\varphi_{\delta,1})^7 (|y|^2 + |z|^2) \right] e^{2s\psi_{\delta,1}} dxdt \\ \leq C \int_{Q_\delta} s\varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dxdt + C(s)B^2.$$

Let us expand the left-hand side of (36). We have

$$\rho_2^2 \partial_t u(x, t) - \rho_1^2 \partial_t^2 u(x, t) + 2\rho_1 \partial_t Lu(x, t) - L^2 u(x, t) = F(x, t), \quad (x, t) \in Q.$$

In particular at $t = t_0$, we have

$$(42) \quad \rho_2^2 \partial_t u(x, t_0) - \rho_1^2 \partial_t^2 u(x, t_0) + 2\rho_1 \partial_t Lu(x, t_0) - L^2 u(x, t_0) = F(x, t_0), \quad x \in \Omega.$$

Taking the weighted L^2 norm of (42) in Ω , we obtain

$$(43) \quad \begin{aligned} & \int_{\Omega} \varphi_{\delta,1}(x, t_0) |F(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq C \sum_{k=1}^3 J_k + C \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 4} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \end{aligned}$$

where

$$\begin{aligned} J_1 &= \int_{\Omega} \varphi_{\delta,1}(x, t_0) |\partial_t u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \\ J_2 &= \int_{\Omega} \varphi_{\delta,1}(x, t_0) |\partial_t^2 u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \\ J_3 &= \int_{\Omega} \varphi_{\delta,1}(x, t_0) |\partial_t Lu(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx. \end{aligned}$$

Let us estimate J_1, J_2, J_3 . We assume that $s > 1$ is large enough to satisfy $s\varphi_{\delta,1} > 1$ in Q . We note that $\partial_t \psi_{\delta,1}(x, t) = (e^{2\lambda(\|d_1\|_{C(\bar{\Omega})} - d_1(x))} - e^{-\lambda d_1(x)})(T - 2t)\varphi_{\delta,1}^2(x, t)$ for $(x, t) \in Q$.

$$\begin{aligned} J_1 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (\varphi_{\delta,1} |y|^2 e^{2s\psi_{\delta,1}}) dx dt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} [\varphi_{\delta,1}^2 |y|^2 + \varphi_{\delta,1} |\partial_t y| |y| + s\varphi_{\delta,1}^3 |y|^2] e^{2s\psi_{\delta,1}} dx dt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^3 (|y|^2 + |z|^2) e^{2s\psi_{\delta,1}} dx dt. \end{aligned}$$

Combining this with (41), we may estimate the right-hand side of the above inequality and we obtain

$$(44) \quad J_1 \leq \frac{C}{s^5} \int_{Q_\delta} \varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dx dt + C(s)B^2.$$

Similarly, we obtain

$$\begin{aligned} J_2 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (\varphi_{\delta,1} |\partial_t y|^2 e^{2s\psi_{\delta,1}}) dx dt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} [\varphi_{\delta,1}^2 |\partial_t y|^2 + \varphi_{\delta,1} |\partial_t^2 y| |\partial_t y| + s\varphi_{\delta,1}^3 |\partial_t y|^2] e^{2s\psi_{\delta,1}} dx dt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^3 (|\partial_t y|^2 + |\partial_t z|^2) e^{2s\psi_{\delta,1}} dx dt. \end{aligned}$$

Putting this together with (41), we see that

$$(45) \quad J_2 \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dx dt + C(s)B^2.$$

Moreover we have

$$\begin{aligned} J_3 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (\varphi_{\delta,1} |Ly|^2 e^{2s\psi_{\delta,1}}) dx dt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} [\varphi_{\delta,1}^2 |Ly|^2 + \varphi_{\delta,1} |\partial_t Ly| |Ly| + s\varphi_{\delta,1}^3 |Ly|^2] e^{2s\psi_{\delta,1}} dx dt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^3 (|Ly|^2 + |Lz|^2) e^{2s\psi_{\delta,1}} dx dt \end{aligned}$$

$$\leq C \int_{Q_\delta} s \varphi_{\delta,1}^3 \left(\sum_{|\alpha| \leq 2} |\partial_x^\alpha y|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha z|^2 \right) e^{2s\psi_{\delta,1}} dx dt$$

This together with (41) gives

$$(46) \quad J_3 \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dx dt + C(s)B^2.$$

By (43) through (46), we have

$$(47) \quad \begin{aligned} & \int_{\Omega} \varphi_{\delta,1}(x, t_0) |F(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dx dt \\ & \quad + C \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 4} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2. \end{aligned}$$

We will estimate the left-hand side of the inequality (47) from below using the Carleman estimate for the elliptic equation stated Lemma 4.6 ($p = 1$). By (37) at $t = t_0$, we have

$$(48) \quad \begin{aligned} & \sum_{i,j=1}^n \partial_i(a_{ij}(x)\partial_j \tilde{f}(x)) \\ & \quad + \frac{1}{R(x, t_0)} \sum_{j=1}^n \left(2 \sum_{i=1}^n a_{ij}(x)\partial_i R(x, t_0) - b_j(x)R(x, t_0) \right) \partial_j \tilde{f}(x) \\ & \quad + \frac{1}{R(x, t_0)} \left[\rho_2 \partial_t^{\frac{1}{2}} R(x, t_0) - \rho_1 \partial_t R(x, t_0) + \sum_{i,j=1}^n \partial_i(a_{ij}(x)\partial_j R(x, t_0)) \right. \\ & \quad \quad \left. - \sum_{j=1}^n b_j(x)\partial_j R(x, t_0) - c(x)R(x, t_0) + \frac{\rho_2 R(x, 0)}{\sqrt{\pi t_0}} \right] \tilde{f}(x) \\ & = \frac{F(x, t_0)}{R(x, t_0)}, \quad x \in \Omega. \end{aligned}$$

We note that $f(x) = 0$, $x \in \partial\Omega$ and $\nabla f(x) = 0$, $x \in \gamma$ are assumed. Applying the Lemma 4.6 to (48) in Ω , we obtain

$$(49) \quad \begin{aligned} & \frac{1}{s} \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 2} |\partial_x^\alpha f(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq \frac{C}{s} \int_{\Omega} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq \frac{C}{s} \int_{\Omega} \left(\sum_{i,j=1}^n |\partial_i \partial_j f(x)|^2 \right. \\ & \quad \left. + (s\varphi_{\delta,1}(x, t_0))^2 |\nabla f(x)|^2 + (s\varphi_{\delta,1}(x, t_0))^4 |f(x)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq C \int_{\Omega} \varphi_{\delta,1}(x, t_0) \left| \frac{F(x, t_0)}{R(x, t_0)} \right|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \end{aligned}$$

$$\leq C \int_{\Omega} \varphi_{\delta,1}(x, t_0) |F(x, t_0)|^2 e^{2s\psi(x, t_0)} dx.$$

By (47) and (49), we obtain

$$(50) \quad \begin{aligned} & \frac{1}{s} \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 2} |\partial_x^\alpha f(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 2} |\partial_x^\alpha f(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx dt \\ & \quad + C \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 4} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2. \end{aligned}$$

Let us estimate the first integral term on the right-hand side of (50).

$$\int_{Q_\delta} \varphi_{\delta,1} \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}} dx dt \leq \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 2} |\partial_x^\alpha f|^2 e^{2s\psi_{\delta,1}(x, t_0)} h_s(x) dx,$$

where

$$h_s(x) = \frac{1}{\varphi_{\delta,1}(x, t_0)} \int_{t_0-\delta}^{t_0+\delta} \varphi_{\delta,1} e^{-2s(\psi_{\delta,1}(x, t_0) - \psi_{\delta,1}(x, t))} dt.$$

Since $\psi_{\delta,1}(x, t_0) - \psi_{\delta,1}(x, t) \geq 0$, $(x, t) \in Q_\delta$, h_s converges pointwise to 0 in Ω as $s \rightarrow \infty$ by Lebesgue's dominated convergence theorem. Moreover by Dini's theorem, we see that h_s converges uniformly to 0 in Ω as $s \rightarrow \infty$. Hence, taking sufficiently large $s > 0$, we can absorb the first term on the right-hand side of (50) into the left-hand side and obtain

$$(51) \quad \begin{aligned} & \frac{1}{s} \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 2} |\partial_x^\alpha f(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx \\ & \leq C \int_{\Omega} \varphi_{\delta,1}(x, t_0) \sum_{|\alpha| \leq 4} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2. \end{aligned}$$

Fix $s > 0$. Noting that $\varphi_{\delta,1}(\cdot, t_0) e^{2s\psi_{\delta,1}(\cdot, t_0)}$ has its upper and lower bound in $\bar{\Omega}$, we see that

$$\|f\|_{H^2(\Omega)} \leq C \|u(\cdot, t_0)\|_{H^4(\Omega)} + CB.$$

Thus we obtain the stability estimate. \square

Proof of Theorem 2.2. We may prove Theorem 2.2 by an argument similar to that used in the proof of Theorem 2.1. In the proof, Theorem 4.3 and Lemma 4.7 are used instead of Theorem 4.2 and Lemma 4.6. \square

5.2. Stability for the diffusion coefficient. Next we prove Theorem 3.3 and Theorem 3.4. The proofs is very similar to the proofs of Theorem 2.1 and Theorem 2.2.

Proof of Theorem 3.3. Applying Lemma 4.1 to (15), we obtain

$$(52) \quad \rho_2^2 \partial_t^2 u(x, t) - (\rho_1 \partial_t - \mathcal{A}_1)^2 u(x, t) = \tilde{F}(x, t), \quad (x, t) \in Q,$$

where

$$(53) \quad \tilde{F}(x, t) = \left[\rho_2 \partial_t^{\frac{1}{2}} - (\rho_1 \partial_t - \mathcal{A}_1) \right] (\operatorname{div}(a(x) \nabla r(x, t))) + \frac{\rho_2 \operatorname{div}(a(x) \nabla r(x, 0))}{\sqrt{\pi t}}$$

$$\begin{aligned}
&= a_1(x) \nabla r(x, t) \cdot \nabla \Delta a(x) + 2a_1(x) \sum_{i,j=1}^n (\partial_i \partial_j r(x, t)) (\partial_i \partial_j a(x)) \\
&\quad + a_1(x) \Delta r(x, t) \Delta a(x) + (\nabla a_1(x) - \mathbf{b}(x)) \cdot (\nabla r(x, t) \cdot \nabla) \nabla a(x) \\
&\quad + \left[(\rho_2 \partial_t^{\frac{1}{2}} - \rho_1 \partial_t) \nabla r(x, t) + 3a_1(x) \nabla \Delta r(x, t) + (\Delta r(x, t)) \nabla a_1(x) \right. \\
&\quad \quad \left. - (\Delta r(x, t)) \mathbf{b}(x) - c(x) \nabla r(x, t) + \frac{\rho_2 \nabla r(x, 0)}{\sqrt{\pi t}} \right] \cdot \nabla a(x) \\
&\quad + (\nabla a_1(x) - \mathbf{b}(x)) \cdot (\nabla a(x) \cdot \nabla) \nabla r(x, t) \\
&\quad + \left[(\rho_2 \partial_t^{\frac{1}{2}} - \rho_1 \partial_t) \Delta r(x, t) + (\nabla a_1(x) \cdot \nabla) \Delta r(x, t) + a_1(x) \Delta^2 r(x, t) \right. \\
&\quad \quad \left. - (\mathbf{b}(x) \cdot \nabla) \Delta r(x, t) - c(x) \Delta r(x, t) + \frac{\rho_2 \Delta r(x, 0)}{\sqrt{\pi t}} \right] a(x), \quad (x, t) \in Q.
\end{aligned}$$

Setting $y = \partial_t u$, $z = \partial_t^2 u$ in Q and differentiating (52) with respect to t , we have

$$(54) \quad \rho_2^2 \partial_t y(x, t) - (\rho_1 \partial_t - \mathcal{A}_1)^2 y(x, t) = \partial_t \tilde{F}(x, t), \quad (x, t) \in Q,$$

$$(55) \quad \rho_2^2 \partial_t z(x, t) - (\rho_1 \partial_t - \mathcal{A}_1)^2 z(x, t) = \partial_t^2 \tilde{F}(x, t), \quad (x, t) \in Q.$$

Since $u(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$, we see that

$$y(x, t) = z(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T).$$

Fixing $\lambda > 0$ and applying Theorem 4.2 ($p = 1$) to (54) and (55) in Q_δ , we have

$$\begin{aligned}
(56) \quad &\int_{Q_\delta} \left[(s\varphi_{\delta,1})^2 (|\nabla \partial_t y|^2 + |\nabla \partial_t z|^2) \right. \\
&\quad + (s\varphi_{\delta,1})^3 (|\nabla(\rho_1 \partial_t - \mathcal{A}_1)y|^2 + |\nabla(\rho_1 \partial_t - \mathcal{A}_1)z|^2) \\
&\quad + (s\varphi_{\delta,1})^4 \left(|\partial_t y|^2 + |\partial_t z|^2 + \sum_{i,j=1}^n |\partial_i \partial_j y|^2 + \sum_{i,j=1}^n |\partial_i \partial_j z|^2 \right) \\
&\quad \left. + (s\varphi_{\delta,1})^6 (|\nabla y|^2 + |\nabla z|^2) + (s\varphi_{\delta,1})^8 (|y|^2 + |z|^2) \right] e^{2s\psi_{\delta,1}} dxdt \\
&\leq C \int_{Q_\delta} (s\varphi_{\delta,1})^2 (|\partial_t \tilde{F}|^2 + |\partial_t^2 \tilde{F}|^2) e^{2s\psi_{\delta,1}} dxdt + C\hat{B},
\end{aligned}$$

where

$$\begin{aligned}
\hat{B} &= \int_{\Sigma_\delta} \left[(s\varphi_{\delta,1})^2 (|\nabla \partial_t y|^2 + |\nabla \partial_t z|^2) \right. \\
&\quad \left. + (s\varphi_{\delta,1})^3 (|\nabla \partial_t^{\frac{1}{2}} y|^2 + |\nabla \partial_t^{\frac{1}{2}} z|^2) + (s\varphi_{\delta,1})^6 (|\nabla y|^2 + |\nabla z|^2) \right] e^{2s\psi_{\delta,1}} dSdt.
\end{aligned}$$

As we have seen in the proof of Theorem 2.1, we may obtain $\hat{B} \leq C(s)B^2$.

Note that

$$\int_{Q_\delta} (s\varphi_{\delta,1})^2 (|\partial_t \tilde{F}|^2 + |\partial_t^2 \tilde{F}|^2) e^{2s\psi_{\delta,1}} dxdt$$

$$\leq C \int_{Q_\delta} (s\varphi_{\delta,1})^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dx dt.$$

This together with (56) gives

$$(57) \quad \int_{Q_\delta} \left[(s\varphi_{\delta,1})^2 (|\nabla \partial_t y|^2 + |\nabla \partial_t z|^2) \right. \\ + (s\varphi_{\delta,1})^3 (|\nabla(\rho_1 \partial_t - \mathcal{A}_1)y|^2 + |\nabla(\rho_1 \partial_t - \mathcal{A}_1)z|^2) \\ + (s\varphi_{\delta,1})^4 \left(|\partial_t y|^2 + |\partial_t z|^2 + \sum_{i,j=1}^n |\partial_i \partial_j y|^2 + \sum_{i,j=1}^n |\partial_i \partial_j z|^2 \right) \\ \left. + (s\varphi_{\delta,1})^6 (|\nabla y|^2 + |\nabla z|^2) + (s\varphi_{\delta,1})^8 (|y|^2 + |z|^2) \right] e^{2s\psi_{\delta,1}} dx dt \\ \leq C \int_{Q_\delta} (s\varphi_{\delta,1})^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dx dt + C(s)B^2.$$

Let us expand the left-hand side of (52). We have

$$\rho_2^2 \partial_t u(x, t) - \rho_1^2 \partial_t^2 u(x, t) + 2\rho_1 \partial_t \mathcal{A}_1 u(x, t) - \mathcal{A}_1^2 u(x, t) = \tilde{F}(x, t), \quad (x, t) \in Q.$$

Moreover we have

$$\rho_2^2 \nabla \partial_t u(x, t) - \rho_1^2 \nabla \partial_t^2 u(x, t) + 2\rho_1 \nabla \partial_t \mathcal{A}_1 u(x, t) - \nabla \mathcal{A}_1^2 u(x, t) = \nabla \tilde{F}(x, t), \quad (x, t) \in Q.$$

In particular at $t = t_0$, we have

$$(58) \quad \rho_2^2 \partial_t u(x, t_0) - \rho_1^2 \partial_t^2 u(x, t_0) + 2\rho_1 \partial_t \mathcal{A}_1 u(x, t_0) - \mathcal{A}_1^2 u(x, t_0) = \tilde{F}(x, t_0), \quad x \in \Omega,$$

and

$$(59) \quad \rho_2^2 \nabla \partial_t u(x, t_0) - \rho_1^2 \nabla \partial_t^2 u(x, t_0) + 2\rho_1 \nabla \partial_t \mathcal{A}_1 u(x, t_0) - \nabla \mathcal{A}_1^2 u(x, t_0) = \nabla \tilde{F}(x, t_0), \quad x \in \Omega.$$

Taking the weighted L^2 norm of (58) and (59) in Ω , we obtain

$$(60) \quad \int_{\Omega} \left(|\tilde{F}(x, t_0)|^2 + |\nabla \tilde{F}(x, t_0)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx \\ \leq C \sum_{k=1}^6 \tilde{J}_k + C \int_{\Omega} \sum_{|\alpha| \leq 5} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx,$$

where

$$\tilde{J}_1 = \int_{\Omega} |\partial_t u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \quad \tilde{J}_2 = \int_{\Omega} |\partial_t^2 u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \\ \tilde{J}_3 = \int_{\Omega} |\partial_t \mathcal{A}_1 u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \quad \tilde{J}_4 = \int_{\Omega} |\nabla \partial_t u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \\ \tilde{J}_5 = \int_{\Omega} |\nabla \partial_t^2 u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx, \quad \tilde{J}_6 = \int_{\Omega} |\nabla \partial_t \mathcal{A}_1 u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx.$$

Henceforth we estimate \tilde{J}_1 through \tilde{J}_6 by using the Carleman estimate. We assume that $s > 1$ is large enough to satisfy $s\varphi_{\delta,1} > 1$ in Q . We note that $\partial_t \psi_{\delta,1}(x, t) =$

$(e^{2\lambda(\|d_1\|_{C(\bar{\Omega})} - d_1(x))} - e^{-\lambda d_1(x)})(T - 2t)\varphi_{\delta,1}^2(x, t)$ for $(x, t) \in Q$.

$$\begin{aligned}\tilde{J}_1 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|y|^2 e^{2s\psi_{\delta,1}}) \, dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\partial_t y| |y| + s\varphi_{\delta,1}^2 |y|^2) e^{2s\psi_{\delta,1}} \, dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|y|^2 + |z|^2) e^{2s\psi_{\delta,1}} \, dxdt.\end{aligned}$$

Combining this with (57), we may estimate the right-hand side of the above inequality and we obtain

$$(61) \quad \tilde{J}_1 \leq \frac{C}{s^5} \int_{Q_\delta} \varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} \, dxdt + C(s)B^2.$$

Similarly, we obtain

$$\begin{aligned}\tilde{J}_2 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|\partial_t y|^2 e^{2s\psi_{\delta,1}}) \, dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\partial_t^2 y| |\partial_t y| + s\varphi_{\delta,1}^2 |\partial_t y|^2) e^{2s\psi_{\delta,1}} \, dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|\partial_t y|^2 + |\partial_t z|^2) e^{2s\psi_{\delta,1}} \, dxdt.\end{aligned}$$

Putting this together with (57), we see that

$$(62) \quad \tilde{J}_2 \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} \, dxdt + C(s)B^2.$$

Moreover we have

$$\begin{aligned}\tilde{J}_3 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|\mathcal{A}_1 y|^2 e^{2s\psi_{\delta,1}}) \, dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\mathcal{A}_1 \partial_t y| |\mathcal{A}_1 y| + s\varphi_{\delta,1}^2 |\mathcal{A}_1 y|^2) e^{2s\psi_{\delta,1}} \, dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|\mathcal{A}_1 y|^2 + |\mathcal{A}_1 z|^2) e^{2s\psi_{\delta,1}} \, dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 \left(\sum_{|\alpha| \leq 2} |\partial_x^\alpha y|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha z|^2 \right) e^{2s\psi_{\delta,1}} \, dxdt\end{aligned}$$

This together with (57) gives

$$(63) \quad \tilde{J}_3 \leq \frac{C}{s} \int_{Q_\delta} \varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} \, dxdt + C(s)B^2.$$

We have

$$\begin{aligned}\tilde{J}_4 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|\nabla y|^2 e^{2s\psi_{\delta,1}}) \, dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\nabla \partial_t y| |\nabla y| + s\varphi_{\delta,1}^2 |\nabla y|^2) e^{2s\psi_{\delta,1}} \, dxdt\end{aligned}$$

$$\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|\nabla y|^2 + |\nabla z|^2) e^{2s\psi_{\delta,1}} dxdt.$$

Combining this with (57), we may estimate the right-hand side of the above inequality and we obtain

$$(64) \quad \tilde{J}_4 \leq \frac{C}{s^3} \int_{Q_\delta} \varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dxdt + C(s)B^2.$$

Similarly, we obtain

$$\begin{aligned} \tilde{J}_5 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|\nabla \partial_t y|^2 e^{2s\psi_{\delta,1}}) dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\nabla \partial_t^2 y| |\nabla \partial_t y| + s\varphi_{\delta,1}^2 |\nabla \partial_t y|^2) e^{2s\psi_{\delta,1}} dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|\nabla \partial_t y|^2 + |\nabla \partial_t z|^2) e^{2s\psi_{\delta,1}} dxdt. \end{aligned}$$

Putting this together with (57), we see that

$$(65) \quad \tilde{J}_5 \leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dxdt + C(s)B^2.$$

Moreover we have

$$\begin{aligned} \tilde{J}_6 &= \int_{t_0-\delta}^{t_0} \int_{\Omega} \partial_t (|\nabla \mathcal{A}_1 y|^2 e^{2s\psi_{\delta,1}}) dxdt \\ &\leq C \int_{t_0-\delta}^{t_0} \int_{\Omega} (|\nabla \mathcal{A}_1 \partial_t y| |\nabla \mathcal{A}_1 y| + s\varphi_{\delta,1}^2 |\nabla \mathcal{A}_1 y|^2) e^{2s\psi_{\delta,1}} dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 (|\nabla \mathcal{A}_1 y|^2 + |\nabla \mathcal{A}_1 z|^2) e^{2s\psi_{\delta,1}} dxdt \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 \left(|\nabla \partial_t y|^2 + |\nabla (\rho_1 \partial_t - \mathcal{A}_1) y|^2 \right. \\ &\quad \left. + |\nabla \partial_t z|^2 + |\nabla (\rho_1 \partial_t - \mathcal{A}_1) z|^2 \right) e^{2s\psi_{\delta,1}} dxdt \end{aligned}$$

This together with (57) gives

$$(66) \quad \tilde{J}_6 \leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dxdt + C(s)B^2.$$

Summing up the estimate of (60) through (66), we have

$$(67) \quad \begin{aligned} &\int_{\Omega} \left(|\tilde{F}(x, t_0)|^2 + |\nabla \tilde{F}(x, t_0)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx \\ &\leq C \int_{Q_\delta} s\varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^\alpha a|^2 \right) e^{2s\psi_{\delta,1}} dxdt \\ &\quad + C \int_{\Omega} \sum_{|\alpha| \leq 5} |\partial_x^\alpha u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2. \end{aligned}$$

Let us estimate the left-hand side of the inequality (67) from below. By (53) at $t = t_0$, we have

$$\begin{aligned}
(68) \quad & a_1(x) \nabla r(x, t_0) \cdot \nabla \Delta a(x) \\
&= \tilde{F}(x, t_0) - 2a_1(x) \sum_{i,j=1}^n (\partial_i \partial_j r(x, t_0)) (\partial_i \partial_j a(x)) \\
&\quad - a_1(x) \Delta r(x, t_0) \Delta a(x) - (\nabla a_1(x) - \mathbf{b}(x)) \cdot (\nabla r(x, t_0) \cdot \nabla) \nabla a(x) \\
&\quad - \left[(\rho_2 \partial_t^{\frac{1}{2}} - \rho_1 \partial_t) \nabla r(x, t_0) + 3a_1(x) \nabla \Delta r(x, t_0) + (\Delta r(x, t_0)) \nabla a_1(x) \right. \\
&\quad \left. - (\Delta r(x, t_0)) \mathbf{b}(x) - c(x) \nabla r(x, t_0) + \frac{\rho_2 \nabla r(x, 0)}{\sqrt{\pi t_0}} \right] \cdot \nabla a(x) \\
&\quad - (\nabla a_1(x) - \mathbf{b}(x)) \cdot (\nabla a(x) \cdot \nabla) \nabla r(x, t_0) \\
&\quad - \left[(\rho_2 \partial_t^{\frac{1}{2}} - \rho_1 \partial_t) \Delta r(x, t_0) + (\nabla a_1(x) \cdot \nabla \Delta r(x, t_0)) + a_1(x) \Delta^2 r(x, t_0) \right. \\
&\quad \left. - (\mathbf{b}(x) \cdot \nabla \Delta r(x, t_0)) - c(x) \Delta r(x, t_0) + \frac{\rho_2 \Delta r(x, 0)}{\sqrt{\pi t_0}} \right] a(x), \quad x \in \Omega.
\end{aligned}$$

Note that

$$|\nabla r(x, t_0) \cdot \nabla d_1(x)| \geq m_1 > 0, \quad x \in \bar{\Omega},$$

and $a \in H^4(\Omega)$ satisfies $a(x) = 0$, $x \in D$. Let us apply the Lemma 4.8 to (68) in Ω . Then we obtain

$$\begin{aligned}
& \int_{\Omega} \left[s\varphi_{\delta,1}(x, t_0) \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k a(x)|^2 + (s\varphi_{\delta,1}(x, t_0))^2 |\nabla \Delta a(x)|^2 \right. \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^3 \sum_{i,j=1}^n |\partial_i \partial_j a(x)|^2 \right. \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^5 (|\nabla a(x)|^2 + |a(x)|^2) \right] e^{2s\psi_{\delta,1}(x, t_0)} dx \\
& \leq \int_{\Omega} \left(|\tilde{F}(x, t_0)|^2 + |\nabla \tilde{F}(x, t_0)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx + \int_{\Omega} \sum_{|\alpha| \leq 3} |\partial_x^\alpha a(x)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx
\end{aligned}$$

Taking sufficiently large $s > 0$, we may absorb the second term on the right-hand side of the above inequality and we get

$$\begin{aligned}
& \int_{\Omega} \left[s\varphi_{\delta,1}(x, t_0) \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k a(x)|^2 + (s\varphi_{\delta,1}(x, t_0))^2 |\nabla \Delta a(x)|^2 \right. \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^3 \sum_{i,j=1}^n |\partial_i \partial_j a(x)|^2 \right. \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^5 (|\nabla a(x)|^2 + |a(x)|^2) \right] e^{2s\psi_{\delta,1}(x, t_0)} dx \\
& \leq \int_{\Omega} \left(|\tilde{F}(x, t_0)|^2 + |\nabla \tilde{F}(x, t_0)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx
\end{aligned}$$

Combining this with (67), we obtain

$$\begin{aligned}
(69) \quad & \int_{\Omega} \left[s\varphi_{\delta,1}(x, t_0) \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k a(x)|^2 + (s\varphi_{\delta,1}(x, t_0))^2 |\nabla \Delta a(x)|^2 \right. \\
& \quad + (s\varphi_{\delta,1}(x, t_0))^3 \sum_{i,j=1}^n |\partial_i \partial_j a(x)|^2 \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^5 (|\nabla a(x)|^2 + |a(x)|^2) \right] e^{2s\psi_{\delta,1}(x, t_0)} dx \\
& \leq C \int_{Q_{\delta}} s\varphi_{\delta,1}^2 \left(|\nabla \Delta a|^2 + \sum_{|\alpha| \leq 2} |\partial_x^{\alpha} a|^2 \right) e^{2s\psi_{\delta,1}} dx dt \\
& \quad + C \int_{\Omega} \sum_{|\alpha| \leq 5} |\partial_x^{\alpha} u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2. \\
& \leq C \int_{\Omega} s\varphi_{\delta,1}^2(x, t_0) \left(|\nabla \Delta a(x)|^2 + \sum_{|\alpha| \leq 2} |\partial_x^{\alpha} a(x)|^2 \right) e^{2s\psi_{\delta,1}(x, t_0)} dx \\
& \quad + C \int_{\Omega} \sum_{|\alpha| \leq 5} |\partial_x^{\alpha} u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2.
\end{aligned}$$

In the last inequality, we used the fact that

$$\varphi_{\delta,1}^2(x, t) e^{2s\psi_{\delta,1}(x, t)} \leq \varphi_{\delta,1}^2(x, t_0) e^{2s\psi_{\delta,1}(x, t_0)}, \quad (x, t) \in Q_{\delta}$$

for large $s > 0$.

Choose sufficiently large $s > 0$ and absorb the first term on the right-hand side of (69) into the left-hand side. Then we obtain

$$\begin{aligned}
(70) \quad & \int_{\Omega} \left[s\varphi_{\delta,1}(x, t_0) \sum_{i,j,k=1}^n |\partial_i \partial_j \partial_k a(x)|^2 + (s\varphi_{\delta,1}(x, t_0))^2 |\nabla \Delta a(x)|^2 \right. \\
& \quad + (s\varphi_{\delta,1}(x, t_0))^3 \sum_{i,j=1}^n |\partial_i \partial_j a(x)|^2 \\
& \quad \left. + (s\varphi_{\delta,1}(x, t_0))^5 (|\nabla a(x)|^2 + |a(x)|^2) \right] e^{2s\psi_{\delta,1}(x, t_0)} dx \\
& \leq C \int_{\Omega} \sum_{|\alpha| \leq 5} |\partial_x^{\alpha} u(x, t_0)|^2 e^{2s\psi_{\delta,1}(x, t_0)} dx + C(s)B^2.
\end{aligned}$$

Fix $s > 0$. Noting that $\varphi_{\delta,1}(\cdot, t_0) e^{2s\psi_{\delta,1}(\cdot, t_0)}$ has its upper and lower bound in $\overline{\Omega}$, we see that

$$\|a\|_{H^3(\Omega)} \leq C \|u(\cdot, t_0)\|_{H^5(\Omega)} + CB.$$

Thus we obtain the stability estimate (19). \square

Proof of Theorem 3.4. We may prove Theorem 3.4 in the same way as Theorem 3.3. \square

ACKNOWLEDGMENTS

The authors acknowledge support from the Japan Society for the Promotion of Science (JSPS) A3 foresight program: Modeling and Computation of Applied Inverse Problems. MM also acknowledges support from Grant-in-Aid for Scientific Research (17K05572 and 17H02081) of JSPS.

REFERENCES

- [1] B. Amaziane, A. Bourgeat, M. Goncharenko and L. Pankratov, Characterization of the flow for a single fluid in an excavation damaged zone *C. R. Mecanique* **332** (2004) 79–84.
- [2] B. Amaziane, L. Pankratov and A. Piatnitski, Homogenization of a single-phase flow through a porous medium in a thin layer, *Math. Models Methods Appl. Sci.* **17** (2007) 1317–1349.
- [3] A. Ashyralyev, Well-posedness of the Basset problem in spaces of smooth functions, *Appl. Math. Lett.* **24** (2011) 1176–1180.
- [4] A. B. Basset, On the descent of a sphere in a viscous liquid, *Q. J. Math.* **42** (1910) 369–381.
- [5] E. Bazhlekova and I. Dimovski, Exact solution of two-term time-fractional Thornley’s problem by operational method, *Integral Transforms Spec. Funct.* **25** (2014) 61–74.
- [6] C. E. Brennen, *Fundamentals of Multiphase Flow* (Cambridge University Press, 2005).
- [7] A. L. Bukhgeim and M. V. Klivanov, Uniqueness in the large of a class of multidimensional inverse problems, *Dokl. Akad. Nauk SSSR*, **260** (1981) 269–272 (in Russian).
- [8] M. Caputo, Linear model of dissipation whose Q is almost frequency independent-II, *Geophys. J. R. Astr. Soc.*, **13** 529–539
- [9] T. Carleman, Sur un problème d’unicité pour les systèmes d’équations aux dérivées partielles à deux variables indépendantes, *Ark. Mat. Astr. Fys.* **26B** (1939) 1–9.
- [10] A. V. Fursikov and O. Y. Imanuvilov, Controllability of Evolution Equations, (Lecture Notes Series vol. 34) Seoul National University, Seoul (Korea), 1996.
- [11] O. Y. Imanuvilov, Controllability of parabolic equations, *Sbornik Math.*, **186** (1995) 879–900.
- [12] O. Y. Imanuvilov and M. Yamamoto, Lipschitz stability in inverse parabolic problems by the Carleman estimate, *Inverse Problems* **14** (1998) 1229–1245.
- [13] V. Isakov, *Inverse Problems for Partial Differential Equations*, 2nd ed., (Springer-Verlag, Berlin, 2006).
- [14] D. Jiang, Z. Li, Y. Liu and M. Yamamoto, Weak unique continuation property and a related inverse source problem for time-fractional diffusion-advection equations, *Inverse Problems*, **33** (2017) 055013.
- [15] A. Kawamoto, Lipschitz stability estimates in inverse source problems for a fractional diffusion equation of half order in time by Carleman estimates, *J. Inverse Ill-Posed Probl.*, **26** (2018) 647–672.
- [16] A. Kawamoto, Hölder stability estimate in an inverse source problem for a first and half order time fractional diffusion equation, *Inverse Probl. Imaging*, **12** (2018) 315–330.
- [17] M. V. Klivanov, Inverse problems in the “large” and Carleman bounds *Diff. Eq.* **20** (1984) 755–60.
- [18] M. V. Klivanov, Inverse problems and Carleman estimates, *Inverse Problems*, **8** (1992) 575–596.
- [19] M. V. Klivanov, Carleman estimates for global uniqueness, stability and numerical methods for coefficient inverse problems, *J. Inverse Ill-Posed Probl.*, **21** (2013) 477–560.
- [20] M. V. Klivanov and A. A. Timonov, *Carleman Estimates for Coefficient Inverse Problems and Numerical Applications* (VSP, Utrecht, 2004).
- [21] G. P. Langlois, M. Farazmand, and G. Haller, Asymptotic dynamics of inertial particles with memory, *J. Nonlinear. Sci.* **25** (2015) 1225–1255.
- [22] Z. Li, X. Huang, and M. Yamamoto, Carleman estimates for the time-fractional advection-diffusion equations and applications, arXiv preprint arXiv:1704.06011.

- [23] Z. Li, Y. Liu, and M. Yamamoto, Initial-boundary value problems for multi-term time-fractional diffusion equations with positive constant coefficients, *Appl. Math. Comput.* **257** (2015) 381–397.
- [24] Z. Li, O. Y. Imanuvilov, and M. Yamamoto, Uniqueness in inverse boundary value problems for fractional diffusion equations, *Inverse Problems* **32** (2016) 015004.
- [25] Z. Li and M. Yamamoto, Uniqueness for inverse problems of determining orders of multi-term time-fractional derivatives of diffusion equation, *Appl. Anal.*, **94** (2015) 570–579.
- [26] Lin C-L and Nakamura G 2018 Unique continuation property for multi-terms time fractional diffusion equations *Mathematische Annalen* 1–24
- [27] Y. Liu, Strong maximum principle for multi-term time-fractional diffusion equations and its application to an inverse source problem, *Comput. Math. Appl.*, **73** (2017) 96–108.
- [28] Y. Luchko, Maximum principle for the generalized time-fractional diffusion equation, *J. Math. Anal. Appl.* **351** (2009) 218–223.
- [29] Y. Luchko, Initial-boundary-value problems for the generalized multi-term time-fractional diffusion equation, *J. Math. Anal. Appl.* **374** (2011) 538–548.
- [30] M. R. Maxey and J. J. Riley, Equation of motion for a small rigid sphere in a nonuniform flow, *Phys. Fluids* **26** (1983) 883–889.
- [31] I. Podlubny, *Fractional Differential Equations*, Academic Press, New York, 1999.
- [32] C. Ren and X. Xu, Local stability for an inverse coefficient problem of a fractional diffusion equation, *Chin. Ann. Math. Ser. B*, **35** (2014), no. 3, 429–446.
- [33] K. Sakamoto and M. Yamamoto, Inverse source problem with a final overdetermination for a fractional diffusion equation, *Math. Control Relat. Fields*, **1** (2011) 509–518.
- [34] X. Xu, J. Cheng and M. Yamamoto, Carleman estimate for fractional diffusion equation with half order and application, *Appl. Anal.*, **90** (2011) 1355–1371.
- [35] M. Yamamoto, Carleman estimates for parabolic equations and applications, *Inverse Problems* **25** (2009) 123013.
- [36] M. Yamamoto and Y. Zhang, Conditional stability in determining a zeroth-order coefficient in a half-order fractional diffusion equation by a Carleman estimate, *Inverse Problems*, **28** (2012) 105010.

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, KOMABA, MEGURO, TOKYO 153-8914, JAPAN

E-mail address: kawamo@ms.u-tokyo.ac.jp

INSTITUTE FOR MEDICAL PHOTONICS RESEARCH, HAMAMATSU UNIVERSITY SCHOOL OF MEDICINE, HAMAMATSU, SHIZUOKA 431-3192, JAPAN

E-mail address: machida@hama-med.ac.jp