

On bound states in Quantum Field Theory

Changyong Liu*

College of Science, Northwest A&F University, Yangling, Shaanxi 712100, China

(Dated: August 14, 2019)

In this paper, we give a new expression to describe the excited spectrum of positronium (e^+e^-) bound state in Quantum Electrodynamics (QED) and the meson in Quantum Chromodynamics (QCD) separately. As we know, the perturbative series for the bound state which is coming from sum of all Feynman diagrams has zero radius of convergence. The new expression has a nonperturbative term which cannot be obtained from the perturbative expansion in a small coupling constant. In this new expression, we can connect the quantum number n of the excited spectrum with the topological winding number. We point out that the physical picture of bound state can be described by dual fields coming from electric - magnetic transformation.

PACS numbers: 11.10.St, 11.15.Bt, 12.20.-m, 12.38.Aw

keywords: Bound States, Positronium, The excited spectrum, Electric - magnetic transformation

I. INTRODUCTION

The perturbative expansion plays an important role in Quantum Field Theory (QFT), but it is difficult to solve the non-perturbative aspects of QFT. Dyson pointed out that the perturbative series has zero radius of convergence in QFT [1]. We need a new method to study the non-perturbative problem especially the bound state which is coming from sum of all Feynman diagrams. Bound states are important objects in quantum theory and have been widely studied in quantum field theory. Although we haven't solve the bound states problem, there are many methods that is aiming to solve it completely, e.g. Bethe-Salpeter equation [2-4] and Lattice QCD [5, 6]. In this paper, we give a new expression to describe the excited spectrum of bound state. We first give a general framework of our method and then discuss the positronium (e^+e^-) in QED and mesons in QCD separately. We end with the conclusions.

II. A GENERAL FRAMEWORK

In order to study the bound states, we define the integral of meromorphic function $f[z]$ along a smooth contour $C[a, b]$ in complex plane.




FIG. 1 – The $n = 3$ case. A smooth contour $C[a, b]$ in complex plane starting from a to b . The red dots denote poles of meromorphic function $f[z]$.

The contour $C[a, b]$ contain n poles of $f[z]$ labeled by z_i , $i = 1, \dots, n$. We suppose $z_i \neq a, b$, for $i = 1, \dots, n$ (see figure 1). Then the integral of $f[z]$ along contour $C[a, b]$ can be calculated as

$$\int_{C[a, b]} f[z] dz = \int_a^b f[z] dz + \sum_{i=1}^n 2\pi i n_i \text{Res} f[z_i]. \quad (1)$$

Where the $\text{Res} f[z_i]$ is the residue of $f[z]$ at z_i . The $\int_a^b f[z] dz$ is the directly calculation which depends only on the boundary value a and b . The winding number n_i is integer Z which is coming from the contour circling n_i times around the pole z_i .


When we study the mesons in QCD, we find the other case that a or b is one of the poles of $f[z]$ that is infrared divergence in quantum field theory. We can give a small mass μ to gauge field to regulate this divergence. There are many good book which discuss the infrared divergence [7, 8]. For recently progress on this topic, we can see the review paper [9].

III. POSITRONIUM (e^+e^-) SYSTEMS IN QED

To study the positronium (e^+e^-) bound states, we review some results of QED which can be found in text book [10, 11]. We define $i\Pi^{\mu\nu}(q)$ to be the sum of all 1-particle-irreducible (1PI) insertions into the photon propagator. The $\Pi^{\mu\nu}(q)$ has the tensor structure:

$$\Pi^{\mu\nu}(q) = (q^2 g^{\mu\nu} - q^\mu q^\nu) \Pi(q^2). \quad (2)$$

The α contribution to $i\Pi^{\mu\nu}(q)$ is $i\Pi_2^{\mu\nu}(q) = i(q^2 g^{\mu\nu} - q^\mu q^\nu) \Pi_2(q^2)$ which is the following electron loop:



Where the fine-structure constant is $\alpha = \frac{e^2}{4\pi} \approx \frac{1}{137}$.

The excited spectrum is the pole of the two-point function, the exact photon two-point function is

* Email: liuchangyong@nwsuaf.edu.cn

$$G_{\mu\nu}(q^2) = \text{diagram} = \frac{-ig_{\mu\nu}}{q^2(1-\Pi(q^2))}$$

The two-point function has ultraviolet divergences. To remove the divergences, we need renormalization. The renormalization method we choose is that we make the replacement

$$\tilde{G}_{\mu\nu}(q^2) = \frac{-ig_{\mu\nu}}{q^2(1-[\Pi(q^2) - \Pi(0)])}. \quad (3)$$

In this paper, we only consider the α contribution to $i\Pi^{\mu\nu}(q)$ that is

$$\begin{aligned} \hat{\Pi}_2(q^2) &\equiv \Pi_2(q^2) - \Pi_2(0) \\ &= -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \log\left(\frac{m_e^2}{m_e^2 - x(1-x)q^2}\right) \\ &= -\frac{2\alpha}{\pi} \int_0^1 dx x(1-x) \log\left(\frac{1}{1-x(1-x)t^2}\right) \\ &= \frac{2\alpha}{\pi} \int_0^1 dx \left(\frac{x^2}{2} - \frac{x^3}{3}\right) \frac{(1-2x)t^2}{1-x(1-x)t^2}. \end{aligned}$$

Where we have defined the $t = \sqrt{\frac{q^2}{m_e^2}}$. When $t > 2$, this is the two or more free particles states which give a branch cut. In this case, we have two poles that is $x_{1,2} = \frac{1}{2} \pm \frac{1}{2}\sqrt{1 - \frac{4}{t^2}}$. Using the formula (1), we obtain

$$\begin{aligned} \hat{\Pi}_2(q^2) &= \\ &= \frac{\alpha(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4)\arctan[\frac{t}{\sqrt{4-t^2}}])}{9\pi t^3\sqrt{4-t^2}} \\ &- \frac{i\alpha}{3}[(n-m) + \sqrt{1-\frac{4}{t^2}}(1+\frac{2}{t^2})(n+m)]. \end{aligned} \quad (4)$$

Where n, m is integer Z . We consider the positronium (e^+e^-) bound states which is $t < 2$. We use matching methods to calculate the positronium (e^+e^-) bound states. We use analytical continuation of (4) and make replacement

$$\alpha \rightarrow \frac{c}{\alpha} \quad (5)$$

in the last term of (4). Where the constant c is $c \approx 2.00408$ for positronium (e^+e^-) bound states which can be decided by the ground state energy of (7). We get the result

$$\begin{aligned} \hat{\Pi}_2(q^2) &= \\ &= \frac{\alpha(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4)\arctan[\frac{t}{\sqrt{4-t^2}}])}{9\pi t^3\sqrt{4-t^2}} \\ &+ \frac{c}{3\alpha}[-i(n-m) + \sqrt{\frac{4}{t^2}-1}(1+\frac{2}{t^2})(n+m)]. \end{aligned}$$

We consider the long-lived bound states, so we set the $n = m$. The excited spectrum of positronium is the pole of two-point function that is solution of

$$1 - \hat{\Pi}_2(q^2) = 0$$

$$1 + \frac{\alpha(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4)\arctan[\frac{t}{\sqrt{4-t^2}}])}{9\pi t^3\sqrt{4-t^2}} - \frac{2cn}{3\alpha}[\sqrt{\frac{4}{t^2}-1}(1+\frac{2}{t^2})] = 0. \quad (6)$$

Where the quantum number n is natural number N . The solution is consistent with the known result of positronium in high precision. The excited spectrum of positronium to order α^2 is [11]

$$E = 2m_e - m_e \frac{\alpha^2}{4n^2}. \quad (7)$$

To illustrate this, we compare the $t - 2$ of the solution of (6) with the result of (7) in Table I.

n	t-2	$-\frac{\alpha^2}{4n^2}$
1	-1.33128×10^{-5}	-1.33128×10^{-5}
2	-3.32830×10^{-6}	-3.32821×10^{-6}
3	-1.47926×10^{-6}	-1.47920×10^{-6}
4	-8.32084×10^{-7}	-8.32052×10^{-7}
5	-5.32534×10^{-7}	-5.32514×10^{-7}

TABLE I – The t is the solution of (6) with $\alpha = \frac{1}{137.03599913}$.

To obtain the equation (6), we have made replacement of (5) which is the nonperturbative effect of bound states coming from sum of all Feynman diagrams. The physical picture of positronium (e^+e^-) bound state can be equivalent to the system of two magnetic monopoles in leading order term FIG. 2. This is also a kind of electric - magnetic transformation. For some particular quantum field, the electric - magnetic duality play an important role[12-14].




FIG. 2 – The constituents of positronium (e^+e^-) bound state include two magnetic monopoles in leading order approximation.

IV. MESONS PROPERTY IN QCD

The calculation of mesons in QCD is the same as the positronium case which we use the first method (6).




FIG. 3 – The gluon propagator in order g^2 .

The $i\Pi_2^{\mu a, \nu b}(q)$ have the following structure $i\Pi_2^{\mu a, \nu b}(q) = i(q^2 g^{\mu\nu} - q^\mu q^\nu) \delta^{ab} \Pi_2(q^2)$.

Where $\Pi_2(q^2)$ is calculated from FIG. 3 [10]

$$\begin{aligned} \Pi_2(q^2, \mu^2) = & \frac{-2C(r)n_f\alpha_g}{\pi} \int_0^1 dx x(1-x) \frac{\Gamma(2 - \frac{d}{2})}{(m_q^2 - x(1-x)q^2)^{2 - \frac{d}{2}}} \\ & + \frac{C_2(G)\alpha_g}{4\pi} \int_0^1 dx \frac{\Gamma(2 - \frac{d}{2})}{(\mu^2 - x(1-x)q^2)^{2 - \frac{d}{2}}} [(1 - \frac{d}{2}) \\ & (1 - 2x)^2 + 2]. \end{aligned} \quad (8)$$

Here α_g is $\alpha_g = \frac{g^2}{4\pi}$, n_f is the number of fermion species, $C(r)$ is $C(r) = \frac{1}{2}$ for fundamental representations N , and $C_2(G)$ is $C_2(G) = N$ for $SU(N)$. We give a small mass μ to gluon to regulate the infrared divergence. For convenient, we define new variables t and \tilde{t}

$$t = \sqrt{\frac{q^2}{m_q^2}}, \quad \tilde{t} = \sqrt{\frac{q^2}{\mu^2}} = \frac{m_q}{\mu} t. \quad (9)$$

We emphasized that the mass m_q is the free quark mass which is larger than the constituent quark mass in hadrons. The result (8) has ultraviolet divergences, we make the replacement and use the formula (1). We obtain

$$\begin{aligned} \hat{\Pi}_2(q^2, \mu^2) &\equiv \Pi_2(q^2, \mu^2) - \Pi_2(0, \mu^2) \\ &= -\frac{2C(r)n_f\alpha_g}{\pi} \int_0^1 dx x(1-x) \log\left(\frac{1}{1-x(1-x)t^2}\right) \\ &+ \frac{C_2(G)\alpha_g}{4\pi} \int_0^1 dx (1+4x-4x^2) \log\left(\frac{1}{1-x(1-x)\tilde{t}^2}\right) \\ &= -\frac{2C(r)n_f\alpha_g}{\pi} \int_0^1 dx x(1-x) \log\left(\frac{1}{1-x(1-x)t^2}\right) \\ &+ \frac{C_2(G)\alpha_g}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} (4\tilde{t}\sqrt{4-\tilde{t}^2}(6+7\tilde{t}^2) \\ &+ 6(-4+\tilde{t}^2)(4+5\tilde{t}^2) \arctan\left[\frac{\tilde{t}}{\sqrt{4-\tilde{t}^2}}\right]) \\ &+ C_2(G)\alpha_g \frac{i}{6} \left[\frac{5}{2}(k-l) + \sqrt{1-\frac{4}{\tilde{t}^2}} \left(\frac{5}{2} + \frac{2}{\tilde{t}^2}\right) (k+l) \right] \\ &= -\frac{2C(r)n_f\alpha_g}{\pi} \int_0^1 dx x(1-x) \log\left(\frac{1}{1-x(1-x)t^2}\right) \\ &+ Re\left[\frac{C_2(G)\alpha_g}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} (4\tilde{t}\sqrt{4-\tilde{t}^2}(6+7\tilde{t}^2) \right. \\ &+ 6(-4+\tilde{t}^2)(4+5\tilde{t}^2) \arctan\left[\frac{\tilde{t}}{\sqrt{4-\tilde{t}^2}}\right]) \\ &+ C_2(G)\alpha_g \frac{i}{6} \left[\frac{5}{2}(k-l) + \sqrt{1-\frac{4}{\tilde{t}^2}} \left(\frac{5}{2} + \frac{2}{\tilde{t}^2}\right) (k+l) \right]. \end{aligned}$$

Where k, l is integer Z . To discuss the excited spectrum of meson, we make the q^2 to be the range of $4\mu^2 < q^2 <$

$4m_q^2$. Then we obtain

$$\begin{aligned} 1 - \hat{\Pi}_2(q^2) &= 1 + C(r)n_f\alpha_g \\ &\frac{(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4) \arctan\left[\frac{t}{\sqrt{4-t^2}}\right])}{9\pi t^3\sqrt{4-t^2}} \\ &- C(r)n_f \frac{c}{3\alpha_g} \left[-i(n-m) + \sqrt{\frac{4}{t^2} - 1} \left(1 + \frac{2}{t^2}\right) (n+m) \right] \\ &- C_2(G)\alpha_g Re\left[\frac{4\tilde{t}\sqrt{4-\tilde{t}^2}(6+7\tilde{t}^2)}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} \right. \\ &\left. + \frac{6(-4+\tilde{t}^2)(4+5\tilde{t}^2) \arctan\left[\frac{\tilde{t}}{\sqrt{4-\tilde{t}^2}}\right]}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} \right] \\ &- C_2(G)\alpha_g \frac{i}{6} \left[\frac{5}{2}(k-l) + \sqrt{1-\frac{4}{\tilde{t}^2}} \left(\frac{5}{2} + \frac{2}{\tilde{t}^2}\right) (k+l+1) \right]. \end{aligned}$$

Where m, n, k, l is integer Z . The Re is the real part function. As the same as the positronium, we also make replacement $\alpha_g \rightarrow \frac{c}{\alpha_g}$ in the third term. The above expression is different with positronium with the imaginary part which can not be zero. Due to the imaginary part, the excited meson will decay quickly. We make $n = m$ and $k = l = 0$ to make a slightly long lived resonances. The result is

$$\begin{aligned} 1 - \hat{\Pi}_2(q^2) &= 1 + C(r)n_f\alpha_g \\ &\frac{(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4) \arctan\left[\frac{t}{\sqrt{4-t^2}}\right])}{9\pi t^3\sqrt{4-t^2}} \\ &- C(r)n_f \frac{2cn}{3\alpha_g} \left[\sqrt{\frac{4}{t^2} - 1} \left(1 + \frac{2}{t^2}\right) \right] - C_2(G)\alpha_g Re\left[\right. \\ &\left. \frac{4\tilde{t}\sqrt{4-\tilde{t}^2}(6+7\tilde{t}^2) + 6(-4+\tilde{t}^2)(4+5\tilde{t}^2) \arctan\left[\frac{\tilde{t}}{\sqrt{4-\tilde{t}^2}}\right]}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} \right. \\ &\left. - C_2(G)\alpha_g \frac{i}{6} \left[\sqrt{1-\frac{4}{\tilde{t}^2}} \left(\frac{5}{2} + \frac{2}{\tilde{t}^2}\right) \right]. \right. \\ \text{Then the excited spectrum of meson is the solution of equation} & \\ &1 + C(r)n_f\alpha_g \\ &\frac{(t\sqrt{4-t^2}(12+5t^2) + 6(-8-2t^2+t^4) \arctan\left[\frac{t}{\sqrt{4-t^2}}\right])}{9\pi t^3\sqrt{4-t^2}} \\ &- C(r)n_f \frac{2cn}{3\alpha_g} \left[\sqrt{\frac{4}{t^2} - 1} \left(1 + \frac{2}{t^2}\right) \right] - C_2(G)\alpha_g Re\left[\right. \\ &\left. \frac{4\tilde{t}\sqrt{4-\tilde{t}^2}(6+7\tilde{t}^2) + 6(-4+\tilde{t}^2)(4+5\tilde{t}^2) \arctan\left[\frac{\tilde{t}}{\sqrt{4-\tilde{t}^2}}\right]}{36\pi\tilde{t}^3\sqrt{4-\tilde{t}^2}} \right] \\ &= 0. \end{aligned}$$

Where the n is natural number N . We give the picture (FIG. 4 and FIG. 5) of the t^2 as a function of n with $n_f = 6$, $\alpha_g = 10$, $c = 2$, $\frac{m_q}{\mu} = 1000$ and $C_2(G) = 3$ for $SU(3)$.

From the FIG. 4, we find the $t^2 \sim n$ which is consistent with the data [15]. This is also known as the Regge behavior [16]. We conclude that the physical picture of confinement can be described by dual field coming from electric - magnetic transformation. This picture




FIG. 4 – t^2 as a function of n , with $n = 1, \dots, 100$, $\alpha_g = 10$, $c = 2$, $n_f = 6$, and $\frac{m_q}{\mu} = 1000$.




FIG. 5 – t^2 as a function of n , with $n = 1, \dots, 600$, $\alpha_g = 10$, $c = 2$, $n_f = 6$, and $\frac{m_q}{\mu} = 1000$.

may be connected with a dual superconductor picture of confinement[17–20].

V. CONCLUSION

In this paper, we give a new expression to calculate the excited spectrum of bound states in Quantum Field Theory. There is no correctly bound states pole in one-loop Feynman diagrams of the two-point function in traditional method. We use the reasonable definition of integral of $f[z]$ along contour $C[a, b]$ (1) and analytical continuation. We find the bound states pole appear. We discuss the positronium (e^+e^-) in QED and meson in QCD separately. Our results agree with the known ones. To obtain the equation (6), we have make replacement of (5) which is the nonperturbative effect of bound states coming from sum of all Feynman diagrams. The physical picture of bound state can be described by dual field coming from electric - magnetic transformation. For the results is consistent with the experimental data, we need to calculate the higher order loops. There are much related work need to do in the future.

ACKNOWLEDGMENTS

This work is supported by Chinese Universities Scientific Fund Grant No. 2452018158.

REFERENCE

-
- [1] F. J. Dyson, Phys. Rev. **85**, 631 (1952). doi:10.1103/PhysRev.85.631
 - [2] E. E. Salpeter and H. A. Bethe, Phys. Rev. **84**, 1232 (1951). doi:10.1103/PhysRev.84.1232
 - [3] N. Nakanishi, Prog. Theor. Phys. Suppl. **43**, 1 (1969). doi:10.1143/PTPS.43.1
 - [4] G. V. Efimov, Few Body Syst. **33**, 199 (2003) doi:10.1007/s00601-003-0015-1 [hep-ph/0304194].
 - [5] K. G. Wilson, Phys. Rev. D **10**, 2445 (1974). doi:10.1103/PhysRevD.10.2445
 - [6] R. Gupta, hep-lat/9807028.
 - [7] S. Weinberg, “The Quantum theory of fields. Vol. 1: Foundations,”
 - [8] G. F. Sterman, “An Introduction to quantum field theory,”
 - [9] A. Strominger, arXiv:1703.05448 [hep-th].
 - [10] M. E. Peskin and D. V. Schroeder, “An Introduction to quantum field theory,”
 - [11] C. Itzykson and J. B. Zuber, “Quantum Field Theory,” New York, Usa: Mcgraw-hill (1980) 705 P.(International Series In Pure and Applied Physics)
 - [12] P. A. M. Dirac, Proc. Roy. Soc. Lond. A **133**, no. 821, 60 (1931). doi:10.1098/rspa.1931.0130

- [13] C. Montonen and D. I. Olive, Phys. Lett. **72B**, 117 (1977). doi:10.1016/0370-2693(77)90076-4
- [14] N. Seiberg and E. Witten, Nucl. Phys. B **426**, 19 (1994) Erratum: [Nucl. Phys. B **430**, 485 (1994)] doi:10.1016/0550-3213(94)90124-4, 10.1016/0550-3213(94)00449-8 [hep-th/9407087].
- [15] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592**, no. 1-4, 1 (2004). doi:10.1016/j.physletb.2004.06.001
- [16] P. D. B. Collins, “An Introduction to Regge Theory and High-Energy Physics,” doi:10.1017/CBO9780511897603
- [17] H. B. Nielsen and P. Olesen, Nucl. Phys. B **160**, 380 (1979). doi:10.1016/0550-3213(79)90065-8
- [18] J. B. Kogut and L. Susskind, Phys. Rev. D **9**, 3391 (1974). doi:10.1103/PhysRevD.9.3391
- [19] G. 't Hooft, Nucl. Phys. B **138**, 1 (1978). doi:10.1016/0550-3213(78)90153-0
- [20] S. Mandelstam, Phys. Rept. **23**, 307 (1976). doi:10.1016/0370-1573(76)90052-1