

Construction of Multi-Bubble Solutions for a System of Elliptic Equations arising in Rank Two Gauge Theory

Hsin-Yuan Huang

*Department of Applied Mathematics, National Chiao-Tung University,
Hsinchu, Taiwan*

Abstract

We study the existence of multi-bubble solutions for the following skew-symmetric Chern–Simons system

$$\begin{cases} \Delta u_1 + \frac{1}{\varepsilon^2} e^{u_2} (1 - e^{u_1}) = 4\pi \sum_{i=1}^{2k} \delta_{p_{1,i}} \\ \Delta u_2 + \frac{1}{\varepsilon^2} e^{u_1} (1 - e^{u_2}) = 4\pi \sum_{i=1}^{2k} \delta_{p_{2,i}} \end{cases} \quad \text{in } \Omega, \quad (0.1)$$

where $k \geq 1$ and Ω is a flat torus in \mathbb{R}^2 . It continues the joint work with Zhang[20], where we obtained the necessary conditions for the existence of bubbling solutions of Liouville type. Under nearly necessary conditions (see Theorem 1.1), we show that there exist a sequence of solutions $(u_{1,\varepsilon}, u_{2,\varepsilon})$ to (0.1) such that $u_{1,\varepsilon}$ and $u_{2,\varepsilon}$ blow up simultaneously at k points in Ω as $\varepsilon \rightarrow 0$.

1 Introduction

Over the past few decades, various Chern–Simons models have been proposed in the study of condensed matter physics and particle physics, including the relativistic Chern–Simons models of high temperature superconductivity [21, 12], Lozano-Marqués-Moreno-Schaposnik model [32] of bosonic sector of $\mathcal{N} = 2$ supersymmetric Chern–Simons–Higgs theory, Gudnason model [14, 15] of $\mathcal{N} = 2$ supersymmetric Yang–Mills–Chern–Simons–Higgs theory, Aharony–Bergman–Jafferis–Maldacena model [1] and so on. The self-dual solutions of these Chern–Simons models often can be reduced to systems of elliptic partial differential equations with exponential nonlinearities. We refer the readers to the [12, 40, 38] for exhaustive bibliography.

For the Abelian Chern–Simons models, Hong, Kim and Pac [18], and Jackiw and Weinberg [21] considered a model with one Chern–Simons gauge field and constructed selfdual Abelian Chern–Simons–Higgs vortices which describe anyonic solitons in 2+1 dimensions. Speilman et al. [34] observed no parity breaking in an experiment with high temperature superconductivity. In [16] and [39], those authors indicated the parity broken may not happen in the a field theory with even number of Chern–Simons gauge fields. One of the simplest models of this kind is the $U(1) \times U(1)$ Chern–Simons model of two Higgs fields, where each of them coupled to one of

two Chern–Simons fields. In this paper, we will investigate the relativistic self-dual $U(1) \times U(1)$ Chern–Simons model proposed by Kim et al[23]. We give a brief description on this model below.

Consider the Lagrangian action density of the $[U(1)]^2$ Chern–Simons model with only mutual Chern–Simons interaction is given in the form

$$\mathcal{L} = -\frac{\varepsilon}{4} \epsilon^{\mu\nu\alpha} \left(A_\mu^{(1)} F_{\mu\nu}^{(2)} + A_\mu^{(2)} F_{\mu\nu}^{(1)} \right) + \sum_{i=1}^2 D_\mu \phi_i \overline{D^\mu \phi_i} - V(\phi_1, \phi_2), \quad (1.1)$$

where $\varepsilon > 0$ is a coupling parameter, $(A_\mu^{(1)})$ and $(A_\mu^{(2)})$ are two associated Abelian gauge fields with the electromagnetic fields $F_{\mu\nu}^{(i)} = \partial_\mu A_\nu^{(i)} - \partial_\nu A_\mu^{(i)}$, ϕ_1 and ϕ_2 are two Higgs scalar fields with the covariant derivatives $D_\mu \phi_i = \partial_\mu \phi_i - i A_\mu^{(i)} \phi_i$ ($\mu = 0, 1, 2$, $i = 1, 2$), and the Higgs potential $V(\phi_1, \phi_2)$ is taken as

$$V(\phi_1, \phi_2) = \frac{1}{4\varepsilon^2} \left(|\phi_2|^2 [|\phi_1|^2 - 1]^2 + |\phi_1|^2 [|\phi_2|^2 - 1]^2 \right). \quad (1.2)$$

Set $u_{i,\varepsilon} = \ln |\phi_i|^2$, and denote the zeros of ϕ_i by $\{p_{i,1}, \dots, p_{i,N_i}\}$, $i = 1, 2$ as in [22]. After a BPS reduction [2, 33], we find that $(u_{1,\varepsilon}, u_{2,\varepsilon})$ satisfies

$$\begin{cases} \Delta u_1 + \frac{1}{\varepsilon^2} e^{u_2} (1 - e^{u_1}) = 4\pi \sum_{i=1}^{N_1} \delta_{p_{1,i}} \\ \Delta u_2 + \frac{1}{\varepsilon^2} e^{u_1} (1 - e^{u_2}) = 4\pi \sum_{i=1}^{N_2} \delta_{p_{2,i}} \end{cases} \quad \text{in } \Omega, \quad (1.3)$$

where δ_p is the Dirac measure at p . See [23, 13, 24] for the details of the derivation of (1.3). In physical literature, Ω here is usually referred to \mathbb{R}^2 or a flat torus in \mathbb{R}^2 . In this paper, we consider the case of flat torus. We refer the readers to [24, 25, 5, 19, 17, 6] and reference therein for the recent developments.

Set $u_1 \equiv u_2$ and $\{p_i^1\}_{i=1}^{N_1} = \{p_j^2\}_{j=1}^{N_2}$, then the system (1.3) is reduced to the Abelian Chern–Simons equation with one Higgs particle proposed by Kim–Pac[18] and Jackiw–Weinberg[21],

$$\Delta u + \frac{1}{\varepsilon^2} e^u (1 - e^u) = 4\pi \sum_{s=1}^N \delta_{p_s}, \quad (1.4)$$

which has been extensively studied for more than twenty years. We refer the readers to [35, 37, 3, 7, 12, 36, 4, 8, 10, 11] and reference therein for more details.

We introduce the Green function $G(x, p)$ to eliminate the Dirac measure in (1.4). Here, the Green function $G(x, p)$ is a doubly periodic function on $\partial\Omega$ and satisfies

$$-\Delta G(x, p) = \delta_p - \frac{1}{|\Omega|}, \quad (1.5)$$

and $|\Omega|$ is the measure of Ω . Let $u_0(x) = -4\pi \sum_{j=1}^N G(x, p_j)$. With the transformation $u \rightarrow u + u_0$, (1.4) becomes

$$\Delta u + \frac{1}{\varepsilon^2} e^{u+u_0} (1 - e^{u+u_0}) = \frac{4N\pi}{|\Omega|}. \quad (1.6)$$

Choe and Kim[9] showed that, there may have a sequence of solutions to (1.6) satisfying the following:

There is finite set $\{x_{1,\varepsilon}, \dots, x_{l,\varepsilon}\}$, $x_{j,\varepsilon} \in \Omega$, $j = 1, \dots, l$, such that, as $\varepsilon \rightarrow 0$

$$u_\varepsilon(x_{j,\varepsilon}) + 2 \ln \frac{1}{\varepsilon} \rightarrow \infty, \quad j = 1, \dots, l, \quad (1.7)$$

and

$$u_\varepsilon + 2 \ln \frac{1}{\varepsilon} \rightarrow -\infty \text{ uniformly on any compact subset of } \Omega \setminus \{q_1, \dots, q_l\}, \quad (1.8)$$

where $q_j = \lim_{\varepsilon \rightarrow 0} x_{j,\varepsilon}$. Furthermore,

$$\frac{1}{\varepsilon^2} e^{u_\varepsilon + u_0} (1 - e^{u_\varepsilon + u_0}) \rightarrow \sum_{j=1}^l M_j \delta_{q_j} \quad \text{as } \varepsilon \rightarrow 0, \quad M_j \geq 8\pi. \quad (1.9)$$

Solutions of (1.6) satisfying (1.7) and (1.8) are called *bubbling solutions* and q_j is called the blow-up point of the bubbling solution.

We can classify these blow-up points as follows: After suitable rescaling at the blow-up point q_j , the bubbling solution u_ε to (1.6) converges to either the entire solution of

$$\Delta u + |x|^{2m} e^u (1 - |x|^{2m} e^u) = 0, \quad (1.10)$$

or the entire solution of

$$\Delta u + |x|^{2m} e^u = 0. \quad (1.11)$$

Here, $m = 0$ if q_j is not a vortex point and $m = \#\{p_i : p_i = q_j\}$ if q_j is a vortex point. The blow-up point is called Chern-Simons type if the limiting equation is (1.10) and is called mean field type if the limiting equation is (1.11). The existence and non-existence of bubbling solutions of (1.6) have been studied in a series of work of Lin and Yan[28, 29, 30].

For (1.3), we may use the following system to construct the bubbling solutions to (1.3) which blow up at the same point q .

(i)

$$\text{Chern-Simons system: } \begin{cases} \Delta u_1 + |x|^{m_2} e^{u_2} (1 - |x|^{m_1} e^{u_1}) = 0 \\ \Delta u_2 + |x|^{m_1} e^{u_1} (1 - |x|^{m_2} e^{u_2}) = 0 \end{cases} \quad \text{in } \mathbb{R}^2. \quad (1.12)$$

(ii)

$$\text{Liouville system: } \begin{cases} \Delta u_1 + |x|^{m_2} e^{u_2} = 0 \\ \Delta u_2 + |x|^{m_1} e^{u_1} = 0 \end{cases} \quad \text{in } \mathbb{R}^2. \quad (1.13)$$

Here, $m_1 = \#\{p_{1,j} : p_{1,j} = q\}$ and $m_2 = \#\{p_{2,j} : p_{2,j} = q\}$. For the case (i), the blow-up point is called the Chern-Simons type; for the case (ii), the blow-up point is called the Liouville type.

For the existence of the bubbling solutions of Chern-Simons type to (1.3), there are two known results. The first one is due to Lin and Yan[28], in which they assume that $N_1 = N_2$ and one of the vortex points $\{p_{1,i}\}_{i=1, \dots, N_1}$ and $\{p_{1,i}\}_{i=1, \dots, N_1}$ coincide, then they constructed a Chern-Simons type bubbling solution blow up at that vortex point.

Theorem A. [28] *Suppose that $N_1 = N_2 = N > 4$ and $p_{1,1} = p_{2,1} = p_1$ and $p_{1,1} \neq p_{1,j}, p_{2,j}$, $j > 1$. Then, there is an $\varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0)$, (1.3) has a solution $(u_{1,\varepsilon}, u_{2,\varepsilon})$, satisfying*

$$u_{1,\varepsilon} - u_{2,\varepsilon} \rightarrow 4\pi \sum_{j=2}^N G(p_{1,1}, p_{1,j}) - 4\pi \sum_{j=2}^N G(p_{2,1}, p_{2,j}) \quad (1.14)$$

and

$$\frac{1}{\varepsilon^2} e^{u_{i,\varepsilon}} (1 - e^{u_{j,\varepsilon}}) \rightarrow 4\pi N \delta_{p_1}, \quad i \neq j, \quad i, j \in \{1, 2\} \quad (1.15)$$

as $\varepsilon \rightarrow 0$.

The idea in the proof of Theorem A is using the ansatz

$$u_1 - u_2 \approx 4\pi \sum_{j=2}^N G(p_{1,1}, p_{1,j}) - 4\pi \sum_{j=2}^N G(p_{2,1}, p_{2,j})$$

to reduce the system(1.3) to a single equation.

In a joint work with Han and Lin[17], we proved the existence of Chern-Simons type bubbling solutions when the blow-up point is not vortex point.

Theorem B. [17] *Suppose $(N_1 - 1)(N_2 - 1) > 1$, and q satisfies*

$$D(N_1 u_{2,0} + N_2 u_{1,0})(q) = 0 \quad (1.16)$$

and

$$\deg(D(N_1 u_{2,0} + N_2 u_{1,0}), q) \neq 0. \quad (1.17)$$

Then the system (1.3) admits Chern-Simons type bubbling solutions $(u_{1,\varepsilon}, u_{2,\varepsilon})$ blowing up at q .

Theorem A and B only discussed the one blow-up case. However, the problem on the existence of multi-bubble solutions of (1.3) is still open. The analysis of linearized system to (1.12) and (1.13) is not fully understood when $(m_1, m_2) \neq (0, 0)$. Thus, we only investigate the issue on the existence of the *bubbling solutions of Liouville type* to (1.3) whose blow up points are regular points. We give more precise description on this type bubbling solutions below.

Let $(u_{1,\varepsilon}, u_{2,\varepsilon})$ be a sequence of solutions to (1.3) blowing up at $\{q_1, \dots, q_k\}$ and $q_i \notin \{p_{1,i}, p_{2,j}\}_{1 \leq i \leq N_1, 1 \leq j \leq N_2}$. For a small constant $d > 0$, we define the local mass of $(u_{1,\varepsilon}, u_{2,\varepsilon})$ at q_i :

$$(m_{1,i,\varepsilon}, m_{2,i,\varepsilon}) = \left(\frac{1}{\varepsilon^2} \int_{B_d(q_i)} e^{u_{2,\varepsilon}} (1 - e^{u_{1,\varepsilon}}), \frac{1}{\varepsilon^2} \int_{B_d(q_i)} e^{u_{1,\varepsilon}} (1 - e^{u_{2,\varepsilon}}) \right).$$

We assume that there exist $\{x_{1,j,\varepsilon}\}_{j=1}^k$ and $\{x_{2,j,\varepsilon}\}_{j=1}^k$ such that

$$(a_1) \quad u_{i,\varepsilon}(x_{i,j,\varepsilon}) + 2 \ln \frac{1}{\varepsilon} \rightarrow +\infty \text{ as } \varepsilon \rightarrow 0, \quad i = 1, 2, \quad j = 1, \dots, k.$$

$$(a_2) \quad u_{i,\varepsilon}(x) + 2 \ln \frac{1}{\varepsilon} \rightarrow -\infty \text{ as } \varepsilon \rightarrow 0 \text{ uniformly on any compact set of } \Omega \setminus \{q_1, \dots, q_k\}, \text{ where } q_j = \lim_{\varepsilon \rightarrow 0} x_{1,j,\varepsilon} = \lim_{\varepsilon \rightarrow 0} x_{2,j,\varepsilon}, \quad j = 1, \dots, k.$$

$$(a_3) \quad \beta_{j,\varepsilon} = \max\{u_{1,\varepsilon}(x_{1,j,\varepsilon}), u_{2,\varepsilon}(x_{2,j,\varepsilon})\} \rightarrow -\infty \text{ as } \varepsilon \rightarrow 0.$$

$$(a_4) \quad |u_{1,j,\varepsilon}(x_{1,j,\varepsilon}) - u_{2,\varepsilon}(x_{2,j,\varepsilon})| = O(1)$$

$$(a_5) \quad |x_{i,j,\varepsilon} - q_j| < C\varepsilon e^{-\frac{1}{2}\beta_{j,\varepsilon}} \text{ for some constant } C > 0.$$

The solutions $(u_{1,\varepsilon}, u_{2,\varepsilon})$ satisfying $(a_1) - (a_5)$ are called *fully bubbling solutions of Liouville type*.

Let $\mu_{j,\varepsilon} = \varepsilon e^{-\frac{1}{2}\beta_{j,\varepsilon}}$ and assume $\max\{u_{1,\varepsilon}(x), u_{2,\varepsilon}(x)\}$ attains its maximum at $x_{j,\varepsilon}$ for x near q_j . Formally,

$$(\tilde{u}_1(y), \tilde{u}_2(y)) = (u_{1,\varepsilon}(\mu_{j,\varepsilon}y + x_{j,\varepsilon}) - \beta_{j,\varepsilon}, u_{2,\varepsilon}(\mu_{j,\varepsilon}y + x_{j,\varepsilon}) - \beta_{j,\varepsilon})$$

converges to the entire solution $(U_{1,j}, U_{2,j})$ of (1.13) with $m_1 = m_2 = 0$. Furthermore, the flux $(M_{1,j}, M_{2,j}) = (\int_{\mathbb{R}^2} e^{U_{2,j}}, \int_{\mathbb{R}^2} e^{U_{1,j}})$ satisfies

$$\frac{1}{M_{1,j}} + \frac{1}{M_{2,j}} = \frac{1}{4\pi}. \quad (1.18)$$

Thus, either $\min\{M_{1,j}, M_{2,j}\} < 8\pi$ or $M_{1,j} = M_{2,j} = 8\pi$. The analysis of bubbling solutions of these two types has different phenomena. We refer the readers to [31] for the related bubbling analysis for Liouville system.

In a joint work with Zhang[20], we obtained necessary conditions for the fully bubbling solutions of Liouville type to (1.3). Before we state the main result in [20], we introduce some notations. Let $\mathbf{q} = (q_1, \dots, q_k)$, $q_i \in \mathbb{R}^2$, $i = 1, \dots, k$,

$$G_1^*(\mathbf{q}) = \sum_{i=1}^k u_{0,1}(q_i) + 8\pi \sum_{1 \leq i < j \leq k} G(q_i, q_j) \quad \text{and} \quad G_2^*(\mathbf{q}) = \sum_{i=1}^k u_{0,2}(q_i) + 8\pi \sum_{1 \leq i < j \leq k} G(q_i, q_j).$$

Denote the function $f_{i,j,\mathbf{q}}(i = 1, 2, j = 1, \dots, k)$ as follows.

$$f_{i,j,\mathbf{q}} = 8\pi(\gamma(y, q_j) - \gamma(q_j, q_j)) + \sum_{l \neq j} (G(y, q_l) - G(q_j, q_l)) + u_{0,i}(y) - u_{0,i}(q_j), \quad (1.19)$$

where $\gamma(y, q)$ is the regular part of $G(y, p)$. So, it is clear that $\frac{\partial G_1^*(\mathbf{x})}{\partial x_{j,h}} = \frac{\partial f_{1,j,\mathbf{x}}(x)}{\partial x_h}$ and $\frac{\partial G_2^*(\mathbf{x})}{\partial x_{j,h}} = \frac{\partial f_{2,j,\mathbf{x}}(x)}{\partial x_h}$. We define the quantity $\mathcal{D}^{(2)}(\mathbf{q})$ as follows

$$\begin{aligned} \mathcal{D}^{(2)}(\mathbf{q}) = \lim_{\delta \rightarrow 0} & \left(\sum_{j=1}^k \frac{\rho_j}{\rho_1} \left(\int_{\Omega_j \setminus B_\delta} \frac{e^{f_{1,j,\mathbf{q}}} - 1}{|x - q_j|^4} - \int_{\mathbb{R}^2 \setminus \Omega_j} \frac{1}{|x - q_j|^4} \right) \right. \\ & \left. + \sum_{j=1}^k \frac{\rho_j^*}{\rho_1^*} \left(\int_{\Omega_j \setminus B_\delta} \frac{e^{f_{2,j,\mathbf{q}}} - 1}{|x - q_j|^4} - \int_{\mathbb{R}^2 \setminus \Omega_j} \frac{1}{|x - q_j|^4} \right) \right) \end{aligned} \quad (1.20)$$

where $\{\Omega_j\}_{j=1, \dots, k}$ is any open set with

$$(1) \quad \Omega_i \cap \Omega_j = \emptyset, \quad i \neq j, \quad i, j \in \{1, \dots, k\},$$

$$(2) \cup_{j=1}^k \overline{\Omega_j} = \overline{\Omega},$$

$$(3) B_{d_j}(q_j) \subset \subset \Omega_j, \quad j = 1, \dots, k,$$

$$\rho_j = e^{8\pi(\gamma(q_j, q_j) + \sum_{l \neq j} G(q_j, q_l)) + u_{0,1}(q_j)}, \text{ and } \rho_j^* = e^{8\pi(\gamma(q_j, q_j) + \sum_{l \neq j} G(q_j, q_l)) + u_{0,2}(q_j)}.$$

In particular, when $k = 1$,

$$\mathcal{D}^{(2)}(q) = \lim_{\delta \rightarrow 0} \left(\int_{\Omega \setminus B_\delta} \frac{e^{u_{0,1}(x) - u_{0,1}(q)} - 1}{|x - q|^4} + \frac{e^{u_{0,1}(x) - u_{0,1}(q)} - 1}{|x - q|^4} - \int_{\mathbb{R}^2 \setminus \Omega} \frac{2}{|x - q|^4} \right). \quad (1.21)$$

When Ω is a rectangle, $k = 1$ and $p_{i,j} = p$ ($i, j = 1, 2$), it was shown [26] that the Green function $G(x, p)$ has three critical points: two of them are saddle points whose corresponding $\mathcal{D}^{(2)}(q) > 0$, and the other is a maximum points whose $\mathcal{D}^{(2)}(q) < 0$. We refer to [26, 27] for the related discussion on this kind quantity.

The necessary conditions for the existence of fully bubbling solutions of Liouville type are given as follows.

Theorem C [20] Suppose $(u_{1,\varepsilon}, u_{2,\varepsilon})$ is a sequence of fully bubbling solution of Liouville type to (1.3) with $N_1 = N_2 = 2k$ and blow up set is $\{q_1, \dots, q_k\} \notin \{p_{1,i}, p_{2,i}\}_{i=1, \dots, 2k}$. Then

- (1) $(u_{0,1} - u_{0,2})(q_i) = (u_{0,1} - u_{0,2})(q_j)$, $1 \leq i, j \leq k$.
- (2) $\mathbf{q} = (q_1, \dots, q_k)$ is a critical point of G_1^* and G_2^* .
- (3) $\mathcal{D}^{(2)}(\mathbf{q}) \leq 0$.

Naturally, we are led to the question whether the conditions obtained in Theorem C are sufficient for the existence of such solutions. In this paper, we will construct a sequence of bubbling solutions whose limiting local masses are $(8\pi, 8\pi)$ and the sufficient condition are nearly necessary. Since we assume that $N_1 = N_2 = 2k$, it is natural to use the entire solution (U_1, U_2) of (1.13) with flux $(M_1, M_2) \approx (8\pi, 8\pi)$ to construct the approximation solutions for (1.3). But (M_1, M_2) must satisfy (1.18), it adds a difficulty in construction of good approximation solutions for multi-bubble case. With the assumption (A.1)(see below), we can choose (U_1, U_2) with $(\int_{\mathbb{R}^2} e^{U_2} dx, \int_{\mathbb{R}^2} e^{U_1} dx) = (8\pi, 8\pi)$ to construct the approximation solutions around each blow-up point and glue them together. It is a crucial step when we apply the invertibility of linear operator in the contraction mapping argument.

Theorem 1.1. *Assume that*

- (A.1) $(u_{0,1} - u_{0,2})(q_i) = (u_{0,1} - u_{0,2})(q_j)$, $1 \leq i, j \leq k$.
- (A.2) \mathbf{q} be a critical point of G_1^* and G_2^*
- (A.3) \mathbf{q} be non-degenerate critical point of $G_1^* + G_2^*$
- (A.4) $\mathcal{D}^{(2)}(\mathbf{q}) < 0$

Then for $\varepsilon > 0$ small, (1.3) has a solution $(u_{1,\varepsilon}, u_{2,\varepsilon})$ satisfies

$$\frac{1}{\varepsilon^2} e^{u_i} (1 - e^{u_j}) \rightarrow 8\pi \sum_{k=1}^N \delta_{q_k}, \quad i \neq j, i, j = 1, 2, l = 1, \dots, k, \quad \text{as } \varepsilon \rightarrow 0. \quad (1.22)$$

When $k = 1$, Theorem 1.1 is reduced to the following theorem.

Theorem 1.2. *Assume q is a critical point of $u_{0,1}$ and $u_{0,2}$ and a non-degenerate critical point of $u_{0,1} + u_{0,2}$, and $\mathcal{D}^{(2)}(q) < 0$, then there exists a sequence of solution $(u_{1,\varepsilon}, u_{2,\varepsilon})$ to (1.3) blowing up at q .*

It will be interesting to consider the Liouville type bubbling solutions with $N_1 \neq N_2$. In this case, $\frac{k}{N_1} + \frac{k}{N_2} = 1$ and $k \geq 2$. For instance, $k = 2$ and $(N_1, N_2) = (3, 6)$. As suggested in the study of Liouville system [31], the quantity of $\mathcal{D}^{(2)}(\mathbf{q})$ should be a slight different form and only depend on $\min\{\frac{4\pi N_1}{k}, \frac{4\pi N_2}{k}\}$. In [31], they only consider the simple bubble case, but in our case, it must be multi-bubble. Furthermore, the invertibility of the corresponding linear operator could be another difficulty. Since the height of each bubble is compatible, the dimension of the kernel space to the corresponding linear operator is $2k + 1$. The kernel with respect to height is not a local condition which makes the analysis of linear operator difficult. We will come back to this issue in the future.

The rest of our paper is organized as follows. In Sec. 2, we construct approximate solutions of (1.3). In Sec. 3, we use the contraction mapping theorem to prove the existence of multi-bubble solutions. Appendix is devoted to the invertibility of the linear operator L_μ .

2 The Approximation Solution

In this section, we will construct an approximation solution $(U_{1,\mu}, U_{2,\mu})$ (see (2.5) below) for (1.3) with the assumptions in Theorem 1.1 and give the estimates on the approximation solution. This construction follows the method in [29]. Without loss of generality, we assume that $|\Omega| = 1$.

We consider the solution of mean field equation

$$V_{x_i, \mu_i}(y) = \ln \frac{8\mu_i^2}{(1 + \mu_i^2 |y - x_i|^2)^2}, \quad x_i \in \mathbb{R}^2, \quad \mu_i > 0, \quad (2.1)$$

which satisfies

$$\begin{cases} \Delta V_{x_i, \mu_i}(y) + e^{V_{x_i, \mu_i}(y)} = 0 \text{ in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^{V_{x_i, \mu_i}(y)} dy = 8\pi. \end{cases} \quad (2.2)$$

In the construction of approximation solution, we always assume that $|x_i - q_i| < \frac{C_1}{\mu_i}$ for some large C_1 and

$$\mu_i \in \left[\frac{\beta_1}{\sqrt{\varepsilon}}, \frac{\beta_2}{\sqrt{\varepsilon}} \right] \text{ for some constants } \beta_1, \beta_2 > 0.$$

Define ω_{x_i, μ_i}^* and $\omega_{x, \mu}^*$ as follows.

$$\omega_{x_i, \mu_i}^*(y) = \begin{cases} V_{x_i, \mu_i}(y) + 8\pi\gamma(y, x_i)(1 - \frac{1}{\theta_i}), & y \in B_{d_i}(x_i), \\ V_{0, \mu_i}(d_i) + 8\pi(G(y, x_i) - \frac{1}{2\pi} \ln \frac{1}{d_i})(1 - \frac{1}{\theta_i}), & y \in \Omega \setminus B_d(x_i), \end{cases} \quad (2.3)$$

where $\frac{1}{\theta_i} = \frac{1}{1 + (\mu_i d_i)^2}$ which makes $\omega_{x_i, \mu_i}^* \in C^1(\Omega)$, and

$$\omega_{x, \mu}^* = \sum_{i=1}^k \omega_{x_i, \mu_i}^*.$$

Set ω_μ

$$\omega_\mu(y) = \omega_{x, \mu}^*(y) - \int_{\Omega} \omega_{x, \mu}^*. \quad (2.4)$$

For $k \geq 2$, we construct the approximation solution $(U_{1, \mu}, U_{2, \mu})$ as follows:

$$(U_{1, \mu}, U_{2, \mu}) = (\omega_\mu + c_{1, \mu}, \omega_\mu + c_{2, \mu}), \quad (2.5)$$

where

$$c_{1, \mu} = \ln \frac{16k\pi\varepsilon^2}{\int_{\Omega} e^{u_{0,1} + \omega_\mu} \left(1 + \sqrt{1 - 32k\pi\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (u_{0,i} + \omega_\mu)}}{\int_{\Omega} e^{u_{0,1} + \omega_\mu} \int_{\Omega} e^{u_{0,2} + \omega_\mu}}} \right)}$$

and

$$c_{2, \mu} = \ln \frac{16k\pi\varepsilon^2}{\int_{\Omega} e^{u_{0,2} + \omega_\mu} \left(1 + \sqrt{1 - 32k\pi\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (u_{0,i} + \omega_\mu)}}{\int_{\Omega} e^{u_{0,1} + \omega_\mu} \int_{\Omega} e^{u_{0,2} + \omega_\mu}}} \right)}.$$

Remark 2.1. When $k = 1$, the approximation solution for (1.3) has a simpler form:

$$\begin{pmatrix} U_{1, \mu} \\ U_{2, \mu} \end{pmatrix} = \begin{pmatrix} \omega_{x_1, \mu_1}^* - u_{0,1}(x_1) \\ \omega_{x_1, \mu_1}^* - u_{0,2}(x_1) \end{pmatrix}. \quad (2.6)$$

In the construction of approximation, the height μ_i of the bubble ω_{x_i, μ_i} cannot be an independent variables. So, we set

$$\mu_1 = \mu, \quad \mu_i = \sqrt{\frac{\rho_1}{\rho_i}} \mu, \quad i = 2, \dots, k, \quad \text{for some } \mu > 0,$$

where

$$\rho_i = e^{8\pi\gamma(x_i, x_i) + 8\pi \sum_{j \neq i} G(x_i, x_j) + u_{0,1}(x_i)}, \quad i = 1, \dots, k.$$

It is clear that

$$\rho_1 \mu_1^2 = \rho_i \mu_i^2, \quad i = 2, \dots, k. \quad (2.7)$$

Denote $\rho_i^* = e^{8\pi\gamma(x_i, x_i) + 8\pi \sum_{j \neq i} G(x_i, x_j) + u_{0,2}(x_i)}$, $i = 1, \dots, k$. By using $(u_{0,1} - u_{0,2}(x_i)) - (u_{0,1} - u_{0,2}(q_i)) = O(\frac{1}{\mu})$, we obtain

$$\rho_1^* \mu_1^2 = \rho_i^* \mu_i^2 \left(1 + O\left(\frac{1}{\mu}\right)\right), \quad i = 1, \dots, k. \quad (2.8)$$

In the rest of this section, we will some properties of the approximation solution $(U_{1,\mu}, U_{2,\mu})$.

For $l = 1, 2$, $j = 1, \dots, k$ and $y \in B_{d_j}(x_j)$, we have

$$\begin{aligned} & u_{0,l}(y) - u_{0,l}(x_i) + 8\pi \left((\gamma(y, x_i) - \gamma(x_i, x_i)) \left(1 - \frac{1}{\theta_i}\right) + \sum_{m \neq j} (G(y, x_m) - G(x_j, x_m)) \left(1 - \frac{1}{\theta_m}\right) \right) \\ &= \left\langle Du_{0,l}(x_i) + 8\pi \sum_{m \neq j} DG(x_i, x_m), y - x_i \right\rangle + O(|y - x_i|^2 + \frac{1}{\mu^2}). \end{aligned} \quad (2.9)$$

By this and (2.1), we find that

$$\begin{aligned} & \int_{B_{d_i}(x_i)} e^{\omega_\mu^* + u_{0,1}} \\ &= \frac{8^{k-1} \rho_i \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(1 + O\left(\frac{1}{\mu^2}\right)\right) \\ & \quad \times \int_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i} + u_{0,i}(y) - u_{0,i}(x_j) + 8\pi((\gamma(y, x_j) - \gamma(x_j, x_j)) + \sum_{m \neq j} (G(y, x_m) - G(x_j, x_m)))} \\ &= \frac{8^{k-1} \rho_i \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8\pi + \int_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} |y - x_i|^2 + O\left(\frac{1}{\mu^2}\right)\right) = \frac{8^{k-1} \rho_i \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8\pi + O\left(\frac{\ln \mu}{\mu^2}\right)\right). \end{aligned} \quad (2.10)$$

Similarly,

$$\int_{B_{d_i}(x_i)} e^{\omega_\mu^* + u_{0,2}} = \frac{8^{k-1} \rho_i^* \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8\pi + O\left(\frac{\ln \mu}{\mu^2}\right)\right). \quad (2.11)$$

By (2.7), (2.8), (2.10) and (2.11), we obtain

$$\int_{\Omega} e^{\omega_\mu^* + u_{0,1}} = \sum_{i=1}^k \frac{8^{k-1} \rho_i \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8\pi + O\left(\frac{\ln \mu}{\mu^2}\right)\right) = \frac{8^{k-1} \rho_i \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8k\pi + O\left(\frac{\ln \mu}{\mu^2}\right)\right). \quad (2.12)$$

Similarly,

$$\int_{\Omega} e^{\omega_\mu^* + u_{0,2}} = \frac{8^{k-1} \rho_i^* \mu_i^2}{\mu_1^2 \cdots \mu_k^2} \left(8k\pi + O\left(\frac{1}{\mu}\right)\right), \quad (2.13)$$

where (2.8) is used. The asymptotic behaviours of ω_μ and $c_{i,\mu}$ ($i=1,2$) will be discussed in the following proposition.

Proposition 2.2. (1) The function ω_μ defined in (2.4) satisfies that, for $\delta > 0$,

$$\begin{cases} \max_{y \in B_\delta(x_i)} \omega_\mu(y) = 2 \ln \frac{1}{\varepsilon} + O(1), \\ \omega_\mu(y) = O(1), y \in \Omega \setminus \cup_{i=1}^k B_\delta(x_i). \end{cases} \quad (2.14)$$

(2) The constants $c_{1,\mu}$ and $c_{2,\mu}$ satisfy

$$c_{i,\mu} = -3 \ln \frac{1}{\varepsilon} + O(1), \quad i = 1, 2. \quad (2.15)$$

Proof. The estimate of ω_μ is exactly the same as Proposition 2.1 in [29]. So, we omit the proof.

By definition of $c_{1,\mu}$, we deduce that, for $\mu > 0$ large,

$$e^{c_{1,\mu}} = \frac{16k\pi\varepsilon^2}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \left(2 - 16k\pi\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \int_{\Omega} e^{\omega_\mu+u_{0,2}}} + O\left(\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \int_{\Omega} e^{\omega_\mu+u_{0,2}}}\right)^2 \right)} \quad (2.16)$$

On the other hand,

$$\frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \int_{\Omega} e^{\omega_\mu+u_{0,2}}} = \frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu^*+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu^*+u_{0,1}} \int_{\Omega} e^{\omega_\mu^*+u_{0,2}}} = O(\mu^2). \quad (2.17)$$

So, by (2.16) and (2.17),

$$e^{c_{1,\mu}} = \frac{8k\pi\varepsilon^2}{\int_{\Omega} e^{\omega_\mu+u_{0,1}}} \left(1 + 8k\pi\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \int_{\Omega} e^{\omega_\mu+u_{0,2}}} + O(\varepsilon^4 \mu^4) \right), \quad (2.18)$$

and thus

$$\begin{aligned} c_{1,\mu} &= \ln(8k\pi) + 2 \ln \varepsilon - \ln \int_{\Omega} e^{\omega_\mu+u_{0,1}} + 8k\pi\varepsilon^2 \frac{\int_{\Omega} e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_{\Omega} e^{\omega_\mu+u_{0,1}} \int_{\Omega} e^{\omega_\mu+u_{0,2}}} + O(\varepsilon^4 \mu^4) \\ &= 2 \ln \varepsilon - 2 \ln \mu + O(1) \\ &= -3 \ln \frac{1}{\varepsilon} + O(1). \end{aligned} \quad (2.19)$$

Similarly, $c_{2,\mu} = -3 \ln \frac{1}{\varepsilon} + O(1)$ □

In the end of this section we estimate

$$-\Delta U_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}} (e^{U_{1,\mu}+u_{0,1}} - 1) + 8k\pi$$

and

$$-\Delta U_{2,\mu} + \frac{1}{\varepsilon^2} e^{U_{1,\mu}+u_{0,1}} (e^{U_{2,\mu}+u_{0,2}} - 1) + 8k\pi$$

which will be useful in the next section.

Proposition 2.3. *Let $(U_{1,\mu}, U_{2,\mu})$ be defined in (2.5). Then*

$$\begin{aligned}
& -\Delta U_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}} (e^{U_{1,\mu}+u_{0,1}} - 1) + 8k\pi \\
&= \sum_{i=1}^k \left(1_{B_{d_i}(x_i)} e^{V_{x_i,\mu_i}} (1 - e^{f_{2,i,\mathbf{x},\mu}}) + \frac{8\pi}{\theta_i} \right) \\
&+ O\left(\frac{\ln \mu}{\mu^2} + \sum_{i=1}^k \left(\frac{1}{\mu^2} e^{V_{x_i,\mu_i}+u_{0,2}} + \frac{1}{\mu^4} e^{2V_{x_i,\mu_i}+u_{0,1}+u_{0,2}} \right) \right)
\end{aligned} \tag{2.20}$$

and

$$\begin{aligned}
& -\Delta U_{2,\mu} + \frac{1}{\varepsilon^2} e^{U_{1,\mu}+u_{0,1}} (e^{U_{2,\mu}+u_{0,2}} - 1) + 8k\pi \\
&= \sum_{i=1}^k \left(1_{B_{d_i}(x_i)} e^{V_{x_i,\mu_i}} (1 - e^{f_{1,i,\mathbf{x},\mu}}) + \frac{8\pi}{\theta_i} \right) \\
&+ O\left(\frac{\ln \mu}{\mu^2} + \sum_{i=1}^k \left(\frac{1}{\mu^2} e^{V_{x_i,\mu_i}+u_{0,1}} + \frac{1}{\mu^4} e^{2V_{x_i,\mu_i}+u_{0,1}+u_{0,2}} \right) \right).
\end{aligned} \tag{2.21}$$

where

$$\begin{aligned}
f_{1,i,\mathbf{x},\mu}(y) &= 8\pi \left((\gamma(y, x_i) - \gamma(x_i, x_i)) \left(1 - \frac{1}{\theta_i}\right) + \sum_{j \neq i} (G(y, x_i) - G(x_i, x_j)) \left(1 - \frac{1}{\theta_j}\right) \right) \\
&+ u_{0,1}(y) - u_{0,1}(x_i)
\end{aligned} \tag{2.22}$$

$$\begin{aligned}
f_{2,i,\mathbf{x},\mu}(y) &= 8\pi \left((\gamma(y, x_i) - \gamma(x_i, x_i)) \left(1 - \frac{1}{\theta_i}\right) + \sum_{j \neq i} (G(y, x_i) - G(x_i, x_j)) \left(1 - \frac{1}{\theta_j}\right) \right) \\
&+ u_{0,2}(y) - u_{0,2}(x_i)
\end{aligned} \tag{2.23}$$

Proof. We only prove (2.20) since the proof of (2.21) is similar.

By the definition of $U_{1,\mu}$, we find that

$$\begin{aligned}
-\Delta U_{1,\mu} &= -\Delta \omega_\mu \\
&= \sum_{i=1}^k \left(1_{B_{d_i}(x_i)} e^{V_{x_i,\mu_i}} - 8\pi \left(1 - \frac{1}{\theta_i}\right) \right)
\end{aligned} \tag{2.24}$$

where $1_A = 1$ in A and $1_A = 0$ otherwise.

Using (2.12) and (2.13), we have

$$\begin{aligned}
& \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}} \\
&= \frac{8k\pi e^{\omega_\mu+u_{0,2}}}{\int_\Omega e^{\omega_\mu+u_{0,2}}} \left(1 + 8k\pi \varepsilon^2 \frac{\int_\Omega e^{\sum_{i=1}^2 (\omega_\mu+u_{0,i})}}{\int_\Omega e^{\omega_\mu+u_{0,1}} \int_\Omega e^{\omega_\mu+u_{0,2}}} + O(\varepsilon^4 \mu^4) \right)
\end{aligned} \tag{2.25}$$

and

$$\begin{aligned} & \frac{1}{\varepsilon^2} e^{\sum_{i=1}^2 (U_{i,\mu} + u_{0,i})} \\ &= 64k^2 \pi^2 \varepsilon^2 \left(\frac{e^{\omega_\mu + u_{0,1}} e^{\omega_\mu + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,1}} \int_\Omega e^{\omega_\mu + u_{0,2}}} \right) \left(1 + 16k\pi \varepsilon^2 \frac{\int_\Omega e^{\sum_{i=1}^2 (\omega_\mu + u_{0,i})}}{\int_\Omega e^{\omega_\mu + u_{0,1}} \int_\Omega e^{\omega_\mu + u_{0,2}}} + O(\varepsilon^4 \mu^4) \right). \end{aligned} \quad (2.26)$$

Combining (2.24), (2.25) and (2.26), we obtain

$$\begin{aligned} & -\Delta U_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + u_{0,2}} (e^{U_{1,\mu} + u_{0,1}} - 1) + 8k\pi \\ &= \sum_{i=1}^k \left(1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} + \frac{8\pi}{\theta_i} \right) - \frac{8k\pi e^{\omega_\mu + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,2}}} + K_{1,u} + R_{1,\mu} \end{aligned} \quad (2.27)$$

where

$$K_{1,\mu} = \frac{64k^2 \pi^2 \varepsilon^2 \int_\Omega e^{2\omega_\mu + u_{0,1} + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,1}} \int_\Omega e^{\omega_\mu + u_{0,2}}} \left[\frac{e^{2\omega_\mu + u_{0,1} + u_{0,2}}}{\int_\Omega e^{2\omega_\mu + u_{0,1} + u_{0,2}}} - \frac{e^{\omega_\mu + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,2}}} \right]$$

and

$$R_{1,\mu} = O \left(\frac{\mu^4 \varepsilon^4 e^{\omega_\mu + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,2}}} + \frac{\varepsilon^4 e^{2\omega_\mu + u_{0,1} + u_{0,2}} \int_\Omega e^{2\omega_\mu + u_{0,1} + u_{0,2}}}{(\int_\Omega e^{\omega_\mu + u_{0,1}})^2 (\int_\Omega e^{\omega_\mu + u_{0,2}})^2} + \frac{\varepsilon^6 \mu^4 e^{2\omega_\mu + u_{0,1} + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,1}} \int_\Omega e^{\omega_\mu + u_{0,2}}} \right).$$

Note that

$$\int_\Omega e^{\sum_{i=1}^2 (\omega_\mu^* + u_{0,i})} = O \left(\frac{\mu^2}{\mu^{4(k-1)}} \right)$$

By this, and (2.12) and (2.13), we obtain

$$K_{1,\mu} = O \left(\frac{1}{\mu^2} e^{V_{x_i, \mu_i} + u_{0,2}} + \frac{1}{\mu^4} e^{2V_{x_i, \mu_i} + u_{0,1} + u_{0,2}} \right)$$

and

$$R_{1,\mu} = O \left(\frac{1}{\mu^4} e^{V_{x_i, \mu_i} + u_{0,2}} + \frac{1}{\mu^6} e^{2V_{x_i, \mu_i} + u_{0,1} + u_{0,2}} \right)$$

Next, by (2.13)

$$\frac{8k\pi e^{\omega_\mu + u_{0,2}}}{\int_\Omega e^{\omega_\mu + u_{0,2}}} = \frac{8k\pi e^{\omega_\mu^* + u_{0,2}}}{\int_\Omega e^{\omega_\mu^* + u_{0,2}}} = \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i} + f_{1,i,\mu}} + O\left(\frac{1}{\mu^2}\right) \quad (2.28)$$

By this, we are led to (2.20). \square

3 The Reduction and The Existence

In this section, we will use the contraction mapping theorem to show that there exist $\mathbf{x} = (x_1, \dots, x_k)$ and $\mu = (\mu_1, \dots, \mu_k)$ such that (1.3) has a solution of this form $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} U_{1,\mu} \\ U_{2,\mu} \end{pmatrix} +$

$\begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix}$ where $\begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix}$ is a perturbation term. For this purpose, we will use the linear operator L_μ (see (3.1) below). However, the linear operator L_μ has non-trivial kernel, hence, we only can use the contraction mapping theorem to solve (1.3) up to its kernel. Then we use the non-degeneracy of $G_1^* + G_2^*$ at the blow-up point \mathbf{q} and $\mathcal{D}^{(2)}(\mathbf{q}) < \mathbf{0}$ to find suitable \mathbf{x} and μ such that (1.3) has a real solution whose blow-up set is $\{q_1, \dots, q_k\}$. Here, we only present the multi-bubble case ($k \geq 2$). For the case $k = 1$, it can be easily deduced from the multi-bubble case by using the approximation solution (2.6).

In view of $U_{1,\mu}, U_{2,\mu}$, we consider the following simplified operator

$$L_\mu \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 & \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} \\ \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \quad (3.1)$$

Then, $(\omega_{1,\mu}, \omega_{2,\mu})$ satisfies

$$L_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} = \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix}, \quad (3.2)$$

where

$$g_{1,\mu}(x, t_1, t_2) = \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} t_2 - \frac{1}{\varepsilon} e^{U_{2,\mu} + u_{0,2} + t_2} (1 - e^{U_{1,\mu} + u_{0,1} + t_1}) - \Delta U_{1,\mu} + 8k\pi,$$

and

$$g_{2,\mu}(x, t_1, t_2) = \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} t_1 - \frac{1}{\varepsilon} e^{U_{1,\mu} + u_{0,1} + t_1} (1 - e^{U_{2,\mu} + u_{0,2} + t_2}) - \Delta U_{2,\mu} + 8k\pi.$$

To apply the contraction argument, we first introduce two function spaces:

$$X_{\alpha,\mu,2} \quad \text{and} \quad Y_{\alpha,\mu,2}.$$

Fix a small fixed constant $\alpha > 0$, we define

$$\rho(x) = (1 + |x|)^{1 + \frac{\alpha}{2}}, \quad \hat{\rho}(x) = \frac{1}{(1 + |x|)(\log(2 + |x|))^{1 + \frac{\alpha}{2}}}.$$

Denote $\Omega' = \cup_{j=1}^k B_{d_j}(x_j)$. We say $\begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix}$ is in $X_{\alpha,\mu,2}$ if

$$\left\| \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \right\|_{X_{\alpha,\mu,2}}^2 = \sum_{j=1}^k \sum_{i=1}^2 \left(\|\Delta \tilde{\xi}_{i,j} \rho\|_{L^2(B_{2d_j \mu_j})}^2 + \|\tilde{\xi}_{i,j} \hat{\rho}\|_{L^2(B_{2d_j \mu_j})}^2 \right) \quad (3.3)$$

$$+ \sum_{i=1}^2 \left(\|\Delta \xi_i\|_{L^2(\Omega')}^2 + \|\xi_i\|_{L^2(\Omega')}^2 \right) < +\infty, \quad (3.4)$$

where $\tilde{\xi}_{i,j}(y) = \xi(q_j + \frac{1}{\mu_j}y)$ and $B_d = B_d(0)$; $\begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix}$ is in $Y_{\alpha,\mu,2}$ if

$$\left\| \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \right\|_{Y_{\alpha,\mu}} = \sum_{j=1}^N \sum_{i=1}^2 \left(\frac{1}{\mu_j^4} \|\tilde{\xi}_{i,j}\rho\|_{L^2(B_{2d_j\mu_j})}^2 \right) + \sum_{i=1}^2 \left(\|\xi_i\|_{L^2(\Omega \setminus \Omega')}^2 \right) < +\infty. \quad (3.5)$$

We also denote $\|\xi\|_{X_{\alpha,\mu}} = \left\| \begin{pmatrix} \xi \\ 0 \end{pmatrix} \right\|_{X_{\alpha,\mu,2}}$ and $\|\xi\|_{Y_{\alpha,\mu}} = \left\| \begin{pmatrix} \xi \\ 0 \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}}$. So, we say $\xi \in X_{\alpha,\mu}$ if $\|\xi\|_{X_{\alpha,\mu}} < +\infty$ and $\xi \in Y_{\alpha,\mu}$ if $\|\xi\|_{Y_{\alpha,\mu}} < +\infty$.

Consider the cut-off function $\chi_j \in C^\infty(\mathbb{R}^2)$ satisfying

$$\chi_j(x) = \begin{cases} 1 & \text{for } |x| \leq d_j \\ 0 & \text{for } |x| \geq 2d_j \end{cases}$$

and $0 \leq \chi_j \leq 1$. Next, we define the approximated kernel for L_μ as follows:

$$\begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\mu_1} + \sum_{i=1}^k \sqrt{\frac{\rho_1}{\rho_i}} \frac{2\chi_i(y-x_i)}{\mu_i(1+\mu_i|y-x_i|^2)} \\ -\frac{1}{\mu_1} + \sum_{i=1}^k \sqrt{\frac{\rho_1}{\rho_i}} \frac{2\chi_i(y-x_i)}{\mu_i(1+\mu_i|y-x_i|^2)} \end{pmatrix}, \quad (3.6)$$

$$\begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix} = \chi_j(|x - p_{j,\varepsilon}|) \begin{pmatrix} \frac{\mu_j^2(y_i - x_{j,i})}{1+\mu_j|y-x_i|^2} \\ \frac{\mu_j^2(y_i - x_{j,i})}{1+\mu_j|y-x_i|^2} \end{pmatrix}, \quad i = 1, 2, \quad j = 1, \dots, k. \quad (3.7)$$

After calculation, it is not difficult to see that

$$L_\mu \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} = O\left(\frac{1}{\mu^3}\right), \quad L_\mu \begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix} = O(1), \quad i = 1, 2, \quad j = 1, \dots, k.$$

Let

$$\begin{pmatrix} Z_{1,0} \\ Z_{2,0} \end{pmatrix} = \begin{pmatrix} \Delta Y_{1,0} \\ \Delta Y_{2,0} \end{pmatrix} \quad (3.8)$$

and

$$\begin{pmatrix} Z_{1,i,j} \\ Z_{2,i,j} \end{pmatrix} = \begin{pmatrix} \Delta Y_{1,i,j} \\ \Delta Y_{2,i,j} \end{pmatrix}, \quad i = 1, 2, \quad j = 1, \dots, k. \quad (3.9)$$

Note that $\begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix}$, $\begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix}$, $\begin{pmatrix} Z_{1,0} \\ Z_{2,0} \end{pmatrix}$ and $\begin{pmatrix} Z_{1,i,j} \\ Z_{2,i,j} \end{pmatrix}$ are doubly periodic. Let

$$E_{\mu,2} = \left\{ \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \in X_{\alpha,\mu,2} : \int_{\Omega} \left\langle \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, \begin{pmatrix} Z_{1,0} \\ Z_{2,0} \end{pmatrix} \right\rangle = \int_{\Omega} \left\langle \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, \begin{pmatrix} Z_{1,i,j} \\ Z_{2,i,j} \end{pmatrix} \right\rangle = 0, \quad i = 1, 2, \quad j = 1, \dots, k. \right\}$$

and

$$F_{\mu,2} = \left\{ \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \in Y_{\alpha,\mu,2} : \int_{\Omega} \left\langle \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle = \int_{\Omega} \left\langle \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, \begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix} \right\rangle = 0, \quad i = 1, 2, \quad j = 1, \dots, k. \right\}.$$

We define the project operator $Q_\mu : Y_{\alpha,\mu,2} \rightarrow F_{\mu,2}$ as follows:

$$Q_\mu \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} - c_0 \begin{pmatrix} Z_{1,0} \\ Z_{2,0} \end{pmatrix} - \sum_{i=1}^2 \sum_{j=1}^k c_{i,j} \begin{pmatrix} Z_{1,i,j} \\ Z_{2,i,j} \end{pmatrix} \quad (3.10)$$

where $c_0, c_{i,j}$ are chosen so that

$$Q_\mu \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \in F_{\mu,2}. \quad (3.11)$$

It is standard to prove the following result(or see Lemma 3.1 in [29]).

Lemma 3.1. *There is a constant $C > 0$, independent of \mathbf{x} and μ , such that*

$$\left\| Q_\mu \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}} \leq C \left\| \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}}. \quad (3.12)$$

By using the contraction mapping argument, we obtain the following theorem.

Theorem 3.2. *Assume (A.1)-(A.4) in Theorem 1.1 hold. There exists $\varepsilon_0 > 0$ such that for $\varepsilon \in (0, \varepsilon_0)$, $|\mathbf{x} - \mathbf{q}| < \frac{C_1}{\mu}$, and $\mu \in (\frac{\beta_1}{\sqrt{\varepsilon}}, \frac{\beta_2}{\sqrt{\varepsilon}})$ for some constants $C_1, \beta_1, \beta_2 > 0$, then there exists $\begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \in E_{\mu,2}$ satisfying*

$$Q_\mu \left(L_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} - \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (3.13)$$

Furthermore, $\begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix}$ is a C^1 map of $(\mathbf{x}, \mu_1, \dots, \mu_k)$ to $X_{\alpha,\mu,2}$ and

$$\left\| Q_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| Q_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \leq \frac{C \ln \mu}{\mu^{2-\frac{\alpha}{2}}}, \quad (3.14)$$

where the constant α is the same constant as in $X_{\alpha,\mu,2}$.

Proof. By the Theorem 4.1, the equation (3.13) can be rewritten as

$$\begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} = B_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} := (Q_\mu L_\mu)^{-1} Q_\mu \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix}. \quad (3.15)$$

Define

$$S_\mu = \left\{ \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} : \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \in E_\mu, \left\| \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \leq \frac{1}{\mu} \right\}.$$

Firstly, we show that B_μ maps S_μ to S_μ . By Lemma 3.1 and (4.2), we have

$$\left\| B_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \leq C \ln \mu \left\| Q_\mu \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}} \leq C \ln \mu \left\| \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}}. \quad (3.16)$$

Thus we need to estimate $\left\| \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}}$.

Note that

$$\begin{aligned} & e^{U_{2,\mu}+u_{0,2}+t_2} (1 - e^{U_{1,\mu}+u_{0,1}+t_1}) \\ &= e^{U_{2,\mu}+u_{0,2}} (1 - e^{U_{1,\mu}+u_{0,1}}) + e^{U_{2,\mu}+u_{0,2}} (1 - e^{U_{1,\mu}+u_{0,1}}) t_2 - e^{\sum_{i=1}^2 U_{i,\mu}+u_{0,i}} t_1 \\ & \quad + O\left((e^{U_{1,\mu}+u_{0,1}} + e^{U_{2,\mu}+u_{0,2}} + e^{\sum_{i=1}^2 U_{i,\mu}+u_{0,i}}) (t_1^2 + t_2^2) \right) \end{aligned} \quad (3.17)$$

Thus, by (3.17), we obtain

$$\begin{aligned} g_{1,\mu} &= h_\mu \omega_{2,\mu} - \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}+\omega_{2,\mu}} (1 - e^{U_{1,\mu}+u_{0,1}+\omega_{1,\mu}}) + 8k\pi - \Delta U_{1,\mu} \\ &= (h_\mu - \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}}) \omega_{2,\mu} + O\left(\frac{1}{\varepsilon^2} e^{\sum_{i=1}^2 U_{i,\mu}+u_{0,i}} (|\omega_{1,\mu}| + |\omega_{2,\mu}|) \right) \\ & \quad + O\left(\left(\frac{1}{\varepsilon^2} e^{U_{1,\mu}+u_{0,1}} + \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}} \right) (|\omega_{1,\mu}|^2 + |\omega_{2,\mu}|^2) \right) \\ & \quad + O\left(\sum_{i=1}^k e^{V_{x_i,\mu_i}} (D(u_{0,2}(x_i) + 8\pi \sum_{j \neq i} G(x_j, x_i)) |y - x_i| + |y - x_i|^2) \right) \\ & \quad + O\left(\frac{\ln \mu}{\mu^2} + \sum_{i=1}^k \left(\frac{1}{\mu^2} e^{V_{x_i,\mu_i}+u_{0,2}} + \frac{1}{\mu^4} e^{2V_{x_i,\mu_i}+u_{0,1}+u_{0,2}} \right) \right). \end{aligned} \quad (3.18)$$

By this, we have

$$\begin{aligned} & \frac{1}{\mu^4} \|g_{1,\mu}(\mu^{-1}y + x_i, \omega_{1,\mu}(\mu^{-1}y + x_i), \omega_{2,\mu}(\mu^{-1}y + x_i)) \rho(y)\|_{L^2(B_{2d_i})}^2 \\ & \leq C \left(\frac{\ln^2 \mu}{\mu^4} + \varepsilon^4 \mu^4 \right) \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)} + \frac{C |DG_2^*(\mathbf{x})|^2}{\mu^2} \\ & \leq \frac{C}{\mu^{4-\alpha}}. \end{aligned} \quad (3.19)$$

On the other hand, by the definition of $(U_{1,\mu}, U_{2,\mu})$

$$\|g_{1,\mu}\|_{L^2(\Omega \setminus \Omega')} \leq \frac{C}{\mu^2}. \quad (3.20)$$

Similarly, we have

$$\frac{1}{\mu^4} \|g_{2,\mu}(\mu^{-1}y + x_i, \omega_{1,\mu}(\mu^{-1}y + x_i), \omega_{2,\mu}(\mu^{-1}y + x_i)) \rho(y)\|_{L^2(B_{2d_i})}^2 \leq \frac{C}{\mu^{4-\alpha}} \quad (3.21)$$

and

$$\|g_{2,\mu}\|_{L^2(\Omega \setminus \Omega')} \leq \frac{C}{\mu^2}. \quad (3.22)$$

By (3.19), (3.20), (3.21) and (3.22),

$$\left\| \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}} \leq \frac{C}{\mu^{2-\frac{\alpha}{2}}} \quad (3.23)$$

Next, we show that B_μ is a contraction map. For any $(\omega_{1,\mu}, \omega_{2,\mu})$ and $(\widetilde{\omega}_{1,\mu}, \widetilde{\omega}_{2,\mu}) \in S_\mu$, as the calculation above, we have

$$\begin{aligned} & \left\| B_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} - B_\mu \begin{pmatrix} \widetilde{\omega}_{1,\mu} \\ \widetilde{\omega}_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| B_\mu \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} - B_\mu \begin{pmatrix} \widetilde{\omega}_{1,\mu} \\ \widetilde{\omega}_{2,\mu} \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \\ & \leq C \ln \mu \left\| \begin{pmatrix} g_{1,\mu}(x, \omega_{1,\mu}, \omega_{2,\mu}) \\ g_{2,\mu}(x, \omega_{1,\mu}, \omega_{2,\mu}) \end{pmatrix} - \begin{pmatrix} g_{1,\mu}(x, \widetilde{\omega}_{1,\mu}, \widetilde{\omega}_{2,\mu}) \\ g_{1,\mu}(x, \widetilde{\omega}_{1,\mu}, \widetilde{\omega}_{2,\mu}) \end{pmatrix} \right\|_{X_{\alpha,\mu,2}}. \end{aligned} \quad (3.24)$$

Combing (3.23) and (3.24), we have proved that B_μ is a contraction map. Furthermore, by the contraction mapping theorem, we know that there exists $(\omega_{1,\mu}, \omega_{2,\mu}) \in S_\mu$ satisfying

$$\left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \leq \left\| \begin{pmatrix} g_{1,\mu} \\ g_{2,\mu} \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}} \leq \frac{C \ln \mu}{\mu^{2-\frac{\alpha}{2}}}. \quad (3.25)$$

□

By the above theorem, we obtain that for $|\mathbf{x} - \mathbf{q}| < \frac{1}{\mu}$ and $\mu \in [\frac{\beta_0}{\sqrt{\varepsilon}}, \frac{\beta_0}{\sqrt{\varepsilon}}]$, then there exists a doubly periodic function $(\omega_{1,\mu}, \omega_{2,\mu})$, c_0 and $c_{i,j}$, $i = 1, 2, j = 1, \dots, k$, such that

$$\begin{aligned} & \Delta \begin{pmatrix} U_{1,\mu} + \omega_{1,\mu} \\ U_{2,\mu} + \omega_{2,\mu} \end{pmatrix} + \begin{pmatrix} \frac{1}{\varepsilon^2} e^{U_{2,\mu} + u_{0,2} + \omega_{2,\mu}} (1 - e^{U_{1,\mu} + u_{0,1} + \omega_{1,\mu}}) - 8k\pi \\ \frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1} + \omega_{1,\mu}} (1 - e^{U_{2,\mu} + u_{0,2} + \omega_{2,\mu}}) - 8k\pi \end{pmatrix} \\ & = c_0 \begin{pmatrix} Z_{1,0} \\ Z_{2,0} \end{pmatrix} + \sum_{i=1}^2 \sum_{j=1}^k \begin{pmatrix} Z_{1,i,j} \\ Z_{2,i,j} \end{pmatrix}. \end{aligned} \quad (3.26)$$

To obtain a true solution for (1.3), we need to choose suitable \mathbf{x} and μ such that

$$c_0 = c_{i,j} = 0, \quad i = 1, 2, \quad j = 1, \dots, k. \quad (3.27)$$

We begin with the following simple observation.

Lemma 3.3. *If*

$$\int_{\Omega} \left\langle \begin{pmatrix} \Delta(U_{1,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}} (1 - e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}}) - 8k\pi \\ \Delta(U_{2,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}} (1 - e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}}) - 8k\pi \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle = 0 \quad (3.28)$$

and for $i = 1, 2, j = 1, \dots, k$,

$$\int_{\Omega} \left\langle \begin{pmatrix} \Delta(U_{1,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}} (1 - e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}}) - 8k\pi \\ \Delta(U_{2,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}} (1 - e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}}) - 8k\pi \end{pmatrix}, \begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix} \right\rangle = 0, \quad (3.29)$$

then $c_0 = c_{i,j} = 0, i = 1, 2, j = 1, \dots, k$.

We calculate the left hand side of (3.28) and (3.29) in the following two lemmas.

Lemma 3.4. *For $i = 1, 2$, $j = 1, \dots, k$, there exist $A_{i,j} > 0$, such that*

$$\begin{aligned} & \int_{\Omega} \left\langle \left(\begin{array}{c} \Delta(U_{1,\mu} + \omega_{1,\mu}) \\ \Delta(U_{2,\mu} + \omega_{1,\mu}) \end{array} \right) + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}} (1 - e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}}) - 8k\pi \right\rangle, \begin{pmatrix} Y_{1,i,j} \\ Y_{2,i,j} \end{pmatrix} \right\rangle \\ & = A_{i,j} \left(\frac{\partial G_1^*(\mathbf{x})}{\partial x_{i,j}} + \frac{\partial G_2^*(\mathbf{x})}{\partial x_{i,j}} \right) + O\left(\frac{\ln \mu}{\mu^{2-\frac{\alpha}{2}}}\right) \end{aligned} \quad (3.30)$$

Proof. We mainly use (2.20), (2.21) and (3.25) to obtain the estimate (3.30). We calculate the case $j = 1$ only. The others are similar.

Step 1. We first calculate this term

$$\int_{\Omega} \left\langle \left(\begin{array}{c} \Delta U_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + u_{0,2}} - 8k\pi \\ \Delta U_{2,\mu} + \frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1}} - 8k\pi \end{array} \right), \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle. \quad (3.31)$$

By the definition of $(U_{1,\mu}, U_{2,\mu})$ and symmetry, we find that

$$\begin{aligned} & \int_{\Omega} \left\langle \left(\begin{array}{c} \Delta U_{1,\mu} - 8k\pi \\ \Delta U_{2,\mu} - 8k\pi \end{array} \right) \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle \\ & = \int_{\Omega} 1_{B_{d_1}(x_1)} e^{V_{x_1, \mu_1}} Y_{1,i,1} + 1_{B_{d_1}(x_1)} e^{V_{x_1, \mu_1}} Y_{2,i,1} + \sum_{i=1}^k \frac{8\pi}{\theta_i} \int_{\Omega} Y_{1,i,1} + Y_{2,i,1} = 0 \end{aligned} \quad (3.32)$$

Next,

$$\begin{aligned} & \int_{\Omega} \left\langle \left(\begin{array}{c} \frac{1}{\varepsilon^2} e^{U_{2,\mu} + u_{0,2}} - 8k\pi \\ \frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1}} - 8k\pi \end{array} \right), \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle \\ & = \frac{8k\pi \int_{\Omega} e^{\omega_{\mu} + u_{0,1}} Y_{1,i,1}}{\int_{\Omega} e^{\omega_{\mu} + u_{0,1}}} + \frac{8k\pi \int_{\Omega} e^{\omega_{\mu} + u_{0,2}} Y_{2,i,1}}{\int_{\Omega} e^{\omega_{\mu} + u_{0,2}}} \\ & = \frac{\mu_2^2 \cdots \mu_k^2 \int_{B_{d_1}(x_1)} e^{\omega_{\mu}^* + u_{0,2}} Y_{1,i,1}}{8^{k-1}(\rho_1^* + O(\frac{1}{\mu}))} + \frac{\mu_2^2 \cdots \mu_k^2 \int_{B_{d_1}(x_1)} e^{\omega_{\mu}^* + u_{0,1}} Y_{2,i,1}}{8^{k-1}(\rho_1 + O(\frac{1}{\mu}))} + O\left(\frac{1}{\mu^2}\right) \\ & = \int_{\mathbb{R}^2} \frac{8}{(1+|y|^2)^2} \frac{|y|^2}{1+|y|^2} dy \left(\frac{\partial G_2^*(\mathbf{x})}{\partial x_{i,2}} + \frac{\partial G_1^*(\mathbf{x})}{\partial x_{i,1}} \right) + O\left(\frac{\ln \mu}{\mu^2}\right), \end{aligned} \quad (3.33)$$

where (A_2) is used. We thus denote $A_{i,j} = \int_{\mathbb{R}^2} \frac{8}{(1+|y|^2)^2} \frac{|y|^2}{1+|y|^2} dy$.

Step 2. Next, we estimate the remainder term. By mean value theorem, there exist t_i between

0 and $\omega_{i,\mu}$ ($i = 1, 2$), such that

$$\begin{aligned}
& \int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \frac{1}{\varepsilon^2}(e^{U_{2,\mu}+\omega_{2,\mu}+u_{0,2}} - e^{U_{2,\mu}+u_{0,2}}) \\ \Delta\omega_{2,\mu} + \frac{1}{\varepsilon^2}(e^{U_{1,\mu}+\omega_{1,\mu}+u_{0,1}} - e^{U_{1,\mu}+u_{0,1}}) \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle \\
&= \int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \frac{1}{\varepsilon^2}e^{U_{2,\mu}+u_{0,2}+t_2}\omega_{2,\mu} \\ \Delta\omega_{2,\mu} + \frac{1}{\varepsilon^2}e^{U_{1,\mu}+u_{0,1}+t_1}\omega_{1,\mu} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle \\
&= \int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \frac{8\pi e^{\omega_{\mu}^*+u_{0,2}}}{\int_{\Omega} e^{\omega_{\mu}^*+u_{0,2}}} \omega_{2,\mu} \\ \Delta\omega_{2,\mu} + \frac{8\pi e^{\omega_{\mu}^*+u_{0,1}}}{\int_{\Omega} e^{\omega_{\mu}^*+u_{0,1}}} \omega_{1,\mu} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle + O\left(\mu \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)}^2\right) \\
&= \int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} \omega_{2,\mu} \\ \Delta\omega_{2,\mu} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} \omega_{1,\mu} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle + O\left(\mu \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)}^2\right)
\end{aligned} \tag{3.34}$$

Since $(\omega_{1,\mu}, \omega_{2,\mu})$ and $(Y_{1,i,1}, Y_{2,i,1})$ are doubly periodic,

$$\begin{aligned}
& \int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} \omega_{2,\mu} \\ \Delta\omega_{2,\mu} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} \omega_{1,\mu} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle \\
&= \int_{\Omega} \left\langle \begin{pmatrix} \Delta Y_{1,i,1} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} Y_{2,i,1} \\ \Delta Y_{2,i,1} + \sum_{m=1}^k 1_{B_{d_m}}(x_m) e^{V_{x_m,\mu m}} Y_{1,i,1} \end{pmatrix}, \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\rangle + O\left(\left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)}\right) \\
&= O\left(\left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)}\right).
\end{aligned} \tag{3.35}$$

Similarly, we have

$$\int_{\Omega} \left\langle \begin{pmatrix} \frac{1}{\varepsilon^2} e^{\sum_{j=1}^2 (U_{j,\mu} + \omega_{j,\mu} + u_{0,j})} \\ \frac{1}{\varepsilon^2} e^{\sum_{j=1}^2 (U_{j,\mu} + \omega_{j,\mu} + u_{0,j})} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle = O\left(\varepsilon^2 \mu^2 + \varepsilon^2 \mu^3 \left\| \begin{pmatrix} \omega_{1,\mu} \\ \omega_{2,\mu} \end{pmatrix} \right\|_{L^\infty(\Omega)}\right). \tag{3.36}$$

By (3.25), (3.34), (3.35) and (3.36), we conclude that

$$\int_{\Omega} \left\langle \begin{pmatrix} \Delta\omega_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu}+\omega_{2,\mu}+u_{0,2}} (1 - e^{U_{1,\mu}+\omega_{1,\mu}+u_{0,1}}) - \frac{1}{\varepsilon^2} e^{U_{2,\mu}+u_{0,2}} \\ \Delta\omega_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{1,\mu}+\omega_{1,\mu}+u_{0,1}} (1 - e^{U_{2,\mu}+\omega_{2,\mu}+u_{0,2}}) - \frac{1}{\varepsilon^2} e^{U_{1,\mu}+u_{0,1}} \end{pmatrix}, \begin{pmatrix} Y_{1,i,1} \\ Y_{2,i,1} \end{pmatrix} \right\rangle = O\left(\frac{\ln \mu}{\mu^{2-\frac{\alpha}{2}}}\right). \tag{3.37}$$

□

Lemma 3.5. *There exists $B > 0$, such that*

$$\begin{aligned}
& \int_{\Omega} \left\langle \begin{pmatrix} \Delta(U_{1,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{2,\mu}+\omega_{2,\mu}+u_{0,2}} (1 - e^{U_{1,\mu}+\omega_{1,\mu}+u_{0,1}}) - 8k\pi \\ \Delta(U_{2,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{1,\mu}+\omega_{1,\mu}+u_{0,1}} (1 - e^{U_{2,\mu}+\omega_{2,\mu}+u_{0,2}}) - 8k\pi \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle \\
&= \frac{8}{\mu^3 \rho_1} \left(\sum_{i=1}^k \rho_i \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{1,i,x}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\
&+ \frac{8}{\mu^3 \rho_1^*} \left(\sum_{i=1}^k \rho_i^* \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{2,i,x}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\
&+ B\varepsilon^2 \mu + O\left(\frac{\delta^2}{\mu^3}\right) + O\left(\frac{\ln \mu}{\mu^{4-\frac{\alpha}{2}}}\right)
\end{aligned} \tag{3.38}$$

Proof. As in the proof of Lemma (3.4), we firstly calculate

$$\begin{aligned}
& \int_{\Omega} \left\langle \left(\Delta(U_{1,\mu} + \omega_{1,\mu}) + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + \omega_{2,\mu} + u_{0,2}} (1 - e^{U_{1,\mu} + \omega_{1,\mu} + u_{0,1}}) - 8k\pi \right), \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle \\
&= \sum_{i=1}^k \int_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{2,i,\mathbf{x},\mu}} - 1) Y_{1,0} + e^{V_{x_i, \mu_i}} (e^{f_{1,i,\mathbf{x},\mu}}) Y_{2,0} \\
&+ \int_{\Omega} \frac{8\pi}{\theta_i} (Y_{1,0} + Y_{2,0}) + \frac{\int_{\Omega \setminus \Omega'} e^{\omega_{\mu}^* + u_{0,2}} Y_{1,0}}{\int_{\Omega} e^{\omega_{\mu}^* + u_{0,2}}} + \frac{\int_{\Omega \setminus \Omega'} e^{\omega_{\mu}^* + u_{0,1}} Y_{2,0}}{\int_{\Omega} e^{\omega_{\mu}^* + u_{0,1}}}
\end{aligned} \tag{3.39}$$

For $i = 1, \dots, n$, $j = 1, 2$, we write

$$e^{f_{j,i,\mathbf{x},\mu}(y)} - 1 = e^{f_{j,i,\mathbf{x}}(y)} - 1 + e^{f_{j,i,\mathbf{x},\mu}(y)} - e^{f_{j,i,\mathbf{x}}(y)} \tag{3.40}$$

Then

$$\begin{aligned}
& \int_{B_{d_i}(x_i)} \left(e^{V_{x_i, \mu_i}} (e^{f_{2,i,\mathbf{x},\mu}(y)} - 1) Y_{1,0} + e^{V_{x_i, \mu_i}} (e^{f_{1,i,\mathbf{x},\mu}(y)} - 1) Y_{2,0} \right) \\
&= \int_{B_{d_i}(x_i)} \left(e^{V_{x_i, \mu_i}} (e^{f_{2,i,\mathbf{x}}} - 1) Y_{1,0} + e^{V_{x_i, \mu_i}} (e^{f_{1,i,\mathbf{x}}} - 1) Y_{2,0} \right) + O\left(\frac{1}{\mu^4}\right)
\end{aligned} \tag{3.41}$$

A straightforward calculation shows that

$$\sum_{i=1}^k \int_{\Omega} \frac{8\pi}{\theta_i} (Y_{1,0} + Y_{2,0}) = -\frac{16\pi}{\mu} \sum_{i=1}^k \frac{1}{\theta_i} + O\left(\frac{\ln \mu}{\mu^5}\right). \tag{3.42}$$

and

$$\begin{aligned}
& \frac{\int_{\Omega \setminus \Omega'} e^{\omega_{\mu} + u_{0,2}} Y_{1,0}}{\int_{\Omega} e^{\omega_{\mu}^* + u_{0,2}}} + \frac{\int_{\Omega \setminus \Omega'} e^{\omega_{\mu}^* + u_{0,1}} Y_{2,0}}{\int_{\Omega} e^{\omega_{\mu}^* + u_{0,1}}} \\
&= -\frac{8}{\rho_1^* \mu^2} \left(1 + O\left(\frac{1}{\mu}\right)\right) \int_{\Omega \setminus \Omega'} e^{u_{0,2} + 8\pi \sum_{i=1}^k G(y, x_i)} \left(-\frac{1}{\mu} + O\left(\frac{1}{\mu^3}\right)\right) \\
&\quad - \frac{8}{\rho_1 \mu^2} \left(1 + O\left(\frac{1}{\mu^2}\right)\right) \int_{\Omega \setminus \Omega'} e^{u_{0,1} + 8\pi \sum_{i=1}^k G(y, x_i)} \left(-\frac{1}{\mu} + O\left(\frac{1}{\mu^3}\right)\right) \\
&= -\frac{8}{\rho_1^* \mu^3} \int_{\Omega \setminus \Omega'} e^{u_{0,2} + 8\pi \sum_{i=1}^k G(y, x_i)} - \frac{8}{\rho_1 \mu^3} \int_{\Omega \setminus \Omega'} e^{u_{0,1} + 8\pi \sum_{i=1}^k G(y, x_i)} + O\left(\frac{1}{\mu^4}\right)
\end{aligned} \tag{3.43}$$

Fix a small positive constant δ with $\delta \ll d_i$, $i = 1, \dots, k$. Note that $\Delta e^{f_{i,j,\mathbf{x}}} = e^{f_{i,j,\mathbf{x}}} |Df_{i,j,\mathbf{x}}|$, $i = 1, \dots, k$, $j = 1, 2$. By the symmetry and Taylor expansion on $(e^{f_{i,j,\mathbf{x}}} - 1)$, we obtain

$$\begin{aligned}
& \int_{B_{\delta}(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{2,i,\mathbf{x}}} - 1) Y_{1,0} + \int_{B_{\delta}(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{1,i,\mathbf{x}}} - 1) Y_{2,0} \\
&= O\left(\frac{1}{\mu} \int_{B_{\delta}(x_i)} e^{V_{x_i, \mu_i}} \left(|Df_{2,i,\mathbf{x}}(x_i)|^2 |y - x_i|^2 + |Df_{1,i,\mathbf{x}}(x_i)|^2 |y - x_i|^2 + |y - x_i|^4 \right)\right) \\
&= O\left(\frac{\delta^2}{\mu^3} + \frac{\ln \mu}{\mu^5}\right)
\end{aligned} \tag{3.44}$$

By this, we find that

$$\begin{aligned}
& \int_{B_\delta(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{2,i,x}} - 1) Y_{1,0} + \int_{B_\delta(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{1,i,x}} - 1) Y_{2,0} \\
&= \int_{B_{d_i}(x_i) \setminus B_\delta(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{2,i,x}} - 1) Y_{1,0} + \int_{B_{d_i}(x_i) \setminus B_\delta(x_i)} e^{V_{x_i, \mu_i}} (e^{f_{1,i,x}} - 1) Y_{2,0} + O\left(\frac{\delta^2}{\mu^3} + \frac{\ln \mu}{\mu^5}\right) \\
&= -\frac{8}{\mu^3} \int_{B_{d_i}(x_i) \setminus B_\delta(x_i)} \frac{e^{f_{2,i,x}} - 1}{|y - x_i|^4} - \frac{8}{\mu^3} \int_{B_{d_i}(x_i) \setminus B_\delta(x_i)} \frac{e^{f_{1,i,x}} - 1}{|y - x_i|^4} + O\left(\frac{\delta^2}{\mu^3} + \frac{\ln \mu}{\mu^5}\right)
\end{aligned} \tag{3.45}$$

For $i = 1, \dots, k$, we have

$$e^{u_{0,1}(y) + 8\pi \sum_{i=1}^k G(y, x_i)} = \frac{\rho_i}{|y - x_i|^4} e^{f_{1,i,x}(y)} \tag{3.46}$$

and

$$e^{u_{0,2}(y) + 8\pi \sum_{i=1}^k G(y, x_i)} = \frac{\rho_i^*}{|y - x_i|^4} e^{f_{2,i,x}(y)} \tag{3.47}$$

Recall that $\frac{1}{\theta_i} = \frac{1}{1 + (\mu_i d_i)^2}$. So, we have

$$\frac{1}{\theta_i} = \int_{\mathbb{R}^2 \setminus B_{d_i}(x_i)} \frac{1}{|y - x_i|^4} dy + O\left(\frac{1}{\mu^2}\right) \tag{3.48}$$

Combining (3.42) (3.43) (3.44) (3.45) (3.46) (3.47) and (3.48), we are led to

$$\begin{aligned}
& \int_{\Omega} \left\langle \begin{pmatrix} \Delta U_{1,\mu} + \frac{1}{\varepsilon^2} e^{U_{2,\mu} + u_{0,2}} \\ \Delta U_{2,\mu} + \frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1}} \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle \\
&= -\frac{8}{\mu^3 \rho_1^*} \left(\sum_{i=1}^k \rho_i^* \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{2,i,x}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\
&\quad - \frac{8}{\mu^3 \rho_1} \left(\sum_{i=1}^k \rho_i \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{1,i,x}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\
&\quad + O\left(\frac{\delta^2}{\mu^3} + \frac{1}{\mu^4}\right)
\end{aligned} \tag{3.49}$$

By the argument in Lemma 3.4,

$$\int_{\Omega} \left\langle \begin{pmatrix} -\frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1} + U_{2,\mu} + u_{0,2}} \\ -\frac{1}{\varepsilon^2} e^{U_{1,\mu} + u_{0,1} + U_{2,\mu} + u_{0,2}} \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle = B\varepsilon^2 \mu + O\left(\frac{1}{\mu^5}\right), \tag{3.50}$$

for some $B > 0$, and

$$\int_{\Omega} \left\langle \begin{pmatrix} \Delta \omega_{1,\mu} + \frac{1}{\varepsilon^2} R_{1,\mu} \\ \Delta \omega_{2,\mu} + \frac{1}{\varepsilon^2} R_{2,\mu} \end{pmatrix}, \begin{pmatrix} Y_{1,0} \\ Y_{2,0} \end{pmatrix} \right\rangle = O\left(\frac{\ln \mu}{\mu^{4 - \frac{\alpha}{2}}}\right), \tag{3.51}$$

where

$$R_{1,\mu} = e^{U_{2,\mu+u_0,2+\omega_{2,\mu}}}(1 - e^{U_{1,\mu+u_0,1+\omega_{1,\mu}}}) - e^{U_{2,\mu+u_0,2}}(1 - e^{U_{1,\mu+u_0,1}})$$

and

$$R_{2,\mu} = e^{U_{1,\mu+u_0,1+\omega_{1,\mu}}}(1 - e^{U_{2,\mu+u_0,2+\omega_{2,\mu}}}) - e^{U_{1,\mu+u_0,1}}(1 - e^{U_{2,\mu+u_0,2}}).$$

□

Proof of Theorem 1.1 From (3.30) and (3.38), we observe that (3.28) and (3.29) are equivalent to

$$DG_i^*(\mathbf{x}) = O\left(\frac{\ln \mu}{\mu^{2-\frac{\alpha}{2}}}\right), \quad i = 1, 2, \quad (3.52)$$

and

$$\begin{aligned} & \frac{8}{\mu^3 \rho_1} \left(\sum_{i=1}^k \rho_i \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{1,i}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\ & + \frac{8}{\mu^3 \rho_1^*} \left(\sum_{i=1}^k \rho_i^* \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{2,i}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) + B\varepsilon^2 \mu \quad (3.53) \\ & = \frac{1}{\mu^3} O\left(\left(|DG_1^*(\mathbf{x})|^2 + |DG_2^*(\mathbf{x})|^2\right) \ln \mu + \delta^2\right) + O\left(\frac{1}{\mu^5}\right) \end{aligned}$$

Since we assume $\mathcal{D}^{(2)}(\mathbf{q}) < 0$, for small $\delta > 0$, there exists \mathbf{x} close to \mathbf{q} such that

$$\begin{aligned} & \sum_{i=1}^k \left(\frac{\rho_i}{\rho_1} \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{1,i}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) + \frac{\rho_i^*}{\rho_1^*} \left(\int_{\Omega_i \setminus B_\delta(x_i)} \frac{e^{f_{2,i}} - 1}{|y - x_i|^4} - \int_{\mathbb{R}^2 \setminus \Omega_i} \frac{1}{|y - x_i|^4} \right) \right) \\ & + O(\delta^2) < 0. \quad (3.54) \end{aligned}$$

From this and (A.2), we find that (3.52) and (3.53) have a solution \mathbf{x}_ε and $(\mu_{1,\varepsilon}, \dots, \mu_{k,\varepsilon})$ satisfies

$$|DG_1^*(\mathbf{x})| + |DG_2^*(\mathbf{x})| \leq C \frac{\ln \mu}{\mu^{2-\frac{\alpha}{2}}}, \quad \mu_{i,\varepsilon} \in \left(\frac{\beta_0}{\sqrt{\varepsilon}}, \frac{\beta_1}{\sqrt{\varepsilon}}\right), \quad i = 1, \dots, k. \quad (3.55)$$

4 Appendix

In this section, we will discuss the invertibility of the linear operator L_μ . From our construction of the approximation solutions, we can split the associated linear operator L_μ into two parts \mathcal{L}_1 and \mathcal{L}_2 (see (4.3) below). Apply Theorem A.2 in [29] and Theorem B.1 in [28] to \mathcal{L}_1 and \mathcal{L}_2 respectively, we obtain the invertibility of L_μ .

Theorem 4.1. (1) *The operator $Q_\mu L_\mu$ is an isomorphism from $E_{\mu,2}$ to $F_{\mu,2}$.*

(2) If $\begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \in E_{\mu,2}$ and $\begin{pmatrix} h_1 \\ h_2 \end{pmatrix} \in F_{\mu,2}$ satisfies

$$L_\mu \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} = \begin{pmatrix} h_1 \\ h_2 \end{pmatrix}. \quad (4.1)$$

Then there exists a constant $C > 0$, independent of \mathbf{x} and μ , such that

$$\left\| \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \right\|_{L^\infty(\Omega)} + \left\| \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} \right\|_{X_{\alpha,\mu,2}} \leq C \ln \mu \left\| \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} \right\|_{Y_{\alpha,\mu,2}}. \quad (4.2)$$

Proof. Since $Z_{1,0} = Z_{2,0}$, $Z_{1,i,j} = Z_{2,i,j}$, $Y_{1,0} = Y_{2,0}$, and $Y_{1,i,j} = Y_{2,i,j}$, $i = 1, 2$, $j = 1, \dots, k$. We denote

$$E_\mu = \left\{ \omega \in X_{\alpha,\mu} : \int_\Omega \omega Z_{1,0} = \int_\Omega \omega Z_{1,i,j} = 0, i = 1, 2, j = 1, \dots, k. \right\}$$

and

$$F_\mu = \left\{ \omega \in Y_{\alpha,\mu} : \int_\Omega \omega Y_{1,0} = \int_\Omega \omega Y_{1,i,j} = 0, i = 1, 2, j = 1, \dots, k. \right\}.$$

We use the same notation for the project operator $Q_\mu : Y_\mu \rightarrow F_\mu$.

We can rewrite (4.1) we as

$$\begin{aligned} \mathcal{L}_1(\omega_1 + \omega_2) &:= \Delta(\omega_1 + \omega_2) + \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} (\omega_1 + \omega_2) = h_1 + h_2 \\ \mathcal{L}_2(\omega_1 - \omega_2) &:= \Delta(\omega_1 - \omega_2) - \sum_{i=1}^k 1_{B_{d_i}(x_i)} e^{V_{x_i, \mu_i}} (\omega_1 - \omega_2) = h_1 - h_2 \end{aligned} \quad (4.3)$$

By Theorem A.2 in [29], $Q_\mu \mathcal{L}_1$ is an isomorphism from E_μ to F_μ ; by Theorem B.1 in [28], \mathcal{L}_2 is an isomorphism from $X_{\alpha,\mu}$ to $Y_{\alpha,\mu}$. Furthermore, by Theorem A.2 in [29] and Theorem B.1 in [28] again, we obtain

$$\|\omega_1 + \omega_2\|_{L^\infty(\Omega)} + \|\omega_1 + \omega_2\|_{X_{\alpha,\mu}} \leq C \ln \mu \|h_1 + h_2\|_{Y_{\alpha,\mu}} \quad (4.4)$$

and

$$\|\omega_1 - \omega_2\|_{L^\infty(\Omega)} + \|\omega_1 - \omega_2\|_{X_{\alpha,\mu}} \leq C \ln \mu \|h_1 - h_2\|_{Y_{\alpha,\mu}} \quad (4.5)$$

which imply (4.2). □

References

- [1] O. Aharony, O. Bergman, D. Jafferis, and J. Maldacena. $N = 6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals. *J. High Energy Phys.*, (10):091, 38, 2008.
- [2] E. B. Bogomolny. The stability of classical solutions. *Jadernaja Fiz.*, 24(4):861–870, 1976.

- [3] L. Caffarelli and Y. Yang. Vortex condensation in the Chern-Simons Higgs model: an existence theorem. *Comm. Math. Phys.*, 168(2):321–336, 1995.
- [4] H. Chan, C. Fu, and C.-S. Lin. Non-topological multi-vortex solutions to the self-dual Chern-Simons-Higgs equation. *Comm. Math. Phys.*, 231(2):189–221, 2002.
- [5] J. Chern, Z. Chen, and C.S. Lin. Uniqueness of topological solutions and the structure of solutions for the Chern-Simons system with two Higgs particles. *Comm. Math. Phys.*, 296(2):323–351, 2010.
- [6] J. Chern and S. Yang. The non-topological fluxes of a two-particle system in the Chern-Simons theory. *J. Differential Equations*, 256(10):3417–3439, 2014.
- [7] K. Choe. Uniqueness of the topological multivortex solution in the self-dual Chern-Simons theory. *J. Math. Phys.*, 46(1):012305, 22, 2005.
- [8] K. Choe. Multiple existence results for the self-dual Chern-Simons-Higgs vortex equation. *Comm. Partial Differential Equations*, 34(10-12):1465–1507, 2009.
- [9] K. Choe and N. Kim. Blow-up solutions of the self-dual Chern-Simons-Higgs vortex equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 25(2):313–338, 2008.
- [10] K. Choe, N. Kim, and C.S. Lin. Existence of self-dual non-topological solutions in the Chern-Simons Higgs model. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 28(6):837–852, 2011.
- [11] M. Del Pino, P. Esposito, P. Figueroa, and M. Musso. Nontopological condensates for the self-dual Chern-Simons-Higgs model. *Communications on Pure and Applied Mathematics*, 2014.
- [12] G. V. Dunne. Aspects of Chern-Simons theory. In *Aspects topologiques de la physique en basse dimension/Topological aspects of low dimensional systems (Les Houches, 1998)*, pages 177–263. EDP Sci., Les Ulis, 1999.
- [13] J. Dziarmaga. Low energy dynamics of $[U(1)]^N$ chern-simons solitons. *Phys. Rev. D*, 49:5469–5479, May 1994.
- [14] S. Gudnason. Non-abelian Chern-Simons vortices with generic gauge groups. *Nuclear Phys. B*, 821(1-2):151–169, 2009.
- [15] S. Gudnason. Fractional and semi-local non-Abelian Chern-Simons vortices. *Nuclear Phys. B*, 840(1-2):160–185, 2010.
- [16] C.R. Hagen. Parity conservation in Chern-Simons theories and the anyon interpretation. *Phys.Rev.Lett.*, 68:3821–3825, 1992.
- [17] X. Han, H. Huang, and C.-S. Lin. Bubbling solutions for a skew-symmetric chernsimons system in a torus. *Journal of Functional Analysis*, 273(4):1354 – 1396, 2017.

- [18] J. Hong, Y. Kim, and P. Pac. Multivortex solutions of the abelian Chern-Simons-Higgs theory. *Phys. Rev. Lett.*, 64(19):2230–2233, 1990.
- [19] H. Huang and C.-S. Lin. Uniqueness of non-topological solutions for the Chern-Simons system with two Higgs particles. *Kodai Math. J.*, 37(2):274–284, 2014.
- [20] H. Huang and L. Zhang. The domain geometry and the bubbling phenomenon of rank two gauge theory. *Comm. Math. Phys.*, 349(1):393–424, 2017.
- [21] R. Jackiw and E. Weinberg. Self-dual Chern-Simons vortices. *Phys. Rev. Lett.*, 64:2234–2237, May 1990.
- [22] A. Jaffe and C. Taubes. *Vortices and monopoles*, volume 2 of *Progress in Physics*. Birkhäuser, Boston, Mass., 1980. Structure of static gauge theories.
- [23] C. Kim, C. Lee, P. Ko, B. Lee, and H. Min. Schrödinger fields on the plane with $[U(1)]^N$ Chern-Simons interactions and generalized self-dual solitons. *Phys. Rev. D*, 48:1821–1840, Aug 1993.
- [24] C.-S. Lin, A. Ponce, and Y. Yang. A system of elliptic equations arising in Chern-Simons field theory. *Journal of Functional Analysis*, 247(2):289 – 350, 2007.
- [25] C.-S. Lin and J. Prajapat. Vortex condensates for relativistic abelian Chern-Simons model with two Higgs scalar fields and two gauge fields on a torus. *Comm. Math. Phys.*, 288(1):311–347, 2009.
- [26] C.-S. Lin and C. Wang. Elliptic functions, Green functions and the mean field equations on tori. *Ann. Math. (2)*, 172(2):911–954, 2010.
- [27] C.-S. Lin and C. Wang. On the minimality of extra critical points of Green functions on flat tori. *Int. Math. Res. Not. IMRN*, (18):5591–5608, 2017.
- [28] C.-S. Lin and S. Yan. Bubbling solutions for relativistic abelian Chern-Simons model on a torus. *Comm. Math. Phys.*, 297(3):733–758, 2010.
- [29] C.-S. Lin and S. Yan. Existence of bubbling solutions for Chern-Simons model on a torus. *Arch. Ration. Mech. Anal.*, 207(2):353–392, 2013.
- [30] C.-S. Lin and S. Yan. On condensate of solutions for the ChernSimonsHiggs equation. *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, 34(5):1329 – 1354, 2017.
- [31] C.-S. Lin and L. Zhang. On liouville systems at critical parameters, part 1: One bubble. *Journal of Functional Analysis*, 264(11):2584 – 2636, 2013.
- [32] G. S. Lozano, D. Marqués, E. F. Moreno, and F. A. Schaposnik. Non-abelian Chern-Simons vortices. *Phys. Lett. B*, 654(1-2):27–34, 2007.
- [33] M. K. Prasad and C. Sommerfield. Exact classical solution for the ’t hooft monopole and the julia-zee dyon. *Phys. Rev. Lett.*, 35:760–762, Sep 1975.

- [34] S. Spielman, K. Fesler, C. B. Eom, T. H. Geballe, M. M. Fejer, and A. Kapitulnik. Test for nonreciprocal circular birefringence in $\text{YBa}_2\text{Cu}_3\text{O}_7$ thin films as evidence for broken time-reversal symmetry. *Phys. Rev. Lett.*, 65:123–126, Jul 1990.
- [35] J. Spruck and Y. Yang. The existence of nontopological solitons in the self-dual Chern-Simons theory. *Comm. Math. Phys.*, 149(2):361–376, 1992.
- [36] G. Tarantello. Multiple condensate solutions for the Chern-Simons-Higgs theory. *J. Math. Phys.*, 37(8):3769–3796, 1996.
- [37] G. Tarantello. Uniqueness of selfdual periodic Chern-Simons vortices of topological-type. *Calc. Var. Partial Differential Equations*, 29(2):191–217, 2007.
- [38] G. Tarantello. *Selfdual gauge field vortices*. Progress in Nonlinear Differential Equations and their Applications, 72. Birkhäuser Boston, Inc., Boston, MA, 2008. An analytical approach.
- [39] F. Wilczek. Disassembling anyons. *Phys. Rev. Lett.*, 69:132–135, Jul 1992.
- [40] Y. Yang. *Solitons in field theory and nonlinear analysis*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2001.