

HFLAV τ branching fractions fit and measurements of $|V_{us}|$ with τ lepton data

A. Lusiani^{1*},

¹ Scuola Normale Superiore and INFN sezione di Pisa, Italy

* alberto.lusiani@pi.infn.it

April 5, 2022



*Proceedings for the 15th International Workshop on Tau Lepton Physics,
Amsterdam, The Netherlands, 24-28 September 2018
scipost.org/SciPostPhysProc.Tau2018*

Abstract

We report the status of the Heavy Flavour Averaging Group (HFLAV) averages of the τ lepton measurements. We then update the latest published HFLAV global fit of the τ lepton branching fractions (Spring 2017) with recent results by *BABAR*. We use the fit results to update the Cabibbo-Kobayashi-Maskawa (CKM) matrix element $|V_{us}|$ measurements with the τ branching fractions. We combine the direct τ branching fraction measurements with indirect predictions using kaon branching fractions measurements to improve the determination of $|V_{us}|$ using τ branching fractions. The $|V_{us}|$ determinations based on the inclusive branching fraction of τ to strange final states are about 3σ lower than the $|V_{us}|$ determination from the CKM matrix unitarity.

Contents

1	Introduction	2
2	New τ branching fraction measurements	2
3	$ V_{us} $ determination including the 2018 <i>BABAR</i> results	2
4	τ branching fraction predictions from kaon measurements	3
5	Consistency of $ V_{us} $ with the CKM matrix unitarity	3
6	Conclusions	4
	References	4

1 Introduction

The τ subgroup of the Heavy Flavour Averaging Group (HFLAV) provides a global fit of the τ branching fractions, the lepton universality tests and the $|V_{us}|$ determination based on τ measurements. The latest published report for the τ lepton is labelled ‘‘Spring 2017’’ [1]. A version of the HFLAV τ branching fractions fit with unitarity constraint is published on the Review of Particle Physics [2] (RPP). There are additional minor differences between the two fits [1, 3]. The fit results are used to test lepton universality and to compute $|V_{us}|$ [1].

The HFLAV-Tau group collects and combines also a list of upper limits set by searches of lepton-flavour-violating τ decays [1].

In the following, we update the HFLAV-Tau global fit inputs with two *BABAR* measurements that became public in 2018 [4, 5] and we update the $|V_{us}|$ determinations based on τ data. The new results have a negligible effect on the lepton universality tests.

Finally, we add to the fit input measurements of three τ branching fractions that are indirectly determined using measurements of kaon branching fractions [6], in order to improve the precision on $|V_{us}|$.

2 New τ branching fraction measurements

Since the last HFLAV report, *BABAR* published [4] a measurement of

$$B(\tau^- \rightarrow K^- K^0 \nu_\tau) = (14.78 \pm 0.22 \pm 0.40) \cdot 10^{-4}$$

and presented [5] preliminary measurements of

$$\begin{aligned} B(\tau^- \rightarrow K^- \nu_\tau) &= (7.174 \pm 0.033 \pm 0.213) \cdot 10^{-3} , \\ B(\tau^- \rightarrow K^- \pi^0 \nu_\tau) &= (5.054 \pm 0.021 \pm 0.148) \cdot 10^{-3} , \\ B(\tau^- \rightarrow K^- 2\pi^0 \nu_\tau \text{ (ex. } K^0)) &= (6.151 \pm 0.117 \pm 0.338) \cdot 10^{-4} , \\ B(\tau^- \rightarrow K^- 3\pi^0 \nu_\tau \text{ (ex. } K^0, \eta)) &= (1.246 \pm 0.164 \pm 0.238) \cdot 10^{-4} , \\ B(\tau^- \rightarrow \pi^- 3\pi^0 \nu_\tau \text{ (ex. } K^0, \eta)) &= (1.168 \pm 0.006 \pm 0.038) \cdot 10^{-2} , \\ B(\tau^- \rightarrow \pi^- 4\pi^0 \nu_\tau \text{ (ex. } K^0, \eta)) &= (9.020 \pm 0.400 \pm 0.652) \cdot 10^{-4} . \end{aligned}$$

3 $|V_{us}|$ determination including the 2018 *BABAR* results

We add the measurements listed in the previous section to the HFLAV-Tau global fit, removing a former *BABAR* measurement of $B(\tau^- \rightarrow K^- \pi^0 \nu_\tau)$ [7] that has been superseded [5]. The new measurements of the branching fractions τ decaying to a kaon and 0–4 π^0 's improve the experimental resolution on several modes that most contribute to the uncertainty on $|V_{us}|$.

We compute $|V_{us}|_{\tau s}$ using the total branching fraction of the τ to strange final states following Ref. [8]:

$$|V_{us}|_{\tau s} = \sqrt{R_s / \left[\frac{R_{VA}}{|V_{ud}|^2} - \delta R_{\text{theory}} \right]} = 0.2195 \pm 0.0019 ,$$

where $|V_{ud}| = 0.97420 \pm 0.00021$ [9], R_s and R_{VA} are the τ hadronic partial widths to strange and to non-strange hadronic final states (Γ_s and Γ_{had}) divided by the universality-improved branching fraction $B(\tau \rightarrow e\nu\bar{\nu}) = B_e^{\text{uni}} = (17.815 \pm 0.023)\%$ [1,3], and the SU(3)-breaking term $\delta R_{\text{theory}} = 0.242 \pm 0.033$ is computed using inputs from Ref. [8] and $m_s = (95.00 \pm 6.70)$ MeV [2] (the uncertainties on m_s have been symmetrized).

We compute also

$$|V_{us}|_{\tau K/\pi} = |V_{ud}| \frac{f_\pi}{f_K} \frac{1 - m_\pi^2/m_\tau^2}{1 - m_K^2/m_\tau^2} \sqrt{\frac{B(\tau^- \rightarrow K^- \nu_\tau)}{B(\tau^- \rightarrow \pi^- \nu_\tau)} \frac{1}{R_{\tau K/\tau\pi}}} = 0.2241 \pm 0.0016 ,$$

where $f_K/f_\pi = 1.1930 \pm 0.0030$ from the FLAG 2016 Lattice averages with $N_f = 2 + 1 + 1$ [10–13] (the same value persists in the FLAG 2017 web update). The radiative correction term is $R_{\tau K/\tau\pi} = 0.9930 \pm 0.0035$ [1], and the other parameters are taken from the Review of Particle Physics (RPP) 2018 [2].

Averaging the two above $|V_{us}|$ determinations, we obtain $|V_{us}|_\tau = 0.2223 \pm 0.0014$.

4 τ branching fraction predictions from kaon measurements

Assuming the validity of the Standard Model (SM), three τ branching fractions have been computed using the precisely measured $K_{\ell 2}$ and $K_{\ell 3}$ branching fractions and the measured $\tau^- \rightarrow (K\pi)^- \nu_\tau$ spectra [6]:

$$\begin{aligned} B(\tau^- \rightarrow K^- \nu_\tau) &= (0.713 \pm 0.003)\% , \\ B(\tau^- \rightarrow K^- \pi^0 \nu_\tau) &= (0.471 \pm 0.018)\% , \\ B(\tau^- \rightarrow K^0 \pi^- \nu_\tau) &= (0.857 \pm 0.030)\% . \end{aligned}$$

The uncertainties on the last two results are fully correlated. It has been observed [6, 14] that all the above indirect values are higher than the corresponding directly measured τ branching fractions. If the indirect values replace the direct ones, $|V_{us}| = 0.2207 \pm 0.027$ [6].

We add the kaon-indirect determinations of the three above τ branching fractions to the data set used in the previous section in order to obtain improved calculations of $|V_{us}|$:

$$\begin{aligned} |V_{us}|_{\tau s} &= 0.2202 \pm 0.0018 , \\ |V_{us}|_{\tau K/\pi} &= 0.22596 \pm 0.00099 , \\ |V_{us}|_\tau &= 0.22475 \pm 0.00089 . \end{aligned}$$

5 Consistency of $|V_{us}|$ with the CKM matrix unitarity

Assuming the CKM matrix unitarity,

$$|V_{us}|_{\text{uni}} = \sqrt{1 - |V_{ud}|^2 - |V_{ub}|^2} = 0.22565 \pm 0.00089 ,$$

using $|V_{ud}| = 0.97420 \pm 0.00021$ [9] and $|V_{ub}| = (0.3940 \pm 0.0360) \cdot 10^{-2}$ [2]. Table 1 summarizes the residuals, expressed as numbers of standard deviations, of the above mentioned $|V_{us}|$ determinations with respect to the $|V_{us}|$ computation from the CKM matrix unitarity. $|V_{us}|$ computed with the τ -inclusive method is significantly lower, but the significance of the discrepancy is mildly reduced alongside a mild progress in the experimental resolution.

Table 1: Deviations of $|V_{us}|$ computed with τ data with respect to $|V_{us}|$ obtained with CKM unitarity. The second and third row use the $|V_{us}|$ determinations performed above.

	$\Delta V_{us} _{\tau s}$ [σ]	$\Delta V_{us} _{\tau K/\pi}$ [σ]	$\Delta V_{us} _{\tau}$ [σ]
HFLAV Spring 2017	-3.0	-1.0	-2.3
HFLAV + <i>BABAR</i> 2018	-2.9	-0.8	-2.0
HFLAV + <i>BABAR</i> + kaon predictions	-2.7	0.2	-0.7

6 Conclusions

Figure 1 reports the $|V_{us}|_{\tau s}$ determinations described above, a determination of $|V_{us}|_{\tau s}$ obtained replacing some τ branching fractions measurements with the indirect predictions based on kaon branching fractions [6], and other more complex determinations that use the τ spectral functions [15] and Lattice QCD techniques [16]. Updates on the last two determinations have been presented at the Tau 2018 workshop [17]. The last four determinations use an older and in some cases partial set of experimental τ branching fractions measurements.

The τ based $|V_{us}|$ determinations use the $|V_{ud}|$ measurements as input. The dependence on $|V_{ud}|$ is however very small, and there is in first approximation negligible correlation between $|V_{us}|$ and $|V_{ud}|$ when doing a simultaneous fit. Figure 2 shows the results of a $|V_{ud}|-|V_{us}|$ simultaneous fit on the τ measurements corresponding to the HFLAV Spring 2017 fit and the *BABAR* 2018 results. The fit results are:

$$\begin{aligned}
 |V_{ud}| &= 0.97420 \pm 0.00021 \\
 |V_{us}| &= 0.2223 \pm 0.0014 \\
 |V_{ud}| - |V_{us}| \text{ correlation} &= 0.035
 \end{aligned}$$

Tables 2 and 3 report the contributions to the $|V_{us}|_{\tau s}$ uncertainty before and after the *BABAR* 2018 results. The largest contributions come from the τ branching fractions to strange final states and from the theory. The *BABAR* 2018 measurements reduced significantly several large contributions. High multiplicity τ decays to strange final states dominate the $|V_{us}|_{\tau s}$ uncertainty. The Belle II super flavour factory will offer the opportunity to improve the experimental precision on the τ strange branching fractions. More precise τ branching fractions and spectral function measurements will help improving also the theory uncertainty.

References

- [1] Y. Amhis *et al.*, *Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2016*, Eur. Phys. J. **C77**(12), 895 (2017), doi:10.1140/epjc/s10052-017-5058-4, 1612.07233.
- [2] M. Tanabashi *et al.*, *Review of Particle Physics*, Phys. Rev. **D98**(3), 030001 (2018), doi:10.1103/PhysRevD.98.030001.




Figure 1: $|V_{us}|_{\tau s}$ determinations obtained in this document, from the top: $|V_{us}|_{\text{uni}}$, $|V_{us}|_{\tau s}$ with the HFLAV Spring 2017 fit, after adding the *BABAR* 2018 data, after adding both the *BABAR* 2018 and the kaon indirect predictions, from Ref. [6], from Ref. [15], and two determinations from Ref. [16].




Figure 2: Results of a $|V_{ud}|-|V_{us}|$ simultaneous fit. The bands describe the constraints corresponding to the $|V_{ud}|$ measurement, the $|V_{us}|_{\tau s}$ and the $|V_{us}|_{\tau K/\pi}$ determinations that use the τ measurements. The oblique line corresponds to the CKM matrix unitarity constraint. The ellipse corresponds to 1σ uncertainty on the $|V_{ud}|$ and $|V_{us}|$ fit results.

Table 2: Contributions to the $|V_{us}|_{\tau s}$ uncertainty in percent before the *BABAR* 2018 results.

$\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau$ (ex. K^0)	0.3963	
$K^- 2\pi^0 \nu_\tau$ (ex. K^0)	0.3789	
$K^- 3\pi^0 \nu_\tau$ (ex. K^0, η)	0.3714	
$\bar{K}^0 h^- h^- h^+ \nu_\tau$	0.3478	
$K^- \pi^0 \nu_\tau$	0.2561	
$K^- \pi^- \pi^+ \pi^0 \nu_\tau$ (ex. K^0, ω, η)	0.2456	
$\pi^- \bar{K}^0 \nu_\tau$	0.2424	
$\pi^- \bar{K}^0 \pi^0 \nu_\tau$	0.2219	
$K^- \nu_\tau$	0.1646	
$K^- \omega \nu_\tau$	0.1585	
$K^- \pi^- \pi^+ \nu_\tau$ (ex. K^0, ω)	0.1157	
$\pi^- \bar{K}^0 \eta \nu_\tau$	0.0256	
$K^- \pi^0 \eta \nu_\tau$	0.0200	
$K^- \eta \nu_\tau$	0.0138	
$K^- \phi \nu_\tau$ ($\phi \rightarrow K^+ K^-$)	0.0138	
$K^- \phi \nu_\tau$ ($\phi \rightarrow K_S^0 K_L^0$)	0.0096	
$K^- 2\pi^- 2\pi^+ \nu_\tau$ (ex. K^0)	0.0021	
$K^- 2\pi^- 2\pi^+ \pi^0 \nu_\tau$ (ex. K^0)	0.0010	
$\tau \rightarrow$ non-strange	0.0896	
B_e^{univ}	0.0045	
theory	0.4861	

- [3] A. Lusiani, *HFAG 2016 and PDG 2016 τ lepton averages and $|V_{us}|$ determination from τ data*, Nucl. Part. Phys. Proc. **287-288**, 29 (2017), doi:10.1016/j.nuclphysbps.2017.03.038.
- [4] J. P. Lees *et al.*, *Measurement of the spectral function for the $\tau^- \rightarrow K^- K_S \nu_\tau$ decay*, Phys. Rev. **D98**(3), 032010 (2018), doi:10.1103/PhysRevD.98.032010, 1806.10280.
- [5] T. Lueck, *Recent results on τ -lepton decays with the BABAR detector*, Talk given at the 'XXXIX International Conference On High Energy Physics, Seoul, South Korea' (2018).
- [6] M. Antonelli, V. Cirigliano, A. Lusiani and E. Passemar, *Predicting the τ strange branching ratios and implications for V_{us}* , JHEP **10**, 070 (2013), doi:10.1007/JHEP10(2013)070, 1304.8134.
- [7] B. Aubert *et al.*, *Measurement of the $\tau^- \rightarrow K^- \pi^0 \nu_\tau$ branching fraction*, Phys. Rev. **D76**, 051104 (2007), doi:10.1103/PhysRevD.76.051104, 0707.2922.
- [8] E. Gamiz, M. Jamin, A. Pich, J. Prades and F. Schwab, *$|V(us)|$ and $m(s)$ from hadronic tau decays*, Nucl. Phys. Proc. Suppl. **169**, 85 (2007), doi:10.1016/j.nuclphysbps.2007.02.053, hep-ph/0612154.
- [9] J. Hardy and I. S. Towner, *$|V_{ud}|$ from nuclear β decays*, PoS **CKM2016**, 028 (2016), doi:10.22323/1.291.0028.
- [10] S. Aoki *et al.*, *Review of lattice results concerning low-energy particle physics*, Eur. Phys. J. **C77**, 112 (2017), doi:10.1140/epjc/s10052-016-4509-7, See also <http://itpwiki.unibe.ch/flag/>, 1607.00299.
- [11] A. Bazavov *et al.*, *Charmed and light pseudoscalar meson decay constants from four-flavor lattice QCD with physical light quarks*, Phys. Rev. **D90**(7), 074509 (2014), doi:10.1103/PhysRevD.90.074509, 1407.3772.
- [12] R. J. Dowdall, C. T. H. Davies, G. P. Lepage and C. McNeile, *V_{us} from π and K decay constants in full lattice QCD with physical u , d , s and c quarks*, Phys. Rev. **D88**, 074504 (2013), doi:10.1103/PhysRevD.88.074504, 1303.1670.
- [13] N. Carrasco *et al.*, *Leptonic decay constants f_K, f_D , and f_{D_s} with $N_f = 2 + 1 + 1$ twisted-mass lattice QCD*, Phys. Rev. **D91**(5), 054507 (2015), doi:10.1103/PhysRevD.91.054507, 1411.7908.
- [14] A. Pich, *Precision Tau Physics*, Prog. Part. Nucl. Phys. **75**, 41 (2014), doi:10.1016/j.pnpnp.2013.11.002, 1310.7922.
- [15] R. J. Hudspith, R. Lewis, K. Maltman and J. Zanotti, *A resolution of the inclusive flavor-breaking $\tau |V_{us}|$ puzzle*, Phys. Lett. **B781**, 206 (2018), doi:10.1016/j.physletb.2018.03.074, 1702.01767.
- [16] P. Boyle, R. J. Hudspith, T. Izubuchi, A. Jüttner, C. Lehner, R. Lewis, K. Maltman, H. Ohki, A. Portelli and M. Spraggs, *$|V_{us}|$ determination from inclusive strange tau decay and lattice HVP*, EPJ Web Conf. **175**, 13011 (2018), doi:10.1051/epjconf/201817513011, 1803.07228.
- [17] K. Maltman, *The status of the inclusive tau determination of V_{us}* , Talk given at the '15th International Workshop on Tau Lepton Physics, Amsterdam, The Netherlands' (2018).

Table 3: Contributions to the $|V_{us}|_{\tau S}$ uncertainty in percent after the *BABAR* 2018 results.

$\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau$ (ex. K^0)	0.3931	
$\bar{K}^0 h^- h^- h^+ \nu_\tau$	0.3450	
$K^- \pi^- \pi^+ \pi^0 \nu_\tau$ (ex. K^0, ω, η)	0.2436	
$\pi^- \bar{K}^0 \nu_\tau$	0.2372	
$\pi^- \bar{K}^0 \pi^0 \nu_\tau$	0.2200	
$K^- \omega \nu_\tau$	0.1572	
$K^- \pi^0 \nu_\tau$	0.1554	
$K^- \nu_\tau$	0.1459	
$K^- \pi^- \pi^+ \nu_\tau$ (ex. K^0, ω)	0.1147	
$K^- 2\pi^0 \nu_\tau$ (ex. K^0)	0.0460	
$K^- 3\pi^0 \nu_\tau$ (ex. K^0, η)	0.0449	
$\pi^- \bar{K}^0 \eta \nu_\tau$	0.0254	
$K^- \pi^0 \eta \nu_\tau$	0.0198	
$K^- \eta \nu_\tau$	0.0137	
$K^- \phi \nu_\tau$ ($\phi \rightarrow K^+ K^-$)	0.0136	
$K^- \phi \nu_\tau$ ($\phi \rightarrow K_S^0 K_L^0$)	0.0095	
$K^- 2\pi^- 2\pi^+ \nu_\tau$ (ex. K^0)	0.0021	
$K^- 2\pi^- 2\pi^+ \pi^0 \nu_\tau$ (ex. K^0)	0.0010	
$\tau \rightarrow$ non-strange	0.0855	
B_e^{univ}	0.0045	
theory	0.4863	