

Star formation in outer rings of S0 galaxies.

I. NGC 6534 and MCG 11-22-015.

O. Sil'chenko¹, I. Kostiuk², A. Burenkov², and H. Parul³

¹ Sternberg Astronomical Institute of the Lomonosov Moscow State University, University av. 13, 119991 Russia
e-mail: olga@sai.msu.su

² Special Astrophysical Observatory of the Russian Academy of Sciences, Nizhnij Arkhyz, 369167 Russia
e-mail: kostiuk@sao.ru, ban@sao.ru

³ Department of Physics and Astronomy, University of Alabama, P.O. Box 870324, Tuscaloosa, AL 35487-0324, USA
e-mail: hparul@crimson.ua.edu

Received ..., 2018; accepted ..., 2018

ABSTRACT

Aims. Though S0 galaxies are usually thought to be ‘red and dead’, they often demonstrate star formation organized in ring structures. We try to clarify the nature of this phenomenon and its difference from star formation in spiral galaxies. Two early-type galaxies with outer rings, NGC 6534 and MCG 11-22-015, are selected to be studied.

Methods. After inspecting the gas excitation in the rings using the Baldwin-Phillips-Terlevich method (Baldwin et al. 1981), we estimate the star formation rates (SFR) in the two outer rings of our galaxies by using several SFR indicators derived from narrow-band photometry in the H α emission line and archival GALEX ultraviolet images of the galaxies.

Results. The ionized gas is excited by young stars in the ring of NGC 6534 and partly by shocks – in MCG 11-22-015. The oxygen abundances in the HII regions of the rings are close to solar. The derived SFRs allow to qualitatively restore star formation histories (SFH) in the rings: in NGC 6534 the SFH is flat during the last 100-200 Myr, and in MCG 11-22-015 the star formation has started only a few Myr ago. We suggest that the rings in NGC 6534 and MCG 11-22-015 have different natures: the former is a resonant one supplied with gas perhaps through tidal effects, and the latter has been produced by a satellite accretion. Recent outer gas accretion is implied in both cases.

Key words. galaxies: structure – galaxies: evolution – galaxies, elliptical and lenticular

1. Introduction

Early-type reddish galaxies possessing large-scale detached outer stellar rings were first noted by de Vaucouleurs (1959). Boris A. Vorontsov-Velyaminov, during his compilation of the ‘Morphological Catalogue of Galaxies’ (MCG), had found the galaxies with large-scale rings to be so numerous that he proposed to introduce a special branch into the Hubble’s fork to describe them (Vorontsov-Velyaminov 1960). Recent surveys reveal outer stellar rings to be found mostly in S0 galaxies – according to Comerón et al. (2014), up to 50% S0–S0/a galaxies have outer rings. A similar fraction of ring structures in early-type disk galaxies, $53 \pm 5\%$, was noted by Laurikainen et al. (2011).

Rings are usually thought to be mostly of resonance nature: they represent the result of gas accumulation at Lindblad resonances of large-scale bars and of consequent star formation. However, not all galaxies with outer rings have large-scale bars; and not all S0s with outer rings demonstrate any gas presence and star formation in the rings. Our estimates based on the listing of the ARRAKIS catalogue (Comerón et al. 2014) and on the GALEX imaging data reveal that about half of all outer rings in S0s have had star formation during the last 100-200 Myr (Kostiuk & Sil’chenko 2015). The problem of gas sources to feed starforming rings in S0 galaxies is still unsolved; and nothing is known about the star formation histories of these spectacular structures. In this Letter we will consider NGC 6534 and

Table 1. Global parameters of the galaxies

Galaxy	NGC 6534	MCG 11-22-015
R_{25} , arcsec (LEDA)	28	24
R_{ring} (outer edge), arcsec	19×13	20×18
M_H (NED)	–23.5	–23.7
M_{NUV} , (our estimate)	–17.2	–16.1
$(u-r)_0$ (SDSS/DR9)	2.55	2.63
V_r (NED), km · s ^{–1}	8118	8064
Distance, Mpc (NED)	110	109
Scale, kpc per arcsec (NED)	0.518	0.514
PA_{phot} (LEDA)	16.5°	75°
PA_{ring} (our estimate)	21.7°	72.5°

MCG 11-22-015 – two early-type disk galaxies belonging to the red sequence with outer rings; their global properties are given in Table 1. The optical-band rings and first spectral results for these galaxies were reported by one of us long ago (Kostiuk 1975; Kostiuk et al. 1981). Later, after the GALEX data releases, we have detected there UV-rings (Ilyina & Sil’chenko 2011) with the radii corresponding to the optical rings. Now we will present new spectral data as well as SFR estimates obtained from the UV images and from our own H α narrow-band imaging.

2. Observations and star formation rate calibrations

We exposed both galaxies with the SCORPIO reducer of the Russian 6m telescope (Afanasiev & Moiseev 2005) operating in the imaging mode, through the narrow-band filter centered onto the redshifted $H\alpha$ line (both galaxies have close radial velocities) and in the neighbouring continuum. A spectrophotometric standard, AGK+81d266, was also exposed the same night through the FN674 filter. We also used the SCORPIO reducer to obtain spectra. We obtained green- and red-band medium-resolution spectra for NGC 6534 and only a red spectrum for MCG 11-22-015. The details of the observations are given in Table 2.

To estimate the SFR over various temporal scales, we have used the GALEX images in the NUV and FUV bands and our calibrated $H\alpha$ images; the formulae for the SFR calculations have been taken from Hao et al. (2011) and from the review by Kennicutt & Evans (2012).

3. The structure of NGC 6534 and MCG 11-22-015.

To study the large-scale structures of the galaxies, we have used the imaging data from the SDSS/DR9. We have made an isophotal analysis, and then we have calculated azimuthally averaged surface brightness profiles by integrating the fluxes within elliptical annuli with running major axes and with a fixed ellipticity and major-axis orientation determined by the outermost isophote measurements. The results are shown in Fig. 1 and Fig. 2, together with the colour maps $g - r$. One can see some differences in the whole structure and ring visibility in NGC 6534 and MCG 11-22-015. NGC 6534 is moderately inclined to our line of sight as demonstrated by the elongated outer isophotes; the outer ring matches the ellipticity of the outermost isophotes being evidently a round structure inside the exponential stellar disk. However, the inner isophotes, $R \leq 8''$, are very round; taking in mind quasi-exponential surface-brightness profile of the inner part of NGC 6534 we can suspect a pseudobulge containing an inner stellar ring slightly elongated orthogonally with respect to the outer isophotes and to the outer ring. MCG 11-22-015, on the contrary, has rather round outer isophotes and two local ellipticity maxima at $R = 5''$ and at $R = 14''$. The latter radius corresponds roughly to the outer ring, so if we assume that the galaxy is seen face-on we must recognize the elliptical shape of its outer ring. The surface-brightness profiles show the prominent ring in NGC 6534 overposed onto the regular exponential disk ($\mu_{r,0} = 19.4$, $h = 9.8''$, or 5.1 kpc) and a Type-II disk, according to the Freeman (1970) classification, in MCG 11-22-015, with the scalelength of $4''.2$, or 2.2 kpc. The elliptical ring in MCG 11-22-015 represents a boundary between the elongated lens and the outer exponential disk. Despite the structural differences, the rings in both galaxies are bluer than the underlying disks, in accordance with the fact that the rings were initially noted as UV-bright features (Ilyina & Sil'chenko 2011).

4. Corotating gas in NGC 6534

It is only for NGC 6534 that we have obtained the spectrum in the green with a spectral resolution of 2 Å, to determine the stellar and ionized-gas rotation curves and velocity dispersion. The velocities are shown in Fig. 3. The gas in NGC 6534 rotates strictly with the stellar component, and we can conclude that probably it is confined to the stellar-disk plane. The profile of the stellar velocity dispersion is typical for lenticular galaxies: it has a moderate maximum in the center, $\sigma_{*,0} = 140 \text{ km s}^{-1}$, and

fast drop at $R > 10''$ corresponding to the onset of disk domination in the total surface brightness.

When we exposed the green spectrum of NGC 6534 with the slit at $PA = 16.5^\circ$, the southern nearby galaxy, twice less luminous than NGC 6534 and also with S0 morphology, was serendipitously caught by the slit. Its radial velocity is not given in the NED database, so NGC 6534 is formally classified as an isolated galaxy. From the absorption-line spectrum of the satellite we have derived the systemic velocity of 2MASX J17560734+6415509, $V_r = 8124 \pm 20 \text{ km s}^{-1}$. Since NGC 6534 has $V_r = 8118 \text{ km s}^{-1}$ (NED), 2MASX J17560734+6415509 is a real and very close satellite of NGC 6534. It is slightly bluer than NGC 6534, according to the SDSS/DR9 data; but any signatures of ionized-gas presence are absent from the spectrum. The projected distance between the galaxies is $75''$, or only 39 kpc. If 2MASX J17560734+6415509 (or LEDA 2666218) has anytime been a gas-rich satellite, it might be a gas donor for NGC 6534.

5. Gas excitation and oxygen abundance in the rings

By inspecting the slit spectra of the galaxies, we have identified radius ranges where the rings are betrayed by strong Balmer emissions; the spectra are shown in Fig. 4. We calculated the ratios of the strong emission lines to check the gas excitation. The excitation can be checked with $\log([\text{NII}]\lambda 6583/H\alpha)$. In both cuts of the ring in NGC 6534 and at the southern side of MCG 11-22-015 this ratio are -0.40 ± 0.03 , -0.37 ± 0.02 , and -0.39 ± 0.20 , correspondingly. In the ring of NGC 6534, for which we have also green spectrum, $\log([\text{OIII}]\lambda 5007/H\beta)$ is -0.12 ± 0.15 and 0.00 ± 0.09 . According to Kewley et al. (2006), such emission-line ratios signify gas excitation by young stars in the southern parts of both rings and possible composite excitation mechanism for the northern half of the NGC 6534 ring. At the northern side of the MCG 11-22-015 where the emission lines are weak $\log([\text{NII}]\lambda 6583/H\alpha) = -0.12 \pm 0.12$ so here some shocks may be present.

When the gas is excited by young stars, one can estimate its oxygen abundance by using empirical calibrations of the strong-line ratios. For the ring regions where the gas is shown to be excited by young stars we have probed two recognized calibrations – that by Pettini & Pagel (2004), through N2 for NGC 6534 and MCG 11-22-015, and that by Dopita et al. (2016), D16 with also the $[\text{SII}]\lambda 6717, 6731$ lines, for NGC 6534. We have obtained $12 + \log(\text{O}/\text{H}) = 8.67 \pm 0.02$ from N2 and $12 + \log(\text{O}/\text{H}) = 8.65 \pm 0.10$ from D16 for the southern tip of the ring in NGC 6534; for the starforming site of the southern part of the ring in MCG 11-22-015 the indicator N2 has given $12 + \log(\text{O}/\text{H}) = 8.68 \pm 0.11$. For the northern tip of the ring in NGC 6534 where the excitation may be composite the Dopita et al. (2016) method gives 8.44 ± 0.06 . However, in any case the metallicity of the starforming gas is comparable with the solar value or slightly lower.

6. $H\alpha$ emission morphology and star formation rates in the rings

By undertaking narrow-band photometry of the galaxies through the filters centered onto the redshifted $H\alpha$ line and onto the nearby continuum, we have tried to subtract the latter images from the former ones to derive net emission-line maps which are calibrated into energetic units with the images of the spectrophotometric standard observed the same night. While subtracting, the off-emission images were normalized to reduce

Table 2. Log of the observations with the Russian 6m telescope BTA

Date	Galaxy	Config/Grism or Filter	Exposure	PA(slits)	Spectral range	Seeing
Oct 07, 2016	MCG 11-22-015	slit/VPHG1800R	60 min	50°	6100–7100 Å	1'5
Oct 07, 2016	NGC 6534	slit/VPHG1800R	60 min	20°	6100–7100 Å	1'5
June 24, 2017	NGC 6534	slit/VPHG2300G	60 min	16.5°	4800–5500 Å	3'3
May 28, 2017	NGC 6534	directimage/FN674	40 min	124°	6703–6763 Å	1'5
May 29, 2017	NGC 6534	directimage/SED608	17.5 min	124°	5976–6144 Å	1'8
May 28, 2017	MCG 11-22-015	directimage/FN674	30 min	124°	6703–6763 Å	1'5
May 29, 2017	MCG 11-22-015	directimage/SED608	15.5 min	124°	5976–6144 Å	1'8
May 28, 2017	AGK+81d266	directimage/FN674	1 min	–	6703–6763 Å	1'5




Fig. 1. SDSS-photometry of NGC 6534: the results of the isophote analysis (*upper row*), the azimuthally averaged surface brightness profiles (*bottom left*), and the colour map (*bottom right*). The fitted exponential laws are also shown. The red dotted verticals mark the borders of the $H\alpha$ ring.

to zero the surrounding foreground stars implied to be pure-continuum sources. The resulting net $H\alpha$ maps are shown in Fig. 5. The morphology of the emission-line image of NGC 6534 is somewhat unexpected: while in the broad bands the galaxy has an evident ring, quite detached from the main body of the galaxy, in the $H\alpha$ emission we see patchy spiral arms starting from the ansae of the inner ring (Fig. 5, left). The $H\alpha$ morphology of MCG 11-22-015 is more traditional: it is a round, strongly inhomogeneous ring, with its southern half more bright in the emission line; it borders the inner lens, also filled by weak $H\alpha$ emission.

To calculate star formation rates (SFR) by using three independent indicators, namely, ultraviolet, NUV and FUV fluxes from the GALEX images as well as our estimates of the fluxes in the $H\alpha$ emission line, we have overposed the same ellipti-

cal apertures onto all 3 images of every galaxy. To determine the borders of the apertures, we have used the $H\alpha$ images because they are the ones with the most contrast. The apertures are shown in Fig. 6. We have measured the fluxes inside the elliptical annuli, just between the inner and outer ellipses of Fig. 6 for the three above-mentioned bands for NGC 6534 and only for the deep NUV/MIS image and the $H\alpha$ image of MCG 11-22-015 – the latter galaxy is weak in the UV, and its shallow FUV/AIS image does not contain any signal in the ring above 3σ of the sky level. Then we have applied the calibrations of the SFR from Kennicutt & Evans (2012) and have corrected the UV-based estimates of SFR for the intrinsic dust as it is described in Kostiuk & Sil'chenko (2018), by using the WISE imaging in $22\mu\text{m}$.




Fig. 2. SDSS-photometry of MCG 11-22-015: the results of the isophote analysis (*upper row*), the azimuthally averaged surface brightness profiles (*bottom left*), and the colour map (*bottom right*). The fitted exponential law valid over two-scalelength extension is also shown. The red dotted verticals mark the borders of the $H\alpha$ ring.




Fig. 3. The line-of-sight velocity profiles for the stars and ionized gas in NGC 6534 taken along its major axis.

We found that over the NGC 6534 ring the star formation proceeded with a roughly constant rate during the last 100–200 Myr: the SFRs from the NUV, FUV (GALEX/AIS), and $H\alpha$ fluxes are 0.13 , 0.12 , and $0.18 M_{\odot} \text{ yr}^{-1}$, respectively, with an individual accuracy of the estimates of $0.03 M_{\odot} \text{ yr}^{-1}$.

In MCG 11-22-015 the situation is more complex. The NUV flux measured in the deep MIS-image gives $\text{SFR} = 0.07 \pm 0.01 M_{\odot} \text{ yr}^{-1}$, while the $H\alpha$ flux gives $\text{SFR} = 0.26 \pm 0.01 M_{\odot} \text{ yr}^{-1}$

with the standard calibration. In this situation we remember the caveat from Calzetti (2013) that the standard UV-SFR calibration is valid only for star formation proceeding over long timescales, ~ 100 Myr. If the SF timescale is shorter, we must increase the SFR estimate; for example, for SF occurring during only the last 2 Myr we must multiply the standard calibration by a factor of 3.45 (Calzetti 2013). By multiplying our UV-based SFR of $0.07 M_{\odot} \text{ yr}^{-1}$ by 3.45, we are obtaining $0.24 M_{\odot} \text{ yr}^{-1}$. The latter estimate leads to consistency with the SFR from the $H\alpha$. So for MCG 11-22-015 we may suggest that its star formation in the ring is quite recent – no older than a few Myr.

7. Conclusions

We have studied the outer starforming rings in two lenticular galaxies, NGC 6534 and MCG 11-22-015: spectral data as well as narrow-band and UV images are analyzed. The ionized gas in the rings is mostly excited by young stars and demonstrates nearly solar metallicity rejecting a possibility of its origin from cosmological filaments. Despite the classification of NGC 6534 as an isolated galaxy in the NED, we have detected a very close companion, PGC 2666218, at some 40 kpc from it in projection, with the similar radial velocity. MCG 11-22-015 has already been known as a member of an X-ray bright group (Henry et al. 1995). So there is no problem with the outer gas sources feeding star formation – it may be tidal harassment of the neighbour by NGC 6534 and a minor merger for MCG 11-22-015. Star formation histories in the rings are also different: the SFR has




Fig. 4. The continuum-subtracted spectra near the $H\alpha$ emission line for NGC 6534 (*left*) and for MCG 11-22-015 (*right*).




Fig. 5. The comparison of the broad-band SDSS images (left plot in every pair) and the narrow-band $H\alpha$ images (right plot in every pair). NGC 6534 is to the left and MCG 11-22-015 is to the right.




Fig. 6. The elliptical-annulus apertures used to integrate the fluxes for SFR derivations are overposed onto the $H\alpha$ and the NUV images (left and right plots in every pair). The size of the fields shown is $50''$; NGC 6534 is to the left and MCG 11-22-015 is to the right.

been constant at least for the last 200 Myr in NGC 6534, in accordance with the possible resonance nature of its ring, and it started recently in MCG 11-22-015 betraying a fresh gas-rich satellite cannibalism. The quasi-solar, or only slightly subsolar metallicity of the gas in the outer starforming rings of S0 galaxies was found by us previously more than once, including the cases of obvious minor mergers – see, for example, NGC 4513 with its counterrotating gas (Ilyina & Sil'chenko 2011). Perhaps, we need some specific effective chemical evolution mechanisms to explain this phenomenon.

Acknowledgements. The study of the galactic rings was supported by the Russian Foundation for Basic Researches, grant no. 18-02-00094a. The work is based on the data obtained at the Russian 6m telescope of the Special Astrophysical Observatory and on the public data of the SDSS (<http://www.sdss3.org>) and GALEX (<http://galex.stsci.edu/GR6/>) surveys. We are grateful to D. Oparin, A. Moiseev, and O. Egorov having supported our observations at the Russian 6m telescope (operating under the financial support of the Ministry of Science and Education of the Russian Federation, agreement No14.619.21.0004, the project RFMEF161914X0004). This research has made use of the NASA/IPAC Extragalactic Database (NED, <http://ned.ipac.caltech.edu>) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration, and of the Lyon Extragalactic Database (HyperLEDA, <http://leda.univ-lyon1.fr>).

References

- Afanasiev, V. L., Moiseev, A. V., 2005, *Astronomy Letters*, 31, 194
 Baldwin, J. A., Phillips, M. M., Terlevich, R. 1981, *PASP*, 93, 5
 Calzetti, D., 2013, in *Secular Evolution of Galaxies*, ed. J. Falc3n-Barroso & J. H. Knapen (Cambridge Univ. Press, Cambridge, UK) p. 419
 Comer3n, S., Salo, H., Laurikainen, E., et al. 2014, *A&A*, 562, A121
 de Vaucouleurs, G. 1959, *Handbuch der Physik*, 53, 275
 Dopita, M. A., Kewley, L. J., Sutherland, R. S., Nicholls, D. C., 2016, *Astrophys. Space Sci.*, 361, 61
 Freeman, K. C. 1970, *ApJ*, 160, 767
 Hao, C.-N., Kennicutt, R. C., Johnson, B. D., et al., 2011, *ApJ*, 741, A124
 Henry, J. P., Gioia, I. M., Huchra, J. P., et al. 1995, *ApJ*, 449, 422
 Ilyina, M.A., Sil'chenko, O.K., 2011, *Astronomy Letters*, 37, 589
 Kennicutt Jr., R. C., Evans II, N. J. 2012, *Ann.Rev. A&A*, 50, 531
 Kewley, L. J., Groves, B., Kauffmann, G., Heckman, T., 2006, *MNRAS*, 372, 961
 Kostyuk, I. P. 1975, *Soobscheniya Spets. Astrofiz. Obs.*, 13, 45 [in Russian].
 Kostyuk, I. P., Karachentsev, I. D., Kopylov, A. I., 1981, *Soviet Astronomy Lett.*, 7, 148
 Kostyuk, I. P., Sil'chenko, O. K. 2015, *Astrophys. Bull.*, 70, 280
 Kostyuk, I., Sil'chenko, O. 2018, *Astrophysics*, 61, 1
 Laurikainen, E., Salo, H., Buta, R., Knapen, J.H., 2011, *MNRAS*, 418, 1452
 Pettini M., Pagel B.E.J., 2004, *MNRAS*, 348, L59
 Vorontsov-Velyaminov, B. A., 1960, *Soviet Astronomy*, 4, 365