On a Cahn-Hilliard-Brinkman model for tumour growth and its singular limits

Matthias Ebenbeck^{*} Harald Garcke[†]

Abstract

In this work, we study a model consisting of a Cahn-Hilliard-type equation for the concentration of tumour cells coupled to a reaction-diffusion type equation for the nutrient density and a Brinkmantype equation for the velocity. We equip the system with Neumann boundary for the tumour cell variable and the chemical potential, Robin-type boundary conditions for the nutrient and a "nofriction" boundary condition for the velocity, which allows us to consider solution dependent source terms. Well-posedness of the model as well as existence of strong solutions will be established for a broad class of potentials. We will show that in the singular limit of vanishing viscosities we recover a Darcy-type system related to Cahn-Hilliard-Darcy type models for tumour growth which have been studied earlier. An asymptotic limit will show that the results are also valid in the case of Dirichlet boundary conditions for the nutrient.

Key words: tumour growth, Cahn-Hilliard equation, Brinkman's law, chemotaxis, Stokes flow, Darcy flow, outflow conditions

AMS-Classification: 35K35, 35Q92, 92C50, 35D30, 76D07

1 Introduction

The growth of living cancer cells is affected by many biological and chemical mechanisms. Although the amount of experimental data coming from clinical experiments is quite extensive, the understanding of involved mechanisms and biological effects is still at an unsatisfying level. In the recent past, several mathematical models for tumour growth have been developed and simulated and some of them seem to compare well with tumour data coming from clinical experiments, see [2, 5, 18]. These models may provide further insights into tumour growth dynamics to understand key mechanisms and to design new treatment strategies.

Many models are based on the hypothesis that different tissue components of the tumour (viable, quiescent, necrotic) are separated by a sharp interfacial layer and therefore can be described by free boundary problems, see [8, 17, 19, 20]. As a young tumour does not have its own vascular system and must therefore consume growth factors like nutrients or oxygen from the surrounding host tissue, in the early stage of growth the tumour may undergo morphological instabilities like fingering or folding (see [12, 13]) to grow without angiogenesis and to overcome diffusional limitations. This leads to highly challenging mathematical problems when modelling the tumour in the context of free boundary problems since changes in topology have to be tracked.

To overcome this difficulties, it has turned out that diffuse interface models, treating the tumour as a collection of cells, are a good strategy to describe the evolution and interactions of different species. These models are typically based on a multiphase approach, on balance laws for the single constituents, like

^{*}Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, Germany, e-mail: matthias.ebenbeck@ur.de

[†]Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, Germany, e-mail: harald.garcke@mathematik.uniregensburg.de

mass and momentum balance, on constitutive laws and on thermodynamic principles. Several additional variables describing the extracellular matrix (ECM), growth factors or inhibitors, can be incorporated into these models, biological mechanisms like chemotaxis, apoptosis or necrosis and effects of stress, plasticity or viscoelasticity can be included, see [4, 27, 34, 36]. Many contributions in the literature consider a mixture of two components (healthy and surrounding tissue), modelled as a two-phase flow and coupled to a reaction-diffusion type equation for an unknown species acting as a nutrient for the tumour, like oxygen or glucose, leading to coupled systems of partial differential equations of Cahn-Hilliard-type, see [10, 21, 25, 26, 30, 31]. Some of them incorporate effects based on the momentum balance equation which leads to models involving e.g. Darcy's law for the velocity or a Stokes like equation, see [16, 23, 24, 32].

In the following, we will consider a Cahn-Hilliard-Brinkman system for tumour growth. For a bounded domain $\Omega \subset \mathbb{R}^d$, d = 2, 3, and a fixed time T > 0, we consider for $Q := \Omega \times (0, T)$ the following system of equations

$$\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v}}(\varphi, \sigma) \qquad \qquad \text{in } Q, \tag{1.1a}$$

$$-\operatorname{div}(\mathbf{T}(\mathbf{v}, p)) + \nu \mathbf{v} = (\mu + \chi \sigma) \nabla \varphi \qquad \text{in } Q, \qquad (1.1b)$$

$$\partial_t \varphi + \operatorname{div}(\varphi \mathbf{v}) = \operatorname{div}(m(\varphi) \nabla \mu) + \Gamma_{\varphi}(\varphi, \sigma) \quad \text{in } Q,$$
(1.1c)

$$\mu = \epsilon^{-1} \psi'(\varphi) - \epsilon \Delta \varphi - \chi \sigma \qquad \text{in } Q, \tag{1.1d}$$

$$0 = \Delta \sigma - h(\varphi)\sigma \qquad \qquad \text{in } Q, \qquad (1.1e)$$

where the viscous stress tensor is defined by

$$\mathbf{T}(\mathbf{v}, p) \coloneqq 2\eta(\varphi)\mathbf{D}\mathbf{v} + \lambda(\varphi)\mathrm{div}(\mathbf{v})\mathbf{I} - p\mathbf{I},$$
(1.2)

and the symmetric velocity gradient is given by

$$\mathbf{D}\mathbf{v} \coloneqq \frac{1}{2}(\nabla \mathbf{v} + \nabla \mathbf{v}^T)$$

In (1.1)-(1.2), v denotes the volume-averaged velocity of the mixture, p denotes the pressure, σ denotes the concentration of an unknown species acting as a nutrient, $\varphi \in [-1,1]$ denotes the difference in volume fractions, with $\{\varphi = 1\}$ representing the unmixed tumour tissue and $\{\varphi = -1\}$ representing the surrounding healthy tissue, and μ denotes the chemical potential for φ . The function $m(\cdot)$ represents the mobility for the phase variable φ . The constant $\epsilon > 0$ is related to the thickness of the diffuse interface, whereas ν is a positive constant representing the fluid permeability. Moreover, the functions $\eta(\cdot)$ and $\lambda(\cdot)$ are non-negative and represent the shear and the bulk viscosity, respectively. The chemotaxis parameter χ is a non-negative constant.

By **n** we will denote the outer unit normal on $\partial \Omega$, and $\partial_{\mathbf{n}}g \coloneqq \nabla g \cdot \mathbf{n}$ is the directional derivative. We equip the system with the following initial and boundary conditions

$$\partial_{\mathbf{n}}\mu = \partial_{\mathbf{n}}\varphi = 0 \qquad \text{on } \partial\Omega \times (0,T) \eqqcolon \Sigma, \tag{1.3a}$$

$$\partial_{\mathbf{n}}\sigma = K(\sigma_{\infty} - \sigma) \quad \text{on } \Sigma, \tag{1.3b}$$
$$\mathbf{T}(\mathbf{y}, p)\mathbf{n} = \mathbf{0} \qquad \text{on } \Sigma, \tag{1.3c}$$

$$\mathbf{T}(\mathbf{v}, p)\mathbf{n} = \mathbf{0} \qquad \text{on } \Sigma, \tag{1.3c}$$

$$\varphi(0) = \varphi_0 \qquad \text{in } \Omega, \tag{1.3d}$$

where φ_0, σ_∞ are given functions and K is a positive permeability constant. We now motivate and compare our model with other models in the literature.

• In [16], the authors of this paper considered a similar model with (1.1b), (1.1e) and (1.3b) replaced by

$$-\operatorname{div}(\mathbf{T}(\mathbf{v},p)) + \nu \mathbf{v} = \mu \nabla \varphi + (\chi_{\sigma} + \chi(1-\varphi)) \nabla \sigma \qquad \text{in } Q, \qquad (1.4a)$$

$$\partial_t \sigma + \operatorname{div}(\sigma \mathbf{v}) = \operatorname{div}(n(\varphi)(\chi_\sigma \nabla \sigma - \chi \nabla \varphi)) - \Gamma_\sigma(\varphi, \sigma, \mu) \quad \text{in } Q,$$
(1.4b)

$$n(\varphi)\chi_{\sigma}\partial_{\mathbf{n}}\sigma = K(\sigma_{\infty} - \sigma) \qquad \text{on } \Sigma, \qquad (1.4c)$$

where χ_{σ} is a given positive constant denoting the nutrient diffusion coefficient, $n(\varphi)$ is a mobility function and Γ_{σ} is a source or sink term similar to ours. Reformulating the pressure as

$$\tilde{p} \coloneqq p - \frac{\chi_{\sigma}}{2} |\sigma|^2 + \chi \sigma (1 - \varphi),$$

we can indeed recover (1.1b) from (1.4a). For a discussion regarding pressure reformulations, we refer to [23]. Following the arguments in [28, Sec. 3.3.3], we introduce an active transport parameter κ , a new mobility function $\mathcal{D}(\varphi)$ and set

$$n(\varphi) = \kappa D(\varphi)\chi^{-1}, \quad \chi_{\sigma} = \kappa^{-1}\chi,$$

which allows us to "decouple" active transport and chemotaxis mechanisms. Neglecting mechanism due to active transport, we can rewrite (1.4b)-(1.4c) as

$$\partial_t \sigma + \operatorname{div}(\sigma \mathbf{v}) = \operatorname{div}(\mathcal{D}(\varphi) \nabla \sigma) - \Gamma_\sigma(\varphi, \sigma, \mu) \quad \text{in } Q, \tag{1.5a}$$

$$\mathcal{D}(\varphi)\partial_{\mathbf{n}}\sigma = K(\sigma_{\infty} - \sigma) \qquad \text{on } \Sigma. \tag{1.5b}$$

Performing a non-dimensionalisation and using that the ratio between nutrient diffusion time-scale (minutes or hours) and cell-proliferation time-scale (days or weeks) is quite small, we can drop the time derivative and the convection term in (1.5a) to obtain

$$0 = \Delta \sigma - \Gamma_{\sigma}(\varphi, \sigma, \mu) \quad \text{in } Q, \tag{1.6a}$$

$$\partial_{\mathbf{n}}\sigma = K(\sigma_{\infty} - \sigma) \qquad \text{on } \Sigma.$$
 (1.6b)

We omit the details and refer to [25, 35, 41].

• Brinkman's law was first proposed in [7] and has been derived rigorously by several authors using a homogenization argument for the Stokes equation, see [3, 37]. It can be interpreted as an interpolation between Stokes flow and Darcy's law since the former one is approximated on small length scales whereas the latter one on large length scales, see [15]. Setting $\sigma = 0$, neglecting source terms, i.e. $\Gamma_{\varphi} = \Gamma_{\mathbf{v}} = 0$, and imposing Dirichlet boundary conditions for the velocity, equations (1.1), (1.3) have been studied in [6] for smooth potentials and in [11] for logarithmic potentials. However, if source terms are present, the Dirichlet boundary condition leads to a compatibility condition since

$$\int_{\Omega} \Gamma_{\mathbf{v}} \, \mathrm{d}x = \int_{\Omega} \operatorname{div}(\mathbf{v}) \, \mathrm{d}x = \int_{\partial \Omega} \mathbf{v} \cdot \mathbf{n} \mathrm{d}\mathcal{H}^{d-1} = 0,$$

which can in general not be fulfilled if $\Gamma_{\mathbf{v}}$ depends on additional variables like φ and σ . The "no-friction" boundary condition does not lead to such an condition and therefore enables us to consider solution dependent source terms.

• In the case $\mathbf{v} = 0$ and with (1.1e) replaced by (1.4b), the model (1.1), (1.3) was analysed in [25] and well-posedness has been shown in the case of Dirichlet boundary conditions for φ, μ, σ in [26], where they could rigorously prove the quasi-static limit. In [21], well-posedness has been established for $\partial_{\mathbf{n}}\varphi = \partial_{\mathbf{n}}\mu = \partial_{\mathbf{n}}\sigma = 0$ on Σ , for a large class of potentials and with source terms of the form

$$\Gamma_{\varphi}(\varphi,\mu,\sigma) = P(\varphi)(\sigma-\mu), \quad \Gamma_{\sigma}(\varphi,\mu,\sigma) = P(\varphi)(\sigma-\mu),$$

where $P(\cdot)$ is a nonlinear proliferation function.

• When setting $\eta(\cdot) = \lambda(\cdot) \equiv 0$, the model reduces to the so called Cahn-Hilliard-Darcy model. In the absence of nutrients, this model has been studied in [32] for $\partial_{\mathbf{n}}\varphi = \partial_{\mathbf{n}}\mu = \mathbf{v} \cdot \mathbf{n} = 0$ on Σ and with prescribed source terms $\Gamma_{\varphi} = \Gamma_{\mathbf{v}} = \mathcal{S}$. Setting $\sigma = 0$, prescribing $\partial_{\mathbf{n}}\varphi = \partial_{\mathbf{n}}\mu = \mathbf{v} = 0$ on Σ and setting

 $\Gamma_{\mathbf{v}} = \Gamma_{\varphi} = 0$, solutions of the Cahn-Hilliard-Darcy system have been established upon considering the zero viscosity limit in the Brinkman equation, see [6]. Furthermore, we want to refer to [14], where they considered a multispecies Cahn-Hilliard-Darcy model for tumour growth with quasi-static nutrient equation.

In the following, we want to outline some challenges arising in the analysis. When testing the Brinkman equation with \mathbf{v} , we have to estimate the term $\int_{\Omega} p \operatorname{div}(\mathbf{v}) \, dx$. Hence, we need to get an estimate on $\|p\|_{L^2}$ in the absence of any a-priori-estimates. To overcome this difficulty, we will use the so-called method of "Subtracting the divergence". More precisely, we choose $\mathbf{v} - \mathbf{u}$ as a test-function in the momentum equation, where \mathbf{u} satisfies

div
$$(\mathbf{u}) = \Gamma_{\mathbf{v}}$$
 in Ω , $\mathbf{u} = \frac{1}{|\partial \Omega|} \left(\int_{\Omega} \Gamma_{\mathbf{v}} \right) \mathbf{n}$ on $\partial \Omega$.

Although this prevents us from estimating the pressure, we now have to handle the term

$$\int_{\Omega} \mu \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x = \int_{\Omega} (\mu - \mu_{\Omega}) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x + \mu_{\Omega} \int_{\Omega} \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x$$
$$= \int_{\Omega} (\mu - \mu_{\Omega}) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x + \mu_{\Omega} \left(\frac{1}{|\partial \Omega|} \int_{\Omega} \Gamma_{\mathbf{v}} \, \mathrm{d}x \int_{\partial \Omega} \varphi \mathrm{d}\mathcal{H}^{d-1} - \int_{\Omega} \varphi \Gamma_{\mathbf{v}} \, \mathrm{d}x \right),$$

where $\mu_{\Omega} = \frac{1}{|\Omega|} \int_{\Omega} \mu \, dx$. To handle the boundary integral, we will derive an estimate for the $L^{\rho}(\partial\Omega)$ -norm for φ , where $\rho \in [2, 6]$ is an exponent connected to the growth rate of the potential $\psi(\cdot)$. Furthermore, we comment on the assumption $\sigma_{\infty} \in L^4(L^2(\partial\Omega))$, which is not needed to prove existence of weak solutions, but crucial to establish well-posedness of the system. Indeed, this enables us to estimate the velocity in $L^{\frac{8}{3}}(0,T;\mathbf{H}^1)$ (see Section 3.3). As a consequence, we can handle the term

$$\int_{\Omega} 2(\eta(\varphi_1) - \eta(\varphi_2)) \mathbf{D} \mathbf{v}_2 \colon \nabla \mathbf{v} \, \mathrm{d} x$$

in the proof for continuous dependence (see Section 5). We remark that this term does not arise in the case of constant viscosity.

Finally, in the proof for existence of strong solutions we will derive an estimate for the time derivative of the nutrient concentration by using a difference quotient method. This is needed due to the fact that the L^2 -orthogonal projection \mathbb{P}_n onto the *n*-dimensional Galerkin solution spaces is not continuous on the whole space H^2 . Indeed, when testing (1.1d) with $\Delta \partial_t \varphi$ in the Galerkin scheme and integrating by parts twice, we encounter the term

$$\int_0^T \int_\Omega \chi \Delta(\mathbb{P}_n \sigma) \partial_t \varphi \mathrm{d}x \mathrm{d}t.$$

Although we can control $\sigma \in L^2(H^2)$, an estimate of $\Delta \mathbb{P}_n \sigma \in L^2(L^2)$ can not be deduced due to (1.3b). If the time derivative of σ fulfils $\partial_t \sigma \in L^2(H^1)$, a control of $\Delta \mathbb{P}_n \sigma \in L^2(L^2)$ is not needed, see Section 6.

1.1 Notation and preliminaries

We first want to fix some notation: For a (real) Banach space X we denote by $\|\cdot\|_X$ its norm, by X^* the dual space and by $\langle \cdot, \cdot \rangle_X$ the duality pairing between X^* and X. For an inner product space X, the inner product is denoted by $(\cdot, \cdot)_X$. We define the scalar product of two matrices by

$$\mathbf{A} \colon \mathbf{B} \coloneqq \sum_{i=1}^{d} a_{jk} b_{jk} \quad \text{for } \mathbf{A}, \mathbf{B} \in \mathbb{R}^{d \times d}.$$

For the standard Lebesgue and Sobolev spaces with $1 \leq p \leq \infty$, k > 0, we use the notation $L^p \coloneqq L^p(\Omega)$ and $W^{k,p} \coloneqq W^{k,p}(\Omega)$ with norms $\|\cdot\|_{L^p}$ and $\|\cdot\|_{W^{k,p}}$ respectively. In the case p = 2 we use $H^k \coloneqq W^{k,2}$ and the norm $\|\cdot\|_{H^k}$. For $\beta \in (0,1)$ and $r \in (1,\infty)$, we will denote the Lebesgue and Sobolev spaces on the boundary by $L^p(\partial\Omega)$ and $W^{\beta,r}(\partial\Omega)$ with corresponding norms $\|\cdot\|_{L^p(\partial\Omega)}$ and $\|\cdot\|_{W^{\beta,r}(\partial\Omega)}$ (see [40, Chap. I.3] for more details). By \mathbf{L}^p , $\mathbf{W}^{k,p}$, \mathbf{H}^k , $\mathbf{L}^p(\partial\Omega)$ and $\mathbf{W}^{\beta,r}(\partial\Omega)$, we will denote the corresponding spaces of vector valued and matrix valued functions. We denote the space H_0^1 as the completion of $C_0^{\infty}(\Omega)$ with respect to the H^1 -norm. For the Bochner spaces, we use the notation $L^p(X) \coloneqq L^p(0,T;X)$ for a Banach space X with $p \in [1,\infty]$. For the dual space X^{*} of a Banach space X, we introduce the (generalised) mean value by

$$v_{\Omega} \coloneqq \frac{1}{|\Omega|} \int_{\Omega} v \, \mathrm{d}x \quad \text{for } v \in L^1, \quad v_{\Omega}^* \coloneqq \frac{1}{|\Omega|} \langle v, 1 \rangle_X \quad \text{for } v \in X^*$$

Moreover, we introduce the function spaces

$$\begin{split} L_0^2 &\coloneqq \{ w \in L^2 \colon w_\Omega = 0 \}, \quad H_N^2 \coloneqq \{ w \in H^2 \colon \partial_{\mathbf{n}} w = 0 \text{ on } \partial\Omega \}, \\ (H^1)_0^* &\coloneqq \{ f \in (H^1)^* \colon f_\Omega^* = 0 \}. \end{split}$$

Then, the Neumann-Laplace operator $-\Delta_N \colon H^1 \cap L_0^2 \to (H^1)_0^*$ is positive definite and self-adjoint. In particular, by the Lax-Milgram theorem and Poincaré's inequality, the inverse operator $(-\Delta_N)^{-1} \colon (H^1)_0^* \to H^1 \cap L_0^2$ is well-defined, and we set $u \coloneqq (-\Delta_N)^{-1} f$ for $f \in (H^1)_0^*$ if $u_\Omega = 0$ and

$$-\Delta u = f \text{ in } \Omega, \quad \partial_{\mathbf{n}} u = 0 \text{ on } \partial \Omega.$$

We have dense and continuous embeddings $H_N^2 \subset H^1 \subset L^2 \simeq (L^2)^* \subset (H^1)^* \subset (H_N^2)^*$ and the identifications $\langle u, v \rangle_{H^1} = (u, v)_{L^2}$, $\langle u, w \rangle_{H^2} = (u, w)_{L^2}$ for all $u \in L^2$, $v \in H^1$ and $w \in H_N^2$. We also want to state the following inequalities resulting from elliptic regularity theory and integration by parts:

$$\|\Delta f\|_{L^{2}} \leq \|\nabla f\|_{\mathbf{L}^{2}}^{\frac{1}{2}} \|\nabla \Delta f\|_{\mathbf{L}^{2}}^{\frac{1}{2}} \leq C \|\varphi\|_{H^{1}}^{\frac{1}{2}} \|\varphi\|_{H^{3}}^{\frac{1}{2}} \qquad \forall f \in H^{2}_{N} \cap H^{3},$$
(1.7a)

$$\|f\|_{H^2} \le C \left(\|f\|_{L^2} + \|\Delta f\|_{L^2}\right) \qquad \forall f \in H_N^2, \tag{1.7b}$$

with a constant C depending only on Ω . Furthermore, we define

$$L^2_{\operatorname{div}}(\Omega) \coloneqq \left\{ \mathbf{f} \in \mathbf{L}^2 \colon \operatorname{div}(\mathbf{f}) \in L^2 \right\}$$

equipped with the norm

$$\|\mathbf{f}\|_{L^2_{\operatorname{div}}(\Omega)} \coloneqq \left(\|\mathbf{f}\|_{\mathbf{L}^2}^2 + \|\operatorname{div}(\mathbf{f})\|_{L^2}^2\right)^{\frac{1}{2}},$$

which is a reflexive Banach space (see [38]). In particular, for $\mathbf{u} \in L^2_{\text{div}}(\Omega)$ we have

$$\langle \mathbf{u} \cdot \mathbf{n}, \Phi \rangle_{H^{\frac{1}{2}}(\partial \Omega)} = \int_{\Omega} \mathbf{u} \cdot \nabla \Phi \mathrm{d}x + \int_{\Omega} \Phi \mathrm{div}(\mathbf{u}) \mathrm{d}x \quad \forall \Phi \in H^1$$
 (1.8)

and

$$\|\mathbf{u}\cdot\mathbf{n}\|_{\left(H^{\frac{1}{2}}(\partial\Omega)\right)^{*}} \leq C_{\mathrm{div}}\|\mathbf{u}\|_{L^{2}_{\mathrm{div}}(\Omega)},\tag{1.9}$$

with a constant C_{div} depending only on Ω (see e.g. [22, Chap. III.2]). Furthermore, we will use the following generalised Gagliardo-Nirenberg inequality:

Lemma 1.1. Let $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$, be a bounded domain with Lipschitz-boundary and $f \in W^{m,r} \cap L^q$, $1 \leq q, r \leq \infty$. For any integer j, $0 \leq j < m$, suppose there is $\alpha \in \mathbb{R}$ such that

$$j - \frac{d}{p} = \left(m - \frac{d}{r}\right)\alpha + (1 - \alpha)\left(-\frac{d}{q}\right), \quad \frac{j}{m} \le \alpha \le 1.$$

Then, there exists a positive constant C depending only on Ω, d, m, j, q, r , and α such that

$$\|D^{j}f\|_{L^{p}} \leq C\|f\|_{W^{m,r}}^{\alpha}\|f\|_{L^{q}}^{1-\alpha}.$$
(1.10)

We will also need the following theorem concerning solvability of the divergence equation:

Lemma 1.2. ([22, Sec. III.3]) Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be a bounded domain with Lipschitz-boundary and let $1 < q < \infty$. Then, for every $f \in L^q$ and $\mathbf{a} \in \mathbf{W}^{1-1/q,q}(\partial \Omega)$ satisfying

$$\int_{\Omega} f \, \mathrm{d}x = \int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} \mathrm{d}\mathcal{H}^{d-1},\tag{1.11}$$

there exists at least one solution $\mathbf{u} \in \mathbf{W}^{1,q}$ of the problem

$$div(\mathbf{u}) = f \quad in \ \Omega,$$
$$\mathbf{u} = \mathbf{a} \quad on \ \partial\Omega.$$

In addition, the following estimate holds

$$\|\mathbf{u}\|_{\mathbf{W}^{1,q}} \le C\left(\|f\|_{\mathbf{L}^{q}} + \|\mathbf{a}\|_{\mathbf{W}^{1-1/q,q}(\partial\Omega)}\right),\tag{1.12}$$

with C depending only on Ω and q.

Finally, in the Galerkin ansatz (see Sec. 3) we will make use of the following lemma (see [1] for a proof):

Lemma 1.3. Let $\Omega \subset \mathbb{R}^d$, d = 2, 3, be a bounded domain with C^2 -boundary and outer unit normal \mathbf{n} and $1 < q < \infty$. Furthermore, assume that $g \in W^{1,q}$, $\mathbf{f} \in \mathbf{L}^q$, $c \in W^{1,r}$ with r > d, and the functions $\eta(\cdot)$, $\lambda(\cdot)$ fulfil (A3) (see Assumptions 2.1 below). Then, there exists a unique solution $(\mathbf{v}, p) \in \mathbf{W}^{2,q} \times W^{1,q}$ of the system

$$-\operatorname{div}(2\eta(c)\mathbf{D}\mathbf{v} + \lambda(c)\operatorname{div}(\mathbf{v})\mathbf{I}) + \nu\mathbf{v} + \nabla p = \mathbf{f} \quad a. \ e. \ in \ \Omega,$$
(1.13a)

 $\operatorname{div}(\mathbf{v}) = g \quad a. \ e. \ in \ \Omega, \tag{1.13b}$

$$(2\eta(c)\mathbf{D}\mathbf{v} + \lambda(c)\operatorname{div}(\mathbf{v})\mathbf{I} - p\mathbf{I})\mathbf{n} = \mathbf{0} \quad a. \ e. \ on \ \partial\Omega, \tag{1.13c}$$

satisfying the following estimate

$$\|\mathbf{v}\|_{\mathbf{W}^{2,q}} + \|p\|_{W^{1,q}} \le C\left(\|\mathbf{f}\|_{\mathbf{L}^{q}} + \|g\|_{W^{1,q}}\right),\tag{1.14}$$

with a constant C depending only on $\eta_0, \eta_1, \lambda_0, q, \|c\|_{W^{1,r}}$ and Ω .

2 Main results

We make the following assumptions:

Assumptions 2.1. (A1) The positive constants ϵ , ν , K, T are fixed, χ is a fixed, non-negative constant. Furthermore, the function $\sigma_{\infty} \in L^2(L^2(\partial\Omega))$ and the initial datum $\varphi_0 \in H^1$ are prescribed.

(A2) The mobility $m(\cdot)$ is continuous on \mathbb{R} and satisfies

$$m_0 \leq m(t) \leq m_1 \quad \forall t \in \mathbb{R},$$

for positive constants m_0, m_1 .

(A3) The viscosities fulfil $\eta, \lambda \in C^2(\mathbb{R})$ with bounded first derivatives and

$$\eta_0 \le \eta(t) \le \eta_1, \quad 0 \le \lambda(t) \le \lambda_0 \quad \forall t \in \mathbb{R},$$
(2.1)

for positive constants η_0, η_1 and a non-negative constant λ_0 .

(A4) The source terms are of the form

$$\Gamma_{\mathbf{v}}(\varphi,\sigma) = b_{\mathbf{v}}(\varphi)\sigma + f_{\mathbf{v}}(\varphi), \quad \Gamma_{\varphi}(\varphi,\sigma) = b_{\varphi}(\varphi)\sigma + f_{\varphi}(\varphi),$$

where $b_{\mathbf{v}}, f_{\mathbf{v}} \in C^1(\mathbb{R})$ are bounded with bounded first derivatives and $b_{\varphi}, f_{\varphi} \in C^0(\mathbb{R})$ are bounded functions. The function $h \in C^0(\mathbb{R})$ is continuous, bounded and non-negative.

(A5) The function $\psi \in C^2(\mathbb{R})$ is non-negative and can be written as

$$\psi(s) = \psi_1(s) + \psi_2(s) \quad \forall s \in \mathbb{R},$$
(2.2)

where $\psi_1, \psi_2 \in C^2(\mathbb{R})$ and

$$R_1(1+|s|^{\rho-2}) \le \psi_1''(s) \le R_2(1+|s|^{\rho-2}) \quad \forall s \in \mathbb{R},$$
(2.3)

$$|\psi_2''(s)| \le R_3 \qquad \forall s \in \mathbb{R}, \tag{2.4}$$

where R_i , i = 1, 2, 3, are positive constants with $R_1 < R_2$ and $\rho \in [2, 6]$. Furthermore, if $\rho = 2$, we assume $2R_1 > R_3$.

Remark 2.2. Using (A5), it is straightforward to check that there exist positive constants R_i , i = 4, 5, 6, such that

$$\psi(s) \ge R_4 |s|^{\rho} - R_5 \quad \forall s \in \mathbb{R}$$

$$(2.5)$$

and

$$|\psi'(s)| \le R_6(1+|s|^{\rho-1}) \quad \forall s \in \mathbb{R}.$$
 (2.6)

We now introduce the weak formulation of (1.1), (1.3):

Definition 2.3. (Weak solution for (1.1),(1.3)) We call a quintuple $(\varphi, \sigma, \mu, \mathbf{v}, p)$ a weak solution of (1.1) and (1.3) if

$$\begin{split} \varphi &\in H^1(0,T;(H^1)^*) \cap L^{\infty}(0,T;H^1) \cap L^2(0,T;H^3), \quad \mu \in L^2(0,T;H^1), \\ \sigma &\in L^2(0,T;H^1), \quad \mathbf{v} \in L^2(0,T;\mathbf{H}^1), \quad p \in L^2(0,T;L^2), \end{split}$$

such that

$$\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v}}(\varphi, \sigma) \text{ a.e. in } Q, \quad \varphi(0) = \varphi_0 \text{ a.e. in } \Omega,$$

and

$$0 = \int_{\Omega} \mathbf{T}(\mathbf{v}, p) \colon \nabla \mathbf{\Phi} + \nu \mathbf{v} \cdot \mathbf{\Phi} - (\mu + \chi \sigma) \nabla \varphi \cdot \mathbf{\Phi} \, \mathrm{d}x,$$
(2.7a)

$$0 = \langle \partial_t \varphi, \Phi \rangle_{H^1, (H^1)^*} + \int_{\Omega} m(\varphi) \nabla \mu \cdot \nabla \Phi + (\nabla \varphi \cdot \mathbf{v} + \varphi \Gamma_{\mathbf{v}}(\varphi, \sigma) - \Gamma_{\varphi}(\varphi, \sigma)) \Phi \, \mathrm{d}x, \qquad (2.7b)$$

$$0 = \int_{\Omega} (\mu + \chi \sigma) \Phi - \epsilon^{-1} \Psi'(\varphi) \Phi - \epsilon \nabla \varphi \cdot \nabla \Phi \, \mathrm{d}x, \qquad (2.7c)$$

$$0 = \int_{\Omega} \nabla \sigma \cdot \nabla \Phi + h(\varphi) \sigma \Phi \, \mathrm{d}x + \int_{\partial \Omega} K(\sigma - \sigma_{\infty}) \Phi \mathrm{d}\mathcal{H}^{d-1}, \qquad (2.7\mathrm{d})$$

for a.e. $t \in (0,T)$ and for all $\Phi \in \mathbf{H}^1$, $\Phi \in H^1$.

Theorem 2.4. Let $\Omega \subset \mathbb{R}^d$, d = 2, 3, be a bounded domain with C^3 -boundary and assume that Assumptions 2.1 is fulfilled. Then there exists a solution quintuple $(\varphi, \mu, \sigma, \mathbf{v}, p)$ of (1.1), (1.3) in the sense of Definition 2.3. Furthermore, we have

$$\begin{split} \varphi &\in H^1(0,T;(H^1)^*) \cap L^\infty(0,T;H^1) \cap L^4(0,T;H^2) \cap L^2(0,T;H^3), \quad \sigma \in L^2(0,T;H^1), \\ \mu &\in L^2(0,T;H^1), \quad \mathbf{v} \in L^2(0,T;\mathbf{H}^1), \quad p \in L^2(0,T;L^2), \end{split}$$

and the following estimate holds:

$$\begin{aligned} \|\varphi\|_{H^{1}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{L^{2}(H^{1})} + \|\mu\|_{L^{2}(H^{1})} \\ &+ \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{\frac{3}{2}})} + \|\mathbf{v}\|_{L^{2}(\mathbf{H}^{1})} + \|p\|_{L^{2}(L^{2})} \leq C, \end{aligned}$$
(2.8)

with a constant C depending only on the system parameters, Ω and T. If in addition $\sigma_{\infty} \in L^4(L^2(\partial \Omega))$, we have

$$\sigma \in L^4(0,T;H^1), \quad \mu \in L^4(0,T;L^2), \quad \mathbf{v} \in L^{\frac{8}{3}}(0,T;\mathbf{H}^1),$$

and

$$\|\sigma\|_{L^{4}(H^{1})} + \|\mu\|_{L^{4}(L^{2})} + \|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})} + \|\operatorname{div}(\varphi\mathbf{v})\|_{L^{2}(L^{2})} \le C.$$
(2.9)

Theorem 2.5. (The limit $K \to \infty$) Let the assumptions of Theorem 2.4 be fulfilled and assume in addition that $\sigma_{\infty} \in L^2(H^{\frac{1}{2}}(\partial\Omega))$. Let K > 0 and denote by $(\varphi_K, \mu_K, \sigma_K, \mathbf{v}_K, p_K)$ a weak solution of (1.1), (1.3) corresponding to φ_0 and K in the sense of Definition (2.3). Then, as $K \to \infty$, we have

$$\begin{array}{lll} \varphi_{K} \rightarrow \varphi & weakly \ast & in \ H^{1}((H^{1})^{\ast}) \cap L^{\infty}(H^{1}) \cap L^{4}(H^{2}) \cap L^{2}(H^{3}), \\ \sigma_{K} \rightarrow \sigma & weakly \ast & in \ L^{2}(H^{1}), \\ \mu_{K} \rightarrow \mu & weakly & in \ L^{2}(H^{1}), \\ p_{K} \rightarrow p & weakly & in \ L^{2}(L^{2}), \\ \mathbf{v}_{K} \rightarrow \mathbf{v} & weakly & in \ L^{2}(\mathbf{H}^{1}), \\ \operatorname{div}(\varphi_{K}\mathbf{v}_{K}) \rightarrow \operatorname{div}(\varphi\mathbf{v}) & weakly & in \ L^{2}(L^{3}), \\ \sigma_{K} \rightarrow \sigma_{\infty} & strongly & in \ L^{2}(L^{2}(\partial\Omega)), \end{array}$$

where $(\varphi, \mu, \sigma, \mathbf{v}, p)$ satisfies

$$\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v}}(\varphi, \sigma) \ a. \ e. \ in \ Q, \quad \varphi(0) = \varphi_0 \ a. \ e. \ in \ \Omega, \quad \sigma \in (\sigma_{\infty} + L^2(0, T; H_0^1))$$

and (2.7) with (2.7d) replaced by

$$0 = \int_{\Omega} \nabla \sigma \cdot \nabla \xi + h(\varphi) \sigma \xi \, \mathrm{d}x, \qquad (2.10)$$

for a. e. $t \in (0,T)$ and for all $\xi \in H_0^1$.

We now introduce the definition of weak solutions of the Cahn-Hilliard-Darcy system endowed with (1.3a)-(1.3b) and

$$p = 0 \quad \text{on } \Sigma. \tag{2.11}$$

Definition 2.6. We call a quintuple $(\varphi, \mu, \sigma, \mathbf{v}, p)$ weak solution of the Cahn-Hilliard-Darcy system endowed with (1.3a)-(1.3b) and (2.11) if

$$\begin{split} \varphi \in W^{1,\frac{8}{5}}(0,T;(H^1)^*) \cap L^{\infty}(0,T;H^1) \cap L^2(0,T;H^3), \quad \mu \in L^2(0,T;H^1), \\ \sigma \in L^2(0,T;H^1), \quad \mathbf{v} \in L^2(0,T;\mathbf{L}^2), \quad p \in L^2(0,T;H_0^1), \end{split}$$

such that

$$\varphi(0) = \varphi_0$$
 a.e. in Ω

and

$$0 = \langle \partial_t \varphi, \phi \rangle_{H^1} + \int_{\Omega} m(\varphi) \nabla \mu \cdot \nabla \phi dx + \int_{\Omega} (\nabla \varphi \cdot \mathbf{v} + \varphi \Gamma_{\mathbf{v}}(\varphi, \sigma) - \Gamma_{\varphi}(\varphi, \sigma)) \phi dx, \qquad (2.12a)$$

$$0 = \int_{\Omega} (\mu + \chi \sigma) \phi - \epsilon^{-1} \psi'(\varphi) \phi - \epsilon \nabla \varphi \cdot \nabla \phi \mathrm{d}x, \qquad (2.12b)$$

$$0 = \int_{\Omega} \nabla \sigma \cdot \nabla \phi + h(\varphi) \sigma \phi dx + \int_{\partial \Omega} K(\sigma - \sigma_{\infty}) \phi d\mathcal{H}^{d-1}, \qquad (2.12c)$$

$$0 = \int_{\Omega} \left(\mathbf{v} + \frac{1}{\nu} \nabla p - \frac{1}{\nu} (\mu + \chi \sigma) \nabla \varphi \right) \cdot \mathbf{\Phi} \mathrm{d}x, \qquad (2.12\mathrm{d})$$

$$0 = \int_{\Omega} \frac{1}{\nu} \nabla p \cdot \nabla \xi - \Gamma_{\mathbf{v}}(\varphi, \sigma) \xi - \frac{1}{\nu} (\mu + \chi \sigma) \nabla \varphi \cdot \nabla \xi \mathrm{d}x, \qquad (2.12e)$$

for a.e. $t\in (0,T)$ and all $\phi\in H^1,\,\xi\in H^1_0,\,\Phi\in \mathbf{L}^2.$

The following theorem states that solutions of the Cahn-Hilliard-Darcy system can be found as the limit of the Cahn-Hilliard-Brinkman system when the viscosities tend to zero.

Theorem 2.7. Let $\Omega \subset \mathbb{R}^d$, d = 2, 3, be a bounded domain with C^3 -boundary and assume that (A1)-(A2), (A4)-(A5) holds. Furthermore, let $\{\eta_n, \lambda_n\}_{n \in \mathbb{N}}$ be a sequence of function pairs fulfilling (A3) such that

$$\|\eta_n(\cdot)\|_{C^0(\mathbb{R})} \to 0, \quad \|\lambda_n(\cdot)\|_{C^0(\mathbb{R})} \to 0 \quad as \ n \to \infty.$$

Let $(\varphi_n, \mu_n, \sigma_n, \mathbf{v}_n, p_n)$ be a sequence of weak solutions of the Cahn-Hilliard-Brinkman system in the sense of Definition 2.3 for $\eta(\cdot) = \eta_n(\cdot), \lambda(\cdot) = \lambda_n(\cdot)$ and originating from $\varphi_0 \in H^1$. Then, at least for a subsequence, $(\varphi_n, \mu_n, \sigma_n, \mathbf{v}_n, p_n)$ converges to a weak solution $(\varphi, \mu, \sigma, \mathbf{v}, p)$ of the Cahn-Hilliard-Darcy system in the sense of Definition 2.6 such that

$$\begin{split} \varphi_n &\to \varphi \quad weakly \ast \quad in \ W^{1,\frac{8}{5}}((H^1)^*) \cap L^{\infty}(H^1) \cap L^4(H^2) \cap L^2(H^3), \\ \sigma_n &\to \sigma \quad weakly \quad in \ L^2(H^1), \\ \mu_n &\to \mu \quad weakly \quad in \ L^2(H^1), \\ p_n &\to p \quad weakly \quad in \ L^2(L^2), \\ \mathbf{v}_n &\to \mathbf{v} \quad weakly \quad in \ L^2(\mathbf{L}^2) \cap L^2\left(L^2_{\mathrm{div}}(\Omega)\right), \\ 2\eta_n(\varphi_n) \mathbf{D} \mathbf{v}_n &\to \mathbf{0} \quad weakly \quad in \ L^2(\mathbf{L}^2), \\ \lambda_n(\varphi_n) \mathrm{div}(\mathbf{v}_n) \mathbf{I} \to \mathbf{0} \quad weakly \quad in \ L^2(\mathbf{L}^2), \end{split}$$

and

$$\varphi_n \to \varphi$$
 strongly in $C^0(L^r) \cap L^2(W^{2,r})$ and a.e. in Q ,

for all $r \in [1, 6)$. Moreover, it holds that

$$\begin{split} \partial_{\mathbf{n}}\varphi &= p = 0 & a. e. \ on \ \Sigma, \\ \operatorname{div}(\mathbf{v}) &= \Gamma_{\mathbf{v}}(\varphi, \sigma) & a. e. \ in \ Q, \\ \mathbf{v} &= \frac{1}{\nu} \left(-\nabla p + (\mu + \chi \sigma) \nabla \varphi \right) & a. e. \ in \ Q, \\ \mu &= \epsilon^{-1} \psi'(\varphi) - \epsilon \Delta \varphi - \chi \sigma & a. e. \ in \ Q, \end{split}$$

and

$$\begin{aligned} \|\varphi\|_{W^{1,\frac{8}{5}}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\mu\|_{L^{2}(H^{1})} + \|\sigma\|_{L^{2}(H^{1})} \\ &+ \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{\frac{8}{5}}((H^{1})^{*})} + \|\mathbf{v}\|_{L^{2}(L^{2}_{\operatorname{div}}(\Omega))} + \|p\|_{L^{2}(H^{1}_{0})} \leq C, \end{aligned}$$
(2.13)

with a constant C depending only on the system parameters and on Ω , T.

To prove continuous dependence on the initial and boundary data, we make the following additional assumptions:

Assumptions 2.8. (B1) The mobility $m(\cdot)$ is a constant, without loss of generality we assume $m(\cdot) \equiv 1$.

- (B2) The functions $b_{\varphi}(\cdot), f_{\varphi}(\cdot)$ and $h(\cdot)$, are Lipschitz continuous with Lipschitz constants L_b, L_f and L_h , respectively.
- (B3) For ψ' and ψ'' , we assume that

$$|\psi'(s_1) - \psi'(s_2)| \le k_1(1 + |s_1|^4 + |s_2|^4)|s_1 - s_2| \quad \forall s_1, s_2 \in \mathbb{R},$$
(2.14)

$$|\psi''(s_1) - \psi''(s_2)| \le k_2(1 + |s_1|^3 + |s_2|^3)|s_1 - s_2| \quad \forall s_1, s_2 \in \mathbb{R},$$
(2.15)

for some positive constants k_1, k_2 .

Theorem 2.9. Let $\Omega \subset \mathbb{R}^d$, d = 2, 3, be a bounded domain with C^3 -boundary and assume that Assumptions 2.1 and 2.8 hold. Then, for any two weak solution quintuples $\{\varphi_i, \mu_i, \sigma_i, \mathbf{v}_i, p_i\}$, i = 1, 2, of (1.1), (1.3) satisfying

$$\begin{split} \varphi &\in H^1(0,T;(H^1)^*) \cap L^{\infty}(0,T;H^1) \cap L^4(0,T;H^2) \cap L^2(0,T;H^3), \quad \sigma \in L^4(0,T;H^1), \\ \mu &\in L^4(0,T;L^2) \cap L^2(0,T;H^1), \quad \mathbf{v} \in L^{\frac{8}{3}}(0,T;\mathbf{H}^1), \quad p \in L^2(0,T;L^2), \end{split}$$

with $\sigma_{i,\infty} \in L^4(L^2(\partial\Omega))$ and $\varphi_i(0) = \varphi_{i,0} \in H^1$ for i = 1, 2, there exists a positive constant C depending only on the system parameters and on Ω, T, L_h, L_b, L_f such that

$$\sup_{t \in (0,T]} \left(\left\| \varphi_{1}(t) - \varphi_{2}(t) \right\|_{H^{1}}^{2} \right) + \left\| \varphi_{1} - \varphi_{2} \right\|_{H^{1}((H^{1})^{*}) \cap L^{2}(H^{3})}^{2} + \left\| \mu_{1} - \mu_{2} \right\|_{L^{2}(H^{1})}^{2} + \left\| \sigma_{1} - \sigma_{2} \right\|_{L^{2}(H^{1})}^{2} + \left\| \mathbf{v}_{1} - \mathbf{v}_{2} \right\|_{L^{2}(\mathbf{H}^{1})}^{2} + \left\| p_{1} - p_{2} \right\|_{L^{2}(L^{2})}^{2} \leq C \left(\left\| \varphi_{1,0} - \varphi_{2,0} \right\|_{H^{1}}^{2} + \left\| \sigma_{1,\infty} - \sigma_{2,\infty} \right\|_{L^{4}(L^{2}(\partial\Omega))}^{2} \right).$$

$$(2.16)$$

We have the following notion of strong solutions:

Definition 2.10. (Strong solution for (1.1),(1.3)) We call a quintuple $(\varphi, \sigma, \mu, \mathbf{v}, p)$ a strong solution of (1.1) and (1.3) if

$$\begin{split} \varphi &\in H^1(0,T;L^2) \cap L^\infty(0,T;H^2) \cap L^2(0,T;H^4), \quad \mu \in L^2(0,T;H^2), \\ \sigma &\in L^2(0,T;H^2), \quad \mathbf{v} \in L^2(0,T;\mathbf{H}^2), \quad p \in L^2(0,T;H^1), \end{split}$$

and (1.1), (1.3) are fulfilled almost everywhere in the respective sets.

For the existence of strong solutions, we make the following additional assumptions:

Assumptions 2.11.

(C1) The mobility $m(\cdot)$ is a constant, without loss of generality we assume $m(\cdot) \equiv 1$. The function $h(\cdot)$ is Lipschitz continuous with Lipschitz constant L_h .

(C2) The boundary datum $\sigma_{\infty} \in H^1(0,T; H^{\frac{1}{2}}(\partial\Omega))$ and the initial datum $\varphi_0 \in H^2_N$ are prescribed.

(C3) The function $\psi \in C^3(\mathbb{R})$ fulfils

$$|\psi'''(s)| \le k_3(1+|s|^3) \quad \forall s \in \mathbb{R},$$
(2.17)

for a positive constant k_3 .

We have the following result concerning strong solutions:

Theorem 2.12. Let $\Omega \subset \mathbb{R}^d$, d = 2, 3, be a bounded domain with C^4 -boundary and assume that Assumptions 2.1 and 2.11 hold. Then there exists a solution quintuple $(\varphi, \mu, \sigma, \mathbf{v}, p)$ of (1.1), (1.3) in the sense of Definition 2.10. Furthermore, we have

$$\varphi \in C^0(\bar{Q}), \quad \mu \in L^\infty(0,T;L^2), \quad \sigma \in H^1(0,T;H^1) \cap L^\infty(0,T;H^2), \quad \mathbf{v} \in L^8(0,T;\mathbf{H}^2), \quad p \in L^8(0,T;H^1), \quad \mu \in L^\infty(0,T;H^1), \quad \mu \in L^$$

and the following estimate holds:

$$\begin{aligned} \|\varphi\|_{H^{1}(L^{2})\cap C^{0}(\bar{Q})\cap L^{\infty}(H^{2})\cap L^{2}(H^{4})} + \|\sigma\|_{H^{1}(H^{1})\cap L^{\infty}(H^{2})} + \|\mu\|_{L^{\infty}(L^{2})\cap L^{2}(H^{2})} \\ + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{2})} + \|\mathbf{v}\|_{L^{8}(\mathbf{H}^{2})} + \|p\|_{L^{8}(H^{1})} \leq C. \end{aligned}$$

$$(2.18)$$

3 Existence of weak solutions

In order to prove the result, we now derive a priori estimates for (2.7a)-(2.7d). By C, we denote a positive constant not depending on $(\varphi, \mu, \sigma, \mathbf{v}, p)$ which may vary from line to line. The duality pairing in (2.7b) can be replaced by the L^2 -product for smooth enough functions which is satisfied for example by a Galerkin-ansatz. Approximating solutions can be constructed by applying a Galerkin approximation with respect to φ and μ and at the same time solving for \mathbf{v} , p and σ in the corresponding whole function spaces (for details, we refer to [16], [23]). In the following, we will write Γ_{φ} , $\Gamma_{\mathbf{v}}$, instead of $\Gamma_{\varphi}(\varphi, \sigma)$, $\Gamma_{\mathbf{v}}(\varphi, \sigma)$.

3.1 A-priori-estimates

3.1.1 Estimating the nutrient concentration

Choosing $\Phi = \sigma$ in (2.7d) and using the non-negativity of $h(\cdot)$, we obtain

$$\int_{\Omega} |\nabla \sigma|^2 \, \mathrm{d}x + K \int_{\partial \Omega} |\sigma|^2 \mathrm{d}\mathcal{H}^{d-1} \le K \int_{\partial \Omega} \sigma \sigma_{\infty} \mathrm{d}\mathcal{H}^{d-1}.$$
(3.1)

Using Hölder's and Young's inequalities, we have

$$\left| K \int_{\partial \Omega} \sigma \sigma_{\infty} \mathrm{d} \mathcal{H}^{d-1} \right| \leq \frac{K}{2} \| \sigma \|_{L^{2}(\partial \Omega)}^{2} + \frac{K}{2} \| \sigma_{\infty} \|_{L^{2}(\partial \Omega)}^{2}.$$
(3.2)

Using (3.1)-(3.2) and Poincaré's inequality, we deduce that

$$\|\sigma\|_{H^1} \le C \|\sigma_\infty\|_{L^2(\partial\Omega)}.$$
(3.3)

Moreover, by the continuous embedding $H^1 \hookrightarrow L^p$, $p \in [2, 6]$, and (A4), we have

$$\|\Gamma_{\varphi}\|_{L^{p}} + \|\Gamma_{\mathbf{v}}\|_{L^{p}} \le C\left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}\right) \quad \forall p \in [2, 6].$$

$$(3.4)$$

3.1.2 An energy identity

Due to (A4), (3.3)-(3.4), there exists a solution $\mathbf{u} \in \mathbf{H}^1$ of the problem

$$\operatorname{div}(\mathbf{u}) = \Gamma_{\mathbf{v}} \quad \text{in } \Omega, \qquad \mathbf{u} = \frac{1}{|\partial\Omega|} \left(\int_{\Omega} \Gamma_{\mathbf{v}} \, \mathrm{d}x \right) \mathbf{n} \eqqcolon \mathbf{a} \quad \text{on } \partial\Omega, \tag{3.5}$$

satisfying the estimate

$$\|\mathbf{u}\|_{\mathbf{H}^{1}} \leq C \|\Gamma_{\mathbf{v}}\|_{L^{2}} \leq C \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}\right).$$

$$(3.6)$$

Choosing $\Phi = \mathbf{v} - \mathbf{u}$ in (2.7a), $\Phi = \mu + \chi \sigma$ in (2.7b), $\Phi = \partial_t \varphi$ in (2.7c), and summing the resulting identities, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} \int_{\Omega} \epsilon^{-1} \psi(\varphi) + \frac{\epsilon}{2} |\nabla\varphi|^2 \,\mathrm{d}x + \int_{\Omega} m(\varphi) |\nabla\mu|^2 + \int_{\Omega} 2\eta(\varphi) |\mathbf{D}\mathbf{v}|^2 + \nu |\mathbf{v}|^2 \,\mathrm{d}x$$

$$= \int_{\Omega} -m(\varphi) \chi \nabla \mu \cdot \nabla \sigma + (\Gamma_{\varphi} - \varphi \Gamma_{\mathbf{v}}) (\mu + \chi \sigma) \,\mathrm{d}x$$

$$+ \int_{\Omega} 2\eta(\varphi) \mathbf{D}\mathbf{v} \colon \nabla \mathbf{u} + \nu \mathbf{v} \cdot \mathbf{u} \,\mathrm{d}x - \int_{\Omega} (\mu + \chi \sigma) \nabla \varphi \cdot \mathbf{u} \,\mathrm{d}x.$$
(3.7)

3.1.3 Estimating the right hand side of the energy identity

Using Hölder's and Young's inequalities together with (3.6), we have

$$\left| \int_{\Omega} 2\eta(\varphi) \mathbf{D} \mathbf{v} \colon \nabla \mathbf{u} + \nu \mathbf{v} \cdot \mathbf{u} \, \mathrm{d} x \right| \leq \left\| \sqrt{\eta(\varphi)} \mathbf{D} \mathbf{v} \right\|_{\mathbf{L}^{2}}^{2} + \frac{\nu}{2} \left\| \mathbf{v} \right\|_{\mathbf{L}^{2}}^{2} + C \left(1 + \left\| \eta(\varphi) \right\|_{L^{\infty}} \right) \left\| \Gamma_{\mathbf{v}} \right\|_{L^{2}}^{2}.$$
(3.8)

We now want to estimate the terms involving $\Gamma_{\mathbf{v}}$ and Γ_{φ} . Using Hölder's, Poincaré's and Young's inequalities together with (3.3)-(3.4), we obtain

$$\left| \int_{\Omega} \Gamma_{\varphi}(\mu + \chi \sigma) \, \mathrm{d}x \right| \leq C_P \|\Gamma_{\varphi}\|_{L^2} \left(\left| (\mu + \chi \sigma)_{\Omega} \right| + \|\nabla(\mu + \chi \sigma)\|_{\mathbf{L}^2} \right) \\ \leq C \left(1 + \|\sigma_{\infty}\|_{L^2(\partial\Omega)}^2 \right) \left(1 + \left| (\mu + \chi \sigma)_{\Omega} \right| \right) + \frac{m_0}{8} \|\nabla\mu\|_{\mathbf{L}^2}^2.$$
(3.9)

With similar arguments, using the continuous embedding $H^1 \hookrightarrow L^6$, we obtain

$$\begin{aligned} \left| \int_{\Omega} \Gamma_{\mathbf{v}} \varphi(\mu + \chi \sigma) \, \mathrm{d}x \right| &\leq C \| \Gamma_{\mathbf{v}} \|_{L^{3}} \| \varphi \|_{L^{2}} \| \mu + \chi \sigma \|_{L^{6}} \\ &\leq C \| \Gamma_{\mathbf{v}} \|_{L^{3}} \| \varphi \|_{L^{2}} \left(|(\mu + \chi \sigma)_{\Omega}| + \| \nabla (\mu + \chi \sigma) \|_{\mathbf{L}^{2}} \right) \\ &\leq C \left(1 + \| \sigma_{\infty} \|_{L^{2}(\partial \Omega)}^{2} \right) \left(1 + \| \varphi \|_{L^{2}}^{2} + \| \varphi \|_{L^{2}} |(\mu + \chi \sigma)_{\Omega}| \right) + \frac{m_{0}}{8} \| \nabla \mu \|_{\mathbf{L}^{2}}^{2}. \tag{3.10}$$

Now, choosing $\Phi = 1$ in (2.7c) and using (2.6), we obtain

$$\left| \int_{\Omega} \mu + \chi \sigma \, \mathrm{d}x \right| = \left| \int_{\Omega} \epsilon^{-1} \psi'(\varphi) \, \mathrm{d}x \right| \le \epsilon^{-1} R_6 \int_{\Omega} 1 + |\varphi|^{\rho-1} \, \mathrm{d}x \le C \left(1 + \|\varphi\|_{L^{\rho-1}}^{\rho-1} \right),$$
$$\left| (\mu + \chi \sigma)_{\Omega} \right| \le C \left(1 + \|\varphi\|_{L^{\rho-1}}^{\rho-1} \right) \le C \left(1 + \|\varphi\|_{L^{\rho}}^{\rho-1} \right).$$

hence

implies

$$|(\mu + \chi_0)_{M_1}| \leq C \left(1 + ||\varphi||_{L^{\rho-1}}\right) \leq C \left(1 + ||\varphi||_{L^{\rho}}\right).$$

In particular, using Young's inequality, the continuous embedding $L^{\rho} \hookrightarrow L^2, \rho \in [2, 6]$, and (2.5), this

$$|(\mu + \chi \sigma)_{\Omega}| \le C \left(1 + \|\varphi\|_{L^{\rho}}^{\rho-1} \right) \le C \left(1 + \|\varphi\|_{L^{\rho}}^{\rho} \right) \le C \left(1 + \|\psi(\varphi)\|_{L^{1}} \right), \tag{3.11}$$

$$\|\varphi\|_{L^{2}}|(\mu + \chi\sigma)_{\Omega}| \le C \left(\|\varphi\|_{L^{2}} + \|\varphi\|_{L^{\rho}}^{\rho}\right) \le C \left(1 + \|\varphi\|_{L^{\rho}}^{\rho}\right) \le C \left(1 + \|\psi(\varphi)\|_{L^{1}}\right).$$
(3.12)

Using (3.11)-(3.12) in (3.9)-(3.10) and applying the continuous embedding $L^{\rho} \hookrightarrow L^2$, $\rho \in [2, 6]$, together with (2.5), we end up with

$$\int_{\Omega} (\Gamma_{\varphi} - \varphi \Gamma_{\mathbf{v}})(\mu + \chi \sigma) \, \mathrm{d}x \bigg| \le C \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} \right) \left(1 + \|\psi(\varphi)\|_{L^{1}} \right) + \frac{m_{0}}{4} \|\nabla \mu\|_{\mathbf{L}^{2}}^{2}. \tag{3.13}$$

For the first term on the r.h.s. of (3.7), applying Hölder's and Young's inequalities, (A2) and (3.3), we obtain

$$\int_{\Omega} m(\varphi) \chi \nabla \mu \cdot \nabla \sigma \, \mathrm{d}x \bigg| \le m_1 \chi \|\nabla \mu\|_{\mathbf{L}^2} \|\nabla \sigma\|_{\mathbf{L}^2} \le \frac{m_0}{8} \|\nabla \mu\|_{\mathbf{L}^2}^2 + C \|\sigma_{\infty}\|_{L^2(\partial\Omega)}^2.$$
(3.14)

3.1.4 Estimating the remaining term

The case $\rho = 2$: Using Hölder's, Young's and Poincaré's inequalities, the continuous embeddings $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$, $H^1 \hookrightarrow L^3$ and (2.5), (3.6), (3.11), we obtain

$$\begin{aligned} \left| \int_{\Omega} (\mu + \chi \sigma) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x \right| &\leq \|\mu + \chi \sigma\|_{L^{3}} \|\nabla \varphi\|_{\mathbf{L}^{2}} \|\mathbf{u}\|_{\mathbf{L}^{6}} \\ &\leq C \left(|(\mu + \chi \sigma)_{\Omega}| + \|\nabla (\mu + \chi \sigma)\|_{\mathbf{L}^{2}} \right) \|\nabla \varphi\|_{\mathbf{L}^{2}} \|\mathbf{u}\|_{\mathbf{H}^{1}} \\ &\leq C \left(1 + \|\varphi\|_{L^{2}} + \|\nabla (\mu + \chi \sigma)\|_{\mathbf{L}^{2}} \right) \|\nabla \varphi\|_{\mathbf{L}^{2}} \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)} \right) \\ &\leq C \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} \right) \left(1 + \|\varphi\|_{L^{2}}^{2} + \|\nabla \varphi\|_{\mathbf{L}^{2}}^{2} \right) + \frac{m_{0}}{8} \|\nabla \mu\|_{\mathbf{L}^{2}}^{2} \\ &\leq C \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} \right) \left(1 + \|\psi(\varphi)\|_{L^{1}} + \|\nabla \varphi\|_{\mathbf{L}^{2}}^{2} \right) + \frac{m_{0}}{8} \|\nabla \mu\|_{\mathbf{L}^{2}}^{2}. \end{aligned}$$
(3.15)

The case $\rho \in (2,6]$: In this case, we need a more subtle argument. Choosing $\Phi = -\Delta \varphi$ in (2.7c), integrating by parts and using (2.2), it holds

$$\epsilon \int_{\Omega} |\Delta \varphi|^2 \, \mathrm{d}x + \epsilon^{-1} \int_{\Omega} \psi_1''(\varphi) |\nabla \varphi|^2 \, \mathrm{d}x = -\epsilon^{-1} \int_{\Omega} \psi_2''(\varphi) |\nabla \varphi|^2 \, \mathrm{d}x + \int_{\Omega} \nabla(\mu + \chi \sigma) \cdot \nabla \varphi \, \mathrm{d}x. \tag{3.16}$$

Neglecting the non-negative term $\epsilon \int_{\Omega} |\Delta \varphi|^2 dx$ on the l.h.s. of this equation and using (2.3)-(2.4) together with Young's inequality, we obtain

$$\int_{\Omega} |\varphi|^{\rho-2} |\nabla\varphi|^2 \, \mathrm{d}x \le \left(1 + \frac{R_3}{R_1} + \frac{\epsilon^2}{4\delta R_1^2}\right) \|\nabla\varphi\|_{\mathbf{L}^2}^2 + \delta \|\nabla(\mu + \chi\sigma)\|_{\mathbf{L}^2}^2,\tag{3.17}$$

with $\delta > 0$ to be chosen later. Observing that

$$\left|\nabla\left(\frac{2|\varphi|^{\frac{\rho}{2}}}{\rho}\right)\right| = |\varphi|^{\frac{\rho-2}{2}}|\nabla\varphi|,$$

from (3.17) we deduce that

$$\left\|\nabla(|\varphi|^{\frac{\rho}{2}})\right\|_{\mathbf{L}^{2}}^{2} \leq \frac{\rho^{2}}{4} \left(1 + \frac{R_{3}}{R_{1}} + \frac{\epsilon^{2}}{4\delta R_{1}^{2}}\right) \left\|\nabla\varphi\right\|_{\mathbf{L}^{2}}^{2} + \frac{\delta\rho^{2}}{4} \left\|\nabla(\mu + \chi\sigma)\right\|_{\mathbf{L}^{2}}^{2}.$$
(3.18)

Now, applying the trace theorem and (3.18) yields

$$\begin{split} \left\| |\varphi|^{\frac{\rho}{2}} \right\|_{L^{2}(\partial\Omega)}^{2} &\leq C_{tr}^{2} \left(\left\| |\varphi|^{\frac{\rho}{2}} \right\|_{L^{2}}^{2} + \left\| \nabla \left(|\varphi|^{\frac{\rho}{2}} \right) \right\|_{\mathbf{L}^{2}}^{2} \right) \\ &\leq C_{tr}^{2} \left(\left\| \varphi \right\|_{L^{\rho}}^{\rho} + \frac{\rho^{2}}{4} \left(1 + \frac{R_{3}}{R_{1}} + \frac{\epsilon^{2}}{4\delta R_{1}^{2}} \right) \left\| \nabla \varphi \right\|_{\mathbf{L}^{2}}^{2} + \frac{\delta \rho^{2}}{4} \left\| \nabla (\mu + \chi \sigma) \right\|_{\mathbf{L}^{2}}^{2} \right), \end{split}$$

hence

$$\|\varphi\|_{L^{\rho}(\partial\Omega)}^{\rho} \le C_{tr}^{2} \left(\|\varphi\|_{L^{\rho}}^{\rho} + \frac{\rho^{2}}{4} \left(1 + \frac{R_{3}}{R_{1}} + \frac{\epsilon^{2}}{4\delta R_{1}^{2}} \right) \|\nabla\varphi\|_{\mathbf{L}^{2}}^{2} + \frac{\delta\rho^{2}}{4} \|\nabla(\mu + \chi\sigma)\|_{\mathbf{L}^{2}}^{2} \right).$$
(3.19)

Now, upon integrating by parts and recalling $(3.5)_2$, we calculate

$$\int_{\Omega} (\mu + \chi \sigma) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x = \int_{\Omega} \left(\mu + \chi \sigma - (\mu + \chi \sigma)_{\Omega} \right) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x + (\mu + \chi \sigma)_{\Omega} \int_{\Omega} \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x$$
$$= \int_{\Omega} \left(\mu + \chi \sigma - (\mu + \chi \sigma)_{\Omega} \right) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x$$
$$+ (\mu + \chi \sigma)_{\Omega} \left(\frac{1}{|\partial \Omega|} \int_{\Omega} \Gamma_{\mathbf{v}} \, \mathrm{d}x \int_{\partial \Omega} \varphi \mathrm{d}\mathcal{H}^{d-1} - \int_{\Omega} \varphi \Gamma_{\mathbf{v}} \, \mathrm{d}x \right).$$
(3.20)

Using Hölder's, Young's and Poincaré's inequalities, the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ and (3.6), it is straightforward to check that

$$\int_{\Omega} \left(\mu + \chi \sigma - (\mu + \chi \sigma)_{\Omega} \right) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x \bigg| \le C_{\delta_1} (1 + \|\sigma_{\infty}\|_{L^2(\partial\Omega)}^2) (1 + \|\nabla\varphi\|_{\mathbf{L}^2}^2) + \delta_1 \|\nabla\mu\|_{\mathbf{L}^2}^2, \tag{3.21}$$

with $\delta_1 > 0$ to be chosen. Using (3.6), (3.12) and Hölder's inequality, we obtain

$$\left| (\mu + \chi \sigma)_{\Omega} \int_{\Omega} \varphi \Gamma_{\mathbf{v}} \, \mathrm{d}x \right| \leq \left| (\mu + \chi \sigma)_{\Omega} \right| \|\Gamma_{\mathbf{v}}\|_{L^{2}} \|\varphi\|_{L^{2}} \leq C \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)} \right) \left(1 + \|\psi(\varphi)\|_{L^{1}} \right)$$
(3.22)

Now, using Hölder's and Young's inequalities, (2.5), (3.6), (3.11), (3.19) and recalling $\rho > 2$, we deduce that

$$\left| (\mu + \chi \sigma)_{\Omega} \frac{1}{|\partial \Omega|} \int_{\Omega} \Gamma_{\mathbf{v}} \, \mathrm{d}x \int_{\partial \Omega} \varphi \mathrm{d}\mathcal{H}^{d-1} \right| \leq C \left| (\mu + \chi \sigma)_{\Omega} |\| \Gamma_{\mathbf{v}} \|_{L^{2}} \|\varphi\|_{L^{\rho}(\partial \Omega)}$$

$$\leq C \left(1 + \|\varphi\|_{L^{\rho}}^{\rho-1} \right) \|\Gamma_{\mathbf{v}}\|_{L^{2}} \|\varphi\|_{L^{\rho}(\partial \Omega)}$$

$$\leq C_{\delta_{2}} \left(1 + \|\varphi\|_{L^{\rho}}^{\rho} \right) \|\Gamma_{\mathbf{v}}\|_{L^{2}}^{\frac{\rho}{\rho-1}} + \delta_{2} \|\varphi\|_{L^{\rho}(\partial \Omega)}^{\rho}$$

$$\leq C_{\delta_{2}} \left(1 + \|\sigma_{\infty}\|_{L^{2}(\partial \Omega)}^{2} \right) \left(1 + \|\psi(\varphi)\|_{L^{1}} \right) + \delta_{2} \|\varphi\|_{L^{\rho}(\partial \Omega)}^{\rho} \quad (3.23)$$

where we used that $\frac{\rho-1}{\rho} + \frac{1}{\rho} = 1$. Now, plugging in (3.21)-(3.23) into (3.20), using (2.5), (3.19) and choosing $\delta, \delta_1, \delta_2$ small enough, we finally obtain

$$\left| \int_{\Omega} (\mu + \chi \sigma) \nabla \varphi \cdot \mathbf{u} \, \mathrm{d}x \right| \le \left(1 + \left\| \sigma_{\infty} \right\|_{L^{2}(\partial \Omega)}^{2} \right) \left(1 + \left\| \psi(\varphi) \right\|_{L^{1}} + \left\| \nabla \varphi \right\|_{\mathbf{L}^{2}}^{2} \right) + \frac{m_{0}}{8} \left\| \nabla \mu \right\|_{\mathbf{L}^{2}}^{2}. \tag{3.24}$$

Plugging in (3.8), (3.13)-(3.15), (3.24) into (3.7), using (2.5) and (3.3), we end up with

$$\frac{\mathrm{d}}{\mathrm{d}t} \int_{\Omega} \epsilon^{-1} \psi(\varphi) + \frac{\epsilon}{2} |\nabla \varphi|^2 \,\mathrm{d}x + \frac{m_0}{4} \|\nabla \mu\|_{\mathbf{L}^2}^2 + \frac{\nu}{2} \|\mathbf{v}\|_{\mathbf{L}^2}^2 + \|\sqrt{\eta(\varphi)} \mathbf{D}\mathbf{v}\|_{\mathbf{L}^2}^2 + \|\sigma\|_{H^1}^2$$

$$\leq \alpha(t) \left(1 + \|\nabla \varphi(t)\|_{\mathbf{L}^2}^2 + \|\psi(\varphi(t))\|_{L^1}\right),$$

where, recalling (A1) and (A3),

$$\alpha(t) \coloneqq C \left(1 + \|\eta(\varphi(t))\|_{L^{\infty}} \right) \left(1 + \|\sigma_{\infty}(t)\|_{L^{2}(\partial\Omega)}^{2} \right) \in L^{1}(0,T).$$
(3.25)

Integrating the last inequality in time from 0 to $s \in (0,T]$ and applying Gronwall's Lemma (see [25, Lemma 3.1]) yields

$$\epsilon^{-1} \|\psi(\varphi(s))\|_{L^{1}} + \frac{\epsilon}{2} \|\nabla\varphi(s)\|_{\mathbf{L}^{2}}^{2} + \int_{0}^{s} \frac{m_{0}}{4} \|\nabla\mu\|_{\mathbf{L}^{2}}^{2} + \frac{\nu}{2} \|\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + \|\sqrt{\eta(\varphi)}\mathbf{D}\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + \|\sigma\|_{H^{1}}^{2} dt$$

$$\leq \left(\epsilon^{-1} \|\psi(\varphi_{0})\|_{L^{1}} + \frac{\epsilon}{2} \|\nabla\varphi_{0}\|_{\mathbf{L}^{2}}^{2} + \int_{0}^{s} \alpha(t) dt\right) \exp\left(\int_{0}^{s} \alpha(t) dt\right) \quad \forall s \in (0, T].$$
(3.26)

Due to (A1), (A5) and the continuous embedding $H^1 \hookrightarrow L^6$, we have $\psi(\varphi_0) \in L^1$, $\varphi_0 \in \mathbf{L}^2$. Then, due to Korn's inequality (see [9, Thm. 6.3-3]) and (A3), taking the supremum over all $s \in (0, T]$ in (3.26) implies

$$\operatorname{ess\,sup}_{s\in(0,T]} \left(\|\psi(\varphi(s))\|_{L^1} + \|\nabla\varphi(s)\|_{\mathbf{L}^2}^2 \right) + \int_0^T \|\mathbf{v}\|_{\mathbf{H}^1}^2 + \|\nabla\mu\|_{\mathbf{L}^2}^2 + \|\sigma\|_{H^1}^2 \,\mathrm{d}t \le C.$$
(3.27)

Recalling (2.5), using Poincaré's inequality, (3.11), (3.27) and the fact that $\rho \ge 2$, this in particular gives

$$\underset{s \in (0,T]}{\text{ess sup}} \|\varphi(s)\|_{H^1}^2 + \int_0^T \|\mu\|_{H^1}^2 \, \mathrm{d}t \le C.$$

Combining the last inequality with (3.27) yields

$$\underset{s \in (0,T]}{\mathrm{ess\,sup}} \left(\left\| \psi(\varphi(s)) \right\|_{L^{1}} + \left\| \varphi(s) \right\|_{H^{1}}^{2} \right) + \int_{0}^{T} \left\| \mathbf{v} \right\|_{\mathbf{H}^{1}}^{2} + \left\| \mu \right\|_{H^{1}}^{2} + \left\| \sigma \right\|_{H^{1}}^{2} \, \mathrm{d}t \le C.$$
 (3.28)

Due to (A4) and the continuous embedding $H^1 \hookrightarrow L^6$, this implies

$$\|\Gamma_{\mathbf{v}}\|_{L^{2}(L^{6})} + \|\Gamma_{\varphi}\|_{L^{2}(L^{6})} \le C.$$
(3.29)

3.1.5 Estimating the pressure

Using Lemma 1.2, we deduce that there is at least one solution $\mathbf{q} \in \mathbf{H}^1$ of the system

$$\operatorname{div}(\mathbf{q}) = p \quad \text{in } \Omega, \qquad \mathbf{q} = \frac{1}{|\partial \Omega|} \left(\int_{\Omega} p \, \mathrm{d}x \right) \mathbf{n} \quad \text{on } \partial \Omega,$$

such that

$$\|\mathbf{q}\|_{\mathbf{H}^{1}} \le C_{d} \|p\|_{L^{2}},\tag{3.30}$$

with C_d depending only on Ω and q = 2. Notice that the compatibility condition (1.11) is satisfied since

$$\int_{\partial\Omega} \mathbf{q} \cdot \mathbf{n} \mathrm{d}\mathcal{H}^{d-1} = \frac{1}{|\partial\Omega|} \left(\int_{\Omega} p \, \mathrm{d}x \right) \int_{\partial\Omega} \mathbf{n} \cdot \mathbf{n} \mathrm{d}\mathcal{H}^{d-1} = \int_{\Omega} p \, \mathrm{d}x.$$

Choosing $\mathbf{\Phi} = \mathbf{q}$ in (2.7a) and using div $(\mathbf{v}) = \Gamma_{\mathbf{v}}$ a.e. in Q, we obtain

$$\int_{\Omega} |p|^2 \, \mathrm{d}x = \int_{\Omega} (2\eta(\varphi)\mathbf{D}\mathbf{v} + \lambda(\varphi)\Gamma_{\mathbf{v}}\mathbf{I}) \colon \nabla \mathbf{q} \, \mathrm{d}x + \int_{\Omega} (\nu \mathbf{v} - (\mu + \chi\sigma)\nabla\varphi) \cdot \mathbf{q} \, \mathrm{d}x.$$
(3.31)

Using (3.30), (A3) and Hölder's and Young's inequalities, an easy calculation shows that

$$\|p\|_{L^{2}}^{2} \leq C\left(\|\eta(\varphi)\|_{L^{\infty}}\|\sqrt{\eta(\varphi)}\mathbf{D}\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + \|\lambda(\varphi)\|_{L^{\infty}}^{2}\|\Gamma_{\mathbf{v}}\|_{L^{2}}^{2} + \|\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + \|\mu + \chi\sigma\|_{L^{3}}^{2}\|\nabla\varphi\|_{\mathbf{L}^{2}}^{2}\right).$$
(3.32)

Integrating this inequality in time from 0 to T and using Hölder's inequality, we arrive at

$$\begin{split} \|p\|_{L^{2}(L^{2})}^{2} &\leq C\left(\|\eta(\cdot)\|_{L^{\infty}(\mathbb{R})}\|\sqrt{\eta(\varphi)}\mathbf{D}\mathbf{v}\|_{L^{2}(\mathbf{L}^{2})}^{2} + \|\lambda(\cdot)\|_{L^{\infty}(\mathbb{R})}^{2}\|\Gamma_{\mathbf{v}}\|_{L^{2}(L^{2})}^{2}\right) \\ &+ C\left(\|\mathbf{v}\|_{L^{2}(\mathbf{L}^{2})}^{2} + \|\mu + \chi\sigma\|_{L^{2}(L^{3})}^{2}\|\nabla\varphi\|_{L^{\infty}(\mathbf{L}^{2})}^{2}\right). \end{split}$$

Using (3.26), (3.28)-(3.29) and (A3), the last inequality implies

$$\|p\|_{L^2(L^2)} \le C. \tag{3.33}$$

3.1.6 Higher order estimates for φ

Our aim is to show that

$$\|\varphi\|_{L^2(H^3)} \le C. \tag{3.34}$$

Choosing $\Phi = -\Delta \varphi$ in (2.7c), integrating by parts and neglecting the non-negative term resulting from ψ_1 (see (A5)), we obtain

$$\epsilon \|\Delta \varphi\|_{L^2}^2 + \int_{\Omega} \epsilon^{-1} \psi_2''(\varphi) |\nabla \varphi|^2 \, \mathrm{d}x \le \int_{\Omega} \nabla (\mu + \chi \sigma) \cdot \nabla \varphi \, \mathrm{d}x.$$

Using Hölder's inequality and the assumptions on ψ_2 , we therefore get

$$\epsilon \|\Delta\varphi\|_{L^2}^2 \le \epsilon^{-1} R_3 \|\nabla\varphi\|_{\mathbf{L}^2}^2 + \|\nabla(\mu + \chi\sigma)\|_{\mathbf{L}^2} \|\nabla\varphi\|_{\mathbf{L}^2}.$$

Taking the square of this inequality and integrating in time from 0 to T, we obtain

$$\epsilon^{2} \int_{0}^{T} \left\| \Delta \varphi \right\|_{L^{2}}^{4} \leq C \int_{0}^{T} \left\| \nabla \varphi \right\|_{\mathbf{L}^{2}}^{4} + \left\| \nabla \varphi \right\|_{\mathbf{L}^{2}}^{2} \left\| \nabla (\mu + \chi \sigma) \right\|_{\mathbf{L}^{2}}^{2} dt$$
$$\leq C \left(\left\| \nabla \varphi \right\|_{L^{\infty}(\mathbf{L}^{2})}^{4} + \left\| \nabla (\mu + \chi \sigma) \right\|_{L^{2}(\mathbf{L}^{2})}^{2} \left\| \nabla \varphi \right\|_{L^{\infty}(\mathbf{L}^{2})}^{2} \right).$$

Applying elliptic regularity theory and (3.28), this gives

$$\|\varphi\|_{L^4(H^2)} \le C. \tag{3.35}$$

Next, we take $\Phi = \Delta^2 \varphi$ in (2.7c), integrate by parts and in time from 0 to T to obtain

$$\epsilon \|\nabla\Delta\varphi\|_{L^2(\mathbf{L}^2)}^2 = -\int_0^T \int_{\Omega} \nabla(\mu + \chi\sigma) \cdot \nabla\Delta\varphi \, \mathrm{d}x \, \mathrm{d}t + \int_0^T \int_{\Omega} \epsilon^{-1} \psi''(\varphi) \nabla\varphi \cdot \nabla\Delta\varphi \, \mathrm{d}x \, \mathrm{d}t.$$
(3.36)

For the first term on the r.h.s. of (3.36), applying Hölder's and Young's inequality gives

$$\left| \int_0^T \int_\Omega \nabla(\mu + \chi \sigma) \cdot \nabla \Delta \varphi \, \mathrm{d}x \, \mathrm{d}t \right| \le C \left(\|\mu\|_{L^2(H^1)}^2 + \|\sigma\|_{L^2(H^1)}^2 \right) + \frac{\epsilon}{4} \|\nabla \Delta \varphi\|_{L^2(\mathbf{L}^2)}^2. \tag{3.37}$$

Due to (1.10), (2.3)-(2.4) and (3.28), using Hölder's and Young's inequalities yields

$$\left| \int_{0}^{T} \int_{\Omega} \epsilon^{-1} \psi''(\varphi) \nabla \varphi \cdot \nabla \Delta \varphi \, \mathrm{d}x \, \mathrm{d}t \right| \leq C \int_{0}^{T} \int_{\Omega} \left(1 + |\varphi||^{\rho-2} \right) |\nabla \varphi| |\nabla \Delta \varphi| \mathrm{d}x \mathrm{d}t$$
$$\leq C \int_{\Omega} \left(1 + \|\varphi\|_{L^{\infty}}^{\rho-2} \right) \|\nabla \varphi\|_{\mathbf{L}^{2}} \|\nabla \Delta \varphi\|_{\mathbf{L}^{2}} \mathrm{d}t$$
$$\leq C \int_{0}^{T} \left(1 + \|\varphi\|_{H^{2}}^{\rho-2} \right) \|\nabla \Delta \varphi\|_{\mathbf{L}^{2}} \mathrm{d}t$$
$$\leq C \left(1 + \|\varphi\|_{L^{\rho-2}(H^{2})}^{\rho-2} \right) + \frac{\epsilon}{4} \|\nabla \Delta \varphi\|_{L^{2}(\mathbf{L}^{2})}^{2}. \tag{3.38}$$

Recalling that $\rho - 2 \leq 4$, collecting (3.36)-(3.38) and using (3.28), (3.35), we see that

$$\frac{\epsilon}{2} \left\| \nabla \Delta \varphi \right\|_{L^2(\mathbf{L}^2)}^2 \le C$$

Together with (3.35) and using again elliptic regularity theory, this implies

$$\|\varphi\|_{L^2(H^3)} \le C. \tag{3.39}$$

3.1.7 Regularity for the convection terms and the time derivatives

By Hölder's and Young's inequalities and the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$, we observe that

$$\|\nabla \varphi \cdot \mathbf{v}\|_{L^{2}(0,T;L^{\frac{3}{2}})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{2}}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{\infty}(\mathbf{L}^{2})}^{2} \|\mathbf{v}\|_{L^{2}(0,T;\mathbf{H}^{1})}^{2}$$

Using (3.28)-(3.29) we see that

$$\|\varphi\Gamma_{\mathbf{v}}\|_{L^{2}(L^{2})}^{2} \leq C \|\varphi\|_{L^{\infty}(H^{1})}^{2} \|\Gamma_{\mathbf{v}}\|_{L^{2}(L^{3})}^{2} \leq C.$$

From the last two inequalities, we deduce that

$$\|\operatorname{div}(\varphi \mathbf{v})\|_{L^2(0,T;L^{\frac{3}{2}})} \le C.$$
 (3.40)

Finally, using (3.28)-(3.29) and (3.40), a comparison argument in (2.7b) shows that

$$\|\partial_t \varphi\|_{L^2((H^1)^*)}.\tag{3.41}$$

Notice that we have lower time regularity for the time derivative of φ compared to the convection term since the regularity of the time derivative depends on the term $\nabla \mu$. Summarising all the estimates, we end up with

$$\begin{aligned} \|\varphi\|_{H^{1}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{L^{2}(H^{1})} + \|\mu\|_{L^{2}(H^{1})} \\ &+ \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{\frac{3}{2}})} + \|\mathbf{v}\|_{L^{2}(\mathbf{H}^{1})} + \|p\|_{L^{2}(L^{2})} \leq C. \end{aligned}$$
(3.42)

3.2 Passing to the limit

The a-priori-estimates (3.42) deduced within the Galerkin scheme are enough to deduce existence of solutions. We refer the reader to [16, 23] for details in passing to the limit in the Galerkin scheme.

3.3 Further results on regularity

In the case when $\sigma_{\infty} \in L^4(L^2(\partial\Omega))$, by (3.3) we obtain

$$\|\sigma\|_{L^4(H^1)} \le C. \tag{3.43}$$

In particular, by (A4) this gives

$$\|\Gamma_{\mathbf{v}}\|_{L^4(L^2)} + \|\Gamma_{\varphi}\|_{L^4(L^2)} \le C.$$
(3.44)

Thanks to Lemma 1.1, we have the continuous embedding

$$L^{\infty}(H^1) \cap L^2(H^3) \hookrightarrow L^{20}(L^{10}).$$

Hence, the assumptions on $\psi(\cdot)$ and (3.42) imply

$$\|\psi'(\varphi)\|_{L^4(L^2)} \le C. \tag{3.45}$$

Taking $\Phi = \mu + \chi \sigma$ in (2.7c) and squaring the resulting identity, an application of Hölder's and Young's inequalities gives

 $\|\mu + \chi\sigma\|_{L^{2}}^{4} \leq C\left(\|\psi'(\varphi)\|_{L^{2}}^{4} + \|\nabla(\mu + \chi\sigma)\|_{\mathbf{L}^{2}}^{2}\|\nabla\varphi\|_{\mathbf{L}^{2}}^{2}\right).$

Integrating this inequality in time from 0 to T and using (3.42), (3.45), we conclude that

$$\|\mu + \chi \sigma\|_{L^4(L^2)} \le C. \tag{3.46}$$

We now choose $\Phi = \mathbf{v}$ in (2.7a), use Young's, Hölder's and Korn's inequality (see [9, Thm. 6.3-3]) together with the Sobolev embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ to obtain

$$\|\mathbf{v}\|_{\mathbf{H}^{1}}^{\frac{8}{3}} \leq C\left(\|p\|_{L^{2}}^{\frac{4}{3}}\|\Gamma_{\mathbf{v}}\|_{L^{2}}^{\frac{4}{3}} + \|(\mu + \chi\sigma)\nabla\varphi\|_{\mathbf{L}^{\frac{8}{5}}}^{\frac{8}{3}}\right).$$

Integrating this inequality in time from 0 to T, by Hölder's and Young's inequalities we get

$$\|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})}^{\frac{8}{3}} \leq C\left(\|p\|_{L^{2}(L^{2})}^{\frac{4}{3}}\|\Gamma_{\mathbf{v}}\|_{L^{4}(L^{2})}^{\frac{4}{3}} + \|\mu + \chi\sigma\|_{L^{4}(L^{2})}^{\frac{8}{3}}\|\nabla\varphi\|_{L^{8}(\mathbf{L}^{3})}^{\frac{8}{3}}\right).$$
(3.47)

Applying Lemma 1.1, we have the continuous embedding

$$L^{\infty}(H^1) \cap L^4(H^2) \hookrightarrow L^8(W^{1,3}).$$
(3.48)

Hence, using (3.42), (3.45) and (3.46), from (3.47) we conclude that

$$\|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^1)} \le C. \tag{3.49}$$

Furthermore, using Hölder's and Young's inequalities, the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ and (3.48)-(3.49) yields

$$\|\nabla \varphi \cdot \mathbf{v}\|_{L^{2}(0,T;L^{2})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{3}}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{8}(\mathbf{L}^{3})}^{2} \|\mathbf{v}\|_{L^{\frac{8}{3}}(0,T;\mathbf{H}^{1})}^{2} \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{2}(0,T;L^{2})}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{2}(0,T;L^{2})}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{2}(0,T;L^{2})}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \leq C \int_{0}^{T} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2} \|\nabla \varphi\|_{\mathbf{L}^{2}(0,T;L^{2})}^{2} \, \mathrm{d}t \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \|\nabla \varphi\|_{L^{2}(0,T;L^{2})}^{2} \leq C \|\nabla \varphi\|_{L^{2}(0,T;L^{2$$

Together with the estimate (see (3.29), (3.42))

$$\|\varphi\Gamma_{\mathbf{v}}\|_{L^{2}(L^{2})}^{2} \leq C \|\varphi\|_{L^{\infty}(H^{1})}^{2} \|\Gamma_{\mathbf{v}}\|_{L^{2}(L^{3})}^{2} \leq C,$$

this implies

$$\|\operatorname{div}(\varphi \mathbf{v})\|_{L^2(L^2)} \le C. \tag{3.50}$$

Using (3.43), (3.46), (3.49)-(3.50) and recalling (3.42), we obtain

$$\begin{aligned} \|\varphi\|_{H^{1}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{L^{4}(H^{1})} + \|\mu\|_{L^{2}(H^{1})\cap L^{4}(L^{2})} \\ + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{2})} + \|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})} + \|p\|_{L^{2}(L^{2})} \leq C. \end{aligned}$$

$$(3.51)$$

3.4 The singular limit of large boundary permeability

Due to Theorem 2.4, for every K > 0 there exists a solution quintuple $(\varphi_K, \mu_K, \sigma_K, \mathbf{v}_K, p_K)$ solving (1.1), (1.3) in the sense of Definition 2.3 and enjoying the regularity properties stated in Theorem 2.4. In the following, we assume without loss of generality that K > 1. Let $E: H^{\frac{1}{2}}(\partial\Omega) \to H^1$ be a bounded, linear extension operator satisfying $(Ef)|_{\partial\Omega} = f$ for all $f \in H^{\frac{1}{2}}(\partial\Omega)$ (see [33, Chap. 2, Thm. 5.7] for details). Then, choosing $\Phi = \sigma_K - E\sigma_\infty$ in (2.7d) (in the following, we will omit the operator E), we obtain

$$\int_{\Omega} |\nabla \sigma_K|^2 + h(\varphi) |\sigma_K|^2 \, \mathrm{d}x + K \int_{\partial \Omega} |\sigma_K - \sigma_\infty|^2 \mathrm{d}\mathcal{H}^{d-1} = \int_{\Omega} \nabla \sigma_K \cdot \nabla \sigma_\infty + h(\varphi) \sigma_K \sigma_\infty \, \mathrm{d}x.$$
(3.52)

For the first term on the r.h.s. of this equation, we use Hölder's and Young's inequalities and the boundedness of the extension operator to obtain

$$\left| \int_{\Omega} \nabla \sigma_K \cdot \nabla \sigma_\infty \, \mathrm{d}x \right| \le \frac{1}{4} \| \nabla \sigma_K \|_{\mathbf{L}^2}^2 + \| \nabla \sigma_\infty \|_{\mathbf{L}^2}^2 \le \frac{1}{4} \| \nabla \sigma_K \|_{\mathbf{L}^2}^2 + C \| \sigma_\infty \|_{H^{\frac{1}{2}}(\partial\Omega)}^2. \tag{3.53}$$

With the same arguments and using the boundedness of $h(\cdot)$, we can estimate the second term on the r.h.s. of (3.52) by

$$\left| \int_{\Omega} h(\varphi) \sigma_K \sigma_{\infty} \, \mathrm{d}x \right| \le \delta \|\sigma_K\|_{L^2}^2 + C_{\delta} \|\sigma_{\infty}\|_{H^{\frac{1}{2}}(\partial\Omega)}^2.$$
(3.54)

From Poincaré's inequality and the boundedness of the extension operator, we know that

$$\|\sigma_{K}\|_{L^{2}}^{2} \leq \tilde{C}\left(\|\nabla\sigma_{K}\|_{\mathbf{L}^{2}}^{2} + \|\sigma_{K} - \sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} + \|\sigma_{\infty}\|_{H^{\frac{1}{2}}(\partial\Omega)}^{2}\right),$$
(3.55)

for a positive constant \tilde{C} independent of K. Choosing δ small enough and using this inequality in (3.54), we obtain

$$\left| \int_{\Omega} h(\varphi) \sigma_K \sigma_{\infty} \, \mathrm{d}x \right| \le \frac{1}{4} \left(\left\| \nabla \sigma_K \right\|_{\mathbf{L}^2}^2 + \left\| \sigma_K - \sigma_{\infty} \right\|_{L^2(\partial\Omega)}^2 \right) + C \left\| \sigma_{\infty} \right\|_{H^{\frac{1}{2}}(\partial\Omega)}^2. \tag{3.56}$$

Plugging in (3.53), (3.56) into (3.52) and neglecting the non-negative term $\int_{\Omega} h(\varphi) |\sigma_K|^2 dx$ on the l.h.s. of (3.52), we arrive at

$$\int_{\Omega} |\nabla \sigma_K|^2 \, \mathrm{d}x + K \int_{\partial \Omega} |\sigma_K - \sigma_\infty|^2 \mathrm{d}\mathcal{H}^{d-1} \le C \|\sigma_\infty\|_{H^{\frac{1}{2}}(\partial\Omega)}^2.$$
(3.57)

Multiplying (3.55) by $\frac{1}{2\tilde{C}}$ and adding the resulting equation to (3.57) yields

$$\|\sigma_K\|_{H^1}^2 + K \int_{\partial\Omega} |\sigma_K - \sigma_\infty|^2 \mathrm{d}\mathcal{H}^{d-1} \le C \|\sigma_\infty\|_{H^{\frac{1}{2}}(\partial\Omega)}^2$$

Integrating this inequality in time from 0 to T and using $\sigma_{\infty} \in L^2(H^{\frac{1}{2}}(\partial\Omega))$, we conclude that

$$\|\sigma_K\|_{L^2(H^1)} + \sqrt{K} \|\sigma_K - \sigma_\infty\|_{L^2(L^2(\partial\Omega))} \le C,$$

where C is independent of K. Then, with exactly the same arguments as above, it follows that

$$\begin{aligned} \|\varphi_K\|_{H^1((H^1)^*)\cap L^{\infty}(H^1)\cap L^4(H^2)\cap L^2(H^3)} + \|\sigma_K\|_{L^2(H^1)} + \sqrt{K} \|\sigma_K - \sigma_\infty\|_{L^2(L^2(\partial\Omega))} \\ + \|\mu_K\|_{L^2(H^1)} + \|\operatorname{div}(\varphi_K \mathbf{v}_K)\|_{L^2(L^{\frac{3}{2}})} + \|\mathbf{v}_K\|_{L^2(\mathbf{H}^1)} + \|p_K\|_{L^2(L^2)} \le C. \end{aligned}$$
(3.58)

Using standard compactness arguments (Aubin-Lions theorem (see [39, Sec. 8, Cor. 4]) and reflexive weak compactness), we obtain exactly the convergence results as stated in Theorem 2.5. Passing to the limit can be carried out with exactly the same arguments as stated in Section 3.2. We will only present the arguments needed for (2.7d). In the following, let $\xi \in H_0^1$ be arbitrary. Multiplying (2.7d) with $\delta \in C_0^{\infty}(0,T)$, integrating in time from 0 to T and noting that $H_0^1 \subset H^1$, we observe that

$$0 = \int_0^T \int_\Omega \delta(\nabla \sigma_K \cdot \nabla \xi + h(\varphi_K) \sigma_K \xi) \, \mathrm{d}x \, \mathrm{d}t \quad \forall \xi \in H_0^1.$$
(3.59)

Since $h(\cdot)$ is a bounded, continuous function, $\delta \xi \in C^{\infty}(H_0^1)$ and $\varphi_K \to \varphi$ a.e. in Q, the Lebesgue theorem gives that

$$\|h(\varphi_K)\delta\xi - h(\varphi)\delta\xi\|_{L^2(Q)} \to 0 \text{ as } K \to \infty.$$

Since $\sigma_K \to \sigma$ weakly in $L^2(Q)$ as $K \to \infty$, by the product of weak-strong convergence we obtain

$$\int_0^T \int_\Omega h(\varphi_K) \sigma_K \xi \, \mathrm{d}x \, \mathrm{d}t \to \int_0^T \int_\Omega h(\varphi) \sigma \xi \, \mathrm{d}x \, \mathrm{d}t \quad \text{as } K \to \infty.$$
(3.60)

Furthermore, since $\sigma_K \to \sigma$ weakly in $L^2(H^1)$ and as $\delta \xi \in L^2(H^1)$, it follows that

$$\int_{0}^{T} \int_{\Omega} \delta \nabla \sigma_{K} \cdot \nabla \xi \, \mathrm{d}x \, \mathrm{d}t \to \int_{0}^{T} \int_{\Omega} \delta \nabla \sigma \cdot \nabla \xi \, \mathrm{d}x \, \mathrm{d}t \quad \text{as } K \to \infty.$$
(3.61)

Due to (3.60)-(3.61), we can pass to the limit in (3.59) to deduce that

$$0 = \int_0^T \int_\Omega \delta(\nabla \sigma \cdot \nabla \xi + h(\varphi)\sigma \xi) \, \mathrm{d}x \, \mathrm{d}t \quad \forall \xi \in H_0^1$$

Since this holds for all $\delta \in C_0^{\infty}(0,T)$, we can recover (2.10). Finally, from (3.58), we know that

$$\|\sigma_K - \sigma_\infty\|_{L^2(L^2(\partial\Omega))} \le \frac{C}{\sqrt{K}},$$

where C is independent of K. Letting $K \to \infty$ and recalling that $\sigma_K \to \sigma$ weakly in $L^2(L^2(\partial \Omega))$ as $K \to \infty$, it follows that

$$\sigma = \sigma_{\infty}$$
 a.e. on Σ_{\pm}

which completes the proof.

4 The singular limit of vanishing viscosities

Let $\{\eta_n, \lambda_n\}_{n \in \mathbb{N}}$ be a sequence of function pairs fulfilling (A3) such that

$$\|\eta_n(\cdot)\|_{C^0(\mathbb{R})} \to 0, \quad \|\lambda_n(\cdot)\|_{C^0(\mathbb{R})} \to 0 \quad \text{as } n \to \infty.$$

$$(4.1)$$

Without loss of generality, we can assume that

$$\|\eta_n(\cdot)\|_{L^{\infty}(\mathbb{R})} \le 1, \quad \|\lambda_n(\cdot)\|_{L^{\infty}(\mathbb{R})} \le 1.$$

$$(4.2)$$

Then, by Theorem 2.4, for every $n \in \mathbb{N}$ there exists a solution quintuple $(\varphi_n, \mu_n, \sigma_n, \mathbf{v}_n, p_n)$ of (1.1), (1.3) in the sense of Definition 2.3 fulfilling

$$\begin{split} \varphi_n &\in H^1(0,T;(H^1)^*) \cap L^{\infty}(0,T;H^1) \cap L^4(0,T;H^2) \cap L^2(0,T;H^3), \quad \mu_n \in L^2(0,T;H^1), \\ \sigma_n &\in L^2(0,T;H^1), \quad \mathbf{v}_n \in L^2(0,T;\mathbf{H}^1), \quad p_n \in L^2(0,T;L^2), \end{split}$$

such that

$$\operatorname{div}(\mathbf{v}_n) = \Gamma_{\mathbf{v}}(\varphi_n, \sigma_n) \text{ a.e. in } Q, \quad \partial_{\mathbf{n}}\varphi_n = 0 \text{ a.e. in } \Sigma, \quad \varphi_n(0) = \varphi_0 \text{ a.e. in } \Omega, \tag{4.3}$$

and

$$0 = \int_{\Omega} \mathbf{T}_n(\mathbf{v}_n, p_n) \colon \nabla \mathbf{\Phi} + \nu \mathbf{v}_n \cdot \mathbf{\Phi} - (\mu_n + \chi \sigma_n) \nabla \varphi_n \cdot \mathbf{\Phi} dx, \qquad (4.4a)$$

$$0 = \langle \partial_t \varphi_n, \Phi \rangle_{H^1, (H^1)^*} + \int_{\Omega} m(\varphi_n) \nabla \mu_n \cdot \nabla \Phi + (\nabla \varphi_n \cdot \mathbf{v}_n + \varphi_n \Gamma_{\mathbf{v}}(\varphi_n, \sigma_n) - \Gamma_{\varphi}(\varphi_n, \sigma_n)) \Phi \mathrm{d}x, \quad (4.4b)$$

$$0 = \int_{\Omega} \nabla \sigma_n \cdot \nabla \Phi + h(\varphi_n) \sigma_n \Phi dx + \int_{\partial \Omega} K(\sigma_n - \sigma_\infty) \Phi d\mathcal{H}^{d-1}, \qquad (4.4c)$$

for a.e. $t \in (0,T)$ and for all $\Phi \in \mathbf{H}^1$, $\Phi \in H^1$, where μ_n is given by

$$\mu_n = \epsilon^{-1} \Psi'(\varphi_n) - \epsilon \Delta \varphi_n - \chi \sigma_n \quad \text{a. e in } Q$$
(4.4d)

and

$$\mathbf{T}_{n}(\mathbf{v}_{n}, p_{n}) \coloneqq 2\eta_{n}(\varphi_{n})\mathbf{D}\mathbf{v}_{n} + \lambda_{n}(\varphi_{n})\operatorname{div}(\mathbf{v}_{n})\mathbf{I} - p_{n}\mathbf{I}$$

We will denote $\Gamma_{\mathbf{v},n} = \Gamma_{\mathbf{v}}(\varphi_n, \sigma_n), \ \Gamma_{\varphi,n} = \Gamma_{\varphi}(\varphi_n, \sigma_n).$

4.1 A-priori-estimates

In the following, we will derive estimates which are independent of $n \in \mathbb{N}$. By C, we will denote a generic constant depending on the system parameters and on Ω , T, but not on $n \in \mathbb{N}$. Furthermore, we will frequently use Hölder's and Young's inequalities.

First, we recall that (A1), (A5) and the continuous embedding $H^1 \hookrightarrow L^6$ imply that $\psi(\varphi_0) \in L^1$, $\nabla \varphi_0 \in L^2$. Then, using (A1), (3.25)-(3.26) and (4.2), taking the supremum over all $s \in (0, T]$ in (3.26) yields

$$\sup_{s \in (0,T]} \left(\|\psi(\varphi_n(s))\|_{L^1} + \|\nabla\varphi_n(s)\|_{\mathbf{L}^2}^2 \right) + \int_0^T \frac{m_0}{4} \|\nabla\mu_n\|_{\mathbf{L}^2}^2 + \frac{\nu}{2} \|\mathbf{v}_n\|_{\mathbf{L}^2}^2 + \|\sqrt{\eta_n(\varphi_n)}\mathbf{D}\mathbf{v}_n\|_{\mathbf{L}^2}^2 + \|\sigma_n\|_{H^1}^2 \mathrm{d}t \le C.$$

$$(4.5)$$

Recalling (2.5), using Poincaré's inequality, (3.11), (4.5) and the fact that $\rho \ge 2$, this in particular gives

$$\underset{s \in (0,T]}{\text{ess}} \sup_{s \in (0,T]} \|\varphi_n(s)\|_{H^1}^2 + \int_0^T \|\mu_n\|_{H^1}^2 \, \mathrm{d}t \le C.$$
(4.6)

Now, using exactly the same arguments as in Subsection 3.1.6, we obtain

$$\|\varphi_n\|_{L^4(H^2)} \le C. \tag{4.7}$$

Then, using (A5), it is straightforward to check that $\psi'(\varphi) \in L^2(H^1)$ with bounded norm. Together with (4.5)-(4.7) and using elliptic regularity theory, this implies

$$\|\varphi_n\|_{L^4(H^2)\cap L^2(H^3)} \le C. \tag{4.8}$$

By the specific form of $\Gamma_{\mathbf{v}}(\cdot,\cdot)$, $\Gamma_{\varphi}(\cdot,\cdot)$ and the continuous embedding $H^1 \hookrightarrow L^6$, an application of (4.3) and (4.5) yields

$$\|\Gamma_{\mathbf{v},n}\|_{L^2(L^6)} + \|\Gamma_{\varphi,n}\|_{L^2(L^6)} \le C.$$
(4.9)

Using (3.32) and (4.2), for every $n \in \mathbb{N}$ we obtain

$$\|p_n\|_{L^2}^2 \le C\left(\|\sqrt{\eta_n(\varphi_n)}\mathbf{D}\mathbf{v}_n\|_{\mathbf{L}^2}^2 + \|\Gamma_{\mathbf{v},n}\|_{L^2}^2 + \|\mathbf{v}_n\|_{\mathbf{L}^2}^2 + \|\mu_n + \chi\sigma_n\|_{L^3}^2\|\nabla\varphi_n\|_{\mathbf{L}^2}^2\right).$$

Integrating this inequality in time from 0 to T and using (4.5)-(4.6), (4.9), we have

$$\|p_n\|_{L^2(L^2)} \le C. \tag{4.10}$$

Using (4.6)-(4.9) and the continuous embedding $H^1 \hookrightarrow L^3$, we calculate

$$\|\varphi_n \Gamma_{\mathbf{v},n}\|_{L^2(L^2)} \le \|\varphi_n\|_{L^{\infty}(L^3)} \|\Gamma_{\mathbf{v},n}\|_{L^2(L^6)} \le C \|\varphi_n\|_{L^{\infty}(H^1)} \|\Gamma_{\mathbf{v},n}\|_{L^2(L^6)} \le C.$$
(4.11)

Now, let $\rho \in L^{\frac{8}{3}}(H^1)$. Then, using (1.10) and (4.5)-(4.8), we obtain

$$\left| \int_{0}^{T} \int_{\Omega} \nabla \varphi_{n} \cdot \mathbf{v}_{n} \rho \, \mathrm{d}x \, \mathrm{d}t \right| \leq \int_{0}^{T} \|\nabla \varphi_{n}\|_{\mathbf{L}^{3}} \|\mathbf{v}_{n}\|_{\mathbf{L}^{2}} \|\rho\|_{H^{1}} \, \mathrm{d}t$$
$$\leq \int_{0}^{T} \|\varphi_{n}\|_{H^{1}}^{\frac{3}{4}} \|\varphi_{n}\|_{H^{3}}^{\frac{1}{4}} \|\mathbf{v}_{n}\|_{\mathbf{L}^{2}} \|\rho\|_{H^{1}} \, \mathrm{d}t$$
$$\leq \|\varphi_{n}\|_{L^{\infty}(H^{1})}^{\frac{3}{4}} \|\varphi_{n}\|_{L^{2}(H^{3})}^{\frac{1}{4}} \|\mathbf{v}_{n}\|_{L^{2}(\mathbf{L}^{2})} \|\rho\|_{L^{\frac{8}{3}}(H^{1})}$$
$$\leq C \|\rho\|_{L^{\frac{8}{3}}(H^{1})},$$

which implies

$$\left\|\nabla\varphi_n\cdot\mathbf{v}_n\right\|_{L^{\frac{8}{5}}((H^1)^*)}\leq C$$

Using the continuous embedding $L^2 \hookrightarrow (H^1)^*$ and (4.3), (4.11), we deduce that

$$\|\operatorname{div}(\varphi_n \mathbf{v}_n)\|_{L^{\frac{8}{5}}((H^1)^*)} \le C.$$
 (4.12)

Finally, a comparison argument in (4.4b) yields

$$\|\partial_t \varphi_n\|_{L^{\frac{8}{5}}((H^1)^*)} \le C,$$
(4.13)

where we used (4.5)-(4.9) and (4.12). Summarising (4.5)-(4.13), we end up with

$$\begin{aligned} \|\varphi_n\|_{W^{1,\frac{8}{5}}((H^1)^*)\cap L^{\infty}(H^1)\cap L^4(H^2)\cap L^2(H^3)} + \|\mu_n\|_{L^2(H^1)} + \|\sigma_n\|_{L^2(H^1)} + \|\operatorname{div}(\varphi_n\mathbf{v}_n)\|_{L^{\frac{8}{5}}((H^1)^*)} \\ + \|\operatorname{div}(\mathbf{v}_n)\|_{L^2(L^6)} + \|\sqrt{\eta_n(\varphi_n)}\mathbf{D}\mathbf{v}_n\|_{L^2(\mathbf{L}^2)} + \|\mathbf{v}_n\|_{L^2(\mathbf{L}^2_{\operatorname{div}}(\Omega))} + \|p_n\|_{L^2(L^2)} \le C. \end{aligned}$$
(4.14)

4.2 Passing to the limit

Recalling (1.9) and (4.14), using standard compactness arguments (Aubin-Lions theorem (see [39, Sec. 8, Cor. 4]) and reflexive weak compactness), the compact embeddings

$$H^{j+1}(\Omega) = W^{j+1,2}(\Omega) \hookrightarrow W^{j,r} \quad \forall j \in \mathbb{Z}, \, j \ge 0, \, 1 \le r < 6,$$

and $L^2 \hookrightarrow \hookrightarrow (H^1)^*$, we obtain, at least for a subsequence which will again be labelled by n, the following convergence results:

 $\begin{array}{lll} \varphi_n \rightarrow \varphi & \mbox{weakly-*} & \mbox{in } W^{1,\frac{8}{5}}((H^1)^*) \cap L^{\infty}(H^1) \cap L^4(H^2) \cap L^2(H^3), \\ \sigma_n \rightarrow \sigma & \mbox{weakly} & \mbox{in } L^2(H^1), \\ \mu_n \rightarrow \mu & \mbox{weakly} & \mbox{in } L^2(L^2), \\ \mathbf{v}_n \rightarrow \mathbf{v} & \mbox{weakly} & \mbox{in } L^2(\mathbf{L}^2) \cap L^2(L^2_{\rm div}(\Omega)), \\ {\rm div}(\mathbf{v}_n) \rightarrow \xi & \mbox{weakly} & \mbox{in } L^2(L^2), \\ \mathbf{v}_n \cdot \mathbf{n} \rightarrow \mathbf{v} \cdot \mathbf{n} & \mbox{weakly} & \mbox{in } L^2((H^{\frac{1}{2}}(\partial\Omega))^*), \\ {\rm div}(\varphi_n \mathbf{v}_n) \rightarrow \tau & \mbox{weakly} & \mbox{in } L^{\frac{8}{5}}((H^1)^*), \end{array}$

for some limit functions $\xi \in L^2(L^2)$, $\tau \in L^{\frac{8}{5}}((H^1)^*)$. Furthermore, by (4.1), we have the strong convergences

$$\begin{split} \varphi_n &\to \varphi \quad \text{strongly} \quad \text{ in } C^0(L^r) \cap L^2(W^{2,r}) \text{ and a.e. in } Q, \\ \eta_n(\varphi_n) &\to 0 \qquad \text{ in } Q, \\ \lambda_n(\varphi_n) &\to 0 \qquad \text{ in } Q, \end{split}$$

for $r \in [1,6)$. In the following, we fix $\delta \in C_0^{\infty}(0,T)$, $\Phi \in H^1$, $\Phi \in \mathbf{H}^1$ and we note that $\delta \Phi \in C^{\infty}(H^1)$, $\delta \Phi \in C^{\infty}(\mathbf{H}^1)$. Multiplying (4.4a)-(4.4d) with δ and integrating in time from 0 to T and multiplying (4.4a) with $\delta \Phi$ and integrating over Q, we obtain

$$0 = \int_{0}^{T} \int_{\Omega} \delta(t) (2\eta_{n}(\varphi_{n}) \mathbf{D} \mathbf{v}_{n} + \lambda_{n}(\varphi_{n}) \operatorname{div}(\mathbf{v}_{n}) \mathbf{I} - p_{n} \mathbf{I}) \colon \nabla \mathbf{\Phi} + \nu \mathbf{v}_{n} \cdot \mathbf{\Phi}) \mathrm{d}x \mathrm{d}t$$

$$- \int_{0}^{T} \int_{\Omega} \delta(t) (\mu_{n} + \chi \sigma_{n}) \nabla \varphi_{n} \cdot \mathbf{\Phi} \mathrm{d}x \mathrm{d}t, \qquad (4.15a)$$

$$0 = \int_{0}^{T} \delta(t) \langle \partial_{t} \varphi_{n}, \Phi \rangle_{H^{1}, (H^{1})^{*}} dt + \int_{0}^{T} \int_{\Omega} \delta(t) (m(\varphi_{n}) \nabla \mu_{n} \cdot \nabla \Phi - \Gamma_{\varphi, n} \Phi) dx dt + \int_{0}^{T} \int_{\Omega} \delta(t) (\nabla \varphi_{n} \cdot \mathbf{v}_{n} + \varphi_{n} \Gamma_{\mathbf{v}, n}) \Phi dx dt,$$
(4.15b)

$$0 = \int_0^T \int_\Omega \delta(t) \mu_n \Phi dx dt - \int_0^T \int_\Omega \delta(t) (\epsilon^{-1} \Psi'(\varphi_n) - \epsilon \Delta \varphi_n - \chi \sigma_n) \Phi dx dt, \qquad (4.15c)$$

$$0 = \int_0^T \int_\Omega \delta(t) (\nabla \sigma_n \cdot \nabla \Phi + h(\varphi_n) \sigma_n \Phi) dx dt + \int_0^T \int_{\partial \Omega} \delta(t) (K(\sigma_n - \sigma_\infty) \Phi) d\mathcal{H}^{d-1} dt.$$
(4.15d)

Furthermore, we multiply $(4.3)_1$ with $\delta\Phi$ and integrate over Q to obtain

$$\int_{0}^{T} \int_{\Omega} \delta \operatorname{div}(\mathbf{v}_{n}) \Phi \mathrm{d}x \mathrm{d}t = \int_{0}^{T} \int_{\Omega} \delta(t) \Gamma_{\mathbf{v},n} \Phi \mathrm{d}x \mathrm{d}t.$$
(4.15e)

We now want to analyse each term individually. For (4.15c)-(4.15d), we omit the details and refer to the arguments used in [16, Sec. 5], [23, Sec. 5].

Step 1 ((4.15e)): Since $\varphi_n \to \varphi$ a.e. in Q and due to the boundedness of $b_{\mathbf{v}}(\cdot)$, $f_{\mathbf{v}}(\cdot)$, Lebesgue dominated convergence theorem implies

$$\|\delta\Phi(b_{\mathbf{v}}(\varphi_n) - b_{\mathbf{v}}(\varphi))\|_{L^2(Q)} \to 0, \quad \|\delta\Phi(f_{\mathbf{v}}(\varphi_n) - f_{\mathbf{v}}(\varphi))\|_{L^2(Q)} \to 0$$

as $n \to \infty$. Together with the weak convergence $\sigma_n \to \sigma$ in $L^2(Q)$ as $n \to \infty$, by the product of weak-strong convergence we obtain

$$\int_{0}^{T} \int_{\Omega} \delta(t) \Gamma_{\mathbf{v},n} \Phi dx dt \to \int_{0}^{T} \int_{\Omega} \delta(t) \Gamma_{\mathbf{v}}(\varphi, \sigma) \Phi dx dt \quad \text{as } n \to \infty.$$
(4.16)

Using (1.8), we see that

$$\int_0^T \int_\Omega \delta \operatorname{div}(\mathbf{v}_n) \Phi \mathrm{d}x \mathrm{d}t = \int_0^T \delta(t) \langle \mathbf{v}_n \cdot \mathbf{n}, \Phi \rangle_{H^{\frac{1}{2}}(\partial \Omega)} \mathrm{d}t - \int_0^T \int_\Omega \delta(t) \mathbf{v}_n \cdot \nabla \Phi \mathrm{d}x \mathrm{d}t.$$

Since $\mathbf{v}_n \cdot \mathbf{n} \to \mathbf{v} \cdot \mathbf{n}$ weakly in $L^2((H^{\frac{1}{2}}(\partial \Omega))^*)$, $\mathbf{v}_n \to \mathbf{v}$ weakly in $L^2(\mathbf{L}^2)$ and $\operatorname{div}(\mathbf{v}_n) \to \xi$ weakly in $L^2(L^2)$ as $n \to \infty$, we can pass to the limit on both sides of this equation to obtain

$$\int_0^T \int_\Omega \delta\xi \Phi \mathrm{d}x \mathrm{d}t = \int_0^T \delta(t) \langle \mathbf{v} \cdot \mathbf{n}, \Phi \rangle_{H^{\frac{1}{2}}(\partial\Omega)} \mathrm{d}t - \int_0^T \int_\Omega \delta(t) \mathbf{v} \cdot \nabla \Phi \mathrm{d}x \mathrm{d}t$$

Since $\mathbf{v} \in L^2_{\text{div}}(\Omega)$, we can again use (1.8) to obtain

$$\int_0^T \int_\Omega \delta\xi \Phi dx dt = \int_0^T \int_\Omega \delta div(\mathbf{v}) \Phi dx dt.$$

This in particular gives

 $\operatorname{div}(\mathbf{v}) = \xi \quad \text{a.e. in } Q. \tag{4.17}$

From this considerations and recalling (4.16), we can pass to the limit $n \to \infty$ in (4.15e) to obtain

$$\int_{0}^{T} \int_{\Omega} \delta \operatorname{div}(\mathbf{v}) \Phi \mathrm{d}x \mathrm{d}t = \int_{0}^{T} \int_{\Omega} \delta \Gamma_{\mathbf{v}}(\varphi, \sigma) \Phi \mathrm{d}x \mathrm{d}t,$$

$$\lim_{t \to 0} \left(\sum_{i=1}^{T} \int_{\Omega} \delta \Gamma_{\mathbf{v}}(\varphi, \sigma) \Phi \mathrm{d}x \mathrm{d}t,$$

$$(1.10)$$

which in particular implies

 $\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v}}(\varphi, \sigma) \quad \text{a.e. in } Q.$ (4.18)

Step 2 ((4.15b)): Since $\delta \Phi \in C^{\infty}(H^1)$ and $\operatorname{div}(\varphi_n \mathbf{v}_n) \to \tau$ weakly in $L^{\frac{8}{5}}((H^1)^*)$, we have

$$\int_{0}^{T} \int_{\Omega} \delta \operatorname{div}(\varphi_{n} \mathbf{v}_{n}) \Phi \mathrm{d}x \mathrm{d}t \to \int_{0}^{T} \delta(t) \langle \tau, \Phi \rangle_{H^{1}} \mathrm{d}t \quad \text{as } n \to \infty.$$

$$(4.19)$$

Moreover, since $\nabla \varphi_n \to \nabla \varphi$ strongly in $L^2(\mathbf{L}^3)$ and due to the continuous embedding $H^1 \hookrightarrow L^6$, we have

$$\int_0^T \int_\Omega |\delta|^2 |\Phi|^2 |\nabla\varphi_n - \nabla\varphi|^2 \mathrm{d}x \mathrm{d}t \le \int_0^T |\delta|^2 ||\Phi||_{L^6}^2 ||\nabla\varphi_n - \nabla\varphi||_{\mathbf{L}^3}^2 \mathrm{d}t$$
$$\le C ||\delta||_{L^\infty(0,T)}^2 ||\Phi||_{H^1}^2 ||\nabla\varphi_n - \nabla\varphi||_{L^2(\mathbf{L}^3)}^2 \to 0$$

as $n \to \infty$. This implies $\delta \Phi \nabla \varphi_n \to \delta \phi \nabla \varphi$ strongly in $L^2(\mathbf{L}^2)$. Together with the weak convergence $\mathbf{v}_n \to \mathbf{v}$ in $L^2(\mathbf{L}^2)$ as $n \to \infty$, by the product of weak-strong convergence we get

$$\int_{0}^{T} \int_{\Omega} \delta \Phi \nabla \varphi_{n} \cdot \mathbf{v}_{n} \mathrm{d}x \mathrm{d}t \to \int_{0}^{T} \int_{\Omega} \delta \Phi \nabla \varphi \cdot \mathbf{v} \mathrm{d}x \mathrm{d}t \quad \text{as } n \to \infty.$$
(4.20)

Since $\varphi_n \to \varphi$ strongly in $L^2(L^3)$ and a.e. in Q as $n \to \infty$, the boundedness of $b_{\mathbf{v}}(\cdot)$, $f_{\mathbf{v}}(\cdot)$ and Lebesgue dominated convergence theorem imply

$$\|(b_{\mathbf{v}}(\varphi_n)\varphi_n - b_{\mathbf{v}}(\varphi)\varphi)\delta\Phi\|_{L^2(Q)} \to 0, \qquad \|(f_{\mathbf{v}}(\varphi_n)\varphi_n - f_{\mathbf{v}}(\varphi)\varphi)\delta\Phi\|_{L^2(Q)} \to 0$$

as $n \to \infty$, where we used that $\Phi \in H^1 \hookrightarrow L^6$. Together with the weak convergence $\sigma_n \to \sigma$ in $L^2(Q)$ as $n \to \infty$, this implies

$$\int_{0}^{T} \int_{\Omega} \delta \Phi \Gamma_{\mathbf{v},n} \varphi_{n} \mathrm{d}x \mathrm{d}t \to \int_{0}^{T} \int_{\Omega} \delta \Phi \Gamma_{\mathbf{v}}(\varphi,\sigma) \varphi \mathrm{d}x \mathrm{d}t \quad \text{as } n \to \infty.$$
(4.21)

Using $(4.3)_1$, we see that

$$\int_0^T \int_\Omega \delta \operatorname{div}(\varphi_n \mathbf{v}_n) \Phi \mathrm{d}x \mathrm{d}t = \int_0^T \int_\Omega \delta \Phi \nabla \varphi_n \cdot \mathbf{v}_n \mathrm{d}x \mathrm{d}t + \int_0^T \int_\Omega \delta \Phi \Gamma_{\mathbf{v},n} \varphi_n \mathrm{d}x \mathrm{d}t.$$

Passing to the limit $n \to \infty$ on both sides of this equation and using (4.20)-(4.21), we obtain

$$\int_{0}^{T} \delta(t) \langle \tau, \Phi \rangle_{H^{1}} \mathrm{d}t = \int_{0}^{T} \int_{\Omega} \delta \Phi \nabla \varphi \cdot \mathbf{v} \mathrm{d}x \mathrm{d}t + \int_{0}^{T} \int_{\Omega} \delta \Phi \Gamma_{\mathbf{v}}(\varphi, \sigma) \varphi \mathrm{d}x \mathrm{d}t.$$
(4.22)

Together with (4.18), this gives

$$\int_{0}^{T} \delta(t) \langle \tau, \Phi \rangle_{H^{1}, (H^{1})^{*}} dt = \int_{0}^{T} \int_{\Omega} \delta(t) \operatorname{div}(\varphi \mathbf{v}) \Phi dx dt,$$
(4.23)

hence $\operatorname{div}(\varphi \mathbf{v}) = \tau$ in the sense of distributions. For the remaining terms in (4.15b), we again refer to [16, Sec. 5], [23, Sec. 5], where they used similar arguments.

Step 3 ((4.15a)): With exactly the same arguments as used for (4.20), it follows that $\delta \Phi \cdot \nabla \varphi_n \to \delta \Phi \cdot \nabla \varphi$ strongly in $L^2(L^2)$ as $n \to \infty$. Then, recalling that $\mu_n + \chi \sigma_n \to \mu + \chi \sigma$ weakly in $L^2(L^2)$ as $n \to \infty$, by the product of weak-strong convergence we obtain

$$\int_{0}^{T} \int_{\Omega} \delta(t)(\mu_{n} + \chi\sigma_{n})\nabla\varphi_{n} \cdot \mathbf{\Phi} dx dt \to \int_{0}^{T} \int_{\Omega} \delta(t)(\mu + \chi\sigma)\nabla\varphi \cdot \mathbf{\Phi} dx dt \quad \text{as } n \to \infty.$$
(4.24)

Now, since $p_n \to p$, $\mathbf{v}_n \to \mathbf{v}$ weakly in $L^2(L^2)$ and $L^2(\mathbf{L}^2)$ respectively and observing

$$\int_0^T \int_\Omega \delta p_n \mathbf{I} \colon \nabla \mathbf{\Phi} \mathrm{d}x \mathrm{d}t = \int_0^T \int_\Omega \delta p_n \mathrm{div}(\mathbf{\Phi}) \mathrm{d}x \mathrm{d}t,$$

we obtain that

$$\int_{0}^{T} \int_{\Omega} \delta(-p_{n} \operatorname{div}(\boldsymbol{\Phi}) + \nu \mathbf{v}_{n} \cdot \boldsymbol{\Phi}) \mathrm{d}x \mathrm{d}t \to \int_{0}^{T} \int_{\Omega} \delta(-p \operatorname{div}(\boldsymbol{\Phi}) + \nu \mathbf{v} \cdot \boldsymbol{\Phi}) \mathrm{d}x \mathrm{d}t \quad \text{as } n \to \infty.$$
(4.25)

Finally, we recall that $\eta_n(\varphi_n) \to 0$ a.e. in Q as $n \to \infty$. Consequently, applying (4.14) yields

$$\left| \int_{0}^{T} \int_{\Omega} \delta(t) 2\eta_{n}(\varphi_{n}) \mathbf{D} \mathbf{v}_{n} \colon \nabla \mathbf{\Phi} dx dt \right| \leq C \|\sqrt{\eta_{n}(\varphi_{n})} \mathbf{D} \mathbf{v}_{n}\|_{L^{2}(\mathbf{L}^{2})} \|\sqrt{\eta_{n}(\varphi_{n})}\|_{L^{\infty}(Q)} \|\delta\|_{L^{\infty}(0,T)} \|\mathbf{\Phi}\|_{\mathbf{H}^{1}}$$
$$\leq C \|\sqrt{\eta_{n}(\varphi_{n})}\|_{L^{\infty}(Q)} \|\delta\|_{L^{\infty}(0,T)} \|\mathbf{\Phi}\|_{\mathbf{H}^{1}}$$
$$\to 0 \quad \text{as } n \to \infty.$$
(4.26)

Using that $\lambda_n(\varphi_n) \to 0$ a.e. in Q as $n \to \infty$ and applying (4.14), it follows that

$$\left| \int_{0}^{T} \int_{\Omega} \delta(t) \lambda_{n}(\varphi_{n}) \operatorname{div}(\mathbf{v}_{n}) \mathbf{I} \colon \nabla \mathbf{\Phi} \mathrm{d}x \mathrm{d}t \right| \leq C \| \operatorname{div}(\mathbf{v}_{n})\|_{L^{2}(L^{2})} \|\lambda_{n}(\varphi_{n})\|_{L^{\infty}(Q)} \|\delta\|_{L^{\infty}(0,T)} \|\mathbf{\Phi}\|_{\mathbf{H}^{1}}$$
$$\leq C \|\lambda_{n}(\varphi_{n})\|_{L^{\infty}(Q)} \|\delta\|_{L^{\infty}(0,T)} \|\mathbf{\Phi}\|_{\mathbf{H}^{1}}$$
$$\to 0 \quad \text{as } n \to \infty.$$
(4.27)

Step 4: Due to (4.16)-(4.27), we have enough to pass to the limit $n \to \infty$ in (4.15) to obtain that

$$0 = \int_{0}^{T} \delta(t) \left(\int_{\Omega} -p \operatorname{div}(\mathbf{\Phi}) + (\nu \mathbf{v} - (\mu + \chi \sigma) \nabla \varphi) \cdot \mathbf{\Phi} \mathrm{d}x \right) \mathrm{d}t,$$
(4.28a)

$$0 = \int_{0}^{T} \delta(t) \left(\langle \partial_{t} \varphi, \Phi \rangle_{H^{1}} + \int_{\Omega} m(\varphi) \nabla \mu \cdot \nabla \Phi + (\nabla \varphi \cdot \mathbf{v} + \varphi \Gamma_{\mathbf{v}}(\varphi, \sigma) - \Gamma_{\varphi}(\varphi, \sigma)) \Phi \mathrm{d}x \right) \mathrm{d}t, \qquad (4.28b)$$

$$0 = \int_{0}^{T} \delta(t) \left(\int_{\Omega} (\mu - \epsilon^{-1} \Psi'(\varphi) + \epsilon \Delta \varphi + \chi \sigma) \Phi dx \right) dt,$$
(4.28c)

$$0 = \int_0^T \delta(t) \left(\int_\Omega \nabla \sigma \cdot \nabla \Phi + h(\varphi) \sigma \Phi dx + \int_{\partial \Omega} K(\sigma - \sigma_\infty) \Phi d\mathcal{H}^{d-1} \right) dt,$$
(4.28d)

for all $\delta \in C_0^\infty(0,T)$ and

$$\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v}}(\varphi, \sigma) \quad \text{a.e. in } Q.$$
(4.29a)

Since (4.28) holds for all $\delta \in C_0^{\infty}(0,T)$, we deduce that

$$0 = \int_{\Omega} -p \operatorname{div}(\mathbf{\Phi}) + (\nu \mathbf{v} - (\mu + \chi \sigma) \nabla \varphi) \cdot \mathbf{\Phi} dx, \qquad (4.29b)$$

$$0 = \langle \partial_t \varphi, \Phi \rangle_{H^1} + \int_{\Omega} m(\varphi) \nabla \mu \cdot \nabla \Phi + (\nabla \varphi \cdot \mathbf{v} + \varphi \Gamma_{\mathbf{v}}(\varphi, \sigma) - \Gamma_{\varphi}(\varphi, \sigma)) \Phi \mathrm{d}x, \qquad (4.29c)$$

$$0 = \int_{\Omega} (\mu - \epsilon^{-1} \Psi'(\varphi) + \epsilon \Delta \varphi + \chi \sigma) \Phi dx, \qquad (4.29d)$$

$$0 = \int_{\Omega} \nabla \sigma \cdot \nabla \Phi + h(\varphi) \sigma \Phi dx + \int_{\partial \Omega} K(\sigma - \sigma_{\infty}) \Phi d\mathcal{H}^{d-1}, \qquad (4.29e)$$

holds for a.e. $t \in (0,T)$ and all $\Phi \in H^1$, $\Phi \in \mathbf{H}^1$. The initial condition is satisfied since $\varphi_n(0) = \varphi_0$ a.e. in Ω and by the strong convergence $\varphi_n \to \varphi$ in $C^0(L^2)$ as $n \to \infty$. By the weak (weak-star) lower semi-continuity of norms and (4.14), we obtain that $(\varphi, \mu, \sigma, \mathbf{v}, p)$ satisfies

$$\begin{aligned} \|\varphi\|_{W^{1,\frac{8}{5}}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\mu\|_{L^{2}(H^{1})} + \|\sigma\|_{L^{2}(H^{1})} + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{\frac{8}{5}}((H^{1})^{*})} \\ &+ \|\operatorname{div}(\mathbf{v})\|_{L^{2}(L^{6})} + \|\mathbf{v}\|_{L^{2}(\mathbf{L}^{2}_{\operatorname{div}}(\Omega))} + \|p\|_{L^{2}(L^{2})} \leq C. \end{aligned}$$

$$(4.30)$$

Step 5: Using (4.29b) and (4.30), we obtain that p has a weak derivative in $L^{\frac{8}{5}}(L^2)$ and it holds

$$\nabla p = -\nu \mathbf{v} + (\mu + \chi \sigma) \nabla \varphi \quad \text{a.e. in } Q.$$
(4.31)

By (1.10), we have

$$\|\nabla\varphi\|_{\mathbf{L}^{3}} \le C \|\nabla\varphi\|_{\mathbf{L}^{2}}^{\frac{3}{4}} \|\nabla\varphi\|_{\mathbf{H}^{2}}^{\frac{1}{4}} \le C \|\varphi\|_{H^{1}}^{\frac{3}{4}} \|\varphi\|_{H^{3}}^{\frac{1}{4}}$$

Using (4.30), this implies

$$\|\nabla\varphi\|_{L^8(\mathbf{L}^3)} \le C.$$

Then, using the continuous embedding $H^1 \hookrightarrow L^6$ and (4.30) again, we obtain

$$\|(\mu + \chi \sigma) \nabla \varphi\|_{L^{\frac{8}{5}}(\mathbf{L}^2)} \le C.$$

Since $\mathbf{v} \in L^2(\mathbf{L}^2)$ and $p \in L^2(L^2)$ with bounded norm, from (4.31) we obtain

$$\|p\|_{L^2(L^2)\cap L^{\frac{8}{5}}(H^1)} \le C.$$
(4.32)

Integrating (4.29b) by parts, we obtain

$$-\int_{\partial\Omega} p\mathbf{\Phi} \cdot \mathbf{n} \mathrm{d}\mathcal{H}^{d-1} = \int_{\Omega} (-\nabla p + (\mu + \chi\sigma)\nabla\varphi - \nu\mathbf{v}) \cdot \mathbf{\Phi} \mathrm{d}x$$

for all $\mathbf{\Phi} \in \mathbf{H}^1$ and a.e. $t \in (0, T)$. Because of (4.31), this implies

$$\int_{\partial\Omega} p \mathbf{\Phi} \cdot \mathbf{n} \mathrm{d} \mathcal{H}^{d-1} = 0$$

for all $\mathbf{\Phi} \in \mathbf{H}^1$ and a.e. $t \in (0, T)$. Therefore, we obtain

$$p = 0 \quad \text{a.e. on } \Sigma. \tag{4.33}$$

With similar arguments, it is straightforward to show that

$$\mu = \epsilon^{-1} \psi'(\varphi) - \epsilon \Delta \varphi - \chi \sigma \quad \text{a.e. in } Q, \qquad \partial_{\mathbf{n}} \varphi = 0 \quad \text{a.e. on } \Sigma, \tag{4.34}$$

which completes the proof.

5 Continuous dependence

In the following, since it has no bearing on the further analysis, we set $\epsilon = 1$. Let $(\varphi_i, \mu_i, \sigma_i, \mathbf{v}_i, p_i)_{i=1,2}$ be two solutions of (1.1), (1.3) in the sense of Definition 2.3. We denote $\Gamma_{\mathbf{v}}(\varphi_i, \sigma_i) \coloneqq \Gamma_{\mathbf{v},i}, \Gamma_{\varphi}(\varphi_i, \sigma_i) \coloneqq \Gamma_{\varphi,i}, i = 1, 2$, and $\sigma_{\infty} \coloneqq \sigma_{1,\infty} - \sigma_{2,\infty}$. Then, the difference $f \coloneqq f_1 - f_2, f_i \in {\varphi_i, \mu_i, \sigma_i, \mathbf{v}_i, p_i}, i = 1, 2$, satisfies

$$\operatorname{div}(\mathbf{v}) = \Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2} \quad \text{a.e. in } Q, \quad \varphi(0) = \varphi_{1,0} - \varphi_{2,0} \eqqcolon \varphi_0 \quad \text{a.e. in } \Omega,$$

and

$$0 = \int_{\Omega} (2\eta(\varphi_1)\mathbf{D}\mathbf{v} + \lambda(\varphi_1)\operatorname{div}(\mathbf{v})\mathbf{I} - p\mathbf{I}) \colon \nabla \Phi + \nu \mathbf{v} \cdot \Phi \, \mathrm{d}x - \int_{\Omega} (\mu + \chi\sigma)\nabla\varphi_1 \cdot \Phi + (\mu_2 + \chi\sigma_2)\nabla\varphi \cdot \Phi \, \mathrm{d}x + \int_{\Omega} (2(\eta(\varphi_1) - \eta(\varphi_2))\mathbf{D}\mathbf{v}_2 + (\lambda(\varphi_1) - \lambda(\varphi_2))\operatorname{div}(\mathbf{v}_2)\mathbf{I}) \colon \nabla \Phi \, \mathrm{d}x,$$
(5.1a)

$$0 = \langle \partial_t \varphi, \Phi \rangle_{H^1, (H^1)^*} + \int_{\Omega} \nabla \mu \cdot \nabla \Phi + (\varphi_2(\Gamma_{\mathbf{v}, 1} - \Gamma_{\mathbf{v}, 2}) - (\Gamma_{\varphi, 1} - \Gamma_{\varphi, 2})) \Phi \, \mathrm{d}x + \int_{\Omega} (\nabla \varphi_1 \cdot \mathbf{v} + \nabla \varphi \cdot \mathbf{v}_2) \Phi + \varphi \Gamma_{\mathbf{v}, 1} \Phi \, \mathrm{d}x,$$
(5.1b)

$$0 = \int_{\Omega} \nabla \sigma \cdot \nabla \Phi \, \mathrm{d}x + \int_{\Omega} (h(\varphi_1)\sigma + \sigma_2(h(\varphi_1) - h(\varphi_2))\Phi \, \mathrm{d}x + \int_{\partial\Omega} K(\sigma - \sigma_\infty)\Phi \mathrm{d}\mathcal{H}^{d-1}, \tag{5.1c}$$

for a.e. $t \in (0,T)$ and for all $\mathbf{\Phi} \in \mathbf{H}^1$, $\Phi \in H^1$, where μ is given by

$$\mu = \psi'(\varphi_1) - \psi'(\varphi_2) - \Delta \varphi - \chi \sigma.$$
(5.2)

Step 1: Taking $\Phi = \sigma$ in (5.1c), we obtain

$$\int_{\Omega} |\nabla \sigma|^2 \, \mathrm{d}x + K \int_{\partial \Omega} |\sigma|^2 \mathrm{d}\mathcal{H}^{d-1} + \int_{\Omega} h(\varphi_1) |\sigma|^2 \, \mathrm{d}x = -\int_{\Omega} \sigma_2 (h(\varphi_1) - (\varphi_2)) \sigma \, \mathrm{d}x + K \int_{\partial \Omega} \sigma \sigma_{\infty} \mathrm{d}\mathcal{H}^{d-1}.$$

Using the non-negativity of $h(\cdot)$, we can neglect the third term on the l.h.s. of this equation. Applying Hölder's and Young's inequalities, we therefore obtain

$$\int_{\Omega} |\nabla \sigma|^2 \, \mathrm{d}x + K \int_{\partial \Omega} |\sigma|^2 \mathrm{d}\mathcal{H}^{d-1} \le \frac{L_h^2}{4\delta} \|\sigma_2\|_{L^3}^2 \|\varphi\|_{L^2}^2 + \frac{K}{2} \left(\|\sigma\|_{L^2(\partial\Omega)}^2 + \|\sigma_\infty\|_{L^2(\partial\Omega)}^2 \right) + \delta \|\sigma\|_{L^6}^2, \quad (5.3)$$

with $\delta > 0$ to be chosen and where we used (B2). Using Poincaré's inequality and the continuous embedding $H^1 \hookrightarrow L^6$, we have

$$\|\sigma\|_{L^{6}}^{2} \leq 2C_{P}^{2}\left(\|\nabla\sigma\|_{\mathbf{L}^{2}}^{2} + \|\sigma\|_{L^{2}(\partial\Omega)}^{2}\right).$$
(5.4)

Choosing

$$\delta = \frac{1}{8C_P^2} \min\{1, K\},$$

and using (5.4), this implies

$$\|\sigma\|_{H^{1}} \le C(K, L_{h}, \Omega) \left(\|\sigma_{2}\|_{L^{6}} \|\varphi\|_{L^{2}} + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)} \right).$$
(5.5)

Step 2: With similar arguments as in Section 3, we deduce the existence of a solution $\mathbf{u} \in \mathbf{H}^1$ of the problem

div
$$(\mathbf{u}) = \Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}$$
 in Ω , $\mathbf{u} = \frac{1}{|\partial\Omega|} \left(\int_{\Omega} \Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2} \, \mathrm{d}x \right) \mathbf{n}$ on $\partial\Omega$,
stimate
 $\|\mathbf{u}\|_{\mathbf{H}^1} \le c \|\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}\|_{L^2},$ (5.6)

satisfying the estimate

with a constant c depending only on Ω . Choosing $\Phi = \mathbf{v} - \mathbf{u}$ in (5.1a) and $\Phi = \varphi - \Delta \varphi$ in (5.1b), integrating by parts, using (5.2) and summing the resulting identities, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t} \frac{1}{2} (\|\nabla\varphi\|_{\mathbf{L}^{2}}^{2} + \|\varphi\|_{L^{2}}^{2}) + \int_{\Omega} 2\eta |\mathbf{D}\mathbf{v}|^{2} + \nu |\mathbf{v}|^{2} + |\Delta\varphi|^{2} + |\nabla\Delta\varphi|^{2} \,\mathrm{d}x$$

$$= \int_{\Omega} \nabla(\psi'(\varphi_{1}) - \psi'(\varphi_{2})) \cdot (\nabla\Delta\varphi - \nabla\varphi) - ((\Gamma_{\varphi,1} - \Gamma_{\varphi,2}) - \varphi_{2}(\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}))\Delta\varphi \,\mathrm{d}x$$

$$+ \int_{\Omega} (\nabla\varphi \cdot \mathbf{v}_{2} + \varphi\Gamma_{\mathbf{v},1})\Delta\varphi + (\psi'(\varphi_{1}) - \psi'(\varphi_{2}))\nabla\varphi_{1} \cdot (\mathbf{v} - \mathbf{u}) \,\mathrm{d}x,$$

$$+ \int_{\Omega} (\mu_{2} + \chi\sigma_{2})\nabla\varphi \cdot (\mathbf{v} - \mathbf{u}) + \Delta\varphi\nabla\varphi_{1} \cdot \mathbf{u} \,\mathrm{d}x + \int_{\Omega} \chi\nabla\sigma \cdot (\nabla\varphi - \nabla\Delta\varphi)\mathrm{d}x$$

$$+ \int_{\Omega} ((\Gamma_{\varphi,1} - \Gamma_{\varphi,2}) - \varphi_{2}(\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}))\varphi - (\nabla\varphi_{1} \cdot \mathbf{v} + \nabla\varphi \cdot \mathbf{v}_{2})\varphi - \Gamma_{\mathbf{v},1}|\varphi|^{2} \,\mathrm{d}x$$

$$+ \int_{\Omega} 2\eta(\varphi_{1})\mathbf{D}\mathbf{v} \colon \nabla\mathbf{u} + \nu\mathbf{v} \cdot \mathbf{u} - 2(\eta(\varphi_{1}) - \eta(\varphi_{2}))\mathbf{D}\mathbf{v}_{2} \colon \nabla(\mathbf{v} - \mathbf{u}) \,\mathrm{d}x.$$
(5.7)

Step 3: We now estimate the terms on the r.h.s. of (5.7). In the following, we will frequently use Young's, Hölder's inequalities and Lemma 1.1. First of all, using (A4) and (B2), we obtain

$$|\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}| \leq C(|\sigma| + |\sigma_2||\varphi| + |\varphi|), \quad |\Gamma_{\varphi,1} - \Gamma_{\varphi,2}| \leq C(|\sigma| + |\sigma_2||\varphi| + |\varphi|).$$

Hence, applying (5.5) and the continuous embedding $H^1 \hookrightarrow L^6$ yields

$$\|\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2}\|_{L^2} + \|\Gamma_{\varphi,1} - \Gamma_{\varphi,2}\|_{L^2} \le C \left(1 + \|\sigma_2\|_{L^6}\right) \|\varphi\|_{H^1} + C\|\sigma_\infty\|_{L^2(\partial\Omega)}.$$
(5.8)

Using (2.8), (5.8) and the continuous embedding $H^1 \hookrightarrow L^6$, we obtain

$$\left| \int_{\Omega} \left((\Gamma_{\varphi,1} - \Gamma_{\varphi,2}) - \varphi_{2} (\Gamma_{\mathbf{v},1} - \Gamma_{\mathbf{v},2})) \Delta \varphi \, \mathrm{d}x \right| \\
\leq C \left(1 + \|\varphi_{2}\|_{L^{\infty}}^{2} \right) \left((1 + \|\sigma_{2}\|_{L^{6}}^{2}) \|\varphi\|_{H^{1}}^{2} + \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} \right) + \frac{1}{8} \|\Delta\varphi\|_{L^{2}}^{2} \\
\leq C \left(1 + \|\varphi_{2}\|_{H^{2}}^{2} + \|\sigma_{2}\|_{H^{1}}^{4} \right) \|\varphi\|_{H^{1}}^{2} + C \left(1 + \|\varphi_{2}\|_{H^{2}}^{2} \right) \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} + \frac{1}{8} \|\Delta\varphi\|_{L^{2}}^{2}.$$
(5.9)

By the specific form of $\Gamma_{\mathbf{v}}$ and (A4), applying the continuous embedding $H^1 \hookrightarrow L^6$ gives

$$\left| \int_{\Omega} \varphi \Gamma_{\mathbf{v},1} \Delta \varphi \, \mathrm{d}x \right| \le C \left(1 + \|\sigma_1\|_{L^3}^2 \right) \|\varphi\|_{H^1}^2 + \frac{1}{8} \|\Delta \varphi\|_{L^2}^2.$$

$$(5.10)$$

Hence, recalling the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ and (1.7), we calculate

$$\left| \int_{\Omega} \nabla \varphi \cdot \mathbf{v}_{2} \Delta \varphi \, \mathrm{d}x \right| \leq \| \nabla \varphi \|_{\mathbf{L}^{3}} \| \mathbf{v}_{2} \|_{\mathbf{L}^{6}} \| \Delta \varphi \|_{L^{2}}$$

$$\leq C \| \mathbf{v}_{2} \|_{\mathbf{H}^{1}} \| \nabla \varphi \|_{\mathbf{L}^{2}} \| \varphi \|_{H^{2}}^{\frac{1}{2}} \| \nabla \Delta \varphi \|_{\mathbf{L}^{2}}^{\frac{1}{2}}$$

$$\leq C \| \mathbf{v}_{2} \|_{\mathbf{H}^{1}} \| \varphi \|_{H^{1}} (\| \varphi \|_{L^{2}} + \| \Delta \varphi \|_{L^{2}} + \| \nabla \Delta \varphi \|_{\mathbf{L}^{2}})$$

$$\leq C \left(1 + \| \mathbf{v}_{2} \|_{\mathbf{H}^{1}}^{2} \right) \| \varphi \|_{H^{1}}^{2} + \frac{1}{8} \| \Delta \varphi \|_{L^{2}}^{2} + \frac{1}{4} \| \nabla \Delta \varphi \|_{\mathbf{L}^{2}}^{2}. \tag{5.11}$$

Due to (B3), (1.10), (2.8) and the continuous embedding $H^1 \hookrightarrow L^6$, we obtain

$$\|\psi'(\varphi_1) - \psi'(\varphi_2)\|_{L^2}^2 \le C\left(1 + \|\varphi_1\|_{H^2}^2 + \|\varphi_2\|_{H^2}^2\right)\|\varphi\|_{H^1}^2.$$
(5.12)

By the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ and (2.8), this gives

$$\left| \int_{\Omega} (\psi'(\varphi_{2}) - \psi'(\varphi_{1})) \nabla \varphi_{1} \cdot \mathbf{v} \, \mathrm{d}x \right| \leq C \|\psi'(\varphi_{1}) - \psi'(\varphi_{2})\|_{L^{2}}^{2} \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}}^{2} + \delta_{1} \|\mathbf{v}\|_{\mathbf{L}^{6}}^{2}$$
$$\leq C \left(1 + \|\varphi_{1}\|_{H^{2}}^{4} + \|\varphi_{2}\|_{H^{2}}^{4} \right) \|\varphi\|_{H^{1}}^{2} + C\delta_{1} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2}, \tag{5.13}$$

with $\delta_1 > 0$ to be chosen later. Recalling (5.6) and (5.8), we obtain with similar arguments that

$$\left| \int_{\Omega} (\psi'(\varphi_{2}) - \psi'(\varphi_{1})) \nabla \varphi_{1} \cdot \mathbf{u} \, \mathrm{d}x \right| \leq C \|\psi'(\varphi_{1}) - \psi'(\varphi_{2})\|_{L^{2}} \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}} \|\mathbf{u}\|_{\mathbf{L}^{6}}$$
$$\leq C \left(1 + \|\varphi_{1}\|_{L^{12}}^{4} + \|\varphi_{2}\|_{L^{12}}^{4} \right) \|\varphi\|_{H^{1}} \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}} \|\mathbf{u}\|_{\mathbf{H}^{1}}$$
$$\leq C \left(1 + \|\varphi_{1}\|_{H^{2}}^{4} + \|\varphi_{2}\|_{H^{2}}^{4} + \|\sigma_{2}\|_{L^{6}}^{4} \right) \|\varphi\|_{H^{2}}^{2} + C \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2}.$$
(5.14)

Again using (5.6) and (5.8) and the continuous embeddings $H^1 \hookrightarrow L^6$, $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$, we obtain

$$\left| \int_{\Omega} (\mu_{2} + \chi \sigma_{2}) \nabla \varphi \cdot (\mathbf{v} - \mathbf{u}) \, \mathrm{d}x \right| \leq C \left(1 + \|\mu_{2}\|_{H^{1}}^{2} + \|\sigma_{2}\|_{H^{1}}^{2} \right) \|\varphi\|_{H^{1}}^{2} + C \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} + \delta_{2} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2}, \quad (5.15)$$

with $\delta_2 > 0$ to be chosen later. Once more using (2.8),(5.6) and (5.8) and the continuous embedding $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$, we deduce that

$$\begin{aligned} \left| \int_{\Omega} \Delta \varphi \nabla \varphi_{1} \cdot \mathbf{u} \, \mathrm{d}x \right| &\leq \|\Delta \varphi\|_{L^{2}} \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}} \|\mathbf{u}\|_{\mathbf{L}^{6}} \\ &\leq C \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}}^{2} \|\mathbf{u}\|_{\mathbf{H}^{1}}^{2} + \frac{1}{8} \|\Delta \varphi\|_{L^{2}}^{2} \\ &\leq C \left(1 + \|\varphi_{1}\|_{H^{2}}^{2} + \|\sigma_{2}\|_{H^{1}}^{4} \right) \|\varphi\|_{H^{1}}^{2} + C \|\varphi_{1}\|_{H^{2}} \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} + \frac{1}{8} \|\Delta \varphi\|_{L^{2}}^{2}. \end{aligned}$$
(5.16)

Using the assumptions on $\eta(\cdot)$, we obtain

$$\left| \int_{\Omega} 2\eta(\varphi_{1}) \mathbf{D} \mathbf{v} \colon \nabla \mathbf{u} + \nu \mathbf{v} \cdot \mathbf{u} \, \mathrm{d} x \right| \leq \frac{\eta_{0}}{2} \| \mathbf{D} \mathbf{v} \|_{\mathbf{L}^{2}}^{2} + \frac{\nu}{2} \| \mathbf{v} \|_{\mathbf{L}^{2}}^{2} + C \| \mathbf{u} \|_{\mathbf{H}^{1}}^{2}$$
$$\leq \frac{\eta_{0}}{2} \| \mathbf{D} \mathbf{v} \|_{\mathbf{L}^{2}}^{2} + \frac{\nu}{2} \| \mathbf{v} \|_{\mathbf{L}^{2}}^{2} + C \left(1 + \| \sigma_{2} \|_{H^{1}}^{2} \right) \| \varphi \|_{H^{1}}^{2}$$
$$+ C \| \sigma_{\infty} \|_{L^{2}(\partial\Omega)}^{2}. \tag{5.17}$$

Furthermore, using the boundedness of $\eta'(\cdot)$, elliptic regularity and (1.10), (5.6), (5.8), we obtain

$$\left| \int_{\Omega} 2(\eta(\varphi_{1}) - \eta(\varphi_{2})\mathbf{D}\mathbf{v}_{2} \colon \nabla(\mathbf{v} - \mathbf{u}) \, \mathrm{d}x \right| \\
\leq C \|\varphi\|_{L^{\infty}} \|\mathbf{D}\mathbf{v}_{2}\|_{\mathbf{L}^{2}} \left(\|\nabla\mathbf{v}\|_{\mathbf{L}^{2}} + \|\nabla\mathbf{u}\|_{\mathbf{L}^{2}} \right) \\
\leq C \|\varphi\|_{H^{1}}^{\frac{3}{4}} \|\varphi\|_{H^{3}}^{\frac{1}{4}} \|\mathbf{D}\mathbf{v}_{2}\|_{\mathbf{L}^{2}} \left(\|\nabla\mathbf{v}\|_{\mathbf{L}^{2}} + \|\nabla\mathbf{u}\|_{\mathbf{L}^{2}} \right) \\
\leq \delta_{3} \|\nabla\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + C \|\nabla\mathbf{u}\|_{\mathbf{L}^{2}}^{2} + C \|\varphi\|_{H^{1}}^{\frac{3}{2}} \|\varphi\|_{H^{3}}^{\frac{1}{2}} \|\mathbf{D}\mathbf{v}_{2}\|_{\mathbf{L}^{2}}^{2} \\
\leq \delta_{3} \|\nabla\mathbf{v}\|_{\mathbf{L}^{2}}^{2} + C \|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2} + \frac{1}{4} \|\nabla\Delta\varphi\|_{\mathbf{L}^{2}}^{2} + C \left(1 + \|\mathbf{D}\mathbf{v}_{2}\|_{\mathbf{L}^{2}}^{\frac{8}{3}} + \|\sigma_{2}\|_{H^{1}}^{2}\right) \|\varphi\|_{H^{1}}^{2},$$
(5.18)

with $\delta_3 > 0$ to be chosen later. Due to (5.8) and since $\varphi_2 \in L^{\infty}(H^1)$ with bounded norm, we get

$$\left| \int_{\Omega} (1 - \varphi_2) (\Gamma(\varphi_1, \sigma_1) - \Gamma(\varphi_2, \sigma_2)) \varphi \, \mathrm{d}x \right| \le C \left(1 + \|\sigma_2\|_{L^6} \right) \|\varphi\|_{H^1}^2 + C \|\sigma_\infty\|_{L^2(\partial\Omega)}^2.$$
(5.19)

Applying (2.8) and the continuous embeddings $H^1 \hookrightarrow L^6$, $\mathbf{H}^1 \hookrightarrow \mathbf{L}^6$ gives

.

$$\left| \int_{\Omega} (\nabla \varphi_{1} \cdot \mathbf{v} + \nabla \varphi \cdot \mathbf{v}_{2}) \varphi \, \mathrm{d}x \right| \leq \|\nabla \varphi_{1}\|_{\mathbf{L}^{3}} \|\mathbf{v}\|_{\mathbf{L}^{6}} \|\varphi\|_{L^{2}} + \|\nabla \varphi\|_{\mathbf{L}^{2}} \|\mathbf{v}_{2}\|_{\mathbf{L}^{6}} \|\varphi\|_{L^{3}}$$
$$\leq C \left(\|\nabla \varphi_{1}\|_{\mathbf{L}^{3}} \|\mathbf{v}\|_{\mathbf{H}^{1}} \|\varphi\|_{H^{1}} + \|\varphi\|_{H^{1}}^{2} \|\mathbf{v}_{2}\|_{\mathbf{H}^{1}} \right)$$
$$\leq C \left(\|\varphi_{1}\|_{H^{2}} + \|\mathbf{v}_{2}\|_{\mathbf{H}^{1}} \right) \|\varphi\|_{H^{1}}^{2} + \delta_{4} \|\mathbf{v}\|_{\mathbf{H}^{1}}^{2}, \tag{5.20}$$

with $\delta_4 > 0$ to be chosen later. For the last term on the r.h.s. of (5.7), we use (A4) and the continuous embedding $H^1 \hookrightarrow L^6$ to obtain

$$\left| \int_{\Omega} \Gamma_{\mathbf{v},1} |\varphi|^2 \, \mathrm{d}x \right| \le C \left(1 + \|\sigma_2\|_{L^6} \right) \|\varphi\|_{H^1}^2.$$
(5.21)

We now estimate the first term on the r.h.s. of (5.7). To this end, we first observe that

$$\nabla(\psi'(\varphi_1) - \psi'(\varphi_2)) = \psi''(\varphi_1)\nabla\varphi_1 - \psi''(\varphi_2)\nabla\varphi_2 = \psi''(\varphi_1)\nabla\varphi + \nabla\varphi_2(\psi''(\varphi_1) - \psi''(\varphi_2)).$$

Due to (2.3)-(2.4) and (2.8), we obtain

. . .

$$\int_{\Omega} |\psi''(\varphi_1)\nabla\varphi|^2 \,\mathrm{d}x \le C \int_{\Omega} (1+|\varphi_1|^8) |\nabla\varphi|^2 \,\mathrm{d}x \le C \left(1+\|\varphi_1\|_{L^{\infty}}^8\right) \|\nabla\varphi\|_{\mathbf{L}^2}^2 \le C \left(1+\|\varphi_1\|_{H^3}^2\right) \|\varphi\|_{H^1}^2.$$

Applying (2.8) and (2.15), we conclude that

$$\begin{split} \int_{\Omega} |\nabla \varphi_2(\psi''(\varphi_1) - \psi''(\varphi_2))|^2 \, \mathrm{d}x &\leq C \int_{\Omega} (1 + |\varphi_1|^6 + |\varphi_2|^6) |\nabla \varphi_2|^2 |\varphi|^2 \, \mathrm{d}x \\ &\leq C \left(1 + \|\varphi_1\|_{L^{18}}^6 + \|\varphi_2\|_{L^{16}}^6 \right) \|\nabla \varphi_2\|_{\mathbf{L}^6}^2 \|\varphi\|_{L^6}^2 \\ &\leq C \left(1 + \|\varphi_1\|_{H^2}^4 + \|\varphi_2\|_{H^2}^4 \right) \|\varphi\|_{H^1}^2. \end{split}$$

Combining the last two estimates, we obtain

$$\left\|\nabla(\psi'(\varphi_1) - \psi'(\varphi_2))\right\|_{\mathbf{L}^2}^2 \le C\left(1 + \|\varphi_1\|_{H^2}^4 + \|\varphi_2\|_{H^2}^4 + \|\varphi_1\|_{H^3}^2\right) \|\varphi\|_{H^1}^2.$$
(5.22)

From this inequality, we deduce that

$$\left| \int_{\Omega} \nabla(\psi'(\varphi_{1}) - \psi'(\varphi_{2})) \cdot (\nabla \Delta \varphi - \nabla \varphi) \, \mathrm{d}x \right| \leq C \left(1 + \|\varphi_{1}\|_{H^{2}}^{4} + \|\varphi_{2}\|_{H^{2}}^{4} + \|\varphi_{1}\|_{H^{3}}^{2} \right) \|\varphi\|_{H^{1}}^{2} + \frac{1}{8} \|\nabla \Delta \varphi\|_{\mathbf{L}^{2}}^{2}.$$
(5.23)

Finally, using (5.5), it is easy to check that

$$\left| \int_{\Omega} \chi \nabla \sigma \cdot (\nabla \varphi - \nabla \Delta \varphi) \mathrm{d}x \right| \le C \left(1 + \|\sigma_2\|_{L^6}^2 \right) \|\varphi\|_{H^1}^2 + C \|\sigma_\infty\|_{L^2(\partial\Omega)}^2 + \frac{1}{8} \|\nabla \Delta \varphi\|_{\mathbf{L}^2}^2.$$
(5.24)

Using (5.9)-(5.24) in (5.7) and choosing

$$\delta_1 = \delta_2 = \delta_3 = \delta_4 = \frac{\min\{\frac{\nu}{2}, \eta\}}{8C_K^2} \eqqcolon C_1,$$

where C_K is Korn's constant, we end up with

$$\frac{\mathrm{d}}{\mathrm{d}t}\frac{1}{2}\left(\|\nabla\varphi\|_{\mathbf{L}^{2}}^{2}+\|\varphi\|_{L^{2}}^{2}\right)+4C_{1}\|\mathbf{v}\|_{\mathbf{H}^{1}}^{2}+\frac{1}{2}\int_{\Omega}|\Delta\varphi|^{2}+|\nabla\Delta\varphi|^{2}\,\mathrm{d}x\leq\alpha_{1}(t)\|\varphi\|_{H^{1}}^{2}+\alpha_{2}(t)\|\sigma_{\infty}\|_{L^{2}(\partial\Omega)}^{2},\tag{5.25}$$

where

$$\begin{aligned} \alpha_1(t) &\coloneqq C \left(1 + \|\varphi_1\|_{H^2}^4 + \|\varphi_2\|_{H^2}^4 + \|\varphi_1\|_{H^3}^2 + \|\varphi_2\|_{H^2}^2 + \|\mathbf{v}_2\|_{\mathbf{H}^1}^{\frac{8}{3}} + \|\mu_2\|_{H^1}^2 + \|\sigma_1\|_{H^1}^2 + \|\sigma_2\|_{H^1}^4 \right) \\ \alpha_2(t) &\coloneqq C \left(1 + \|\varphi_1\|_{H^2} + \|\varphi_2\|_{H^2} \right). \end{aligned}$$

Due to (2.8)-(2.9), it follows that $\alpha_1 \in L^1(0,T)$, $\alpha_2 \in L^4(0,T)$. We remark that $\alpha_1 \in L^1(0,T)$ only holds provided $\sigma_2 \in L^4(H^1)$ with bounded norm. Then, using a Gronwall argument (see [25, Lemma 3.1]) in (5.25) yields

$$\sup_{s \in (0,T]} \|\varphi(s)\|_{H^1}^2 + \int_0^T \|\mathbf{v}\|_{\mathbf{H}^1}^2 \mathrm{d}s + \int_0^T \int_{\Omega} |\Delta\varphi|^2 + |\nabla\Delta\varphi|^2 \,\mathrm{d}x \mathrm{d}s \le C \left(\|\varphi_0\|_{H^1}^2 + \|\sigma_\infty\|_{L^4(L^2(\partial\Omega))}^2 \right).$$

Together with elliptic regularity theory, this gives

$$\|\varphi\|_{L^{\infty}(H^{1})\cap L^{2}(H^{3})} + \|\mathbf{v}\|_{L^{2}(\mathbf{H}^{1})} \leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right).$$
(5.26)

Now, from (5.5) and (5.26), we immediately obtain

$$\|\sigma\|_{L^{2}(H^{1})} \leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right).$$
(5.27)

Using (1.10), (5.12), (5.22), (5.26) and the boundedness of $\varphi_1, \varphi_2 \in L^{\infty}(H^1) \cap L^4(H^2) \cap L^2(H^3)$, it is straightforward to check that

$$\|\psi'(\varphi_1) - \psi'(\varphi_2)\|_{L^2(H^1)} \le C\left(\|\varphi_0\|_{H^1} + \|\sigma_\infty\|_{L^4(L^2(\partial\Omega))}\right).$$

Together with (5.26)-(5.27), a comparison argument in (5.2) yields

$$\|\mu\|_{L^{2}(H^{1})} \leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right).$$
(5.28)

Using (5.26)-(5.28), a comparison argument in (5.1b) yields

$$\|\varphi\|_{H^{1}((H^{1})^{*})} \leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right).$$
(5.29)

Combining (5.26)-(5.29), we deduce the estimate

$$\begin{aligned} \|\varphi\|_{H^{1}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{2}(H^{3})} + \|\mu\|_{L^{2}(H^{1})} + \|\sigma\|_{L^{2}(H^{1})} + \|\mathbf{v}\|_{L^{2}(\mathbf{H}^{1})} \\ &\leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right). \end{aligned}$$
(5.30)

Step 4: It remains to get an estimate on the pressure. To this end, let $\mathbf{q} \in \mathbf{H}^1$ be a solution of

$$\operatorname{div}(\mathbf{q}) = p \quad \text{in } \Omega, \quad \mathbf{q} = \frac{1}{|\partial \Omega|} \left(\int_{\Omega} p \, \mathrm{d}x \right) \mathbf{n} \quad \text{on } \partial \Omega,$$

such that

$$\|\mathbf{q}\|_{\mathbf{H}^{1}} \le c \|p\|_{L^{2}},\tag{5.31}$$

with c depending only on Ω . Then, choosing $\mathbf{\Phi} = \mathbf{q}$ in (5.1a), we obtain

$$\|p\|_{L^{2}}^{2} = \int_{\Omega} (2\eta(\varphi_{1})\mathbf{D}\mathbf{v} + \lambda(\varphi_{1})\operatorname{div}(\mathbf{v})\mathbf{I}) \colon \nabla\mathbf{q} + \nu\mathbf{v} \cdot \mathbf{q} \, \mathrm{d}x - \int_{\Omega} (\mu + \chi\sigma)\nabla\varphi_{1} \cdot \mathbf{q} + (\mu_{2} + \chi\sigma_{2})\nabla\varphi \cdot \mathbf{q} \, \mathrm{d}x + \int_{\Omega} (2(\eta(\varphi_{1}) - \eta(\varphi_{2}))\mathbf{D}\mathbf{v}_{2} + (\lambda(\varphi_{1}) - \lambda(\varphi_{2}))\operatorname{div}(\mathbf{v}_{2})\mathbf{I}) \colon \nabla\mathbf{q} \, \mathrm{d}x.$$
(5.32)

Using (2.8)-(2.9) and (A3), a straightforward calculation shows that

$$\begin{aligned} \left| \int_{\Omega} (2\eta(\varphi_1) \mathbf{D} \mathbf{v} + \lambda(\varphi_1) \operatorname{div}(\mathbf{v}) \mathbf{I}) \colon \nabla \mathbf{q} + \nu \mathbf{v} \cdot \mathbf{q} \, \mathrm{d}x - \int_{\Omega} (\mu + \chi \sigma) \nabla \varphi_1 \cdot \mathbf{q} + (\mu_2 + \chi \sigma_2) \nabla \varphi \cdot \mathbf{q} \, \mathrm{d}x \right| \\ & \leq C \left(\|\mathbf{v}\|_{\mathbf{H}^1}^2 + \|\mu + \chi \sigma\|_{H^1}^2 + \|\mu_2 + \chi \sigma_2\|_{H^1}^2 \|\varphi\|_{H^1}^2 \right) + \frac{1}{4} \|p\|_{L^2}^2. \end{aligned}$$

For the remaining terms, we use again (2.8)-(2.9) and (A3) to obtain

$$\left| \int_{\Omega} (2(\eta(\varphi_1) - \eta(\varphi_2)) \mathbf{D} \mathbf{v}_2 + (\lambda(\varphi_1) - \lambda(\varphi_2)) \operatorname{div}(\mathbf{v}_2) \mathbf{I}) \colon \nabla \mathbf{q} \, \mathrm{d}x \right| \le C \|\mathbf{v}_2\|_{\mathbf{H}^1}^2 \|\varphi\|_{L^{\infty}}^2 + \frac{1}{4} \|p\|_{L^2}^2$$

Using the last two inequalities in (5.32), integrating the resulting estimate in time from 0 to T and using Young's generalised inequality, we deduce that

$$\|p\|_{L^{2}(L^{2})}^{2} \leq C\left(\|\mathbf{v}\|_{L^{2}(\mathbf{H}^{1})}^{2} + \|\mu + \chi\sigma\|_{L^{2}(H^{1})}^{2} + \|\mu_{2} + \chi\sigma_{2}\|_{L^{2}(H^{1})}^{2} \|\varphi\|_{L^{\infty}(H^{1})}^{2} + \|\mathbf{v}_{2}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})}^{2} \|\varphi\|_{L^{8}(L^{\infty})}^{2}\right).$$

Therefore, using (2.8)-(2.9), (5.30) and the continuous embedding $L^8(L^{\infty}) \hookrightarrow L^{\infty}(H^1) \cap L^2(H^3)$, the last inequality implies

$$\|p\|_{L^{2}(L^{2})} \leq C\left(\|\varphi_{0}\|_{H^{1}} + \|\sigma_{\infty}\|_{L^{4}(L^{2}(\partial\Omega))}\right).$$
(5.33)

Together with (5.30), we obtain (2.16), whence the proof is complete.

6 Existence of strong solutions

In this section, we will prove Theorem 2.12. The testing procedure can again be justified by a Galerkin scheme. In the following, we assume for simplicity and as it has no further consequence for the analysis that $\epsilon = 1$. Then, with similar arguments as before, we can arrive at

$$\begin{aligned} \|\varphi\|_{H^{1}((H^{1})^{*})\cap L^{\infty}(H^{1})\cap L^{4}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{L^{4}(H^{1})} + \|\mu\|_{L^{2}(H^{1})\cap L^{4}(L^{2})} \\ &+ \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{2})} + \|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})} + \|p\|_{L^{2}(L^{2})} \leq C. \end{aligned}$$

$$(6.1)$$

We will now show the result in a series of higher order estimates.

Step 1: Observing that (2.7d) is the weak formulation of

$$\begin{aligned} -\Delta \sigma + h(\varphi)\sigma &= 0 & \text{a. e. in } \Omega, \\ \partial_{\mathbf{n}}\sigma + K\sigma &= K\sigma_{\infty} & \text{a. e. on } \partial\Omega, \end{aligned}$$

due to the assumptions on $h(\cdot)$ and using [29, Thm. 2.4.2.6] we deduce that

$$\|\sigma\|_{H^2} \le C \|K\sigma_\infty\|_{H^{\frac{1}{2}}(\partial\Omega)}.$$

Therefore, using the continuous embedding $H^2 \hookrightarrow L^{\infty}$ and the fact that $\sigma_{\infty} \in H^1(H^{\frac{1}{2}}(\partial\Omega)) \hookrightarrow C^0(H^{\frac{1}{2}}(\partial\Omega))$, we have

$$\|\sigma\|_{L^{\infty}(H^2)\cap L^{\infty}(L^{\infty})} \le C. \tag{6.2}$$

By (A3), this yields

$$\|\operatorname{div}(\mathbf{v})\|_{L^{\infty}(L^{\infty})} + \|\Gamma_{\varphi}\|_{L^{\infty}(L^{\infty})} \le C.$$
(6.3)

Now, for h > 0 we introduce the incremental ratio

$$\partial_t^h u(t) = \frac{1}{h} (u(t+h) - u(t)).$$

Then, using (2.7d), we see that

$$0 = \int_{\Omega} \nabla \partial_t^h \sigma(t) \cdot \nabla \Phi + \left(\partial_t^h (h(\varphi(t))) \sigma(t+h) + \partial_t^h \sigma(t) h(\varphi(t)) \right) \Phi \, \mathrm{d}x + \int_{\partial \Omega} K(\partial_t^h \sigma(t) - \partial_t^h \sigma_\infty(t)) \Phi \mathrm{d}\mathcal{H}^{d-1}$$

holding for almost every $t \in (0, T - h]$. Testing this equation with $\Phi = \partial_t^h \sigma(t)$, integrating in time from 0 to T - h and using the non-negativity of $h(\cdot)$, we obtain

$$\int_{0}^{T-h} \|\nabla \partial_{t}^{h} \sigma(t)\|_{\mathbf{L}^{2}}^{2} dt + K \int_{0}^{T-h} \|\partial_{t}^{h} \sigma(t)\|_{L^{2}(\partial\Omega)}^{2} dt \leq \int_{0}^{T-h} \int_{\Omega} \partial_{t}^{h} (h(\varphi(t))) \sigma(t+h) \partial_{t}^{h} \sigma(t) dx dt + \int_{0}^{T-h} \int_{\partial\Omega} K \partial_{t}^{h} \sigma(t) \partial_{t}^{h} \sigma_{\infty}(t) d\mathcal{H}^{d-1} dt. \quad (6.4)$$

To estimate the r.h.s. of this equation, we use (6.2) and the Lipschitz-continuity of $h(\cdot)$ to obtain

$$\left| \int_{0}^{T-h} \int_{\Omega} \partial_{t}^{h}(h(\varphi(t)))\sigma(t+h)\partial_{t}^{h}\sigma(t)\mathrm{d}x\mathrm{d}t \right| \leq C \int_{0}^{T-h} \int_{\Omega} |\partial_{t}^{h}\varphi(t)\partial_{t}^{h}\sigma(t)|\mathrm{d}x\mathrm{d}t$$
$$\leq C \|\partial_{t}^{h}\varphi\|_{L^{2}(0,T-h;(H^{1})^{*})} \|\partial_{t}^{h}\sigma\|_{L^{2}(0,T-h;H^{1})}$$
$$\leq C \|\partial_{t}\varphi\|_{L^{2}(0,T;(H^{1})^{*})} \|\partial_{t}^{h}\sigma\|_{L^{2}(0,T-h;H^{1})}. \tag{6.5}$$

With similar arguments and using the trace theorem, the remaining term can be estimated by

$$\left| \int_{0}^{T-h} \int_{\partial\Omega} K \partial_t^h \sigma(t) \partial_t^h \sigma_\infty(t) \mathrm{d}\mathcal{H}^{d-1} \mathrm{d}t \right| \le C \|\partial_t \sigma_\infty\|_{L^2(0,T;H^{\frac{1}{2}}(\partial\Omega))} \|\partial_t^h \sigma\|_{L^2(0,T-h;H^1)}.$$
(6.6)

Using (6.4)-(6.6) together with (6.1) and (C2), an application of Poincaré's inequality implies

$$\|\partial_t^h \sigma\|_{L^2(0,T-h;H^1)} \le C \left(\|\partial_t \varphi\|_{L^2(0,T;(H^1)^*)} + \|\partial_t \sigma_\infty\|_{L^2(0,T;H^{\frac{1}{2}})} \right) \le C.$$
(6.7)

Since the constant C is independent of h > 0, this yields

$$\|\partial_t \sigma\|_{L^2(H^1)} \le C.$$

Combining this inequality with (6.2) and using the continuous embedding $H^1(H^1) \hookrightarrow C^0(H^1)$, we obtain

$$\|\sigma\|_{H^1(H^1)\cap C^0(H^1)\cap L^\infty(H^2)} \le C.$$
(6.8)

Step 2: Choosing $\Phi = \partial_t \varphi$ in (2.7b) and $\Phi = \Delta \partial_t \varphi$ in (2.7c) and integrating by parts, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}t}\frac{1}{2}\int_{\Omega}|\Delta\varphi|^{2}\,\mathrm{d}x + \int_{\Omega}|\partial_{t}\varphi|^{2} = -\int_{\Omega}(\mathrm{div}(\varphi\mathbf{v}) - \Gamma_{\varphi})\partial_{t}\varphi\,\mathrm{d}x + \chi\int_{\Omega}\nabla\sigma\cdot\nabla\partial_{t}\varphi\,\mathrm{d}x + \int_{\Omega}\psi'''(\varphi)|\nabla\varphi|^{2}\partial_{t}\varphi + \psi''(\varphi)\Delta\varphi\partial_{t}\varphi\,\mathrm{d}x.$$
(6.9)

We recall that Γ_{φ} , $\Gamma_{\mathbf{v}} \in L^2(L^2)$ with bounded norm. Then, using Hölder's and Young's inequalities, we can estimate the first three terms on the r.h.s. of (6.9) by

$$\left| \int_{\Omega} (\operatorname{div}(\varphi \mathbf{v}) - \Gamma_{\varphi}) \partial_t \varphi \, \mathrm{d}x \right| \le C \left(\left\| \operatorname{div}(\varphi \mathbf{v}) \right\|_{L^2}^2 + \left\| \Gamma_{\varphi} \right\|_{L^2}^2 \right) + \frac{1}{4} \left\| \partial_t \varphi \right\|_{L^2}^2.$$
(6.10)

For the last term on the r.h.s. of (6.9), we use Hölder's and Young's inequalities together with (1.10), (2.3)-(2.4) and (6.1) to obtain

$$\left| \int_{\Omega} \psi''(\varphi) \Delta \varphi \partial_t \varphi \, \mathrm{d}x \right| \le C \left(1 + \|\varphi\|_{L^{\infty}}^4 \right) \|\Delta \varphi\|_{L^2} \|\partial_t \varphi\|_{L^2} \le \frac{1}{4} \|\partial_t \varphi\|_{L^2}^2 + C \left(1 + \|\varphi\|_{H^3}^2 \right) \|\Delta \varphi\|_{L^2}^2.$$
(6.11)

Now, using (C3), Hölder's and Young's inequalities, (1.7), (1.10) and (6.1), we obtain

$$\left| \int_{\Omega} \psi'''(\varphi) |\nabla \varphi|^{2} \partial_{t} \varphi \, \mathrm{d}x \right| \leq C \left(1 + \|\varphi\|_{L^{\infty}}^{3} \right) \|\nabla \varphi\|_{L^{4}}^{2} \|\partial_{t} \varphi\|_{L^{2}}$$

$$\leq C \left(1 + \|\varphi\|_{H^{3}}^{\frac{3}{4}} \right) \|\varphi\|_{H^{2}}^{\frac{3}{2}} \|\partial_{t} \varphi\|_{L^{2}}$$

$$\leq C \left(1 + \|\varphi\|_{H^{3}} \right) (\|\varphi\|_{L^{2}} + \|\Delta \varphi\|_{L^{2}}) \|\partial_{t} \varphi\|_{L^{2}}$$

$$\leq C \left(1 + \|\varphi\|_{H^{3}}^{2} \right) \left(1 + \|\Delta \varphi\|_{L^{2}}^{2} \right) + \frac{1}{4} \|\partial_{t} \varphi\|_{L^{2}}^{2}. \tag{6.12}$$

For the remaining term on the r.h.s. of (6.9), we observe that

$$\chi \int_{\Omega} \nabla \sigma \cdot \nabla \partial_t \varphi dx = \frac{\mathrm{d}}{\mathrm{d}t} \chi \int_{\Omega} \nabla \sigma \cdot \nabla \varphi dx - \int_{\Omega} \nabla \partial_t \sigma \cdot \nabla \varphi \quad \text{for a. e. } t \in (0, T).$$

Collecting (6.9)-(6.11), (6.12) and using the last identity, we end up with

$$\frac{\mathrm{d}}{\mathrm{d}t}\frac{1}{2}\int_{\Omega}|\Delta\varphi|^{2}\,\mathrm{d}x + \frac{1}{4}\int_{\Omega}|\partial_{t}\varphi|^{2} \leq C\left(1 + \|\mathrm{div}(\varphi\mathbf{v})\|_{L^{2}}^{2} + \|\Gamma_{\varphi}\|_{L^{2}}^{2} + \|\varphi\|_{H^{3}}^{2}\right) + C\left(1 + \|\varphi\|_{H^{3}}^{2}\right)\|\Delta\varphi\|_{L^{2}}^{2} + \frac{\mathrm{d}}{\mathrm{d}t}\chi\int_{\Omega}\nabla\sigma\cdot\nabla\varphi\mathrm{d}x - \int_{\Omega}\nabla\partial_{t}\sigma\cdot\nabla\varphi\mathrm{d}x.$$

Integrating this inequality in time from 0 to $s \in (0, T]$, we obtain

$$\frac{1}{2} \|\Delta\varphi(s)\|_{L^{2}}^{2} + \frac{1}{4} \|\partial_{t}\varphi\|_{L^{2}(0,s;L^{2})}^{2} \leq \|\Delta\varphi_{0}\|_{L^{2}}^{2} + \int_{0}^{s} \alpha_{1}(t) + \alpha_{2}(t) \|\Delta\varphi(t)\|_{L^{2}}^{2} dt
+ \chi \int_{\Omega} \nabla\sigma(s) \cdot \nabla\varphi(s) dx - \chi \int_{\Omega} \nabla\sigma(0) \cdot \nabla\varphi_{0} dx
- \chi \int_{0}^{s} \int_{\Omega} \nabla\partial_{t}\sigma \cdot \nabla\varphi dx dt,$$
(6.13)

where

 $\alpha_1(t) \coloneqq C\left(1 + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^2}^2 + \|\Gamma_{\varphi}\|_{L^2}^2 + \|\varphi\|_{H^3}^2\right), \qquad \alpha_2(t) \coloneqq C\left(1 + \|\varphi\|_{H^3}^2\right).$

Now, using (6.1), (6.8) and $\varphi_0 \in H^2_N$, we obtain

$$\begin{aligned} \left| \chi \int_{\Omega} \nabla \sigma(0) \cdot \nabla \varphi_0 \mathrm{d}x \right| &= \left| \chi \int_{\Omega} \sigma(0) \Delta \varphi_0 \mathrm{d}x \right| \le C \left(1 + \left\| \Delta \varphi_0 \right\|_{L^2}^2 \right), \\ \left| \chi \int_{\Omega} \nabla \sigma(s) \cdot \nabla \varphi(s) \mathrm{d}x \right| \le \left\| \sigma \right\|_{C^0(H^1)} \sup_{s \in (0,T]} \left\| \nabla \varphi(s) \right\| \le C, \\ \left| \chi \int_0^s \int_{\Omega} \nabla \partial_t \sigma \cdot \nabla \varphi \mathrm{d}x \mathrm{d}t \right| \le C \int_0^s \left\| \partial_t \sigma(t) \right\|_{H^1} \mathrm{d}t. \end{aligned}$$

Together with (6.13), this implies

$$\frac{1}{2} \|\Delta\varphi(s)\|_{L^2}^2 + \frac{1}{4} \|\partial_t\varphi\|_{L^2(0,s;L^2)}^2 \le C\left(1 + \|\Delta\varphi_0\|_{L^2}^2\right) + \int_0^s \beta_1(t) + \beta_2(t) \|\Delta\varphi(t)\|_{L^2}^2 \mathrm{d}t, \tag{6.14}$$

where

$$\beta_1(t) \coloneqq C\left(1 + \|\partial_t \sigma\|_{H^1} + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^2}^2 + \|\Gamma_\varphi\|_{L^2}^2 + \|\varphi\|_{H^3}^2\right), \qquad \beta_2(t) \coloneqq C\left(1 + \|\varphi\|_{H^3}^2\right).$$

Due to (6.1), (6.3) and (6.8), we know that $\beta_1, \beta_2 \in L^1(0,T)$. Together with the assumption $\varphi_0 \in H^2_N$, an application of Gronwall's lemma in (6.14) yields

$$\|\Delta\varphi\|_{L^{\infty}(L^{2})} + \|\partial_{t}\varphi\|_{L^{2}(L^{2})} \le C.$$
(6.15)

Step 3: Combining (6.1) and (6.15), from elliptic regularity theory we obtain

$$\begin{aligned} \|\varphi\|_{H^{1}(L^{2})\cap L^{\infty}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{H^{1}(H^{1})\cap C^{0}(H^{1})\cap L^{\infty}(H^{2})} + \|\mu\|_{L^{2}(H^{1})\cap L^{4}(L^{2})} \\ + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{2})} + \|\mathbf{v}\|_{L^{\frac{8}{3}}(\mathbf{H}^{1})} + \|p\|_{L^{2}(L^{2})} \leq C. \end{aligned}$$

$$(6.16)$$

Using (6.16) and applying elliptic regularity theory in (2.7b), we obtain

$$\|\mu\|_{L^2(H^2)} \le C. \tag{6.17}$$

Using a comparison argument in (2.7c) and (6.16)-(6.17), it follows that

$$\|\mu\|_{L^{\infty}(L^2)} \le C. \tag{6.18}$$

Step 4: We now want to apply Lemma 1.3 with q = 2. Using the generalised chain rule for Sobolev functions, (6.16) and the assumptions on $\Gamma_{\mathbf{v}}$, it is straightforward to check that

$$\|\Gamma_{\mathbf{v}}\|_{L^{\infty}(H^1)} \le C. \tag{6.19}$$

Furthermore, since $\nabla \varphi \in L^4(\mathbf{L}^{\infty})$, $\mu + \chi \sigma \in L^{\infty}(L^2)$ with bounded norm, we observe that

$$\|(\mu + \chi \sigma) \nabla \varphi\|_{L^4(\mathbf{L}^2)} \le C. \tag{6.20}$$

Hence, using the assumptions on $\lambda(\cdot)$, $\eta(\cdot)$, an application of (1.14) yields

$$\|\mathbf{v}\|_{\mathbf{H}^{2}} + \|p\|_{H^{1}} \le C(\eta_{0}, \eta_{1}, \lambda_{0}, \|\varphi\|_{W^{1,4}}) (\|(\mu + \chi\sigma)\nabla\varphi\|_{\mathbf{L}^{2}} + \|\Gamma_{\mathbf{v}}\|_{H^{1}})$$

Integrating this inequality in time from 0 to T, using (6.19)-(6.20) and recalling $\varphi \in L^{\infty}(W^{1,4})$ (since $H^2 \hookrightarrow W^{1,4}$), we obtain

$$\|\mathbf{v}\|_{L^4(\mathbf{H}^2)} + \|p\|_{L^4(H^1)} \le C.$$
(6.21)

Step 5: Finally, due to the compact embedding $H^2 \hookrightarrow C^0(\overline{\Omega})$ and because of (6.16), we obtain

$$\|\varphi\|_{C^0(\bar{Q})} \le C.$$

For completeness, we summarize all the estimates we deduced in this Section, given by

$$\begin{aligned} \|\varphi\|_{H^{1}(L^{2})\cap C^{0}(\bar{Q})\cap L^{\infty}(H^{2})\cap L^{2}(H^{3})} + \|\sigma\|_{H^{1}(H^{1})\cap C^{0}(H^{1})\cap L^{\infty}(H^{2})} + \|\mu\|_{L^{\infty}(L^{2})\cap L^{2}(H^{2})} \\ + \|\operatorname{div}(\varphi \mathbf{v})\|_{L^{2}(L^{2})} + \|\mathbf{v}\|_{L^{4}(\mathbf{H}^{2})} + \|p\|_{L^{4}(H^{1})} \leq C. \end{aligned}$$

$$(6.22)$$

These a-priori-estimates together with a Galerkin-scheme are enough to pass to the limit in the weak formulation to show existence of strong solutions. For the details when passing to the limit, we again refer to [16], [23].

Step 6: Since (1.1d) holds a.e. in Q, we see that φ is a solution of

$$\Delta \varphi = \psi'(\varphi) - \mu - \chi \sigma \quad \text{a.e. in } Q,$$
$$\nabla \varphi \cdot \mathbf{n} = 0 \qquad \qquad \text{a.e. on } \Sigma.$$

Since $\psi'(\varphi) - \mu - \chi \sigma \in L^2(H^2)$ with bounded norm, elliptic regularity theory implies

$$\|\varphi\|_{L^2(H^4)} \le C.$$
 (6.23)

Due to the continuous embedding $L^{\infty}(H^1) \cap L^2(H^3) \hookrightarrow L^8(L^{\infty})$ and by (6.22), this implies $(\mu + \chi \sigma) \nabla \varphi \in L^8(\mathbf{L}^2)$ with bounded norm. Consequently, with the same arguments as used for (6.21), we deduce that

$$\|\mathbf{v}\|_{L^{8}(\mathbf{H}^{2})} + \|p\|_{L^{8}(H^{1})} \le C,$$
(6.24)

which completes the proof.

Acknowledgements

This work was supported by the RTG 2339 "Interfaces, Complex Structures, and Singular Limits" of the German Science Foundation (DFG). The support is gratefully acknowledged.

References

- H. Abels and Y. Terasawa. "On Stokes operators with variable viscosity in bounded and unbounded domains". In: *Math. Ann.* 344.2 (2009), pp. 381–429.
- [2] A. Agosti, C. Cattaneo, C. Giverso, D. Ambrosi, and P. Ciarletta. "A computational framework for the personalized clinical treatment of glioblastoma multiforme". In: ZAMM - Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik (2018).
- G. Allaire. "Homogenization of the Navier-Stokes equations and derivation of Brinkman's law". In: Mathématiques appliquées aux sciences de l'ingénieur (Santiago, 1989). Cépaduès, Toulouse, 1991, pp. 7–20.
- [4] S. Astanin and L. Preziosi. "Multiphase models of tumour growth". In: Selected topics in cancer modeling. Model. Simul. Sci. Eng. Technol. Birkhäuser Boston, Boston, MA, 2008, pp. 223–253.
- [5] E.L. Bearer et al. "Multiparameter computational modeling of tumor invasion." In: *Cancer Research* 69.10 (2009), pp. 4493–4501.
- S. Bosia, M. Conti, and M. Grasselli. "On the Cahn-Hilliard-Brinkman system". In: Commun. Math. Sci. 13.6 (2015), pp. 1541–1567.
- [7] H. C. Brinkman. "A calculation of the viscous force exerted by a flowing fluid on a dense swarm of particles". In: Appli. Sci. Res. A.1 (1947), pp. 27–36.
- [8] H. Byrne and M. Chaplain. "Free boundary value problems associated with the growth and development of multicellular spheroids". In: *Euro. Jnl. of Applied Mathematics* 8 (1997), pp. 639– 658.
- P. G. Ciarlet. *Mathematical elasticity. Vol. I.* Vol. 20. Studies in Mathematics and its Applications. Three-dimensional elasticity. North-Holland Publishing Co., Amsterdam, 1988, pp. xlii+451.
- [10] P. Colli, G. Gilardi, and D. Hilhorst. "On a Cahn-Hilliard type phase field system related to tumor growth". In: Discrete Contin. Dyn. Syst. 35.6 (2015), pp. 2423–2442.
- [11] M. Conti and A. Giorgini. "The three-dimensional Cahn-Hilliard-Brinkman system with unmatched viscosities". preprint. Mar. 2018. URL: https://hal.archives-ouvertes.fr/hal-01559179.
- [12] V. Cristini, J. Lowengrub, and Q. Nie. "Nonlinear simulation of tumor growth". In: J. Math. Biol. 46.3 (2003), pp. 191–224.
- [13] V. Cristini et al. "Morphologic Instability and Cancer Invasion". In: Clin. Cancer Res. 11.19 (2005), pp. 6772–6779.
- [14] M. Dai, E. Feireisl, E. Rocca, J. Schimperna, and M.E. Schonbek. "Analysis of a diffuse interface model of multispecies tumor growth". In: *Nonlinearity* 30.4 (2017), pp. 1639–1658.
- [15] L. Durlofsky and J. F. Brady. "Analysis of the Brinkman equation as a model for flow in porous media". In: *Phys. Fluids* 30.11 (1987), pp. 3329–3341.
- [16] M. Ebenbeck and H. Garcke. "Analysis of a Cahn-Hilliard-Brinkman model for tumour growth with chemotaxis". In: J. Differential Equations (2018). DOI: https://doi.org/10.1016/j.jde.2018.10.045.
- [17] S.J. Franks and J.R. King. "Interactions between a uniformly proliferating tumour and its surroundings: uniform material properties". In: Math. Med. Biol. 20.1 (2003), pp. 47–89.
- [18] H.B. Frieboes et al. "Computer Simulation of Glioma Growth and Morphology". In: NeuroImage 37 Suppl 1 (Feb. 2007), pp. 59–70.
- [19] A. Friedman. "A free boundary problem for a coupled system of elliptic, hyperbolic, and Stokes equations modeling tumor growth". In: *Interfaces Free Bound.* 8.2 (2006), pp. 247–261.

- [20] A. Friedman. "Free boundary problems for systems of Stokes equations". In: Discrete Contin. Dyn. Syst. Ser. B 21.5 (2016), pp. 1455–1468.
- [21] S. Frigeri, M. Grasselli, and E. Rocca. "On a diffuse interface model of tumour growth". In: European J. Appl. Math. 26.2 (2015), pp. 215–243.
- [22] G. P. Galdi. An introduction to the mathematical theory of the Navier-Stokes equations. Second. Springer Monographs in Mathematics. Steady-state problems. Springer, New York, 2011, pp. xiv+1018.
- [23] H. Garcke and K. F. Lam. "Global weak solutions and asymptotic limits of a Cahn-Hilliard-Darcy system modelling tumour growth". In: AIMS Mathematics 1(3) (2016), pp. 318–360.
- [24] H. Garcke and K. F. Lam. "On a Cahn-Hilliard-Darcy system for tumour growth with solution dependent source terms". In: *Trends in applications of mathematics to mechanics*. Vol. 27. Springer INdAM Ser. Springer, Cham, 2018, pp. 243–264.
- [25] H. Garcke and K. F. Lam. "Well-posedness of a Cahn-Hilliard system modelling tumour growth with chemotaxis and active transport". In: European J. Appl. Math. 28.2 (2017), pp. 284–316.
- [26] H. Garcke and K.F. Lam. "Analysis of a Cahn-Hilliard system with non-zero Dirichlet conditions modeling tumor growth with chemotaxis". In: *Discrete Contin. Dyn. Syst.* 37.8 (2017), pp. 4277– 4308.
- [27] H. Garcke, K.F. Lam, R. Nürnberg, and E. Sitka. "A multiphase Cahn-Hilliard-Darcy model for tumour growth with necrosis". In: *Math. Models Methods Appl. Sci.* 28.3 (2018), pp. 525–577.
- [28] H. Garcke, K.F. Lam, E. Sitka, and V. Styles. "A Cahn-Hilliard-Darcy model for tumour growth with chemotaxis and active transport". In: *Math. Models Methods Appl. Sci.* 26.6 (2016), pp. 1095– 1148.
- [29] P. Grisvard. Elliptic problems in nonsmooth domains. Vol. 69. Classics in Applied Mathematics. Reprint of the 1985 original. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011, pp. xx+410.
- [30] A. Hawkins-Daarud, K. G. van der Zee, and J. T. Oden. "Numerical simulation of a thermodynamically consistent four-species tumor growth model". In: Int. J. Numer. Methods Biomed. Eng. 28.1 (2012), pp. 3–24.
- [31] D. Hilhorst, J. Kampmann, T. N. Nguyen, and K. G. Van Der Zee. "Formal asymptotic limit of a diffuse-interface tumor-growth model". In: *Math. Models Methods Appl. Sci.* 25.6 (2015), pp. 1011– 1043.
- [32] J. Jiang, H. Wu, and S. Zheng. "Well-posedness and long-time behavior of a non-autonomous Cahn-Hilliard-Darcy system with mass source modeling tumor growth". In: J. Differential Equations 259.7 (2015), pp. 3032–3077.
- [33] J. Necas. Direct methods in the theory of elliptic equations. Springer Monographs in Mathematics. Translated from the 1967 French original by G. Tronel and A. Kufner. Springer, Heidelberg, 2012, pp. xvi+372.
- [34] J. T. Oden, A. Hawkins, and S. Prudhomme. "General diffuse-interface theories and an approach to predictive tumor growth modeling". In: *Math. Models Methods Appl. Sci.* 20.3 (2010), pp. 477–517.
- [35] L. Preziosi, ed. Cancer modelling and simulation. Chapman & Hall/CRC Mathematical Biology and Medicine Series. Chapman & Hall/CRC, Boca Raton, FL, 2003, pp. xxvi+426.
- [36] L. Preziosi and A. Tosin. "Multiphase modelling of tumour growth and extracellular matrix interaction: mathematical tools and applications". In: J. Math. Biol. 58.4-5 (2009), pp. 625–656.
- [37] E. Sánchez-Palencia. "On the asymptotics of the fluid flow past an array of fixed obstacles". In: Internat. J. Engrg. Sci. 20.11 (1982), pp. 1291–1301.

- [38] C. G. Simader and H. Sohr. "A new approach to the Helmholtz decomposition and the Neumann problem in L^q-spaces for bounded and exterior domains". In: *Mathematical problems relating to* the Navier-Stokes equation. Vol. 11. Ser. Adv. Math. Appl. Sci. World Sci. Publ., River Edge, NJ, 1992, pp. 1–35.
- [39] J. Simon. "Compact sets in the space $L^p(0,T,B)$ ". In: Annali di Matematica Pura ed Applicata 146.1 (1986), pp. 65–96.
- [40] H. Sohr. The Navier-Stokes equations. Modern Birkhäuser Classics. An elementary functional analytic approach, 2013 reprint of the 2001 original. Birkhäuser/Springer Basel AG, Basel, 2001, pp. x+367.
- [41] S. M. Wise, J. S. Lowengrub, H. B. Frieboes, and V. Cristini. "Three-dimensional multispecies nonlinear tumor growth—I: Model and numerical method". In: J. Theoret. Biol. 253.3 (2008), pp. 524–543.