Equivalence of helicity and Euclidean self-duality for gauge fields $*$

Leonard Gross Department of Mathematics Cornell University Ithaca, NY 14853-4201 gross@math.cornell.edu

November 20, 2018

Abstract

In the canonical formalism for the free electromagnetic field a solution to Maxwell's equations is customarily identified with its initial gauge potential (in Coulomb gauge) and initial electric field, which together determine a point in phase space. The solutions to Maxwell's equations, all of whose plane waves in their plane wave expansions have positive helicity, thereby determine a subspace of phase space. We will show that this subspace consists of initial gauge potentials which lie in the positive spectral subspace of the operator curl together with initial electric fields conjugate to such potentials. Similarly for negative helicity. Helicity is thereby characterized by the spectral subspaces of curl in configuration space. A gauge potential on three-space has a Poisson extension to a four dimensional Euclidean half space, defined as the solution to the Maxwell-Poisson equation whose initial data is the given gauge potential. We will show that the extension is anti-self dual if and only if the gauge potential lies in the positive spectral subspace of curl. Similarly for self dual extension and negative spectral

[∗]Key words and phrases. Helicity, self duality, Yang-Mills, non-abelian gauge theories 2010 Mathematics Subject Classification. Primary; 81T13, 78A25, Secondary; 78A40.

subspace. Helicity is thereby characterized for a normalizable electromagnetic field by the canonical formalism and (anti-)self duality.

For a non-abelian gauge field on Minkowski space a plane wave expansion is not gauge invariant. Nor is the notion of positive spectral subspace of curl. But if one replaces the Maxwell-Poisson equation by the Yang-Mills-Poisson equation then (anti-)self duality on the Euclidean side induces a decomposition of (now non-linear) configuration space similar to that in the electromagnetic case. The strong analogy suggests a gauge invariant definition of helicity for non-abelian gauge fields. We will provide further support for this view.

Contents

8 Bibliography [47](#page-46-0)

1 Introduction

A workable notion of helicity for quantized Yang-Mills fields has been sought in many works in recent years. For a broad perspective on this work see the extensive review by Leader and Lorcé $[32]$. Any notion of helicity for Yang-Mills fields must be gauge invariant, Lorentz invariant and reduce to the standard notion of helicity for electromagnetic fields. This paper proposes a notion of helicity for non-abelian gauge fields based on a Euclidean interpretation of helicity in electromagnetism. For the latter, it will first be shown that the Poisson extension of the initial gauge potential of an electromagnetic field to a half space in Euclidean space-time sets up an equivalence between helicity, which is a Minkowski space notion, and (anti-)self duality, which is a Euclidean space notion.

For classical electromagnetic fields helicity is customarily defined in terms of plane waves, which do not mesh well with a non-linear theory such as Yang-Mills, [\[1\]](#page-46-1). To make the jump from electromagnetism to Yang-Mills fields, first classically (in this paper) and then quantum mechanically (in $[17]$), we are going to describe helicity in the electromagnetic case in a manner that allows a natural extension to the Yang-Mills case. To begin, we will first describe the precise configuration space for the free electromagnetic field that reflects the unique Lorentz invariant norm of Bargmann and Wigner, [\[5\]](#page-46-2). Second, given a real valued gauge potential $A(x)$ on \mathbb{R}^3 we will show that its Poisson extension to a half space in \mathbb{R}^4 not only captures the Lorentz invariant norm of Bargmann and Wigner, but is also self-dual if and only if A is the initial data in Minkowski space of some solution to Maxwell's equations of negative helicity. Similarly for anti-self dual and positive helicity.

To be more explicit, suppose that $A(x) := \sum_{j=1}^{3} A_j(x) dx^j$ is a real valued gauge potential on \mathbb{R}^3 and that, for each $s \geq 0$, $a(x, s) = \sum_{j=1}^3 a_j(x, s) dx^j$ is another gauge potential on \mathbb{R}^3 such that the function a satisfies the MaxwellPoisson equation with initial value A:

$$
\mathsf{a}''(s) = \operatorname{curl}^2 \mathsf{a}(s), \quad \mathsf{a}(0) = A. \tag{1.1}
$$

We may regard **a** as a 1-form in temporal gauge on the half-space $\mathbb{R}^4_+ :=$ $\mathbb{R}^3 \times [0, \infty)$. Its four dimensional curvature F is then given by $F = ds \wedge a' + da$, where d denotes the three dimensional exterior derivative. Equation (1.1) is the Euler equation for minimization of $\int_{\mathbb{R}^4_+} |F(x,s)|^2 d^3x ds$, subject to the initial condition $a(0) = A$. It happens that this minimum is the square of the unique Lorentz compatible norm on configuration space. We will show that A is the initial potential of some solution to Maxwell's equations with only positive helicity plane waves in its plane wave decomposition if and only if F is anti-self-dual. Similarly for negative helicity and self-duality. The Maxwell-Poisson equation [\(1.1\)](#page-3-0) thereby sets up a correspondence between helicity on the Minkowski side and (anti-)self-duality on the Euclidean side. The result is a decomposition of the electromagnetic configuration space $\mathcal C$ into two orthogonal subspaces \mathcal{C}_{\pm} , determined by helicity, or equivalently, by (anti-)self duality.

For a non-abelian gauge field the corresponding Poisson-like equation is the Yang-Mills-Poisson equation. It is a nonlinear, degenerate elliptic differential equation for which existence and uniqueness of solutions has not yet been proven. We are going to assume both, however, and show how the Yang-Mills-Poisson equation sets up a similar decomposition of the Yang-Mills configuration space \mathcal{C} , which is no longer a linear space, into two submanifolds C_{\pm} corresponding to anti-self dual and self dual solutions. The similarity of this procedure to the electromagnetic case suggests that the submanifolds \mathcal{C}_\pm are the "correct" non-abelian analogs of the classical helicity subspaces for Maxwell's theory. But we will give more support to this interpretation in Section [6](#page-37-0) by showing that this decomposition exhibits two additional properties which are definitive in the electromagnetic case.

The representation of a gauge potential as the initial data of a Poisson-like equation automatically induces a Riemannian metric on the set of potentials (modulo gauge transformations), whose Poisson action $\int_{\mathbb{R}^4_+} |F(x, s)|^2 d^3x ds$ is finite. This in turn gives a quantitative meaning to phase space $T^*(\mathcal{C})$ for both electromagnetic and Yang-Mills fields. It turns out, remarkably, that the norm defined on the electromagnetic phase space in this way is exactly the unique Lorentz invariant norm first discovered by Bargmann and Wigner [\[5\]](#page-46-2) in their fundamental work on unitary representations of the inhomogeneous

Lorentz group. The norm induced on configuration space and phase space by our procedure is therefore forced on us if we wish to have a Lorentz invariant theory. We will show this in the course of establishing the equivalence of helicity with (anti-)self duality for electromagnetism. In the non-abelian case the Riemannian metric induced from the Euclidean side by the same procedure is presumably Lorentz invariant in some appropriate sense. But this has not been explored.

In accordance with the canonical formalism (as in, say, [\[7,](#page-47-0) Chapter 14]) the electric field E at time zero is canonically conjugate to A and therefore the pair $\{A, E\}$ defines a point in phase space. We will show that if A is in \mathcal{C}_+ and E is in $T_A^*(\mathcal{C}_+)$ then the unique solution to Maxwell's equations with initial data A, E contains only positive helicity plane waves in its plane wave decomposition (and conversely). Similarly $T^*(\mathcal{C}_-)$ corresponds to negative helicity plane waves. In this way Euclidean (anti-)self duality provides a completely geometric characterization of helicity in electromagnetic theory

We will reinforce this geometric interpretation of helicity further by starting over again in [\[17\]](#page-48-0) with the quantized theories, each of which has a generally accepted "forward component of angular momentum" operator used for defining helicity in their respective quantum field theories. In the electromagnetic case we will see that the operator simply "quantizes" the decomposition C_{+} . The Schrödinger representation of the quantized electromagnetic field will be instrumental in implementing this view. In the non-abelian case, where configuration space is not a linear space, we will see that the apparent "forward component of angular momentum" operator similarly quantizes the two submanifolds C_+ , thereby reinforcing again the link between helicity and Euclidean (anti-)self-duality for non-abelian gauge fields. The Schrödinger representation of the quantized Yang-Mills field will again be indispensable in implementing this view, and for this, the Riemannian metric introduced in this paper will enable us to use the gradient operator on functions over configuration space instead of momentum space annihilation operators, which have no gauge invariant meaning for the intermediate non-abelian fields.

There are many unsettled purely mathematical issues in the body of this paper that we will ignore. They are in the nature of existence and uniqueness theorems. In the last section we will discuss the open problems raised by these structures.

2 Plane waves, helicity, configuration space in electromagnetism

2.1 Review of helicity for plane waves

This subsection is a review of circularly polarized light in a form that will be useful for our purposes. Nothing in this subsection should be construed to be less than 165 years old. See for example Born and Wolfe [\[8\]](#page-47-1) or Jackson [\[27\]](#page-49-1).

Notation 2.1 For $0 \neq \mathbf{k} \in \mathbb{R}^3$ define

$$
\mathbf{k}^{\perp} = \{ u \in \mathbb{C}^3 : u \cdot \mathbf{k} = 0 \}. \tag{2.1}
$$

 k^{\perp} is a two dimensional complex vector space closed under complex conjugation $u \mapsto \bar{u}$. Its real part $\mathbf{k}_{real}^{\perp} := \mathbf{k}^{\perp} \cap \mathbb{R}^3$ is a two dimensional real vector space. Define

$$
C_{\mathbf{k}}u = i\mathbf{k} \times u, \text{ for } u \in \mathbb{C}^3. \tag{2.2}
$$

 $C_{\mathbf{k}}u$ is in \mathbf{k}^{\perp} for all $u \in \mathbb{C}^{3}$ because $\mathbf{k} \times u$ is perpendicular to \mathbf{k} . The significance of the operator $C_{\mathbf{k}}$ is that it implements curl under Fourier transformation in accordance with the easily verified identity

$$
\operatorname{curl}(ue^{i\mathbf{k}\cdot x}) = (C_{\mathbf{k}}u)e^{i\mathbf{k}\cdot x} \quad \text{for any vector } u \in \mathbb{C}^3. \tag{2.3}
$$

Discussions of circularly polarized light and, more generally, elliptically polarized light amount to an analysis of the operator $C_{\mathbf{k}}$ acting in the two dimensional subspace \mathbf{k}^{\perp} . See for example [\[27,](#page-49-1) Chapter 7] or [\[8,](#page-47-1) Section 1.4].

Lemma 2.2 (Properties of $C_{\mathbf{k}}$). The restriction of $C_{\mathbf{k}}$ to \mathbf{k}^{\perp} satisfies

a) $C_{\mathbf{k}}$ is Hermitian in \mathbf{k}^{\perp} . b) $C_{\mathbf{k}}^2 = |\mathbf{k}|^2 on \mathbf{k}$ \perp (2.4)

c) $C_{\mathbf{k}}$ decomposes \mathbf{k}^{\perp} into two one dimensional complex subspaces \mathbf{k}^{\perp}_{\pm} corresponding to eigenvalues $\pm |\mathbf{k}|$. A vector $u \in \mathbf{k}^{\perp}$ is in \mathbf{k}_{+}^{\perp} if and only if its complex conjugate $\bar{u} \in \mathbf{k}_{-}^{\perp}$.

Proof. Using the triple product identity $(\mathbf{k} \times u) \cdot v = -u \cdot (\mathbf{k} \times v)$, which is valid for u and v in \mathbb{C}^3 , we see that

$$
(C_{\mathbf{k}}u,v)_{\mathbb{C}^3} = (i\mathbf{k}\times u)\cdot \bar{v} = u\cdot(\overline{i\mathbf{k}\times v}) = (u,C_{\mathbf{k}}v)_{\mathbb{C}^3},
$$

which proves a). Item b) follows from the identity $C_{\mathbf{k}}^2 u = -\mathbf{k} \times (\mathbf{k} \times u) = |\mathbf{k}|^2 u$ when $u \cdot \mathbf{k} = 0$. To prove c) let $\hat{\mathbf{k}} = \mathbf{k}/|\mathbf{k}|$ and let e_1 be any unit vector in $\mathbf{k}_{real}^{\perp}$. Define $e_2 = \hat{\mathbf{k}} \times e_1$. Then $\hat{\mathbf{k}}, e_1, e_2$ form a right handed basis of \mathbb{R}^3 . In particular $\hat{\mathbf{k}} \times e_2 = -e_1$. Thus

$$
C_{\mathbf{k}}(e_1 + ie_2) = i|\mathbf{k}|\{e_2 - ie_1\} = |\mathbf{k}|(e_1 + ie_2)
$$
\n(2.5)

Hence $(e_1 + ie_2)$ is an eigenvector for C_k with eigenvalue |k|. Using the same vectors one can compute similarly that $e_1 - ie_2$ is an eigenvector of $C_{\mathbf{k}}$ with eigenvalue $-|\mathbf{k}|$. Moreover, if $u \in \mathbf{k}_+^{\perp}$ then $C_{\mathbf{k}}\bar{u} = \overline{(-C_{\mathbf{k}}u)} = -|\mathbf{k}| \bar{u} = -|\mathbf{k}| \bar{u}$. This proves c). \blacksquare

Definition 2.3 (Plane waves) Let $0 \neq k \in \mathbb{R}^3$. A plane wave with wave vector **k** is a function on \mathbb{R}^4 of the form

$$
A_{\mathbf{k}}(x,t) = ae^{i(\mathbf{k}\cdot x - |\mathbf{k}|t)} + \bar{a}e^{-i(\mathbf{k}\cdot x - |\mathbf{k}|t)}, \ x, t \in \mathbb{R}^{3+1}, \ a \in \mathbf{k}^{\perp}
$$
 (2.6)

Clearly $A_{\mathbf{k}}(x,t) \in \mathbb{R}^3$ for each x, t and

$$
div A_{\mathbf{k}}(x,t) = 0 \tag{2.7}
$$

because of the identity

$$
div (ae^{i\mathbf{k}\cdot x}) = (i\mathbf{k}\cdot a)e^{i\mathbf{k}\cdot x}.
$$
 (2.8)

Lemma 2.4 For the plane wave (2.6) define

$$
B_{\mathbf{k}}(x,t) = curl A_{\mathbf{k}}(x,t)
$$
\n(2.9)

$$
E_{\mathbf{k}}(x,t) = -(\partial/\partial t)A_{\mathbf{k}}(x,t).
$$
 (2.10)

Then $B_{\mathbf{k}}, E_{\mathbf{k}}$ is a solution to Maxwell's equations,

$$
\nabla \cdot B = 0, \quad \nabla \cdot E = 0, \quad \dot{B} = -\text{curl } E, \quad \dot{E} = \text{curl } B. \tag{2.11}
$$

Proof. Use the identities [\(2.8\)](#page-6-1) and [\(2.3\)](#page-5-2) along with [\(2.6\)](#page-6-0) and [\(2.4\)](#page-5-3). \blacksquare

Definition 2.5 (Helicity for plane waves) The standard definition of helicity can be stated thus. The plane wave [\(2.6\)](#page-6-0) has *positive helicity* if $C_{\mathbf{k}}a = |\mathbf{k}|a$. It has negative helicity if $C_{\mathbf{k}}a = -|\mathbf{k}|a$.

2.2 The Lorentz invariant norm

A small portion of the classic paper [\[5\]](#page-46-2) of Bargmann and Wigner will be needed to explain our choice of the norm [\(2.13\)](#page-7-1), which we will later take to be the norm on configuration space in [\(2.35\)](#page-13-0). The Coulomb gauge fixing that we are going to use here for convenience will be removed when we discuss Yang-Mills fields.

Definition 2.6 (Spaces) The operator curl on vector fields over \mathbb{R}^3 acts in the real Hilbert space $L^2(\mathbb{R}^3; \mathbb{R}^3) \cap \{div u = 0\}$ because $div curl = 0$. It is easily seen to be self-adjoint by an integration by parts. Moreover in this space it has a zero nullspace. We will denote by C the operator curl acting in this Hilbert space or some closely related spaces. The square root of $C²$ will be denoted $|C|$: Thus

$$
C = curl, \qquad |C| = \sqrt{C^2}.
$$
\n(2.12)

For a vector field u on \mathbb{R}^3 with $div u = 0$ let

$$
||u||_{H_{1/2}}^2 = |||C|^{1/2}u||_{L^2}^2
$$
\n(2.13)

$$
||u||_{H_{-1/2}}^2 = |||C|^{-1/2}u||_{L^2}^2.
$$
\n(2.14)

Define Hilbert spaces $H_{\pm 1/2}$ by the condition that the corresponding norm in [\(2.13\)](#page-7-1) or [\(2.14\)](#page-7-2) is finite.

For vector fields B and E on \mathbb{R}^3 with divergence zero define

$$
\|\{B, E\}\|_{bw}^2 = \|B\|_{H_{-1/2}}^2 + \|E\|_{H_{-1/2}}^2.
$$
\n(2.15)

The main assertion of this section is that the norm $\|\cdot,\cdot\|_{bw}$ is invariant under Lorentz transformations in the sense that the unique solution to Maxwell's equations with B and E as initial data has, after any Lorentz transformation, initial data with the same norm. The following computations reduce this assertion to a theorem in [\[5\]](#page-46-2) by showing that $\|\cdot,\cdot\|_{bw}$ is equal to the norm introduced in [\[5\]](#page-46-2) by Bargmann and Wigner.

Theorem 2.7 (Plane wave decomposition) Suppose that $a(\mathbf{k})$ is a \mathbb{C}^3 valued function on \mathbb{R}^3 such that

$$
a. \quad \mathbf{k} \cdot a(\mathbf{k}) = 0 \text{ for all } \mathbf{k} \in \mathbb{R}^3 \text{ and } \tag{2.16}
$$

$$
b. \qquad \int_{\mathbb{R}^3} |a(\mathbf{k})|^2 d^3 \mathbf{k} / |\mathbf{k}| < \infty. \tag{2.17}
$$

Define

$$
A(x,t) = \int_{\mathbb{R}^3} \left(a(\mathbf{k}) e^{i(\mathbf{k} \cdot x - |\mathbf{k}|t)} + \overline{a(\mathbf{k})} e^{-i(\mathbf{k} \cdot x - |\mathbf{k}|t)} \right) d^3 \mathbf{k} / |\mathbf{k}|.
$$
 (2.18)

$$
B(x,t) = curl A(x,t).
$$
\n(2.19)

$$
E(x,t) = -(\partial/\partial t)A(x,t). \tag{2.20}
$$

Then A, B and E have divergence zero for each t and $B(\cdot), E(\cdot)$ is a solution to Maxwell's equations [\(2.11\)](#page-6-2). Moreover for all real t there holds

$$
\begin{aligned} \|\{B(t), E(t)\}\|_{bw}^2 &= \|A(t)\|_{H_{1/2}}^2 + \|E(t)\|_{H_{-1/2}}^2 \\ &= 2(2\pi)^3 \int_{\mathbb{R}^3} |a(\mathbf{k})|^2 d^3 \mathbf{k} / |\mathbf{k}|. \end{aligned} \tag{2.21}
$$

Conversely, suppose that B_0 and E_0 are divergence free vector fields on \mathbb{R}^3 and each is in $H_{-1/2}(\mathbb{R}^3)$. Then there is a unique solution $B(t)$, $E(t)$ to Maxwell's equations [\(2.11\)](#page-6-2) and a unique (up to a set of measure zero) function $a(\cdot)$: $\mathbb{R}^3 \to \mathbb{C}^3$ satisfying [\(2.16\)](#page-7-3) and [\(2.17\)](#page-7-4) such that $B(t)$ and $E(t)$ are given by $(2.18) - (2.20).$ $(2.18) - (2.20).$ $(2.18) - (2.20).$ $(2.18) - (2.20).$

Proof. Upon differentiating under the integral in [\(2.18\)](#page-8-0) we see from Lemma [2.4](#page-6-3) that $A(t)$, $B(t)$ and $E(t)$ all have divergence zero and the pair $\{B(t), E(t)\}$ satisfies Maxwell's equations.

For the proof of (2.21) observe first that, with $C = \text{curl}$ we have $||B(t)||_{H_{-1/2}} = ||A(t)||_{H_{1/2}}$ because $||B(t)||_{H_{-1/2}} = ||C A(t)||_{H_{-1/2}} =$ $|||C|^{-1/2}CA(t)||_{L^2} = |||C|^{1/2}A(t)||_{L^2}$. It suffices therefore to prove the sec-ond equality in line [\(2.21\)](#page-8-2). Moreover it suffices to prove it just at $t = 0$ because replacing $a(\mathbf{k})$ by $a(\mathbf{k})e^{-i|\mathbf{k}|t}$ in the $t=0$ equality gives the equality at time t. Let

$$
\alpha(\mathbf{k}) = a(\mathbf{k}) + \overline{a(-\mathbf{k})} \quad \text{and} \quad (2.22)
$$

$$
e(\mathbf{k}) = i|\mathbf{k}| \left(a(\mathbf{k}) - \overline{a(-\mathbf{k})} \right). \tag{2.23}
$$

Then from (2.18) and (2.20) we find

$$
A(x,0) = \int_{\mathbb{R}^3} \frac{\alpha(\mathbf{k})}{|\mathbf{k}|} e^{i\mathbf{k} \cdot x} d\mathbf{k}.
$$
 (2.24)

$$
E(x,0) = \int_{\mathbb{R}^3} \frac{e(\mathbf{k})}{|\mathbf{k}|} e^{i\mathbf{k} \cdot x} d\mathbf{k}.
$$
 (2.25)

Therefore

$$
||A(\cdot,0)||_{H_{1/2}}^{2} + ||E(\cdot,0)||_{H_{-1/2}}^{2}
$$

\n
$$
= (2\pi)^{3} \int_{\mathbb{R}^{3}} \left(\left| |\mathbf{k}|^{1/2} \frac{\alpha(\mathbf{k})}{|\mathbf{k}|} \right|^{2} + \left| |\mathbf{k}|^{-1/2} \frac{e(\mathbf{k})}{|\mathbf{k}|} \right|^{2} \right) d^{3} \mathbf{k}
$$

\n
$$
= (2\pi)^{3} \int_{\mathbb{R}^{3}} \left(|a(\mathbf{k}) + \overline{a(-\mathbf{k})}|^{2} + |a(\mathbf{k}) - \overline{a(-\mathbf{k})}|^{2} \right) d^{3} \mathbf{k} / |\mathbf{k}|
$$

\n
$$
= (2\pi)^{3} \int_{\mathbb{R}^{3}} \left(|a(\mathbf{k})|^{2} + |\overline{a(-\mathbf{k})}|^{2} \right) d^{3} \mathbf{k} / |\mathbf{k}|
$$

\n
$$
= (2\pi)^{3} 2 \int_{\mathbb{R}^{3}} |a(\mathbf{k})|^{2} d^{3} \mathbf{k} / |\mathbf{k}|.
$$
 (2.26)

This proves the equality in line [\(2.21\)](#page-8-2).

Conversely, suppose that B_0 and E_0 are given, that div $B_0 = div E_0 = 0$ and that $||B_0||^2_{H_{-1/2}} + ||E_0||^2_{H_{-1/2}} < \infty$. Let $A_0 = (curl)^{-1}B_0$. This defines A₀ as a potential with $div A_0 = 0$ and $||A_0||_{H_{1/2}} = ||B_0||_{H_{-1/2}} < \infty$. Define $\alpha(\mathbf{k})$ and $e(\mathbf{k})$ from A_0 and E_0 by [\(2.24\)](#page-8-3) and [\(2.25\)](#page-8-4) respectively. Since A_0 and E_0 are "real", i.e. both take their values in \mathbb{R}^3 , we have $\alpha(-\mathbf{k}) = \overline{\alpha(\mathbf{k})}$ and $e(-\mathbf{k}) = \overline{e(\mathbf{k})}$. With [\(2.22\)](#page-8-5) and [\(2.23\)](#page-8-6) in mind, define

$$
a(\mathbf{k}) = (1/2) \Big(\alpha(\mathbf{k}) + \frac{e(\mathbf{k})}{i|\mathbf{k}|} \Big). \tag{2.27}
$$

From the hermiticity of $\alpha(\cdot)$ and $e(\cdot)$ it follows that

$$
\overline{a(-\mathbf{k})} = (1/2) \left(\alpha(\mathbf{k}) - \frac{e(\mathbf{k})}{i|\mathbf{k}|} \right) \tag{2.28}
$$

and from this the relations [\(2.22\)](#page-8-5) and [\(2.23\)](#page-8-6) follow. The computation [\(2.26\)](#page-9-0) now shows that

$$
||A_0||_{H_{1/2}}^2 + ||E_0||_{H_{-1/2}}^2 = (2\pi)^3 2 \int_{\mathbb{R}^3} |a(\mathbf{k})|^2 d^3 \mathbf{k} / |\mathbf{k}| \tag{2.29}
$$

Hence (2.17) holds for the constructed function $a(\mathbf{k})$. Of course (2.16) follows because A_0 and E_0 are divergence free. The solution $B(t)$, $E(t)$ with initial values B_0, E_0 can now be constructed from $a(\mathbf{k})$ by (2.18) - (2.20) .

Remark 2.8 (Integral formula for $\|\{B, E\}\|_{bw}^2$) Since the Laplacian ($\Delta \equiv$ ∇^2) on vector fields $u(x)$ over \mathbb{R}^3 is given by $-\Delta u(x) = \operatorname{curl}^2 u(x) - \operatorname{grad} \operatorname{div} u(x)$

we may write $-\Delta u(x) = \frac{curl^2 u(x)}{\Delta x}$ in the space of divergence free vector fields. The space of divergence free vector fields is invariant under Δ and consequently $|curl|u = (-\Delta)^{1/2}u$ on this space. One therefore has on this space $|curl|^{-1} = (-\Delta)^{-1/2}$, which, by Fourier transform, is easily seen to be given by convolution by $const.(1/|x|^2)$. Since $||u||_{H_{-1/2}}^2 = (|C|^{-1}u, u)_{L^2(\mathbb{R}^3)}$ on the space of divergence free vector fields u , the Bargmann-Wigner norm [\(2.15\)](#page-7-5) can be written

$$
\|\{B, E\}\|_{bw}^2 = const. \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{B(x) \cdot B(y) + E(x) \cdot E(y)}{|x - y|^2} dx dy.
$$
 (2.30)

This representation of the Lorentz invariant norm for the electromagnetic field was derived in [\[18\]](#page-48-1) and used there to show that the norm is invariant under the 15 dimensional conformal group of Minkowski space, which contains transformations to uniformly accelerated coordinate systems. (See e.g. [\[18,](#page-48-1) footnote 2]). It was shown in [\[18\]](#page-48-1) that each of the two terms in [\(2.30\)](#page-10-1) is separately invariant under the inversion $x \mapsto x/|x|^2$ of \mathbb{R}^3 , which, together with dilations of \mathbb{R}^4 and the inhomogeneous Lorentz group, generate the conformal group. Unitarity of the representation of the conformal group, as opposed to just norm invariance, requires also proof of invariance of the complex structure, which will be described on the configuration space side in the proof of Theorem [2.9.](#page-10-2) Invariance of the complex structure under the conformal group was first proved in [\[29\]](#page-49-2).

2.3 Helicity and sgn(curl) for solutions of finite action

Our goal in this subsection is to take a step away from the plane wave decomposition [\(2.18\)](#page-8-0) by formulating helicity entirely on the position space side with the help of the operator curl. This is reflected explicitly in the equivalence between assertions 1.) and 2.) in the next theorem

Theorem 2.9 Suppose that $B(t)$, $E(t)$ is a solution to Maxwell's equations (2.11) with finite Bargmann-Wigner norm. Let (2.18) - (2.20) be its plane wave decomposition. Then the following are equivalent.

1.) Every plane wave in its plane wave decomposition has positive helicity.

2.) $A(0)$ and $E(0)$ are in the positive spectral subspace of curl.

3.) $B(t)$ and $E(t)$ are both in the positive spectral subspace of curl for some t.

4.) $B(t)$ and $E(t)$ are both in the positive spectral subspace of curl for all t

5.) $A(t)$ is in the positive spectral subspace of curl for all t. These statements also hold with positive replaced by negative everywhere.

Proof. The space of functions $a(\mathbf{k})$ on momentum space satisfying (2.16) and [\(2.17\)](#page-7-4) represents Bargmann and Wigner's original description of the one-particle photon space, [\[5\]](#page-46-2). Denote it by BW_{mom} . On the position space side denote by BW the space of divergence free real fields B, E over \mathbb{R}^3 with finite norm [\(2.15\)](#page-7-5). Define

$$
W: BW_{mom} \to BW \tag{2.31}
$$

by $W: a(\mathbf{k}) \mapsto {B(x, 0), E(x, 0)}$ where B and E are given by [\(2.18\)](#page-8-0) - [\(2.20\)](#page-8-1). Then Theorem [2.7](#page-7-6) shows that W is a norm preserving transformation from BW_{mom} onto the direct sum space BW if one uses $2(2\pi)^3 \int_{\mathbb{R}^3} |a(\mathbf{k})|^2 d^3\mathbf{k}/|\mathbf{k}|$ for the norm squared on BW_{mom} . Actually, since there is no hermiticity condition on $a(\mathbf{k})$, BW_{mom} is a complex Hilbert space under the usual operation of multiplication by $i \equiv \sqrt{-1}$. BW itself has a natural complex structure j given by $j\{B, E\} = \{sgn(C)E, -sgn(C)B\}$, where $sgn(C) = |C|^{-1}C$. The latter complex structure is more often expressed in terms of the pair $\{A, E\}$ in the form $j\{A, E\} = \{|C|^{-1}E, -|C|A\}$, which is an equivalent description since $B = CA$. BW is therefore also a complex Hilbert space. Comparing (2.22) and (2.23) with (2.24) and (2.25) one sees easily that $Wia = jWa$. W is therefore unitary. Furthermore (2.3) , $(2.18)-(2.20)$ $(2.18)-(2.20)$, show that W intertwines curl with the operator $C_{(\cdot)}$, which is multiplication by $C_{\mathbf{k}}$.

$$
curl W = WC_{(\cdot)}.
$$
\n(2.32)

Here we are writing $curl \{B, E\} = \{curl B, curl E\}.$

Now suppose that $B(t)$, $E(t)$ is a solution to Maxwell's equations [\(2.11\)](#page-6-2) with finite Bargmann-Wigner norm and with plane wave decomposition given by (2.18) - (2.20) . Suppose also that it has only positive helicity plane waves in its plane wave decomposition. According to Definition [2.5](#page-6-4) $C_{\bf k} a({\bf k})$ = $|\mathbf{k}|a(\mathbf{k})$ for every $\mathbf{k} \neq 0$ in \mathbb{R}^3 . Therefore $a(\cdot)$ is in the positive spectral subspace for the operator multiplication by $C_{\mathbf{k}}$. Since W is unitary, $\{B(0), E(0)\}$ is in the positive spectral subspace of curl on BW, (which is easily seen to be equivalent to saying that $B(0)$ and $E(0)$ are each in the positive spectral subspace of *curl.*) Since $|C| : H_{1/2} \to H_{-1/2}$ is an orthogonal transformation and commutes with curl it preserves all spectral properties of curl. In particular $A(0)$ is also in the positive spectral subspace of curl because $B_0 = \text{curl } A_0$. Therefore statement 1.) implies statement 2.). Conversely, if $A(0)$ and $E(0)$ are in the positive spectral subspace of $curl$ then $B(0)$ is in the positive spectral subspace of curl and so therefore is the pair $\{B(0), E(0)\}\$. By (2.32) and the unitarity of W it follows that $a(\cdot)$ is in the positive spectral subspace of the operator of multiplication by $C_{\mathbf{k}}$. Hence $C_{\mathbf{k}}a(\mathbf{k}) = |\mathbf{k}|a(\mathbf{k})$ for all $\mathbf{k} \neq 0$. Therefore statement 1.) holds. This proves the equivalence of statements 1.) and 2.). The proof shows that the pair $\{A(0), E(0)\}\)$ could be replaced by the pair ${B(0), E(0)}$ in the statement of item 2.). Now for any fixed t the map $W_t: a(\cdot) \mapsto {B(t), E(t)}$ is also unitary and intertwines *curl* and C_k . Therefore the proof of equivalence of 1.) with 2.) also applies to the proof of equivalence of 1.) with 3.) and 1.) with 4.).

Concerning the condition 5.), we see that if 1.) holds then by 2.) $A(0)$, and similarly $A(t)$, is in the positive spectral subspace of curl in its own space $H_{1/2}$. Conversely, if 5.) holds then $(d/dt)A(t)$ is in the positive spectral subspace of curl. In particular $A(0)$ and its time derivative $-E(0)$ are in the positive spectral subspace of curl and we can apply 2.) to see that 1.) holds. $\mathcal{L}_{\mathcal{A}}$

Remark 2.10 The operator of multiplication by C_k on BW_{mom} decomposes BW_{mom} into two orthogonal subspaces given respectively by the two identities by $C_{\mathbf{k}}a(\mathbf{k}) = |\mathbf{k}|a(\mathbf{k})$ and $C_{\mathbf{k}}a(\mathbf{k}) = -|\mathbf{k}|a(\mathbf{k})$. These are the two spectral subspaces of curl on the momentum space side.

2.4 Configuration space, phase space and helicity

The Lorentz invariant norm [\(2.21\)](#page-8-2) has a geometric interpretation based on the canonical formalism for the electromagnetic field: As in [\[7,](#page-47-0) Chapter 14], we take configuration space for the free electromagnetic field to be a set of divergence free potentials A on \mathbb{R}^3 with size of A yet to be specified. This corresponds to the radiation gauge. The momentum canonically conjugate to A is $-\dot{A}$, which is E. See [\[7,](#page-47-0) Eq. (14.10)] for a derivation of this from the Lagrangian formalism. The canonical formalism therefore dictates that the pair A, E is a point in phase space. Thus if A lies in some configuration space C (still to be determined, quantitatively) and $T_A(\mathcal{C})$ denotes the tangent space to C at A, then E is a point in its dual space $T_A^*(\mathcal{C})$, as is customary for momentum in classical mechanics. In the next definition we will make a quantitative choice for $\mathcal C$ and show that the norm on phase space which is automatically induced by this choice of configuration space is the Bargmann-Wigner norm [\(2.21\)](#page-8-2).

Definition 2.11 (Configuration space) Let

 $C = \{$ real valued 1-forms A on \mathbb{R}^3 such that (2.33)

a. $div A = 0$ (2.34)

$$
b. \quad ||A||_{\mathcal{C}} := ||C|^{1/2} A||_{L^2(\mathbb{R}^3)} < \infty \}. \tag{2.35}
$$

Here $C = \text{curl}$ as before. Define also

 $C_+ = \{A \in \mathcal{C} : A \text{ is in the positive spectral subspace of } curl\}$ (2.36)

$$
C_{-} = \{ A \in \mathcal{C} : A \text{ is in the negative spectral subspace of } curl \}. \tag{2.37}
$$

Then $\mathcal C$ is a real Hilbert space and

$$
\mathcal{C} = \mathcal{C}_+ \oplus \mathcal{C}_-.\tag{2.38}
$$

The subspaces intersect only at $A = 0$ because curl does not have a zero eigenvalue in \mathcal{C} .

Since $\mathcal C$ is a vector space its tangent space at a point A can be identified with C itself. The dual space can therefore be identified with \mathcal{C}^* : $T_A^*(\mathcal{C}) \equiv \mathcal{C}^*$. Consequently the entire phase space can be identified as

$$
T^*(\mathcal{C}) \equiv \mathcal{C} \oplus \mathcal{C}^*.
$$
\n
$$
(2.39)
$$

The electric field E is to be identified with an element of the dual space via the pairing

$$
\langle E, A \rangle = \int_{\mathbb{R}^3} E(x) \cdot A(x) d^3 x,\tag{2.40}
$$

which is the field theoretic analog of $\sum_{i=1}^{n} p_i dq_i$, when one identifies $T_A(\mathcal{C})$ with \mathcal{C} . Combined with our choice of norm (2.35) , this pairing induces on E the norm $||E||_{\mathcal{C}^*} = |||C|^{-1/2}E||_{L^2(\mathbb{R}^3)}$. Since C involves the restriction $div A = 0$, so does the dual space. We can therefore make the natural identification

$$
C^* = \{ \text{real valued vector fields } E \text{ on } \mathbb{R}^3 \text{ such that } (2.41)
$$

$$
a. \quad \operatorname{div} E = 0 \tag{2.42}
$$

$$
b. \quad ||E||_{\mathcal{C}^*} := ||C|^{-1/2} E||_{L^2(\mathbb{R}^3)} < \infty \}. \tag{2.43}
$$

In the canonical formalism a point in phase space is thereby specified by a pair $\{A, E\}$ with $||A||_{\mathcal{C}}^2 + ||E||_{\mathcal{C}^*}^2 < \infty$ and $div A = div E = 0$.

We see from (2.21) at $t = 0$ that this is precisely the Bargmann-Wigner norm on the initial data when we take $B = \text{curl } A$.

$$
\|\{B, E\}\|_{bw}^2 = \|A\|_{\mathcal{C}}^2 + \|E\|_{\mathcal{C}^*}^2, \quad \text{div}\,A = 0, \quad \text{div}\,E = 0. \tag{2.44}
$$

Since C commutes with $|C|$ the annihilator in \mathcal{C}^* of the positive spectral subspace of curl in \mathcal{C} is the negative spectral subspace of curl in \mathcal{C}^* . Similarly with positive and negative reversed. Consequently we have the natural identifications

$$
\mathcal{C}_{+}^{*} = \{ E \in \mathcal{C}^{*} : E \text{ is in the positive spectral subspace of } curl \}. \tag{2.45}
$$

$$
\mathcal{C}_{-}^{*} = \{ E \in \mathcal{C}^{*} : E \text{ is in the negative spectral subspace of } curl \}. \tag{2.46}
$$

Analogous to the identification [\(2.39\)](#page-13-1) we therefore have the identifications

$$
T^*(\mathcal{C}_+) = \mathcal{C}_+ \oplus \mathcal{C}_+^*, \quad T^*(\mathcal{C}_-) = \mathcal{C}_- \oplus \mathcal{C}_-^*, \quad T^*(\mathcal{C}) = T^*(\mathcal{C}_+) + T^*(\mathcal{C}_-). \tag{2.47}
$$

Theorem 2.12 Suppose that the pair $\{A, E\}$ is a point in phase space $T^*(\mathcal{C})$. Let $A(t)$, $E(t)$ be the unique solution to Maxwell's equations with initial data $A(0) = A, -A(0) = E(0) = E$. Then the plane wave decomposition of the solution consists of

- a) positive helicity plane waves if and only if $\{A, E\}$ is in $T^*(\mathcal{C}_+)$.
- b) negative helicity plane waves if and only if $\{A, E\}$ is in $T^*(\mathcal{C}_-)$.

Proof. The assumption that $\{A, E\}$ lies in $T^*(\mathcal{C})$ is equivalent, by [\(2.44\)](#page-14-0), to the assumption that the pair has finite Bargmann-Wigner norm. In view of (2.36) , (2.37) , (2.45) and (2.46) , the present theorem just restates the equivalence between items 1.) and 2.) in Theorem [2.9.](#page-10-2) \blacksquare

Remark 2.13 (Primacy of the configuration space decomposition) The helicity character of a solution to Maxwell's equations is not determined by the value of $A(0)$ alone but also requires knowledge of $A(0)$. To conclude, for example, that a solution has only positive helicity plane waves in its plane wave decomposition it is not sufficient to know that $A(0)$ is in the positive spectral subspace of curl. One must know also that $A(0)$ is in the positive spectral subspace of curl. And yet the decomposition (2.38) of C is responsible for determining helicity in the sense that once this decomposition is known it

automatically determines the decomposition of $T^*(\mathcal{C})$ as in [\(2.47\)](#page-14-3), allowing Theorem [2.12](#page-14-4) to be applied. The decomposition of $T^*(\mathcal{C})$ thereby plays a secondary role compared to the decomposition [\(2.38\)](#page-13-4). This is important to observe for two disparate reasons. First, in the quantized theory, where a state of the field is given by a function ψ on C, the Heisenberg field expectations $(B(x, t)\psi, \psi)$ and $(E(x, t)\psi, \psi)$ in the state ψ will be shown in [\[17\]](#page-48-0) to have only positive helicity plane waves in their plane wave decomposition when ψ depends only on \mathcal{C}_+ . Second, in the non-abelian theory, where the configuration space is a non-linear manifold, the electromagnetic decomposition [\(2.38\)](#page-13-4) has a natural analog with properties that well imitate those of the electromagnetic case. This will be shown in Section [6.](#page-37-0)

Remark 2.14 (A bit of history) Immediately after Heisenberg and Pauli published their fundamental paper [\[20\]](#page-48-2) in 1929, describing the quantized electromagnetic field, Landau and Peierls [\[31\]](#page-49-3), in 1930, rederived some of the formalism of Heisenberg and Pauli, emphasizing the role of the electromagnetic configuration space. In the formulation of Landau and Peierls E and B were not independent. They introduced a norm equal to one of the two terms in (2.15) (or equivalently (2.30)) for the normalization condition for their configuration space. See e.g. $[31, \text{ Equations } (7)-(10)]$. It is now understood that in the canonical formalism A and E are to be regarded as independent variables in phase space. So their suggestion to allow either term in [\(2.30\)](#page-10-1) as the norm on their configuration space obscured the distinction between configuration space and phase space. Interestingly, they arrived at their norm, not by citing Lorentz invariance, but by arguing that a unit vector in this particular norm represents a single photon. An updated version of their "number of photons" argument for this norm, leading to the full phase space norm [\(2.30\)](#page-10-1), can be found in Jackson [\[27,](#page-49-1) Problem 7.30].

3 Equivalence of helicity and Euclidean selfduality in electromagnetism

3.1 The Poisson action and Maxwell-Poisson semigroup

The helicity decomposition [\(2.38\)](#page-13-4) and it's associated decomposition of phase space [\(2.47\)](#page-14-3) are based on use of the spectral decomposition of the operator curl and were shown to be equivalently characterizable in terms of plain wave expansions. (cf. Theorems [2.9](#page-10-2) and [2.12.](#page-14-4)) But neither plain wave expansions nor the operator curl mesh well with non-abelian gauge fields. In this section we are going to show that the helicity decomposition of configuration space can be reproduced without using either plane wave expansions or the spectral decomposition of curl. Instead, we will extend the gauge potential A to a half-space in \mathbb{R}^4 and show that the decomposition is equivalent to (anti-)selfduality of the extension. This method for implementing the decomposition [\(2.38\)](#page-13-4) in electromagnetism will be shown in succeeding sections to go over to non-abelian gauge fields.

To simplify otherwise cumbersome calculations we will make use henceforth of differential form notation. Recall that if (u_1, u_2, u_3) is a vector field on \mathbb{R}^3 then the corresponding differential form is given by $u = \sum_{i=1}^3 u_i dx^i$, the Hodge $*$ operator is given on 1-forms by $*u = (1/2) \sum_{i,j,k} \epsilon_{ijk} u_i dx^j \wedge dx^k$ and on 2-forms by $*\sum_{jk} v_{jk} dx^j \wedge dx^k = \sum_{ijk} \epsilon_{ijk} v_{jk} dx^i$. The exterior derivative is given by $du = \sum_{ji} (\partial u_i / \partial x_j) dx^j \wedge dx^i$ and the action of curl on 1-forms is given by $curl u = *du$.

Notation 3.1 Denote by \mathbb{R}^4_+ the Euclidean half space $\mathbb{R}^3 \times [0, \infty)$ with coordinates (x, s) . Let a be a real valued 1-form on \mathbb{R}^4_+ in temporal gauge. That is,

$$
\mathsf{a}(x,s) = \sum_{j=1}^{3} \mathsf{a}_j(x,s) dx^j, \quad x \in \mathbb{R}^3, \quad s \in [0,\infty). \tag{3.1}
$$

Then the four dimensional curvature of **a** is given by

$$
F = ds \wedge \mathbf{a}' + d\mathbf{a},\tag{3.2}
$$

where $' = (\partial/\partial s)$ and d is again the three dimensional exterior derivative operator.

We wish to minimize the functional

$$
\int_{\mathbb{R}_+^4} |F(x,s)|^2 dx ds \tag{3.3}
$$

subject to the initial condition

$$
a(x, 0) = A(x), \quad x \in \mathbb{R}^3,
$$
 (3.4)

wherein A is a given real valued 1-form on \mathbb{R}^3 . Observe first that

$$
|F(x,s)|_{\Lambda^2}^2 = |\mathbf{a}'(x,s)|_{\Lambda^1}^2 + |d\mathbf{a}(x,s)|_{\Lambda^2}^2
$$
\n(3.5)

for each (x, s) because the first 2-form on the right side of (3.2) is orthogonal to the second one. In [\(3.5\)](#page-17-0) Λ ¹ denotes, as usual, the three dimensional space of linear functionals on \mathbb{R}^3 and Λ^2 denotes the three dimensional space of skew symmetric bilinear functionals on \mathbb{R}^3 . The variational equation for the minimization problem can be derived in the usual way as follows. Let $u(x,s) = \sum_{j=1}^{3} u_j(x,s) dx^j$ with each coefficient lying in $C_c^{\infty}(\mathbb{R}^3 \times (0,\infty)).$ Then we find

$$
\partial_u \int_{\mathbb{R}_+^4} \left\{ |\mathbf{a}'(x,s)|^2 + |d\mathbf{a}(x,s)|^2 \right\} dx ds
$$

= $2 \int_{\mathbb{R}_+^4} \left\{ - (\mathbf{a}''(x,s), u(x,s)) + (d^* d\mathbf{a}(x,s), u(x,s)) \right\} dx ds$ (3.6)

after doing an integration by parts in both time and space. The variational equation is therefore

$$
\mathsf{a}'' = d^*d\mathsf{a}.\tag{3.7}
$$

This is not quite Poisson's equation because half of the Laplacian, $-\Delta =$ $d^*d + dd^*$, is missing from the right hand side. But it is intimately related to Maxwell's theory and we will refer to [\(3.7\)](#page-17-1) as the Maxwell-Poisson equation. It is easily solved. Since $d^* = *d^*$ on 2-forms over \mathbb{R}^3 we see that $d^*d = (*d)(*d) = curl^2$. The equation [\(3.7\)](#page-17-1) may therefore be written as $a''(s) = C^2 a(s)$ with $C = curl$. The function $a(s) = e^{sC} A$ clearly solves this equation as does also $a(s) = e^{-sC}A$. But if A is in the strictly positive spectral subspace of curl then $\|\mathsf{a}'(s)\|_{L^2(\mathbb{R}^3)}$ grows exponentially in the first case, making (3.3) infinite, while if A is in the strictly negative spectral subspace of curl then [\(3.3\)](#page-16-1) is infinite in the second case. The solution in both cases is given, for arbitrary $A \in L^2(\mathbb{R}^3, \Lambda^1)$, by

$$
\mathsf{a}(s) := e^{-s|C|} A, \quad s \ge 0. \tag{3.8}
$$

It solves the Maxwell-Poisson equation [\(3.7\)](#page-17-1) because $a'' = |C|^2 a = C^2 a$ d^*d and also satisfies the initial condition $a(0) = A$. Under reasonable growth restrictions on $a(\cdot)$ the solution given by [\(3.8\)](#page-17-2) is unique. We will always assume uniqueness. The size of the minimum in [\(3.3\)](#page-16-1) will be discussed in the following. It bears emphasizing that we are not making an assumption that A is in Coulomb gauge in this discussion.

Terminology 3.2 The unique solution to the Maxwell-Poisson equation [\(3.7\)](#page-17-1) with initial value A will be referred to as the Poisson extension of A. We define the Poisson action of A by

$$
\mathcal{P}(A) = \int_0^\infty \left(\| \mathbf{a}'(s) \|_2^2 + \| d\mathbf{a}(s) \|_2^2 \right) ds,\tag{3.9}
$$

where a is the Poisson extension of A . We are only interested in those potentials A for which $\mathcal{P}(A) < \infty$. We reiterate that $\mathcal{P}(A)$ is the minimum of the integral in [\(3.9\)](#page-18-0) subject to the condition that $a(0) = A$.

Remark 3.3 The Poisson action of a gauge potential A can be computed explicitly. We need not assume that A is in Coulomb gauge. From (3.8) we find

$$
\mathbf{a}'(s) = -|C|e^{-s|C|}A
$$

\n
$$
\|\mathbf{a}'(s)\|_2^2 = \|\ |C|e^{-s|C|}A\|_2^2 = (|C|^2e^{-2s|C|}A, A)
$$

\n
$$
\|d\mathbf{a}(s)\|_2^2 = \|\ * d\mathbf{a}(s)\|_2^2 = \|Ce^{-s|C|}A\|_2^2 = (|C|^2e^{-2s|C|}A, A).
$$

Therefore

$$
\mathcal{P}(A) = 2 \int_0^\infty (|C|^2 e^{-2s|C|} A, A) ds
$$

$$
= (|C|A, A).
$$
(3.10)

A reader concerned about the lack of exponential decrease in [\(3.8\)](#page-17-2) when A is in the null space of curl should observe that in this case the integrand in [\(3.9\)](#page-18-0) is identically zero.

Remark 3.4 The identity (3.10) holds even if A is longitudinal, that is, $A = d\lambda$ for some real scalar function λ . In this case A is in the null space of C and therefore also in the nullspace of $|C|$. It follows then from [\(3.10\)](#page-18-1) that

$$
\mathcal{P}(d\lambda) = 0 \text{ for any function } \lambda : \mathbb{R}^3 \to \mathbb{R}.
$$
 (3.11)

This could also be derived directly from the definition [\(3.9\)](#page-18-0) because the solution to the Maxwell-Poisson equation [\(3.7\)](#page-17-1) is easily verified to be $a(s) =$ $d\lambda$.

On the other hand, for A in Coulomb gauge, [\(3.10\)](#page-18-1) and Definition [2.11](#page-13-5) show that

$$
\mathcal{P}(A) = (|C|A, A)_2 = ||A||_C^2 \quad \text{if } div A = 0.
$$
 (3.12)

In particular if A is in Coulomb gauge then $\mathcal{P}(A) < \infty$ if and only if $A \in$ $H_{1/2}(\mathbb{R}^3)$.

Remark 3.5 A general gauge potential A can be decomposed into its longitudinal part and transverse part:

$$
A = A_{long} + A_{trans}, \qquad A_{long} = d\lambda, \qquad div \, A_{trans} = 0. \tag{3.13}
$$

Its Poisson extension is then given by

$$
\mathsf{a}(s) = A_{long} + e^{-s|C|} A_{trans}.\tag{3.14}
$$

Since $|C|$ has zero nullspace among the transverse potentials, the second term goes to zero as $s \to \infty$. Hence

$$
\lim_{s \to \infty} \mathsf{a}(s) = A_{long}.\tag{3.15}
$$

The Maxwell-Poisson equation therefore filters out the transverse part of A and leaves the longitudinal part (i.e. pure gauge part) in the limit. For a non-abelian gauge field the pure gauge part cannot be separated out from the initial potential A. Instead we will use the non-abelian analog of the Maxwell-Poisson equation to separate out a "pure gauge piece" of the initial data by an analog of [\(3.15\)](#page-19-1).

Remark 3.6 If A is in Coulomb gauge then (3.12) shows that the first term in the Bargmann-Wigner norm [\(2.44\)](#page-14-0) is equal to the Poisson action of A. Since the full Bargmann-Wigner norm is then determined by the dual space norm as in [\(2.44\)](#page-14-0), the Poisson action actually determines the full Lorentz invariant norm on the initial data space for Maxwell's equations.

3.2 Self duality and helicity for the classical electromagnetic field

The four dimensional Euclidean Hodge star operation $*_e$ is given on the four dimensional curvature F defined in (3.2) by

$$
*_e F = ds \wedge * d\mathbf{a} + *\mathbf{a}',\tag{3.16}
$$

wherein ∗ denotes the three dimensional Hodge star operation. Therefore the four dimensional curvature F is self-dual if and only if

$$
ds \wedge \mathbf{a}' + d\mathbf{a} = ds \wedge *d\mathbf{a} + *\mathbf{a}' \quad \text{(self-duality)}.
$$
 (3.17)

That is

$$
a' = *da \qquad \text{(self-duality)}.\tag{3.18}
$$

(Keep in mind that $*^2 = Identity$ over \mathbb{R}^3 .) Similarly F is anti-self-dual if and only if

$$
\mathbf{a}' = - * d\mathbf{a} \quad \text{(anti-self-duality)}.\tag{3.19}
$$

It is clear that [\(3.18\)](#page-20-0) and [\(3.19\)](#page-20-1) each imply the Maxwell-Poisson equation [\(3.7\)](#page-17-1)

Theorem 3.7 Suppose that div $A = 0$ and that $A \in H_{1/2}(\mathbb{R}^3)$. Let $\mathsf{a}(s)$ be the Poisson extension of A. Then

A is in the negative spectral subspace of curl if and only if $a(\cdot)$ is self-dual. A is in the positive spectral subspace of curl if and only if $a(\cdot)$ is antiself-dual.

Proof. Writing $C = \text{curl} = *d$ we may write [\(3.18\)](#page-20-0) and [\(3.19\)](#page-20-1) in the form

$$
a' = Ca \qquad \text{(self-dual)} \tag{3.20}
$$

$$
\mathbf{a}' = -C\mathbf{a} \quad \text{(anti-self-dual)}.\tag{3.21}
$$

If A lies in the negative spectral subspace of curl then $e^{-s|C|}A = e^{sC}A$ and therefore the solution [\(3.8\)](#page-17-2) reduces to $a(s) = e^{sC}A$. Consequently $a'(s) =$ $Ca(s)$ and therefore $a(\cdot)$ is self-dual.

Conversely, suppose that $a(\cdot)$ is self-dual. Then by [\(3.20\)](#page-20-2) we have $a(s)$ = $e^{sC}A$ and therefore $\|\mathbf{a}'(s)\|_{L^2(\mathbb{R}^3)} = \|Ce^{sC}A\|_{L^2(\mathbb{R}^3)}$. If A has a non-zero spectral component for C in the spectral subspace $[\epsilon, \infty)$ with some $\epsilon > 0$ then, by the spectral theorem, $||Ce^{sC}A||_{L^2(\mathbb{R}^3)} \geq ce \exp(\epsilon s)$ for some strictly positive constant c. Therefore $a(\cdot)$ does not have finite Poisson action and A is not in $H_{1/2}$. So A must lie in the spectral subspace $(-\infty, 0]$ for C if a is self-dual.

A similar argument applies to show that the Poisson extension of A is anti-self-dual if and only if A lies in the positive spectral subspace of curl. \blacksquare

The following corollary is the main result of this section.

Corollary 3.8 Suppose that A is in the electromagnetic configuration space ^C and ^a is its Poisson extension. Then

A is in C_+ if and only if a is anti-self-dual.

A is in \mathcal{C}_- if and only if a is self-dual.

Proof. This just restates Theorem [3.7](#page-20-3) in view of Definition [2.11.](#page-13-5) \blacksquare

4 Yang-Mills fields

Both the Lorentz invariant norm and helicity for the electromagnetic field can be characterized in terms of either plane waves, the operator curl, or the Maxwell-Poisson semigroup, as we saw in Sections [2](#page-5-0) and [3.](#page-15-0) Since the classical Yang-Mills equations are non-linear, there is no useful gauge invariant representation of these fields in terms of plane waves. In this section we are going to describe the Yang-Mills analog of the Maxwell-Poisson semigroup and then use it to produce both a Riemannian metric on configuration space and a decomposition of configuration space into positive and negative "helicity" submanifolds. Here is an outline of the transition steps from electromagnetism to Yang-Mills.

i) The norm on the (linear) configuration space of the electromagnetic field will be replaced by a Riemannian metric on the (non-linear) configuration space $\mathcal C$ of the Yang-Mills field. The Yang-Mills-Poisson equation will be instrumental in this construction. (Section [5.](#page-22-0))

ii) The helicity subspaces \mathcal{C}_{\pm} in the linear decomposition [\(2.38\)](#page-13-4) will be replaced by submanifolds \mathcal{C}_{\pm} determined by (anti-)self duality of solutions to the Yang-Mills-Poisson equation. (Section [6.1.](#page-37-1))

iii) The orthogonal projections onto the helicity subspaces \mathcal{C}_\pm in the linear decomposition [\(2.38\)](#page-13-4) will be replaced by flows along orthogonal vector fields over the manifold \mathcal{C} , yielding non-linear analogs of these projections. (Section [6.2.](#page-39-0))

iv) The spectral behavior of the operator curl in the electromagnetic helicity spaces will be replaced by a corresponding behavior of the gauge covariant curl operator acting not on the configuration space $\mathcal C$ but on the tangent spaces $T(\mathcal{C}_\pm)$. (Section [6.2.](#page-39-0))

5 The configuration space for Yang-Mills fields

5.1 The Poisson action and configuration space

Denote by K a compact Lie group contained in the orthogonal or unitary group of a finite dimensional inner product space V. \mathfrak{k} will denote its Lie algebra, on which we assume given an $Ad K$ invariant real inner product $\langle \cdot, \cdot \rangle_{\mathfrak{k}}$. Let

$$
\mathsf{a}(s) := \sum_{j=1}^{3} \mathsf{a}_j(s) dx^j, \quad s \ge 0 \tag{5.1}
$$

be a \mathfrak{k} valued 1-form on \mathbb{R}^3 for each (Euclidean) time $s \geq 0$. Here each coefficient $a_j(x, s)$ lies in $\mathfrak k$. Denote by

$$
\mathsf{b}(s) := d\mathsf{a}(s) + \mathsf{a}(s) \land \mathsf{a}(s)
$$
\n
$$
(5.2)
$$

its three dimensional curvature and denote by $a'(s)$ its derivative with respect to s. We may regard **a** as a 1-form on a half space of \mathbb{R}^4 in temporal gauge. That is, it has no ds component. Its four dimensional curvature is then given by

$$
F = ds \wedge \mathbf{a}'(s) + \mathbf{b}(s). \tag{5.3}
$$

Since the two summands are mutually orthogonal at each point (x, s) we have

$$
|F(x,s)|^2_{\Lambda^2 \otimes \mathfrak{k}} = |\mathsf{a}'(x,s)|^2_{\Lambda^1 \otimes \mathfrak{k}} + |\mathsf{b}(x,s)|^2_{\Lambda^2 \otimes \mathfrak{k}} \tag{5.4}
$$

in analogy with [\(3.5\)](#page-17-0). We are interested in the minimization of the functional

$$
\int_0^\infty \left(\| \mathbf{a}'(s) \|_{L^2(\mathbb{R}^3)}^2 + \| \mathbf{b}(s) \|_{L^2(\mathbb{R}^3)}^2 \right) ds \tag{5.5}
$$

subject to an initial condition $a(0) = A$, where A is a given \mathfrak{k} valued 1-form on \mathbb{R}^3 . In view of [\(5.4\)](#page-22-2) this functional of **a** is also given by the familiar expression (3.3) but with F now given by (5.3) .

The Euler equation for this minimization problem is the Yang-Mills-Poisson equation

$$
\mathsf{a}''(s) = d^*_{\mathsf{a}(s)} \mathsf{b}(s), \qquad \mathsf{a}(0) = A. \tag{5.6}
$$

Here we are using d_A to denote the gauge covariant exterior derivative, which is given on any $\mathfrak k$ valued p-form on $\mathbb R^3$ by $d_A\omega = d\omega + [A \wedge \omega]$, while $d_{\mathsf{a}(s)}^*$: $L^2(\mathbb{R}^3; \Lambda^2 \otimes \mathfrak{k}) \to L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})$ denotes the adjoint of $d_{\mathsf{a}(s)}$ in [\(5.6\)](#page-22-4). The suggestive notation $[u \wedge v]$ for \mathfrak{k} valued forms means to use commutator of the coefficients with wedge product of the differentials. For example $[(\sum_i u_i dx^i) \wedge$ $(\sum_j v_j dx^j)$ = $\sum_{ij} [u_i, v_j] dx^i \wedge dx^j$. Later we will also need to use the adjoint of this product, given by $\langle [u_{\perp} v], \omega \rangle = \langle v, [u \wedge \omega] \rangle$, which must hold for each point in \mathbb{R}^3 and for all $\mathfrak k$ valued forms ω such that $deg u + deg \omega = deg v$. $\langle \cdot, \cdot \rangle$ is the inner product on forms induced by the given inner product on \mathfrak{k} .

The Yang-Mills-Poisson equation [\(5.6\)](#page-22-4) is equivalent to the elliptic Yang-Mills equation for forms in temporal gauge since $d_{a}^{(*e)}F = -a'' + d_{a}^{*}b$, where $*_e$ is the Euclidean Hodge star operator. This is an elliptic non-linear boundary value problem which has been investigated in ground breaking work by A. Marini and T. Isobe. [\[33,](#page-49-4) [34,](#page-49-5) [35,](#page-49-6) [24,](#page-48-3) [36,](#page-49-7) [37,](#page-50-0) [38,](#page-50-1) [25,](#page-48-4) [26\]](#page-48-5). For a bounded open set in \mathbb{R}^4 they have proven existence and non-uniqueness of solutions to the minimization problem if one allows the solution to take on the prescribed boundary values in a weak sense, namely equality only up to local gauge transformations. Our half space problem can be conformally transformed into their setting. But we need a stronger notion of attainment of the boundary value at $s = 0$, namely actual equality. However we can allow a singularity at the point of their boundary corresponding to $s = \infty$. Thus at the present time it is not inconsistent with their work to assume existence and uniqueness for solutions to the Yang-Mills-Poisson equation [\(5.6\)](#page-22-4) under some appropriate conditions on A. We will assume throughout the remainder of this paper that for the potentials A of interest to us existence and uniqueness holds for the boundary value problem [\(5.6\)](#page-22-4) with finite value of the action functional in [\(5.5\)](#page-22-5). In Example [5.4](#page-26-1) we will see how instantons give rise to such solutions.

Notation 5.1 (Poisson action) Define

$$
\mathcal{P}(A) = \int_0^\infty \left(\| \mathbf{a}'(s) \|_{L^2(\mathbb{R}^3)}^2 + \| \mathbf{b}(s) \|_{L^2(\mathbb{R}^3)}^2 \right) ds,\tag{5.7}
$$

wherein **a** is the solution to the boundary value problem (5.6) . We will refer to a as the Poisson extension of A and to $\mathcal{P}(A)$ as the Poisson action of A.

A gauge function $g : \mathbb{R}^3 \to K$ acts on a potential A via the action $A \mapsto A^g := g^{-1}Ag + g^{-1}dg$. The Poisson action is a gauge invariant function of A. That is,

$$
\mathcal{P}(A^g) = \mathcal{P}(A). \tag{5.8}
$$

To see this, suppose that a is a solution to the Yang-Mills-Poisson equation [\(5.6\)](#page-22-4). Then the function

$$
\mathbf{a}^g(s) := g^{-1}\mathbf{a}(s)g + g^{-1}dg\tag{5.9}
$$

is again a solution to [\(5.6\)](#page-22-4), as we can see from the identities $(d/ds)^n a^g(s) =$ $g^{-1}(d/ds)^n \mathsf{a}(s)g, n = 1, 2 \text{ and } d^*_{\mathsf{a}^g(s)} \mathsf{b}^g(s) = g^{-1}\Big(d^*_{\mathsf{a}(s)}b(s)\Big)g. \text{ Of course } \mathsf{a}^g(0) =$ A^g . And since $b^g(s) = g⁻¹b(s)g$ we see also that the functional in [\(5.7\)](#page-23-0) has the same value for a^g . This proves [\(5.8\)](#page-23-1).

As in the electromagnetic case $\mathcal{P}(A)$ is zero on pure gauges. (cf. [\(3.11\)](#page-18-2).) Indeed, if A is a pure gauge, say $A = g^{-1}dg$, then its Poisson extension is given by $a(s) = g^{-1} dg$ for all $s \ge 0$ since both $a'(s) = 0$ and $b(s) = 0$ for this function, which shows that [\(5.6\)](#page-22-4) holds and also that

$$
\mathcal{P}(g^{-1}dg) = 0.\tag{5.10}
$$

We wish to use the function $\mathcal{P}(A)$ to construct the relativistically correct configuration space as we did in the electromagnetic case. Nominally, the configuration space is given by $\mathcal{C} = \mathcal{A}/\mathcal{G}$ for some appropriate set $\mathcal A$ of gauge potentials on \mathbb{R}^3 and corresponding gauge group \mathcal{G} . But [\(5.10\)](#page-24-0) shows that the restriction $\mathcal{P}(A) < \infty$ provides no control over A in some directions (the longitudinal directions). Defining A simply by the condition $\mathcal{P}(A) < \infty$ would lead therefore to an unsatisfactory candidate for A , a circumstance which we avoided in the electromagnetic case by imposing the Coulomb gauge on A. In our non-abelian case we want to avoid such a gauge choice because of the ever lurking Gribov ambiguity. We will instead put an indirect size restriction on the gauge potentials allowed into our space A by making use of the filtering action of the Yang-Mills-Poisson equation.

Let

$$
\|\omega\|_{H_{1/2}} = \|(-\Delta)^{1/4}\omega\|_{L^2(\mathbb{R}^3)}
$$

for any \mathfrak{k} valued 1-form on \mathbb{R}^3 . Here we are using the Laplacian $-\Delta :=$ $d^*d + dd^*$ on \mathfrak{k} valued 1-forms over \mathbb{R}^3 . We have chosen the Sobolev index 1/2 because in the electromagnetic case the size restriction $\mathcal{P}(A) < \infty$ gives exactly $H_{1/2}$ gauge potentials when the Coulomb gauge is imposed, as we saw in [\(3.10\)](#page-18-1). We are going to impose an $H_{1/2}$ restriction on the "filtered" longitudinal part" of A as follows.

Suppose that $a(s)$ is a solution to [\(5.6\)](#page-22-4) and that

$$
\mathsf{a}_{\infty} := \lim_{s \to \infty} \mathsf{a}(s) \tag{5.11}
$$

exists (for example in the sense of $L^2(\mathbb{R}^3)_{loc}$). We saw in [\(3.15\)](#page-19-1) that a_{∞} exists in the electromagnetic case and is precisely the longitudinal part (i.e. the pure gauge part) of the initial potential A.

Notation 5.2 $(\mathcal{R}, \mathcal{A} \text{ and } \mathcal{C})$ Let

$$
\mathcal{R} = \{\text{solutions a to the differential equation in (5.6) such that} \quad \mathcal{P}(\mathsf{a}(0)) + ||\mathsf{a}_{\infty}||_{H_{1/2}}^2 < \infty\}. \tag{5.12}
$$

The gauge potentials of interest to us will be the set of initial values of such solutions:

$$
\mathcal{A} = \{ \mathsf{a}(0) : \mathsf{a} \in \mathcal{R} \}. \tag{5.13}
$$

We will say that a gauge potential A on \mathbb{R}^3 is in the *soft Coulomb gauge* if it lies in A, and in the asymptotic Coulomb gauge if in addition $a_{\infty} = 0$. The examples we construct from instantons below will be in the asymptotic Coulomb gauge.

Having now chosen the space $\mathcal A$ in a way that parrots the electromagnetic case, the gauge group $\mathcal G$ appropriate for this choice must be chosen so as to preserve A as a set and also to be isometric for the as yet to be determined Riemannian metric on \mathcal{A} . If a is the Poisson extension of A then a^g is the Poisson extension of A^g as we have seen in the proof of [\(5.8\)](#page-23-1). Since $\lim_{s\to\infty} \mathsf{a}^g(s) = (\mathsf{a}_{\infty})^g$ we need

$$
\mathsf{a}^g_\infty \in H_{1/2} \quad \text{whenever} \quad \mathsf{a}_\infty \in H_{1/2}
$$

in order for A to be a gauge invariant set under the gauge function g. In view of [\(5.9\)](#page-24-1) we therefore need $g^{-1}dg$ to be itself in $H_{1/2}$. With this as motivation we define

$$
\mathcal{G} = \{\text{gauge functions } g : \mathbb{R}^3 \to K \text{ such that } g^{-1} dg \text{ is in } H_{1/2} \text{ and } g(x) \to I \text{ as } x \to \infty\}. \tag{5.14}
$$

The second condition reflects the fact that we are interested in charge zero. The relation between the behavior at infinity of a gauge function on the one hand and total charge on the other results from the well known relation between gauge invariance and conservation of charge. This will be elaborated on in a future work.

It happens that $\mathcal G$ is a complete topological group in its natural metric under pointwise multiplication. It is also the critical gauge group for three spatial dimensions in the sense that for any $\epsilon > 0$, the set of gauge functions for which $g^{-1}dg$ lies in the (locally) larger space $H_{1/2-\epsilon}$ is no longer a group: It is not closed under multiplication. See [\[19\]](#page-48-6) for further properties and proofs. Combining the set A of " $H_{1/2}$ " gauge potentials with the matching group $\mathcal G$ of " $H_{3/2}$ " gauge functions, we define the configuration space for the non-abelian gauge field to be

$$
C = \mathcal{A}/\mathcal{G}.\tag{5.15}
$$

We will make A into a Riemannian manifold with a G invariant metric in the next two sections. To this end we will gather information on the derivatives of P in the next section. This produces, of course, a Riemannian metric on $\mathcal C$ also.

Remark 5.3 In addition to the fundamental mathematical question as to the existence and uniqueness of solutions to the Yang-Mills-Poisson equation (5.6) , these definitions raise the question as to whether the limit a_{∞} actually exists for some appropriate class of gauge potentials A. For the Yang-Mills heat equation a similar question has been addressed in many works. See e.g. [\[22,](#page-48-7) [21,](#page-48-8) [23\]](#page-48-9), [\[9\]](#page-47-2), [\[28\]](#page-49-8), [\[43\]](#page-50-2) and references therein.

5.2 Restriction of instantons

Example 5.4 (Restriction of instantons) Finite action solutions to the Yang-Mills-Poisson equation can be constructed from instantons by gauge transforming an instanton to the Euclidean temporal gauge and then restricting it to a half-space. Take, for example, $K = SU(2)$. The simplest instanton is given by

$$
A_{\mu}(x) = \sigma_{\mu,\nu} x^{\nu} / (|x|^2 + \rho^2), \quad x = (\mathbf{x}, s) \in \mathbb{R}^4.
$$

where $\sigma_{\mu,\nu} = -\sigma_{\nu,\mu} \in \mathfrak{k} = su(2)$ and $\rho > 0$ and constant. Its curvature is given by

$$
F_{\mu,\nu} = -4\sigma_{\mu,\nu} \frac{\rho^2}{(|x|^2 + \rho^2)^2}.
$$

See e.g. [\[6\]](#page-47-3) or [\[4\]](#page-46-3) or [\[54,](#page-51-0) Eq. (21.26)] or [\[58\]](#page-52-0).

We want to gauge transform A into the Euclidean temporal gauge, i.e., so that its new fourth component is equal to zero. For a gauge function $g: \mathbb{R}^4 \to SU(2)$ we have $(A^g)_4 = g^{-1}A_4g + g^{-1}\partial_s g$. So we want to choose g so that $A_4 + (\partial_s g)g^{-1} = 0$. We may write

$$
A_4(\mathbf{x},s) = \frac{\mathbf{u}}{s^2 + a^2},
$$

where $\mathbf{u} = \sum_{j=1}^{3} \sigma_{4,j} \mathbf{x}^{j} \in \mathfrak{k}$ and $a^2 = |\mathbf{x}|^2 + \rho^2$. The solution which is the identity element of $SU(2)$ at $s = 0$ is given by Bitar and Chang [\[6,](#page-47-3) Equ. (4.4)] as $g_{bc}(\mathbf{x}, s) = \exp(-(\mathbf{u}/a)tan^{-1}(s/a))$. For $0 \leq s < \infty$, $g_{bc}(\cdot, s)$ lies in our critical gauge group \mathcal{G} , but $g_{bc}(\cdot,\infty)$ does not. We need to use the solution which is the identity element at $s = \infty$. It is given by

$$
g(\mathbf{x}, s) = \exp\left(\frac{\mathbf{u}}{a}\left\{(\pi/2) - \tan^{-1}(s/a)\right\}\right), \quad (\mathbf{x}, s) \in \mathbb{R}^4.
$$

Then $(A^g)_4 = 0$ and

$$
\mathbf{a}_{j}(\mathbf{x}, s) \equiv (A^{g})_{j}(\mathbf{x}, s)
$$

= $g(\mathbf{x}, s)^{-1} A_{j}(\mathbf{x}, s) g(\mathbf{x}, s) + g(\mathbf{x}, s)^{-1} \partial_{j} g(\mathbf{x}, s), j = 1, 2, 3.$

Since $g(\mathbf{x}, \infty) = I$ and $A_j(\mathbf{x}, \infty) = 0$ we see that $(A^g)_j(\mathbf{x}, \infty) = 0$. So A^g is in Euclidean temporal gauge and also in asymptotic Coulomb gauge. Since it is the four dimensional gauge transform of an instanton it satisfies $d_{A^g}^{*_e}F(A^g) = 0$, which is Poisson's equation [\(5.6\)](#page-22-4) for a. Its curvature is square integrable over all of \mathbb{R}^4 and therefore over the half-space \mathbb{R}^4_+ . Hence a has finite action and is in asymptotic coulomb gauge. In particular its initial value, $a(0)$, lies in A.

Since, by definition, an instanton has finite action, this construction applies to any instanton as long as the time dependent gauge transformation $g(\mathbf{x}, s)$ needed to transform the instanton to temporal gauge can be chosen to be the identity of the group K at $s = \infty$.

5.3 Derivatives of the Poisson action.

Notation 5.5 (Variational equation) If $A(t)$ is a time dependent family of gauge potentials on \mathbb{R}^3 and if $u := (d/dt)A(t)\Big|_{t=0}$ then u is a \mathfrak{k} valued 1-form on \mathbb{R}^3 and measures the variation of $A(t)$ at $t = 0$. For any function f on A

the derivative of f at A in the direction u is given, in accordance with the chain rule, by

$$
(\partial_u f)(A) = (d/dt) f(A(t))|_{t=0}, \tag{5.16}
$$

where $A = A(0)$.

Suppose that for each t the Poisson extension of $A(t)$ is $a_t(s)$. The Poisson equation $a''_t(s) = d^*_{a_t(s)} b_t(s)$ can be differentiated with respect to t at $t = 0$ to find the variational equation associated to the variation u. Define $u(s) =$ $(\partial/\partial t)\mathsf{a}_t(s)\Big|_{t=0}$. Then we find for $\mathsf{u}(s)$ the variational equation

$$
\mathbf{u}''(s) = d^*_{\mathbf{a}(s)} d_{\mathbf{a}(s)} \mathbf{u}(s) + [\mathbf{u}(s) \lrcorner \ \mathbf{b}(s)]. \tag{5.17}
$$

Here $a(s) \equiv a_0(s)$ is the Poisson extension of $A \equiv A(0)$ and $u : [0, \infty) \rightarrow$ $L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})$ is a function satisfying $\mathsf{u}(0) = u$ in addition to [\(5.17\)](#page-28-0). The last term in [\(5.17\)](#page-28-0) has been defined in the paragraph after Eq. [\(5.6\)](#page-22-4). In the following we will always assume that for a variation u of interest to us there is a unique solution to the variational equation (5.17) with initial value u and such that $u(s) \to 0$ as $s \to \infty$. We will refer to $u(s)$ as the *Poisson extension* of u along a.

We can compute the derivative of the Poisson action as follows.

Lemma 5.6 Let u be a variation of a point A in \mathcal{A} , i.e. u is a \mathfrak{k} valued 1-form on \mathbb{R}^3 . If $a(s)$ is the Poisson extension of A then

$$
\partial_u \mathcal{P}(A) = -2(u, \mathbf{a}'(0))_{L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})}.\tag{5.18}
$$

Proof. Denote by $u(s)$ the Poisson extension of u. Suppose that $A(t)$ is a curve with $A(0) = A$ and $\left(\frac{d}{dt}A(t)\right)|_{t=0} = u$ and that $a_t(s)$ is the Poisson extension of $A(t)$. Then we have, at $t = 0$,

$$
(d/dt) \|\mathbf{a}'_t(s)\|^2 = 2(\mathbf{a}'(s), \mathbf{u}'(s)) \text{ and } (d/dt) \|\mathbf{b}_t(s)\|^2 = 2(\mathbf{b}(s), d_{\mathbf{a}(s)}\mathbf{u}(s)).
$$

Therefore, using the Poisson equation [\(5.6\)](#page-22-4) in the third line below, we find

$$
\partial_u \mathcal{P}(A) = 2 \int_0^\infty \left((\mathbf{a}'(s), \mathbf{u}'(s)) + (\mathbf{b}(s), d_{\mathbf{a}(s)} \mathbf{u}(s)) \right) ds \tag{5.19}
$$

= $2 \int_0^\infty \left(\left\{ (d/ds)(\mathbf{a}'(s), \mathbf{u}(s)) \right\} - (\mathbf{a}''(s), \mathbf{u}(s)) + (d_\mathbf{a}(s)^* \mathbf{b}(s), \mathbf{u}(s)) \right\} ds$
= $2(\mathbf{u}(s), \mathbf{a}'(s)) \Big|_0^\infty$
= $-2(\mathbf{u}(0), \mathbf{a}'(0)).$ (5.19)

Remark 5.7 (Longitudinal and transverse) The decomposition of an electromagnetic field into longitudinal and transverse parts is not a gauge invariant decomposition. Nor is there any gauge invariant analog of this decomposition for non-abelian gauge fields. But for the tangent spaces to the manifold A there is a well known gauge invariant decomposition analogous to the Coulomb gauge. The terminology, vertical and horizontal, for the differential analogs of longitudinal and transverse goes back to Ambrose and Singer [\[2\]](#page-46-4). This will be reviewed here in our special context to help establish notation.

Notation 5.8 (Vertical and horizontal) If $\lambda : \mathbb{R}^3 \to \mathfrak{k}$ is a reasonable function then $q(t, x) := \exp(t\lambda(x))$ defines a curve $t \to q(t)$ in the gauge group G. Given a point $A \in \mathcal{A}$ the curve $t \mapsto A^{g(t)}$ is a curve in A. Since, at $t = 0$ one has $(d/dt)A^{g(t)} = [A, \lambda] + d\lambda$, the tangent to this curve in A at $t = 0$ is given by the 1-form $d_A \lambda := d\lambda + [A, \lambda]$. The vertical subspace of $T_A(\mathcal{A})$ is by definition the set of tangent vectors to these curves at A. They consist therefore of $\mathfrak k$ valued 1-forms $d_A\lambda$ with λ running over the Lie algebra of G. The subspace of L^2 orthogonal to the vertical subspace is called the *horizontal* subspace. Thus a $\mathfrak k$ valued 1-form u on \mathbb{R}^3 is

vertical at A if
$$
u = d_A \lambda
$$
 for some scalar function λ . (5.20)
horizontal at A if $d_A^* u = 0$. (5.21)

This is the standard terminology associated to the Coulomb connection for the G bundle A , [\[56\]](#page-51-1). In this discussion we have made the usual identification of a tangent vector to A with a $\mathfrak k$ valued 1-form on $\mathbb R^3$. The corresponding action of a gauge function on such a form is given by $u \mapsto u^g := g^{-1}ug$.

Lemma 5.9 If $f : A \to \mathbb{C}$ is gauge invariant then

$$
\partial_v f(A) = 0 \tag{5.22}
$$

for any vertical vector $v = d_A \lambda$.

Proof. Let $g(t, x)$ = $\exp(t\lambda(x))$. By assumption $f(A^{g(t)}) = f(A)$. But at $t = 0$, $(d/dt)A^{g(t)} = v$. Therefore $(\partial_v f)(A) = 0$ by the chain rule.

Corollary 5.10 If $a(\cdot)$ is a solution to the Yang-Mills-Poisson equation of finite action and $a(0) = A$ then $a'(0)$ is horizontal at A. That is,

$$
d_A^* \mathbf{a}'(0) = 0. \tag{5.23}
$$

Moreover

$$
d_{\mathsf{a}(s)}^* \mathsf{a}'(s) = 0 \quad \forall s \ge 0. \tag{5.24}
$$

Proof. Since $\mathcal P$ is gauge invariant Lemma [5.9](#page-29-0) shows that $\partial_v \mathcal P(A) = 0$ for any vertical vector v, say $v = d_A \lambda$. But the derivative of P is given by [\(5.18\)](#page-28-1), from which we can conclude that $(d_A\lambda, \mathsf{a'}(0))_{L^2} = 0$ for all λ . That is, $(\lambda, d_A^* \mathsf{a}'(0))_{L^2} = 0$ for all λ . Hence [\(5.23\)](#page-29-1) holds. Now for any $\mathfrak k$ valued 2-form ω we have the identity $(d_A^*)^2 \omega = [B \cup \omega]$, which follows by duality from the Bianchi identity $(d_A)^2 \phi = [B, \phi]$ for scalar functions ϕ . Hence

$$
(d/ds)\left(d_{\mathsf{a}(s)}^*\mathsf{a}'(s)\right) = d_{\mathsf{a}(s)}^*\mathsf{a}''(s) + [\mathsf{a}'(s)\mathsf{a}'(s)]
$$

$$
= d_{\mathsf{a}(s)}^*d_{\mathsf{a}(s)}^*\mathsf{b}(s) + 0
$$

$$
= [\mathsf{b}(s)\mathsf{a}]\mathsf{b}(s)]
$$

$$
= 0.
$$

Therefore $d_{\mathsf{a}(s)}^* \mathsf{a}'(s)$ is constant in s. Since, by [\(5.23\)](#page-29-1), it is zero at $s = 0$ it is identically zero. \blacksquare

Theorem 5.11 (Second derivatives of the Poisson action) Suppose that A has finite Poisson action and that u and v are $\mathfrak k$ valued 1-forms on $\mathbb R^3$. Denote by a the Poisson extension of A and by μ , ν the respective Poisson extensions of u and v along a. Then

$$
(1/2)\partial_v \partial_u \mathcal{P}(A) = \int_0^\infty \left\{ \left(\mathbf{u}'(s), \mathbf{v}'(s) \right)_{L^2} + \left(d_{\mathbf{a}(s)} \mathbf{u}(s), d_{\mathbf{a}(s)} \mathbf{v}(s) \right)_{L^2} + \left([\mathbf{u}(s) \wedge \mathbf{v}(s)], \mathbf{b}(s) \right)_{L^2} \right\} ds \tag{5.25}
$$

and also

$$
-(1/2)\partial_v \partial_u \mathcal{P}(A) = (u, v'(0)) = (v, u'(0)).
$$
\n(5.26)

In particular,

$$
(1/2)\partial_u^2 \mathcal{P}(A) = \int_0^\infty \left\{ \|u'(s)\|_{L^2}^2 + \|d_{a(s)}u(s)\|_2^2 + \left([u(s) \wedge u(s)], b(s) \right) \right\} ds
$$
\n(5.27)

and also

$$
\partial_u^2 \mathcal{P}(A) = -2(u, u'(0)) = -(d/ds) ||u(s)||_{L^2}^2 \Big|_{s=0}.
$$
 (5.28)

Proof. Differentiate (5.18) with respect to A in the direction v to find

$$
-(1/2)\partial_v \partial_u \mathcal{P}(A) = \left(u, v'(0)\right)_{L^2}.
$$
\n(5.29)

Using the variational equation [\(5.17\)](#page-28-0) we get

$$
(d/ds)(u(s), v'(s))
$$

= (u(s), v''(s)) + (u'(s), v'(s))
= (u(s), d_a^*d_a v(s) + [v(s) \cup b(s)]) + (u'(s), v'(s))
= (d_a u(s), d_a v(s)) + (u(s), [v(s) \cup b(s)]) + (u'(s), v'(s))
= (d_a u(s), d_a v(s)) + ([v(s) \wedge u(s)], b(s)) + (u'(s), v'(s)). \t(5.30)

Assuming that $(u(s), v'(s))$ goes to zero as $s \to \infty$ we may integrate this identity over $[0, \infty)$ to find, using $[u \wedge v] = [v \wedge u]$,

$$
-(\mathsf{u}(0),\mathsf{v}'(0))\tag{5.31}
$$

= $\int_0^\infty \left\{ (d_\mathsf{a}\mathsf{u}(s), d_\mathsf{a}\mathsf{v}(s)) + ([\mathsf{u}(s) \wedge \mathsf{v}(s)], \mathsf{b}(s)) + (\mathsf{u}'(s), \mathsf{v}'(s)) \right\} ds.$

 (5.25) and (5.26) now follow from (5.29) , (5.31) and the symmetry in u and v in [\(5.31\)](#page-31-1). Set $v = u$ in [\(5.25\)](#page-30-0) and [\(5.26\)](#page-30-1) to find [\(5.27\)](#page-30-2) and [\(5.28\)](#page-30-3).

Remark 5.12 (Expansion of P near zero.) The behavior of $P(A)$ for small A is given by

$$
\mathcal{P}(u) = (|C|u, u)_{L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})} + O(u^3), \tag{5.32}
$$

where u is a small variation of A at $A = 0$. To see this observe that the Poisson extension of $A = 0$ is $a(s) = 0$. The variational equation [\(5.17\)](#page-28-0) is therefore $u''(s) = d^* du(s)$, which is the Maxwell-Poisson equation for the $\mathfrak k$ valued 1-form $u(s)$. The solution, as in the real valued case, [\(3.8\)](#page-17-2), is $u(s) = e^{-s|C|}u$, where C is the curl acting on each \mathfrak{k} component of u. Hence $u'(0) = -|C|u$. We find therefore, $\mathcal{P}(0) = 0$, $(\partial_u \mathcal{P})(0) = -2(u, a'(0)) = 0$ and $(1/2)(\partial_u^2 \mathcal{P})(0) = (|C|u, u)$, from which [\(5.32\)](#page-31-2) follows.

Remark 5.13 It may be useful to observe that $a'(s)$ is itself a variational field along a, and if u is also a variational field along a then

$$
(\mathsf{u}(0),\mathsf{a}''(0)) = (\mathsf{u}'(0),\mathsf{a}'(0)).\tag{5.33}
$$

Proof. Let $v(s) = a'(s)$. Then

$$
\mathbf{v}''(s) = (d/ds)\mathbf{a}''(s)
$$

= $(d/ds)d_{\mathbf{a}}^*\mathbf{b}$
= $d_{\mathbf{a}}^*d_{\mathbf{a}}\mathbf{a}'(s) + [\mathbf{a}'(s) \bot \mathbf{b}(s)]$
= $d_{\mathbf{a}}^*d_{\mathbf{a}}\mathbf{v}(s) + [\mathbf{v}(s) \bot \mathbf{b}(s)].$

Therefore a' is a variational field along a. Upon taking $v(s) = a'(s)$ in [\(5.26\)](#page-30-1) we find (5.33) .

5.4 The Riemannian metrics on A and C

We want to define a Riemannian metric on A which is gauge invariant, captures a non-linear version of the electromagnetic norm identity [\(3.12\)](#page-19-2), and, unlike [\(3.12\)](#page-19-2), does not require the Coulomb gauge. We aim to do this by using the first variation of the Poisson action to define a norm on horizontal vectors and then using an explicitly defined norm on vertical vectors that's commensurate with our choice of gauge group, \mathcal{G} .

Notation 5.14 (Riemannian metric on A) Suppose that A is in A and that **a** is its Poisson extension. If u is any \mathfrak{k} valued 1-form on \mathbb{R}^3 and u is its Poisson extension along a define

$$
{A}||u||^{2} = \int{0}^{\infty} (||u'(s)||_{2}^{2} + ||d_{a(s)}u(s)||_{2}^{2}) ds.
$$
 (5.34)

Suppose that w is a $\mathfrak k$ valued 1-form on $\mathbb R^3$ which decomposes into horizontal and vertical components:

$$
w = u + v, \quad d_A^* u = 0, \quad v = d_A \lambda.
$$

Define

$$
||w||_A^2 = ||u + v||_A^2 = A||u||^2 + ||(-\Delta_A)^{1/4}v||_{L^2(\mathbb{R}^3)}^2,
$$
\n(5.35)

where $\Delta_A = \sum_{j=1}^3 (\partial_j^A)^2$ and $\partial_j^A = \partial_j + [A_j, \cdot]$. We define the horizontal subspace at A by

$$
u \in H_A \quad \text{if} \quad \begin{cases} d_A^* u = 0 \quad \text{and} \\ \|u\|_A < \infty. \end{cases} \tag{5.36}
$$

It seems likely that the norm $||u||_A$ is non-degenerate on H_A and we will assume this in the following. But the integral in [\(5.34\)](#page-32-1) is meaningful even if u is not horizontal at A. However $_A||u||$ is not a norm since it can be degenerate on non-horizontal elements, as will be shown in the next example. For this reason we have added the second term in [\(5.35\)](#page-32-2). We will see that it is commensurate with our choice of \mathcal{G} , defined in [\(5.14\)](#page-25-0).

In the case of electromagnetism, where $K = U(1)$, [\(3.12\)](#page-19-2) shows that $\mathcal{P}(A) = ||A||_{H_{1/2}}^2$ when A is in Coulomb gauge. Identifying, as usual, the tangent space to A with A itself in this linear theory, we see that the norm $||u||_A$ in [\(5.35\)](#page-32-2) reduces to the $H_{1/2}$ norm of electromagnetism when u is horizontal. One should think of $||u||_A$ as an " $H_{1/2}$ "-like norm on H_A in the non-abelian case.

Example 5.15 (Degeneracy) If $A = 0$ then, for any \mathfrak{k} valued function λ on \mathbb{R}^3 , $d\lambda$ is a vertical vector at A and we have

$$
A||d\lambda|| = 0.\tag{5.37}
$$

This is easily seen because the Poisson extension of $A := 0$ is $a(s) \equiv 0$. Therefore if we define $u(s) = d\lambda$ for all $s \geq 0$ then $u'(s) = u''(s) = 0$, while $d^*d\mu(s) = d^*d^2\lambda = 0$ and $b(s) = d^2\lambda = 0$, from which we see that the variational equation [\(5.17\)](#page-28-0) is satisfied. Since also $u(0) = d\lambda$, $u(s)$ is the Poisson extension of $d\lambda$. But then the right hand side of [\(5.34\)](#page-32-1) is zero.

Lemma 5.16 (Gauge invariance of the metric) Let $w \in T_A(\mathcal{A})$ and let $g \in$ G. Suppose that $w = u + v$ with u horizontal at A and v vertical at A. Then $w^g = u^g + v^g$ with u^g horizontal at A^g and v^g vertical at A^g . Moreover

$$
||w^g||_{A^g} = ||w||_A \tag{5.38}
$$

Proof. Horizontal and vertical are preserved under gauge transformations because $d^*_{A^g}u^g = (d^*_Au)^g$ and $(d_A\lambda)^g = d_{A^g}\lambda^g$. In view of the definition [\(5.35\)](#page-32-2) it suffices to show that (5.38) holds for u and v separately.

Concerning the horizontal component, let a be the Poisson extension of A and let u be the Poisson extension of u along a. Then a^g is the Poisson extension of A^g , as we saw in Section [\(5.1\)](#page-22-1), and $u^g(s) := g^{-1}u(s)g$ is the Poisson extension of u^g , as one sees by a similar argument. Since $||(u^g)'(s)||_2 = ||g^{-1}u'(s)g||_2 = ||u'(s)||_2$ and $d_{a^g(s)}u^g(s) = g^{-1}d_{a(s)}u(s)g$, [\(5.38\)](#page-33-0) follows from (5.34) for the horizontal component of w.

Concerning the vertical component, observe that the gauge covariant Laplacian Δ_A commutes with gauge transform in the sense that $\Delta_{A^g} v^g =$ $g^{-1}(\Delta_A v)g$ for any $\mathfrak k$ valued 1-form v. Since $Ad g^{-1}$ is unitary in $L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak k)$, it follows from the spectral theorem that $(-\Delta_{A^g})^{1/4}v^g = g^{-1}((-\Delta_A)^{1/4}v)g$. Therefore $\|(-\Delta_{A^g})^{1/4}v^g\|_{L^2} = \|(-\Delta_A)^{1/4}v\|_{L^2}$.

Corollary 5.17

$$
|\partial_u \mathcal{P}(A)| \le 2\sqrt{\mathcal{P}(A)} \|u\|_A, \quad u \in H_A(\mathcal{A})
$$
\n(5.39)

$$
|(u, \mathsf{a}'(0))| \le \sqrt{\mathcal{P}(A)} \|u\|_{A}, \quad u \in H_A(\mathcal{A}). \tag{5.40}
$$

Proof. From (5.19) and the Schwarz inequality we see that

$$
|\partial_u \mathcal{P}(A)| \le 2\sqrt{\mathcal{P}(A)} \sqrt{\int_0^\infty \left(||\mathbf{u}'(s)||_2^2 + ||d_{\mathbf{a}(s)}\mathbf{u}(s)||_2^2 \right) ds},\tag{5.41}
$$

which is (5.39) . (5.40) follows from (5.18) .

Note: Both inequalities hold for all $u \in T_A$ because $\partial_v \mathcal{P}(A) = 0$ for vertical v , by Lemma [5.9.](#page-29-0)

Notation 5.18 (Riemannian metric on C) Since G acts isometrically on A it induces a metric on $\mathcal{C} := \mathcal{A}/\mathcal{G}$, which makes the projection into a submersion of Riemannian manifolds. We take this induced metric as the Riemannian metric on \mathcal{C} .

Remark 5.19 (Kinematic and dynamic norms) A point q in configuration space is a G orbit, $q := AG$, through a point in A. The tangent space to such an orbit is the set of vertical vectors $d_A\lambda$ at A and the tangent space to C at the orbit q can be identified with the set of horizontal vectors at A . Since the norm defined in (5.35) is gauge invariant in the sense of (5.38) , it descends to a norm on $T_q(\mathcal{C})$. But the simple L^2 norm $||u||_{L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})}$ on H_A is also gauge invariant and also descends to a norm on $T_q(\mathcal{C})$. Unlike the first norm the L^2 norm does not depend in any way on the dynamics, which was used e.g. in (5.6) . The L^2 norm is entirely kinematic and should appropriately be referred to as the kinematic Riemannian metric for configuration space, in contrast to the dynamic Riemannian metric [\(5.35\)](#page-32-2). We already saw in the electromagnetic case that the norm on phase space [\(2.15\)](#page-7-5), induced from the dynamic norm on configuration space by the duality set up by the kinematic norm, gives the unique Lorentz invariant norm. Both the kinematic and dynamic norms thereby appear naturally in considerations of phase space. The kinematic metric is needed to construct the Coulomb connection. See e.g. [\[55,](#page-51-2) [56,](#page-51-1) [40\]](#page-50-3). Other Sobolev norms have been used in some works to ensure that only irreducible connections need be considered, thereby ensuring in turn that Green functions exist. See e.g. [\[39,](#page-50-4) [3\]](#page-46-5). These strong norms enter in a technical way only.

5.5 Phase space

The phase space for the gauge field is $T^*(\mathcal{C})$, as usual in classical mechanics. The natural Riemannian metric on $\mathcal C$ to use here is the one defined in Notation [5.18,](#page-34-2) which gives the Lorentz invariant norm in the electromagnetic case. An electric field is a member of $T^*_A(\mathcal{A})$ for some point $A \in \mathcal{A}$. In the electromagnetic case the space $T_A^*(\mathcal{A})$ can be taken to be independent of A because A is a linear space. The electric field can therefore be, and customarily is, taken to be independent of A. But not in a non-abelian gauge theory.

If the electric field E annihilates all vertical vectors at A, that is, $\langle d_A \lambda, E \rangle_{L^2(\mathbb{R}^3)} =$ 0 for all $\mathfrak k$ valued scalar functions λ , then $d_A^*E = 0$. Otherwise one has $d_A^* E = -4\pi \rho$ for some $\mathfrak k$ valued charge distribution ρ . One can see already from this that charge is connected to vertical vectors by some duality and therefore also to the dual space of the Lie algebra of the gauge group. At an informal level this statement just reflects the well recognized fact that conservation of charge is a consequence of gauge invariance. See e.g. [\[46,](#page-50-5) pages 210-211]. But the charge will not show up in the structure of $T^*(\mathcal{C})$ because the projection of E into $T^*(\mathcal{C})$ depends on evaluation of E on horizontal vectors only. To incorporate charge, from either external sources or additional fields, one must make further use of A and the gauge group G , beyond their roles in forming the configuration space \mathcal{A}/\mathcal{G} . The incorporation of charge into our phase space framework will be addressed quantitatively in a future work. But in the present paper we are only concerned with helicity and therefore we will always take the charge to be zero.

As already noted in Remark [2.13,](#page-14-5) only the decomposition of configuration space must be specified in order to define helicity, whether in the classical or quantum theory, because the associated decomposition of $T^*(\mathcal{C})$ is automatically determined by this decomposition in the classical theory and is not needed in the quantum theory. In the remainder of this paper we will be concerned only with decomposition of the non-abelian configuration space C into submanifolds C_{\pm} analogous to those of the electromagnetic case. We limit further discussion of phase space to the following remarks.

Remark 5.20 (Lorentz invariance) In the case of electromagnetism we saw how the phase space norm gives the Bargmann-Wigner Lorentz invariant norm. (cf. (2.21) .) For non-abelian fields the configuration space C is not linear and one must replace what was previously a norm on $\mathcal{C} \oplus \mathcal{C}^*$ (namely (2.21)) by a Riemannian metric on $T^*(\mathcal{C})$. This space automatically inherits a Riemannian metric from the Riemannian metric on \mathcal{C} , defined in Notation [5.18.](#page-34-2) It remains to be seen whether this Riemannian metric is invariant under the action of the Lorentz group. Invariance under spatial rotations and spatial translations is clear. But invariance under time translation is not. It is not at all clear that the Riemannian metric on $T^*(\mathcal{C})$ needs to be Lorentz invariant in order for the configuration space itself to play its usual role in the quantized theory, where quantum time evolution is not so simply related to the non-linear classical evolution.

Remark 5.21 (History, Symplectic form, Complex structure) Candidates for the phase space associated to linear and non-linear wave equations have been investigated from many different viewpoints. One expects a symplectic structure on the phase space and in some cases one can hope also for a Riemannian structure on this infinite dimensional manifold. In field theory the symplectic structure is an infinite dimensional analog of the canonical 2-form $\sum_{j=1}^{n} dp_j \wedge dq_j$. There is a large literature devoted to the role of the symplectic structure in quantization of the field theory. See e.g. [\[47,](#page-50-6) [52,](#page-51-3) [44,](#page-50-7) [48,](#page-51-4) [49,](#page-51-5) [51,](#page-51-6) [45,](#page-50-8) [59,](#page-52-1) [12,](#page-47-4) [14,](#page-47-5) [13,](#page-47-6) [15,](#page-47-7) [16\]](#page-48-10) for samples. The phase space is usually not chosen in a quantitative way, however, as we have done above. But the references [\[44,](#page-50-7) [48,](#page-51-4) [49,](#page-51-5) [45\]](#page-50-8) do discuss the symplectic form in combination with a Riemannian metric on the phase space. See especially [\[49\]](#page-51-5) for a discussion of phase space for Yang-Mills fields and a proposed quantitative candidate for it. In the context of the Yang-Mills theory the symplectic form is given by the 2-form,

$$
\omega \langle X_1, X_2 \rangle = \int_{\mathbb{R}^3} \left(\langle u_1(x), v_2(x) \rangle - \langle u_2(x), v_1(x) \rangle \right) d^3x \tag{5.42}
$$

where $X_j := u_j, v_j, j = 1, 2$ are two elements of $T_{A,E}T^*(\mathcal{A})$ and $\langle u(x), v(x) \rangle =$ $\langle u(x), v(x)\rangle_{\Lambda^1\otimes \mathfrak{k}}$. If A evolves by the hyperbolic Yang-Mills equation and $X_i(t)$ evolves by the induced flow then it is known that

$$
(d/dt)\omega\langle X_1(t), X_2(t)\rangle = 0.
$$

Moreover ω is gauge invariant, Lorentz invariant and descends to a nondegenerate symplectic form on $T^*(\mathcal{C})$. See e.g. [\[47\]](#page-50-6)-[\[52\]](#page-51-3) and especially [\[49\]](#page-51-5) for early discussions of these structures. The 2-form ω is continuous in the u_i and v_k because they are in dual spaces. But our particular choice of the metric on configuration space gives the Sobolev indices $\pm 1/2$ for these spaces, which match fortuitously with the expectation that the electric field should be one derivative less regular than the potential.

The complex structure on the linear phase space for electromagnetism, used in the proof of Theorem [2.9,](#page-10-2) has a natural analog in the non-abelian theory. If u varies A and v varies E then the map $j : \{u, v\} \mapsto {\hat{v}, -\hat{u}}$, where $T_{A,E} \leftrightarrow T_{A,E}^*$ is the natural isometry, generalizes the electromagnetic complex structure given in the proof of Theorem [2.9.](#page-10-2)

6 Decomposition of Yang-Mills configuration space by (anti-) self-duality

6.1 Definition of \mathcal{C}_\pm

The four dimensional curvature of the 1-form a on the half space $\mathbb{R}^3 \times [0, \infty)$ is given by [\(5.3\)](#page-22-3). Its Euclidean dual is given by $*_e F = ds \wedge *b(s) + *a'(s)$, where ∗ is the three dimensional Hodge star operator. As in the discussion in Section [3.2](#page-19-0) the conditions for Euclidean self-duality or anti-self-duality, $*_eF = \pm F$, of the curvature of $a(\cdot)$ can be written in terms of a as follows.

Definition 6.1 A function $\mathsf{a}: (0, \infty) \to L^2(\mathbb{R}^3; \Lambda^1 \otimes \mathfrak{k})$ is self-dual if

$$
\mathsf{a}'(s) = * \mathsf{b}(s), \qquad 0 < s < \infty.
$$

It is anti-self-dual if

$$
\mathsf{a}'(s) = - * \mathsf{b}(s), \qquad 0 < s < \infty.
$$

The following lemma restates for temporal gauge a well known fact.

Lemma 6.2 If $a(\cdot)$ is self-dual or anti-self dual then it satisfies the Yang-Mills-Poisson equation [\(5.6\)](#page-22-4).

Proof. If $a'(s) = *b(s)\epsilon$ with $\epsilon = \pm 1$ then

$$
\begin{aligned} \mathsf{a}'' &= (d/ds) * \mathsf{b}(s)\epsilon \\ &= *d_{\mathsf{a}(s)}\mathsf{a}'(s)\epsilon \\ &= *d_{\mathsf{a}(s)} * \mathsf{b}(s)\epsilon^2 \\ &= d_{\mathsf{a}(s)}^*\mathsf{b}(s) \end{aligned}
$$

since $*d_{\mathsf{a}(s)} * = d^*_{\mathsf{a}(s)}$ when acting on two forms.

Lemma 6.3 Let $g : \mathbb{R}^3 \to K$ be a gauge function. If a is $(anti-)self-dual$ then so is a^g .

Proof. Since the three dimensional curvature of $a^g(s)$ is $g^{-1}b(s)g$ we have, for $\epsilon = \pm 1$,

$$
(d/ds)\mathbf{a}^g(s) = g^{-1}\mathbf{a}'(s)g = g^{-1} * \mathbf{b}(s)\epsilon g = *g^{-1}\mathbf{b}(s)g\epsilon = *Curv\,\,\mathbf{a}^g(s)\,\epsilon.
$$

where Curv means the three dimensional curvature.

Notation 6.4 $(\mathcal{R}_{\pm}, \mathcal{A}_{\pm}, \mathcal{C}_{\pm})$ Denote by \mathcal{R}_{+} the set of anti-self-dual solutions in R and by R_− the set of self-dual solutions in R. \mathcal{R}_+ and R_− are each gauge invariant sets by Lemma [6.3.](#page-38-0) Let

$$
\mathcal{A}_+ := \{ \mathsf{a}(0) : \mathsf{a} \in \mathcal{R}_+ \},\tag{6.1}
$$

$$
\mathcal{A}_{-} := \{ \mathsf{a}(0) : \mathsf{a} \in \mathcal{R}_{-} \}. \tag{6.2}
$$

 \mathcal{A}_{\pm} are gauge invariant sets because if a is the Poisson extension of A then a^g is the Poisson extension of A^g . We define submanifolds of configuration space by

$$
\mathcal{C}_{+} = \mathcal{A}_{+} / \mathcal{G}, \quad \mathcal{C}_{-} = \mathcal{A}_{-} / \mathcal{G}
$$
\n
$$
(6.3)
$$

The definitions of \mathcal{C}_\pm given in Notation [6.4](#page-38-1) in the non-abelian case are precisely analogous to those for the electromagnetic field and reduce to them when $K = U(1)$. We saw that in the electromagnetic case these manifolds control helicity in the sense that a solution to Maxwell's equations with initial data in $T^*(\mathcal{C}_+)$ has only positive helicity plane waves in its plane wave decomposition. Similarly for $T^*(\mathcal{C}_-)$. In the next section and in [\[17\]](#page-48-0) we will provide further justification for the interpretation of the manifolds \mathcal{C}_\pm as characterizers of helicity in the non-abelian case .

Remark 6.5 In the Riemannian metric on $\mathcal A$ defined in Section [5.4](#page-32-0) the subsets \mathcal{A}_{\pm} are closed in \mathcal{A} . Since $\mathcal G$ acts as isometries on $\mathcal A$ the quotient spaces \mathcal{C}_{\pm} are also closed submanifolds of \mathcal{C} .

6.2 Flows of the helicity vector fields and positivity of $curl_A$

We will give support in this section for the interpretation of the submanifolds \mathcal{C}_{\pm} as the non-abelian analogs of the electromagnetic helicity subspaces. For non-abelian fields we have used (anti-)self duality to define the submanifolds C_{+} in Section [6.1.](#page-37-1) Now we are going to show how this decomposition of configuration space relates to the signature of the covariant curl. We will also show that the orthogonal direct sum [\(2.38\)](#page-13-4), that holds in electromagnetism, has an analogous product decomposition in the non-abelian case.

The transition from the electromagnetic linear space theory to the Yang-Mills non-linear theory will be guided by the following observations, which will gradually shift the paradigm from functional analysis over the electromagnetic Hilbert space $\mathcal C$ to infinite dimensional differential geometry over the analogous non-abelian configuration space. Denote again by C the operator curl.

a. In electromagnetism the positive spectral subspace of C is the null space of the operator $|C| - C$.

b. The function

$$
h_{+}(A) := -(|C| - C)A \tag{6.4}
$$

can be interpreted as a vector field on the electromagnetic Hilbert space C. As such, it has a flow $\exp(th_+)$. It will be shown that the flow converges as $t \to \infty$ to the orthogonal projection P_+ of C onto C_+

c. The operator |C|, whose definition as $\sqrt{C^2}$ has a functional analytic character, can be replaced in electromagnetism by use of the Maxwell-Poisson semigroup, which has a PDE character. Indeed we see from [\(3.8\)](#page-17-2) that $|C|A =$ $-a'(0)$, where $a(s)$ is the solution to the Maxwell-Poisson equation with initial value A.

d. The second term in the vector field $h_{+}(A)$ in [\(6.4\)](#page-39-1) is just the magnetic field $curl A$ in its usual representation as a 1-form (which is better written as $*B$ with $B = dA$). Thus the vector field h_+ can be specified in electromagnetism by the expression

$$
h_{+}(A) = \mathsf{a}'(0) + *B,\tag{6.5}
$$

thereby bypassing the functional analytic construction $|C| = \sqrt{C^2}$.

e. (Decomposition of configuration space) In the non-abelian theory we want to avoid making any gauge choice, such as the Coulomb gauge. The

expression (6.5) defines a vector field on $\mathcal A$ in the non-abelian case. We will show that it flows exactly onto A_+ as $t \to \infty$, analogous to the abelian case, thereby producing a non-linear map $P_+ : A \to A_+$. Similarly for A_- . Upon quotienting by the gauge group, we arrive at a decomposition $C = C_+ \times C_-$. The one-to-one ness of this map needs to be verified, however.

f. (Positivity of the covariant curl.) In electromagnetism one has $(curlA, A)_{L^2(\mathbb{R}^3)} \geq 0$ for all $A \in \mathcal{C}_+$ by (2.36) and Theorem [2.9.](#page-10-2) In the non-abelian case $curl_A$, the gauge covariant curl, does not act on connection forms A but on tangent vectors to A. We will show that $(curl_Au, u)_{L^2(\mathbb{R}^3)} \geq 0$ for all $A \in \mathcal{A}_+$ and for all tangent vectors u to \mathcal{A}_+ at A. The form of the last statement reflects the change often needed when replacing a linear space by a non-linear manifold, where one can no longer identify the manifold with its tangent spaces.

Definition 6.6 For a point A in A denote by B its curvature: $B = dA +$ $A \wedge A$ and by $a(s)$ the Poisson extension of A. Define

$$
h_{+}(A) = \mathsf{a}'(0) + *B \tag{6.6}
$$

$$
h_{-}(A) = \mathbf{a}'(0) - *B. \tag{6.7}
$$

Note that $h_{\pm}(A)$ are 1-forms and therefore can be identified as tangent vectors to A at A. h_{\pm} therefore can and will be identified as vector fields on A. Moreover they are both gauge covariant:

$$
h_{\pm}(A^{g}) = g^{-1}h_{\pm}(A)g.
$$

Theorem 6.7

a. The vector fields $h_{\pm}(A)$ are horizontal at A and mutually orthogonal in $L^2(\mathbb{R}^3)$ for each point $A \in \mathcal{A}$.

- b. h_{\pm} is zero exactly on \mathcal{A}_{\pm} .
- c. The flows $\exp th_{\pm}$ generated by the vector fields h_{\pm} for $t \geq 0$ satisfy

$$
P_{+}(A) := \lim_{t \to \infty} (\exp th_{+})A \quad \text{lies in} \quad \mathcal{A}_{+}.
$$
 (6.8)

$$
P_{-}(A) := \lim_{t \to \infty} (\exp th_{-})A \quad lies \quad in \quad \mathcal{A}_{-}.
$$
 (6.9)

$$
P_{\pm}(A) = A \text{ if } A \in \mathcal{A}_{\pm}.\tag{6.10}
$$

d. Suppose that the Poisson action is convex at A. That is,

$$
\partial_u^2 \mathcal{P}(A) \ge 0 \text{ for all } u. \tag{6.11}
$$

If $A \in \mathcal{A}_+$ and u is tangential to \mathcal{A}_+ at A then

$$
(curl_A u, u)_{L^2} \ge 0. \tag{6.12}
$$

If $A \in \mathcal{A}_-$ and u is tangential to \mathcal{A}_- at A then

$$
(curl_A u, u)_{L^2} \le 0. \tag{6.13}
$$

The proof depends on the following lemmas.

Lemma 6.8

$$
a.) \quad d_A^* \ h_{\pm}(A) = 0 \quad \forall \ A \in \mathcal{A}.
$$

$$
b.) \quad (h_{+}(A), h_{-}(A))_{L^{2}(\mathbb{R}^{3}; \Lambda^{1} \otimes \mathfrak{k})} = 0 \quad \forall \quad A \in \mathcal{A}.
$$
 (6.15)

c.)
$$
h_+(A) = 0
$$
 if and only if $A \in \mathcal{A}_+$ and
\n $h_-(A) = 0$ if and only if $A \in \mathcal{A}_-$.
\n(6.16)

Proof. From [\(5.23\)](#page-29-1) we see that $d_A^* \mathbf{a}'(0) = 0$. Moreover $d_A^* (*B) = -(*d_A*)$ $B = - * d_A B = 0$ by the Bianchi identity. We have used here the identity $d_A^* = - * d_A *$ when acting on 1-forms. Both terms in $h_{\pm}(A)$ are therefore horizontal at A , which proves (6.14) .

To prove [\(6.15\)](#page-41-1) we will first establish the identity

$$
\|\mathbf{a}'(s)\|_2^2 = \|\mathbf{b}(s)\|_2^2 \quad \forall \quad s \ge 0. \tag{6.17}
$$

The computation

$$
(1/2)(d/ds) (||a'(s)||_2^2 - ||b(s)||_2^2)
$$

= $(a''(s), a'(s)) - (b'(s), b(s))$
= $(d_{a(s)}^*b(s), a'(s)) - (d_{a(s)}a'(s), b(s))$
= $(b(s), d_{a(s)}a'(s)) - (d_{a(s)}a'(s), b(s))$
= 0

shows that $\|\mathbf{a}'(s)\|_2^2 - \|\mathbf{b}(s)\|_2^2$ is constant in s. Since $\mathcal{P}(A) < \infty$ it follows that the constant is zero. This proves (6.17) . Using (6.17) we find

$$
(h_{+}(A), h_{-}(A))_{2} = (\mathbf{a}'(0), \mathbf{a}'(0))_{2} - (*\mathbf{b}(0), *b(0))_{2}
$$

= $||\mathbf{a}'(0)||_{2}^{2} - ||\mathbf{b}(0)||_{2}^{2}$
= 0.

This proves [\(6.15\)](#page-41-1).

For the proof of [\(6.16\)](#page-41-3) observe that if $A \in \mathcal{A}_+$ and a is its Poisson extension then $a'(s) + *b(s) = 0$ for all $s \ge 0$ and in particular $h_+(A) =$ $\mathsf{a}'(0) + \ast \mathsf{b}(0) = 0$. Similarly, if $A \in \mathcal{A}_-$ then $h_-(A) = 0$.

To prove the converses let $\alpha(s) = \mathsf{a}'(s) + \ast \mathsf{b}(s)$. Then

$$
(d/ds)\alpha(s) = \mathbf{a}''(s) + *d_{\mathbf{a}}\mathbf{a}'(s)
$$

$$
= d_{\mathbf{a}}^*\mathbf{b}(s) + *d_{\mathbf{a}}\mathbf{a}'(s)
$$

$$
= *d_{\mathbf{a}}(*\mathbf{b}(s) + \mathbf{a}'(s))
$$

$$
= *d_{\mathbf{a}(s)}\alpha(s).
$$

This is a linear homogeneous first order differential equation in $\alpha(s)$. Therefore if $\alpha(0) = 0$ then $\alpha(s) \equiv 0$. So if $h_+(A) = 0$ then $a'(s) + *b(s) = 0$ for all $s \geq 0$. Therefore a is anti-self-dual and $A \in \mathcal{A}_{+}$. Similarly if $h_{-}(A) = 0$ then the function $\beta(s) \equiv a'(s) - *b(s)$ is zero at $s = 0$ and, since it satisfies the differential equation $(d/ds)\beta(s) = - * d_{a(s)}\beta(s)$, it is identically zero. So **a** is self-dual and $A \in \mathcal{A}_-$. \blacksquare

Lemma 6.9 For $t \geq 0$, $\mathcal{P}(\exp(th_{\pm})A)$ is a non-increasing function of t for each $A \in \mathcal{A}$ and for each of the two flows.

Proof. It suffices to compute the derivative of $\mathcal{P}(\exp(th_{\pm})A)$ at $t=0$ because $\exp(th_{\pm})$ are semigroups. At $t = 0$ we have, by [\(5.18\)](#page-28-1),

$$
(1/2)(d/dt)\mathcal{P}(\exp(th_{+})A)\Big|_{t=0} = (1/2)\partial_{u}\mathcal{P}(A)\Big|_{u=h_{+}(A)}
$$

= $-(u, \mathbf{a}'(0))\Big|_{u=h_{+}(A)}$
= $-(\mathbf{a}'(0) + *B, \mathbf{a}'(0))$
= $-||\mathbf{a}'(0)||_{2}^{2} - (*\mathbf{b}(0), \mathbf{a}'(0)).$

But from the Schwarz inequality and [\(6.17\)](#page-41-2) we find

$$
|(*b(0),\mathsf{a}'(0))| \le ||\mathsf{b}(0)||_2 ||\mathsf{a}'(0)||_2
$$

$$
\le ||\mathsf{a}'(0)||_2^2.
$$

Hence

$$
(d/dt)\mathcal{P}(\exp(th_+)A)\Big|_{t=0}\leq 0.
$$

A similar proof applies to $\exp(th_-)$. ■

Lemma 6.10 Let $A \in \mathcal{A}$. If $h_{+}(A)\mathcal{P}(A) = 0$ then $h_{+}(A) = 0$. If $h_-(A)\mathcal{P}(A) = 0$ then $h_-(A) = 0$.

Proof. The hypotheses are short for $\partial_u \mathcal{P}(A) \Big|_{u=h_{\pm}(A)} = 0$. Using [\(5.18\)](#page-28-1) we see that

$$
0 = (1/2)h_{+}(A)\mathcal{P}(A) = -(h_{+}(A), \mathbf{a}'(0))
$$
\n(6.18)

$$
= -(a'(0) + *B, a'(0))
$$
 (6.19)

$$
= -\|\mathbf{a}'(0)\|_2^2 - (*\mathbf{b}(0), \mathbf{a}'(0)). \tag{6.20}
$$

Therefore

$$
-(\ast \mathsf{b}(0), \mathsf{a}'(0)) = \|\mathsf{a}'(0)\|_2^2 \tag{6.21}
$$

But $\|\cdot\|$ + b(0) $\|_2 = \|a'(0)\|_2$ by [\(6.17\)](#page-41-2). Therefore $-*b(0) = a'(0)$ by saturation of the Schwarz inequality. That is, $h_{+}(A) = 0$. In the case of h_{-} one finds $(*b(0), a'(0)) = ||a'(0)||_2^2$ and therefore $*b(0) = a'(0)$. Hence $h_-(A) = 0$.

Proof of Theorem [6.7.](#page-40-0) Items a) and b) of the theorem are proved in Lemma [6.8.](#page-41-4)

Heuristic argument for [\(6.8\)](#page-40-1): The existence of

$$
A_{+} := \lim_{t \to \infty} (\exp th_{+})A \tag{6.22}
$$

in the $L^2(\mathbb{R}^3)$ sense of convergence is reasonable because $\mathcal P$ is decreasing on the orbit of the flow by Lemma [6.9,](#page-42-0) while $\{A : \mathcal{P}(A) \leq const.\}$ (modulo gauge transformations) is compact in L^2 norm over each bounded set in \mathbb{R}^3 , since P has nominally a Sobolev $H_{1/2}$ strength (modulo gauge transformations).

Assuming then the existence of the limit, $\mathcal{P}(A_+)$ cannot decrease anymore under the flow. Hence $h_+(A_+) \mathcal{P}(A_+) = 0$. Therefore $h_+(A_+) = 0$ by Lemma [6.10.](#page-43-0) So $A_+ \in \mathcal{A}_+$ by [\(6.16\)](#page-41-3). Conversely, if $A \in \mathcal{A}_+$ then $h_+(A) = 0$ by [\(6.16\)](#page-41-3). Therefore $\exp(th_+)A = A$. So $A_+ = A$ if $A \in \mathcal{A}_+$. A similar argument applies to $A_-\$. This proves item c) of the theorem. Of course this ensures that the maps $P_{\pm} : A \to A_{\pm}$ are surjective.

For the proof of [\(6.12\)](#page-41-5) suppose that $A \in \mathcal{A}_+$ and that u is tangential to \mathcal{A}_+ at A. Since h_+ is identically zero on \mathcal{A}_+ its tangential derivative $\partial_u h_+(A)$ is zero. That is, by (6.6) ,

$$
\mathsf{u}'(0) + *d_A u = 0,\tag{6.23}
$$

where $u(s)$ is the Poisson extension of u along a. So $curl_A u = *d_A u = -u'(0)$. From the second derivative formula [\(5.28\)](#page-30-3) we find $\partial_u^2 \mathcal{P}(A) = -2(u, u'(0))$ for all A and u . In particular

$$
\partial_u^2 \mathcal{P}(A) = 2(u, \operatorname{curl}_A u) \quad \text{if } A \in \mathcal{A}_+ \text{ and } u \text{ is tangential to } \mathcal{A}_+. \tag{6.24}
$$

From our convexity assumption on the Poisson action at A it now follows that [\(6.12\)](#page-41-5) holds. In case $A \in \mathcal{A}_-$ we find $curl_A u = u'(0)$ and therefore (6.13) holds. \blacksquare

Remark 6.11 The intersection $\mathcal{A}_+ \cap \mathcal{A}_-$ consists of pure gauges. Indeed if A lies in this intersection then $h_{+}(A) = h_{-}(A) = 0$. Therefore $*B = 0$ and $a'(0) = 0$. Since $B = 0$, A is a pure gauge.

6.3 The flows in the electromagnetic case

In the electromagnetic case the vector fields h_{\pm} are given by

$$
h_{\pm}(A) = (-|C| \pm C)A, \tag{6.25}
$$

as we see from the argument showing equality between [\(6.4\)](#page-39-1) and [\(6.5\)](#page-39-2). We can operate in Coulomb gauge in this example. The limits of the flows of h_{\pm} are given in the following theorem.

Theorem 6.12 (Electromagnetic case.) Denote by P_{\pm} the orthogonal projections of C onto \mathcal{C}_\pm respectively. Then

$$
\lim_{t \to \infty} \exp(th_{\pm})A = P_{\pm}A \quad \text{for} \ \ A \in \mathcal{C}.
$$
 (6.26)

Proof. The flow equations for these vector fields are the linear differential equations

$$
(d/dt)A(t) = (-|C| \pm C)A(t),
$$

whose solutions with initial value A are correctly given by

$$
A(t) = e^{t(-|C| \pm C)} A, \quad t \ge 0
$$

because $-|C| \pm C \leq 0$ in both cases. But $-|C| + C$ is zero on C_+ and is $-2|C|$ on \mathcal{C}_- . Therefore, writing $A = P_+A + P_-A$, we find

$$
A(t) = e^{t(-|C|+C)}(P_+A + P_-A) = P_+A + e^{-2t|C|}P_-A,
$$

which converges in C norm to $P_{+}A$ as $t \to \infty$ because $|C| \geq 0$ and has trivial null space in our Coulomb gauge. This proves (6.26) in the case of h_{+} . The case of $h_-\$ is similar because $-|C|-C$ is zero on $\mathcal{C}_-\$ and is $-2|C|$ on $\mathcal{C}_+\$. ■

7 Open questions

1. Existence and uniqueness of solutions to the Euler equation [\(5.6\)](#page-22-4) for the minimization of the Poisson action have been assumed throughout the sections on the Yang-Mills helicity theory. A slight variant of this non-linear boundary value problem has been investigated in deep work by Marini and Isobe cited in Section [5.1.](#page-22-1) The form of the existence and uniqueness theorem we need is still open.

2. The soft Coulomb gauge and asymptotic Coulomb gauge were defined in Section [5.1](#page-22-1) under the assumption that $\lim_{s\to\infty} a(s)$ exists in some useful sense. It needs to be proven that this limit actually exists for $H_{1/2}$ initial data for the Yang-Mills-Poisson equation and is a pure gauge. Such a limit theorem has been established for the Yang-Mills heat equation in various cases. See Remark [5.3.](#page-26-2)

3. The flows of the vector fields h_{\pm} have been assumed to exist in Theorem [6.7](#page-40-0) for all positive time. This needs to be proven along with the existence of the limits as time goes to infinity. The resulting limit maps P_{\pm} are gauge invariant and therefore produce a map $P_+ \times P_- : \mathcal{C} \to \mathcal{C}_+ \times \mathcal{C}_-$. For electromagnetism this gives the linear decomposition $\mathcal{C} = \mathcal{C}_+ \oplus \mathcal{C}_-$. In the non-linear case it remains to be seen whether this map is one-to-one.

4. The region of convexity of the Poisson action function $\mathcal P$ should be understood better. Is there a neighborhood of $A = 0$ on which $\mathcal P$ is convex? Is there a boundary to this region which somehow reflects the Gribov ambiguity?

5. The Lorentz invariance of the natural metric on phase space should be proven or disproven in the non-abelian theory, as already pointed out in Remark [5.20.](#page-36-0) The first step would be to prove invariance under time translation, just at an informal level for, say, smooth initial data. If such invariance should hold then the metric itself would be a candidate for a useful invariant for proving short and long time existence of solutions to the hyperbolic Yang-Mills equation with $H_{1/2} \oplus H_{-1/2}$ initial data. Existence of solutions to this equation has not yet been proven for these critical initial data over three dimensional space. See e.g. [\[50,](#page-51-7) [10,](#page-47-8) [11,](#page-47-9) [30,](#page-49-9) [57,](#page-51-8) [41,](#page-50-9) [42,](#page-50-10) [53\]](#page-51-9). Invariance under Lorentz boosts requires gauge transforming a boosted solution to temporal gauge as part of the boost transformation.

8 Bibliography

References

- [1] Alfred Actor, Classical solutions of SU(2) Yang-Mills theories, Rev. Modern Phys. 51 (1979), no. 3, 461–525. MR 541884
- [2] W. Ambrose and I. M. Singer, A theorem on holonomy, Trans. Amer. Math. Soc. 75 (1953), 428–443. MR 0063739
- [3] M. Asorey and P. K. Mitter, Regularized, continuum Yang-Mills process and Feynman-Kac functional integral, Comm. Math. Phys. 80 (1981), no. 1, 43–58. MR 623151
- [4] M. F. Atiyah, Geometry of Yang-Mills fields, Scuola Normale Superiore Pisa, Pisa, 1979. MR 554924
- [5] V. Bargmann and E. P. Wigner, Group theoretical discussion of relativistic wave equations, Proc. Nat. Acad. Sci. U. S. A. 34 (1948), 211–223. MR 0024827 (9,553d)
- [6] Khalil M. Bitar and Shau Jin Chang, Vacuum tunneling of gauge theory in Minkowski space, Phys. Rev. D (3) 17 (1978), no. 2, 486–497. MR 0462356
- [7] James D. Bjorken and Sidney D. Drell, Relativistic quantum fields, McGraw-Hill Book Co., New York-Toronto-London-Sydney, 1965. MR 0187642 (32 #5092)
- [8] Max Born and Emil Wolf, Principles of optics: Electromagnetic theory of propagation, interference and diffraction of light, With contributions by A. B. Bhatia, P. C. Clemmow, D. Gabor, A. R. Stokes, A. M. Taylor, P. A. Wayman and W. L. Wilcock. Third revised edition, Pergamon Press, Oxford, 1965. MR 0198807 (33 #6961)
- [9] Nelia Charalambous and Leonard Gross, Neumann domination for the Yang-Mills heat equation, J. Math. Phys. 56 (2015), no. 7, 073505, 21. MR 3405967
- [10] Douglas M. Eardley and Vincent Moncrief, The global existence of Yang-Mills-Higgs fields in 4-dimensional Minkowski space. I. Local existence and smoothness properties, Comm. Math. Phys. 83 (1982), no. 2, 171– 191. MR 649158
- [11] , The global existence of Yang-Mills-Higgs fields in 4-dimensional Minkowski space. II. Completion of proof, Comm. Math. Phys. 83 (1982), no. 2, 193–212. MR 649159
- [12] P. L. García and A. Pérez-Rendón, *Symplectic approach to the theory of* quantized fields. I, Comm. Math. Phys. 13 (1969), 24–44. MR 0256681
- [13] Pedro L. García, *Gauge algebras, curvature and symplectic structure*, (1974), 63–71. MR 0448418
- $[14]$ $_____\$ nfilterianarriant in the calculus of variations, (1974), 219–246. MR 0406246
- [15] _____, Reducibility of the symplectic structure of classical fields with gauge-symmetry, (1977), 365–376. Lecture Notes in Math., Vol. 570. MR 0445542
- [16] , Tangent structure of Yang-Mills equations and Hodge theory, Differential geometrical methods in mathematical physics (Proc. Conf., Aix-en-Provence/Salamanca, 1979), Lecture Notes in Math., vol. 836, Springer, Berlin-New York, 1980, pp. 292–312. MR 607703
- [17] Leonard Gross, Equivalence of helicity and Euclidean self-duality for quantized gauge fields.
- [18] $\frac{1}{s}$, Norm invariance of mass-zero equations under the conformal group, J. Mathematical Phys. 5 (1964), 687–695. MR 0164632
- [19] , The Yang-Mills heat equation with finite action, (2016), 137 pages, http://arxiv.org/abs/1606.04151.
- [20] Werner Heisenberg and Wolfgang Pauli, Zur Quantendynamic der *Wellenfelder*, Zeits. f. Phys. **56** (1929), no. 1, $1 - 61$.
- [21] Min-Chun Hong and Gang Tian, Asymptotical behaviour of the Yang-Mills flow and singular Yang-Mills connections, Math. Ann. 330 (2004), no. 3, 441–472. MR MR2099188 (2006h:53063)
- [22] , Global existence of the m-equivariant Yang-Mills flow in four dimensional spaces, Comm. Anal. Geom. 12 (2004), no. 1-2, 183–211. MR MR2074876 (2005e:53103)
- [23] Min-Chun Hong, Gang Tian, and Hao Yin, The Yang-Mills α -flow in vector bundles over four manifolds and its applications, Comment. Math. Helv. 90 (2015), no. 1, 75–120. MR 3317334
- [24] Takeshi Isobe and Antonella Marini, On topologically distinct solutions of the Dirichlet problem for Yang-Mills connections, Calc. Var. Partial Differential Equations 5 (1997), no. 4, 345–358. MR MR1450715 (99f:58045)
- [25] Small coupling limit and multiple solutions to the Dirichlet problem for Yang-Mills connections in four dimensions. I, J. Math. Phys. 53 (2012), no. 6, 063706, 39. MR 2977701
- [26] _____, Small coupling limit and multiple solutions to the Dirichlet problem for Yang-Mills connections in four dimensions. II, J. Math. Phys. 53 (2012), no. 6, 063707, 39. MR 2977702
- [27] John David Jackson, Classical electrodynamics, third ed., John Wiley & Sons Inc., New York, 1999. MR MR0436782 (55 #9721)
- $|28|$ Adam Jacob, The limit of the Yang-Mills flow on semi-stable bundles, J. Reine Angew. Math. 709 (2015), 1–13. MR 3430873
- [29] Hans Plesner Jakobsen and Michele Vergne, Wave and Dirac operators, and representations of the conformal group, J. Functional Analysis 24 (1977), no. 1, 52–106. MR 0439995
- [30] S. Klainerman and M. Machedon, Finite energy solutions of the Yang-Mills equations in \mathbb{R}^{3+1} , Ann. of Math. (2) 142 (1995), no. 1, 39–119. MR MR1338675 (96i:58167)
- [31] L. D. Landau and R. Peierls, Quantum electrodynamics in configuration space, Zeits. f. Phys. 62 (1930), 188–199 (Trans. in ter Haar (1965) pp. 19–30.
- [32] Elliot Leader and Cédric Lorcé, *The angular momentum controversy:* Whats it all about and does it matter?, Physics Reports 541 (2014), no. 3, 163–248.
- [33] Antonella Marini, Boundary value problems for Yang-Mills connections, C. R. Acad. Sci. Paris Sér. I Math. 312 (1991), no. 7, 503-508. MR MR1099681 (92c:58020)
- [34] $____\$ n Dirichlet and Neumann boundary value problems for Yang-Mills connections, Comm. Pure Appl. Math. 45 (1992), no. 8, 1015–1050. MR MR1168118 (93k:58059)
- [35] $______\$ Elliptic boundary value problems for connections: a nonlinear Hodge theory, Mat. Contemp. 2 (1992), 195–205, Workshop on the Geometry and Topology of Gauge Fields (Campinas, 1991). MR MR1303162 (95k:58162)
- [36] $________\$ A topological method for finding non-absolute minima for the Yang-Mills functional with Dirichlet data, Geometry, topology and physics (Campinas, 1996), de Gruyter, Berlin, 1997, pp. 191–199. MR MR1605228 (99k:58042)
- [37] , The generalized Neumann problem for Yang-Mills connections, Comm. Partial Differential Equations 24 (1999), no. 3-4, 665–681. MR MR1683053 (2000c:58025)
- [38] $_____\$ n, Regularity theory for the generalized Neumann problem for Yang-Mills connections—non-trivial examples in dimensions 3 and 4, Math. Ann. 317 (2000), no. 1, 173–193. MR MR1760673 (2001i:58020)
- [39] P. K. Mitter and C.-M. Viallet, On the bundle of connections and the gauge orbit manifold in Yang-Mills theory, Comm. Math. Phys. 79 (1981), no. 4, 457–472. MR 623962 (83f:81056)
- [40] M. S. Narasimhan and T. R. Ramadas, Geometry of SU(2) gauge fields, Comm. Math. Phys. 67 (1979), no. 2, 121–136. MR MR539547 (84k:58050)
- [41] Sung-Jin Oh, Gauge choice for the Yang-Mills equations using the Yang-Mills heat flow and local well-posedness in H^1 , J. Hyperbolic Differ. Equ. 11 (2014), no. 1, 1–108. MR 3190112
- [42] ____, Finite energy global well-posedness of the Yang-Mills equations on \mathbb{R}^{1+3} : an approach using the Yang-Mills heat flow, Duke Math. J. 164 (2015), no. 9, 1669–1732. MR 3357182
- [43] Sung-Jin Oh and Daniel Tataru, The Yang-Mills heat flow and the caloric gauge, (2017) , http://arxiv.org/abs/1709.08599.
- [44] S. M. Paneitz and I. E. Segal, Quantization of wave equations and Hermitian structures in partial differential varieties, Proc. Nat. Acad. Sci. U.S.A. 77 (1980), no. 12, part 1, 6943–6947. MR 603063
- [45] Stephen M. Paneitz, *Essential unitarization of symplectics and applica*tions to field quantization, J. Funct. Anal. 48 (1982), no. 3, 310–359. MR 678176
- [46] Silvan S. Schweber, An introduction to relativistic quantum field theory, Foreword by Hans A. Bethe, Row, Peterson and Company, Evanston, Ill.-Elmsford, N.Y., 1961. MR 0127796
- [47] I. E. Segal, Quantization of nonlinear systems, J. Mathematical Phys. 1 (1960), 468–488. MR 0135093
- [48] $_____$, *Stability theory and quantization*, Differential geometrical methods in mathematical physics (Proc. Conf., Aix-en-Provence/Salamanca, 1979), Lecture Notes in Math., vol. 836, Springer, Berlin-New York, 1980, pp. 375–382. MR 607709
- [49] \ldots , The phase space for the Yang-Mills equations, Differential geometric methods in mathematical physics (Proc. Internat. Conf., Tech. Univ. Clausthal, Clausthal-Zellerfeld, 1978), Lecture Notes in Phys., vol. 139, Springer, Berlin-New York, 1981, pp. 101–109. MR 612994
- [50] Irving Segal, The Cauchy problem for the Yang-Mills equations, J. Funct. Anal. 33 (1979), no. 2, 175–194. MR 546505
- [51] , Quantization of symplectic transformations, Mathematical analysis and applications, Part B, Adv. in Math. Suppl. Stud., vol. 7, Academic Press, New York-London, 1981, pp. 749–758. MR 634267
- [52] Irving E. Segal, Mathematical problems of relativistic physics, With an appendix by George W. Mackey. Lectures in Applied Mathematics (proceedings of the Summer Seminar, Boulder, Colorado, vol. 1960, American Mathematical Society, Providence, R.I., 1963. MR 0144227
- [53] Sigmund Selberg and Achenef Tesfahun, Null structure and local wellposedness in the energy class for the Yang-Mills equations in Lorenz gauge, J. Eur. Math. Soc. (JEMS) 18 (2016), no. 8, 1729–1752. MR 3519539
- [54] M. Shifman, Advanced topics in quantum field theory, Cambridge University Press, Cambridge, 2012, A lecture course. MR 2896444
- [55] I. M. Singer, Some remarks on the Gribov ambiguity, Comm. Math. Phys. 60 (1978), no. 1, 7–12. MR MR500248 (80d:53025)
- [56] $\frac{1}{\sqrt{56}}$, The geometry of the orbit space for nonabelian gauge theories, Phys. Scripta 24 (1981), no. 5, 817–820. MR MR639408 (83a:81055)
- [57] Terence Tao, Local well-posedness of the Yang-Mills equation in the temporal gauge below the energy norm, J. Differential Equations 189 (2003), no. 2, 366–382. MR MR1964470 (2003m:58016)
- [58] Stefan Vandoren and Peter van Nieuwenhuisen, Lectures on instantons, hep-th (2008), 118 pages, http://arxiv.org/abs/0802.1862v1.
- [59] N. M. J. Woodhouse, Geometric quantization, second ed., Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1992, Oxford Science Publications. MR 1183739