

# Black hole evaporation in Conformal (Weyl) gravity

Hao Xu\*

*Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China and  
Department of Physics, University of Science and Technology of China, Hefei 230026, China*

Man-Hong Yung

*Institute for Quantum Science and Engineering and Department of Physics,  
Southern University of Science and Technology, Shenzhen 518055, China and  
Shenzhen Key Laboratory of Quantum Science and Engineering, Shenzhen, 518055, China*  
(Dated: April 16, 2019)

In this work, we explore all possible scenarios in the evaporation process of a spherical neutral AdS black hole in four-dimensional conformal gravity, where the equations of states are branched. In one branch, the final states correspond to the extremal black hole where the total decay time is divergent given any initial mass. In the other branch, the black holes always evaporate completely within a finite time; the total decay time depends linearly on the AdS radius, when the mass is taken to be infinity.

Recently, conformal (Weyl) gravity, which is described by a pure Weyl squared action, has attracted a considerable amount of interest as an alternative theory to Einstein gravity. From the equation of motion and the symmetry of conformal gravity, any conformal class of the Einstein solution arises naturally as a solution to the conformal gravity. In particular, subject to the Neumann boundary condition, conformal gravity can single out Einstein's solution [1, 2]. Moreover, unlike Einstein's gravity, conformal gravity has been shown to be perturbatively renormalizable in four dimensions [3], leading to many interesting alternative approaches to quantum gravity [4].

Another appealing aspect of conformal gravity comes from cosmology. Although Einstein's gravity can describe the physics within the scale of the Solar system perfectly, there are still many unsolved puzzles when applied to a larger scale, such as the inconsistency with the observation of galactic rotation curves and accelerating universe. Consequently, unknown entities namely "dark matter and dark energy", have to be introduced in order to fix the inconsistency problems. Thus, one may wonder whether it is possible to modify the nature of gravity to explain the physics at a larger scale, while maintaining the correct behaviors at the scale of the Solar system. Since conformal gravity admits more solutions than Einstein's gravity, it can produce the effective potential consistent with the observed phenomenon, making it a compelling modified gravity theory [5–8].

Although conformal gravity and Einstein's gravity may share the same space-time solutions, thermodynamic quantities of the black hole, such as the entropy and mass, depend on the action rather than the metric. The difference in the actions means that the black-hole thermodynamics of the two gravity theories would be different. The thermodynamic phase structure for the conformal gravity in the four-dimensional AdS space-time

has been explored [9–12]; the phase structure of the conformal gravity includes two branches of equations of states, zeroth-order phase transition, and Hawking-Page-like phase transition.

However, there are still some puzzles left unsolved. More than forty years after the discovery of Hawking radiation, its precise nature and consequences are still, to a large extent, open questions. The radiation causes a black hole to lose mass and thermal entropy, if there is not sufficient incoming radiation to balance it. For a non-rotating neutral Schwarzschild black hole with mass  $M_0$  in four-dimensional asymptotically flat space-time without incoming radiation, we expect it has a lifetime  $t \sim M_0^3$ , which grows as the cube of the mass and diverges when  $M_0 \rightarrow +\infty$  [13, 14]. Similar analysis was also carried out for a spherical black hole in AdS space-time [15] and was later extended to flat and hyperbolic black holes [16]. The gravitational potential of AdS space-time on the boundary acts like a finite confining box for the black hole. Massless radiation particles can escape to infinity and reflect back in a finite time. However, if one chooses an absorbing boundary condition [17] so that Hawking radiation is absorbed by the AdS boundary preventing the returning of the radiation, then one can calculate the time for the black-hole mass to drop from one initial value to another, which is of great interest in holography [18, 19]. In particular, Page made a counter-intuitive discovery [15] that this time does not diverge even when the initial black hole mass is taken to be infinity.

Given the thermodynamic phase structure of conformal gravity in AdS space-time, it is natural to investigate the black-hole evaporation process, which points to an area where extensive research should be carried out. In particular, we are interested in answering the question, whether the total decay time is also finite when the black hole mass is taken to infinity.

We start by giving a brief review of the black hole thermodynamics in conformal gravity [9–12]. The gravity

\* xuh5@sustc.edu.cn

action is proportional to the square of the Weyl tensor,

$$S = \alpha \int d^4x \sqrt{-g} C^{\mu\nu\rho\sigma} C_{\mu\nu\rho\sigma}, \quad (1)$$

where the coupling constant  $\alpha$  is associated with a dimension of [length]<sup>2</sup>. Notice that some thermodynamic quantities, such as the black hole mass, entropy, and Gibbs free energy, are proportional to  $\alpha$ . However, the exact value of  $\alpha$  does not change any qualitative features of the thermodynamics. Without loss of generality, we set  $\alpha = 1$  in the present work. In other words, the black hole thermodynamic quantities are described in unit of  $\alpha$ .

The simplest form of spherical black hole solution to four-dimensional conformal gravity [20, 21] is

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (2)$$

where

$$f(r) = c_0 + c_1 r + \frac{d}{r} - \frac{1}{3}\Lambda r^2. \quad (3)$$

Due to the conformal symmetry, the Weyl rescaling of the metric above remains a static and spherically-symmetric solution. Here the four integral constants,  $c_0, c_1, d, \Lambda$ , are subject to the following constraint:

$$c_0^2 = 3c_1 d + 1, \quad (4)$$

which implies that we have a discrete freedom in choosing the constant  $c_0$ ,

$$c_0 = \pm \sqrt{3c_1 d + 1}. \quad (5)$$

If we take  $c_1 = 0$  and  $c_0 = 1$ , the solution reduces to the well-known Schwarzschild (A)dS black hole.

Here the  $\Lambda$  plays the role of ‘‘cosmological constant’’, which comes from the integral instead of the action. Furthermore, we adopt the language of extended phase space and treat it as effective pressure  $P = -\frac{\Lambda}{8\pi}$ , which is allowed to vary [22–51].

The black hole mass is defined to be the conserved charge of the time-like killing vector [9]

$$M = \frac{(c_1 c_0 - c_1 - 16\pi P d)}{12\pi}. \quad (6)$$

The temperature is proportional to the surface gravity at the black hole horizon  $r_0$

$$T = \frac{8\pi P r_0^3 - 3c_0 r_0 - 6d}{12\pi r_0^2}, \quad (7)$$

and its conjugate, i.e. the entropy is

$$S = \frac{(r_0 - c_0 r_0 - 3d)}{3r_0}. \quad (8)$$

Note that the entropy depends on a combination of  $c_0, r_0$  and  $d$ , rather than just being proportional to the area

of the horizon [9]. Furthermore, the first law of black hole thermodynamics and the Smarr formula in extended phase space have been investigated in [9–12] and will not be covered here.

Now, applying  $f(r_0) = 0$  and (4) to eliminate other unnecessary coefficients, we found that the equations of states are branched [11]. One set of the solutions is given by,

$$\begin{aligned} T_1 &= 2Pr_0 - \frac{\sqrt{-32\pi P c_1 r_0^3 - 3c_1^2 r_0^2 + 4}}{8\pi r_0} + \frac{c_1}{8\pi}, \\ S_1 &= \frac{8}{3}\pi P r_0^2 - \frac{1}{3}\sqrt{-32\pi P c_1 r_0^3 - 3c_1^2 r_0^2 + 4} + \frac{1}{3}. \end{aligned} \quad (9)$$

The other one is given by,

$$\begin{aligned} T_2 &= 2Pr_0 + \frac{\sqrt{-32\pi P c_1 r_0^3 - 3c_1^2 r_0^2 + 4}}{8\pi r_0} + \frac{c_1}{8\pi}, \\ S_2 &= \frac{8}{3}\pi P r_0^2 + \frac{1}{3}\sqrt{-32\pi P c_1 r_0^3 - 3c_1^2 r_0^2 + 4} + \frac{1}{3}. \end{aligned} \quad (10)$$

Here we use  $r_0$  as an intermediate parameter to avoid the overwhelming complexity of formula  $T(P, S)$ .

Next, we investigate the possible critical behavior [10, 11] by considering the following two conditions:

$$\frac{\partial T}{\partial S} = 0, \quad \frac{\partial^2 T}{\partial S^2} = 0, \quad (11)$$

yielding the critical point,

$$\begin{aligned} P_c &= \frac{(2 - 3X^2 - \sqrt{3}X)c_1^2}{24\pi X^3}, \quad T_c = \frac{c_1}{6\pi X^2}, \\ S_c &= \frac{\sqrt{3}X + 3X + 2}{9X}, \end{aligned} \quad (12)$$

where  $X = \sqrt{2} - 2\sqrt{3}/3$ . This critical point only appears for  $c_1 > 0$  and corresponds to the zeroth-order phase transition.

Now we concentrate on the black hole evaporation process. The black hole mass  $M$  and entropy  $S$  should be monotonically-decreasing functions of time. We shall apply the geometrical optics approximation, which assumes massless quanta move along null geodesics [15]. If we orient the angular coordinates and normalize the affine parameter  $\lambda$ , the geodesic equation takes the form

$$\left(\frac{dr}{d\lambda}\right)^2 = E^2 - J^2 \frac{f(r)}{r^2}, \quad (13)$$

where  $E = f(r)\frac{dt}{d\lambda}$  and  $J = r^2\frac{d\theta}{d\lambda}$  are the energy and angular momentum of the massless quanta respectively. Considering a null geodesic comes from the black hole horizon, it will return to the hole when there is a turning point satisfying  $(\frac{dr}{d\lambda})^2 = 0$  thus cannot be detected by the observer on the AdS boundary. If we introduce  $b \equiv \frac{J}{E}$ , then the massless quanta can reach infinity if and only if  $b$  satisfies

$$\frac{1}{b^2} \geq \frac{f(r)}{r^2} \quad (14)$$

for any  $r \geq r_0$ . Therefore, we need to find the maximal value of  $\frac{f(r)}{r^2}$ . Depending on the exact form of  $f(r)$ , there are different cases. The  $\frac{f(r)}{r^2}$  may admit a maximal value corresponding to an unstable photon orbit  $r_p$ , such as in the spherical AdS black hole case [15]. Or it may be monotonically increasing and approaches  $\frac{8\pi P}{3}$  near the boundary, such as in the flat AdS black hole case [16].

For the conformal gravity with metric (2), we can investigate the  $\frac{f(r)}{r^2}$  for different signs of  $c_1$  and  $c_0$ . In TABLE I, we list all possible values of  $r_p$  so that we can find the impact factor  $b_c^2 = \frac{r_p^2}{f(r_p)}$ . Here the *none* means the  $\frac{f(r)}{r^2}$  is a monotonic function of  $r_0$  and the impact factor  $b_c^2 = \frac{3}{8\pi P}$ .

	$c_1 = 0$	$c_1 > 0$	$c_1 < 0$
$c_0 > 0$	$-\frac{3d}{2}$	$\frac{1-\sqrt{1+3c_1d}}{c_1}$	$\frac{1-\sqrt{1+3c_1d}}{c_1}$
$c_0 < 0$	<i>none</i>	$\frac{1+\sqrt{1+3c_1d}}{c_1}$	<i>none</i>

TABLE I. The value of the photon orbit radius  $r_p$ .

When  $c_1 = 0$  and  $c_0 > 0$ , the black hole metric reduces to the Schwarzschild AdS metric, and the photon orbit radius  $r_p = -\frac{3d}{2}$ . When  $c_1 \neq 0$  and  $c_0 > 0$ ,

$$r_p = \frac{1 - \sqrt{1 + 3c_1d}}{c_1} \quad (15)$$

$$\sim -\frac{3d}{2} \left(1 - \frac{3d}{4}c_1 + \frac{9d^2}{8}c_1^2 + O(d^3c_1^3)\right),$$

which goes to  $-\frac{3d}{2}$  as  $c_1 \rightarrow 0$ . When  $c_1 > 0$  and  $c_0 < 0$ ,

$$r_p = \frac{1 + \sqrt{1 + 3c_1d}}{c_1} \quad (16)$$

$$\sim \frac{2}{c_1} + \frac{3d}{2} - \frac{9d^2}{8}c_1 + \frac{27d^3}{16}c_1^2 + O(d^4c_1^3),$$

which goes to  $+\infty$  as  $c_1 \rightarrow 0$ . In  $c_1 \leq 0$  and  $c_0 < 0$ , the  $\frac{f(r)}{r^2}$  is a monotonic function of  $r_0$ .

According to the Stefan-Boltzmann law, we conclude that in four dimensional space-time, the Hawking emission power (luminosity) is

$$\frac{dM}{dt} = -Cb_c^2T^4 \quad (17)$$

with some numerical constants  $C$ . Since we are only concerned about the qualitative features of the evaporation process, we shall ignore this constant. The qualitative features of the thermodynamic phase structure and the impact factors depend on the signs of  $c_1$  and  $c_0$ . We will investigate them in all different situations.

1.  $c_1 = 0$ . In this case, the equations of states for the two solutions become

$$T_1 = 2Pr_0 - \frac{1}{4\pi r_0}, \quad S_1 = \frac{8}{3}\pi Pr_0^2 - \frac{1}{3}, \quad (18)$$

and

$$T_2 = 2Pr_0 + \frac{1}{4\pi r_0}, \quad S_2 = \frac{8}{3}\pi Pr_0^2 + 1, \quad (19)$$

which correspond to  $c_0 = -1$  and  $c_0 = 1$  respectively. The isobaric curves are presented in FIG.1. For  $c_0 = -1$  (the first figure in in FIG.1), the curves are monotonically increasing, and  $T \rightarrow 0$  as  $S \rightarrow 0$ , which corresponds to an extremal black hole. The radius of the extremal black hole reads

$$r_0 = \frac{1}{\sqrt{8\pi P}}. \quad (20)$$

When the black hole evaporates and loses mass and entropy, the temperature decreases so the evaporation process is increasingly difficult.

For  $c_0 = 1$  (the second figure in in FIG.1), the black hole is in an analogy with the Hawking-Page phase transition in Schwarzschild AdS black hole from Einstein gravity [52]. For each  $P$  the black hole admits a minimal temperature. The curve with negative slope to the left of the minimum corresponds to small black holes that are thermodynamically unstable. Black hole that reaches these states will soon be evaporated completely. Notice that the black hole entropy includes a pure constant 1 even in  $r_0 = 0$ . It can be removed by introducing a Gauss-Bonnet term in the action [9].

In FIG.2 we provide some numerical examples of the black hole evolution. We introduce  $\Delta M$ , defined as  $\Delta M = M_0 - M_f$ .  $M_0$  is the initial mass.  $M_f$  is the final mass, which corresponds to the  $T \rightarrow 0$  in  $c_0 = -1$  and  $T \rightarrow \infty$  on the left side of the minimum in  $c_0 = 1$ . In the first figure, we can observe the black hole mass decreases rapidly in the beginning but the evaporation slows down quickly and it will take infinite time to reach zero temperature, which obeys the third law of thermodynamics. However, in the second figure, although we set the initial mass to be very large, the total decay time to Hawking emission is still *finite*. This result agrees with the case of the Schwarzschild AdS black hole in Einstein gravity.

It is worth emphasizing that from FIG.2 we can observe the smaller pressure  $P$  corresponds the larger total decay time. In [15] Page proved in four dimensional AdS space-time the decay time is bounded by a time of the order of  $\ell^3$ , where  $\ell = \sqrt{\frac{3}{8\pi P}}$  is the AdS radius. However, although the temperature  $T$  in  $c_1 = 0$  and  $c_0 = 1$  shares the same formula with the Schwarzschild AdS black hole, the black hole mass (6) is defined as  $M = -\frac{4\pi Pd}{3} \sim -\frac{d}{\ell^2}$  (the scaling can be compensated by the coupling constant  $\alpha$  in the action), rather than  $-d$ . We can check that the decay time in conformal gravity is in the *linear* order of  $\ell$ , not  $\ell^3$ . See FIG.3 for numerical result.

2.  $c_1 > 0$ . The exact value of  $c_1$  does not change the qualitative features of the thermodynamic phase structure. Since the thermodynamic quantities are described




FIG. 1. The isobaric curves in  $T - S$  planes for  $c_1 = 0$ . The two figures correspond to  $c_0 = -1$  and  $c_0 = 1$  respectively. In each figure the pressure  $P$  corresponds to  $P = 1, 2, 3$  from bottom to top.

as functions of  $P$  and  $r_0$ , we can introduce a reduced parameter space  $\tilde{r}_0 = r_0 c_1$  and  $\tilde{P} = \frac{P}{c_1^2}$ , so that all the thermodynamic quantities, such as  $\tilde{M} = \frac{M}{c_1}$  and  $\tilde{T} = \frac{T}{c_1}$ , have [length]<sup>0</sup>(constant) scaling. The  $c_1$  is used as a unit to describe the quantities. Now the equations of states become

$$\begin{aligned} \tilde{T}_1 &= 2\tilde{P}\tilde{r}_0 - \frac{\sqrt{-32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4}}{8\pi\tilde{r}_0} + \frac{1}{8\pi}, \\ S_1 &= \frac{8}{3}\pi\tilde{P}\tilde{r}_0^2 - \frac{1}{3}\sqrt{-32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4} + \frac{1}{3}, \end{aligned} \quad (21)$$

and

$$\begin{aligned} \tilde{T}_2 &= 2\tilde{P}\tilde{r}_0 + \frac{\sqrt{-32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4}}{8\pi\tilde{r}_0} + \frac{1}{8\pi}, \\ S_2 &= \frac{8}{3}\pi\tilde{P}\tilde{r}_0^2 + \frac{1}{3}\sqrt{-32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4} + \frac{1}{3}. \end{aligned} \quad (22)$$

In FIG.4 we present the isobaric curves in the  $\tilde{T} - S$  plane. The solid and dotted lines correspond to the  $c_0 > 0$  and  $c_0 < 0$  cases respectively. Notice that there is also a region of  $(\tilde{T}_2, S_2)$  with negative  $c_0$ . See [11] for




FIG. 2. The evolution of the black hole in  $c_1 = 0$ . The two figures correspond to  $c_0 = -1$  and  $c_0 = 1$  respectively. In each figure the pressure  $P$  corresponds to  $P = 1, 2, 3$  from top to bottom.




FIG. 3. The relationship between the total decay time  $t_d$  and AdS radius  $\ell = \sqrt{\frac{3}{8\pi P}}$  in  $c_1 = 0$  and  $c_0 = 1$ .

detailed discussion. We can observe that the radius (and also mass and entropy) is bounded from above for each  $P$  due to the  $\sqrt{-32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4}$  term.

Solving the equations of  $\tilde{T}_1 = 0$  and  $S_1 = 0$ , we can find there is also an extremal black hole for each  $P$  in  $c_0 < 0$ . The radius of the extremal black hole reads

$$\tilde{r}_0 = \frac{\sqrt{32\pi P + 1} - 1}{16\pi P}. \quad (23)$$

For  $c_0 > 0$ , the temperature  $\tilde{T}_2 \rightarrow \infty$  as  $\tilde{r}_0 \rightarrow 0$ .

We give a brief discussion about the black hole evaporation in AdS space-time. Unlike asymptotically flat




FIG. 4. The isobaric curves in  $\frac{T}{c_1} - S$  planes for  $c_1 > 0$ . The dotted and solid line correspond to  $c_0 < 0$  and  $c_0 > 0$  respectively. The pressure  $P$  corresponds to  $P = 1, 2, 3$  from bottom to top.

space-time, in AdS space-time the black hole mass is determined by both the cosmological constant and the black hole radius, thus it seems unnatural to compared the black hole evaporation process in different cosmological constants. The best we can do is to take the limit  $r_0 \rightarrow \infty$  and then study the relationship between the total decay time and the cosmological constant. In Einstein gravity, the total decay time is in the order of  $\ell^3$ , while in conformal gravity with  $c_1 = 0$  and  $c_0 = 1$ , it is in the order of  $\ell$ . However, in  $c_1 > 0$  the black hole mass and radius are bounded from above for each  $P$ , so we cannot take the limit  $r_0 \rightarrow \infty$ . The numeral analysis shows in  $c_0 > 0$  the black hole can evaporate completely in finite time, depending on the explicit values of the  $P$  and initial mass. In  $c_0 < 0$  the total decay time is divergent in any  $P$  and any initial black hole radius. We omit the numerical example.

3.  $c_1 < 0$ . In this case, similarly we can also introduce the dimensionless quantities  $\tilde{r}_0 = r_0|c_1|$  and  $\tilde{P} = \frac{P}{c_1^2}$ , and the black hole mass  $\tilde{M} = \frac{M}{|c_1|}$  and temperature  $\tilde{T} = \frac{T}{|c_1|}$ . The equations of states now take the form

$$\begin{aligned} \tilde{T}_1 &= 2\tilde{P}\tilde{r}_0 - \frac{\sqrt{32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4}}{8\pi\tilde{r}_0} - \frac{1}{8\pi}, \\ S_1 &= \frac{8}{3}\pi\tilde{P}\tilde{r}_0^2 - \frac{1}{3}\sqrt{32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4} + \frac{1}{3}, \end{aligned} \quad (24)$$

and

$$\begin{aligned} \tilde{T}_2 &= 2\tilde{P}\tilde{r}_0 + \frac{\sqrt{32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4}}{8\pi\tilde{r}_0} - \frac{1}{8\pi}, \\ S_2 &= \frac{8}{3}\pi\tilde{P}\tilde{r}_0^2 + \frac{1}{3}\sqrt{32\pi\tilde{P}\tilde{r}_0^3 - 3\tilde{r}_0^2 + 4} + \frac{1}{3}. \end{aligned} \quad (25)$$




FIG. 5. The isobaric curves in  $\frac{T}{|c_1|} - S$  planes for  $c_1 < 0$ . The two figures correspond to  $c_0 < 0$  and  $c_0 > 0$  respectively. In each figure the pressure  $P$  corresponds to  $P = 1, 2, 3$  from bottom to top.

Now the black hole radius and mass are unbound. In FIG.5 we can observe the the isobaric curves are in an analogy the the case of  $c_1 = 0$ . For  $c_0 < 0$  (the first figure in FIG.5), we can find there is also an extremal black hole for each  $P$  whose radius reads

$$\tilde{r}_0 = \frac{\sqrt{32\pi P + 1} + 1}{16\pi P}. \quad (26)$$

For  $c_0 > 0$  (the second figure in FIG.5) the black hole is in an analogy with the the Schwarzschild AdS black hole. The  $c_1$  which acts as a correction can be ignored in large mass limit, thus does not change the qualitative features. In FIG.6 we present some numerical examples of the black hole evolution. In  $c_0 < 0$  the total decay time is divergent and in  $c > 0$  it reminds finite. The relationship between the total decay time and AdS radius is presented in FIG.7.

In this paper we explore the spherical neutral AdS black hole evaporation process in four dimensional conformal gravity. We investigate the system in three cases:  $c_1 = 0$ ,  $c_1 > 0$  and  $c_1 < 0$ . In each case we have two branches of equations of states, which corresponds to  $c_0 < 0$  and  $c_0 > 0$ . For  $c_0 < 0$ , the equations of states ad-




FIG. 6. The evolution of the black hole in  $c_1 < 0$ . The two figures correspond to  $c_0 < 0$  and  $c_0 > 0$  respectively. In each figure the pressure  $P$  corresponds to  $P = 1, 2, 3$  from top to bottom.




FIG. 7. The relationship between the total decay time and AdS radius  $\ell = \sqrt{\frac{3}{8\pi P}}$  in  $c_1 < 0$  and  $c_0 > 0$ .

mit an extremal black hole with zero temperature and entropy in all three cases. In  $c_1 = 0$  the extremal black hole radius  $r_0 = \frac{1}{\sqrt{8\pi P}}$ , while in  $c_1 > 0$  the  $\tilde{r}_0 = \frac{\sqrt{32\pi P - 1} + 1}{16\pi P}$ , and in  $c_1 < 0$  the  $\tilde{r}_0 = \frac{\sqrt{32\pi P + 1} + 1}{16\pi P}$ . When the black hole

evaporates and loses mass, the black hole temperature decreases so the evaporation process is increasingly difficult. Near the extremal black hole the evaporation slows down quickly, and it will take infinite time to reach it, which obeys the third law of thermodynamics.

For  $c_0 > 0$ , the black hole is in an analogy with the Hawking-Page phase transition in Schwarzschild AdS black hole. When  $r_0 \rightarrow 0$ ,  $T \rightarrow \infty$  in all three cases. The black hole can always evaporate away in a finite time. In the cases of  $c_1 = 0$  and  $c_1 < 0$ , the black hole mass is unbounded. When the initial mass is taken to infinity, we can observe the decay time is also finite and is in the *linear* order of  $\ell$ , which is different from the  $\ell^3$  in the Einstein gravity. This is due to the different formulas of the black hole mass in two gravity theories.

Similar analysis may also be extended to charged and/or rotating black holes. In Einstein gravity, Page proved an asymptotically flat rotating neutral black hole always loses angular momentum more rapidly than it loses mass [53]. For the charged black hole, the production of charged particles should be treated separately from the massless particles, but they are all part of Hawking emission. The explicit evolution process depends on both the initial black hole mass and charge [54–56]. It would be interesting to consider such examples in conformal gravity. Furthermore, in the present work we only concentrate on the qualitative features of the black hole evaporation. To get a precise lifetime one has to compute the grey body factors of all the particles in the theory [57–72]. More importantly, near the end of the evaporation, the black hole gets extremely hot and effective field theory starts to fail. New physics, maybe quantum gravity, could affect the subsequent evolution. See e.g. [73–76] for discussion and references in this field. This process has yet to be fully understood. We hope to be able to consider these questions in our future work.

## ACKNOWLEDGMENTS

We acknowledge the support by Natural Science Foundation of Guangdong Province (2017B030308003) and the Guangdong Innovative and Entrepreneurial Research Team Program (No.2016ZT06D348), and the Science Technology and Innovation Commission of Shenzhen Municipality (ZDSYS20170303165926217, JCYJ20170412152620376).

- 
- [1] J. Maldacena, “Einstein Gravity from Conformal Gravity,” [arXiv:1105.5632].  
 [2] G. Anastasiou and R. Olea, “From conformal to Einstein Gravity,” Phys. Rev. D **94**, no. 8, 086008 (2016) [arXiv:1608.07826].  
 [3] P. D. Mannheim, “Comprehensive Solution to the

- Cosmological Constant, Zero-Point Energy, and Quantum Gravity Problems,” Gen. Rel. Grav. **43**, 703 (2011) [arXiv:0909.0212].  
 [4] K. S. Stelle, “Renormalization of Higher Derivative Quantum Gravity,” Phys. Rev. D **16**, 953 (1977).  
 [5] P. D. Mannheim and D. Kazanas, “Exact Vacuum

- Solution to Conformal Weyl Gravity and Galactic Rotation Curves,” *Astrophys. J.* **342**, 635 (1989).
- [6] P. D. Mannheim, “Alternatives to dark matter and dark energy,” *Prog. Part. Nucl. Phys.* **56**, 340 (2006) [[astro-ph/0505266](#)].
- [7] P. D. Mannheim and J. G. O’Brien, “Impact of a global quadratic potential on galactic rotation curves,” *Phys. Rev. Lett.* **106**, 121101 (2011) [[arXiv:1007.0970](#)].
- [8] P. D. Mannheim, “Making the Case for Conformal Gravity,” *Found. Phys.* **42**, 388 (2012) [[arXiv:1101.2186](#)].
- [9] H. Lu, Y. Pang, C. N. Pope and J. F. Vazquez-Poritz, “AdS and Lifshitz Black Holes in Conformal and Einstein-Weyl Gravities,” *Phys. Rev. D* **86**, 044011 (2012) [[arXiv:1204.1062](#)].
- [10] W. Xu and L. Zhao, “Critical phenomena of static charged AdS black holes in conformal gravity,” *Phys. Lett. B* **736**, 214 (2014) [[arXiv:1405.7665](#)].
- [11] H. Xu, Y. Sun and L. Zhao, “Black hole thermodynamics and heat engines in conformal gravity,” *Int. J. Mod. Phys. D* **26**, no. 13, 1750151 (2017) [[arXiv:1706.06442](#)].
- [12] H. Xu and M. H. Yung, “On the thermodynamic phase structure of conformal gravity,” *Phys. Lett. B* **783**, 36 (2018) [[arXiv:1804.10446](#)].
- [13] D. N. Page, “Particle Emission Rates from a Black Hole: Massless Particles from an Uncharged, Nonrotating Hole,” *Phys. Rev. D* **13**, 198 (1976).
- [14] D. N. Page, “Hawking radiation and black hole thermodynamics,” *New J. Phys.* **7**, 203 (2005) [[hep-th/0409024](#)].
- [15] D. N. Page, “Finite upper bound for the Hawking decay time of an arbitrarily large black hole in anti-De Sitter spacetime,” *Phys. Rev. D* **97**, no. 2, 024004 (2018) [[arXiv:1507.02682](#)].
- [16] Y. C. Ong, “Hawking Evaporation Time Scale of Topological Black Holes in Anti-de Sitter Spacetime,” *Nucl. Phys. B* **903**, 387 (2016) [[arXiv:1507.07845](#)].
- [17] S. J. Avis, C. J. Isham and D. Storey, “Quantum Field Theory in anti-De Sitter Space-Time,” *Phys. Rev. D* **18**, 3565 (1978).
- [18] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” [*Adv. Theor. Math. Phys.* **2**, 231 (1998)] [[hep-th/9711200](#)].
- [19] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [[hep-th/9802150](#)].
- [20] R. J. Riegert, “Birkhoff’s Theorem in Conformal Gravity,” *Phys. Rev. Lett.* **53**, 315 (1984).
- [21] D. Klemm, “Topological black holes in Weyl conformal gravity,” *Class. Quant. Grav.* **15**, 3195 (1998) [[gr-qc/9808051](#)].
- [22] G. W. Gibbons, R. Kallosh and B. Kol, “Moduli, scalar charges, and the first law of black hole thermodynamics,” *Phys. Rev. Lett.* **77**, 4992 (1996) [[hep-th/9607108](#)].
- [23] D. Kastor, S. Ray and J. Traschen, “Smarr Formula and an Extended First Law for Lovelock Gravity,” *Class. Quant. Grav.* **27**, 235014 (2010) [[arXiv:1005.5053](#)].
- [24] W. Xu, H. Xu and L. Zhao, “Gauss-Bonnet coupling constant as a free thermodynamical variable and the associated criticality,” *Eur. Phys. J. C* **74**, 2970 (2014) [[arXiv:1311.3053](#)].
- [25] A. Anabalón, D. Astefanesei, D. Choque and C. Martínez, “Trace Anomaly and Counterterms in Designer Gravity,” *JHEP* **1603**, 117 (2016) [[arXiv:1511.08759](#)].
- [26] D. Astefanesei, R. Ballesteros, D. Choque and R. Rojas, “Scalar charges and the first law of black hole thermodynamics,” *Phys. Lett. B* **782**, 47 (2018) [[arXiv:1803.11317](#)].
- [27] D. Kastor, S. Ray and J. Traschen, “Enthalpy and the Mechanics of AdS Black Holes,” *Class. Quant. Grav.* **26**, 195011 (2009) [[arXiv:0904.2765](#)].
- [28] D. Kubiznak and R. B. Mann, “P-V criticality of charged AdS black holes,” *JHEP* **1207**, 033 (2012) [[arXiv:1205.0559](#)].
- [29] R. B. Mann, “Black Holes: Thermodynamics, Information, and Firewalls,” Springer International Publishing, 2015:1-95
- [30] R. Gregory, D. Kastor and J. Traschen, “Black Hole Thermodynamics with Dynamical Lambda,” *JHEP* **1710**, 118 (2017) [[arXiv:1707.06586](#)].
- [31] S. Gunasekaran, R. B. Mann and D. Kubiznak, “Extended phase space thermodynamics for charged and rotating black holes and Born-Infeld vacuum polarization,” *JHEP* **1211**, 110 (2012) [[arXiv:1208.6251](#)].
- [32] R. G. Cai, L. M. Cao, L. Li and R. Q. Yang, “P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space,” *JHEP* **1309**, 005 (2013) [[arXiv:1306.6233](#)].
- [33] N. Altamirano, D. Kubiznak, R. B. Mann and Z. Sherkatghanad, “Kerr-AdS analogue of triple point and solid/liquid/gas phase transition,” *Class. Quant. Grav.* **31**, 042001 (2014) [[arXiv:1308.2672](#)].
- [34] D. C. Zou, S. J. Zhang and B. Wang, “Critical behavior of Born-Infeld AdS black holes in the extended phase space thermodynamics,” *Phys. Rev. D* **89**, no. 4, 044002 (2014) [[arXiv:1311.7299](#)].
- [35] N. Altamirano, D. Kubiznak, R. B. Mann and Z. Sherkatghanad, “Thermodynamics of rotating black holes and black rings: phase transitions and thermodynamic volume,” *Galaxies* **2**, 89 (2014) [[arXiv:1401.2586](#)].
- [36] S. W. Wei and Y. X. Liu, “Triple points and phase diagrams in the extended phase space of charged Gauss-Bonnet black holes in AdS space,” *Phys. Rev. D* **90**, no. 4, 044057 (2014) [[arXiv:1402.2837](#)].
- [37] C. V. Johnson, “Holographic Heat Engines,” *Class. Quant. Grav.* **31**, 205002 (2014) [[arXiv:1404.5982](#)].
- [38] H. Xu, W. Xu and L. Zhao, “Extended phase space thermodynamics for third order Lovelock black holes in diverse dimensions,” *Eur. Phys. J. C* **74**, no. 9, 3074 (2014) [[arXiv:1405.4143](#)].
- [39] A. M. Frassino, D. Kubiznak, R. B. Mann and F. Simovic, “Multiple Reentrant Phase Transitions and Triple Points in Lovelock Thermodynamics,” *JHEP* **1409**, 080 (2014) [[arXiv:1406.7015](#)].
- [40] B. P. Dolan, A. Kostouki, D. Kubiznak and R. B. Mann, “Isolated critical point from Lovelock gravity,” *Class. Quant. Grav.* **31**, no. 24, 242001 (2014) [[arXiv:1407.4783](#)].
- [41] J. Xu, L. M. Cao and Y. P. Hu, “P-V criticality in the extended phase space of black holes in massive gravity,” *Phys. Rev. D* **91**, no. 12, 124033 (2015) [[arXiv:1506.03578](#)].
- [42] H. Xu and Z. M. Xu, “Maxwells equal area law for Lovelock thermodynamics,” *Int. J. Mod. Phys. D* **26**,

- no. 04, 1750037 (2016) [arXiv:1510.06557].
- [43] H. Liu and X. h. Meng, “PCV criticality in the extended phase space of charged accelerating AdS black holes,” *Mod. Phys. Lett. A* **31**, no. 37, 1650199 (2016) [arXiv:1607.00496].
- [44] B. R. Majhi and S. Samanta, “P-V criticality of AdS black holes in a general framework,” *Phys. Lett. B* **773**, 203 (2017) [arXiv:1609.06224].
- [45] Y. B. Ma, L. C. Zhang, J. Liu, R. Zhao and S. Cao, “The thermodynamic relationship between the RN-AdS black holes and the RN black hole in canonical ensemble,” *Adv. High Energy Phys.* **2017**, 3812063 (2017) [arXiv:1612.03518].
- [46] D. C. Zou, R. Yue and M. Zhang, “Reentrant phase transitions of higher-dimensional AdS black holes in dRGT massive gravity,” *Eur. Phys. J. C* **77**, no. 4, 256 (2017) [arXiv:1612.08056].
- [47] K. Bhattacharya and B. R. Majhi, “Thermogeometric description of the van der Waals like phase transition in AdS black holes,” *Phys. Rev. D* **95**, no. 10, 104024 (2017) [arXiv:1702.07174].
- [48] D. C. Zou, Y. Liu and R. H. Yue, “Behavior of quasinormal modes and Van der Waals-like phase transition of charged AdS black holes in massive gravity,” *Eur. Phys. J. C* **77**, no. 6, 365 (2017) [arXiv:1702.08118].
- [49] K. Bhattacharya, B. R. Majhi and S. Samanta, “Van der Waals criticality in AdS black holes: a phenomenological study,” *Phys. Rev. D* **96**, no. 8, 084037 (2017) [arXiv:1709.02650].
- [50] W. Xu, “ $\lambda$  phase transition in Horava gravity,” [arXiv:1804.03815].
- [51] B. Liu, Z. Y. Yang and R. H. Yue, “Tricritical point and solid/liquid/gas phase transition of higher dimensional AdS black hole in massive gravity,” [arXiv:1810.07885].
- [52] S. W. Hawking and D. N. Page, “Thermodynamics of Black Holes in anti-De Sitter Space,” *Commun. Math. Phys.* **87**, 577 (1983).
- [53] D. N. Page, “Particle Emission Rates from a Black Hole. 2. Massless Particles from a Rotating Hole,” *Phys. Rev. D* **14**, 3260 (1976).
- [54] W. A. Hiscock and L. D. Weems, “Evolution of Charged Evaporating Black Holes,” *Phys. Rev. D* **41**, 1142 (1990).
- [55] Y. C. Ong, B. McInnes and P. Chen, “Cold black holes in the Harlow-Chayden approach to firewalls,” *Nucl. Phys. B* **891**, 627 (2015) [arXiv:1403.4886].
- [56] Y. C. Ong and P. Chen, “Charge Loss (or the Lack Thereof) for AdS Black Holes,” *JHEP* **1406**, 061 (2014) [arXiv:1404.5215].
- [57] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “String theory and classical absorption by three-branes,” *Nucl. Phys. B* **499**, 217 (1997) [hep-th/9703040].
- [58] S. S. Gubser, “Can the effective string see higher partial waves?,” *Phys. Rev. D* **56**, 4984 (1997) [hep-th/9704195].
- [59] S. D. Mathur, “Absorption of angular momentum by black holes and D-branes,” *Nucl. Phys. B* **514**, 204 (1998) [hep-th/9704156].
- [60] D. Klemm and L. Vanzo, “Quantum properties of topological black holes,” *Phys. Rev. D* **58**, 104025 (1998) [gr-qc/9803061].
- [61] P. Kanti and J. March-Russell, “Calculable corrections to brane black hole decay. 1. The scalar case,” *Phys. Rev. D* **66**, 024023 (2002) [hep-ph/0203223].
- [62] P. Kanti and J. March-Russell, “Calculable corrections to brane black hole decay. 2. Greybody factors for spin 1/2 and 1,” *Phys. Rev. D* **67**, 104019 (2003) [hep-ph/0212199].
- [63] C. M. Harris and P. Kanti, “Hawking radiation from a (4+n)-dimensional black hole: Exact results for the Schwarzschild phase,” *JHEP* **0310**, 014 (2003) [hep-ph/0309054].
- [64] P. Kanti, “Black holes in theories with large extra dimensions: A Review,” *Int. J. Mod. Phys. A* **19**, 4899 (2004) [hep-ph/0402168].
- [65] P. Kanti, J. Grain and A. Barrau, “Bulk and brane decay of a (4+n)-dimensional Schwarzschild-de-Sitter black hole: Scalar radiation,” *Phys. Rev. D* **71**, 104002 (2005) [hep-th/0501148].
- [66] V. Cardoso, M. Cavaglia and L. Gualtieri, “Black Hole Particle Emission in Higher-Dimensional Spacetimes,” *Phys. Rev. Lett.* **96**, 071301 (2006) Erratum: [*Phys. Rev. Lett.* **96**, 219902 (2006)] [hep-th/0512002].
- [67] V. Cardoso, M. Cavaglia and L. Gualtieri, “Hawking emission of gravitons in higher dimensions: Non-rotating black holes,” *JHEP* **0602**, 021 (2006) [hep-th/0512116].
- [68] T. Harmark, J. Natario and R. Schiappa, “Greybody Factors for d-Dimensional Black Holes,” *Adv. Theor. Math. Phys.* **14**, no. 3, 727 (2010) [0708.0017].
- [69] P. Boonserm and M. Visser, “Bounding the greybody factors for Schwarzschild black holes,” *Phys. Rev. D* **78**, 101502 (2008) [0806.2209].
- [70] S. Chen and J. Jing, “Greybody factor for a scalar field coupling to Einstein’s tensor,” *Phys. Lett. B* **691**, 254 (2010) [1005.5601].
- [71] Y. G. Miao and Z. M. Xu, “Hawking Radiation of Five-dimensional Charged Black Holes with Scalar Fields,” *Phys. Lett. B* **772**, 542 (2017) [1704.07086].
- [72] C. Y. Zhang, P. C. Li and B. Chen, “Greybody factors for a spherically symmetric Einstein-Gauss-Bonnet-de Sitter black hole,” *Phys. Rev. D* **97**, no. 4, 044013 (2018) [1712.00620].
- [73] R. J. Adler, P. Chen and D. I. Santiago, “The Generalized uncertainty principle and black hole remnants,” *Gen. Rel. Grav.* **33**, 2101 (2001) [gr-qc/0106080].
- [74] P. Chen, Y. C. Ong and D. h. Yeom, “Black Hole Remnants and the Information Loss Paradox,” *Phys. Rept.* **603**, 1 (2015) [1412.8366].
- [75] Y. C. Ong, “An effective black hole remnant via infinite evaporation time due to generalized uncertainty principle,” *JHEP* **1810**, 195 (2018) [1806.03691].
- [76] Y. Yao, M. S. Hou and Y. C. Ong, “A Complementary Third Law for Black Hole Thermodynamics,” [1812.03136].