

Functions with constant Laplacian satisfying Robin boundary conditions on an ellipse

Grant Keady and Benchawan Wiwatanapataphee
Department of Mathematics and Statistics
Curtin University, Bentley, 6102
Western Australia

July 16, 2020

Abstract

We study the problem of finding functions, defined within and on an ellipse, whose Laplacian is -1 and which satisfy a homogeneous Robin boundary condition on the ellipse. The parameter in the Robin condition is denoted by β . The integral of the solution over the ellipse, denoted by Q , is a quantity of interest in some physical applications. The dependence of Q on β and the ellipse's geometry is found. Several methods are used.

- Part I: To find the general solution the boundary value problem is formulated in elliptic cylindrical coordinates. Truncated Fourier series solution are then used to achieve good numerical approximations. The truncation is higher order than earlier papers and also totally systematic. Various asymptotic approximations are found directly from the pde formulations, this being far easier than from our series solution. An ad-hoc approximation that the solutions might have nearly elliptic level curves also leads to estimates. Numerical results from the various approaches are compared.
- Part II. Fourier series are much more directly applicable to a Problem $(P(\infty))$, this problem first arising in connection with asymptotics as

β tends to infinity. This problem is solved by Fourier series and a difference equation which would facilitate rapid computation of the coefficients is studied.

- Part III. A variational approach applicable to any cross-section – requiring however information on the solution of Problem (P(∞)) – yields simple and accurate lower bounds. The method was first illustrated with the rectangle as an example as for that cross-section Problem (P(∞)) is easy to solve. (This material is now published in *J. Fluids Engineering*.)
- Part IV. The approximation – lower bound – described in Part III is further approximated in the case of an ellipse. A variational approach to Problem (P(∞)) is used, and connections made to earlier approximations in Part I and to the Fourier Series in Part II.

One notational blemish is that our original usual notation for the perimeter $|\partial\Omega|$ is L but a referee requested, for engineering journals, using P . The latter is used consistently in Part IV.

This arXiv preprint will be referenced by journal articles we submit on parts of the work. The arXiv version contains material, e.g. codes for calculating Q , not in the very much shorter journal version.

Part I is joint work. For the remaining parts GK is the author.

Part I: Approximations by Fourier series, asymptotics and an ad hoc variational guess

1 Introduction

1.1 The pde problem

With Ω a plane domain and $\beta \geq 0$ given, we seek the solution of

$$-\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = 1 \quad \text{in } \Omega, \quad u + \beta \frac{\partial u}{\partial n} = 0 \quad \text{on } \partial\Omega. \quad (\text{P}(\beta))$$

Under reasonable conditions on the boundary $\partial\Omega$ (classical) solutions exist and, by Maximum Principle arguments, are unique and positive in Ω . [37] surveys some of the easier results.

In the context of flows, this boundary condition is called a *slip boundary condition* or *Navier's boundary condition*; in the wider mathematical literature it is called a *Robin boundary condition*. The $\beta = 0$ case is, in the fluid mechanics context, called a ‘no-slip boundary condition’: in elasticity the $\beta = 0$ case is called ‘the elastic torsion problem’. A functional of interest, in the context of flows, is the volume flow rate

$$Q := \int_{\Omega} u. \quad (1.1)$$

We remark that the same pde problem arises in contexts other than slip flow, for example, heat-flow with ‘Newton’s law of cooling’, e.g. [37, 48]. While there are other applications of this pde problem we will, on occasions, use the fluid flow terminology for u , Q and β .

There are many inequalities established for Problem (P(β)). There are isoperimetric inequalities, reviewed in [40] and others, for example, from [37]

$$\max \left(\beta \frac{|\Omega|^2}{|\partial\Omega|} + Q_0, \beta \frac{|\Omega|^2}{|\partial\Omega|} + \Sigma_{\infty} + \frac{\Sigma_1}{\beta} \right) \leq Q(\beta) \leq \beta \frac{|\Omega|^2}{|\partial\Omega|} + \Sigma_{\infty} = U(\beta), \quad (1.2)$$

where $\Sigma_{\infty} > 0$ and $\Sigma_1 \leq 0$ are defined in equation (4.2) at the beginning of §4 and used associated with asymptotics for β large. $U(\beta)$ is defined as the expression immediately to its left. These old inequalities (1.2) are established using the variational formulation of §4 in [37] : see inequalities (4.4) and (4.9).

The inequality (4.4) of [37] is well-known, (4.9) less so. The same methods lead to an improvement given in inequality (14.4).

Except in this subsection and a very small number of introductory subsections, e.g. §4.1, §4.2 and §6.1 where we allow Ω to be more general, we will be treating Ω as the elliptical domains within ellipses described by

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} \leq 1 \quad \text{with } a \geq b. \quad (1.3)$$

In Part III we apply the same methods to rectangular domains.

In this Part, we derive, for elliptical domains, a Fourier series form of the analytical solution of the problem and provide many checks on it. Others, e.g. [19], have used similar separation of variables approaches but have chosen to approximate the Fourier series of the function g occurring in the boundary condition, equation (3.11) by its first two terms, whereas we treat this more carefully.

Some recent papers treat the problem in the context of slip flows in elliptic microchannels, e.g. [17, 19, 20]. (Again in the context of flows in microchannels, time-dependent and other aspects are treated in various papers. [28, 29, 57, 58] study pulsatile flows. Papers [18, 64] are relevant to studies of the decay of transients and starting flows. Other physical effects are studied in [24, 31]. The steady flow case is, for each of these, a special case when a parameter is set to zero.) We make considerable effort to check our results. For example, there are a number of rigorously proved bounds. An old variational result, a lower bound on Q is equation (4.4) of [37],

$$\begin{aligned} Q(\beta) &\geq Q(\beta = 0) + \frac{\beta |\Omega|^2}{|\partial\Omega|} \quad \text{for general } \Omega, \\ &= \frac{\pi a^3 b^3}{4(a^2 + b^2)} + \frac{\pi^2 \beta a^2 b^2}{|\partial\Omega|} \quad \text{for ellipse } \Omega. \end{aligned} \quad (1.4)$$

Inequality (1.4) and the more general form preceding it are very easy to establish: see §4. Inequality (1.4) becomes an equality when the ellipse is a circle, i.e. $b = a$. The left-hand term in the right-hand side of inequality (1.4) gives the dominant term in the asymptotic expansion of Q as β tends to 0. The right-hand term in inequality (1.4) gives the dominant term in the asymptotic expansion of Q as β tends to infinity. However (except when $b = a$), in both asymptotic limits, the other term is not the next term in the asymptotic expansion.

We sometimes denote the area $|\Omega|$ by A and the perimeter $|\partial\Omega|$ by L . For an ellipse, the formula for L is given in equation (6.1).

A much more recent inequality, a generalization of the St Venant Inequality (see [52]), is given in [9] (with related work in [7]):

$$Q \leq Q_{\circ}(\beta) := \frac{\pi(ab)^{3/2}}{8} \left(\sqrt{ab} + 4\beta \right). \quad (1.5)$$

In words: amongst all ellipses with a given area that which has the greatest Q is the circular disk. See equation (2.2). (This, and other isoperimetric inequalities, are reviewed, in the context of microchannels in [38, 40].)

1.2 The structure of this Part

The rest of this part is organized as follows.

- In §2 we summarize the very well-known elementary solutions, polynomial in Cartesian coordinates, for the simplest special cases against which to check our general solution. Each of them forms the lowest order term in asymptotics developed later in this part.
- In §3, the boundary value problem is formulated in elliptic cylindrical coordinates, and then the series representation is approximated (to any accuracy desired).
- The next sections provide many checks. To reduce the number of parameters, in these sections, we take $b = 1/a$, so the area of the ellipse is π .
 - In §4 we present a variational formulation and associated bounds.
 - In §5 we present asymptotics for the situation where the ellipse is nearly circular.
 - In §6 we give asymptotics for β small.
 - In §7 we study asymptotics for β large.
- Finally, a conclusion is given in §8.

- §9, Appendix A, treats various geometric matters.
- §10, Appendix B, has some notes on elliptic integrals. These arise in the Fourier series for a functions g and \hat{g} used at several sections of this work, notably §3 and developments from §7 given in Part II.
- §11, Appendix C, gives some numerical examples, its first parts comparing our numerics with some previously published examples. We conclude this appendix with an attempt to suggest numbers which might arise in connection with possible applications involving blood flow.

2 Some simple explicit solutions:

u quadratic polynomial in x, y

2.1 Circular cross-section with $\beta \geq 0$

When Ω is circular, radius a , in polar coordinates, $r = \sqrt{x^2 + y^2}$, the solution is

$$u_{\beta\odot} = u = \frac{1}{4}(a^2 - r^2) + \frac{\beta a}{2}. \quad (2.1)$$

In the context of fluid flows, when $\beta = 0$ this is Poiseuille flow.

$$Q_{\beta\odot} = Q(\beta) = \frac{\pi a^3}{8}(a + 4\beta). \quad (2.2)$$

See also [43]§331, p586.

In later sections the circle will have radius $a = 1$, area π .

2.2 Elliptic cross-section with $\beta = 0$

In this subsection (and in sections §4 to §7) we consider ellipses with $b = 1/a$ and $a \geq 1$. The $\beta = 0$ solution dates back at least to St Venant. In polar coordinates with the origin at the centroid of the ellipse the velocity u_e is, in Cartesian coordinates, a quadratic polynomial in x and y while, in polar coordinates, it is

$$u_{0e} = \frac{1}{4} \left(\sqrt{1 - \epsilon^2} - r^2 + \epsilon r^2 \cos(2\theta) \right).$$

Here

$$\epsilon = \frac{a^2 - a^{-2}}{a^2 + a^{-2}}.$$

In Cartesian coordinates

$$u_0 = \frac{1 - (x/a)^2 - (y/b)^2}{2/a^2 + 2/b^2}. \quad (2.3)$$

Also

$$Q_{0e} = Q(\text{ellipse}, \beta = 0) = \frac{\pi}{4(a^2 + a^{-2})} = \frac{\pi}{8} \sqrt{1 - \epsilon^2}. \quad (2.4)$$

Also relevant is e the eccentricity as defined in equation (9.3).




Figure 1: $\beta = 0$: plot of $\text{Qratio} = (Q_{\odot} - Q(e))/Q_{\odot}$ against eccentricity, e .

There are various interesting alternative expressions for Q . The ellipse's moments of inertia are

$$I_{xx} = \frac{\pi a^2}{4}, \quad I_{yy} = \frac{\pi}{4a^2} \text{ so } Q = J := \frac{I_{xx}I_{yy}}{I_{xx} + I_{yy}}.$$

See [52] p112. For any domain (and $\beta = 0$) it is known that $Q \leq J$.

3 Fourier series solution

3.1 Using elliptic coordinates

Formulae when $a \geq b > 0$

For the geometry of elliptic domains specified as in equation (1.3) with, as always, $a > b$, it is convenient here to use elliptic coordinates (η, ψ) which are related to the rectangular coordinates (as in [17] equation (18)) by

$$x = c \cosh(\eta) \cos(\psi); \quad y = c \sinh(\eta) \sin(\psi);, \quad (3.1)$$

where $0 \leq \eta \leq \infty$, $0 \leq \psi \leq 2\pi$, and c and $-c$ are two common foci of the ellipse. The origin of coordinates in the Cartesian system (the centre of our ellipse) is at $\eta = 0$, $\psi = \pi/2$. The upper side of the positive semi-major axis is $\eta = 0$, $0 < \psi < \pi/2$: the lower side is $\eta = 0$, $0 > \psi > -\pi/2$. Let a and b denote lengths of the semi-axes of the ellipse and $a > b > 0$. Except in this subsection, and elsewhere when so specified, we take $b = 1/a$. Returning to the general situation,

$$a = c \cosh(\eta) \quad \text{and} \quad b = c \sinh(\eta), \quad \text{and} \quad c = \sqrt{a^2 - b^2}.$$

Defining η_0 (as in [17] equation (19)) by

$$\eta_0 = \ln \frac{1 + b/a}{\sqrt{1 - (b/a)^2}} = \operatorname{arctanh}\left(\frac{b}{a}\right),$$

the boundary of the ellipse can be represented by

$$x = c \cosh(\eta_0) \cos(\psi), \quad y = \sinh(\eta_0) \sin(\psi),$$

so that

$$c = \frac{a}{\cosh \eta_0} = \frac{b}{\sinh \eta_0}.$$

The eccentricity is

$$e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \tanh^2(\eta_0)} = \frac{1}{\cosh(\eta_0)}.$$

(Nearly circular ellipses have η_0 large and c near 0.)

The Jacobian J of the transformation from (x, y) -coordinates to (η, ψ) coordinates is

$$J(\eta, \psi) = c^2 (\cosh^2(\eta) - \cos^2(\psi)) = \frac{c^2}{2} (\cosh(2\eta) - \cos(2\psi)).$$

This is needed in the derivation of equation (3.33). The Jacobian will be used in other places in this work, e.g. §12.1.

Formulae when $b = 1/a$

In all major calculations henceforth we scale distances so that we have $b = 1/a$. Formulae used later include the following:

$$c^2 = \frac{2}{\sinh(2\eta_0)} = a^2 - \frac{1}{a^2} = \frac{e^2}{\sqrt{1 - e^2}}, \quad (3.2)$$

$$\tanh(\eta_0) = \frac{1}{a^2}, \quad (3.3)$$

$$\cosh(\eta_0) = \frac{1}{e}, \quad (3.4)$$

$$e^2 = 1 - \frac{1}{a^4}. \quad (3.5)$$

3.2 The pde and boundary conditions

The pde of Problem (P(β)) is recast in these coordinates as

$$\begin{aligned} \frac{\partial^2 u}{\partial \eta^2} + \frac{\partial^2 u}{\partial \psi^2} &= -c^2 (\cosh^2(\eta) - \cos^2(\psi)), \\ &= -\frac{c^2}{2} (\cosh(2\eta) - \cos(2\psi)). \end{aligned} \quad (3.6)$$

We will solve the pde (3.6) subject to boundary conditions (3.7), (3.8) and (3.9). The solution is assumed to be symmetric about the mid-plane for $\psi = 0, \frac{\pi}{2}$. Therefore, symmetric boundary condition is applied, i.e.,

$$\frac{\partial u}{\partial \psi} = 0 \text{ on } \psi = 0, \frac{\pi}{2}. \quad (3.7)$$

In addition, we set

$$\frac{\partial u}{\partial \eta} = 0 \text{ on } \eta = 0. \quad (3.8)$$

The boundary condition on the ellipse, i.e. at those points when $\eta = \eta_0$ is

$$u(\eta_0, \psi) + \frac{\beta}{c\sqrt{\cosh^2 \eta_0 - \cos^2 \psi}} \frac{\partial u}{\partial \eta}(\eta_0, \psi) = 0. \quad (3.9)$$

Define

$$g(\psi) = \frac{\cosh(\eta_0)}{(\cosh^2(\eta_0) - \cos^2(\psi))^{1/2}}. \quad (3.10)$$

Boundary condition (3.9) becomes

$$u(\eta_0, \psi) + \frac{\beta}{c \cosh(\eta_0)} g(\psi) \frac{\partial u}{\partial \eta}(\eta_0, \psi) = 0. \quad (3.11)$$

In summary, the problem in elliptic coordinates is given as the following boundary value problem (BVP).

BVP: Find $u(\eta, \psi)$ such that the above governing equation (3.6) and associated boundary conditions (3.7), (3.8) and (3.11) are satisfied.

3.3 The Fourier series for g

Ellipses in general

The definition of g from (3.10) can be re-cast in several ways:

$$\begin{aligned} g(\psi) &= \left(1 - \frac{\cos^2(\psi)}{\cosh^2(\eta_0)}\right)^{-1/2}, \\ &= C g_*(\psi) \text{ with } g_*(\psi) = \left(1 - \frac{\cos(2\psi)}{\cosh(2\eta_0)}\right)^{-1/2} \text{ and } C = \left(\frac{1 + \cosh(2\eta_0)}{\cosh(2\eta_0)}\right)^{1/2}. \end{aligned}$$

We will, later, use a more economic notation using

$$q = \cosh^2(\eta_0), \quad (3.12)$$

and the form for g_* is consistent with the occurrence, in Fourier coefficients, of polynomials in $\cosh(2\eta_0) = 2q - 1$.

The Fourier series for g is used. Define

$$g_n = \frac{2}{\pi} \int_0^\pi g(\psi) \cos(2n\psi) d\psi, \quad (3.13)$$

so

$$g(\psi) = \frac{g_0}{2} + \sum_{n=1}^{\infty} g_n \cos(2n\psi),$$

In various approximations the first few terms in the Fourier series are used. For these define

$$\text{EllipticE}_0 = \text{EllipticE}\left(\frac{1}{\cosh(\eta_0)}\right) \quad \text{and} \quad \text{EllipticK}_0 = \text{EllipticK}\left(\frac{1}{\cosh(\eta_0)}\right). \quad (3.14)$$

We have

$$g_0 = \frac{4}{\pi} \text{EllipticK}_0, \quad (3.15)$$

$$g_1 = -\frac{8 \cosh(\eta_0)^2}{\pi} \text{EllipticE}_0 + \frac{4(2 \cosh(\eta_0)^2 - 1)}{\pi} \text{EllipticK}_0. \quad (3.16)$$

All the terms g_n can be written in terms of Legendre Q functions though we first encountered them in the form presented by maple,

$$g_n = \frac{4}{\pi} (E_n \text{EllipticE}_0 + K_n \text{EllipticK}_0), \quad (3.17)$$

where E_n and K_n are polynomials of degree n in $\cosh(\eta_0)^2$ with rational number coefficients.

Should future work require more terms we remark that a three term recurrence relation exists to determine the polynomials E_n and K_n . See Part II § 13. In the notation of equation (3.12), the K_n sequence of polynomials starts with

$$K_0 = 1, \quad K_1 = 2q - 1 = q_2.$$

The E_n sequence of polynomials starts with

$$E_0 = 0, \quad E_1 = 2q = q_2 - 1.$$

There are also other representations: see Part II § 13. There are several very elementary facts which have some use. These include that

$$\cos(2k\psi) = \text{ChebyshevT}(k, \cos(2\psi)).$$

Also, the Taylor series

$$(1 - X)^{-1/2} = \sum_{k=0}^{\infty} \frac{(2k)!}{2^{2k} (k!)^2} X^k,$$

can be applied to the preceding expressions for g to obtain series which are sums of powers of $\cos(\psi)^2$ or of $\cos(2\psi)$. To date, our only use of these has been in checking the asymptotic approximations, at small eccentricity, to the Fourier coefficients of g and of $1/g$.

The reciprocal of g also occurs in later calculations (and also in [17] equation (61) though they use a Taylor series rather than Fourier series used here).

Approximating the Fourier series for nearly circular ellipses

Nearly circular ellipses have η_0 large. Then the lowest approximation is

$$g(\psi) \sim g_\epsilon(\psi) = 1 + \frac{\cos^2(\psi)}{2 \cosh^2(\eta_0)} = 1 + \frac{1}{4 \cosh^2(\eta_0)} + \frac{\cos(2\psi)}{4 \cosh^2(\eta_0)},$$

and

$$\frac{1}{g(\psi)} \sim 1 - \frac{1}{4 \cosh^2(\eta_0)} - \frac{\cos(2\psi)}{4 \cosh^2(\eta_0)}. \quad (3.18)$$

We will use the next approximation after this, and will use $e = 1/\cosh(\eta_0)$:

$$\frac{1}{g(\psi)} \sim 1 - \frac{1}{4}e^2 - \frac{3}{64}e^4 - \left(\frac{1}{4}e^2 + \frac{1}{16}e^4\right) \cos(2\psi) - \frac{1}{64}e^4 \cos(4\psi). \quad (3.19)$$

We have, for e small the following Fourier coefficients of $\hat{g} = 1/g$:

$$\hat{g}_0 \sim 2 - \frac{1}{2}e^2 - \frac{3}{32}e^4 \quad (3.20)$$

$$\hat{g}_1 \sim -\frac{1}{4}e^2 - \frac{1}{16}e^4 \quad (3.21)$$

$$\hat{g}_2 \sim -\frac{1}{64}e^4 \quad (3.22)$$

(We will need these in the treatment of u_∞ . See equation (12.1).)

3.4 Completing the Fourier series solution

The general solution of equation (3.6) can be expressed in the form of

$$u(\eta, \psi) = v_h + v_p, \quad (3.23)$$

where v_h and v_p denote the homogeneous and the particular solution of the pde (3.6) subjected to the boundary conditions (3.7), (3.8) and (3.11).

For the particular solution v_p , from the right hand side of the equation (3.6), we can express v_p in the form of

$$v_p = \hat{g}_1 \cosh(2\eta) + \hat{g}_2 \sinh(2\eta) + \hat{g}_3 \cos(2\psi) + \hat{g}_4 \sin(2\psi), \quad (3.24)$$

which gives

$$\frac{\partial^2 v_p}{\partial \eta^2} = 2\hat{g}_1 \sinh(2\eta) + 2\hat{g}_2 \cosh(2\eta); \quad (3.25)$$

$$\frac{\partial^2 v_p}{\partial \psi^2} = -2\hat{g}_3 \sin(2\psi) + 2\hat{g}_4 \cos(2\psi). \quad (3.26)$$

Substituting v_p into the pde (3.6) and comparing the coefficients of $\cosh(2\eta)$ and $\cos(2\psi)$ yields

$$v_p = -\frac{c^2}{8}(\cosh(2\eta) + \cos(2\psi)) \left(= -\frac{1}{4}(x^2 + y^2) \right). \quad (3.27)$$

By separation of variables, the solution v_h of the Laplace equation subject to above boundary conditions (3.7) and (3.8) is

$$v_h = \sum_{n=0}^{\infty} A_n \cosh(2n\eta) \cos(2n\psi). \quad (3.28)$$

Hence, the general solution can be expressed in the form of

$$u(\eta, \psi) = v_h + v_p = \sum_{n=0}^{\infty} A_n \cosh(2n\eta) \cos(2n\psi) - \frac{c^2}{8}(\cosh(2\eta) + \cos(2\psi)), \quad (3.29)$$

where term A_n can be determined by using the boundary condition (3.11). We then obtain

$$\begin{aligned} \sum_{n=0}^{\infty} A_n \cosh(2n\eta_0) \cos(2n\psi) - \frac{c^2}{8}(\cosh(2\eta_0) + \cos(2\psi)) + \frac{\beta g(\psi)}{c \cosh(\eta_0)} \\ \sum_{n=0}^{\infty} n 2A_n \sinh(2n\eta_0) \cos(2n\psi) - \frac{c^2 \beta g(\psi)}{4c \cosh(\eta_0)} \sinh(2\eta_0) = 0. \end{aligned} \quad (3.30)$$

or

$$\begin{aligned} \sum_{n=0}^{\infty} A_n \left(\cosh(2n\eta_0) + \frac{2n\beta g(\psi)}{c \cosh(\eta_0)} \sinh(2n\eta_0) \right) \cos(2n\psi) \\ = \frac{c^2}{8}(\cosh(2\eta_0) + \cos(2\psi)) + \frac{c^2 \beta g(\psi)}{4c \cosh(\eta_0)} \sinh(2\eta_0) = f(\psi). \end{aligned} \quad (3.31)$$

As g is an even function so is f and h_n where

$$h_n(\psi) = \cosh(2n\eta_0) + \frac{2n\beta g(\psi)}{c \cosh(\eta_0)} \sinh(2n\eta_0).$$

(The Fourier coefficients of f and of h_n are very simply related to the Fourier coefficients of g .) We then have

$$\sum_{n=1}^{\infty} A_n \int_{-\pi}^{\pi} h_n(\psi) \cos(2n\psi) \cos(2m\psi) d\psi = - \int_{-\pi}^{\pi} f(\psi) \cos(2m\psi) d\psi. \quad (3.32)$$

Hence if we truncate the summation at n_{\max} we have a system of linear equations for the A_n for n up to n_{\max} . Using the A_n so found, we obtain an approximation to the Laplace solution v_h and then steady-state fluid flow, $u(\eta, \psi) = v_h + v_p$ in the elliptical channel. Finally, the volume flow rate of fluid passing through a elliptical cross sectional area, Q , can determined by

$$Q = c^2 \int_0^{2\pi} \int_0^{\eta_0} u(\eta, s) (\cosh^2(\eta) - \cos^2(s)) d\eta ds. \quad (3.33)$$

3.5 Q from the Fourier series of v_h

We will substitute $u = v_p + v_h$ with v_h given by its Fourier series into the formula for Q given in (3.33). Note that all the Fourier components with $n > 1$ contribute nothing to the integral. First

$$\begin{aligned} Q_p &= c^2 \int_0^{2\pi} \int_0^{\eta_0} v_p (\cosh^2(\eta) - \cos^2(s)) d\eta ds, \\ &= -\frac{1}{4} I_2 = -\frac{1}{4} \frac{\pi}{4} \left(a^2 + \frac{1}{a^2} \right) = -\frac{\pi}{16} \frac{2 - e^2}{\sqrt{1 - e^2}}, \end{aligned}$$

I_2 being the polar area moment of inertia about the centroid. On using $c^2 = 2/\sinh(2\eta_0)$,

$$\begin{aligned} Q &= Q_p + c^2 \int_0^{2\pi} \int_0^{\eta_0} (A_0 + A_1 \cosh(2\eta) \cos(2s)) (\cosh^2(\eta) - \cos^2(s)) d\eta ds, \\ &= -\frac{\pi}{8} \frac{\cosh(2\eta_0)}{\sinh(2\eta_0)} + \pi A_0 - \frac{1}{2} \pi A_1. \end{aligned} \quad (3.34)$$

```

uA := A0+A1*cosh(2*eta)*cos(2*psi)+A2*cosh(4*eta)*cos(4*psi);
  'cSqr Assign;1/'(sqrt(1/e^2-1)/e):
hypToe := -> subs(cosh(eta0) = 1/e, sinh(eta0) = sqrt(1/e), u) :
tmp := subs(cSqr = 1/(cosh(eta0)*sinh(eta0)), int(int(uA*cSqr*(cosh(eta)^2-cos(psi)
  A0*Pi -A1*Pi/2

```

The terms involving the A_j are checked in the following code:

4 Inequalities from a variational formulation

Variational formulations are available. For example, in [37] equation (4.1), the functional \mathcal{J} is defined as

$$\mathcal{J}(v) = \mathcal{A}(v) - \frac{1}{\beta}\mathcal{B}(v) \quad \text{where} \quad \mathcal{A}(v) = \int_{\Omega} (2v - |\nabla v|^2), \quad \mathcal{B}(v) = \int_{\partial\Omega} v^2. \quad (4.1)$$

For any simply-connected domain with sufficiently smooth boundary, the maximiser u of this functional solves Problem (P(β)). Furthermore, $\mathcal{J}(u)$ is $Q = \mathcal{Q}(u)$ where

$$\mathcal{Q}(v) = \int_{\Omega} v.$$

Thus we can find lower bounds for Q by evaluating $\mathcal{J}(v)$ for particular choices of v .

In this work, we have two distinct variational approximations. In the order of importance, though not of our discoveries, these are as follows.

- The more important of these – applying to general domains, not just ellipses and now published in [39] – is included here, in §4.1, as motivation for further work on Problem (P(∞)) for the ellipse. We have, in Part II, a series solution for Σ_{∞} for the ellipse involving Legendre functions but have yet to find Σ_1 . The lower bound is more easily illustrated with the rectangle (see [39]) here treated in Part III. For the rectangle (and some other domains, see [35, 36], u_{∞} is available analytically.
- The second of these is ad hoc, good when the solutions of Problem (P(β)) have nearly elliptical level curves. This is the case for the ellipse, and also, *though only for large β* for a number of other domains, including rectangles. General domains are treated in §4.2 and specialised to ellipses in §4.3.

4.1 A development from [37]

Our notation is as in [37]. For large β

$$u \sim \beta \frac{|\Omega|}{|\partial\Omega|} + u_{\infty} + o(1),$$

where u_{∞} solves

$$-\frac{\partial^2 u_{\infty}}{\partial x^2} - \frac{\partial^2 u_{\infty}}{\partial y^2} = 1 \text{ in } \Omega, \quad \frac{\partial u_{\infty}}{\partial n} = -\frac{|\Omega|}{|\partial\Omega|} \text{ on } \partial\Omega \text{ and } \int_{\partial\Omega} u_{\infty} = 0. \quad (\text{P}(\infty))$$

Define

$$\Sigma_\infty = \int_\Omega u_\infty, \quad \text{and} \quad \Sigma_1 = - \int_{\partial\Omega} u_\infty^2, \quad (4.2)$$

with u_∞ satisfying Problem (P(∞)). A lower bound on $Q(\beta)$ using geometric quantities – area and perimeter – and Q_0 and the two Σ functionals is presented in Part III §14.1.

4.2 Quadratic test functions, and a simple first case

For domains like ellipses symmetric about both the x -axis and y -axis one nice choice for v are the quadratic functions

$$(i) \quad v = c_0 + c_2 \left(\left(\frac{x}{a} \right)^2 + a^2 y^2 \right),$$

and, more generally (and with a different c_0)

$$(ii) \quad v = c_0 + c_{xx}x^2 + c_{yy}y^2.$$

(Case (ii) is motivated by the likelihood that for a range of β and a range of a the level curves of u look to be like ellipses. See, for example, Figure 4 of [60] and our own contour plots shown in Appendix C § 11.)

We briefly return to general domains. An application of the Divergence Theorem give

$$\text{if } -\Delta v = 1 \text{ then } \mathcal{J}(v) - \mathcal{Q}(v) = \int_{\partial\Omega} v \left(v + \beta \frac{\partial v}{\partial n} \right) \quad (4.3)$$

Case (i)

We begin with case (i). While the beginning of the study with case (ii) will allow Ω to be more general than merely an ellipse, for case (i) we restrict to the ellipse. The final result, lower bound, will give the same result as we reported in §1.1 from equation (4.4) of [37]. Write J or $J(c_0, c_2)$ for the value of \mathcal{J} evaluated at the quadratic function (i)

$$J = 2c_0A + 2c_2 \left(\frac{1}{a^2} \left(1 - \frac{2c_2}{a^2} \right) I_{XX} + a^2 \left(1 - 2a^2c_2 \right) I_{YY} \right) - (c_0 + c_2)^2 \frac{L}{\beta}.$$

Finding the gradient of J with respect to $[c_0, c_2]$ we find the maximum of J will occur when the gradient is 0, i.e.

$$\begin{aligned} Lc_0 + Lc_2 &= \pi\beta, \\ 2Lc_0 + c_2 \left(2L + \pi\beta \left(a^2 + \frac{1}{a^2} \right) \right) &= \pi\beta. \end{aligned}$$

This is readily solved, and on integrating, yields the result given before at inequality (1.4).

It is straightforward to show that the quadratic u which is the variational winner in Case (i) is such that the result in (4.3) can be applied to establish $\mathcal{J}(u) = \mathcal{Q}(u)$ (a fact which can be established by other means). The application of (4.3) is facilitated by the fact that the quadratic $u = u_0 + \beta|\Omega|/|\partial\Omega|$ is constant on $\partial\Omega$.

Case (ii), an introduction

Next we begin to treat case (ii). Write J or $J(c_0, c_{xx}, c_{yy})$ for the value of \mathcal{J} evaluated at the quadratic function (ii).

$$\begin{aligned} J &= 2(c_0A + c_{xx}(1 - 2c_{xx})I_{xx} + c_{yy}(1 - 2c_{yy})I_{yy}) - \\ &\quad \frac{1}{\beta} (c_0^2L + 2c_0c_{xx}i_{xx} + 2c_0c_{yy}i_{yy} + 2c_{xx}c_{yy}i_{xxyy} + c_{xx}^2i_{xxxx} + c_{yy}^2i_{yyyy}) . \end{aligned}$$

where A is the area, L the perimeter and the I are area moments and i boundary moments

$$I_{xx} = \int_{\Omega} x^2, \quad I_{yy} = \int_{\Omega} y^2,$$

and

$$i_{xx} = \int_{\partial\Omega} x^2, \quad i_{yy} = \int_{\partial\Omega} y^2,$$

and

$$i_{xxyy} = \int_{\partial\Omega} x^2y^2, \quad i_{xxxx} = \int_{\partial\Omega} x^4, \quad i_{yyyy} = \int_{\partial\Omega} y^4. \quad (4.4)$$

Define also

$$Q_q = c_0A + c_{xx}I_{xx} + c_{yy}I_{yy}, \quad (4.5)$$

the subscript q reminding us of the quadratic approximation to the velocity field. The gradient of J , here denoted \mathbf{g} , is

$$\mathbf{g} = -\frac{2}{\beta} \begin{pmatrix} c_0L + c_{xx}i_{xx} + c_{yy}i_{yy} - \beta A \\ c_0i_{xx} + c_{xx}(4\beta I_{xx} + i_{xxxx}) + c_{yy}i_{xxyy} - \beta I_{xx} \\ c_0i_{yy} + c_{xx}i_{xxyy} + c_{yy}(4\beta I_{yy} + i_{yyyy}) - \beta I_{yy} \end{pmatrix} .$$

We remark that

$$2(J - Q_q) = \mathbf{g}[1]c_0 + \mathbf{g}[2]c_{xx} + \mathbf{g}[3]c_{yy}. \quad (4.6)$$

This gradient \mathbf{g} is zero when

$$\begin{aligned} c_0 L + c_{xx} i_{xx} + c_{yy} i_{yy} &= \beta A, \\ c_0 i_{xx} + c_{xx}(4\beta I_{xx} + i_{xxxx}) + c_{yy} i_{xxyy} &= \beta I_{xx}, \\ c_0 i_{yy} + c_{xx} i_{xxyy} + c_{yy}(4\beta I_{yy} + i_{yyyy}) &= \beta I_{yy}. \end{aligned}$$

Up until now the c_0 , c_{xx} , c_{yy} have been general. Henceforth we use the same letters to denote the c obtained by solving $\mathbf{g} = 0$, the gradient of J equals zero. We remark that after finding c_0 , c_{xx} , c_{yy} there are two methods of estimating Q . On the one hand, we can estimate Q from

$$Q_c := c_0 A + c_{xx} I_{xx} + c_{yy} I_{yy}, \quad (4.7)$$

while, on the other hand, we have the lower bound

$$J_c := J(c_0, c_{xx}, c_{yy}) \leq Q.$$

From identity (4.6) we have $Q_c = J_c$.

For general Ω inequalities on moments (such as some in references in [34]) may lead to lower bounds for Q in terms of simpler geometric quantities. Also for some domains, polygons for example, the various moments could be calculated. In general the restriction of the test functions to quadratics isn't likely to give very tight lower bounds though there may be exceptions. Rectangles at large β might be one such as u_∞ (see [37] or equations (12.5)) is, for rectangles, quadratic.

For an ellipse with $a > 1$ (and $b = 1/a$) the area moments are elementary, but the perimeter and other boundary moments involve elliptic integrals: we present these in the next subsection. For a circle, all the moments are elementary. For a circle of radius 1, the moments are:

$$\begin{aligned} A &= \pi, \quad I_{xx} = \frac{\pi}{4}, \quad I_{yy} = \frac{\pi}{4}, \\ L &= 2\pi, \quad i_{xx} = \pi = i_{yy}, \quad i_{xxxx} = \frac{3\pi}{4} = i_{yyyy}, \quad i_{xxyy} = \frac{\pi}{4}. \end{aligned}$$

For the circle, the variational winner has c_0 , $c_{xx} = c_{yy}$ giving the exact solution previously presented in equation (2.1): both Q_c and J_c evaluate to the formula for Q given in equation (2.2).

A note of caution concerning the behaviour when $\beta = 0$ is appropriate here. If one naively sets $\beta = 0$ in the system of equations $\text{grad}(J) = 0$ the right-hand sides are all zero. The solution with all the c equal to zero incontrovertably gives a lower bound on Q : the lower bound being zero. Of course if $\beta = 0$ the term $\mathcal{B}(v)/\beta$ is the difficulty for functions v not vanishing identically over the whole boundary. As a quadratic function vanishing on the boundary provides the minimizer for ellipses with $\beta = 0$ it seems appropriate, at this stage, to return to the specifics for an ellipse.

4.3 Variational methods for the ellipse, Case (ii) continued

For the ellipse we can write i_{xx} and i_{yy} in terms of the polar second moment about the centre, $i_2 = i_{xx} + i_{yy}$, and the perimeter L . Similarly the 4-th moments can all be written in terms of the polar fourth moment about the centre, i_4 and i_2 and L . Each of L , i_2 and i_4 can be integrated in terms of elliptic integrals.

The variational winner over the quadratics gives lower bounds on Q as in the following code.

```

# FILE varlGenQfnmpl.txt

Qu0fn0:= proc(a,beta) # simplest as in KM93
  local Q0, ecc, L;
  Q0:= Pi/(4*(1/a^2 + a^2));
  ecc:= sqrt(1 - 1/a^4);
  L:= 4*a*EllipticE(ecc);
  Q0 + beta*Pi^2/L
end proc:

Qu0fn:= proc(a,beta) # quadratic test functions
  local ecc, Eecc, Kecc, dQu0s, nQu0s;
  ecc:= sqrt(1 - 1/a^4);
  Eecc:= EllipticE(ecc);
  Kecc:= EllipticK(ecc);
  dQu0s:= 64*a*(4*a^12-15*a^8+4-15*a^4)*Eecc^2+(512*a*(a^8+2*a^4+1)*Kecc+
          720*a^2*Pi*beta*(1-2*a^4+a^8))*Eecc-320*a*(a^4+1)*Kecc^2;
  nQu0s:= 16*a^3*(-19*a^4+4*a^8+4)*Pi*Eecc^2+
          (128*a^3*(a^4+1)*Pi*Kecc+3*Pi^2*beta*(-55*a^4-55*a^8+23+
          23*a^12))*Eecc-80*a^3*Pi*Kecc^2+
          24*Pi^2*beta*(7+7*a^8-6*a^4)*Kecc+180*Pi^3*beta^2*a*(1-2*a^4+a^8);
  nQu0s/dQu0s
end proc:

Qfn:= proc(a,beta) # both the above
  local Q0, ecc, Eecc, Kecc, QKM93, nQu0, dQu0beta0, dQu0;
  Q0:= Pi/(4*(1/a^2 + a^2));
  ecc:= sqrt(1 - 1/a^4);
  Eecc:= EllipticE(ecc);
  Kecc:= EllipticK(ecc);
  QKM93:= Q0 + beta*Pi^2/(4*a*Eecc);
  nQu0:= (4*(a^4+1)*Kecc+(a^8-10*a^4+1)*Eecc)^2;
  dQu0beta0:= (4*a^8-19*a^4+4)*Eecc^2+8*(a^4+1)*Eecc*Kecc-5*Kecc^2;
  dQu0:= 45*a^2*Pi*beta*(a^4-1)^2*(a^4+1)*Eecc+ 4*a*(a^4+1)^2*dQu0beta0;
  [QKM93, QKM93+ 5*Pi^2*beta*nQu0/(16*Eecc*dQu0)]
end proc:

```

5 Q for nearly circular ellipses

5.1 Q at $\beta = 0$

Now, at $\beta = 0$, the explicit solution as given in §2, yields asymptotics for $\epsilon \rightarrow 0$:

$$\begin{aligned} Q &= \frac{\pi}{4(a^2 + a^{-2})} = \frac{\pi}{8} \sqrt{1 - \epsilon^2}, \\ &\sim \frac{\pi}{8} (1 - 2(a - 1)^2 + O((a - 1)^3)) \quad \text{as } a \rightarrow 1, \\ &\sim \frac{\pi}{8} \left(1 - \frac{1}{2}\epsilon^2 + O(\epsilon^3) \right) \quad \text{as } \epsilon \rightarrow 0, \end{aligned} \quad (5.1)$$

5.2 Nearly circular ellipses with $\beta \geq 0$

The explicit solution available for when $\beta = 0$ is presented above, and is useful to check against our asymptotics for the solution when $\beta \geq 0$. When $\epsilon \rightarrow 0$, equation (2.4) agrees with the asymptotics given earlier in equation (5.1).

Concerning the ellipse when $\beta > 0$, when ϵ is small, u can be approximated in the form

$$u = \frac{1}{4} (1 - r^2) + \frac{\beta}{2} + \epsilon^2 t_{02} + \epsilon t_{11} r^2 \cos(2\theta) + \epsilon^2 t_{22} r^4 \cos(4\theta). \quad (5.2)$$

Details of the perturbation analysis are in [38]. The result of the perturbation analysis is

$$\begin{aligned} t_{11} &= \frac{1}{4} \frac{1 + \beta}{1 + 2\beta}, \\ t_{02} &= -\frac{1}{32} \frac{4 + 5\beta + 6\beta^2}{1 + 2\beta} = -\frac{1}{32} \left(1 + 3\beta + \frac{3}{1 + 2\beta} \right), \\ t_{22} &= -\frac{1}{32} \frac{\beta(1 - 2\beta)}{(1 + 4\beta)(1 + 2\beta)}. \end{aligned}$$

Integrating u with these parameters over the ellipse gives the expansion for $Q(\text{ellipse})$:

$$Q(\text{ellipse}) \sim \frac{\pi}{8} (1 + 4\beta) + q_1 \epsilon^2 \quad (5.3)$$

$$q_1 = -\frac{\pi}{16} (1 - 8t_{11} - 16t_{02}) = -\frac{\pi}{16} \left(1 + \frac{\beta(1 + 6\beta)}{2(2\beta + 1)} \right). \quad (5.4)$$

Maple code for the near-circular approximation is as follows:

```
# FILE nearCircApproxQfnmpl.txt
nearCircApproxQfn:= proc(a,beta)
  local eps,q1;
  eps:= (a^2-1/a^2)/(a^2+1/a^2);
  q1:= -(Pi/16)*(1+beta*(1+6*beta)/(2*(2*beta+1)));
  Pi*(1+4*beta)/8 +q1*eps^2
end proc;
```

Near circular, β small

When $\beta = 0$, this agrees with the asymptotics given in equation (5.1). When β is small but non-zero, asymptotics given in §6.2 check: see equation (6.3).

Near circular, β large

When we first take ϵ small, then let β tend to infinity we have the approximation

$$Q \sim \frac{\pi}{2}\beta + \frac{\pi}{8} - \frac{3\pi}{32}\beta\epsilon^2 - \frac{\pi}{32}\epsilon^2.$$

This is used as a check in §12.1.

Numerical results

Table 5.2 below compares the previous asymptotics (column A) with variational lower bound (column V) and with the Fourier series solution (column F).

$a = 5/4$			
β	A	V	F
1/64	.3825119944	.3807427556	.3807440330
1/16	.4551128646	.4530404184	.4530434724
1/4	.7437766793	.7406928885	.7406944966
1	1.888863782	1.882275712	1.882277797
4	6.450075877	6.429038519	6.429053564
16	24.68100694	24.60087273	24.60089638
64	97.59955268	97.28234644	97.28237296
$a = 17/16$			
β	A	V	F
1/64	.4143605368	.4143493072	.4143493804
1/16	.4879061200	.4878930227	.4878930894
1/4	.7819440804	.7819247200	.7819248140
1	1.957301880	1.957260968	1.957261120
4	6.657144997	6.657014853	6.657015098
16	25.45536249	25.45486728	25.45486742
64	100.6478027	100.6458431	100.6458432
$a = 65/64$			
β	A	V	F
1/64	.4170525380	.4170525625	.4170524896
1/16	.4906779729	.4906782880	.4909655446
1/4	.7851701838	.7851700411	.8350137728
1	1.963086617	1.963085377	1.963085489
4	6.674647535	6.674645198	6.675381808
16	25.52081497	25.52081128	25.51131048
64	100.9054563	100.9054466	100.9054478




Figure 2: §5.2. Plot of $Q_{\text{ratio}} = (Q_{\odot} - Q(e, \beta)) / Q_{\odot}$ at various values of β for small eccentricity.

A simpler, but less accurate, variational approach




Figure 3: §5.2. Plot of small β approximation term Q_1 and a variational lower bound.

The lower bound shown in Figure 3 comes from a trial function which is just the $\beta = 0$ quadratic in Cartesians plus the (best) constant. Of course this is exact for eccentricity zero. Of course we do better with a more general quadratic but the integrals are more ugly.

6 Flows with β small

6.1 General Ω

The asymptotics are described in [37]. The problem is

$$\begin{aligned} -\Delta u &= 1 && \text{in } \Omega, \\ \beta \frac{\partial u}{\partial n} + u &= 0 && \text{on } \partial\Omega. \end{aligned}$$

When β is small, the solution u_β is asymptotically

$$u_\beta \sim u_{\beta=0} + \sum_{j=1}^n \beta^j u_j,$$

with each u_j ($j \geq 1$) harmonic (and independent of β). Our interest is in Q , here denoted Q_β ,

$$Q_\beta = \int_{\Omega} u_\beta,$$

so

$$Q_\beta \sim Q_0 + \beta Q_1 \quad \text{where } Q_1 = \int_{\Omega} u_1,$$

and the integral of u_1 over Ω can be found explicitly (at least up to a single-variable integral) even when u_1 cannot be found explicitly. Now

$$\begin{aligned} -\Delta u_1 &= 0 && \text{in } \Omega, \\ \beta u_1 &= -\beta \frac{\partial u_0}{\partial n} && \text{on } \partial\Omega. \end{aligned}$$

The Divergence Theorem gives

$$\begin{aligned} Q_1 = \int_{\Omega} u_1 &= \int_{\Omega} (u_0 \Delta u_1 - u_1 \Delta u_0) = \int_{\Omega} \operatorname{div} (u_0 \nabla u_1 - u_1 \nabla u_0), \\ &= \int_{\partial\Omega} \left(u_0 \frac{\partial u_1}{\partial n} - u_1 \frac{\partial u_0}{\partial n} \right) = \int_{\partial\Omega} \left(\frac{\partial u_0}{\partial n} \right)^2 ds. \end{aligned}$$

We remark

$$\int_{\partial\Omega} ds = |\partial\Omega|, \quad \int_{\partial\Omega} \left(\frac{\partial u_0}{\partial n} \right) ds = -|\Omega|.$$

This yields, via Cauchy-Schwarz,

$$|\Omega|^2 \leq |\partial\Omega| Q_1,$$

an inequality which becomes more accurate the closer the ellipse is to a circle.

6.2 Ω an ellipse

In general, and in the notation of the preceding subsection,

$$Q_1 = \int_{\partial\Omega} \left(\frac{\partial u_0}{\partial n} \right)^2 ds = \int_{\partial\Omega} |\nabla u_0|^2 ds.$$

For an ellipse $u_0 = u_{0e}$ with u_{0e} given in §2: thus

$$Q_{1,e} = \int_{\partial\Omega} |\nabla u_{0e}|^2 ds.$$

Although the calculation can be done in polar coordinates, perhaps Cartesians lead to simpler integrals:

$$|\nabla u_{0e}|^2 = \frac{a^{-4}x^2 + a^4y^2}{(a^{-2} + a^2)^2}.$$

The Cartesian equation for that part of the ellipse in the first quadrant is

$$Y(x) = \frac{1}{a} \sqrt{1 - \left(\frac{x}{a}\right)^2}, \quad \text{for } 0 < x < a.$$

The perimeter is

$$\begin{aligned} |\partial\Omega| &= 4 \int_0^a \sqrt{1 + Y'(x)^2} dx, \\ &= 4a \text{EllipticE} \left(\frac{\sqrt{a^4 - 1}}{a^2} \right) \quad \text{for } a \geq 1, \\ &= 4a \text{EllipticE}(e) \quad \text{for } a \geq 1, \\ &= 4(1 - e^2)^{-1/4} \text{EllipticE}(e), \end{aligned} \tag{6.1}$$

where e is the eccentricity as defined in equation (3.5).

We now calculate Q_1 , beginning with

$$|\nabla u_{0e}|^2 = \frac{a^2 - a^{-2}(a^2 - a^{-2})x^2}{(a^{-2} + a^2)^2} \quad \text{on } y = Y(x).$$

On integrating this Q_1 is given by

$$Q_1 = \frac{4}{3} \frac{a^3}{(1 + a^4)^2} (2(1 + a^4) \text{EllipticE}(e) - \text{EllipticK}(e)) \quad \text{for } a \geq 1. \tag{6.2}$$

There are many equivalent ways to write this. With $i_2 = i_{xx} + i_{yy}$ we have

$$Q_1 = \frac{L}{a^2 + \frac{1}{a^2}} - \frac{i_2}{(a^2 + \frac{1}{a^2})}.$$

```
# FILE QsmallBetampl.txt
betaSmallQfn:= proc(a,beta)
  local ecc, Q0, Q1;
  ecc:=sqrt(1-1/a^4);
  Q0:= Pi/(4*(a^2 +1/a^2));
  Q1:= (4/3)*(a^3/((1+a^4)^2))*(2*(1+a^4)*EllipticE(ecc)-EllipticK(ecc));
  Q0+beta*Q1
end proc;
```

β small, near circular

The asymptotic expansion for nearly circular ellipse, a near 1, has

$$Q_1 \sim \frac{\pi}{2} - \frac{\pi}{8}(a-1)^2 \quad \text{for } a \rightarrow 1. \quad (6.3)$$

This agrees with the earlier expansion given in equations (5.3,5.4) where, for β small

$$Q \sim \frac{\pi}{8}(1+4\beta) - \frac{\pi}{4}\left(1+\frac{\beta}{2}\right)(a-1)^2 \quad \text{for } a \rightarrow 1, \beta \rightarrow 0.$$

Numerical results

In Table 6.2 below we compare with the numerical values given by the lower-bound variational winner amongst quadratics.

$\beta = 1/4$			
a	A	V	F
65/64	.7851858133	.7851702110	.8350137728
17/16	.7821607512	.7819248090	.7819248140
5/4	.7432200201	.7406928855	.7406944966
2	.4953609194	.4909982744	.4910897996
4	.2147581278	.2139371008	.2142092926
16	.4473378672e-1	.4463047672e-1	.4473097574e-1
$\beta = 1/16$			
a	A	V	F
65/64	.4906792259	.4906780838	.4909655446
17/16	.4879127252	.4878930761	.4878930894
5/4	.4532504638	.4530404192	.4530434724
2	.2624399058	.2620398525	.2621170972
4	.9036181975e-1	.9023065471e-1	.9032360830e-1
16	.1348438275e-1	.1345885965e-1	.1348419675e-1
$\beta = 1/64$			
a	A	V	F
65/64	.4170525792	.4170526698	.4170524896
17/16	.4143507188	.4143493767	.4143493804
5/4	.3807580746	.3807427560	.3807440330
2	.2042096524	.2041611381	.2041881644
4	.5926274272e-1	.5923472697e-1	.5926025316e-1
16	.5672031762e-2	.5665670014e-2	.5671955380e-2

6.3 u_1 for β small: Fourier series?

We begin by recalling (refeq:12b)

$$v_p = -\frac{c^2}{8}(\cosh(2\eta) + \cos(2\psi)) = -\frac{1}{4}(x^2 + y^2).$$

Several items relating to v_p are used, in particular its normal derivative at the boundary, the η -derivative being closely related.

The diagonal matrix M_0 has as its (n, n) entry

$$m_{0,0} = 2\pi, \quad \text{and for } n \geq 1 \quad m_{0,n} = \pi \cosh(2n\eta_0).$$

The first subscript, here 0, reminds us that the entries come from M_0 . The vector F_0 has nonzero entries only at $n = 0$ and $n = 1$ when

$$f_{0,0} = \frac{c^2\pi}{4} \cosh(2\eta_0) \quad \text{and} \quad f_{0,1} = \frac{c^2\pi}{8}.$$

Again, the first subscript, here 0, reminds us that the entries come from f_0 .

The symmetric matrix M_1 is full with entries involving `EllipticK` and `EllipticE`. The vector f_1 , related to the Fourier series of g , similarly involves `EllipticK` and `EllipticE`.

For β positive but small, we seek a representation of the form

$$u \sim u_0 + \beta u_1 \quad \text{for } \beta \rightarrow 0.$$

Our main interest is checking with an asymptotic result obtained using Cartesian coordinates:

$$Q \sim Q_0 + \beta Q_1 \quad \text{for } \beta \rightarrow 0.$$

Our calculation in Cartesian coordinates was completed a year earlier than the elliptic coordinate work here.

$\beta = 0$ re-visited

As one small check on the equations determining the A_n , we note that when $\beta = 0$ the equations are very easy to solve. The matrix $M = M_0$ is diagonal. The only nonzero A_n are those corresponding to $n = 0$ and $n = 1$.

When $\beta = 0$, and subscripting with the first 0 in a pair merely reminding us that the values are for $\beta = 0$, $u_0 = v_p + A_{0,0} + A_{0,1} \cosh(2\eta) \cos(2\psi)$ and the equation $M_0 A = f_0$ gives

$$\begin{aligned} A_{0,0} &= \frac{f_0}{m_{0,0}} = \frac{c^2 \cosh(2\eta_0)}{8} = \frac{\cosh(2\eta_0)}{4 \sinh(2\eta_0)} \\ A_{0,1} &= \frac{f_1}{m_{0,1}} = \frac{c^2}{8 \cosh(2\eta_0)} = \frac{1}{4 \cosh(2\eta_0) \sinh(2\eta_0)}. \end{aligned}$$

As we have already calculated Q_0 from other methods, the calculation here merely serves as a check on $A_{0,0}$ and $A_{0,1}$.

$\beta > 0$ **small**

Consider now approximations when β is small. In §6 we find

$$Q \sim Q_0 + \beta Q_1 \quad \text{as } \beta \rightarrow 0,$$

with Q_0 as in equation (2.4). The approach began with assuming

$$u \sim u_0 + \beta u_1 \quad \text{as } \beta \rightarrow 0,$$

but did not require finding u_1 . If one wished to find u_1 , a Fourier series approach as in this section might be appropriate. Approximate the vector A of Fourier coefficients by

$$A \sim A_0 + \beta A_1,$$

where the vector A_0 is that already calculated in the preceding subsection concerning $\beta = 0$. Then we require, to order β ,

$$(M_0 + \beta M_1)(A_0 + \beta A_1) = (f_0 + \beta f_1),$$

i.e., on approximating to order β ,

$$M_0 A_1 = f_1 - M_1 A_0.$$

Equivalently, using, as noted previously, that M_0 is diagonal,

$$\begin{aligned} m_{0,0} A_{1,0} &= f_{1,0} - m_{1,0,0} A_{0,0} - m_{1,0,1} A_{0,1}, \\ m_{0,1} A_{1,1} &= f_{1,1} - m_{1,1,0} A_{0,0} - m_{1,1,1} A_{0,1} \\ &\vdots = \vdots \\ m_{0,n} A_{1,n} &= f_{1,n} - m_{1,n,0} A_{0,0} - m_{1,n,1} A_{0,1} \end{aligned}$$

Again the first subscript denotes the appropriate order in β .

While the calculations are routine, it may be that the solution A_1 is ugly, and, in any event, the human effort at keeping all the details correct is considerable. We suspect that, at this stage in the study and use of microchannels, the greater accuracy for u is not needed. A calculation which would provide an additional check is to finding $A_{1,0}$ and $A_{1,1}$ which is all that is needed to check against the result for Q_1 given in equation (6.2).

7 β large

7.1 The dominant term

For general Ω

$$Q(\beta) \sim \beta \frac{|\Omega|^2}{|\partial\Omega|} \quad \text{for } \beta \rightarrow \infty.$$

For our ellipses of area π the perimeter is given by equation (6.1), Hence

$$Q(\beta) \sim \beta \frac{\pi^2}{4a \text{EllipticE}(e)} \quad \text{for } \beta \rightarrow \infty.$$

Maple code for this is:

```
betaLargeQfn:= (a,beta) -> beta*Pi^2/(4*a*EllipticE(sqrt(1-1/a^4)));
```

β large, nearly circular ellipses

Further asymptotic approximation of that above for nearly circular ellipses is:

$$Q(\beta) \sim \beta \left(\frac{\pi}{2} - \frac{3\pi}{8}(a-1)^2 + O((a-1)^3) \right) \quad \text{for } \beta \rightarrow \infty, a \rightarrow 1.$$

This may be compared with the asymptotics found in equations (5.3,5.4) further approximated for β large. Then $q_1 \sim -\pi\beta/32$ so

$$Q \sim \frac{\pi\beta}{2} + q_1\epsilon^2 \sim \frac{\pi\beta}{2} - \frac{3\pi\beta}{32}\epsilon^2 \sim \frac{\pi\beta}{2} - \frac{3\pi\beta}{8}(a-1)^2,$$

which agrees with the result of the preceding paragraph.

We have yet to find higher order approximations.

The variational approximation using quadratic test functions agrees with the lowest order term.

The first term $O(\beta)$ is, at fixed β a decreasing function of a on $a > 1$. Note that the quadratic lower bound in column V in the following table is a better approximation than $\beta|\Omega|^2/|\partial\Omega|$ which is less than the lower bound.

Numerical results β large

$\beta = 4$			
a	A	V	F
65/64	6.282052744	6.674647024	6.675381808
17/16	6.265911941	6.657014849	6.657015098
5/4	6.056752387	6.429038511	6.429053564
2	4.602060688	4.869625399	4.870154656
4	2.449872080	2.627235185	2.627747294
16	.6168200091	.6676825774	.6691074764
$\beta = 16$			
a	A	V	F
65/64	25.12821097	25.52081287	25.51131048
17/16	25.06364776	25.45486727	25.45486742
5/4	24.22700955	24.60087275	24.60089638
2	18.40824275	18.69310262	18.69427704
4	9.799488320	10.09964112	10.10542728
16	2.467280036	2.656743920	2.661123514
$\beta = 64$			
a	A	V	F
65/64	100.5128439	100.9054480	100.9054478
17/16	100.2545911	100.6458431	100.6458432
5/4	96.90803820	97.28234641	97.28237296
2	73.63297100	73.92334640	73.92489450
4	39.19795328	39.57699513	39.59760516
16	9.869120146	10.54995058	10.55945235

Further work concerning large β behaviour is given, via Fourier series, in Part II, and, by variational techniques in Part IV.

8 Conclusion and Open Problems

The solution to the Robin boundary problem has been approached from several directions: Fourier series, variational bounds, and asymptotic approximations.

There are several questions that remain.

1. What can be said about asymptotics when the eccentricity tends to 1? The small β and the large β approximations yield some information but more should be possible. (We remark that at large β , small eccentricity will cause the Fourier coefficients V_n of Part II to decay more slowly.) Matched asymptotic approximations may be appropriate.
2. Can u_∞ be found explicitly, or at least the integrals Σ_∞ and Σ_1 ? Given that there are 3-term recurrence relations for the Fourier coefficients of both g and $1/g$, might there be a recurrence relation for the coefficients V_n ? One might combine equation (12.11) with items from Part II § 13. See also Part IV.
3. In the asymptotics for β tending to zero, we found how Q changed, but can one find tidy formulae giving the departure of u from u_0 ?
4. Might it be possible to make better use of the Fourier series for g and $1/g$?

In connection with items 2 and 3 above, we remark that there is a ‘Poisson Integral Formula’ for solving the Dirichlet problem for Laplace’s equation in an ellipse. This follows from the conformal map between ellipse and disk: see [42], p177 and [61]. See also [54, 50, 46, 55].

9 Appendix A: Geometry of ellipses

9.1 General ellipses

9.1.1 Polar coordinates

Consider the ellipse $\frac{x^2}{a^2} + a^2 y^2 \leq 1$. In polar coordinates relative to the centre the boundary curve is

$$\begin{aligned} r &= \frac{1}{\sqrt{\frac{\cos(\theta)^2}{a^2} + a^2 \sin(\theta)^2}} = \sqrt{\frac{2}{a^2 + a^{-2} - (a^2 - a^{-2}) \cos(2\theta)}} \\ &= r\left(\frac{\pi}{4}\right) \left(1 - \frac{a^2 - a^{-2}}{a^2 + a^{-2}} \cos(2\theta)\right)^{-1/2} \quad \text{where } r\left(\frac{\pi}{4}\right) = \sqrt{\frac{2}{a^2 + a^{-2}}} \end{aligned} \quad (9.1)$$

$$= \sqrt{\frac{1 + \tan^2(\theta)}{a^{-2} + a^2 \tan^2(\theta)}}. \quad (9.2)$$

In our computations we take $a \geq 1$.

Without any assumptions on a , the coefficients in the Fourier series for $r(\theta)$ involve elliptic integrals. Our interest in some later sections will be in a near 1. However $(a - 1)$ might not be the best perturbation parameter. We have also used, in computations,

$$\epsilon = \frac{a^2 - a^{-2}}{a^2 + a^{-2}} \sim 2(a - 1) \quad \text{as } a \rightarrow 1,$$

The ellipse is

$$r^2 = \frac{\sqrt{1 - \epsilon^2}}{1 - \epsilon \cos(2\theta)}.$$

We have also used (as have others, e.g. [18]), the eccentricity, as in equation (3.5),

$$e = \sqrt{1 - \frac{1}{a^4}} \sim 2(a - 1)^{1/2} \quad \text{as } a \rightarrow 1. \quad (9.3)$$

As both e^2 and ϵ are relatively similar low-order rational functions of a^2 , one readily finds that

$$\frac{\epsilon}{e^2} = \frac{1}{2}(1 + \epsilon).$$

We remark that the polar equation in terms of the eccentricity is

$$r = \frac{1}{a\sqrt{1 - (e \cos(\theta))^2}} = \frac{g(\theta)}{a}.$$

Including an a dependence in $r(a, \theta)$ we remark that $r(1/a, \theta) = r(a, \theta + \pi/2)$ and $\epsilon(1/a) = -\epsilon(a)$. The binomial expansions

$$\begin{aligned} \frac{r(0)}{r(\pi/4)} &= (1 - \epsilon)^{-1/2} = 1 + \sum_{k=1}^{\infty} \frac{(2k)!}{2^{2k}(k!)^2} \epsilon^k, \\ \frac{r(\theta)}{r(\pi/4)} &= (1 - \epsilon \cos(2\theta))^{-1/2} = 1 + \sum_{k=1}^{\infty} \frac{(2k)!}{2^{2k}(k!)^2} \cos(2\theta)^k \epsilon^k, \end{aligned} \quad (9.4)$$

may be useful in finding higher terms in the perturbation expansions (for a near 1) of some domain functionals. The symmetries of the ellipse explain the form of the expansion in (5.2):

- It is symmetric about $\theta = 0$, hence only cosine terms.
- It is symmetric about $\theta = \pi/2$ and hence only the even order cosine terms $\cos(2m\theta)$.
- When ϵ is replaced by $-\epsilon$ and θ by $\theta + \pi/2$ the expression is unchanged and hence the form of the polynomial coefficients in ϵ forming the Fourier coefficients. For m is odd, only odd powers of ϵ appear: for m is even, only even powers of ϵ appear.

We see these symmetries in connection the solutions of our pde problem. Returning to the study of the boundary curve, we also need the expansion for $r(\pi/4)$:

$$a^2 = \sqrt{\frac{1 + \epsilon}{1 - \epsilon}}, \quad \frac{2}{r(\pi/4)} = a^2 + a^{-2} = \frac{2}{\sqrt{1 - \epsilon^2}}, \quad r\left(\frac{\pi}{4}\right) = (1 - \epsilon^2)^{1/4}.$$

There are a few items of undergraduate calculus that are used. Let s

denote arclength measured around the curve. Then

$$\begin{aligned}
\frac{ds}{d\theta} &= \sqrt{r(\theta)^2 + \left(\frac{dr(\theta)}{d\theta}\right)^2} && \text{in general,} \\
&= 2a\sqrt{\frac{(1+a^8) - (a^8-1)\cos(2\theta)}{((1+a^4) - (a^4-1)\cos(2\theta))^3}} && \text{for an ellipse,} \\
&= 2a\sqrt{\frac{(1+\epsilon)(1+\epsilon^2 - 2\epsilon\cos(2\theta))}{(1-\epsilon\cos(2\theta))^3}} && \text{for an ellipse.}
\end{aligned}$$

We remark that, for the ellipse the normal can be found from the gradient of the ellastic torsion function, $\nabla u_{0e} = \nabla u(\text{ellipse}, \beta = 0)$.

9.1.2 The usual parametric description of an ellipse

The boundary of the ellipse can be described by

$$x = a \cos(\psi), \quad y = \frac{1}{a} \sin(\psi). \quad (9.5)$$

(The relation between the parameter ψ and the polar angle θ is $\tan(\theta) = \tan(\psi)/a^2$.)

$$\begin{aligned}
\left(\frac{ds}{d\psi}\right)^2 &= \left(\frac{dx}{d\psi}\right)^2 + \left(\frac{dy}{d\psi}\right)^2, \\
&= a^2 \sin(\psi)^2 + \frac{1}{a^2} \cos(\psi)^2, \\
&= a^2 (1 - e^2 \cos(\psi)^2)
\end{aligned}$$

From this the perimeter is calculated:

$$\begin{aligned}
|\partial\Omega| &= 4a \int_0^{\pi/2} \sqrt{1 - (e \cos(\psi))^2} d\psi, \\
&= 4a \int_0^{\pi/2} \sqrt{1 - (e \sin(\hat{\psi}))^2} d\hat{\psi}, \\
&= 4a \text{EllipticE}(e).
\end{aligned}$$

The support function, h

There are several equivalent definitions of the support function h which we consider as a function of points on the boundary. Thus at the point associated with the value ψ in definition (9.5)

$$h = \frac{1}{a\sqrt{1 + e^2 \cos^2(\psi)}}.$$

As noted in [41], the curvature is h^3 . Once again, the functions g and \hat{g} make an appearance.

9.1.3 Elliptic coordinates

The elliptic coordinates are related to Cartesians by

$$x = c \cosh(\eta) \cos(\psi), \quad y = c \sinh(\eta) \sin(\psi),$$

and we will set the parameter c by

$$c = \sqrt{a^2 - a^{-2}}.$$

The boundary of the ellipse can be represented, with fixed η_0 ,

$$x = c \cosh(\eta_0) \cos(\psi), \quad y = c \sinh(\eta_0) \sin(\psi).$$

In this

$$a = c \cosh(\eta_0), \quad a^{-1} = c \sinh(\eta_0),$$

so that

$$\tanh(\eta_0) = a^{-2}, \quad (c^2 = a^2 - a^{-2}), \quad \epsilon = \frac{1}{\cosh(2\eta_0)}.$$

Nearly circular ellipses will have η_0 large.

The perimeter is calculated:

$$\begin{aligned} |\partial\Omega| &= 4 \int_0^{\pi/2} \sqrt{\left(\frac{dx}{d\psi}\right)^2 + \left(\frac{dy}{d\psi}\right)^2} d\psi, \\ &= 4c \int_0^{\pi/2} \sqrt{(\cosh(\eta_0))^2 - (\cos(\psi))^2} d\psi, \\ &= 4c \cosh(\eta_0) \text{EllipticE}\left(\frac{1}{\cosh(\eta_0)}\right), \\ &= 4a \text{EllipticE}(e). \end{aligned}$$

We remark that toroidal functions occur elsewhere in this study and that [1] gives a formula, symmetric in a and b ,

$$|\partial\Omega| = 2\pi\sqrt{ab}\text{LegendreP}_{1/2}\left(\frac{a^2 + b^2}{2ab}\right). \quad (9.6)$$

9.2 Nearly circular ellipses, polar coordinates

The first few terms in the expansion of $r(\theta)$ are

$$r(\theta) = 1 + \frac{1}{2} \cos(2\theta)\epsilon + \left(\frac{3}{16} \cos(4\theta) - \frac{1}{16}\right)\epsilon^2 + O(\epsilon^3).$$

Alternatively, we can consider asymptotics as $a \rightarrow 1$. Then $\rho = r_{\text{ellipse}} - 1$ satisfies

$$\begin{aligned} \rho(\theta) &\sim (a-1)\cos(2\theta) - \left(\frac{1}{4} + \frac{1}{2}\cos(2\theta) - \frac{3}{4}\cos(4\theta)\right)(a-1)^2 + O((a-1)^3) \\ &= -\frac{1}{4}(a-1)^2 + \left((a-1) - \frac{1}{2}(a-1)^2\right)\cos(2\theta) + \frac{3}{4}(a-1)^2\cos(4\theta) + O((a-1)^3). \end{aligned} \quad (9.7)$$

For calculations extending the use of higher order terms in the Fourier series (9.7), one might need to replace x^{2n} terms using formulae like

$$x^{2n} = 2^{1-2n} \left(\frac{1}{2} \binom{2n}{n} + \sum_{j=1}^n \binom{2n}{n-j} T_{2j}(x) \right),$$

and, at some future date, we may for elliptical Ω use this but have yet to do so.

9.3 Moments of inertia

9.3.1 General Ω

Area moments

The polar moment of inertia, taking the origin at the centroid, is

$$I_c = \int_{\Omega} ((x - x_c)^2 + (y - y_c)^2),$$

where $z_c = (x_c, y_c)$ is the centroid of Ω . When the boundaries are given in polar coordinates, this is

$$I_c = \frac{1}{4} \int_0^{2\pi} r(\theta)^4 d\theta.$$

Higher order moments arise in connection with calculations based on polynomial test functions in the variational approach of §4.

Boundary moments

See §4.

9.3.2 Ellipse

Area moments

For our disk and ellipse these are

$$I_c(\text{disk}) = \frac{\pi}{2} a^4, \quad I_c(\text{ellipse}) = \frac{\pi}{4} (a^2 + a^{-2}) = \frac{\pi}{2\sqrt{1-\epsilon^2}}. \quad (9.8)$$

The asymptotics below check with the entries in the table of [52] treating domain functionals for nearly circular domains:

$$\begin{aligned} \frac{2I_c(\text{ellipse})}{\pi} &\sim 1 + 2(a-1)^2 \quad \text{as } a \rightarrow 1, \\ \left(\frac{2I_c(\text{ellipse})}{\pi} \right)^{1/4} &\sim 1 + \frac{1}{2}(a-1)^2 + o((a-1)^2) \quad \text{as } a \rightarrow 1, \\ &\sim 1 + a_0 + \frac{3}{4}a_2^2 + o((a-1)^2) \quad \text{as } a \rightarrow 1. \end{aligned}$$

Boundary moments

For doubly-symmetric domains like our ellipse, our notation in §4 is

$$i_{2n} = \int_{\partial\Omega} r^{2n} ds.$$

For the ellipse, these can be evaluated in terms of elliptic integrals.

9.4 Ellipse geometry: miscellaneous

The modulus of asymmetry for an ellipse is calculated in [23]. The calculation involves calculating \hat{g}_0 .

In the case $\beta = 0$ there are improvements to the St Venant inequality in terms of the modulus of asymmetry. (For the St Venant inequality, see [52], and, for ellipses, trivially the $\beta = 0$ case in inequality (1.5).)

10 Appendix B: Elliptic integrals

The functions denoted by `EllipticE` and `EllipticK` in this work are, in this appendix, written as $E(k)$ and $K(k)$. Our notation follows that of many (but not all!) authors, in particular [44]§3.8 and [26]:

$$\begin{aligned} E(k) &= \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2(\theta)} d\theta, \\ K(k) &= \int_0^{\pi/2} \frac{1}{\sqrt{1 - k^2 \sin^2(\theta)}} d\theta. \end{aligned}$$

For $0 \leq z < 1$

$$\begin{aligned} K(k) &= \int_0^1 \frac{1}{\sqrt{(1-t^2)(1-k^2t^2)}} dt, \\ E(k) &= \int_0^1 \sqrt{\frac{1-k^2t^2}{1-t^2}} dt. \end{aligned}$$

The Cauchy-Schwarz inequality yields

$$\frac{\pi^2}{4} \leq K(k) E(k).$$

The functions can be written as hypergeometric functions:

$$\begin{aligned} K(k) &= \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; k^2\right), \\ E(k) &= \frac{\pi}{2} {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; k^2\right). \end{aligned}$$

In showing, in Part II §13 equation (13.16), that two different representations of \hat{g}_n are equivalent, we needed to tell maple that, as in [26] 8.126, with $k' = \sqrt{1 - k^2}$, (and, in a different notation in [2]17.3.29)

$$\begin{aligned} K\left(\frac{1 - k'}{1 + k'}\right) &= \frac{1 + k'}{2} K(k), \\ E\left(\frac{1 - k'}{1 + k'}\right) &= \frac{1}{1 + k'} (E(k) + k' K(k)). \end{aligned}$$

(The first of these is given, in a different notation, on `functions.wolfram.com`.) In our application $k = 1/\sqrt{q} = e$.

Another identity (which we will see in connection with toroidal functions equation (13.53)) is ‘Legendre’s relation’ ([2]17.3.13):

$$K(k)E(k') - K(k)K(k') + K(k')E(k) = \frac{\pi}{2}. \quad (10.1)$$

10.1 Complex k

The EllipticK function has a mirror symmetry:

$$K(\bar{k}) = \bar{K}(k).$$

The real and imaginary parts of $K(\exp(i\phi))$ are given, for example, at the ‘Complex Characteristics’ section of `functions.wolfram.com`. From this one finds

$$|K(\exp(i\phi))|^2 = \frac{1}{4} (K(\cos(\phi))^2 + K(\sin(\phi))^2). \quad (10.2)$$

See equation (13.76).

11 Appendix C: Result Validation and Numerical Examples

11.1 Slip flow in Elliptic Channel with $a = 2$, $b = \frac{1}{2}$

Figure 4 gives, at left, contour plots of u for $\beta = \frac{1}{4}$, $\frac{1}{16}$ and $\frac{1}{64}$, and, at right, those for $\beta = 4$, 16, 64. The contours are very similar to those in Figure

4 of [60]. It is noted that when $\beta > 0$ on the boundary (as well as in the interior) $u > 0$. At any given fixed position z on the boundary, $u(z)$ increases when β value increases.




Figure 4: $a = 2$. Contour plot of axial velocity on the ellipse for six different β values.

11.2 Comparing the results with Ritz Method for Slip Flow

Here we compare with results given in [68].

λ	c	$a = \frac{1}{\sqrt{c}}$	$\beta = \frac{\lambda}{\sqrt{c}}$	A	V	F	Ritz method
0.1	.25	2.000000	.200000	.027080	.026893	.0268994	.0268994
	.5	1.414213	.141421	.131628	.131224	.1312286	.1312286
	.75	1.154700	.115470	.313497	.313320	.3133204	.3133204
0.2	.25	2.000000	.400000	.042611	.041996	.0419990	.0419990
	.5	1.414213	.282843	.184716	.183372	.1833741	.1833741
	.75	1.154700	.230940	.414937	.414338	.4143388	.4143388
0.5	.25	2.000000	1.000000	.071907	.086515	.0865169	.0865167
	.5	1.414214	.707107	.343981	.338284	.3382847	.3382847
	.75	1.154700	.577350	.719257	.716699	.7166986	.7166986
1.0	.25	2.000000	2.000000	.143814	.159568	.1595802	.1595797
	.5	1.414214	1.414214	.509349	.594535	.5945397	.5945397
	.75	1.154701	1.154701	1.004665	1.219750	1.2197505	1.2197505
2.0	.25	2.000000	4.000000	.287629	.304352	.3043838	.3043838
	.5	1.414214	2.828427	1.018698	1.105106	1.1051201	1.1051201
	.75	1.154700	2.309401	2.009330	2.224974	2.2249748	2.2249748
5.0	.25	2.000000	10.000000	.719072	.736625	.7366869	.7366851
	.5	1.414214	7.071068	2.546745	2.634130	2.6341541	2.6341541
	.75	1.154701	5.773503	5.023326	5.239416	5.2394175	5.2394175
10	.25	2.000000	20.000000	1.438144	1.456036	1.4561139	1.4561106
	.5	1.414214	14.142136	5.093491	5.181257	5.1812861	5.1812861
	.75	1.154701	11.547005	10.046652	10.262916	10.2629186	10.2629186

11.3 Blood flow problem

Most of the veins and arteries in our bodies can be taken as having circular cross-section. However in places where the vein or artery has to go through a region which is squeezed in by muscle or bone one might expect the cross-section of the vein or artery to depart somewhat from circular. In this appendix, we take fluid density, ρ , of 1.05 g/mL and fluid viscosity, μ , of 0.04 Poise . See Table 2 in [33]. The geometry of elliptical cross-section of the channel is described by coordinates (x, y) with

$$x = \bar{r}(\psi) \cos(\psi); \quad y = \bar{r}(\psi) \sin(\psi),$$




Figure 5: The cross sections of elliptical shapes with the same length of minor axis $b = 4.9 \mu m$. and various lengths of major axis $a = 6.6, 7.1, 7.6, 8.1$ and $8.5 \mu m$.

where

$$\bar{r}(\psi) = \lambda k(1 + \kappa(k, \psi)), \quad 0 \leq \psi \leq 2\pi$$

and

$$\kappa = -\frac{1}{4}(k-1)^2 + (k-1 - \frac{1}{2}(k-1)^2) \cos(2\psi) + \frac{3}{4}(k-1)^2 \cos(k\psi).$$

The tube ellipticity $\varepsilon = \sqrt{1 - b^2/a^2}$ with the lengths of the half-axes $a = \bar{r}(0) \cos(0)$ and $b = \bar{r}(\pi/2) \sin(\pi/2)$ is determined by setting $\lambda = 0.005$ and $k = 1.15$. Thus, the tube has the ellipticity of 0.6720 with $a = 6.6 \mu m$, $b = 4.9 \mu m$, $\eta_0 = 0.9518$ and $c = 4.4 \mu m$. The boundary of the elliptical cross-section is described by

$$\partial\Omega : (x, y) = (c \cosh \eta_0 \cos \psi, c \sinh \eta_0 \sin \psi).$$

The investigate the influence of elliptical shapes on constant pressure-driven flow of fluid, various sizes of the major axis a are chosen to be vary between $6.6 \mu m$. to $8.6 \mu m$. while the length of minor axis is fixed as $b = 4.9 \mu m$.

To demonstrate the impact of the slip length ℓ on the constant pressure-driven flow of fluid, values of slip length ℓ are chosen to be vary from 0.0 to $50 \mu m$.. The results show that the slip length has a direct influence on the axial velocity. Larger slip length provides higher velocity as shown in Figures 6, 7 and 8.




Figure 6: Axial velocity profiles in x and y directions obtained from the model with no slip lengths: $\ell = 0.0$, and various shapes of cross section.




Figure 7: Axial velocity profiles obtained from the model with $a = 6.6 \mu m$. and $b := 4.9 \mu m$.




Figure 8: Axial velocity profiles in x and y directions obtained from the model having $a = 6.5 \mu\text{m}$. and $b = 4.9 \mu\text{m}$ and various slip lengths $\ell(\mu\text{m}.)$ of 0.0 (solid line), 6.25(dotted line), 25 (dashed line), 37.5 (dash-dotted line) and 50 (long dashed line).

Part II: u_∞ by Fourier series

By G. Keady

12 More about u_∞

In Part I we only noted the dominant approximation to u when β tends to infinity.

12.1 The next term, and u_∞ , for ellipses in general

We seek the term Σ_∞ (independent of β) in the asymptotic expansion

$$Q \sim \frac{\beta |\Omega|^2}{|\partial\Omega|} + \Sigma_\infty \quad \text{for } \beta \rightarrow \infty.$$

The notation here is, as in [37],

$$\Sigma_\infty = \int_{\Omega} u_\infty,$$

with u_∞ satisfying equations (12.5).

The approach here is to seek a Fourier series solution for u_∞ , investigate its contour plot, and after this to integrate u_∞ thereby finding the next term in the expansion for Q . The integration task only requires the constant ($n = 0$) and the next ($n = 1$) Fourier coefficients of u_∞ . The tasks have been completed for nearly circular ellipses, but the general case, specifically determining the $n = 0$ constant coefficient of u_∞ , remains ‘work in progress’.

We recast the elliptic coordinates Fourier series approach so that one uses the Fourier series of $1/g$ rather than g .

Fourier series of $\hat{g} = 1/g$

Write the Fourier series of $1/g$ as

$$\frac{1}{g(s)} = \frac{\hat{g}_0}{2} + \sum_{n=1}^{\infty} \hat{g}_n \cos(2ns). \quad (12.1)$$

As with the earlier calculation of the Fourier series of g , the coefficients \hat{g}_n can be found in terms of `EllipticE` and `EllipticK` functions. In particular,

with the notation of (3.14), we have

$$\begin{aligned} |\partial\Omega| &= 4c \cosh(\eta_0) \text{EllipticE}_0, \\ \hat{g}_0 &= \frac{4}{\pi} \text{EllipticE}_0 = \frac{1}{\pi c \cosh(\eta_0)} |\partial\Omega| \end{aligned} \quad (12.2)$$

$$\hat{g}_1 = \frac{4}{3\pi} (-\cosh(2\eta_0)\text{EllipticE}_0 + (\cosh(2\eta_0) - 1)\text{EllipticK}_0) \quad (12.3)$$

and all the higher \hat{g}_n can also be expressed similarly in terms of EllipticE_0 and EllipticK_0 :

$$\hat{g}_n = \frac{4}{\pi} \left(\hat{E}_n \text{EllipticE}_0 + \hat{K}_n \text{EllipticK}_0 \right), \quad (12.4)$$

where \hat{E}_n and \hat{K}_n are polynomials of degree n in $q = \cosh(\eta_0)^2 = 1/e^2$ with rational number coefficients.

Again there is a three term recurrence relation exists to determine the polynomials \hat{E}_n and \hat{K}_n . In the notation of equation (3.12), the \hat{K}_n sequence of polynomials starts with

$$\hat{K}_0 = 0, \quad \hat{K}_1 = \frac{2}{3}q = \frac{1}{3}(q_2 - 1).$$

The \hat{E}_n sequence of polynomials starts with

$$\hat{E}_0 = 1, \quad \hat{E}_1 = -\frac{1}{3}(2q - 1) = -\frac{1}{3}q_2.$$

See § 13, and the recurrence (13.8).

Calculating u_∞ (at least up to an additive constant)

As in [37], $u \sim \beta|\Omega|/|\partial\Omega| + u_\infty$ where

$$-\Delta u_\infty = 1 \text{ and } \frac{\partial u_\infty}{\partial n} = -\frac{|\Omega|}{|\partial\Omega|}, \quad \int_{\partial\Omega} u_\infty = 0. \quad (12.5)$$

Set

$$u_\infty = v_\infty + v_p,$$

with v_p as in (3.27) before. Once again the harmonic function v_∞ can be represented as a Fourier series

$$v_\infty = \sum_{n=0}^{\infty} V_n \cosh(2n\eta) \cos(2n\psi). \quad (12.6)$$

To determine V_n for $n \geq 1$ we will need

$$\frac{\partial v_\infty}{\partial \eta}(\eta_0, \psi) = \sum_{n=1}^{\infty} 2n V_n \sinh(2n\eta_0) \cos(2n\psi) \quad \text{on } \eta = \eta_0.$$

The Neumann boundary condition is

$$\frac{g(\psi)}{c \cosh(\eta_0)} \frac{\partial u_\infty}{\partial \eta}(\eta_0, \psi) = \frac{1}{c \sqrt{\cosh^2 \eta_0 - \cos^2 \psi}} \frac{\partial u_\infty}{\partial \eta}(\eta_0, \psi) = -\frac{|\Omega|}{|\partial \Omega|}.$$

In terms of the harmonic function v_∞ this is

$$\frac{\partial v_\infty}{\partial \eta} = -\frac{\partial v_p}{\partial \eta} - \frac{|\Omega|}{|\partial \Omega|} \frac{c \cosh(\eta_0)}{g(\psi)} \quad \text{on } \eta = \eta_0. \quad (12.7)$$

Now one finds

$$\frac{\partial v_p}{\partial \eta} = -\frac{1}{2} \quad \text{on } \eta = \eta_0. \quad (12.8)$$

(In Cartesian coordinates we have $\nabla v_p \cdot \nabla u_0$ is constant around $\partial \Omega$, but we seek u_∞ such that $\nabla u_\infty \cdot \nabla u_0$ is a constant multiple of $|\nabla u_0|$ there.) The integral around the boundary of the normal derivative of a harmonic function must be 0, and this accords with the value of \hat{g}_0 , the constant term in the Fourier series of $1/g$ to ensure this as combining (12.8) and

$$\frac{|\Omega|}{|\partial \Omega|} c \cosh(\eta_0) \hat{g}_0 = \frac{\pi}{|\partial \Omega|} \frac{1}{2\pi} |\partial \Omega| = \frac{1}{2}$$

in (12.7) we have the required result.

From the Fourier series of $1/g(\psi)$ we can find all the Fourier coefficients (12.6), except the constant term, in the series for v_∞ . In particular the term V_1 is found from (12.7) (on using (12.8), contributing 0)

$$2V_1 \sinh(2\eta_0) = 0 - \frac{|\Omega|}{|\partial \Omega|} c \cosh(\eta_0) \hat{g}_1 = -\frac{\pi \hat{g}_1}{4\text{EllipticE}_0}.$$

Hence

$$V_1 = -\frac{\pi \hat{g}_1}{8\text{EllipticE}_0 \sinh(2\eta_0)}. \quad (12.9)$$

Similarly

$$V_2 = -\frac{\pi \hat{g}_2}{16 \text{EllipticE}_0 \sinh(4\eta_0)}, \quad (12.10)$$

and, more generally, for $n > 0$,

$$V_n = -\frac{\pi \hat{g}_n}{8n \operatorname{EllipticE}_0 \sinh(2n\eta_0)}. \quad (12.11)$$

Contour plots for u_∞

As we have all the Fourier coefficients V_n except V_0 we can produce contour plots. We show one of these, that for $a = 2$.

The contourplot (at $a = 2$) is visually indistinguishable from that produced from the quadratic found using the $c_0, c_X X, c_Y Y$ values found in §4.3. Except when η_0 is small, the V_n will decrease rapidly with n , and the plots are essentially those from truncating the Fourier series so that it only includes the V_1 term. The numerical solutions given in Figure 4 of Part I Appendix C are, for $\beta \geq 4$, very much like that of Figure 9.




Figure 9: $a = 2$. Contours of u_∞

Finding V_0 and Q ?

The calculation of V_0 remains. We will outline how it can be found as an infinite series, but shall only complete the details for nearly circular ellipses.

The condition that the boundary integral of u_∞ is zero remains to be used. Now, with, as usual, s for arc length around the boundary, $\eta = \eta_0$,

$$\left(\frac{ds}{d\psi}\right)^2 = \left(\frac{dx}{d\psi}\right)^2 + \left(\frac{dy}{d\psi}\right)^2 = c^2 (\cosh(\eta_0)^2 - \cos(\psi)^2) = J(\eta_0, \psi).$$

The square root of the Jacobian is related to the function g :

$$\sqrt{J(\eta_0, \psi)} = \frac{c \cosh(\eta_0)}{g(\psi)} = \frac{a}{g(\psi)}.$$

We need to satisfy

$$0 = \int_{\partial\Omega} u_\infty = 4 \int_0^{\pi/2} (v_p + v_\infty) \sqrt{J(\eta_0, \psi)} d\psi.$$

The integral $I_{p,\partial}$ around the boundary of v_p is $-i_2/4$ where i_2 is the boundary moment of inertia and can also be evaluated directly,

$$\begin{aligned} I_{p,\partial} &= 4 \int_0^{\pi/2} v_p \sqrt{J(\eta_0, \psi)} d\psi = 4a \int_0^{\pi/2} v_p \hat{g}(\psi) d\psi, \\ &= -\frac{a^3}{3} ((2 - e^2)\text{EllipticE}_0 + (1 - e^2)\text{EllipticK}_0). \end{aligned}$$

When evaluating the boundary integral of v_∞ , the square root brings in all the V_n .

- It may be that there are useful Mean Value results applying to ellipses, and harmonic functions in them, of

$$\int_{\partial\Omega} v_\infty - v_\infty(0) |\partial\Omega| ?$$

It is, of course, zero for a circle. In general it wouldn't be zero, as, for example,

$$\int_{\partial\Omega} (x^2 - y^2) = i_X - i_Y \neq 0 \quad \text{when } a > 1.$$

Some boundary integrals can be related to integrals along the major axis of the ellipse, and [54] is given as a reference in papers by Symeonidis. For any function h harmonic in the ellipse, with c as before $c = \sqrt{a^2 - 1/a^2}$,

$$\int_{-c}^c \frac{h(x, 0)}{\sqrt{c^2 - x^2}} dx = \frac{1}{2} \int_{-\pi}^{\pi} h(a \cos(\psi), \frac{\sin(\psi)}{a}) d\psi.$$

(The right-hand side is not the integral of h with respect to arc length.)

- The integral

$$I_{\infty, \partial} = 4 \int_0^{\pi/2} v_{\infty} \sqrt{J(\eta_0, \psi)} d\psi = 4a \int_0^{\pi/2} v_{\infty} \hat{g}(\psi) d\psi,$$

can be evaluated, with V_0 left as an unknown using Parseval's Theorem. We could find V_0 if we could evaluate

$$\sum_{n=1}^{\infty} \frac{\hat{g}_n^2}{n \tanh(2n\eta_0)},$$

which can be recast as a series involving Legendre functions $Q_{n-1/2}^{-1}$: see §13. At each fixed $a > 1$ (or equivalently η_0 or eccentricity e) this would be a simple float numerical task. However, we have not yet found a simple formula for V_0 , and hence Σ_{∞} , as simple as that we have for Q_1 , in equation (6.2).

Finally to determine Σ_{∞} using $u_{\infty} = v_p + v_{\infty}$ and recalling that we will only need, as in (3.34), the $n = 0$ and $n = 1$ Fourier coefficients of v_{∞} :

$$\Sigma_{\infty} = -\frac{\pi}{8} \frac{\cosh(2\eta_0)}{\sinh(2\eta_0)} + \pi V_0 - \frac{1}{2} \pi V_1. \quad (12.12)$$

Combining this with the dominant term we have

$$Q \sim \frac{\beta |\Omega|^2}{|\partial\Omega|} + \Sigma_{\infty} \quad \text{as } \beta \rightarrow \infty.$$

We remark that for a large Σ_1 becomes large. The asymptotics for $Q(a, \beta)$ for β large should only include the Σ_1 term when Σ_1/β is smaller than Σ_{∞} . In practice this means that, for a large, the large β asymptotics are good only for β very large. (When the domain is a rectangle, rather than an ellipse, there is the same behaviour and this is very easy to demonstrate as u_{∞} is, in Cartesian coordinates, a simple quadratic.)

β large for nearly circular ellipses

Recall that the eccentricity $e = 1/\cosh(\eta_0)$. From equation (3.18)

$$\hat{g}_0 = 1 - \frac{e^2}{4}, \quad \hat{g}_1 = -\frac{e^2}{4}.$$

Also

$$\text{EllipticE}_0 \sim \frac{\pi}{2} - \frac{\pi}{8} e^2 - \frac{3\pi}{128} e^4, \quad \text{EllipticK}_0 \sim \frac{\pi}{2} + \frac{\pi}{8} e^2 + \frac{9\pi}{128} e^4,$$

and

$$a = \left(\frac{1}{1-e^2} \right)^{1/4} \sim 1 + \frac{1}{4} e^2 + \frac{5}{32} e^4, \quad c = a e,$$

and

$$|\partial\Omega| = 4a \text{EllipticE}_0 \sim 2\pi + \frac{3\pi}{32} e^4.$$

(The terms in e^4 in the series for EllipticE_0 and a are only there to check against the series for $|\partial\Omega|$.) We also note

$$\sinh(2\eta_0) = 2\frac{1}{e} \sqrt{\frac{1}{e^2} - 1} \sim \frac{2}{e^2} - 1 - \frac{1}{4} e^2.$$

The Fourier coefficients of v_∞ are needed. From equation (12.9)

$$V_1 = -\frac{\pi \hat{g}_1}{8 \text{EllipticE}_0 \sinh(2\eta_0)} \sim \frac{1}{32} e^4.$$

Comparison with earlier near-circular results

Because, for nearly circular ellipses, the asymptotics for all the V_n , $n \geq 1$ are uncomplicated, and the terms get rapidly smaller in n , we can find V_0 .

Maple code gives, for $\beta \rightarrow \infty$ and $e \rightarrow 0$,

$$Q \sim \left(\frac{1}{2} \pi - \frac{3}{128} \pi e^4 \right) \beta + \frac{1}{8} \pi - \frac{1}{128} \pi e^4 - \frac{1}{256} \frac{\pi e^4}{\beta}.$$

Another quick look at the ϵ small solution of §5.2, in η, ψ coordinates, produces the same result.

12.2 u_∞ for other domains

It seems rarely to happen that both u_0 and u_∞ have simple explicit formulae representing them. An exception is the equilateral triangle: see [37, 48]. And, of course, the circular disk has $u_0 = u_\infty$ and is the only domain for which there is equality. (There may well be other less trivial domains, for example, the semicircle.)

For our ellipse, u_0 is simply, in Cartesians, a quadratic polynomial, or equivalently has a Fourier series (in elliptic coordinates) with just 2 terms, while u_∞ has a Fourier series with all terms nonzero.

The situation with a rectangle is as follows. There is an elaborate Fourier series in Cartesian coordinates for u_0 . However u_∞ is simply a quadratic in the Cartesian coordinates. Indeed our function u_0 , given in equation (2.3), is u_∞ for the rectangle $(-a^2, a^2) \times (-b^2/b^2)$. The methods in §4.2 would give good lower bounds for $Q(\text{rectangle})$ when β is large. See Part III § 15.2. (For large β asymptotics for a rectangle, see [40].)

13 A difference equation – and toroidal functions

The Fourier coefficients g_n and \hat{g}_n are central to our separation of variables solution of Problems $(P(\beta))$ and $(P(\infty))$ respectively. On using maple to find these, maple returned (instances of) them as the sum of two terms: the first term is the product of a polynomial with an EllipticE function, and the second term is the product of a polynomial with an EllipticK function. The dependence on n is only in the polynomials. See equations (3.17), (12.4), repeated at equations (13.9), (13.10). The polynomials satisfy the same recurrence relations as g_n and \hat{g}_n : see $(RE(\alpha))$ below where $\alpha = 1/2$ for g_n and $\alpha = 3/2$ for \hat{g}_n .

It happens that g_n and \hat{g}_n can also be expressed in terms of LegendreQ functions – specifically toroidal functions. Again sequences of polynomials satisfying $(RE(1/2))$ and $(RE(3/2))$ arise: see equations (13.61), (13.62). More generally, for any α , these polynomial sequences are present in the entries on Legendre functions at functions.wolfram.com. There is also some literature concerning the polynomials in families of recurrence relations which include our case $\alpha = 1/2$: see [6, 27] and subsection 13.9 below. It isn't obvious what, for general α , the polynomials should be called, though for

$\alpha = 0$ and $\alpha = 1$ they are Chebyshev polynomials. In the case $\alpha = 1/2$, the sequence of polynomials (particularly those labelled p_{01}) might be called ‘associated toroidal Legendre polynomials’, and we now refer to all the sequences of polynomials $p_{01}(n)$, $p_{10}(n)$ of equation (13.61) and $\hat{p}_{01}(n)$, $\hat{p}_{10}(n)$ of equation (13.62) (or linear combinations thereof, e.g. $p_{10} + x p_{01}$ and, more generally, α half an odd integer) by this name. Though others have used the adjective ‘associated’ in this context, we remark that the adjective is used in several different ways. While the previously cited [6, 27] on ‘associated Legendre functions’ includes our case of $\alpha = 1/2$ it doesn’t include $\alpha = 3/2$. ‘Associated Gegenbauer polynomials’ as in [10]§3 or, more generally, the family of Pollaczek polynomials introduced in [51] (see [22]p220) cover general α but have more parameters than our simpler cases. (See also our §13.9.)

Questions which arise include the following.

- Are there simple formulae for these sequences of polynomials? E.g. (i) Can each sequence be expressed as a single hypergeometric function (perhaps a ${}_3F_2$)? (ii) What are the formulae for coefficients?
- Finally, how might the information on the polynomials be best applied to progress the tasks associated with the pde questions which we have listed below in a subsection concerning ‘Hopes’.

A related problem, also involving sequences of polynomials satisfying (RE(1/2)) or (RE(3/2)) is to discover formulae, for general n , agreeing with maple’s output for the g_n and \hat{g}_n involving EllipticE and EllipticK functions. This also seems to be an old problem: finding formulae for the toroidal functions $Q_{n-1/2}$ involving EllipticE and EllipticK functions. We will do relatively little on this last problem.

13.1 Notation

We denote the reciprocal of g by \hat{g} :

$$\hat{g}(\psi) = \frac{1}{g(\psi)} = \sqrt{1 - \frac{\cos^2(\psi)}{q}}.$$

As described earlier, the Fourier cosine coefficients of g are denoted g_n (see equation (3.13)), and those of \hat{g} by \hat{g}_n :

$$g_n = \frac{2}{\pi} \int_0^\pi g(\psi) \cos(2n\psi) d\psi, \quad (13.1)$$

$$\hat{g}_n = \frac{2}{\pi} \int_0^\pi \hat{g}(\psi) \cos(2n\psi) d\psi. \quad (13.2)$$

Both g_n and \hat{g}_n can be written concisely in terms of toroidal functions, Legendre functions: see equations (13.17), (13.18). The sequences g_n and \hat{g}_n are used, so far, just in numerics. In both sequences our interest is in $q > 1$ but, the integral for \hat{g}_n evaluates at other values of q , for example at $q = 1$ to

$$\hat{g}_n(q = 1) = \frac{2}{\pi} \int_0^\pi \sin(\psi) \cos(2n\psi) d\psi = \frac{1}{\pi} \frac{1}{n^2 - 1/4}. \quad (13.3)$$

Though the integrals are complex, one finds

$$\frac{g_{n+1}(q = 1/2)}{g_{n-1}(q = 1/2)} = -\frac{n - 1/2}{n + 1/2},$$

and a similar formula for the \hat{g}_n sequence.

13.2 Hopes for application to the pde problems

We have various hopes for future results.

- One hope is that further results on \hat{g}_n may ultimately lead to further analytical results concerning u_∞ and, in particular, its values on the boundary, and thence to analytical expressions for Σ_∞ and Σ_1 .
- A more modest goal, also as yet not implemented, is that it may be possible to improve the efficiency of the numerical calculations of g_n and of \hat{g}_n as the present codes call on maple to evaluate each of the integrals separately for each n and uses maple's results given in terms of EllipticE and EllipticK functions.
- There also remains the prospect, should there be any demand for codes in languages used by engineers, e.g. matlab, to provide such codes. (We remark that as at 2019a matlab's legendre function does not compute

toroidal functions. We have yet to check the matlab File Exchange site.) We remark that there are numerical stability issues with recurrence relations when performed purely within fixed precision float computation but these can be circumvented using the Symbolic Toolbox. It isn't essential for the code to be entirely within matlab as it would also be possible to use existing Fortran or C codes (such as described in [25]) mex-ed into matlab. (Matlab codes and related algorithms for toroidal functions are mentioned by several authors, Majic is one, Gil and Segura others, e.g [25]. There is a library, POLPAK, implementing algorithms in various languages, including matlab.)

Noting the present limitations of matlab's legendre function, our present suggestion is to compute $Q_{-1/2}$, $Q_{1/2}$ and $Q_{-1/2}^{-1}$, $Q_{1/2}^{-1}$ from their expressions in terms of EllipticE and EllipticK functions, and then the toroidal Q functions using the polynomials p_{01} , p_{10} and \hat{p}_{01} , \hat{p}_{10} defined below. (Matlab's Symbolic Toolbox might have a role in connection with the polynomials.)

13.3 Introducing the difference equation (RE(α))

In the body of this paper we noted, at equations (3.17) and (12.4), that maple, when asked to perform the integrals defining g_n and \hat{g}_n returned a result in which the terms multiplying elliptic integrals are polynomials in q or rather $q_2 = 2q - 1$. We denoted the polynomials by $E_n(q_2)$, $K_n(q_2)$ (these satisfying the recurrence (RE(1/2)) defined below) and by $\hat{E}_n(q_2)$, $\hat{K}_n(q_2)$ (satisfying the recurrence (RE(3/2))).

While we have yet to make any significant use of the items here for our Robin boundary condition problems, we have discovered that the Fourier series of g and of \hat{g} , and related studies of toroidal functions, have been studied in connection with various different problems since at least the 1880s. In spite of the bewilderingly large literature both on Legendre functions and on orthogonal polynomials, we have not seen a systematic account of the few results we present concerning sequences of polynomials satisfying the simple recurrences satisfied by each of g_n and of \hat{g}_n .

Our sequences of Fourier coefficients satisfy difference equations of the form

$$(n + \alpha)u_{n+1} = 2q_2nu_n - (n - \alpha)u_{n-1}, \quad (\text{RE}(\alpha))$$

where $q_2 = 2q - 1$, and where conformity to conventions in the orthogonal

polynomial literature seems appropriate, we write x rather than q_2 . The g_n satisfy this with $\alpha = 1/2$ while the \hat{g}_n satisfy it with $\alpha = 3/2$.

The recurrence relation (RE(α)) satisfied by Legendre functions is given in standard references such as [2] 8.5.3 and [21] §3.8 (2). The order μ of the Legendre functions, $Q_\nu^\mu(q_2)$ is determined by $\alpha = \frac{1}{2} - \mu$: the degrees in the sequence are of the form $\nu = n - 1/2$.

Our interest is in $\alpha = 1/2$ (for which we give later polynomial sequences $p_{01}(n)$ and $p_{10}(n)$) and $\alpha = 3/2$ (for which we give later polynomial sequences $\hat{p}_{01}(n)$ and $\hat{p}_{10}(n)$) but results at other α might be useful in contexts other than the pde problem of this paper.

13.4 An aside: (RE(0)) and (RE(1)): Chebyshev

At $\alpha = 0$ the n cancels and the recurrence is constant coefficient, so solvable by $u_n = r^n$ for appropriate r . The general solution is a linear combination of the Chebyshev polynomial solutions

$$u_n(\alpha = 0) = c_T(q_2)T_n(q_2) + c_U(q_2)U_n(q_2),$$

where c_T and c_U are independent of n . As n no longer occurs in the coefficients in the recurrence (RE(0)), equally good as a polynomial solution is $u_n = T_{n-m}$ for integer m .

Chebyshev polynomials also arise in solving the recurrence when $\alpha = 1$: solutions are

$$u_n(\alpha = 1) = \frac{u_n(\alpha = 0)}{n}.$$

As remarked before, solutions can be expressed in terms of Legendre functions. For the $\alpha = 0$, $\mu = 1/2$ case, see [21]3.6.1(12). Also we have

$$\begin{aligned} P_{n-1/2}^{1/2}(\cos(\theta)) &= \sqrt{\frac{2}{\pi \sin(\theta)}} \cos(n\theta) = \sqrt{\frac{2}{\pi \sin(\theta)}} T_n(\cos(\theta)), \\ Q_{n-1/2}^{1/2}(\cos(\theta)) &= -\sqrt{\frac{\pi}{2 \sin(\theta)}} \sin(n\theta), \\ P_{n-1/2}^{1/2}(\cosh(\eta)) &= \sqrt{\frac{2}{\pi \sinh(\eta)}} \cosh(n\eta) = \sqrt{\frac{2}{\pi \sinh(\eta)}} T_n(\cosh(\eta)), \\ Q_{n-1/2}^{1/2}(\cosh(\eta)) &= -\sqrt{\frac{\pi}{2 \sinh(\eta)}} \exp(-n\eta). \end{aligned}$$

We remark that

$$\begin{aligned} U_n(x) &= \frac{x T_{n+1}(x) - T_{n+2}(x)}{1 - x^2}, \\ U_n(\cosh(\eta)) &= \frac{\sinh((n+1)\eta)}{\sinh(\eta)}, \\ \exp(-n\eta) &= T_n(\cosh(\eta)) - \sinh(\eta)U_{n-1}(\cosh(\eta)), \end{aligned}$$

and $u_n = \exp(-n \cosh^{-1}(q_2))$ solves $(\text{RE}(0))$. Similar formulae are available for the case $\alpha = 1$, order $\mu = -1/2$.

Maple's `rsolve` solves $(\text{RE}(m))$ for other integer m .

None of the items in this elementary subsection are relevant to the $\alpha = 1/2$ and $\alpha = 3/2$ cases which arise in our partial differential equation problem. However, relations between families of polynomials and various connections between Legendre functions will arise in the context of the $\alpha = 1/2$ and $\alpha = 3/2$ cases and the easier situation sometimes suggests possible results.

13.5 Deriving $(\text{RE}(\alpha))$ from integrals defining Fourier coefficients

While the expressions for g_n and \hat{g}_n in terms of toroidal functions have value, much can be derived, simply from their definitions, including the recurrences $(\text{RE}(\alpha))$ and inter-relationships. The simple relationships between g and \hat{g} lead to relationships between their Fourier coefficients.

- Since

$$\hat{g}(\psi) = \left(1 - \frac{\cos^2(\psi)}{q}\right) g(\psi) = \frac{1}{2q} ((2q-1) - \cos(2\psi)) g(\psi),$$

we have by multiplying both sides by $\cos(2n\psi)$, using cosine formulae, and integrating

$$4q\hat{g}_n = 2(2q-1)g_n - g_{n+1} - g_{n-1}. \quad (13.4)$$

- Since

$$2q \frac{d\hat{g}(\psi)}{dq} + \hat{g}(\psi) - g(\psi) = 0,$$

the Fourier coefficients g_n can be found from those of \hat{g} :

$$g_n = \hat{g}_n + 2q \frac{d\hat{g}_n}{dq} = 2\sqrt{q} \frac{d}{dq}(\sqrt{q}\hat{g}_n). \quad (13.5)$$

• Since

$$\begin{aligned} \frac{d\hat{g}(\psi)}{dq} &= \frac{\cos(\psi)^2 g}{2q^2} = \frac{(\cos(2\psi) + 1) g}{4q^2}, \\ \frac{d\hat{g}_n}{dq} &= \frac{1}{8q^2}(2g_n + g_{n-1} + g_{n+1}). \end{aligned} \quad (13.6)$$

(Equations (13.5) and (13.6) check against (13.4).)

Define also

$$q_2 = 2q - 1.$$

There are three-term recurrence relations between the coefficients:

$$(n - \frac{1}{2})g_n = 2q_2(n - 1)g_{n-1} - (n - \frac{3}{2})g_{n-2}, \quad (13.7)$$

$$(n + \frac{1}{2})\hat{g}_n = 2q_2(n - 1)\hat{g}_{n-1} - (n - \frac{5}{2})\hat{g}_{n-2}. \quad (13.8)$$

The derivation is treated below, where we derive the following (RE(1/2)) equivalent to (13.7):

$$(n + \frac{1}{2})g_{n+1} = 2q_2 n g_n - (n - \frac{1}{2})g_{n-1}. \quad (\text{RE}(1/2))$$

Corresponding to (13.8) we have (RE(3/2))

$$(n + \frac{3}{2})\hat{g}_{n+1} = 2q_2 n \hat{g}_n - (n - \frac{3}{2})\hat{g}_{n-1}. \quad (\text{RE}(3/2))$$

We now repeat equations given in the main part of the paper. In these the argument of the elliptic integrals is $1/\sqrt{q}$ and we wrote,

$$g_n = \frac{4}{\pi} (E_n \text{EllipticE}_0 + K_n \text{EllipticK}_0), \quad (13.9)$$

$$\hat{g}_n = \frac{4}{\pi} (\hat{E}_n \text{EllipticE}_0 + \hat{K}_n \text{EllipticK}_0), \quad (13.10)$$

with the E_n , K_n , \hat{E}_n , \hat{K}_n polynomials in q_2 of degree at most n . The polynomials E_n , K_n satisfy the recurrence (13.7), (RE(1/2)): the polynomials

\hat{E}_n , \hat{K}_n satisfy the recurrence (13.8), (RE(3/2)). The starting values for the iterations to determine the polynomials are given, for the g_n iteration by (3.15), (3.16) and for the \hat{g}_n iteration by (12.2), (12.3). Maple's `rsolve` will solve (RE(α)) in the special cases when $q_2 = 0$ and $q_2 = 1$. When $q_2 = 1$ there are constant solutions ($g_n = 1$ for all n), and reduction of order gives the other.

The method to establish (13.7), (13.8) involves a further relation between the coefficients g_n , \hat{g}_n of the two series, and is suggested in a posting by Jack D'Aurizio on math.stackexchange.com/questions/930003/fourier-series-of-sqrt1-k2-sin2t. We have

$$g_{m+1} = (2q - 1)g_m + (4m - 2)q\hat{g}_m, \quad (13.11)$$

and also, exactly as in D'Aurizio's post,

$$8mq\hat{g}_m = g_{m+1} - g_{m-1}. \quad (13.12)$$

Eliminating \hat{g}_m between equations (13.11), (13.12) we find the recurrence (RE(1/2)) with n there replaced by m . Also, equations (13.11) and (13.12) together yield

$$g_{m-1} = (2q - 1)g_m - (4m + 2)q\hat{g}_m, \quad (13.13)$$

It may be that, for $q_2 > 1$, all the g_m are positive, and the sequence is decreasing.

We have $\hat{g}_0 > 0$ and it may be that, for $q_2 > 1$ and $m \geq 1$ that the terms \hat{g}_m are all negative and form an increasing sequence.

13.5.1 Generating functions?

Correcting a misprint in [11] p202, equation (12.5), (which reports [6]) for $\alpha = 1/2$, a generating function is

$$G(x, w) = \frac{1}{2} \frac{1}{\sqrt{w(1 - 2xw + w^2)}} \int_0^w \frac{1}{\sqrt{t(1 - 2xt + t^2)}} dt, \quad (13.14)$$

$$G(x, w) = \sum_{n=0}^{\infty} p_{01}(n+1, x)w^n. \quad (13.15)$$

The $p_{01}(n)$ are polynomials of degree $n-1$ satisfying (RE(1/2)) starting from $p_{01}(0) = 0$, $p_{01}(1) = 1$. These are studied extensively in later parts of this section. (Equation (13.74) follows from this: see [11] equation (12.6).)

This subsection is ‘incomplete’, with everything after this paragraph merely an early personal attempt predating finding the result reported in equation (13.14) above. It may be that, for general α , elaborate special functions, e.g. Appell hypergeometrics, may be needed as they were in the more general situation in [10]. At least for our special case of $\alpha = 1/2$ incomplete elliptic integrals suffice: see also equations (1.10), (1.11) of [62].

The stackexchange posting suggests there might be reasonably neat expressions for generating functions,

$$G(X) = g_0 + g_1 X + \sum_{n=2}^{\infty} g_n X^n, \quad \hat{G}(X) = \hat{g}_0 + \hat{g}_1 X + \sum_{n=2}^{\infty} \hat{g}_n X^n,$$

and that these functions satisfy first order linear differential equations. The differential equation satisfied by G is

$$\mathcal{L}G = (X - 2q_2 X^2 + X^3) \frac{dG}{dX} - \frac{1}{2}(1 - X^2)G = \frac{1}{2}(-g_0 + g_1 X).$$

$X = 0$ is a singular point. The homogeneous de $\mathcal{L}G_h = 0$ is solved by a function G_h with $G_h(0) = 0$,

$$G_h(X) = \sqrt{\frac{X}{1 - 2q_2 X + X^2}}.$$

We would be content with representations of G valid for $q_2 > 1$ and $0 \leq X < 1/(2q_2)$. Applying the ‘variation of parameters’ formula to the nonhomogeneous de above, using G_h , leads to a messy particular solution in terms of incomplete elliptic integrals. (Though possibly unrelated, we will see elliptic integrals in other contexts: see §13.9.)

The corresponding de for \hat{G} is:

$$\hat{\mathcal{L}}\hat{G} = (X - 2q_2 X^2 + X^3) \frac{d\hat{G}}{dX} + \frac{1}{2}(1 - X^2)\hat{G} = \frac{1}{2}(\hat{g}_0 + 3\hat{g}_1 X).$$

$X = 0$ is a singular point. The homogeneous de $\hat{\mathcal{L}}\hat{G}_h = 0$ is solved by a function \hat{G}_h with $\hat{G}_h(0) = 0$,

$$\hat{G}_h(X) = \sqrt{\frac{1 - 2q_2 X + X^2}{X}},$$

(which is the reciprocal of G_n). The relations (13.4,13.5) lead to relations between G and \hat{G} , for example

$$G = 2\sqrt{q} \frac{d}{dq} (\sqrt{q} \hat{G}).$$

One can derive a differential equation for $\hat{G}(q)$ involving derivatives with respect to q . It may be that if one starts from a closed form for $\hat{G}(1)$, i.e. at $q = 1$, solving the initial value problem might be useful.

13.5.2 Hypergeometric representations

Also presented on stackexchange is a hypergeometric formula for \hat{g}_n . See also [15] at Lemma 1 which gives a slightly different representation. Define

$$\lambda = \frac{1}{\sqrt{q} + \sqrt{q-1}} = \sqrt{q} - \sqrt{q-1},$$

from which

$$\lambda^2 = \frac{1}{q_2 + \sqrt{q_2^2 - 1}}.$$

We have

$$\hat{g}_n = -\frac{2}{\lambda\sqrt{q}} \frac{(2n)!}{(2n-1) 2^{2n+1} (n!)^2} (\lambda^2)^n {}_2F_1\left(-\frac{1}{2}, n - \frac{1}{2}; n + 1; \lambda^4\right). \quad (13.16)$$

(Quick check, $\hat{g}_0 > 0$ and, for $n > 0$, $\hat{g}_n < 0$, though we only have numeric evidence for this.)

13.5.3 Negative n

We remark that from the $\cos(2n\psi)$ in the defining integrals we have

$$g_{-n} = g_n, \quad \hat{g}_{-n} = \hat{g}_n.$$

See also equations (13.25), (13.52). It is also more generally true that if u_n is a solution of $(\text{RE}(\alpha))$ and $\alpha \neq 0$ then $u_{-n} = u_n$.

13.6 Toroidal functions, LegendreQ functions

13.6.1 Main facts concerning Q functions

We defer to a later subsection items on LegendreP functions. LegendreP functions are equally appropriate as solutions of (RE(1/2)) and (RE(3/2)) but our initial concern is with the Fourier coefficients g_n and \hat{g}_n and these are LegendreQ functions. Also in this subsection we are interested in the functions defined on the interval $(1, \infty)$ though in some later sections we are also concerned with the interval $(-1, 1)$. (Branch points will then need to be considered even though our application involves real-valued functions of a real argument.)

Toroidal functions are special cases of Legendre functions, specifically $Q_{n-1/2}^\mu$ and $P_{n-1/2}^\mu$ with n integer, and, sufficient for our purposes also μ integer. We have, as in [21]§3.10(3) and other references such as [14]

$$g_n = \frac{4\sqrt{q}}{\pi} Q_{n-1/2}(q_2), \quad (13.17)$$

$$\hat{g}_n = \frac{2\sqrt{q-1}}{\pi} Q_{n-1/2}^{-1}(q_2). \quad (13.18)$$

(See equations (43) and (44) of [14].)

Toroidal functions can be related to Elliptic Integrals. In particular

$$Q_{-1/2}(q_2) = \frac{1}{\sqrt{q}} \text{EllipticK}\left(\frac{1}{\sqrt{q}}\right), \quad (13.19)$$

$$Q_{1/2}(q_2) = q_2 \frac{1}{\sqrt{q}} \text{EllipticK}\left(\frac{1}{\sqrt{q}}\right) - 2\sqrt{q} \text{EllipticE}\left(\frac{1}{\sqrt{q}}\right). \quad (13.20)$$

That for $Q_{-1/2}$ is [2] 8.13.3: that for $Q_{1/2}$ is [2] 8.13.7. The corresponding formulae for the Legendre P functions, $P_{-1/2}(q_2)$ and $P_{1/2}(q_2)$, though not immediately relevant to the g_n and \hat{g}_n are relevant to solving (RE(α)) and are treated in a later section. We also remark, as in [12] p339,

$$Q_{-1/2}^1(q_2) = -\frac{1}{2\sqrt{q-1}} \text{EllipticE}\left(\frac{1}{\sqrt{q}}\right), \quad (13.21)$$

$$Q_{1/2}^1(q_2) = -\frac{q_2}{2\sqrt{q-1}} \text{EllipticE}\left(\frac{1}{\sqrt{q}}\right) + \sqrt{q-1} \text{EllipticK}\left(\frac{1}{\sqrt{q}}\right). \quad (13.22)$$

and, looking ahead, from (13.31)

$$Q_{-1/2}^{-1}(q_2) = -4 Q_{-1/2}^1(q_2), \quad (13.23)$$

$$Q_{1/2}^{-1}(q_2) = \frac{4}{3} Q_{1/2}^1(q_2). \quad (13.24)$$

[2] 8.2.2 gives, for integer μ ,

$$Q_{-n-1/2}^\mu(z) = Q_{n-1/2}^\mu(z). \quad (13.25)$$

[21]§3.8(9) gives

$$\frac{dQ_{n-1/2}(z)}{dz} = \frac{n^2 - 1/4}{\sqrt{z^2 - 1}} Q_{n-1/2}^{-1}. \quad (13.26)$$

As noted above, the g_n are given by a function of q_2 times $Q_{n-1/2}(q_2)$, the latter functions solving (RE(1/2)). The \hat{g}_n are given by a function of q_2 times $Q_{n-1/2}^{-1}(q_2)$, the latter functions solving (RE(3/2)). Where it seems appropriate we prefer to arrange our formulae so that the $Q_{n-1/2}^{-1}$ are written in terms of $Q_{m-1/2}$, for example from [21]§3.8(5),

$$Q_{n-1/2}^{-1}(z) = \frac{z Q_{n-1/2}(z) - Q_{n-3/2}(z)}{(n + 1/2)\sqrt{z^2 - 1}}. \quad (13.27)$$

(This agrees with (13.13).) Also, as in [21]§3.8(3)

$$Q_{n-1/2}^{-1}(z) = \frac{Q_{n+1/2}(z) - Q_{n-3/2}(z)}{2n\sqrt{z^2 - 1}}. \quad (13.28)$$

(See also Theorem 1 below.) Particular cases of equation (13.27), at $n = 0$ and $n = 1$, are:

$$Q_{-1/2}^{-1}(z) = \frac{z Q_{-1/2}(z) - Q_{1/2}(z)}{(1/2)\sqrt{z^2 - 1}}, \quad (13.29)$$

$$Q_{1/2}^{-1}(z) = \frac{z Q_{1/2}(z) - Q_{-1/2}(z)}{(3/2)\sqrt{z^2 - 1}}. \quad (13.30)$$

We note, as in [2] 8.2.6,

$$Q_{n-1/2}^1(z) = (n^2 - 1/4)Q_{n-1/2}^{-1}(z) \quad (13.31)$$

which can be deduced from [21] §3.8 equations (5) and (7). (This accords with Theorem 3 given in the next section.)

As noted above Conway [14] and various other authors give formulae for g_n (see [14] equation (43)) and for \hat{g}_n in terms of Legendre Q functions.

Concerning the latter, see [14] equation (44),

$$\hat{g}_n = \frac{2(\sqrt{q-1})}{\pi} Q_{n-1/2}^{-1}(q_2) = \frac{2(\sqrt{q-1})}{\pi} \frac{1}{n^2 - 1/4} Q_{n-1/2}^1(q_2), \quad (13.32)$$

$$= \frac{2(\sqrt{q-1})}{\pi} \left(\frac{q_2 Q_{n-1/2}(q_2) - Q_{n-3/2}(q_2)}{(n+1/2)\sqrt{q_2^2-1}} \right). \quad (13.33)$$

(See also [12] p339. It also accords with the corollary to Theorem 1.)

As a check we recall (13.3)

$$\hat{g}_n(q_2 = 1) = -\frac{1}{\pi(n^2 - 1/4)}.$$

[21] §3.9(6) gives asymptotics for q_2 tending down to 1:

$$Q_{n-1/2}^{-1}(q_2) \sim -\frac{1}{\sqrt{2}} \frac{\Gamma(n-1/2)}{\Gamma(n+3/2)} \frac{1}{\sqrt{q_2-1}} = -\frac{1}{\sqrt{2}} \frac{1}{n^2-1/4} \frac{1}{\sqrt{2q-2}}.$$

Using this last expression in (13.32) checks with the elementary result that $\hat{g}_n(q_2 = 1) = -1/(\pi(n^2 - 1/4))$.

13.6.2 Integrals of sums of squares of our Fourier coefficients

Using Parseval's Theorem it is easy to find $\sum_0^\infty \hat{g}_n^2$, etc.

See also [14]§5.

13.7 (RE(α)) and polynomial solutions

The earlier subsections have had a narrow focus on the g_n and \hat{g}_n Fourier coefficients. In this subsection we will focus on polynomial solutions of (RE(α)). Since the general solution of the (RE(α)) difference equation can be formed from a linear combination of $Q_{n-1/2}^\mu$ and $P_{n-1/2}^\mu$ we will, later in this section, be led to consider the Legendre P functions.

13.7.1 Easy results

While too elementary to suggest immediate generalization, we note that when we set $x = q_2 = 1$ in (RE(α)), the recurrence is easy to solve. For all α one solution is $u_n = 1$ for all n . Reduction of order can be used to find another solution.

In particular (RE(1/2)) is then solved in terms of the digamma function ψ , specifically by $u_n = \psi(n + 1/2)$. We have

$$\psi\left(n + \frac{3}{2}\right) = \psi\left(n + \frac{1}{2}\right) + \frac{1}{n + 1/2}.$$

Forming the corresponding equation with n replaced by $n-1$, and subtracting the two equations we find a solution. With

$$\begin{aligned} u_n &= (\psi(n + 1/2) - \psi(1/2))/2, & u_0 &= 0, & u_1 &= 1, \\ u_n &= 1 - (\psi(n + 1/2) - \psi(1/2))/2, & u_0 &= 1, & u_1 &= 0, \end{aligned}$$

we have solutions to the case $x = q_2 = 1$ of (RE(1/2))

When $\alpha = 3/2$ the formulae are simpler. Then we have that the sequence $\hat{u}_n = (n^2 - 1)/(4n^2 - 1)$ satisfies (RE(3/2)) and has $\hat{u}_0 = 1$ and $\hat{u}_1 = 0$. A simple relationship between the \hat{u}_n of this paragraph and the u_n of the preceding is

$$\frac{2}{4n^2 - 1} = \frac{\psi(n + 3/2) - \psi(n - 1/2)}{n}.$$

This can be regarded as a simple lead-in to Theorem 1.

Next some results follow from very routine algebraic manipulation. Define

$$\mathcal{L}(\alpha, u, n) := 2nq_2u_n - (n + \alpha)u_{n+1} - (n - \alpha)u_{n-1}.$$

Theorem 1 *If u_n solves (RE(1/2)) and*

$$\hat{u}_n = \frac{1}{n}(u_{n+1} - u_{n-1}),$$

then \hat{u}_n solves (RE(3/2)).

Proof.

$$\begin{aligned} \mathcal{L}\left(\frac{3}{2}, \hat{u}, n\right) &= 2nq_2 \frac{u_{n+1} - u_{n-1}}{n} - \left(n + \frac{3}{2}\right) \frac{u_{n+2} - u_n}{n+1} - \left(n - \frac{3}{2}\right) \frac{u_n - u_{n-2}}{n-1}, \\ &= \frac{1}{n^2 - 1} \left((n-1)\mathcal{L}\left(\frac{1}{2}, u, n+1\right) + (n+1)\mathcal{L}\left(\frac{1}{2}, u, n-1\right) \right), \\ &= 0. \end{aligned}$$

This establishes the theorem.

Corollary. *If u_n solves (RE(1/2)) and*

$$\hat{u}_n = \frac{xu_n - u_{n-1}}{2n + 1},$$

then \hat{u}_n solves (RE(3/2)), and so too does

$$\hat{u}_n = \frac{xu_n - u_{n+1}}{2n - 1}.$$

(The latter is suggested by [22]3.8(6).)

Remarks.

- We have already seen in (13.12) that

$$\hat{g}_n = \frac{1}{4n(q_2 + 1)} (g_{n+1} - g_{n-1}).$$

See also [21] 3.8.3

- With the definitions in the later subsection on ‘Polynomial sequences’ we have the further examples

$$-4\hat{p}_{01}(n) + 12x\hat{p}_{10}(n) = \frac{3}{n} (p_{10}(n + 1) - p_{10}(n - 1)), \quad (13.34)$$

$$4x\hat{p}_{01}(n) - 12\hat{p}_{10}(n) = \frac{3}{n} (p_{01}(n + 1) - p_{01}(n - 1)). \quad (13.35)$$

(Substituting $x = 1$ and adding the equations has each side summing to 0 which provides one simple check.)

Theorem 2 *If u_n solves (RE(1/2)) and*

$$\hat{u}_n = (2q_2u_n - u_{n+1} - u_{n-1}),$$

then \hat{u}_n solves (RE(3/2)).

Remark. We have already seen in (13.4) that

$$\hat{g}_n = \frac{1}{2(q_2 + 1)} (2q_2g_n - g_{n+1} - g_{n-1}).$$

There are many miscellaneous results. For example a routine calculation shows that

Theorem 3 *If \hat{u}_n satisfies (RE(3/2)), then $(n^2 - 1/4)\hat{u}_n$ satisfies (RE(-1/2)).*

(See also [22]3.8(1) with $\mu = -1$.)

13.7.2 Equivalent recurrence relations

There are many equivalent forms of recurrence relations.

- Defining, from a sequence satisfying (RE(3/2)), the a_n sequence via

$$\hat{u}_{n+1} = a_{n+1} (2q_2)^n \frac{\Gamma(n+1)\Gamma(5/2)}{\Gamma(n+5/2)} = a_{n+1} (2q_2)^n \frac{3 \cdot 2^n n!}{(2n+3)!!},$$

we have, for $n > 1$

$$a_{n+1} = a_n + \frac{(n-3/2)(n+1/2)}{4q_2^2 n(n-1)} a_{n-1}. \quad (\text{RE11}(3/2))$$

(See also [62] equations (5.12) and (5.17) for (RE(1/2)).) While \hat{u}_n might, for some initial conditions be polynomial in q_2 degree $n-1$, the a_n are no longer polynomial, merely rational. Also, we cannot set $n=1$ in the last equation.

- Some studies of recurrence equations for sequences of polynomials involve recurrences for monic polynomials, of the form

$$v_{n+1} = x v_n - f(n) v_{n-1}. \quad (\text{RE1xf})$$

Starting with a sequence of polynomials u_n satisfying (RE(α)), one sets $v_n = u_n/g(n)$ and proceeds as follows:

$$g(n+1)(n+\alpha)v_{n+1} = 2nxg(n)v_n - g(n-1)(n-\alpha)v_{n-1}.$$

For this to be of the required form we must have

$$g(n+1)(n+\alpha) = 2ng(n),$$

which solves to

$$g(n) = c \frac{n!}{\Gamma(n+\alpha)}.$$

From this

$$f(n) = \frac{g(n-1)(n-\alpha)}{g(n+1)(n+\alpha)} = \frac{(n-\alpha)(n-1+\alpha)}{4(n-1)n}.$$

Thus, for $n > 1$

$$v_{n+1} = x v_n + \frac{(n - \alpha)(n - 1 + \alpha)}{4(n - 1)n} v_{n-1}. \quad (\text{RE1x}(\alpha))$$

Again we cannot set $n = 0$ or $n = 1$ in the preceding (except, taking appropriate limits, in the cases $\alpha = 0$ and $\alpha = 1$). For the $\alpha = 1/2$ case see [11] Chapter 6, equation (12.9). See also [62] equations (5.12) and (5.17), but note, as occurs with some other references, the indexing is different from ours, in this case their S_n is our v_{n+1} (monic form of p_{01}).

There is another well-known approach to the calculation of f for (RE1xf). Let (v_n) be a sequence of monic polynomials, mutually orthogonal. We do not, at this stage, need to know the weight function but we will assume it is positive, even in x and the interval is symmetric about $x = 0$. Taking inner products of each side of (RE1xf) with v_{n-1} gives

$$0 = \langle x v_n, v_{n-1} \rangle - f_n \langle v_{n-1}, v_{n-1} \rangle.$$

Shifting the subscripts of the terms in (RE1xf) down by 1, i.e. n replaced by $n - 1$, and then taking the inner product with v_n gives

$$\langle v_n, v_n \rangle = \langle x v_{n-1}, v_n \rangle.$$

The preceding two equations yield

$$f_n = \frac{\langle v_n, v_n \rangle}{\langle v_{n-1}, v_{n-1} \rangle}.$$

We can apply this at $\alpha = 1/2$ to our polynomial sequence $p_{01}(n)$ defined later, see §13.8 and (13.58). The weight function is known explicitly and calculations (from [6], reported in various places including [11]) give

$$\frac{\langle p_{01}(n), p_{01}(n) \rangle}{\langle p_{01}(n-1), p_{01}(n-1) \rangle} = \frac{n-1}{n}.$$

We need monic polynomials. So

$$f_n = \frac{n-1}{n} \left(\frac{\text{lcoeff}(p_{01}(n-1))}{\text{lcoeff}(p_{01}(n))} \right)^2.$$

Using the result on the leading coefficient given in equation (13.63) we find the formula gives

$$f_n = \frac{n-1}{n} \left(\frac{2n-1}{4(n-1)} \right)^2,$$

which agrees with the $\alpha = 1/2$ case given in (RE1x(α)) above.

- A different transformation $v_n = \psi(n, \alpha)\sqrt{n}u_n$, with ψ specified later, but with $\psi(n, 1/2) = 1$, leads to another family of recurrence relations, of the form

$$a_n v_{n+1} = x v_n - a_{n-1} v_{n-1}. \quad (\text{RE}a_n x(\alpha))$$

The recurrence (RE $a_n x(\alpha)$) can be written

$$\begin{pmatrix} v_{n+1} \\ a_n v_n \end{pmatrix} = A(n) \begin{pmatrix} v_n \\ a_{n-1} v_{n-1} \end{pmatrix} \text{ where } A(n) = \begin{bmatrix} x & -\frac{1}{a_n} \\ a_n & 0 \end{bmatrix}. \quad (13.36)$$

(We remark that (RE $a_n x(\alpha)$) can be transformed to (RE1xf), the latter with a different v of course, with $f(n) = a_{n-1}^2$.) The fact that $\det(A(n)) = 1$ leads to generalizations of the result we have at equation (13.67). Specialise now to our (RE(α)). For (RE(1/2)) we have

$$a_n = \frac{n+1/2}{2\sqrt{n(n+1)}}. \quad (13.37)$$

More generally, for (RE(α)) we have

$$\psi(n, \alpha) = \sqrt{\frac{\Gamma(n+\alpha)}{\Gamma(n+1-\alpha)}}, \quad a_n = \frac{\sqrt{\Gamma(n+1+\alpha)\Gamma(n+2-\alpha)}}{\sqrt{n(n+1)\Gamma(n+1-\alpha)\Gamma(n+\alpha)}}.$$

Returning to the case $\alpha = 1/2$, on using a_n of equation (13.37) in $A(n)$ of equation (13.36), we find that the product of matrices

$$A(n) A(n-1) \dots A(2) A(1) = \begin{bmatrix} \sqrt{n+1} p_{01}(n+1) & 4\sqrt{n+1} p_{10}(n+1) \\ \sqrt{n} a(n) p_{01}(n) & 4\sqrt{n} a(n) p_{10}(n) \end{bmatrix},$$

where p_{01} and p_{10} are the polynomials satisfying (RE(1/2)) starting 0,1 and 1,0 respectively.

- It is possible that further work related to the items in this paragraph might be productive. We record them in the hope that some reader might suggest how it might be developed.

- Next a definition. See [11]§5.5. A sequence of positive numbers (f_n) is called a *chain sequence* if there exists another sequence (g_n) such that

$$f_n = g_n(1 - g_{n-1}), \text{ with } 0 \leq g_0 < 1, 0 < g_n < 1 \text{ for } n = 1, 2, \dots$$

The $f(n, \alpha) = a_{n-1}^2$ above is a chain sequence. Use

$$g_n = \frac{1}{2} - \frac{1}{4n} \text{ for } \alpha = \frac{1}{2}.$$

- In some papers the a_n is written as a ratio $a_n = \gamma(n)/\gamma(n-1)$. For $\alpha = 1/2$ the a_n of equation (13.37) has

$$a_n = \frac{\gamma(n)}{\gamma(n-1)} \text{ with } \gamma(n) = \frac{2^{n-1} n!}{\sqrt{n}} \frac{\Gamma(1/2)}{\Gamma(n+1/2)}.$$

13.7.3 Legendre $P_{n-1/2}$ functions

Formulae involving Legendre $P_{n-1/2}$ functions, similar to those in (13.19,13.20,13.21,13.22) can be found.

$$P_{-1/2}(\cos(\theta)) = \frac{2}{\pi} \text{EllipticK}\left(\sin\left(\frac{\theta}{2}\right)\right), \quad (13.38)$$

$$P_{1/2}(\cos(\theta)) = \frac{2}{\pi} \left(2\text{EllipticE}\left(\sin\left(\frac{\theta}{2}\right)\right) - \text{EllipticK}\left(\sin\left(\frac{\theta}{2}\right)\right) \right), \quad (13.39)$$

$$Q_{-1/2}(\cos(\theta)) = \text{EllipticK}\left(\cos\left(\frac{\theta}{2}\right)\right), \quad (13.40)$$

$$Q_{1/2}(\cos(\theta)) = \text{EllipticK}\left(\cos\left(\frac{\theta}{2}\right)\right) - 2\text{EllipticE}\left(\cos\left(\frac{\theta}{2}\right)\right), \quad (13.41)$$

$$P_{-1/2}(\cosh(\xi)) = \frac{2}{\pi \cosh\left(\frac{\xi}{2}\right)} \text{EllipticK}\left(\tanh\left(\frac{\xi}{2}\right)\right), \quad (13.42)$$

$$P_{1/2}(\cosh(\xi)) = \frac{2 \exp(\xi/2)}{\pi} \text{EllipticE}\left(\sqrt{1 - \exp(-2\xi)}\right), \quad (13.43)$$

$$Q_{-1/2}(\cosh(\xi)) = \frac{2}{\sqrt{\pi}} \exp\left(-\frac{\xi}{2}\right) \text{EllipticK}\left(\exp(-\xi)\right), \quad (13.44)$$

$$Q_{1/2}(\cosh(\xi)) = \frac{2}{\sqrt{\pi}} \frac{\cosh(\xi)}{\cosh\left(\frac{\xi}{2}\right)} \text{EllipticK}\left(\frac{1}{\cosh\left(\frac{\xi}{2}\right)}\right) - \quad (13.45)$$

$$\frac{4}{\sqrt{\pi}} \cosh\left(\frac{\xi}{2}\right) \text{EllipticE}\left(\frac{1}{\cosh\left(\frac{\xi}{2}\right)}\right). \quad (13.46)$$

See [2]8.13. There are many equivalent ways of expressing the formulae, and we will use, in a later section,

$$P_{-1/2}(x) = \frac{2}{\pi} \text{EllipticK}\left(\sqrt{\frac{1-x}{2}}\right), \quad Q_{-1/2}(x) = \text{EllipticK}\left(\sqrt{\frac{1+x}{2}}\right) \quad \text{for } |x| < 1.$$

Though we don't explicitly use the following it does relate to the weight w in a later section being even, so we note

$$\frac{\pi}{2} P_{-1/2}(-x) = Q_{-1/2}(x), \quad \text{and similarly } \frac{\pi}{2} P_{1/2}(-x) = -Q_{1/2}(x).$$

The functions with order $\mu = -1$ arise in the case $\alpha = 3/2$. The left-hand parts of the following formulae, corresponding to $\mu = 1$ in [2]8.2.5, parallel

those given earlier, (13.23,13.24),

$$P_{-1/2}^{-1}(q_2) = -4 P_{-1/2}^1(q_2), \quad (13.47)$$

$$P_{1/2}^{-1}(q_2) = \frac{4}{3} P_{1/2}^1(q_2). \quad (13.48)$$

$$Q_{-1/2}^{-1}(q_2) = -4 Q_{-1/2}^1(q_2) = \frac{4}{\sqrt{2(q_2-1)}} \text{EllipticE} \left(\sqrt{\frac{2}{q_2+1}} \right) \quad (13.49)$$

$$Q_{1/2}^{-1}(q_2) = \frac{4}{3} \frac{-q_2}{\sqrt{2(q_2-1)}} \text{EllipticE} \left(\sqrt{\frac{2}{q_2+1}} \right) + \quad (13.50)$$

$$\frac{4}{3} \sqrt{\frac{q_2-1}{2}} \text{EllipticK} \left(\sqrt{\frac{2}{q_2+1}} \right). \quad (13.51)$$

There are many remarkable formulae involving both P and Q functions, including Whipple's, which for integer n and μ is, for $z > 1$,

$$P_{n-\frac{1}{2}}^\mu(z) = \frac{(-1)^n}{\Gamma(n-\mu+\frac{1}{2})} \sqrt{\frac{2}{\pi}} (z^2-1)^{-1/4} Q_{\mu-\frac{1}{2}}^n\left(\frac{z}{\sqrt{z^2-1}}\right)$$

$$Q_{n-\frac{1}{2}}^\mu(z) = \frac{(-1)^n \pi}{\Gamma(n-\mu+\frac{1}{2})} \sqrt{\frac{\pi}{2}} (z^2-1)^{-1/4} P_{\mu-\frac{1}{2}}^n\left(\frac{z}{\sqrt{z^2-1}}\right).$$

When both n and μ are zero, we have

$$P_{-\frac{1}{2}}(z) = \frac{\sqrt{2}}{\pi} (z^2-1)^{-1/4} Q_{-\frac{1}{2}}\left(\frac{z}{\sqrt{z^2-1}}\right),$$

$$P_{-\frac{1}{2}}(\cosh(\eta)) = \frac{\sqrt{2}}{\pi} (\sinh(\eta))^{-1/2} Q_{-\frac{1}{2}}(\coth(\eta)),$$

$$Q_{-\frac{1}{2}}(z) = \frac{\pi}{\sqrt{2}} (z^2-1)^{-1/4} P_{-\frac{1}{2}}\left(\frac{z}{\sqrt{z^2-1}}\right),$$

$$Q_{-\frac{1}{2}}(\cosh(\eta)) = \frac{\pi}{\sqrt{2}} (\sinh(\eta))^{-1/2} P_{-\frac{1}{2}}(\coth(\eta)).$$

Another notable formula is from [22]3.3.1(1)

$$P_{-n-1/2}^\mu(z) = P_{n-1/2}^\mu(z). \quad (13.52)$$

The corresponding formula for Q is (13.25) given in [22]3.3.1(3).

There are formulae involving Wronskians and similar determinantal forms, e.g. for $z > 0$

$$P_{1/2}(z)Q_{-1/2}(z) - Q_{1/2}(z)P_{-1/2}(z) = 2. \quad (13.53)$$

On substituting from equations (13.38)–(13.41) one finds that equation (13.53) is equivalent to Legendre’s relation (10.1) for the EllipticE and EllipticK functions.

[2]8.1.8 gives, when $\mu = 0$,

$$\text{Wronskian}(P_\nu(z), Q_\nu(z)) = \frac{-1}{z^2 - 1}.$$

(See also [21]3.4(25).)

We also have from [21] 3.8 (9) and [2]8.5.2

$$\frac{dP_{n-1/2}(z)}{dz} = \frac{n^2 - 1/4}{\sqrt{z^2 - 1}} P_{n-1/2}^{-1}. \quad (13.54)$$

Using this and the corresponding formula for the Q , namely (13.26)

$$P_{n-1/2}(z)Q_{n-1/2}^{-1}(z) - P_{n-1/2}^{-1}(z)Q_{n-1/2}(z) = -\frac{1}{(n^2 - 1/4)\sqrt{z^2 - 1}}.$$

(See also [25] equation (7), as in (13.57) below.)

The Wronskian expression and [2] 8.5.4

$$P_{n+1/2}(z)Q_{n-1/2}(z) - Q_{n+1/2}(z)P_{n-1/2}(z) = \frac{1}{n + 1/2}. \quad (13.55)$$

(See also equation (13.67).) Equation (13.55) is the $m = 0$ case of equation (6) of [25]:

$$P_{n+1/2}^m(z)Q_{n-1/2}^m(z) - Q_{n+1/2}^m(z)P_{n-1/2}^m(z) = \frac{\Gamma(n + \frac{1}{2} + m)}{\Gamma(n + \frac{3}{2} - m)} (-1)^m. \quad (13.56)$$

Equation (7) of [25] allows for orders to be changed by 1

$$P_{n-1/2}^m(z)Q_{n-1/2}^{m+1}(z) - Q_{n-1/2}^m(z)P_{n-1/2}^{m+1}(z) = \frac{\Gamma(n + \frac{1}{2} + m)}{\Gamma(n + \frac{1}{2} - m)} \frac{(-1)^m}{\sqrt{z^2 - 1}}. \quad (13.57)$$

Writing z where formerly we wrote q_2 , the general solution of (RE(1/2)) is

$$u_n(z) = f_P(z)P_{n-1/2}(z) + f_Q(z)Q_{n-1/2}(z),$$

where f_P and f_Q are independent of n . We can prescribe u_0 and u_1 and determine f_P and f_Q . Using equation (13.53) we have

$$f_P = -\frac{1}{2}(u_0 Q_{1/2} - u_1 Q_{-1/2}), \quad f_Q = \frac{1}{2}(u_0 P_{1/2} - u_1 P_{-1/2}).$$

In particular, the sequence starting with $u_0 = 0$, $u_1 = 1$, and denoted by $p_{01}(n)$ in the next subsection is

$$p_{01}(n) = \frac{1}{2}(Q_{-1/2} P_{n-1/2} - P_{-1/2} Q_{n-1/2}). \quad (13.58)$$

Similarly

$$p_{10}(n) = -\frac{1}{2}(Q_{1/2} P_{n-1/2} - P_{1/2} Q_{n-1/2}). \quad (13.59)$$

While these must be the polynomial sequences tabulated in the next subsection, it isn't immediately obvious how to extract information about the polynomials from the formula above. (For general α , see [10] equation (3.5).) One item that follows readily is that the above with equations (13.25) and (13.52) imply $p_{01}(-n) = p_{01}(n)$, etc.

13.7.4 Reduction of order

We have already noted that reduction of order can be used when $x = q_2 = 1$ producing from the obvious constant solution to $(\text{RE}(\alpha))$ a second linearly independent solution. This subsection fails to help so far in our quest to find more information on the polynomial solutions. However reduction of order gives equation (13.60) and this can be used to give a relation between toroidal P and Q functions.

Assume u_n is one solution of $(\text{RE}(\alpha))$ and seek a second linearly independent solution v_n with

$$v_n = w_n u_n \quad \text{with } w_1 \neq w_0 \quad \text{and with all } u_n \neq 0\text{-function.}$$

Writing out the equation stating that $w_n u_n$ satisfies $(\text{RE}(\alpha))$ and then eliminating u_n from the fact that u_n also satisfies $(\text{RE}(\alpha))$ gives

$$(n + \alpha)u_{n+1}(w_{n+1} - w_n) + (n - \alpha)u_{n-1}(w_{n-1} - w_n) = 0.$$

Let $\theta_n = w_{n+1} - w_n$. Then

$$(n + \alpha)u_{n+1}\theta_n - (n - \alpha)u_{n-1}\theta_{n-1} = 0.$$

Hence

$$\frac{\theta_n}{\theta_{n-1}} = \frac{(n - \alpha)u_{n-1}}{(n + \alpha)u_{n+1}},$$

and hence

$$\frac{\theta_n}{\theta_{n-2}} = \frac{(n - \alpha)(n - 1 - \alpha)u_{n-1}u_{n-2}}{(n + \alpha)(n - 1 + \alpha)u_{n+1}u_n}.$$

Now set $\alpha = 1/2$. The last equation becomes

$$\frac{\theta_n}{\theta_{n-2}} = \frac{(n - 2 + \frac{1}{2})u_{n-1}u_{n-2}}{(n + \frac{1}{2})u_{n+1}u_n}.$$

Continuing to multiply the ratios of θ we get

$$\frac{\theta_n}{\theta_0} = \frac{(\frac{1}{2})u_1u_0}{(n + \frac{1}{2})u_{n+1}u_n}.$$

Our concern is with w rather than θ and

$$w_{n+1} - w_0 = \sum_{k=0}^n \theta_k.$$

Now θ_0 is known and nonzero as $\theta_0 = w_1 - w_0$. Hence

$$w_{n+1} = w_0 + \theta_0 u_0 u_1 \sum_{k=0}^n \frac{1}{(2k + 1)u_{k+1}u_k}. \quad (13.60)$$

The preceding equation concludes the general reduction of order. However, a quick check is to return to the case $x = q_2 = 1$ and the known solution $u_k = 1$ for all k . The reduction of order formula for w_{n+1} involves the sum of the reciprocals of odd integers, and this is consistent with our previous formula involving the digamma function ψ .

We remark that one could view one instance as $w_n = P_{n-1/2}/Q_{n-1/2}$ and then one has an identity involving Legendre functions. It presumably is a much disguised consequence of the recurrence relations (RE(1/2)) satisfied by the P and Q .

We remark also that if the input u_n to the reduction of order is polynomial in $x = q_2$, the output v_n is rational but not necessarily polynomial.

For a formula similar to equation (13.60) see [27] equation (14).

13.8 Polynomial sequences relating to LegendreQ functions

Clearly we can write

$$Q_{n-1/2}(z) = p_{10}(\alpha = 1/2, n) Q_{-1/2}(z) + p_{01}(\alpha = 1/2, n) Q_{1/2}(z), \quad (13.61)$$

where the polynomials p satisfy (RE(1/2)) and

$$p_{10}(\alpha = 1/2, 0) = 1, \quad p_{10}(\alpha = 1/2, 1) = 0, \quad p_{01}(\alpha = 1/2, 0) = 0, \quad p_{01}(\alpha = 1/2, 1) = 1.$$

When space is too limited, we drop from the argument list the $\alpha = 1/2$, leaving n as the sole argument.

n	$p_{01}(n)$	$p_{10}(n)$
0	0	1
1	1	0
2	$\frac{4}{3}x$	$-\frac{1}{3}$
3	$\frac{32}{15}x^2 - \frac{3}{5}$	$-\frac{8}{15}x$
4	$\frac{128}{35}x^3 - \frac{208}{105}x$	$-\frac{32}{35}x^2 + \frac{5}{21}$
5	$\frac{2048}{315}x^4 - \frac{544}{105}x^2 + \frac{7}{15}$	$-\frac{512}{315}x^3 + \frac{88}{105}x$

Plots of the p_{01} polynomials are given in Figure 10.




Figure 10: Plots of $p_{01}(n)$ for $n = 1$ to 5 .

Similarly, we write

$$Q_{n-1/2}^{-1}(z) = \hat{p}_{10}(\alpha = 1/2, n) Q_{-1/2}(z) + \hat{p}_{01}(\alpha = 1/2, n) Q_{1/2}(z), \quad (13.62)$$

where the polynomials \hat{p} satisfy (RE(3/2)) and

$$\hat{p}_{10}(\alpha = 3/2, 0) = 1, \quad \hat{p}_{10}(\alpha = 3/2, 1) = 0, \quad \hat{p}_{01}(\alpha = 3/2, 0) = 0, \quad \hat{p}_{01}(\alpha = 3/2, 1) = 1.$$

Thus, p_{01} and p_{10} are the polynomials corresponding to $\alpha = 1/2$ and \hat{p}_{01} and \hat{p}_{10} are those corresponding to $\alpha = 3/2$.

n	$\hat{p}_{01}(n)$	$\hat{p}_{10}(n)$
0	0	1
1	1	0
2	$\frac{4}{5}x$	$\frac{1}{5}$
3	$\frac{32}{35}x^2 - \frac{1}{7}$	$\frac{8}{35}x$
4	$\frac{128}{105}x^3 - \frac{16}{35}x$	$\frac{32}{105}x^2 - \frac{1}{45}$
5	$\frac{2048}{1155}x^4 - \frac{416}{385}x^2 + \frac{5}{77}$	$\frac{512}{1155}x^3 - \frac{232}{1155}x$

We have seen in earlier sections that there are many connections between the polynomials at $\alpha = 1/2$ and $\alpha = 3/2$. For example, defining

$$p_{1x}(n) = p_{10}(n) + x p_{01}(n),$$

Theorem 1 suggests relations with the \hat{p} . We have

$$\frac{p_{1x}(n+1) - p_{1x}(n-1)}{x^2 - 1} = \frac{3\hat{p}_{01}(n)}{4n}.$$

This also follows from equations (13.34) (13.35).)

Various quantities, whose notation has been standardised, as in [22]§10, are associated with sequences of orthogonal polynomials. These include the leading coefficient, often denoted k_n , moments, often denoted c_n and discussed in the later subsection with the weight function, and the norm squared of the polynomial, denoted h_n . (The quantities k_n and h_n occur in [22]10.3(11) in connection with a Christoffel-Darboux formula, a formula we will present in a later subsection.) For the family p_{01} we have

$$\text{lcoeff}(p_{01}(n)) = k_n = \frac{2^{2n-2}(n-1)!}{(2n-1)!}. \quad (13.63)$$

The leading coefficients of the other sequences of polynomials satisfy, for $n \geq 2$,

$$\begin{aligned} \text{lcoeff}(p_{01}(n)) &= -4 \text{lcoeff}(p_{10}(n)), \\ \text{lcoeff}(\hat{p}_{01}(n)) &= 4 \text{lcoeff}(\hat{p}_{10}(n)), \\ \text{lcoeff}(\hat{p}_{01}(n)) &= \frac{3}{2n+1} \text{lcoeff}(p_{01}(n)), \\ \text{lcoeff}(\hat{p}_{10}(n)) &= -\frac{3}{2n+1} \text{lcoeff}(p_{10}(n)). \end{aligned}$$

[22]10.3(7,8) writes the recurrence relation in the form

$$p_{n+1} = A_n x p_n - C_n p_{n-1},$$

and notes that

$$A_n = \frac{k_{n+1}}{k_n}, \quad C_n = \frac{k_{n+1} k_{n-1}}{k_n^2} \frac{h_n}{h_{n-1}}.$$

For us at $\alpha = 1/2$,

$$A_n = \frac{4n}{2n+1}, \quad \frac{h_n}{h_m} = \frac{m}{n}, \quad C_n = \frac{2n-1}{2n+1}.$$

Our interest is in g_n and \hat{g}_n , so $\alpha = 1/2$ and $\alpha = 3/2$. There remains the possibility that investigating other values of α might ultimately be useful, and, perhaps lead to the discovery of further ‘contiguous relations’ such as those in μ for the Legendre functions. In this connection we record some results on $\alpha = -1/2$.

n	$p_{01}(\alpha = -\frac{1}{2}, n)$	$p_{10}(\alpha = -\frac{1}{2}, n)$
0	0	1
1	1	0
2	$4x$	-3
3	$\frac{32}{3}x^2 - \frac{5}{3}$	$-8x$
4	$\frac{128}{5}x^3 - \frac{48}{5}x$	$-\frac{96}{5}x^2 + \frac{21}{5}$
5	$\frac{2048}{35}x^4 - \frac{1248}{35}x^2 + \frac{15}{7}$	$-\frac{1536}{35}x^3 + \frac{696}{35}x$

We focus on connections between the solutions of (RE(-1/2)) and those of (RE(1/2)) and (RE(3/2)). As noted in Theorem 3,

$$p_{01}(\alpha = -1/2, n) = \frac{4n^2 - 1}{3} \hat{p}_{01}(n).$$

13.8.1 Derivatives of p_{01} and of p_{10}

Ordinary differential equations

The Legendre differential equation satisfied by $Q_{n-1/2}(z)$ leads to coupled d.e.s for $p_{01}(n)$ and $p_{10}(n)$ (where we drop the remainder $\alpha = 1/2$).

$$(1-x^2)\frac{d^2p_{10}(n)}{dx^2} - x\frac{dp_{10}(n)}{dx} + p_{10}(n)n^2 + \frac{dp_{01}(n)}{dx} = 0, \quad (13.64)$$

$$(1-x^2)\frac{d^2p_{01}(n)}{dx^2} - 3x\frac{dp_{01}(n)}{dx} + (n^2-1)p_{01}(n) - \frac{dp_{10}(n)}{dx} = 0. \quad (13.65)$$

From these one can find 4-th order differential equations satisfied by the polynomials. For example $p_{01}(n)$, satisfies, with

$$\begin{aligned} P &= (1-x^2)^2, \\ Q &= 10x(1-x^2), \\ R &= 2n^2(1-x^2) - 9 + 25x^2, \\ S &= -3x(2n^2-5), \end{aligned}$$

the differential equation

$$P\frac{d^4p_{01}(n)}{dx^4} + Q\frac{d^3p_{01}(n)}{dx^3} + R\frac{d^2p_{01}(n)}{dx^2} + S\frac{dp_{01}(n)}{dx} + (n^2-1)^2p_{01}(n) = 0.$$

The notation is as in [49]. The conditions in the theorems in [49] are *not* satisfied, but some equations similar to his are. We have, with $W_M = \sqrt{1-x^2}$ which is *not* the weight function for us,

$$\begin{aligned} W_M Q &= 2\frac{d}{dx}(W_M P), \\ W_M S &= \frac{d}{dx}(W_M(R-1)) - \frac{d^3}{dx^3}(W_M P). \end{aligned}$$

The remainder of this subsection is speculative (and won't be included in any paper). We also don't see any use for the main goals of the paper. We have yet to write the d.e. in the form

$$\mathcal{L}_4 u = f(x, n)\mathcal{L}_2 u,$$

with, in some appropriate sense, the 4th order \mathcal{L}_4 and the 2nd order \mathcal{L}_2 being somehow 'self-adjoint'. Associated with this effort, we would like to be able

to express derivatives of the $w(x)$ defined in equation (13.76) in terms of a rational function of $w(x)$, x and $\sqrt{1-x^2}$ (or something similarly simple).

We believe there is much more to be discovered, e.g. in connection with variational characterisations of the (generalized eigen)solutions of the fourth order operator. We expect that the qualitative behaviour would be akin to the generalized matrix eigenvalue problem

$$Au = \lambda Bu,$$

with matrices A and B self-adjoint (real-symmetric). The function spaces involved might include the space of all polynomials. Having found (eigen)solutions polynomial of degree n , Rayleigh Ritz might be appropriate to produce solutions of degree $n + 1$.

Different inner products may be appropriate in different circumstances.

The Wronskian of p_{01} and p_{10}

We find, at any fixed n , using equations (13.69), (13.70) to eliminate the derivative terms and then equation (13.67) to eliminate the polynomials indexed by $n - 1$,

$$\begin{aligned} \text{Wronskian}(p_{01}, p_{10}) &= (p_{01} p'_{10} - p_{10} p'_{01}), \\ &= \frac{(p_{01} + p_{10})^2 + 2(x-1)p_{01}p_{10} - 1}{2(x^2 - 1)}. \end{aligned} \quad (13.66)$$

13.8.2 Contiguous relations

We have already seen several instances of ‘contiguous relations’.

- Some of these involve only the polynomials at a fixed value of α as in (RE(α)) and, in this category, we mention (13.67) and (13.68) below.
- Some of these involve polynomials at a fixed n and the different sequences p at $\alpha = 1/2$ and \hat{p} at $\alpha = 3/2$. One such, namely (13.34), (13.35), was a simple consequence of Theorem 1.
- Others involve not merely the polynomials, but also their first derivatives.

We now see an identity (13.67) for the polynomial solutions of (RE(1/2)) which is similar to (13.55) applying to LegendreQ. At $\alpha = 1/2$ we have

$$p_{10}(n) p_{01}(n+1) - p_{10}(n+1) p_{01}(n) = \frac{1}{2n+1}. \quad (13.67)$$

(See also the preceding subsection on alternative formulations of the recurrence relation, specifically (RE1x a_n).) At $\alpha = 3/2$ we have

$$\hat{p}_{10}(n)\hat{p}_{01}(n+1) - \hat{p}_{10}(n+1)\hat{p}_{01}(n) = \frac{-3}{(2n-1)(2n+1)(2n+3)}. \quad (13.68)$$

(There is a similar result for $\alpha = 0$:

$$T_n(x)U_{n+1}(x) - T_{n+1}(x)U_n(x) = 1.$$

And, again, another similar result for $\alpha = 1$.)

These are easily proved by induction, as given in the following generalization which applies not just to polynomial solutions of the difference equations.

Theorem 4 *Let u_n and U_n be two sequences satisfying (RE(α)). Then*

$$u_n U_{n+1} - u_{n+1} U_n = c(x) \frac{\Gamma(n+1-\alpha)}{\Gamma(n+1+\alpha)},$$

where

$$c(x) = (u_0 U_1 - u_1 U_0) \frac{\Gamma(1+\alpha)}{\Gamma(1-\alpha)}.$$

Proof. The proof is by induction. Define

$$\mathcal{S}(n) = u_n U_{n+1} - u_{n+1} U_n - c(x) \frac{\Gamma(n+1-\alpha)}{\Gamma(n+1+\alpha)},$$

with $c(x)$ as in the statement of the theorem. We have, from the definition of $c(x)$, that $\mathcal{S}(0) = 0$. We will next show that $\mathcal{S}(n-1) = 0$ implies that $\mathcal{S}(n) = 0$. Let $\mathcal{S}^*(n)$ be the expression obtained from $\mathcal{S}(n)$ on using (RE(α)) to write u_{n+1} in terms of u_n and u_{n-1} and, similarly, to write U_{n+1} in terms of U_n and U_{n-1} . One finds

$$\mathcal{S}^*(n) = \frac{n-\alpha}{n+\alpha} \mathcal{S}(n-1).$$

Thus $\mathcal{S}(n-1) = 0$ implies $\mathcal{S}(n) = 0$ so, as asserted, the theorem is established by induction.

Formulae involving the derivatives of the polynomials can also be found.

The starting point in deriving (13.69) and (13.70) below is

$$(x^2 - 1) \frac{dQ_\nu}{dx} = \nu (xQ_\nu - Q_{\nu-1}).$$

See [21] 3.8.10. Also

$$\begin{aligned} (x^2 - 1) \frac{dQ_{1/2}}{dx} &= \frac{1}{2} (xQ_{1/2} - Q_{-1/2}), \\ (x^2 - 1) \frac{dQ_{-1/2}}{dx} &= -\frac{1}{2} (xQ_{-1/2} - Q_{1/2}). \end{aligned}$$

This leads to the somewhat lengthy identities

$$2(x^2 - 1) \frac{dp_{10}(j)}{dx} = 2jxp_{10}(j) + p_{01}(j) - (2j - 1)p_{10}(j - 1), \quad (13.69)$$

$$2(x^2 - 1) \frac{dp_{01}(j)}{dx} = (2j - 2)xp_{01}(j) - p_{10}(j) - (2j - 1)p_{01}(j - 1). \quad (13.70)$$

There are also formulae involving the four sequences of polynomials, both the p sequences and the \hat{p} sequences. The starting point for equations (13.71) and (13.72) below is that derivatives with respect to z of $Q_{n-1/2}^\mu(z)$ can be expressed in terms of the undifferentiated function at a different value of μ . In particular, begin with equation (13.26). After some calculation one finds

$$(4n^2 - 1) \hat{p}_{01}(n) = 6 \frac{d}{dx} (x p_{01}(n) + p_{10}(n)) - 3p_{01}(n), \quad (13.71)$$

$$(4n^2 - 1) \hat{p}_{10}(n), = 2 \frac{d}{dx} (x p_{10}(n) + p_{01}(n)) - 3p_{10}(n). \quad (13.72)$$

13.8.3 Christoffel-Darboux (CD)

As in [22]10.3(11) we have

$$\sum_{j=1}^n j p_{01}(j)^2 = \frac{2n+1}{4} \left(p_{01}(n) \frac{dp_{01}(n+1)}{dx} - \frac{dp_{01}(n)}{dx} p_{01}(n+1) \right).$$

The CD kernel, defined by

$$K(x, y) = \sum_{j=1}^n j p_{01}(j, x) p_{01}(j, y),$$

has the reproducing property

$$K(x, y) = \langle K(x, \cdot), K(\cdot, y) \rangle.$$

13.8.4 A convolution identity involving Legendre polynomials

A special case of [10] equation (3.13) is

$$p_{10}(n) = - \sum_{k=0}^n \frac{\text{LegendreP}(k, x) \text{LegendreP}(n - k, x)}{2k - 1}. \quad (13.73)$$

Another identity (also in [11] (12.6) is

$$p_{01}(n) = - \sum_{k=0}^{n-1} \frac{\text{LegendreP}(k, x) \text{LegendreP}(n - 1 - k, x)}{2k + 1}. \quad (13.74)$$

(Equation (13.74) can also be found using the generating function given in (13.14).)

We have yet to find similar concise identities for $\alpha = 3/2$ and the \hat{p} polynomials. Of course, using equations (13.74), (13.73) in equation (13.35) gives the $4x\hat{p}(n) - 12\hat{p}(n)$ in terms of Legendre polynomials.

Presumably related to $\alpha = 0$ or $\alpha = 1$ we have the well-known identity ([11] (12.7)):

$$\text{ChebyshevU}(n, x) = \sum_{k=0}^n \text{LegendreP}(k, x) \text{LegendreP}(n - k, x).$$

We have yet to find a similar identity for ChebyshevT.

There are, of course, many other ways the identities can be written as, if f is any real-valued function odd about 0 (and any functions of (k, x) not just LegendreP),

$$0 = \sum_{k=0}^n f\left(k - \frac{n}{2}\right) \text{LegendreP}(k, x) \text{LegendreP}(n - k, x).$$

A typical example would be $f\left(k - \frac{n}{2}\right) = k - \frac{n}{2}$.

Allowing other Gegenbauer polynomials than the Legendre P polynomials above, we remark that one obtains convolution identities on using $\gamma = -\beta = \alpha - 1$ in [10] equation (3.13).

13.9 $\alpha = 1/2$: orthogonality and related results

Results from [6] are reported in [11] and [27]. Equation (12.3) of [11] and equation(1) of [27] defines for the interval $(-1, 1)$, a weight function w (in

the case $\nu = 1/2$ appropriate to our problem) by

$$w(x) = \frac{4}{Q_{-1/2}(x)^2 + (\pi P_{-1/2}(x)/2)^2}. \quad (13.75)$$

Using representations of the toroidal functions in terms of complete elliptic integrals, `EllipticK`, this rewrites to

$$w(x) = \frac{4}{K\left(\sqrt{\frac{1-x}{2}}\right)^2 + K\left(\sqrt{\frac{1+x}{2}}\right)^2}. \quad (13.76)$$

(See also [62] where, with A_n satisfying their (1.5), $k^n A_n$ satisfies (RE(1/2)). Also their ψ_n satisfying their (4.10) satisfy our (RE(1/2)), except for a shift. Weight functions involving `EllipticK` occur in their (3.7), (3.9), (5.19) and, close to the form we report in (5.20) and (5.21). See also [53].) We have yet to investigate (RE(1/2)) in connection with study of examples of ‘polynomials orthogonal on the (boundary of the) unit disk’: see equation (10.2).

Define the inner product

$$\langle u, v \rangle = \int_{-1}^1 u(x) v(x) w(x) dx.$$

Direct calculation (via `wolframalpha`) for several (m, n) pairs confirms

$$\langle p_{01}(m), p_{01}(n) \rangle = \frac{\delta_{mn}}{n}. \quad (13.77)$$

Shortly we will give an elementary derivation of this without reference to the details of the inner product.

The moments are defined by

$$c_n = \langle 1, x^n \rangle,$$

and are obviously 0 when n is odd. We have

$$c_0 = 1, \quad c_2 = \frac{9}{2^5}, \quad c_4 = \frac{39}{2^8}, \quad c_6 = \frac{6633}{2^{16}},$$

but have yet to find the formula for c_{2n} . Once all the c_n are calculated, [22] 10.3(4) gives a determinantal formula for the orthogonal polynomials.

Our derivation of equation (13.77) involves considering

$$\phi(n) = \langle x^{n-1}, p_{01}(n) \rangle.$$

(See also [11] Theorems 3.2 and 4.2.) Multiplying the recurrence

$$2nxp_{01}(n) = (n + \frac{1}{2})p_{01}(n+1) + (n - \frac{1}{2})p_{01}(n-1),$$

by x^{n-2} and taking the inner product gives

$$2n\phi(n) = (n - \frac{1}{2})\phi(n-1).$$

If one agrees that $\phi(1) = \langle 1, 1 \rangle = 1$, this first order recurrence for ϕ solves to

$$\phi(n) = \frac{(2n-1)!!}{4^{n-1}n!} = \frac{1}{n \text{lcoeff}(p_{01}(n))},$$

where the leading coefficient lcoeff is given by equation (13.63). Next consider $\langle p_{01}(n), p_{01}(n) \rangle$ beginning with considering

$$p_{01}(n) = \text{lcoeff}(p_{01}(n))x^{n-1} + \text{lower order},$$

and noting that the inner product of the lower order monomials with $p_{01}(n)$ is 0. Thus

$$\langle p_{01}(n), p_{01}(n) \rangle = \text{lcoeff}(p_{01}(n)) \langle x^{n-1}, p_{01}(n) \rangle = \text{lcoeff}(p_{01}(n))\phi(n).$$

From the preceding calculated value of $\phi(n)$ we find the value stated in (13.77).

13.9.1 α generally

The orthogonality result is also available, in a more general form in [22], p220, §10.21 where one of the Pollaczek sequences is treated. The weight function, for our $\alpha = 1/2$ case is given in [22]10.21(14) when one sets

$$a = b = 0 = t, \quad c = \lambda = \frac{1}{2}.$$

These values in (13) yield our (RE(1/2)) while, in (14), the weight function involves, as before, EllipticK functions as the hypergeometric expression in (14) is

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; z\right) = \frac{2}{\pi}K(\sqrt{z}).$$

For $\alpha = 3/2$ we have, in both (13) and (14),

$$a = b = 0 = t, \quad c = \frac{3}{2}, \quad \lambda = -\frac{1}{2}.$$

This time the weight function given in (14) is ${}_2F_1(\frac{3}{2}, \frac{3}{2}; 1; z)$ which, again, can be expressed in terms of EllipticE and EllipticK. We comment further on the $\alpha = 3/2$ case below.

13.9.2 More concerning $\alpha = 3/2$

We noted, in connection with Pollaczek polynomials above, that the weight function for our $\alpha = 3/2$ case is available. However, the setting is considerably more general than appropriate for our very special cases.

Gegenbauer polynomials (themselves special cases of Jacobi polynomials) satisfy the recurrence

$$(n+1)C_{n+1}^\lambda(x) = 2(n+\lambda)x C_n^\lambda(x) - (n+2\lambda-1)C_{n-1}^\lambda(x). \quad (\text{REG})$$

- Our (RE(1)), satisfied by scaled Chebyshev polynomials, is the $\lambda = 0$ case of this.
- (RE(0)) is the $\lambda = 1$ case of (REG).

However (REG) does not include the (RE(α)) needed in our paper. The case $\lambda = 1/2$ in (REG) gives Legendre polynomials.

[10]§3, equation (3.1) modify this to

$$(n+\lambda+1)C_{n+1}^\lambda(x, \beta) = 2x(n+\beta+\lambda)x C_n^\lambda(x, \beta) - (2\beta+n+\lambda-1)C_{n-1}^\lambda(x, \beta). \quad (\text{BI3.2})$$

With $\lambda = \alpha - 1 = -\beta$ we have our (RE(α)). In particular

- $\lambda = \beta = 0$ is our (RE(1)).
- $\lambda = -1 = -\beta$ is our (RE(0)).
- $\lambda = -1/2 = -\beta$ is our (RE(1/2)).
- $\lambda = 1/2 = -\beta$ is our (RE(3/2)).

13.9.3 An aside on other cases in [27]

This is an aside as the only case which is amongst our $(\text{RE}(\alpha))$ is the case $\alpha = 1/2$.

The recurrence relation at the beginning of [27] is, with a shift so as to accord with our indexing,

$$(n + \nu)p_{n+1} = (2n + 2\nu - 1)x p_n - (n + \nu - 1)p_{n-1}.$$

We have already used that when $\nu = 1/2$ this is our $(\text{RE}(1/2))$. The weight function for orthogonality is

$$w(\nu, x) = \frac{1}{(\nu Q_{\nu-1}(x))^2 + (2\nu P_{\nu-1}(x)/\pi)^2}.$$

Now, the limit of $w(\nu, x)$ as ν tends to 0 is 1, so we will write

$$w(0, x) = 1.$$

The recurrence is solved by the Legendre polynomials and these are orthogonal with respect to the weight $w(0, x)$.

We also have

$$\begin{aligned} w(1, x) &= \frac{1}{Q_0(x)^2 + (\pi P_0(x)/2)^2}, \\ &= \frac{1}{\left(\frac{1}{2} \log \left| \frac{1+x}{1-x} \right| \right)^2 + \frac{\pi^2}{4}}. \end{aligned}$$

Repeating work in [27], the usual notation for the LegendreQ functions is

$$Q_n(x) = P_n(x)Q_0(x) - W_{n-1}(x).$$

In [27] there is the demonstration that the polynomials W_n are orthogonal with respect to the weight $w(1, x)$.

13.10 Bessel functions

In view of one of our goals being to sum series such as that for u_∞ there is a case for investigating representations of the toroidal functions.

Connections with Bessel functions are available. See [56] where the reference is to Watson's classic book on Bessel functions. See also [13] equation

(3) and references there. The conditions on the parameters are not those we need here. The formulae below are sometimes approached via Laplace transforms. We have, for $q > 1$,

$$\sqrt{q} \int_0^\infty \exp(-t\sqrt{q}) \text{BesselI}(0, t \cos(\psi)) dt = g(\psi) = \frac{1}{\sqrt{1 - \frac{\cos(\psi)^2}{q}}}. \quad (13.78)$$

The Fourier coefficients of g , $Q_{n-1/2}(q_2)$ can also be written in terms of integrals of Bessel functions. We have, for $s > 1$,

$$\int_0^\infty \exp(-st) \text{BesselI}(n, \frac{t}{2})^2 dt = \frac{2}{\pi} Q_{n-1/2}(2s^2 - 1),$$

and its application here would have $s = \sqrt{q}$. See [26] 6.612.3.

13.11 Boundary moments revisited

We remark that the perimeter can be expressed in terms of a toroidal function: see equation (9.6).

Having introduced the g_n and \hat{g}_n , we might note that boundary moments can be expressed in terms of them. For example, as

$$r^2 = x^2 + y^2 = a^2(1 - e^2) \left(1 - \frac{e^2}{2} + \frac{e^2}{2} \cos(2\psi) \right),$$

we have

$$\begin{aligned} i_2 &= a \int_{-\pi}^{\pi} r^2 \hat{g}(\psi) d\psi \\ &= a^3(1 - e^2) \left(\left(1 - \frac{e^2}{2}\right) \frac{\hat{g}_0}{2} + \frac{e^2}{2} \hat{g}_1 \right). \end{aligned}$$

There may be useful recurrence relations involving the i_n .

It may be that there are neat expressions for any moment, not merely the ‘polar’ moments of the preceding paragraph, involving toroidal functions, both of the LegendreP and LegendreQ kinds.

Part III: A general setting for variational approximation

$$R \leq Q$$

By G. Keady

14 A general lower bound

A published form of the important results of this part is [39]. Further applications are given in [35, 36]

14.1 A development from [37]

Our notation is as in [37]. For large β

$$u \sim \beta \frac{|\Omega|}{|\partial\Omega|} + u_\infty + o(1),$$

where u_∞ solves

$$-\frac{\partial^2 u_\infty}{\partial x^2} - \frac{\partial^2 u_\infty}{\partial y^2} = 1 \text{ in } \Omega, \quad \frac{\partial u_\infty}{\partial n} = -\frac{|\Omega|}{|\partial\Omega|} \text{ on } \partial\Omega \text{ and } \int_{\partial\Omega} u_\infty = 0. \quad (\text{P}(\infty))$$

Define

$$\Sigma_\infty = \int_{\Omega} u_\infty, \quad \text{and} \quad \Sigma_1 = - \int_{\partial\Omega} u_\infty^2, \quad (14.1)$$

with u_∞ satisfying Problem (P(∞)). The terms $\Sigma_\infty > 0$ and $\Sigma_1 \leq 0$ (both obviously independent of β) give coefficients in the next terms in the asymptotic expansion

$$Q(\beta) \sim \frac{\beta |\Omega|^2}{|\partial\Omega|} + \Sigma_\infty + \frac{\Sigma_1}{\beta} \quad \text{for } \beta \rightarrow \infty.$$

Inequality (4.10) of [37] (also (4.12)) states

$$\frac{\beta |\Omega|^2}{|\partial\Omega|} + \Sigma_\infty + \frac{\Sigma_1}{\beta} \leq Q(\beta). \quad (14.2)$$

We now improve on the lower bounds on $Q(\beta)$ given in (1.2) and in (14.2). A reasonable choice for test functions v to insert into $\mathcal{J}(v)$ is

$$v = c_0 + t_0 u_0 + t_\infty u_\infty. \quad (14.3)$$

In substituting the functions as in (14.3) into $\mathcal{J}(v)$ it is useful to note that simple applications of the Divergence Theorem give

$$Q_0 = \int_{\Omega} |\nabla u_0|^2 \quad \text{and} \quad \Sigma_\infty = \int_{\Omega} |\nabla u_\infty|^2.$$

The Divergence Theorem also gives

$$\int_{\Omega} \operatorname{div}(u_0 \nabla u_\infty) = 0 \quad \text{whence} \quad Q_0 = \int_{\Omega} \nabla u_0 \cdot \nabla u_\infty.$$

Now

$$\begin{aligned} \mathcal{A}(v) &= 2c_0|\Omega| + t_0(2-t_0)Q_0 + t_\infty(2-t_\infty)\Sigma_\infty - 2t_0t_\infty \int_{\Omega} \nabla u_0 \cdot \nabla u_\infty, \\ &= 2c_0|\Omega| + t_0(2-t_0-2t_\infty)Q_0 + t_\infty(2-t_\infty)\Sigma_\infty, \\ \mathcal{B}(v) &= c_0^2|\partial\Omega| - t_\infty^2\Sigma_1. \end{aligned}$$

Next we notice that we have quadratics in c_0 and separately in (t_0, t_∞) to maximize. Maximizing over c_0 gives

$$c_{0,\max} = \beta \frac{|\Omega|}{|\partial\Omega|}.$$

Substituting for c_0 and considering the function so formed, quadratic in the t variables, we find that its hessian is negative semidefinite, and that there is a unique maximum, at which $t_0 + t_\infty = 1$. We calculate the maximizing t and denoting them by t_{\max} , then use $c_{0,\max}$ and $t_{0,\max}, t_{\infty,\max}$ to define, by equation (14.3), a function v_{\max} . Then $\mathcal{J}(v_{\max})$ is a rational function of β . Written in a form appropriate for β not close to 0,

$$\begin{aligned} Q(\beta) \geq \mathcal{J}(v_{\max}) &= R\left(\frac{|\Omega|^2}{|\partial\Omega|}, \Sigma_\infty, \Sigma_1, Q_0; \beta\right), \quad (14.4) \\ R\left(\frac{|\Omega|^2}{|\partial\Omega|}, \Sigma_\infty, \Sigma_1, Q_0; \beta\right) &= \left(\beta \frac{|\Omega|^2}{|\partial\Omega|} + \Sigma_\infty + \frac{1}{\beta}\Sigma_1\right) + \frac{\Sigma_1^2}{\beta^2(\Sigma_\infty - Q_0) - \beta\Sigma_1}. \end{aligned}$$

(On using $\Sigma_1 \leq 0$ we see that this is an improvement on the left-hand side of [37] (4.12).) In a form appropriate for β small, the rational function R can be rewritten to give:

$$Q(\beta) \geq \mathcal{J}(v_{\max}) = Q_0 + \beta \frac{|\Omega|^2}{|\partial\Omega|} + \frac{\beta(\Sigma_\infty - Q_0)^2}{\beta(\Sigma_\infty - Q_0) - \Sigma_1}. \quad (14.5)$$

This inequality improves on (4.4) of [37].

The function $\mathcal{J}(v_{\max})$ is increasing in β , a property which it shares with $Q(\beta)$. $\mathcal{J}(v_{\max})$ is concave in β :

$$\frac{\partial^2 \mathcal{J}(v_{\max})}{\partial \beta^2} = \frac{2\Sigma_1(\Sigma_\infty - Q_0)^3}{(\beta(\Sigma_\infty - Q_0) - \Sigma_1)^3}.$$

15 Rectangular domains

15.1 Rectangles: introduction

In this subsection we will consider the rectangle

$$\Omega = (-a, a) \times (b, b) \quad a = rh, \quad b = \frac{h}{r},$$

and usually treat the case $h = 1$ when, of course we have the area of the rectangle as 4. (Outside the context of flows with slip, the corresponding problem for box shapes in R^n can be treated in exactly the same way. See [48].)

We believe the value of our approximation, good lower bound (14.5,14.5), is that it applies to all simply-connected domains. We expect that it will be relatively uncommon for all five functionals involved in the inequality to be explicitly available. A more common situation would be that some would involve computation. In this respect rectangles, besides being a geometry that is relatively common in practice, is reasonably typical.

15.1.1 The functionals Q_0 , Σ_∞ and Σ_1 , etc.

Before presenting the functionals which depend on solving pde problems we note the geometric functionals which we use:

$$|\Omega| = 4ab = 4h^2, \quad |\partial\Omega| = 4(a+b) = 4h\left(r + \frac{1}{r}\right).$$

B defined in [52] is

$$B = 4(a/b + b/a) = 4(r^2 + \frac{1}{r^2}).$$

We use this expression for B below in deriving a very simple lower bound for $Q(\beta)$.

The calculation of Q_0 is an undergraduate exercise in separation of variables: see [45] Problem 133, p74. The quantity Q_0 is available only as an infinite series,

$$Q_0 = \frac{4a^3b}{3} \left(1 - \frac{192}{\pi^5} \frac{a}{b} \sum_{n=1}^{\infty} \frac{\tanh(\pi(2n-1)b/(2a))}{(2n-1)^5} \right).$$

We don't know a closed form for its sum. Also, indicating dependence on the sides as $Q_0(a, b)$, it is 'physically' evident that $Q_0(a, b) = Q_0(b, a)$. Also it is known (by different arguments in [52]) that $Q_0(r, 1/r)$ attains its maximum for a square, $r = 1$. Neither of the last two sentences are immediate from the series. However computation of $Q_0(a, b)$ isn't difficult, just one line of code. The other quantities in the inequality (14.5,14.5) are extremely easy to calculate, and have very simple formulae, just rational functions of the rectangle's side lengths. Once again it is clear that $\mathcal{J}(v_{\max})(a, b)$ is unchanged if we swap a and b as this is true of all five functionals.

There are also series solutions for the case $\beta > 0$ in R^n as well as the R^2 case relevant to flow in ducts: see [48, 19, 16, 70, 38]. The series are very substantially messier than the $\beta = 0$ case, and require about 20 lines of code. Once again $Q_\beta(a, b) - Q_\beta(b, a)$ should be zero and we have found for rectangles with largish aspect ratios and truncated sums there can be noticeable differences. If, however, an approximation is adequate, it is clear that the rational function of β is far easier code. Having said this, nevertheless, the value of our approximation is that it is simple, and general, not merely for rectangular ducts.

Let's start the rectangle calculation. The well-known series solution for Q_0 is easily coded:

Much easier are the calculations associated with Problem (P(∞)). It is easily verified that

$$u_\infty = \frac{ab(a^2 + 6ab + b^2)}{6(a+b)^2} - \frac{bx^2 + ay^2}{2(a+b)},$$

and hence

$$\Sigma_\infty = \frac{8a^3b^3}{3(a+b)^2}.$$

From [40] we note the easy result:

Amongst rectangles with a given area, that which maximizes Σ_∞ is the square. These results (and further terms in the expansion for $Q_r(\beta)$ at large β) are presented in [40]. Also

$$\Sigma_1 = -\frac{4}{45} \frac{a^2b^2(a^4 + 6ba^3 - 10a^2b^2 + 6ab^3 + b^4)}{(a+b)^3}$$

We remark that, both for ellipses and rectangles, when β is moderately large but the aspect ratio a/b is very large, the Σ_1/β term can be relatively large compared to Σ_∞ . Asymptotics at large β require care when the aspect ratio is large.

15.1.2 Comparison of the lower bound with the series

We now compare the lower bound with the elaborate series solution: see Figure 11.




Figure 11: A plot at $a = 5/4$, $b = 4/5$ of the ratio $(S(\beta) - \mathcal{J}(v_{\max}))/\mathcal{J}(v_{\max})$ as a function of β . This tends to 0 both as $\beta \rightarrow 0$ and as $\beta \rightarrow \infty$.

15.2 Using quadratic test functions

As we did with the ellipse, once again choose quadratic test functions. This time we cannot expect useful results for β very small. Indeed, at $\beta = 0$ we obtain just the trivial lower bound of 0. However the results are good for β large.

$$Q_{r,LB} = \frac{4a^2b^2\beta (15a\beta + b^2 + 5ab) (5ab + a^2 + 15b\beta)}{3(30ab^3\beta + 5b^4\beta + 30a^3b\beta + 5a^4\beta + 75a^2b\beta^2 + 75ab^2\beta^2 + 2a^2b^3 + 2b^2a^3)} \quad (15.1)$$

With $b = 1/a$ when one plots both this lower bound and the series solution, the curves are very close for all $\beta > 1$.

Part IV: R approximated for ellipse

By G. Keady

Abstract, Part IV

Consider steady flows in a channel whose cross-section Ω is an ellipse, flows with the Navier slip boundary condition. Denote the volume flow rate by Q . We apply to elliptic cross-sections a recent simple approximation, a rigorous lower bound R on Q , requiring, along with the channel's area and perimeter, the calculation of just the torsional rigidity and two other domain functionals. Here we make a further approximation, R_a , still, however finding a rigorous lower bound for Q . R_a is a good approximation to Q : using it avoids the need for solving the partial differential equation repeatedly for differing values of the slip parameter.

16 Introduction

Some of the papers concerning steady flows under a constant pressure gradient down microchannels are [4, 5, 63, 66, 67] and, for channels with elliptical cross-section [20, 60, 68, 17].

This paper is a sequel to [39] concerning approximating the steady slip flow down a channel with a given cross-section, Ω . As noted in the abstract, just a few parameters are needed, and in cases tested to date, the approximation is remarkably accurate: see [39] for rectangles and [35, 36] for regular polygons and isocles triangles. For the preceding shapes the ‘two other domain functionals’, denoted below by Σ_∞ and Σ_1 , had simple exact formulae. For the ellipse this is no longer the case (as the exact formulae are infinite series with no obvious closed form), and this paper is a check that further approximation still leads to adequate results.

As in previous papers [39, 37, 35, 36] the slip-parameter is denoted by β . The scaled velocity is denoted by $u(\beta)$; the boundary-value problem for $u(\beta)$ is given in Problem (P(β)) in [39] and in an earlier paper [37]. The volume flow rate is defined by

$$Q(\beta) = \int_{\Omega} u(\beta). \quad (16.1)$$

For sufficiently smooth functions v , define, as in [39], and [36], [37], with $\partial\Omega$ denoting the boundary of Ω , \mathcal{A} , \mathcal{B} and \mathcal{J} by

$$\mathcal{A}(v) = \int_{\Omega} (2v - |\nabla v|^2), \quad \mathcal{B}(v) = \int_{\partial\Omega} v^2, \quad \mathcal{J}(v) = \mathcal{A}(v) - \frac{1}{\beta}\mathcal{B}(v). \quad (16.2)$$

The maximiser of $\mathcal{J}(v)$ over all v is $u(\beta)$ and

$$Q(\beta) = \mathcal{J}(u(\beta)) = \int_{\Omega} |\nabla u(\beta)|^2 - \frac{1}{\beta}\mathcal{B}(u(\beta)).$$

This variational characterisation of solutions $u(\beta)$ led to the lower bound to and approximation of $Q(\beta)$ by $R(\beta)$ defined, as in [39], by

$$R(\beta) = Q_0 + \beta \frac{A^2}{P} + \frac{\beta(\Sigma_{\infty} - Q_0)^2}{\beta(\Sigma_{\infty} - Q_0) - \Sigma_1}. \quad (16.3)$$

The arguments of R depending on the domain are: A the area of Ω , P its perimeter, Q_0 the torsional rigidity, and two other quantities Σ_1 and Σ_{∞} . The Σ quantities were defined in [37]: the definition is repeated in [39] and here in §16.2. (Another use of the approximation is that, from an experiment in which the flow rate Q is measured, the slip parameter can be estimated by simply solving, for β , the polynomial equation $R(\beta)$ equals the measured flow Q .)

For ellipses of given area A and eccentricity e , the quantity Q_0 is elementary: see equation (17.8). The other quantities P and approximations $\Sigma_{a,\infty}$ and $\Sigma_{a,1}$ are given in terms of elliptic integrals at equation (17.2) and (on using (17.4), (17.10)) equation (17.11) and (on using (17.5), (17.12)) equation (17.13). We emphasise that the elliptical microchannels have been studied extensively. There is no claim that the numerical results from this paper are better than earlier ones. What is new is the uniform treatment of potentially all cross-sections by the approximation given in R , adding elliptical cross-sections to the list where this has been done.

16.1 The torsion pde, Problem (P(0))

The elastic torsion problem is to find a function u_0 satisfying

$$-\frac{\partial^2 u_0}{\partial x^2} - \frac{\partial^2 u_0}{\partial y^2} = 1 \text{ in } \Omega, \quad u_0 = 0 \text{ on } \partial\Omega. \quad (\text{P}(0))$$

Channel flows with no slip have $\beta = 0$. The *torsional rigidity* of Ω (volume flow in the fluids context) is,

$$Q_0 := \int_{\Omega} u_0. \quad (16.4)$$

With \mathcal{A} defined as in equation (16.2) the solution u_0 maximizes \mathcal{A} over functions vanishing on the boundary $\partial\Omega$. Also $\mathcal{A}(u_0) = Q_0$.

Define also

$$Q_1 := \int_{\partial\Omega} \left(\frac{\partial u_0}{\partial n} \right)^2. \quad (16.5)$$

For small β , as noted in equation (4.5) of [37],

$$Q(\beta) \sim Q_0 + \beta Q_1 \quad \text{as } \beta \rightarrow 0.$$

This suggested the following inequality (proved in [36]) which is used in §17.4 as a check on the quantities occurring in our approximation:

$$\frac{A^2}{P} - \frac{(\Sigma_{\infty} - Q_0)^2}{\Sigma_1} \leq Q_1. \quad (16.6)$$

16.2 Problem (P(∞))

Problem (P(∞)) was defined in [39] and, before that in [37]. Our notation here is as before. The function u_{∞} solves

$$-\frac{\partial^2 u_{\infty}}{\partial x^2} - \frac{\partial^2 u_{\infty}}{\partial y^2} = 1 \text{ in } \Omega, \quad \frac{\partial u_{\infty}}{\partial n} = -\frac{|\Omega|}{|\partial\Omega|} \text{ on } \partial\Omega \text{ and } \int_{\partial\Omega} u_{\infty} = 0. \quad (\text{P}(\infty))$$

Define, as in [39], [37]

$$\Sigma_{\infty} = \int_{\Omega} u_{\infty}, \quad \text{and } \Sigma_1 = - \int_{\partial\Omega} u_{\infty}^2, \quad (16.7)$$

with u_{∞} satisfying Problem (P(∞)).

Once again there is a variational characterisation of the solutions. This time u_{∞} is the maximizer of $\mathcal{A}(v)$ as one varies over functions v for which the integral of v over the boundary $\partial\Omega$ is zero. The inequality $\Sigma_{\infty} \geq Q_0$ is an immediate consequence of the variational approach. (See also [37] equation (4.9).) Also $\mathcal{A}(u_{\infty}) = \Sigma_{\infty}$.

As noted in [37], for large β ,

$$Q(\beta) \sim \frac{\beta A^2}{P} + \Sigma_{\infty} + \frac{\Sigma_1}{\beta} \quad \text{for } \beta \rightarrow \infty.$$

17 The ellipse

Consider ellipses with area $A = \pi$, boundary specified, with $a \geq 1$, by

$$\frac{x^2}{a^2} + a^2 y^2 = 1, \quad A = \pi. \quad (17.1)$$

The eccentricity e of the ellipse satisfies $e^2 = 1 - 1/a^4$.

The elliptic integral EllipticE is denoted by EllipticE : EllipticK is denoted by EllipticK . The perimeter of the ellipse, P , is

$$P = 4a\text{EllipticE}(e) \quad \text{for } a \geq 1. \quad (17.2)$$

The approximations to the Σ quantities (and Q_1) are readily expressed in terms of boundary moments, these being defined by

$$i_{2k} = \int_{\partial\Omega} (x^2 + y^2)^k. \quad (17.3)$$

Evaluating these for the ellipse gives

$$i_2 = \frac{4}{3a} ((1 + a^4) \text{EllipticE}(e) + \text{EllipticK}(e)), \quad (17.4)$$

$$i_4 = \frac{4}{5a^3} ((1 - a^4 + a^8) \text{EllipticE}(e) + 2(1 + a^4) \text{EllipticK}(e)) \quad (17.5)$$

A well-known special case is the circular disk of radius 1 where $u_{\odot}(\beta) = (1 - x^2 - y^2)/4 + \beta/2$ and $Q_{\odot}(\beta) = \pi(1 + 4\beta)/8$. It is also known that for any ellipse with area π , $Q(\beta) \leq Q_{\odot}(\beta)$.

17.1 Q_0 and Q_1

We have

$$u_0 = \frac{1 - \frac{x^2}{a^2} - a^2 y^2}{2(\frac{1}{a^2} + a^2)}, \quad (17.6)$$

$$\frac{\partial u_0}{\partial n} = - \frac{\sqrt{(\frac{1}{a^2} + a^2) - (x^2 + y^2)}}{\frac{1}{a^2} + a^2} \quad \text{on } \partial\Omega. \quad (17.7)$$

and hence

$$Q_0 = \frac{\pi a^4}{4(1+a^4)}, \quad (17.8)$$

$$\begin{aligned} Q_1 &= \frac{a^2 P}{1+a^4} - \frac{a^4 i_2}{(1+a^4)^2}, \\ &= \frac{4}{3} \frac{a^3}{(1+a^4)^2} (2(1+a^4)\text{EllipticE}(e) - \text{EllipticK}(e)) \quad \text{for } a \geq 1 \end{aligned} \quad (17.9)$$

17.2 Approximations $\Sigma_{a,\infty}$ and $\Sigma_{a,1}$

The quantities Σ_∞ and Σ_1 can be approximated by terms denoted below by $\Sigma_{a,\infty}$ and $\Sigma_{a,1}$ which, as mentioned above, are calculated in terms of boundary moments.

We seek an approximation to u_∞ by using test functions of the form

$$v = u_0 + t \left(x^2 - y^2 - \frac{C_1}{P} \right),$$

where the constant C_1 is chosen so the boundary integral of v is zero, i.e.

$$C_1 = \int_{\partial\Omega} (x^2 - y^2) = \frac{-2a^2 P + (1+a^4)i_2}{(-1+a^4)}. \quad (17.10)$$

The quantity $\mathcal{A}(v)$ is quadratic in t

$$\mathcal{A}(v) = -\frac{\pi (4(1+a^4)^2 t^2 - 2(1+a^4)(-1+a^4 - 4a^2 \frac{C_1}{P})t - a^4)}{4a^2(1+a^4)}.$$

The maximum of this quadratic occurs at $t = t_*$ with

$$t_* = \frac{-1 + a^4 - 4a^2 \frac{C_1}{P}}{4(1+a^4)}.$$

Then, at t_* ,

$$\mathcal{A}(v_*) = \frac{\pi (a^4 + \frac{1}{4}(-1 + a^4 - 4a^2 \frac{C_1}{P})^2)}{4a^2(1+a^4)}. \quad (17.11)$$

Define the approximation to Σ_∞ by

$$\Sigma_{a,\infty} = \int_{\Omega} v_*,$$

and then, after performing the integrals, note the remarkable fact that

$$\Sigma_{a,\infty} = \mathcal{A}(v_*) = \int_{\Omega} |\nabla v_*|^2.$$

(The second equality follows immediately from the first. The first equality is checked by performing the integrals. We note it as ‘remarkable’ because, although we can prove using applications of the Divergence Theorem that $\Sigma_{\infty} = \mathcal{A}(u_{\infty})$, it isn’t obvious there would be a parallel result for v_* .) As $\mathcal{A}(u_{\infty}) > \mathcal{A}(v_*)$, our approximation underestimates Σ_{∞} , i.e. $\Sigma_{\infty} > \Sigma_{a,\infty}$.

Higher boundary moments enter in the approximation to Σ_1 . We have

$$\int_{\partial\Omega} v^2 = t^2 \left(C_2 - \frac{C_1^2}{P} \right).$$

Substituting $t = t_*$ gives, with

$$C_2 = \frac{4a^4P - 4a^2(1+a^4)i_2 + (1+a^4)^2i_4}{(-1+a^4)^2}, \quad (17.12)$$

$$\Sigma_{a,1} = - \int_{\partial\Omega} v_*^2 = - \frac{(-4a^2\frac{C_1}{P} + a^4 - 1)^2 \left(C_2 - \frac{C_1^2}{P} \right)}{16(a^4 + 1)^2}. \quad (17.13)$$

We have checked that the approximations $\Sigma_{a,\infty}(a)$, $\Sigma_{a,1}(a)$ when used in the obvious places in inequality (16.6) do satisfy the inequality.

17.3 Comparison of $R_a(\beta)$ with $Q(\beta)$

Write $R_a(\beta)$ as the expression obtained when one uses $\Sigma_{a,\infty}$ and $\Sigma_{a,1}$ in the obvious places in the parameter list of R . The numeric values for $Q(\beta)$ used in Figure 12 have been obtained by several methods. The PDE Toolbox of Matlab is one source (and could be, and has been, used with other cross-sections). The variational principle used for the theory in [36] has been used for numerical values for $Q(\beta)$ for ellipses in [68]. Fourier series for $u(\beta)$ is used in [20] with, however, a further approximation: that approximation is avoided in numerics performed by BW reported in Part I.




Figure 12: For an ellipse with $a = 5/4$, the quantity $R_a(\beta) - \beta A^2/P$ plotted against β . The plot goes between Q_0 and $\Sigma_{a,\infty}$. The solid dots are values when one replaces R_a with Q calculated numerically.

17.4 Further remarks

17.4.1 Approximation by quadratics for $\beta \geq 0$

The numeric values of the approximation coincide with those of the quadratic approximation of Part I.

There we used test functions of the form

$$v_{\text{quad}} = c + c_{xx}x^2 + c_{yy}y^2,$$

in maximizing $\mathcal{J}(v)$, \mathcal{J} defined at equations (16.2). This was completely ad hoc: it was suggested by the contours ‘looking elliptical’ e.g. in [60]. Write $v_{\text{quad,max}}$ for the maximiser. A calculation shows

$$\mathcal{J}v_{\text{quad,max}} = R_a(\beta).$$

Thus our approximate $R_a(\beta)$, as with the $R(\beta)$ involving the true Σ values, is a lower bound for $Q(\beta)$.

17.4.2 Higher order approximation for u_∞

We suspect that finding higher approximations to u_∞ and the Σ quantities is likely to be unimportant for the ellipse, but they could be found. One method begins by defining an inner product

$$\langle \phi, \psi \rangle_{1\Omega} = \int_{\Omega} \nabla \phi \cdot \nabla \psi,$$

and noting that $\langle u_0, h \rangle_{1\Omega} = 0$ for any harmonic function. Suppose one has a set of N orthogonal harmonic functions h_j each of which has boundary integral zero. Use test functions v of the form

$$v = u_0 + \sum t_j h_j,$$

in the variational characterisation. It is easy to maximize $\mathcal{A}(v)$ over the t , and denoting the maximizers with a subscript star

$$t_{j*} = \frac{\int_{\Omega} h_j}{\langle h_j, h_j \rangle_{1\Omega}}.$$

Return now to the ellipse. Take as the h_j

$$h_j = \operatorname{Re} \left(T_{2j} \left(\frac{x + iy}{ae} \right) \right) + C_{1,j}, \quad (17.14)$$

where T_{2j} denote Chebyshev polynomials and the $C_{1,j}$ are chosen so that the boundary integrals are zero. The h_j are orthogonal. Also

$$\int_{\Omega} h_1 = -\frac{\pi}{2} + A C_{1,1}, \quad \int_{\Omega} h_j = A C_{1,j} \text{ for } j > 1,$$

so that the integrals defining the t_{j*} can be evaluated.

The series one gets approximating u_{∞} using this variational method is the same as that found using separation of variables in elliptic coordinates and the Fourier series which follows from that.

18 Conclusion and future prospects

The approximation $R(\beta)$ presented in [39] for the flow $Q(\beta)$ has been determined to be good for elliptical cross-sections.

For any domain it is easy to use finite elements or other numerical methods to find numerical approximations to the five parameters occurring in R . However, having a few cross-sections where some or all of the parameters are available analytically is useful in checking numerics.

The ellipse now joins the list initiated in [39] for rectangles, and continued in [35, 36] for shapes like regular polygons, triangles, rhombi, a list where all the parameters of R are known, or estimated. For many years there have been

tables for $Q(0)$, the torsional rigidity, one of the parameters in R . Looking to the future, a similar table – online lookup – of all five of the parameters would enable experimenters to calculate quickly an approximation to the flow for any value of β .

Outline of appendices

The notes in the appendices were prepared for a student of BW further developing knowledge of u_∞ for the ellipse. This also influenced how GK wrote papers, in order to leave a clear plan for where the student's paper might be submitted.

A Quadratic test functions, general β

B ($P(\infty)$), more terms: agreement with Fourier Series

B.1-B.3 General Ω , including basics of the variational principle

B.4-B.5 More on polynomial test functions

C Conformal maps, ellipses

D More recurrences, moments

There are very many open questions concerning u_∞ for the ellipse.

- We believe series expansions for small eccentricity should be found.
- For eccentricities tending to 1, matched asymptotic expansions might be useful.
- It might be possible to make use of the known conformal map of the ellipse to unit disk.
- Recurrences for t_{j*} , or V_j of Part II might be found.
The geometric quantities I_{2k} and i_{2k} may have other uses and the recurrences explored in a later Appendix may have interest. (GK has more results not presented in this report.)

A Agreement with quadratic approximation

In Part I we used test functions of the form

$$v_{\text{quad}} = c + c_{xx}x^2 + c_{yy}y^2,$$

in maximizing $\mathcal{J}(v)$, \mathcal{J} defined at equations (16.2). The values of the c such that $\mathcal{J}(v_{\text{quad}})$ is maximized were found. This was completely ad hoc: it was

suggested by the contours ‘looking elliptical’ e.g. in [60]. Write $v_{\text{quad,max}}$ for the maximiser. We did this for any geometry, but it probably only works well, at least for small β , for the ellipse. For the ellipse the relations between the moments (recorded at D.1, ensure that the c come out so that the Laplacian of $v_{\text{quad,max}}$ is minus one. Thus, relating the two approaches, we have, for some c and τ depending on β :

$$v_{\text{quad,max}} = c + \tau u_0 + (1 - \tau)u_{a,\infty}.$$

Next, if we maximise \mathcal{J} over functions as on the right of the preceding equation, maximising over (c, τ) one finds a maximiser v_* and

$$\mathcal{J}v_* = R_a(\beta).$$

Thus our approximate $R_a(\beta)$, like the $R(\beta)$ involving the true Σ values, is a lower bound for $Q(\beta)$.

B More terms, agreement with Fourier Series

B.1 Introduction, and $(\mathbf{P}(\infty))$ for general Ω

One of the many questions yet to be treated well is how close are GK’s approximations $\Sigma_{a,\infty}$ and $\Sigma_{a,1}$ to the true values of Σ_∞ (which we know is bigger than $\Sigma_{a,\infty}$) and of Σ_1 ?

It isn’t clear how best to represent higher approximations. The methods include

- Fourier series which is equivalent to the variational characterization of u_∞ using Chebyshev polynomials, and
- the variational characterization of u_∞ using harmonic polynomials, thus the next term after that used in R_a might be approached as

$$v = u_0 + \tilde{t}_1 \left(x^2 - y^2 - \frac{\int_{\partial\Omega}(x^2 - y^2)}{P} \right) + \tilde{t}_2 \left(x^4 - 6x^2y^2 + y^4 - \frac{\int_{\partial\Omega}(x^4 - 6x^2y^2 + y^4)}{P} \right).$$

An outline of how this might be done is given, in the context of general domains, below in §B.3.

Of course, when there are several methods, one must cross-check.

B.2 The variational principle for $(\mathbf{P}(\infty))$

The basics of the variational principle (somewhat formally, as in the usual undergraduate treatment of Euler-Lagrange d.e.) is as follows.

The standard approach to variational methods for applied mathematicians (as opposed to more pure-maths approaches involving Frechet derivatives) starts with a linearization, ϵ small

$$\mathcal{A}(u_\infty + \epsilon\phi) - \mathcal{A}(u_\infty) \sim 2\epsilon\mathcal{I}(u_\infty, \phi) \quad \text{where} \quad \mathcal{I}(u_\infty, \phi) = \int_{\Omega} (\phi - \nabla u_\infty \cdot \nabla \phi),$$

with

$$\int_{\partial\Omega} \phi = 0 \quad \text{and} \quad \int_{\partial\Omega} u_\infty = 0.$$

(The boundary integrals have to be zero in order that the functions are allowed in the variational competition.)

For u_∞ to be the maximum, we must have the $O(\epsilon)$ term, \mathcal{I} , is 0. There are two parts to showing that u_∞ satisfies the pde Problem $(\mathbf{P}(\infty))$, namely (i) the pde (Laplacian is -1) and (ii) the boundary condition that the normal derivative is constant.

(i)

$$\operatorname{div}(\phi\nabla u_\infty) = \phi\Delta u_\infty + \nabla u_\infty \cdot \nabla \phi,$$

so

$$0 = \mathcal{I}(u_\infty, \phi) = \int_{\Omega} (\phi(1 + \Delta u_\infty) - \operatorname{div}(\phi\nabla u_\infty)). \quad (\text{B.1})$$

Now consider smooth functions ϕ which vanish on the boundary. We have $\int_{\Omega} (\phi(1 + \Delta u_\infty)) = 0$ for all such ϕ so we must have $-\Delta u = 1$. (For suppose not and $(1 + \Delta u_\infty) > 0$ in some domain. Then choose a nonzero function $\phi \geq 0$ which has its support in that domain. The contradiction is clear.)

(ii) Now consider maximisers satisfying $-\Delta u = 1$, so

$$\int_{\partial\Omega} \frac{\partial u_\infty}{\partial n} = -A, \quad \text{and as before} \quad \int_{\partial\Omega} \phi = 0.$$

From equation (B.1)

$$0 = \int_{\Omega} \operatorname{div}(\phi\nabla u_\infty) = \int_{\partial\Omega} \phi \frac{\partial u_\infty}{\partial n} = \int_{\partial\Omega} \phi \left(\frac{\partial u_\infty}{\partial n} + \frac{A}{P} \right),$$

as the integral of ϕ around the boundary is 0. Now

$$\int_{\partial\Omega} \left(\frac{\partial u_\infty}{\partial n} + \frac{A}{P} \right) = 0,$$

so if we take

$$\phi = \left(-\frac{\nabla u_\infty \cdot \nabla u_0}{|\nabla u_0|} + \frac{A}{P} \right)$$

which is a function whose boundary integral is 0, we must then have

$$\int_{\partial\Omega} \left(\frac{\partial u_\infty}{\partial n} + \frac{A}{P} \right)^2 = 0,$$

and hence the integrand 0, which establishes the boundary condition of Problem (P(∞)).

B.3 Better approximation, general Ω

Define an inner product

$$\langle f_1, f_2 \rangle_{1,\Omega} = \int_{\Omega} \nabla f_1 \cdot \nabla f_2.$$

For any harmonic function h , using the divergence theorem on $\text{div}(u_0 \nabla h)$, one finds

$$\langle u_0, h \rangle_{1,\Omega} = 0.$$

Given any sequence of harmonic functions \tilde{h}_k , for each of which the boundary integral is zero, one can apply Gram-Schmidt to find an orthogonal sequence of harmonic functions h_k :

$$h_k = \tilde{h}_k - \sum_{j=1}^{k-1} \frac{\langle \tilde{h}_k, h_j \rangle_{1,\Omega}}{\langle h_j, h_j \rangle_{1,\Omega}} h_j.$$

If necessary change the sign of h_k so that $\int_{\Omega} h_k > 0$. The boundary integrals of the h_k are also zero.

The function

$$v = u_0 + t_1 h_1 + t_2 h_2 + \dots,$$

is suitable as a test function in the problem of maximizing $\mathcal{A}(v)$. Now

$$\begin{aligned}\mathcal{A}(v) &= 2 \int_{\Omega} v - \langle v, v \rangle_{1,\Omega}, \\ &= 2 \int_{\Omega} v - \langle u_0, u_0 \rangle_{1,\Omega} - \sum_{j=1}^N t_j^2 \langle h_j, h_j \rangle_{1,\Omega},\end{aligned}$$

on using the orthogonality properties we have noted above. Thus

$$\mathcal{A}(v) = Q_0 - \sum_{j=1}^N \left(t_j^2 \langle h_j, h_j \rangle_{1,\Omega} - 2 t_j \int_{\Omega} h_j \right).$$

There are no ‘cross terms’ $t_j t_k$ with $j \neq k$. It is easy to maximize over the t , and denoting the maximizers with a subscript star

$$t_{j*} = \frac{\int_{\Omega} h_j}{\langle h_j, h_j \rangle_{1,\Omega}}.$$

We expect the N -th approximation to be

$$\Sigma_{\infty,N} = \mathcal{A}(v_{*N}) = Q_0 + \sum_{j=1}^N t_{j*} \int_{\Omega} h_j.$$

B.4 Harmonic polynomials for higher approximations via the variational principle?

We now suggest how the general set-up described in §B.3 might be applied to the case of the ellipse.

Without the orthogonality described in §B.3 the approximations would be in the form

$$u_N = u_0 + \sum_{j=1}^N c_j(N) \hat{h}_j(x, y),$$

with the \hat{h}_j appropriate harmonic functions with boundary integrals being 0. The $c_j(N)$ are then chosen so that \mathcal{A} is maximized. Amongst the possibilities for the \hat{h}_j are the following

- $\hat{h}_j = \operatorname{Re}(x + iy)^{2j} - c_j$ with the c_j chosen so the boundary integral of h_j is 0.

- $\hat{h}_j = c \cosh(2j\eta) \cos(2j\psi) - \hat{c}_j$ as before with $c^2 = a^2 - 1/a^2$, and again with \hat{c}_j such that the boundary integral of \hat{h}_j is zero. This is equivalent to the Chebyshev polynomials treated in the next subsection.

We discovered in §B.3 that it will be possible to have a sequence h_j which has the $c_j(N)$ independent of N . The Chebyshev polynomials of the next subsection provide such a sequence.

B.5 Chebyshev polynomials with complex argument

B.5.1 The polynomials

We begin with a few facts. As before $c = \sqrt{a^2 - 1/a^2}$. Then

$$\int_{\Omega} T_2\left(\frac{x+iy}{c}\right) = -\frac{\pi}{2}, \quad \int_{\Omega} T_{2j}\left(\frac{x+iy}{c}\right) = 0 \text{ for } j > 1.$$

Also

$$\frac{dT_n\left(\frac{z}{c}\right)}{dz} = \frac{n}{c} U_{n-1}\left(\frac{z}{c}\right).$$

Refer now to equation (3.1) of Part I concerning elliptic coordinates.

$$T_n(\cosh(\eta) \cos(\psi) + i \sinh(\eta) \sin(\psi)) = T_n(\cosh(\eta + i\psi)) = \cosh(n(\eta + i\psi)),$$

so

$$\operatorname{Re}(T_n(\cosh(\eta) \cos(\psi) + i \sinh(\eta) \sin(\psi))) = \cosh(n\eta) \cos(n\psi).$$

B.5.2 Integrals over Ω

Let $z = x + iy$. Since

$$z^m \bar{z}^n - \bar{z}^m z^n = |z|^n (z^{m-n} - \bar{z}^{m-n}),$$

it is imaginary: thus we have $\Re((x + iy)^m (x - iy)^n)$ is even in y . Similarly

$$z^m \bar{z}^n + \bar{z}^m z^n = |z|^n (z^{m-n} + \bar{z}^{m-n}),$$

it is real: thus we have $\Im((x + iy)^m (x - iy)^n)$ is odd in y . Thus

$$\int_{\Omega} U_m\left(\frac{z}{c}\right) \bar{z} \text{ is real,}$$

and hence

$$\int_{\Omega} U_m\left(\frac{z}{c}\right) U_n\left(\frac{\bar{z}}{c}\right) \text{ is real.}$$

The value of the integral is, with $r = a + 1/a$, is given by

$$\int_{\Omega} U_m\left(\frac{x+iy}{c}\right) U_n\left(\frac{x-iy}{c}\right) = \frac{\pi c^2}{4(1+n)} \left(\left(\frac{r}{c}\right)^{2n+2} - \left(\frac{c}{r}\right)^{2n+2} \right) \delta_{n,m}. \quad (\text{B.2})$$

(See [3] equation (4.25).) When $m = n$ we can write the right-hand side of (B.2) as follows. First

$$\frac{r^2}{c^2} = \frac{a + 1/a}{a - 1/a} = \exp(2\eta_0),$$

on using $a = c \cosh(\eta_0)$, $1/a = c \sinh(\eta_0)$. Hence

$$\begin{aligned} \int_{\Omega} \left| U_n\left(\frac{x+iy}{c}\right) \right|^2 &= \frac{\pi c^2}{4(1+n)} \left(\left(\frac{r}{c}\right)^{2n+2} - \left(\frac{c}{r}\right)^{2n+2} \right) \\ &= \frac{\pi c^2}{2(1+n)} \sinh(2(n+1)\eta_0), \\ &= \frac{\pi}{\sinh(2\eta_0)(1+n)} \sinh(2(n+1)\eta_0) \end{aligned} \quad (\text{B.3})$$

We now establish the orthogonality of the h_j of equation (17.14). Now

$$\begin{aligned} \langle h_j, h_k \rangle_{1,\Omega} &= \int_{\Omega} \nabla h_j \cdot \nabla h_k, \\ &= \int_{\Omega} \nabla \left(\operatorname{Re}(T_{2j}\left(\frac{x+iy}{c}\right)) \right) \cdot \nabla \left(\operatorname{Re}(T_{2k}\left(\frac{x+iy}{c}\right)) \right). \end{aligned} \quad (\text{B.4})$$

Next, for holomorphic functions f_j and f_k ,

$$\begin{aligned} \operatorname{Re} \int_{\Omega} f_j(z) \overline{f_k(z)} &= \operatorname{Re} \int_{\Omega} (f_{jr} + i f_{ji})(f_{kr} - i f_{ki}), \\ &= \int_{\Omega} (f_{jr} f_{kr} + f_{ji} f_{ki}), \end{aligned}$$

where the r and i in the subscripts indicate real and imaginary parts respectively. Applying this to derivatives and using the Cauchy-Riemann equations gives

$$\operatorname{Re} \int_{\Omega} \frac{df_j}{dz} \overline{\frac{df_k}{dz}} = \int_{\Omega} \nabla f_{jr} \cdot \nabla f_{kr}.$$

Using this with the Chebyshev T functions in equation (B.4) gives

$$\begin{aligned}
\langle h_j, h_k \rangle_{1,\Omega} &= \int_{\Omega} \nabla \left(\operatorname{Re}(T_{2j}(\frac{x+iy}{c})) \right) \cdot \nabla \left(\operatorname{Re}(T_{2k}(\frac{x+iy}{c})) \right), \\
&= \operatorname{Re} \int_{\Omega} \frac{dT_{2j}(\frac{x+iy}{c})}{dz} \overline{\left(\frac{dT_{2k}(\frac{x+iy}{c})}{dz} \right)}, \\
&= \frac{4jk}{c^2} \operatorname{Re} \int_{\Omega} \left(U_{2j-1}(\frac{x+iy}{c}) \right) \overline{\left(U_{2k-1}(\frac{x-iy}{c}) \right)}.
\end{aligned}$$

This final expression is zero for $j \neq k$ and when $j = k$ its value is given as in equation (B.2).

$$\langle h_j, h_j \rangle_{1,\Omega} = \pi j \sinh(4j \eta_0) \quad (\text{B.5})$$

B.6 Integrals over $\partial\Omega$

We have

$$\begin{aligned}
L C_{1,j} &= - \int_{\partial\Omega} \operatorname{Re}(T_{2j}(\frac{x+iy}{c})) ds, \\
&= -2 \int_0^\pi \operatorname{Re}(T_{2j}(\cosh(\eta_0) \cos(\psi) + i \sinh(\eta_0) \sin(\psi))) \sqrt{a^2 \sin(\psi)^2 + \frac{\cos(\psi)^2}{a^2}} d\psi, \\
&= -2a \cosh(2j\eta_0) \int_0^\pi \cos(2j\psi) \sqrt{1 - e^2 \cos(\psi)^2} d\psi, \\
&= -\pi a \cosh(2j\eta_0) \hat{g}_j.
\end{aligned} \quad (\text{B.6})$$

From equations (B.6) and (B.5) one finds t_{j*} and finds that for $j > 1$ it agrees with the Fourier series calculation at (12.11).

C Conformal maps, ellipses

We remark that the normal to $\partial\Omega$ is

- in the direction of ∇u_0 ,
- in the direction of $\log(|f(z)|)$ where f is the conformal map from Ω to the unit disk in which the origin is mapped to the origin.
The function $\log(|f(z)/z|)$ is harmonic in Ω .

We remark that there is a ‘Poisson Integral Formula’ for solving the Dirichlet problem for Laplace’s equation in an ellipse. This follows from the conformal map between ellipse and disk: see [42], p177 and [61]. See also [54, 50, 46, 55].

D More recurrences

D.1 Geometric quantities: moments

The quantities Σ_∞ and Σ_1 can be expressed in terms of boundary moments – moments about the incentre – defined by

$$\begin{aligned} i(m, n) &= \int_{\partial\Omega} (x^{2m} y^{2n}) & (D.1) \\ i_{2k} &= i(k, k) = \int_{\partial\Omega} (x^2 + y^2)^k. \end{aligned}$$

We remark that the Cauchy-Schwarz inequality for the integrals implies that $i_4 \geq i_2^2/P$.

We now record results needed in the approximation leading to v_{max} in §17.2.

- For second moments we have the following:

$$i(1, 0) + i(0, 1) = i_2, \quad \frac{i(1, 0)}{a^2} + a^2 i(0, 1) = P.$$

From this

$$i(1, 0) = \frac{(a^2(a^2 i_2 - P))}{(-1 + a^4)}, \quad i(0, 1) = \frac{(-i_2 + a^2 P)}{(-1 + a^4)},$$

and

$$C_1 = i(1, 0) - i(0, 1) = \frac{-2a^2 P + (1 + a^4) i_2}{(-1 + a^4)}.$$

- For fourth moments we have the following:

$$\begin{aligned} i(2, 0) + 2i(1, 1) + i(0, 2) &= i_4, \\ \frac{i(2, 0)}{a^2} + (a^2 + \frac{1}{a^2})i(1, 1) + a^2 i(0, 2) &= i_2, \\ \frac{i(2, 0)}{a^4} + 2i(1, 1) + a^4 i(0, 2) &= P. \end{aligned}$$

A quantity occurring in the approximation $\Sigma_{a,1}$ is

$$C_2 = i(2, 0) - 2i(1, 1) + i(0, 2) = \frac{2a^4P - 4a^2(1 + a^4)i_2 + (1 + a^4)^2i_4}{(-1 + a^4)^2}.$$

The Cauchy-Schwarz inequality gives $C_2 \geq C_1^2/P$.

D.2 Further work to be done

The further work will need more of the boundary moments i_{2k} . We expect that there is a 3-term recurrence relation.

If (as we expect) all the i_{2k} can be written in the form

$$i_{2k} = r_K(k, a)\text{EllipticK}(e) + r_E(k, a)\text{EllipticE}(e),$$

one can eliminate the EllipticK and EllipticE from pairs of such equations to get the 3-term equation

$$i_{2k} = f_+(k, a)i_{2(k+1)} + f_-(k, a)i_{2(k-1)}.$$

It remains to find the functions f , i.e. the recurrence.

It may also be possible to get to the recurrence from the integrals defining i_{2k} as done for g_k and \hat{g}_k .

The ellipse is given parametrically by

$$x = a \cos(\psi), \quad y = \frac{1}{a} \sin(\psi).$$

Then

$$r^2 = x^2 + y^2 = \frac{1}{a^2} + (a^2 - \frac{1}{a^2}) \cos^2(\psi) = a^2 \left(\frac{1}{a^4} + e^2 \cos^2(\psi) \right),$$

and

$$\left(\frac{ds}{d\psi} \right)^2 = a^2 (1 - e^2 \cos^2(\psi)).$$

So,

$$\frac{ds}{d\psi} = a \hat{g}(\psi),$$

so

$$i_{2k} = 4a^{2k+1} \int_0^{\pi/2} \left(\frac{1}{a^4} + e^2 \cos^2(\psi) \right)^k \hat{g}(\psi).$$

Setting $u = \cos(\psi)^2$

$$j(m,n) = \int_0^1 u^m (1-u)^n \sqrt{\frac{1-e^2u}{u(1-u)}} du.$$

It is trivial that

$$j(m+1, n) + j(m, n+1) = j(m, n). \quad (\text{D.3})$$

Each of the $j(m, n)$ has the form

$$j(m, n) = f_K(m, n, e^2)\text{EllipticK}(e) + f_E(m, n, e^2)\text{EllipticE}(e),$$

for rational functions f_E and f_K . Defining

$$u_m = j(m, 0), \quad v_n = j(0, n),$$

one can find the first couple of terms and recurrences to determine the rest. This enables one to determine the entries at the ends of the rows in the triangle below

$$\begin{array}{ccccccc} & & & & j(0, 0) & & \\ & & & & & & \\ & & & & j(1, 0) & & j(0, 1) \\ & & & & j(2, 0) & & j(1, 1) & & j(0, 2) \\ & & & & j(3, 0) & & j(2, 1) & & j(1, 2) & & j(0, 3) \end{array}$$

From the entries at the ends of the rows one can use equation (D.3) rearranged to determine all the other entries.

The recurrences for the u_n and v_n are as follows.

$$(2m-1)u_{m-1} - ((2m+2)e^2 + 2m)u_m + (2m+3)e^2u_{m+1} = 0,$$

$$-(2m-1)(1-e^2)v_{m-1} - 2((2m+1)e^2 - m)v_m + (2m+3)e^2v_{m+1} = 0.$$

There are various identities that might be used to check, for example (just from $\sin^2 + \cos^2 = 1$)

$$v_m = \sum_{j=0}^m (-1)^j \text{Binomial}(m, j) u_j.$$

References

- [1] P. Abbott, On the perimeter of an ellipse, *The Mathematica Journal*, **11**(2), (2009).
- [2] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions*, (Dover: 1965).
- [3] G. Akemann, T. Nagao, I. Parra, and G. Vernizzi, Gegenbauer and other planar orthogonal polynomials on an ellipse in the complex plane. arXiv (2019).
<http://arxiv.org/1905.02397>
- [4] M. Bahrami, M. Yovanovich and J. Culham, Pressure drop of fully- developed, laminar flow in microchannels of arbitrary cross-section, *Journal of Fluids Engineering*, **128**(5), (2006) pp. 1036–1044.
doi:10.1115/1.2234786
- [5] M. Bahrami, A. Tamayol and P. Taheri, Slip-flow pressure drop in microchannels of general cross section, *Journal of Fluids Engineering*, **131**(3), (2009) pp. 031201–031201–8.
doi:10.1115/1.3059699
- [6] P. Barracund and D. Dickinson, On the associated Legendre polynomials. In *Orthogonal expansions and their continuous analogues* (Southern Illinois University, Carbondale, 1967), 43–50.
- [7] M. van den Berg and D. Bucur, On the torsion function with Robin or Dirichlet boundary conditions, arXiv 1305.2137 (2013).
- [8] R.H. Boyer, Discrete Bessel functions, *J. Math. Analysis and Applications*, **2** (1961) 509-524.
- [9] D. Bucur and A. Giacomini, The St Venant inequality for the Laplace operator with Robin boundary conditions, *Milan J. Math* **83** (2015), 327-343.
doi:10.1007/s00032-015-0243-0
- [10] J. Bustiz and M.E.H. Ismail, The associated ultraspherical polynomials and their q -analogues, *Canadian J. Math* **34**(3) (1982), 718-736

- [11] C.S. Chihara, *An Introduction to Orthogonal Polynomials*, (Gordon and Breach: 1978).
- [12] H. S. Cohl and D. E. Dominici, Generalized Heine's identity for complex Fourier series of binomials, *Proc. Roy. Soc. A* **467** (2010)
<https://doi.org/10.1098/rspa.2010.0222>
- [13] H. S. Cohl and H. Volkmer, Definite integrals using orthogonality and integral transforms. *SIGMA* **8** (2012)
- [14] J. T. Conway, Fourier and other series containing associated Legendre functions for incomplete Epstein-Hubbell integrals and functions related to elliptic integrals, *Integral Transforms and Special Functions*, **18:3** (2007), 179-191.
doi: 10.1080/10652460701210284
- [15] D. Cvijovic, The Fourier series expansions of the Legendre incomplete elliptic integrals of the first and second kind, *Integral Transforms and Special Functions*, **21(3)** (2009), 235-242.
<http://dx.doi.org/10.1080/10652460903178552>
- [16] Y. Damiano and G.C. Georgiou, Viscoplastic Poiseuille flow in a rectangular duct with wall slip, *Journal of NonNewtonian Fluid Mech.* **214** (2014).
doi:10.1016/j.jnnfm.2014.10.002
- [17] S.K. Das and F. Tahmouresi, Analytical solution of fully developed gaseous slip flow in elliptic microchannel, *Int. J. Adv. Appl. Math. and Mech.* **3i** (2016) 1 - 15.
- [18] S. D. Daymond, The principal frequencies of vibrating systems with elliptic boundaries, *Quart. J. Mech. Appl. Math.*, **8**, 1955, pp361-372.
- [19] Z. Duan and Y.S. Muzychka, Slip flow in non-circular microchannels, *Microfluidics and Nanofluidics* **3**, (2007), 473-484.
- [20] Z. Duan, Y.S. Muzychka, Slip flow in elliptic micro-channels, *International Journal of Thermal Sciences* **46**, 1104-1111 (2007).
- [21] A. Erdelyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions, Vol 1*, (McGraw-Hill: 1953)

- [22] A. Erdelyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions, Vol 2*, (McGraw-Hill: 1953)
- [23] S. Finch, Fraenkel asymmetry, (2014).
- [24] A.K. Ghosh, A. R. Khan, L. Debnath, On a pulsatile flow of a two phase viscous fluid in a tube of elliptic cross section, *Internat. J. Math. & Math. Sci.* **13**(4), 669-676 (1990).
- [25] A. Gil, J. Segura, and N.M. Temme, Computing Toroidal Functions for Wide Ranges of the Parameters, *Journal of Computational Physics* **161** (2000), 204217.
doi:10.1006/jcph.2000.6498
- [26] I.S. Gradshteyn and I.W. Ryzhik, *Tables of Integrals, Series and Products*, (Academic Press: 1965).
- [27] G.C. Grosjean, Theory of recursive generation of systems of orthogonal polynomials: An illustrative example. *Journal of Computational and Applied Mathematics* **12-13**, 299-318 (1985).
- [28] S. Gupta, D. Poulikakos and V. Kurtcuoglu. Analytical solution for pulsatile viscous flow in a straight elliptic annulus and application to the motion of the cerebrospinal fluid, *Phys. Fluids* **20**, 093607 (2008);
<http://dx.doi.org/10.1063/1.2988858>
- [29] H. Haslam and M. Zamir. Pulsatile flow in tubes of elliptic cross sections, *Annals of Biomedical Eng.* **26**, 780–787 (1998).
- [30] W.M. Hicks, On toroidal functions. XIV *Proc. Roy. Soc. London* (1881).
<https://royalsocietypublishing.org/doi/pdf/10.1098/rstl.1881.0014>
- [31] Jyh-Ping Hsu, Chen-Yuan Kao, Shiojenn Tseng, and Chur-Jen Chen, Electrokinetic Flow through an Elliptical Microchannel: Effects of Aspect Ratio and Electrical Boundary Conditions, *Journal of Colloid and Interface Science* **248**, 176–184 (2002).
- [32] D. Jackson, The Harmonic Boundary Value Problem for an Ellipse or an Ellipsoid, *American Mathematical Monthly*, **51**(10) (1944), 555-563.

- [33] M. Kaewbumrung, S. Orankitjaroen, P. Boonkrong, B. Nuntadilok and B. Wiwatanapataphee. Numerical Simulation of Dispersed Particle-Blood Flow in the Stenosed Coronary Arteries, *International Journal of Differential Equations* (2018), Article ID 2593425, 16 pages
<https://doi.org/10.1155/2018/2593425>
- [34] G. Keady, On Hadwiger's results concerning Minkowski sums and isoperimetric inequalities for moments of inertia, *RGMIA Research Report Collection* **9**(4) (2006).
- [35] G. Keady, Steady slip flow of newtonian fluids through tangential polygonal microchannels, Submitted (2020) to *IMA Journal of Applied Mathematics*.
- [36] G. Keady, Functions with constant laplacian satisfying homogeneous Robin boundary conditions: tangential polygonal domains, (2020) *Preprint in preparation for arXiv*.
 Draft at
<https://sites.google.com/site/keadyperthunis/home/papers/tangpolygcode>
- [37] G. Keady and A. McNabb, Functions with constant Laplacian satisfying homogeneous Robin boundary conditions, *I.M.A. Jnl of Applied Mathematics*, **50** (1993), 205-224.
[doi:10.1093/imamat/50.3.205](https://doi.org/10.1093/imamat/50.3.205)
- [38] G. Keady and B. Wiwatanapataphee, 2016, Some isoperimetric results concerning unidirectional flows in microchannels, arXiv. See also URL
<http://arxiv.org/abs/1604.03394>
- [39] G. Keady and B. Wiwatanapataphee, Variational approximations for steady unidirectional slip flows in microchannels, *Journal of Fluids Engineering*, **142**, (2020) p. 074501.
<https://doi.org/10.1115/1.4046405>
- [40] G. Keady, B. Wiwatanapataphee and N. Khajohnsaksumeth, On Unidirectional Flows in Microchannels with Slip at the Boundary In *Proceedings of The International Conference on Engineering and Applied Sciences*, 2018 (TICEAS). (Most conveniently available via ResearchGate.)

- [41] D.S. Kim and Y.H. Kim, A characterization of ellipses, *American Math. Monthly* **144**(1) (2007), 66-70.
- [42] H. Kober, *Dictionary of Conformal Representations*, (Dover: 1957)
- [43] H. Lamb, *Hydrodynamics*, (Dover printing of 1932 edition.)
- [44] D.F. Lawden, *Elliptic Functions and Applications*, (Springer, 1989).
- [45] N.N. Lebedev, L.P. Skalskaya and Y.S. Uflyand, *Worked Problems in Applied Mathematics*, (Dover: 1979)
- [46] A. Liemert and A. Kienle, Exact solution of Poisson's equation with an elliptical boundary, *Appl. Math. and Computation* **238** (2014), 123-131.
- [47] J.C. Mason and D. Handscomb, *Chebyshev Polynomials*, Chapman and Hall/CRC (2003).
- [48] A. McNabb and G.Keady, Diffusion and the torsion parameter, *J. Australian Math. Soc.*, **35B**, (1994), 289-301.
doi:10.1017/S0334270000009309
- [49] J.W. Meux, Ordinary differential equations of the fourth order with orthogonal polynomial solutions, *Amer. Math. Monthly* **73**(4), (1966), 104-110.
- [50] R.F. Millar, An integral equation solution to the Dirichlet problem for Laplace's equation in an ellipse, *J. Math. Anal. Applns* **147** (1990), 154-170.
- [51] F. Pollaczek, Sur une famille de polynomes orthogonaux a quatre parametres. *C R Acad Sci. Paris.* (1950) **230**22542256.
- [52] G. Polya and G.Szego. *Isoperimetric Inequalities in Mathematical Physics*, Ann. of Math. Studies no. 27, (Princeton University Press, Princeton, 1951).
- [53] C.J. Rees, Elliptic orthogonal polynomials, *Duke Math. J.* **12** (1945), 173-187.
- [54] W.C. Royster, A Poisson integral formula for the ellipse and some applications, *Proc. Amer. Math. Soc.* **15**(4) (1964), 661-670.
DOI: 10.2307.2034769

- [55] N.B. Savaseneril and H. Delibas, Analytic solution for two-dimensional heat equation for an ellipse region, *New Trends in Math. Sciences, NTM-Sci* **4**(1) (2016), 65-70.
doi:<http://dx.doi.org/10.20852/ntmsci.2016115614>
- [56] B.O.U. Sonnerup, Expression as a Legendre function, of an elliptic integral occurring in wing theory, TAR-TN 59-1, November 1959.
- [57] S. Suharsono, Analytical study of fluid flows with slip boundary, Ph.D. Thesis, (2012)
<https://espace.curtin.edu.au/handle/20.500.11937/2281>
- [58] S. Suharsono, A study slip flow through elliptic microchannels, *Sci.Int(Lahore)*, **30**(3),423-426 (2018).
- [59] Q. Sun, Y. Wu, L. Liu, B. Wiwatanapataphee. Study of a Newtonian fluid through circular channels with slip boundary taking into account electrokinetic effect, *Abstract and Applied Analysis*, 2013.
- [60] M. Spiga and P. Vocale, Slip flow in elliptic microducts with constant heat flux, *Advances in Mechanical Engineering* (2012), Article ID 481280.
doi:10.1155/2012/481280
- [61] G. Szego, Conformal mapping of the interior of an ellipse onto a circle, *American Mathematical Monthly*, **57**(7), (1950), 474-478.
- [62] L. Vinet and A. Zhedanov, Elliptic biorthogonal polynomials connected with Hermite's continued fraction, *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)* **3** (2007)
- [63] A. Tamayol and K. Hooman, Slip-flow in microchannels of non-circular cross sections, *Journal of Fluids Engineering*, **133**(9), (2011) pp. 091202–091202–8.
doi: 10.1115/1.4004591
- [64] B.A. Troesch and H.R. Troesch, Eigenfrequencies of an elliptic membrane, *Mathematics of Computation* **27**(124):755-755 (1973).
doi: 10.1090/S0025-5718-1973-0421276-2

- [65] J.L. Walsh, Note on the orthogonality of Tchebycheff polynomials on confocal ellipses, *Bull. Amer. Math. Soc.* **40** (1934), 84-88.
- [66] C.Y. Wang, Discussion: Slip-flow pressure drop in microchannels of general cross section, *Journal of Fluids Engineering* **134**(5), (2012) pp. 055501–055501–2.
doi:10.1115/1.4005724
- [67] C.Y. Wang, Brief review of exact solutions for slip-flow in ducts and channels, *Journal of Fluids Engineering*, **134**(9), (2012) pp. 094501–094501–3.
doi:10.1115/1.4007232
- [68] C.Y. Wang, Ritz method for slip flow in super-elliptic ducts, *European J. of Mechanics B/Fluids* **43**, (2014), 85-89.
- [69] Z.M. Wilmott, C.J.W. Breward and S.J. Chapman, Slip flow through channels with varying elliptic cross section, *IMA Journal of Applied Mathematics*, **83**, (2018) 874893.
- [70] B. Wiwatanapataphee, Y.H. Wu, and S. Suharnso, Transient flows of Newtonian fluid through a rectangular microchannel with slip, *Abstract and Applied Analysis* paper 530605 (2014).