

DYNAMICS OF THE FOCUSING 3D CUBIC NLS WITH SLOWLY DECAYING POTENTIAL

QING GUO, HUA WANG AND XIAOHUA YAO

ABSTRACT. In this paper, we consider a 3d cubic focusing nonlinear schrödinger equation (NLS) with slowly decaying potentials. Adopting the variational method of Ibrahim-Masmoudi-Nakanishi [9], we obtain a condition for scattering. It is actually sharp in some sense since the solution will blow up if it's false. The proof of blow-up part relies on the method of Du-Wu-Zhang [4].

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Variational characterization	13
4. Criteria for global well-posedness and blow-up	20
5. Scattering result	23
5.1. Existence and compactness of critical element	24
5.2. Precluding the critical element	28
References	28

1. INTRODUCTION

In this paper, we consider a 3d cubic focusing NLS with slowly decaying potentials (NLS_k)

$$(1.1) \quad \begin{cases} i\partial_t u - H_\alpha u + |u|^2 u = 0, & (t, x) \in \mathbf{R} \times \mathbf{R}^3, \\ u(0, x) = u_0(x) \in H^1(\mathbf{R}^3), \end{cases}$$

where $u : \mathbf{R} \times \mathbf{R}^3 \rightarrow \mathbf{C}$ is a complex-valued function, $H_\alpha = -\Delta + V(x)$ and $V(x) = \frac{k}{|x|^\alpha}$ with $k > 0$ and $1 < \alpha \leq 2$. Throughout this paper, we use the symbol $V(x)$ instead of $\frac{k}{|x|^\alpha}$ since we frequently use the general property of V : $V \geq 0$, $x \cdot \nabla V \leq 0$, $2V + x \cdot \nabla V \geq 0$ and $3x \cdot \nabla V + x \nabla^2 V x^T \leq 0$. As $\frac{k}{|x|^\alpha} > 0$ and $\frac{k}{|x|^\alpha} \in L^1_{loc}$, H_α is defined as a unique self-adjoint operator associated with the non-negative quadratic form $\langle (-\Delta + \frac{k}{|x|^\alpha})f, f \rangle$ on $C_0^\infty(\mathbf{R}^3)$. Moreover, H_α is purely absolutely

Date: December 4, 2018.

2000 Mathematics Subject Classification. 35P25; 35Q55; 47J35.

Key words and phrases. Cubic NLS; Focusing; Slowly decaying potential; Scattering; Blow-up.

continuous and has no eigenvalue. Since $k > 0$, the kernel $e^{-tH_\alpha}(x, y)$ of e^{-tH_α} satisfies the upper Gaussian estimate [18] i.e., for $\forall t > 0, \forall x, y \in \mathbf{R}^3$,

$$(1.2) \quad 0 \leq e^{-tH_\alpha}(x, y) \leq e^{t\Delta}(x, y) = (4\pi t)^{-\frac{3}{2}} e^{-\frac{|x-y|^2}{4t}},$$

which implies that Hardy inequality (2.2), Mihklin multiplier theorem and Littlewood-Paley theory (Bernstein inequalities Lemma 2.6, Littlewood-Paley decomposition Lemma 2.4 and square function estimates Lemma 2.5) associated with H_α . Hence it follows from Hardy inequality and Stein complex interpolation that the standard Sobolev norms and the Sobolev norms associated with H_α are equivalent (see Lemma 2.1). Recently, Mizutani [16] showed that e^{-itH_α} satisfies global-in-time Strichartz estimates for any admissible pairs. Combining the Sobolev norm equivalence and the Strichartz estimates and following the same line of the proof of Theorem 2.15 and Remark 2.16 of [14] yield that (NLS_k) is locally well-posed and scatters in $H^1(\mathbf{R}^3)$.

Theorem 1.1. [14] *Let $u_0 \in H^1(\mathbf{R}^3)$. Then the following are true.*

(i) *There exist $T = T(\|u_0\|_{H^1}) > 0$ and a unique solution $u \in C((-T, T), H^1(\mathbf{R}^3))$ of (NLS_k) .*

(ii) *There exists $\epsilon_0 > 0$ such that if for $0 < \epsilon < \epsilon_0$,*

$$\|e^{-itH_\alpha}u_0\|_{L_{t,x}^5(\mathbf{R}^+ \times \mathbf{R}^3)} < \epsilon,$$

then the solution u of (NLS_k) is global in the positive time direction and satisfies

$$\|u\|_{L_{t,x}^5(\mathbf{R}^+ \times \mathbf{R}^3)} \lesssim \epsilon.$$

The similar result holds in the negative time direction.

(iii) *For any $\phi \in H^1(\mathbf{R}^3)$, then there exist $T > 0$ and a solution $u \in C((T, +\infty), H^1(\mathbf{R}^3))$ of (NLS_k) such that*

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{-itH_\alpha}\phi\|_{H^1(\mathbf{R}^3)} = 0.$$

The similar result holds in the negative time direction.

(iv) *If $u : \mathbf{R} \times \mathbf{R}^3 \rightarrow \mathbf{C}$ is a global solution of (NLS_k) with*

$$(1.3) \quad \|u\|_{L_{t,x}^5(\mathbf{R} \times \mathbf{R}^3)} < +\infty,$$

then the solution $u(t)$ scatters in H^1 . That is, there exists $\phi_\pm \in H^1(\mathbf{R}^3)$ such that

$$\lim_{t \rightarrow \pm\infty} \|u(t) - e^{-itH_\alpha}\phi_\pm\|_{H^1(\mathbf{R}^3)} = 0.$$

Moreover, the H^1 solution u obeys the mass and energy conservation laws:

$$(1.4) \quad M(u) = \int_{\mathbf{R}^3} |u(t, x)|^2 dx = M(u_0),$$

and

$$(1.5) \quad E(u) = E_k(u) = \frac{1}{2} \int_{\mathbf{R}^3} |\nabla u(x)|^2 dx + \frac{1}{2} \int_{\mathbf{R}^3} V(x)|u(x)|^2 dx - \frac{1}{4} \int_{\mathbf{R}^3} |u(x)|^4 dx = E(u_0).$$

In the case $k = 0$, Holmer-Roudenko [7] and Duyckaerts-Holmer-Roudenko [5] employed the concentration-compactness approach of Kenig-Merle [11] to obtain sharp criteria between scattering and blow up for (NLS_0) in terms of conservation laws ((1.4) and (1.5)) and the ground state Q , which is the unique positive radial exponential decaying solution of the elliptic equation

$$(1.6) \quad \Delta Q - Q + Q^3 = 0.$$

Fang-Xie-Canzenave [6] and Akahor-Nawa [1] extended the result in [7, 5] to the general power and dimensions. Subsequently, Killip-Murphy-Visan-Zhang [14] established a corresponding sharp threshold between scattering and blow up for (NLS_k) with $k > -\frac{1}{4}$ and $\alpha = 2$. Recently, Miao-Zhang-Zheng [15] used the interaction Morawetz-type estimates and the equivalence of Sobolev norms to prove all solutions scatter for (NLS_k) with $k > 0$, $\alpha = 1$ and $-|u|^{p-1}u$ ($\frac{7}{3} < p < 5$) in place of $|u|^2u$ (i.e., nonlinear Schrödinger equation with repulsive Coulomb potential in the defocusing).

The goal of this paper is to extend the sharp scattering criterion in [14] from $\alpha = 2$ to $1 < \alpha \leq 2$ when $k > 0$ in some sense. Obviously, for $1 < \alpha < 2$, the equation (NLS_k) doesn't enjoy scaling invariant. Therefore, we cannot apply scaling as indicated in [14] to get a critical element (a minimal blow up solution). Hence, we shall adopt the variational argument based on the work of Ibrahim-Masmoudi-Nakanishi [9] to overcome the difficulty. Recently, the same argument have been applied to the focusing mass-supercritical nonlinear Schrödinger equation with repulsive Dirac delta potential on the real line (see [10]).

To state our main result, we introduce some notation now. We define the functional S_k as

$$(1.7) \quad S_k(\varphi) := E(\varphi) + \frac{1}{2}M(\varphi) = \frac{1}{2}\|\varphi\|_{\mathcal{H}_k^1}^2 - \frac{1}{4} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx,$$

where

$$(1.8) \quad \|\varphi\|_{\mathcal{H}_k^1}^2 = \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx + \int_{\mathbf{R}^3} V(x)|\varphi(x)|^2 dx + \int_{\mathbf{R}^3} |\varphi(x)|^2 dx,$$

which is equivalent to $\|\varphi\|_{H^1}$ by $k > 0$ by Hardy's inequality. Denote the scaling quantity $\varphi_\lambda^{a,b}$ by

$$(1.9) \quad \varphi_\lambda^{a,b} := e^{a\lambda} \varphi(e^{-b\lambda} x),$$

where (a, b) satisfies the condition

$$(1.10) \quad a > 0, \quad b \leq 0, \quad 2a + b > 0, \quad 2a + 3b \geq 0, \quad (a, b) \neq (0, 0).$$

We define the scaling derivative of $S_k(\varphi_\lambda^{a,b})$ at $\lambda = 0$ by $K_k^{a,b}(\varphi)$.

$$(1.11) \quad \begin{aligned} K_k^{a,b}(\varphi) &:= \mathcal{L}^{a,b} S_k(\varphi) = \frac{d}{d\lambda} \Big|_{\lambda=0} S_k(\varphi_\lambda^{a,b}) \\ &= \frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx + \frac{2a+3b}{2} \int_{\mathbf{R}^3} V|\varphi(x)|^2 dx + \frac{b}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V)|\varphi(x)|^2 dx \\ &\quad + \frac{2a+3b}{2} \int_{\mathbf{R}^3} |\varphi(x)|^2 dx - \frac{4a+3b}{4} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \end{aligned}$$

In particular, when $(a, b) = (3, -2)$,

$$(1.12) \quad P_k(\varphi) := K_k^{3,-2}(\varphi) = 2 \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx - \int_{\mathbf{R}^3} (x \cdot \nabla V)|\varphi(x)|^2 dx - \frac{3}{2} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx,$$

which is related with the Virial identity of (NLS_k) (4.3) with $\phi(x) = |x|^2$, and when $(a, b) = (3, 0)$,

$$(1.13) \quad I_k(\varphi) := \frac{1}{3} K_k^{3,0}(\varphi) = \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx + \int_{\mathbf{R}^3} V|\varphi(x)|^2 dx + \int_{\mathbf{R}^3} |\varphi(x)|^2 dx - \int_{\mathbf{R}^3} |\varphi(x)|^4 dx.$$

We note that, to get existence of minimal blow-up solutions, we need to use the functional I_k instead of P_k . so that we can apply the linear profile decomposition Lemma 2.12.

The sharp threshold quantity n_k are determined by the following minimizing problem

$$(1.14) \quad n_k = \inf\{S_k(\varphi) : \varphi \in H^1(\mathbf{R}^3) \setminus \{0\}, P_k(\varphi) = 0\}.$$

When $k = 0$, n_0 is positive and is achieved by Q , which is a unique radial solution of (1.6) (see [1]). The sharp criteria between scattering and blow up as mentioned above can be described by n_0 . We state the results of [7, 5, 6, 1] in terms of n_0 as follows.

Theorem 1.2. [7, 5, 6, 1] *Let $u_0 \in H^1(\mathbf{R}^3)$ satisfy $S_0(u_0) < n_0$.*

(i) *If $P_0 \geq 0$, then the solution u of (NLS₀) is global and scatters.*

(ii) *If $P_0 < 0$ and u_0 is radial or $xu_0 \in L^2(\mathbf{R}^3)$, then the solution u of (NLS₀) blows up in finite time in both time directions.*

Furthermore, if $\psi \in H^1(\mathbf{R}^3)$ satisfying $\frac{1}{2}\|\psi\|_{H^1}^2 < n_0$, then there exists a global solution of (INLS₀) that scatters to ψ in the positive time direction. The analogous statement holds in the negative time direction.

When $k > 0$, we prove that $n_k = n_0$ and n_k is never attained (see Lemma 3.6). For succinctness, we next define two subsets of $H^1(\mathbf{R}^3)$ as follows:

$$(1.15) \quad \mathcal{N}^+ := \{\varphi \in H^1(\mathbf{R}^3) : S_k(\varphi) < n_0, P_k(\varphi) \geq 0\}$$

and

$$(1.16) \quad \mathcal{N}^- := \{\varphi \in H^1(\mathbf{R}^3) : S_k(\varphi) < n_0, P_k(\varphi) < 0\}.$$

Now we state our main result.

Theorem 1.3. *Let u be the solution of (NLS_k) on $(-T_{min}, T_{max})$, where $(-T_{min}, T_{max})$ is the maximal life-span.*

(i) *If $u_0 \in \mathcal{N}^+$, then u is global well-posedness, $u(t) \in \mathcal{N}^+$ for any $t \in \mathbf{R}$ and scatters.*

(ii) *If $u_0 \in \mathcal{N}^-$, then $u(t) \in \mathcal{N}^-$ for any $t \in (-T_{min}, T_{max})$ and one of the following four statements holds true:*

(a) *$T_{max} < +\infty$ and $\lim_{t \uparrow T_{max}} \|\nabla u(t)\|_{L^2} = +\infty$.*

(b) *$T_{min} < +\infty$ and $\lim_{t \downarrow -T_{min}} \|\nabla u(t)\|_{L^2} = +\infty$.*

(c) *$T_{max} = +\infty$ and there exists a sequence $\{t_n\}_{n=1}^{+\infty}$ such that $t_n \rightarrow +\infty$ and $\lim_{t_n \uparrow +\infty} \|\nabla u(t)\|_{L^2} = +\infty$.*

(d) *$T_{min} = +\infty$ and there exists a sequence $\{t_n\}_{n=1}^{+\infty}$ such that $t_n \rightarrow -\infty$ and $\lim_{t_n \downarrow -\infty} \|\nabla u(t)\|_{L^2} = +\infty$.*

Here the blow-up result is proved by the method of Du-Wu-Zhang [4].

This present paper is organized as follows. We fix notations at the end of Section 1. In Section 2, as preliminaries, we state some our required lemmas, including Sobolev norm equivalence, Strichartz estimates, stability theory, Littlewood-Paley theory, some limit lemmas between H_α^n and H_α^∞ , linear profile decomposition and nonlinear profiles for $|x_n| \rightarrow +\infty$. In Section 3, using the variational idea of Ibrahim-Masmoudi-Nakanishi [9], we obtain that if $\psi \in \mathcal{N}^+$, $P_k(\psi)$ and $I_k(\psi)$ have the same sign and $S_k(\psi)$ is equivalent to $\|\psi\|_{H^1}$ and if $\psi \in \mathcal{N}^\pm$, $P_k(\psi)$ has the uniform

bounds, which play a vital role to get blow-up and scattering results. In Section 4, using the upper bound of $P_k(\psi)$ for $\psi \in \mathcal{N}^-$ and adopting the method of Du-Wu-Zhang [4], we establish blow-up part of Theorem 1.3. Global part of Theorem 1.3 can be obtained by the lower bound of $P_k(\psi)$ for $\psi \in \mathcal{N}^+$ and local well-posedness i of Theorem 1.1. In the last section, we show the scattering part of Theorem 1.3 in two steps. In Step 1, by contradiction, if scattering fails, then a critical element must exist. In Step 2, we utilize lower bound of $P_k(\psi)$ for $\psi \in \mathcal{N}^+$ to preclude the critical element. Putting the last two sections together completes the proof of Theorem 1.3.

Notations:

We fix notations used throughout the paper. In what follows, we write $A \lesssim B$ to signify that there exists a constant c such that $A \leq cB$, while we denote $A \sim B$ when $A \lesssim B \lesssim A$. Given a real number α , $\alpha- = \alpha - \epsilon$ for $0 < \epsilon \ll 1$.

Let $L_t^q L_x^r$ be the space of measurable functions from an interval $I \subset \mathbf{R}$ to L_x^r whose $L_t^q L_x^r$ - norm $\|\cdot\|_{L_t^q L_x^r}$ is finite, where

$$\|u\|_{L_t^q L_x^r} = \left(\int_I \|u(t)\|_{L_x^r}^q dt \right)^{\frac{1}{q}}.$$

When $I = \mathbf{R}$, we may use $L_t^q L_x^r$ instead of $L_I^q L_x^r$, respectively. In particular, when $q = r$, we may simply write them as $L_{t,x}^q$, respectively.

Moreover, the Fourier transform on \mathbf{R}^3 is defined by $\hat{f}(\xi) = (2\pi)^{-\frac{3}{2}} \int_{\mathbf{R}^3} e^{-ix \cdot \xi} f(x) dx$. Define the inhomogeneous Sobolev space $H^s(\mathbf{R}^3)$ and the homogeneous Sobolev space $\dot{H}^s(\mathbf{R}^3)$, respectively, by norms

$$\|f\|_{H^s(\mathbf{R}^3)} = \|(1 + |\xi|^2)^{\frac{s}{2}} \hat{f}(\xi)\|_{L^2(\mathbf{R}^3)} = \|(1 + \Delta)^{\frac{s}{2}} f\|_{L^2(\mathbf{R}^3)}$$

and

$$\|f\|_{\dot{H}^s(\mathbf{R}^3)} = \||\xi|^s \hat{f}(\xi)\|_{L^2(\mathbf{R}^3)} = \|\Delta^{\frac{s}{2}} f\|_{L^2(\mathbf{R}^3)}.$$

Denote the inhomogeneous Sobolev space and homogeneous Sobolev space adapted to H_α by $\mathcal{H}_k^{s,p}(\mathbf{R}^3)$ and $\dot{\mathcal{H}}_k^{s,p}(\mathbf{R}^3)$, respectively, with norms

$$\|f\|_{\mathcal{H}_k^{s,p}(\mathbf{R}^3)} = \|(1 + H_\alpha)^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)}$$

and

$$\|f\|_{\dot{\mathcal{H}}_k^{s,p}(\mathbf{R}^3)} = \|H_\alpha^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)}.$$

$\mathcal{H}_k^s(\mathbf{R}^3)$ and $\dot{\mathcal{H}}_k^s(\mathbf{R}^3)$ are shorthand by $\mathcal{H}_k^{s,2}(\mathbf{R}^3)$ and $\dot{\mathcal{H}}_k^{s,2}(\mathbf{R}^3)$, respectively.

Given $p \geq 1$, let p' be the conjugate of p , that is $\frac{1}{p} + \frac{1}{p'} = 1$.

Acknowledgement The first author is financially supported by the China National Science Foundation (No.11301564, 11771469), the second author is financially supported by the China National Science Foundation (No. 11771165 and 11571131), and the third author is financially supported by the China National Science Foundation(No. 11771165).

2. PRELIMINARIES

As we've mentioned in the introduction, the heat kernel associated with H_α satisfies (1.2), so Mihlin multiplier theorem holds, which implies that for $\forall 1 < p < \infty$,

$$(2.1) \quad \|f\|_{L^p} \lesssim \|(1 + H_\alpha)f\|_{L^p}.$$

And we have the following Hardy type inequality for H_α (e.g., see [12] for $\alpha = 2$).

$$(2.2) \quad \left\| |x|^{-s} f \right\|_{L^p(\mathbf{R}^3)} \lesssim \|H_\alpha^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)} \lesssim \|(1 + H_\alpha)^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)},$$

where $0 < s < 3$ and $1 < p < \frac{3}{s}$.

Using (2.1) and Stein complex interpolation yields the following Sobolev norm equivalence (see [8] for $V \geq 0$ and $V \in L^{\frac{3}{2}}$ and [19, 12, 15] for $\alpha = 2$).

Lemma 2.1. *Let $k > 0$ and $0 < \alpha < 2$, $1 < p < \frac{3}{s}$ and $0 \leq s \leq 2$, then*

$$(2.3) \quad \|(1 + H_\alpha)^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)} \sim \|(1 - \Delta)^{\frac{s}{2}} f\|_{L^p(\mathbf{R}^3)}.$$

Proof. As the heat kernel associated with H_α satisfies (1.2), the kernel of Riesz potentials $(1 + H_\alpha)^{-\frac{s}{2}}$

$$(1 + H_\alpha)^{-\frac{s}{2}}(x, y) = \frac{1}{\Gamma(\frac{s}{2})} \int_0^{+\infty} e^{-t(1+H_\alpha)}(x, y) t^{\frac{s}{2}-1} dt$$

satisfies

$$|(1 + H_\alpha)^{-\frac{s}{2}}(x, y)| \lesssim |x - y|^{s-3},$$

which, by Hardy-Littlewood-Sobolev inequality, implies that

$$(2.4) \quad \|(1 + H_\alpha)^{-\frac{s}{2}} f\|_{L^{\frac{3p}{3-ps}}} \lesssim \|f\|_{L^p}.$$

It suffices to prove that (2.3) with $s = 2$ and $1 < p < \frac{3}{2}$ holds. Indeed, if it is true, then (2.3) follows from Stein complex interpolation and the L^p -boundedness of $(1 + H_\alpha)^{iy}$ with $\forall y \in \mathbf{R}$ and $1 < p < +\infty$ (which can be obtained by (1.2) and Sikora-Wright [17]).

Let $\chi(x)$ be a smooth compact supported function such that $\chi(x) = 1$ for $|x| \leq 1$ and $\chi(x) = 0$ for $|x| \geq 2$. On one hand, using Hölder inequality and Sobolev embedding yields that

$$\begin{aligned} \|(1 + H_\alpha)f\|_{L^p} &\leq \|(1 - \Delta)f\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} f \right\|_{L^p} \\ &\leq \|(1 - \Delta)f\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} \chi f \right\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} (1 - \chi)f \right\|_{L^p} \\ &\lesssim \|(1 - \Delta)f\|_{L^p} + \left\| |x|^{-\alpha} \chi \right\|_{L^{\frac{3}{2}}} \|f\|_{L^{\frac{3p}{3-2p}}} + \|f\|_{L^p} \\ &\lesssim \|(1 - \Delta)f\|_{L^p}. \end{aligned}$$

On the other hand, using Hölder inequality, (2.1) and (2.4) with $s = 2$ gives that

$$\begin{aligned} \|(1 - \Delta)f\|_{L^p} &\leq \|(1 + H_\alpha)f\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} f \right\|_{L^p} \\ &\leq \|(1 + H_\alpha)f\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} \chi f \right\|_{L^p} + k \left\| \frac{1}{|x|^\alpha} (1 - \chi)f \right\|_{L^p} \\ &\lesssim \|(1 + H_\alpha)f\|_{L^p} + \left\| |x|^{-\alpha} \chi \right\|_{L^{\frac{3}{2}}} \|f\|_{L^{\frac{3p}{3-2p}}} + \|f\|_{L^p} \\ &\lesssim \|(1 + H_\alpha)f\|_{L^p}. \end{aligned}$$

Thus, we get (2.3) with $s = 2$ and $1 < p < \frac{3}{2}$ and then conclude the proof. \square

Recently, Mizutani [16] proved that the solution to free Schrödinger equations with a class of slowly decaying repulsive potentials including $k|x|^{-\alpha}$ with $k > 0$ and $0 < \alpha < 2$ satisfies global-in-time Strichartz estimates for any admissible pairs. Besides, it is well known that Strichartz estimates for free Schrödinger equation with inverse-square potentials were established by Burq-Planchon-Stalker-Tahvildar-Zadeh [2]. Hence, we have the following global-in-time Strichartz estimate.

Lemma 2.2. [2, 16] *Let $k > 0$ and $0 < \alpha \leq 2$. Then the solution u of $iu_t - H_\alpha u = F$ with initial data u_0 obeys*

$$(2.5) \quad \|u\|_{L_t^q L_x^r} \lesssim \|u_0\|_{L_x^2} + \|F\|_{L_t^{\tilde{q}} L_x^{\tilde{r}}}$$

for any $2 \leq q, \tilde{q} \leq \infty$ with $\frac{2}{q} + \frac{3}{r} = \frac{2}{\tilde{q}} + \frac{3}{\tilde{r}} = \frac{3}{2}$.

Once we have Strichartz estimates Lemma 2.2 and Sobolev norm equivalence Lemma 2.1, the local well-posedness Theorem 1.1 and stability result Lemma 2.3 for (NLS_k) can be obtained by the same proofs as in Theorem 2.15 and Theorem 2.17 of [14], respectively.

Lemma 2.3. [14] *For $k > 0$ and $0 < \alpha \leq 2$. Let \tilde{u} be the solution of*

$$(2.6) \quad \begin{cases} i\tilde{u}_t - H_\alpha \tilde{u} + |\tilde{u}|^2 \tilde{u} = e, & (t, x) \in I \times \mathbf{R}^3, \\ \tilde{u}(0, x) = \tilde{u}_0(x) \in H^1(\mathbf{R}^3), \end{cases}$$

for some 'error' e . Given $u_0 \in H^1(\mathbf{R}^3)$ and assume

$$(2.7) \quad \|u_0\|_{H^1} + \|\tilde{u}_0\|_{H^1} \leq A \quad \text{and} \quad \|\tilde{u}\|_{L_{t,x}^5} \leq M$$

for some $A, M > 0$. For any given $\frac{1}{2} \leq s < 1$, there exists $\epsilon_0 = \epsilon_0(A, M)$ such that if $0 < \epsilon < \epsilon_0$ and

$$(2.8) \quad \|u_0 - \tilde{u}_0\|_{H^s} + \left\| (1 - \Delta)^{\frac{s}{2}} e \right\|_{N(I)} < \epsilon,$$

where

$$N(I) := L_{t,x}^{\frac{10}{7}} + L_I^{\frac{5}{3}} L_x^{\frac{30}{23}} + L_I^1 L_x^2,$$

then there exists a solution u of (NLS_k) such that

$$(2.9) \quad \|u - \tilde{u}\|_{S_a^s(I)} \lesssim_{A,M} \epsilon,$$

$$(2.10) \quad \|u\|_{S_a^{\frac{1}{2}}(I)} \lesssim_{A,M} 1,$$

where

$$S_a^s(I) = L_I^2 \mathcal{H}_\alpha^{s,6} \cap L_I^\infty \mathcal{H}_\alpha^s.$$

Since Mihlin multiplier theorem for H_α holds, naturally, we have the Littlewood-Paley theory associated with H_α (see [12] for $\alpha = 2$). We first give the definition of Littlewood-Paley projection via the heat kernel as follows: For $N \in 2^{\mathbf{Z}}$,

$$(2.11) \quad P_N := e^{-\frac{1}{N^2} H_\alpha} - e^{-\frac{4}{N^2} H_\alpha}.$$

We next state Littlewood-Paley decomposition, square function estimate and Bernstein estimates (see [12] for $\alpha = 2$).

Lemma 2.4. *Let $1 < p < \infty$. If $k > 0$ and $0 < \alpha \leq 2$, then*

$$(2.12) \quad f = \sum_{N \in 2^{\mathbb{Z}}} P_N f$$

as elements of $L^p(\mathbf{R}^3)$. In particular, the sum converges in $L^p(\mathbf{R}^3)$.

Lemma 2.5. *Let $0 \leq s < 2$ and $1 < p < +\infty$. If $k > 0$ and $0 < \alpha \leq 2$, then*

$$(2.13) \quad \left\| \left(\sum_{N \in 2^{\mathbb{Z}}} N^{2s} |P_N f|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbf{R}^3)} \sim \|(H_\alpha)^{\frac{s}{2}}\|_{L^p(\mathbf{R}^3)}.$$

Lemma 2.6. *Let $1 < p \leq q < +\infty$ and $s \in \mathbf{R}$. If $k > 0$ and $0 < \alpha \leq 2$, then*

(i)

$$(2.14) \quad \|P_N f\|_{L^p(\mathbf{R}^3)} \lesssim \|f\|_{L^p(\mathbf{R}^3)}.$$

(ii)

$$(2.15) \quad \|P_N f\|_{L^q(\mathbf{R}^3)} \lesssim N^{3(\frac{1}{p} - \frac{1}{q})} \|f\|_{L^p(\mathbf{R}^3)}.$$

(iii)

$$(2.16) \quad \|(H_\alpha)^{\frac{s}{2}} P_N f\|_{L^p(\mathbf{R}^3)} \sim N^s \|f\|_{L^p(\mathbf{R}^3)}.$$

In order to establish linear profile decomposition associated with e^{-itH_α} and find a critical element, we apply the argument of [13] to get some convergence results. For convenience, define two operators: Let $\{x_n\}_{n=1}^{+\infty} \subset \mathbf{R}^3$,

$$(2.17) \quad H_\alpha^n := -\Delta + V(x + x_n) \quad \text{and} \quad H_\alpha^\infty := \begin{cases} -\Delta + V(x + \bar{x}) & x_n \rightarrow \bar{x} \in \mathbf{R}^3 \\ -\Delta & |x_n| \rightarrow +\infty \end{cases}$$

Obviously, $\tau_{x_n} H_\alpha^n \psi = H_\alpha \tau_{x_n} \psi$, where $\tau_y \psi(x) = \psi(x - y)$. So H_α doesn't commute with τ_{x_n} . H_α^∞ can be regarded as limits of H_α^n in the following sense (see [13] for $\alpha = 2$).

Lemma 2.7. *Let $k > 0$ and $1 < \alpha < 2$. Assume $t_n \rightarrow \bar{t} \in \mathbf{R}$ and $\{x_n\}_{n=1}^{+\infty} \rightarrow \bar{x} \in \mathbf{R}^3$ or $|x_n| \rightarrow +\infty$. Then*

$$(2.18) \quad \lim_{n \rightarrow +\infty} \|(H_\alpha^n - H_\alpha^\infty)\psi\|_{H^{-1}} = 0, \quad \forall \psi \in H^1.$$

$$(2.19) \quad \lim_{n \rightarrow +\infty} \|(e^{-it_n H_\alpha^n} - e^{-i\bar{t} H_\alpha^\infty})\psi\|_{H^{-1}} = 0, \quad \forall \psi \in H^{-1}.$$

$$(2.20) \quad \lim_{n \rightarrow +\infty} \|(1 + H_\alpha^n)^{\frac{1}{2}} - (1 + H_\alpha^\infty)^{\frac{1}{2}}\|\psi\|_{L^2} = 0, \quad \forall \psi \in H^1.$$

For any $2 < q \leq +\infty$ and $\frac{2}{q} + \frac{3}{r} = \frac{3}{2}$.

$$(2.21) \quad \lim_{n \rightarrow +\infty} \|(e^{-it H_\alpha^n} - e^{-it H_\alpha^\infty})\psi\|_{L_t^q L_x^r} = 0, \quad \forall \psi \in L^2.$$

If $\bar{x} \neq 0$, then for any $t > 0$,

$$(2.22) \quad \lim_{n \rightarrow +\infty} \|(e^{-it H_\alpha^n} - e^{-it H_\alpha^\infty})\delta_0\|_{H^{-1}} = 0.$$

Proof. Here we only give the proof of (2.21) because the others can be obtained by using the same method of Lemma 3.3 in [13], where in the proof of (2.18)-(2.20) we need to replace \dot{H}^1 , \dot{H}^{-1} , H_α^n , H_α^∞ and homogeneous Sobolev equivalence Theorem 2.2 in [13] with H^1 , H^{-1} , $1 + H_\alpha^n$, $1 + H_\alpha^\infty$ and inhomogeneous Sobolev equivalence Lemma 2.1, respectively and in the proof of (2.22) we need to replace \dot{H}^{-1} by H^{-1} . It only suffices to prove (2.21) in the case $(q, r) = (\infty, 2)$, since the general case can be obtained by interpolating with the end-point Strichartz estimates (i.e., $(q, r) = (2, 6)$). By density and Strichartz estimates, let ψ be a smooth function with compact support, so that for $\forall M > 0$,

$$(2.23) \quad |(e^{-it\Delta}\psi)(x)| \lesssim_\psi \langle t \rangle^{-\frac{3}{2}} \left(1 + \frac{|x|}{\langle t \rangle}\right)^{-M}.$$

By the definition of H_α^∞ , we consider two cases: $|x_n| \rightarrow +\infty$ and $x_n \rightarrow \bar{x}$.

For the first case $|x_n| \rightarrow +\infty$, we have

$$\begin{aligned} e^{it\Delta}\psi &= e^{-itH_\alpha^n}\psi + i \int_0^t e^{-i(t-s)H_\alpha^n} V(x+x_n) e^{is\Delta}\psi ds \\ &= e^{-itH_\alpha^n}\psi + i \int_0^t e^{-i(t-s)H_\alpha^n} (\chi_{|x+x_n|\leq R} + \chi_{|x+x_n|>R}) V(x+x_n) e^{is\Delta}\psi ds, \end{aligned}$$

where R is a sufficiently large number that will be chosen later and χ_A is a characteristic function on a set A . Hence by Strichartz estimates (2.5) and dispersive estimates (2.23), we find that

$$\begin{aligned} \|(e^{-itH_\alpha^n} - e^{-itH_\alpha^\infty})\psi\|_{L_t^\infty L_x^2} &\lesssim \|\chi_{|x+x_n|\leq R} V(x+x_n) e^{it\Delta}\psi\|_{L_{t,x}^{\frac{10}{7}}} + \|\chi_{|x+x_n|>R} V(x+x_n) e^{it\Delta}\psi\|_{L_t^q L_x^r} \\ &\lesssim \|\chi_{|x+x_n|\leq R} V(x+x_n) \langle t \rangle^{-\frac{3}{2}} \left(1 + \frac{|x|}{\langle t \rangle}\right)^{-M}\|_{L_{t,x}^{\frac{10}{7}}} \\ &\quad + \|\chi_{|x+x_n|>R} V(x+x_n) \langle t \rangle^{-\frac{3}{2}} \left(1 + \frac{|x|}{\langle t \rangle}\right)^{-M}\|_{L_t^q L_x^r} \\ (2.24) \quad &:= I_1 + I_2, \end{aligned}$$

where $(q, r) = (1, 2)$ if $1 < \alpha \leq \frac{3}{2}$ and $(q, r) = (\frac{10}{7}, \frac{10}{7})$ if $\frac{3}{2} < \alpha \leq 2$.

We note that $|x_n| \geq R$ for n large enough. So for the first part I_1 , $R < |x| \sim |x_n|$. If $|x| \leq \langle t \rangle$, then $|x_n| \lesssim \langle t \rangle$. Therefore,

$$(2.25) \quad I_1 \lesssim \|V(x+x_n)\|_{L_x^{\frac{10}{7}}(|x+x_n|\leq R)} \left\| \langle t \rangle^{-\frac{3}{2}} \right\|_{L_t^{\frac{10}{7}}(|t|\geq|x_n|)} \lesssim R^{-\alpha+\frac{21}{10}} |x_n|^{-\frac{4}{5}}.$$

If $|x| > \langle t \rangle$,

$$I_1 \lesssim \|\chi_{|x+x_n|\leq R} V(x+x_n) \langle t \rangle^{-\frac{3}{2}+M} |x|^{-M}\|_{L_{t,x}^{\frac{10}{7}}}.$$

When $|t| \leq 1$, it is easy to get

$$(2.26) \quad I_1 \lesssim R^{-\alpha+\frac{21}{10}} |x_n|^{-M}.$$

When $|t| > 1$ and M sufficiently large,

$$\|\langle t \rangle^{-\frac{3}{2}+M}\|_{L_t^{\frac{10}{7}}(|t|\leq|x|)} \lesssim |x|^{-\frac{3}{2}+M+\frac{7}{10}},$$

so

$$(2.27) \quad I_1 \lesssim R^{-\alpha+\frac{21}{10}} |x_n|^{-\frac{4}{5}}.$$

Putting (2.25), (2.26) and (2.27) together gives that I_1 tends to 0 as n approaches $+\infty$, regardless of $R > 0$.

For the second part I_2 , we consider two subcases: $1 < \alpha \leq \frac{3}{2}$ and $\frac{3}{2} < \alpha \leq 2$. If $1 < \alpha \leq \frac{3}{2}$, $(q, r) = (1, 2)$. When $|x| \leq \langle t \rangle$, using Hölder inequality, we have

$$(2.28) \quad I_2 \lesssim \|V(x + x_n)\|_{L_x^{\frac{3}{\alpha}}(|x+x_n|>R)} \left\| |t|^{-\frac{3}{2}} \right\|_{L_t^1 L_x^{\frac{6}{3-2(\alpha)}}(|x| \leq \langle t \rangle)} \lesssim R^{1-\frac{\alpha}{2}}.$$

When $|x| > \langle t \rangle$,

$$(2.29) \quad I_2 \lesssim \|V(x + x_n)\|_{L_x^{\frac{3}{\alpha}}(|x+x_n|>R)} \left\| |t|^{-\frac{3}{2}} |x|^{-M} \langle t \rangle^M \right\|_{L_t^1 L_x^{\frac{6}{3-2(\alpha)}}(|x|>\langle t \rangle)} \lesssim R^{1-\frac{\alpha}{2}}.$$

It follows from (2.28) and (2.29) that I_2 can be chosen arbitrarily small if we take R sufficiently large.

If $\frac{3}{2} < \alpha \leq 2$, $(q, r) = (\frac{10}{7}, \frac{10}{7})$. When $|x| \leq \langle t \rangle$, using Hölder inequality, we have

$$(2.30) \quad I_2 \lesssim \|V(x + x_n)\|_{L_x^2(|x+x_n|>R)} \left\| |t|^{-\frac{3}{2}} \right\|_{L_t^{\frac{10}{7}} L_x^{\frac{5}{3}}(|x| \leq \langle t \rangle)} \lesssim R^{-\alpha+\frac{3}{2}}.$$

When $|x| > \langle t \rangle$,

$$(2.31) \quad I_2 \lesssim \|V(x + x_n)\|_{L_x^2(|x+x_n|>R)} \left\| |t|^{-\frac{3}{2}} |x|^{-M} \langle t \rangle^M \right\|_{L_t^{\frac{10}{7}} L_x^{\frac{5}{3}}(|x|>\langle t \rangle)} \lesssim R^{-\alpha+\frac{3}{2}}.$$

It follows from (2.30) and (2.31) that I_2 can be chosen arbitrarily small if we take R sufficiently large. So we get (2.21) in the case $|x_n| \rightarrow +\infty$.

Now we turn to the other case $x_n \rightarrow \bar{x}$. Here we obtain that

$$e^{-itH_\alpha^\infty} \psi = e^{-itH_\alpha^n} \psi + i \int_0^t e^{-i(t-s)H_\alpha^n} (V(x + x_n) - V(x + \bar{x})) e^{-isH_\alpha^\infty} \psi ds$$

Hence by Strichartz estimates (2.5), we have

$$\begin{aligned} \|(e^{-itH_\alpha^n} - e^{-itH_\alpha^\infty})\psi\|_{L_t^\infty L_x^2} &\lesssim \|(V(x + x_n) - V(x + \bar{x}))e^{-itH_\alpha^\infty} \psi\|_{L_t^{\frac{6}{5}} L_x^{\frac{6}{5}}} \\ &= \|(V(x + x_n - \bar{x}) - V(x))e^{-itH_\alpha} \tau_{\bar{x}} \psi\|_{L_t^{\frac{6}{5}} L_x^{\frac{6}{5}}} \end{aligned}$$

Replacing $x_n - \bar{x}$ by x_n and $\tau_{\bar{x}} \psi$ by ψ , so we can suppose that $\bar{x} = 0$. Hence, for $\forall \epsilon > 0$, $|x_n| < \epsilon$ when n is large enough. Besides, by Newton-Leibniz formula,

$$|V(x + x_n) - V(x)| \lesssim |x_n| \int_0^1 |x + (1-t)x_n|^{-\alpha-1} dt.$$

Using Hölder inequality and Strichartz estimates (2.5) yields that

$$\begin{aligned} \|(e^{-itH_\alpha^n} - e^{-itH_\alpha^\infty})\psi\|_{L_t^\infty L_x^2} &\lesssim \|(V(x + x_n) - V(x))\|_{L_x^{\frac{15}{11}}(|x| \leq 2\epsilon)} \|e^{-itH_\alpha} \psi\|_{L_t^2 L_x^{10}} \\ &\quad + \|(V(x + x_n) - V(x))\|_{L_x^{\frac{3}{2}}(|x| > 2\epsilon)} \|e^{-itH_\alpha} \psi\|_{L_t^2 L_x^6} \\ &\lesssim \epsilon^{\frac{1}{5}} \|\psi\|_{\mathcal{H}_k^{\frac{10}{5}}} + |x_n| \epsilon^{-(\alpha-1)}, \end{aligned}$$

which implies that

$$\lim_{n \rightarrow +\infty} \|(e^{-itH_\alpha^n} - e^{-itH_\alpha^\infty})\psi\|_{L_t^\infty L_x^2} \lesssim \epsilon^{\frac{1}{5}}.$$

Since ϵ can be chosen arbitrarily,

$$\lim_{n \rightarrow +\infty} \|(e^{-itH_\alpha^n} - e^{-itH_\alpha^\infty})\psi\|_{L_t^\infty L_x^2} = 0.$$

Thus, we get (2.21) in the case $x_n \rightarrow \bar{x}$.

□

Once getting (2.20) and (2.21), we follow the same proof of Corollary 3.4 for $\alpha = 2$ in [13] and replace \dot{H}^1 and H_α^n with H^1 and $1 + H_\alpha^n$ in the procedure of the proof, respectively, to get the following decaying estimates.

Lemma 2.8. *Let $k > 0$, $1 < \alpha < 2$. Given $\psi \in H^1$, $t_n \rightarrow \pm\infty$ and any sequence $\{x_n\}_{n=1}^{+\infty} \subset \mathbf{R}^3$. We have*

$$(2.32) \quad \lim_{n \rightarrow +\infty} \|e^{-it_n H_\alpha^n} \psi\|_{L_x^6} = 0.$$

Moreover, if $\psi \in H^1$, then for $2 < p < \infty$,

$$(2.33) \quad \lim_{n \rightarrow +\infty} \|e^{-it_n H_\alpha^n} \psi\|_{L_x^p} = 0.$$

Using (2.21), Sobolev equivalence Lemma 2.1 and interpolation yields the following convergence (see the same result and the detailed proof for $\alpha = 2$ in [14]).

Lemma 2.9. *Let $x_n \rightarrow \bar{x} \in \mathbf{R}^3$ or $x_n \rightarrow \pm\infty$. Then for $\forall \psi \in H^1$, $k > 0$, $1 < \alpha < 2$,*

$$(2.34) \quad \lim_{n \rightarrow +\infty} \|(e^{-it_n H_\alpha^n} - e^{-it_n H_\alpha^\infty}) \psi\|_{L_{t,x}^2} = 0.$$

To get the parameters of linear profile decomposition are asymptotically orthogonal, we finally need two weak convergence results Lemma 2.10 and Lemma 2.11. Since they are the direct consequences of Lemma 2.7 and the detailed proof can be found in Lemma 3.8 and Lemma 3.9 for $\alpha = 2$ in [13] with a small modification by replacing \dot{H}^1 and Δ with H^1 and $1 + \Delta$, respectively, and using the inequality $\|\sqrt{H_\alpha^n} \psi_n\|_{L^2} \lesssim \|\psi_n\|_{H^1}$, where the implicit constant is independent of n , we omit their proof here.

Lemma 2.10. *Let $\psi_n \in H^1(\mathbf{R}^3)$ satisfy $\psi_n \rightharpoonup 0$ in $H^1(\mathbf{R}^3)$ and let $t_n \rightarrow \bar{t} \in \mathbf{R}$. Then for $k > 0$, $1 < \alpha \leq 2$,*

$$(2.35) \quad e^{-it_n H_\alpha^n} \psi_n \rightharpoonup 0 \text{ in } H^1(\mathbf{R}^3).$$

Lemma 2.11. *Let $\psi \in H^1(\mathbf{R}^3)$ and let $\{(t_n, y_n)\} \subset \mathbf{R} \times \mathbf{R}^3$, $|t_n| \rightarrow \infty$ or $|x_n| \rightarrow \infty$. Then for $k > 0$, $1 < \alpha \leq 2$,*

$$(2.36) \quad (e^{-it_n H_\alpha^n} \psi)(\cdot + y_n) \rightharpoonup 0 \text{ in } H^1(\mathbf{R}^3).$$

Use Lemma 2.7, Lemma 2.8, Lemma 2.10, Lemma 2.11 and Littlewood-Paley theory Lemma 2.4- Lemma 2.6 and follow the proof of Proposition 5.1 for $\alpha = 2$ in [14] to give the following linear profile decomposition. Similar to the above, we need to use \mathcal{H}_k^1 , $1 + H_\alpha^n$, $1 + H_\alpha^\infty$, and H^{-1} to replace $\dot{\mathcal{H}}_k^1$, H_α^n , H_α^∞ and \dot{H}^{-1} , respectively, in some appropriate places (e.g. in the proof of (2.44)).

Lemma 2.12. *Let $\{\phi_n\}_{n=1}^{+\infty}$ be a uniformly bounded sequence in $H^1(\mathbf{R}^3)$. Then there exist $M^* \in \mathbf{N} \cup \{+\infty\}$, a subsequence of $\{\phi_n\}_{n=1}^{M^*}$, which is denoted by itself, such that for $k > 0$, $1 < \alpha \leq 2$, the following statements hold.*

(1) *For each $1 \leq j \leq M \leq M^*$, there exist (fixed in n) a profile ψ^j in $H^1(\mathbf{R}^3)$, a sequence (in n) of time shifts t_n^j and a sequence (in n) of space shifts x_n^j , and there exists a sequence (in n) of*

remainder W_n^M in $H^1(\mathbf{R}^3)$ such that

$$(2.37) \quad \phi_n = \sum_{j=1}^M e^{it_n^j H_\alpha} \tau_{x_n^j} \psi^j + W_n^M := \sum_{j=1}^M \psi_n^j + W_n^M.$$

(2) For each $1 \leq j \leq M$,

$$(2.38) \quad \text{either } t_n^j = 0 \text{ for any } n \in \mathbf{N} \text{ or } \lim_{n \rightarrow +\infty} t_n^j = \pm\infty$$

$$(2.39) \quad \text{either } x_n^j = 0 \text{ for any } n \in \mathbf{N} \text{ or } \lim_{n \rightarrow +\infty} |x_n^j| = +\infty.$$

(3) The time and space sequence have a pairwise divergence property. Namely, for $1 \leq j \neq k \leq M$,

$$(2.40) \quad \lim_{n \rightarrow +\infty} (|t_n^j - t_n^k| + |x_n^j - x_n^k|) = +\infty.$$

(4) The remainder sequence has the following asymptotic smallness property and weak convergence property:

$$(2.41) \quad \lim_{M \rightarrow M^*} \overline{\lim}_{n \rightarrow +\infty} \|e^{-itH_\alpha} W_n^M\|_{L_{t,x}^5} = 0$$

and

$$(2.42) \quad \tau_{-x_n^M} e^{-it_n^M H_\alpha} W_n^M \rightharpoonup 0 \text{ in } H^1, \text{ as } n \rightarrow +\infty.$$

(5) For each fixed M , we have the asymptotic Pythagorean expansion as follows:

$$(2.43) \quad \|\phi_n\|_{L^2}^2 = \sum_{j=1}^M \|\psi^j\|_{L^2}^2 + \|W_n^M\|_{L^2}^2 + o_n(1),$$

$$(2.44) \quad \|\phi_n\|_{\mathcal{H}_k^1}^2 = \sum_{j=1}^M \|\tau_{x_n^j} \psi^j\|_{\mathcal{H}_k^1}^2 + \|W_n^M\|_{\mathcal{H}_k^1}^2 + o_n(1)$$

and

$$(2.45) \quad \|\phi_n\|_{L^4}^4 = \sum_{j=1}^M \|\psi_n^j\|_{L^4}^4 + \|W_n^M\|_{L^4}^4 + o_n(1),$$

where $o_n(1) \rightarrow 0$ as $n \rightarrow +\infty$. Specially,

$$(2.46) \quad S_k(\phi_n) = \sum_{j=1}^M S_k(\psi_n^j) + S_k(W_n^M) + o_n(1)$$

and

$$(2.47) \quad I_k(\phi_n) = \sum_{j=1}^M I_k(\psi_n^j) + I_k(W_n^M) + o_n(1).$$

Following the proof of Theorem 6.1 for $\alpha = 2$ in [14] and using Theorem 1.2, Lemma 2.3, Lemma 2.7 and Lemma 2.9, replacing $\frac{1}{|x|^2}$ and homogenous Sobolev spaces with $\frac{1}{|x|^\alpha}$ inhomogeneous Sobolev spaces, respectively, in some appropriate places gives the following lemma on nonlinear profiles when $|x_n| \rightarrow +\infty$.

Lemma 2.13. *Let $k > 0$, $1 < \alpha \leq 2$, $\psi \in H^1(\mathbf{R}^3)$ and the time sequence $t_n \equiv 0$ or $t_n \rightarrow \pm\infty$ such that*

$$(2.48) \quad \text{if } t_n \equiv 0, \quad S_0(\psi) < n_0 \quad \text{and} \quad P_0(\psi) \geq 0$$

and

$$(2.49) \quad \text{if } t_n \rightarrow \pm\infty, \quad \frac{1}{2}\|\psi\|_{H^1}^2 < n_0.$$

Let

$$\psi_n = e^{-it_n H_\alpha} \tau_{x_n} \psi,$$

where the space sequence x_n satisfies $|x_n| \rightarrow +\infty$. Then taking n large enough, we have that the solution $u(t) := NLS_k(t)\psi_n$ of (NLS_k) with initial data $u_0 = \psi_n$ is global and satisfies

$$\|NLS_k(t)\psi_n\|_{S_v^1(t)} \lesssim \|\psi\|_{H^1} \quad 1.$$

Moreover, for $\forall \epsilon > 0$, there exist a positive number $N = N(\epsilon)$ and a smooth compact supported function χ_ϵ on $\mathbf{R} \times \mathbf{R}^3$ satisfying

$$(2.50) \quad \|NLS_k(t)\psi_n(x) - \chi_\epsilon(t + t_n, x - x_n)\|_Z < \epsilon \quad \text{for } n \geq N,$$

where the norm

$$\|f\|_Z := \|f\|_{L_{t,x}^5} + \|f\|_{L_{t,x}^{\frac{10}{3}}} + \|f\|_{L_t^{\frac{30}{7}} L_x^{\frac{90}{31}}} + \|f\|_{L_t^{\frac{30}{7}} \mathcal{H}_k^{\frac{31}{60}, \frac{90}{31}}}$$

3. VARIATIONAL CHARACTERIZATION

We start with proving the positivity of $K_k^{a,b}$ near 0 in the $H^1(\mathbf{R}^3)$.

Lemma 3.1. *If the uniform L^2 -bounded sequence $\varphi_n \in H^1(\mathbf{R}^3) \setminus \{0\}$ satisfies $\lim_{n \rightarrow +\infty} \|\nabla \varphi_n\|_{L^2} = 0$, then for sufficiently large $n \in \mathbf{N}$, $K_k^{a,b}(\varphi_n) > 0$.*

Proof. By the fact that $-2V \leq x \cdot \nabla V \leq 0$ and $V \geq 0$, we always have that for large enough n

$$\begin{aligned} K_k^{a,b}(\varphi_n) &= \frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx + \frac{2a+3b}{2} \int_{\mathbf{R}^3} V|\varphi_n(x)|^2 dx + \frac{b}{2} \int_{\mathbf{R}^3} x \cdot \nabla V |\varphi_n(x)|^2 dx \\ &\quad + \frac{2a+3b}{2} \int_{\mathbf{R}^3} |\varphi_n(x)|^2 dx - \frac{4a+3b}{4} \int_{\mathbf{R}^3} |\varphi_n(x)|^4 dx \\ &\geq \frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx + \frac{2a+3b}{2} \int_{\mathbf{R}^3} |\varphi_n(x)|^2 dx - \frac{4a+3b}{4} \int_{\mathbf{R}^3} |\varphi_n(x)|^4 dx \\ &\geq \frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx - \frac{4a+3b}{4} \int_{\mathbf{R}^3} |\varphi_n(x)|^4 dx \\ &\geq \frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx - \frac{4a+3b}{4} c \left(\int_{\mathbf{R}^3} |\varphi_n(x)|^2 dx \right) \left(\int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx \right)^{\frac{3}{2}} \\ &\geq \frac{2a+b}{4} \int_{\mathbf{R}^3} |\nabla \varphi_n(x)|^2 dx > 0, \end{aligned}$$

where we have used the Gagliardo-Nirenberg inequality in the line before last and the assumptions $\lim_{n \rightarrow +\infty} \|\nabla \varphi_n\|_{L^2} = 0$ and uniform boundedness of $\|\varphi_n\|_{L^2}$ in the last line. \square

Simple computation gives

$$\|\nabla\varphi_\lambda^{a,b}\|_{L^2} = e^{\frac{1}{2}(2a+b)\lambda}\|\nabla\varphi\|_{L^2},$$

which implies that

$$(3.1) \quad \lim_{\lambda \rightarrow -\infty} \|\nabla\varphi_\lambda^{a,b}\|_{L^2} = 0.$$

So it follows from Lemma 3.1 that for sufficiently small $\lambda < 0$,

$$(3.2) \quad K_k^{a,b}(\varphi_\lambda^{a,b}) > 0.$$

For brevity, let $\bar{\mu} = 2a + b$ and $\underline{\mu} = 2a + 3b$. Next we introduce the functional

$$J_k^{a,b}(\varphi) := \frac{1}{\bar{\mu}}(\bar{\mu} - \mathcal{L}^{a,b})S_k(\varphi) = \frac{1}{\bar{\mu}}(\bar{\mu}S_k(\varphi) - K_k^{a,b}(\varphi)).$$

The lemma shows that the positivity of $J_k^{a,b}(\varphi)$ and the monotonicity of $J_k^{a,b}(\varphi_\lambda^{a,b})$ in λ .

Lemma 3.2. *For any $\varphi \in H^1(\mathbf{R}^3)$, we have the following two identities:*

$$(3.3) \quad \begin{aligned} \bar{\mu}J_k^{a,b}(\varphi) &= (\bar{\mu} - \mathcal{L}^{a,b})S_k^{a,b}(\varphi) \\ &= -b \int_{\mathbf{R}^3} V|\varphi|^2 dx - \frac{b}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V)|\varphi|^2 dx - b\|\varphi\|_{L^2}^2 + \frac{a+b}{2}\|\varphi\|_{L^4}^4. \end{aligned}$$

and

$$(3.4) \quad (\mathcal{L}^{a,b} - \bar{\mu})(\underline{\mu} - \mathcal{L}^{a,b})S_k^{a,b}(\varphi) = b^2 \int_{\mathbf{R}^3} (-3x \cdot \nabla V - x\nabla^2 V x^T)|\varphi|^2 dx + (a+b)a\|\varphi\|_{L^4}^4,$$

where $\nabla^2 V$ is Hessian matrix of V .

Proof. By simple computations, we have

$$\mathcal{L}^{a,b}\|\nabla\varphi\|_{L^2}^2 = \bar{\mu}\|\nabla\varphi\|_{L^2}^2, \quad \mathcal{L}^{a,b}\|\varphi\|_{L^2}^2 = \underline{\mu}\|\varphi\|_{L^2}^2, \quad \mathcal{L}^{a,b}\|\varphi\|_{L^4}^4 = (4a+3b)\|\varphi\|_{L^4}^4,$$

$$\mathcal{L}^{a,b}\left(\int_{\mathbf{R}^3} V|\varphi|^2 dx\right) = \underline{\mu} \int_{\mathbf{R}^3} V|\varphi|^2 dx + b \int_{\mathbf{R}^3} (x \cdot \nabla V)|\varphi|^2 dx,$$

and

$$\mathcal{L}^{a,b}\left(\int_{\mathbf{R}^3} (x \cdot \nabla V)|\varphi|^2 dx\right) = 2(a+b) \int_{\mathbf{R}^3} x \cdot \nabla V|\varphi|^2 dx + b \int_{\mathbf{R}^3} x\nabla^2 V x^T|\varphi|^2 dx,$$

which imply that (3.3) and (3.4). We conclude the proof. \square

Remark 3.3. *By the definition of V , we have $2V + x \cdot V \geq 0$ and $3x \cdot \nabla V + x\nabla^2 V x^T \leq 0$, which together with (3.3) and (3.4) yield that $J_k^{a,b}(\varphi) > 0$ and $\mathcal{L}^{a,b}J_k^{a,b}(\varphi) \geq 0$ for any $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ which implies that the function $\lambda \mapsto J_k^{a,b}(\varphi_\lambda^{a,b})$ is increasing.*

Using the functional $K_k^{a,b}$ with (a, b) satisfying (1.10), we introduce a general minimizing problem:

$$(3.5) \quad n_k^{a,b} = \inf\{S_k(\varphi) : \varphi \in H^1(\mathbf{R}^3) \setminus \{0\}, K_k^{a,b}(\varphi) = 0\}.$$

In particular, when $(a, b) = (3, -2)$, it is namely n_k in (1.14). In fact, we shall show that

$$(3.6) \quad n_k^{a,b} = n_0^{a,b} = n_k = n_0.$$

To this end, we introduce another one general minimizing problem in terms of the positive functional $J_0^{a,b}$ for any a, b satisfying (1.10):

$$(3.7) \quad j^{a,b} = \inf\{J_0^{a,b}(\varphi) : \varphi \in H^1(\mathbf{R}^3) \setminus \{0\}, K_0^{a,b}(\varphi) \leq 0\}.$$

In the following lemma, we shall show a relation between the two minimizers $n_0^{a,b}$ in (3.5) (i.e., $k = 0$) and $j^{a,b}$ in (3.7).

Lemma 3.4.

$$(3.8) \quad j^{a,b} = n_0^{a,b}.$$

Proof. On one hand, for any $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ with $K_0^{a,b}(\varphi) \leq 0$, there are two possibilities: $K_0^{a,b}(\varphi) = 0$ and $K_0^{a,b}(\varphi) < 0$. When $K_0^{a,b}(\varphi) = 0$, $S_0(\varphi) = J_0^{a,b}(\varphi)$ by the definition of $J_k^{a,b}$. Hence,

$$(3.9) \quad n_0^{a,b} \leq J_0^{a,b}(\varphi).$$

When $K_0^{a,b}(\varphi) = K_0^{a,b}(\varphi_0^{a,b}) < 0$, using the continuity of $K_0^{a,b}(\varphi_\lambda^{a,b})$ in λ and the fact that for sufficiently small $\lambda < 0$ such that $K_0^{a,b}(\varphi_\lambda^{a,b}) > 0$ holds by (3.2) yields that there exists a $\lambda_0 < 0$ such that $K_0^{a,b}(\varphi_{\lambda_0}^{a,b}) = 0$. Using Remark 3.3, we get

$$S_0(\varphi_{\lambda_0}^{a,b}) = J_0^{a,b}(\varphi_{\lambda_0}^{a,b}) \leq J_0^{a,b}(\varphi_0^{a,b}) = J_0^{a,b}(\varphi)$$

Hence, we still have (3.9). Altogether, for any $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ with $K_0^{a,b}(\varphi) \leq 0$, we have (3.9). By the definition of $j^{a,b}$, we have $n_0^{a,b} \leq j^{a,b}$.

On the other hand, for any $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ with $K_0^{a,b}(\varphi) = 0$. Of course, $K_0^{a,b}(\varphi) \leq 0$ and $S_0(\varphi) = J_0^{a,b}(\varphi)$, which implies that $j^{a,b} \leq n_0^{a,b}$. Thus, we complete the proof. \square

Next we shall apply Lemma 3.4 to get $n_k^{a,b} = n_0^{a,b}$.

Lemma 3.5.

$$(3.10) \quad n_k^{a,b} = n_0^{a,b}.$$

Proof. On one hand, for any $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ with $K_k^{a,b}(\varphi) = 0$, we have $S(\varphi) = J^{a,b}(\varphi)$ by the definition of $J_k^{a,b}$. Since $V \geq 0$ and $2V + x \cdot \nabla V \geq 0$, we get $K_0^{a,b}(\varphi) \leq K_k^{a,b}(\varphi) = 0$ and $J_0^{a,b}(\varphi) \leq J_k^{a,b}(\varphi)$, which together with (3.8) implies that

$$n_0^{a,b} = j^{a,b} \leq J_k^{a,b}(\varphi).$$

Also as $K^{a,b}(\varphi) = 0$, taking infimum on both sides of the above inequality yields that

$$n_0^{a,b} \leq n_k^{a,b}.$$

On the other hand, we review that Q is the positive radial exponential decaying solution of the elliptic equation (1.6), so there exists a constant c such that $Q(x) \leq e^{-c|x|}$ for any $x \in \mathbf{R}^3$ (e.g. See Theorem 8.1.1 in [3]). Here we only need the decaying property of Q , not necessarily exponential decaying). Let x_n be a sequence satisfying $|x_n| \rightarrow +\infty$. Hence, for any given $R > 0$, there exists $N = N(R) > 0$, for any $n \geq N$, we have $|x_n| \geq 2R$. we claim that

$$(3.11) \quad \int_{\mathbf{R}^3} V(x)|Q(x-x_n)|^2 dx \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Indeed,

$$\begin{aligned} \int_{\mathbf{R}^3} V(x)|Q(x-x_n)|^2 dx &\leq \int_{|x|\leq R} V(x)|Q(x-x_n)|^2 dx + \int_{|x|>R} V(x)|Q(x-x_n)|^2 dx \\ &:= I_1 + I_2. \end{aligned}$$

For the first part I_1 , when $n \geq N$, $|x-x_n| \geq \frac{|x_n|}{2}$ and then

$$|Q(x-x_n)| \leq \sup_{|x|\geq \frac{|x_n|}{2}} |Q(x)| := C(|x_n|) \rightarrow 0$$

as n tends to $+\infty$. Therefore,

$$I_1 \lesssim C(|x_n|) \int_{|x|\leq R} V(x) dx \lesssim R^{(3-\alpha)} C(|x_n|).$$

For the second part I_2 , $V(x) \lesssim R^{-\alpha}$, which implies that

$$I_2 \lesssim R^{-\alpha} \int_{\mathbf{R}^3} |Q(x)|^2 dx.$$

Combining the above two parts yields that the claim (3.11) holds true. Noticing that $x \cdot \nabla V = (-\alpha)V$, thus we also have

$$(3.12) \quad \int_{\mathbf{R}^3} (x \cdot \nabla V) |Q(x-x_n)|^2 dx \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Using the definition of $K_k^{a,b}$ (1.11) and $K_0^{a,b}(Q) = 0$ and taking θ sufficiently large, we have

$$K_k^{a,b}(\tau_{x_n} Q) > K_0^{a,b}(\tau_{x_n} Q) = K_0^{a,b}(Q) = 0$$

and

$$K_k^{a,b}(\theta \tau_{x_n} Q) < 0,$$

from which it follows that there must be $\theta_n > 1$ satisfying $K_k^{a,b}(\theta_n \tau_{x_n} Q) = 0$. We claim that

$$(3.13) \quad \lim_{n \rightarrow +\infty} \theta_n = 1.$$

In fact, by $K_0^{a,b}(Q) = 0$, we have

$$\frac{2a+b}{2} \int_{\mathbf{R}^3} |\nabla Q(x)|^2 dx + \frac{2a+3b}{2} \int_{\mathbf{R}^3} |Q(x)|^2 dx = \frac{4a+3b}{4} \int_{\mathbf{R}^3} |Q(x)|^4 dx,$$

which implies that

$$\begin{aligned} K_k^{a,b}(\theta_n \tau_{x_n} Q) &= \theta_n^2 \left[\frac{2a+3b}{2} \int_{\mathbf{R}^3} V |Q(x-x_n)|^2 dx + \frac{b}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |Q(x-x_n)|^2 dx \right. \\ &\quad \left. + \frac{4a+3b}{4} (1-\theta_n^2) \int_{\mathbf{R}^3} |Q(x)|^4 dx \right] = 0. \end{aligned}$$

Hence,

$$\frac{2a+3b}{2} \int_{\mathbf{R}^3} V |Q(x-x_n)|^2 dx + \frac{b}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |Q(x-x_n)|^2 dx + \frac{4a+3b}{4} (1-\theta_n^2) \int_{\mathbf{R}^3} |Q(x)|^4 dx = 0.$$

Taking limit in the above quality and using (3.11) and (3.12) gives the claim (3.13).

Using (3.11) and (3.13) yields that

$$\lim_{n \rightarrow +\infty} S_k(\theta_n \tau_{x_n} Q) = S_0(Q) = n_0^{a,b},$$

which together with $K_k^{a,b}(\theta_n \tau_{x_n} Q) = 0$ implies that

$$n_k^{a,b} \leq n_0^{a,b}.$$

We conclude the proof. \square

Lemma 3.5 shows that $n_k^{a,b} = n_0^{a,b} = S_0(Q)$, so $n_k^{a,b}$ and $n_0^{a,b}$ don't depend on the parameters a and b . Therefore, (3.6) holds true. It is known that n_0 is attained by Q . However, the following lemma shows that n_k is never attained for any $k > 0$.

Lemma 3.6. *n_k is never attained for any $k > 0$.*

Proof. Suppose that there exists a $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$ such that n_k is attained by φ . Namely, $P_k(\varphi) = 0$ and $S_k(\varphi) = n_k$. As $\varphi \in H^1(\mathbf{R}^3) \setminus \{0\}$, $\lim_{|x| \rightarrow +\infty} \varphi(x) = 0$. Following the proof of (3.11), we have

$$(3.14) \quad \int_{\mathbf{R}^3} V(x)|\varphi(x - x_n)|^2 dx \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

where $|x_n| \rightarrow +\infty$ as $n \rightarrow +\infty$. Hence, we have

$$- \int_{\mathbf{R}^3} (x \cdot \nabla V)|\tau_{x_n} \varphi|^2 dx \quad \text{is positive and tends to zero as } n \rightarrow +\infty$$

and

$$2 \int_{\mathbf{R}^3} V|\tau_{x_n} \varphi|^2 dx + \int_{\mathbf{R}^3} (x \cdot \nabla V)|\tau_{x_n} \varphi|^2 dx \quad \text{is positive and tends to zero as } n \rightarrow +\infty.$$

Therefore, for n sufficiently large, we have

$$(3.15) \quad P_k(\tau_{x_n} \varphi) < P_k(\varphi) = 0$$

and

$$(3.16) \quad J_k^{3,-2}(\tau_{x_n} \varphi) < J_k^{3,-2}(\varphi) = S_k(\varphi) = n_k.$$

By (3.2), for sufficiently small $\lambda < 0$, we have $P_k((\tau_{x_n} \varphi)_\lambda^{3,-2}) > 0$, which combined with (3.15) implies that there exists a $\lambda_0 < 0$ such that $P_k((\tau_{x_n} \varphi)_{\lambda_0}^{3,-2}) = 0$. Using Remark 3.3 and (3.16), we get

$$n_k = S_k((\tau_{x_n} \varphi)_{\lambda_0}^{3,-2}) = J_k^{3,-2}((\tau_{x_n} \varphi)_{\lambda_0}^{3,-2}) < J_k^{3,-2}((\tau_{x_n} \varphi)_0^{3,-2}) = J_k^{3,-2}(\tau_{x_n} \varphi) < n_k,$$

which is impossible. Thus we complete the proof. \square

To get the fact that $P_k(\varphi)$ in (1.12) and $I_k(\varphi)$ in (1.13) have the same sign under the condition $S_k(\varphi) < n_0$ with $\varphi \in H^1(\mathbf{R}^3)$, we introduce $\mathcal{N}_{a,b}^\pm \subset H^1(\mathbf{R}^3)$ defined by

$$(3.17) \quad \mathcal{N}_{a,b}^+ := \{\varphi \in H^1(\mathbf{R}^3) : S_k(\varphi) < n_0, K_k^{a,b}(\varphi) \geq 0\}$$

and

$$(3.18) \quad \mathcal{N}_{a,b}^- := \{\varphi \in H^1(\mathbf{R}^3) : S_k(\varphi) < n_0, K_k^{a,b}(\varphi) < 0\}.$$

It is easy to see that $\mathcal{N}_{3,-2}^\pm = \mathcal{N}^\pm$ in (1.15) and (1.16). The above fact can be obtained if we show that $\mathcal{N}_{a,b}^\pm$ are independent of (a, b) (i.e., $\mathcal{N}_{a,b}^\pm = \mathcal{N}^\pm$), which follows from the contractivity of $\mathcal{N}_{a,b}^+$.

Lemma 3.7. *Let (a, b) satisfy (1.10), then $\mathcal{N}_{a,b}^\pm$ are independent of (a, b) .*

Proof. The proof is similar to the one of Lemma 2.9 in [9] (see also Lemma 2.15 in [10]), so we omit it. \square

The following lemma shows that for any element φ in \mathcal{N}^+ , $S_k(\varphi) \sim \|\varphi\|_{\mathcal{H}_k^1} \sim \|\varphi\|_{H^1}$.

Lemma 3.8. *Let $\varphi \in \mathcal{N}^+$, then*

$$(3.19) \quad \frac{1}{4}\|\varphi\|_{\mathcal{H}_k^1}^2 \leq S_k(\varphi) \leq \frac{1}{2}\|\varphi\|_{\mathcal{H}_k^1}^2.$$

Proof. By Lemma 3.7, we have $I_k(\varphi) \geq 0$, which implies that

$$\int_{\mathbf{R}^3} |\varphi(x)|^4 dx \leq \|\varphi\|_{\mathcal{H}_k^1}^2.$$

Thus, we have

$$\frac{1}{2}\|\varphi\|_{\mathcal{H}_k^1}^2 \geq S_k(\varphi) = \frac{1}{2}\|\varphi\|_{\mathcal{H}_k^1}^2 - \frac{1}{4} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \geq \frac{1}{4}\|\varphi\|_{\mathcal{H}_k^1}^2,$$

which is namely (3.19). We complete the proof. \square

Finally, we obtain the corresponding uniform bounds on $P_k(\varphi)$ when $\varphi \in \mathcal{N}^\pm$, which plays a vital role in the proof of Theorem 1.3.

Lemma 3.9. *1. Let $\varphi \in \mathcal{N}^-$, then*

$$(3.20) \quad P_k(\varphi) \leq -4(n_0 - S_k(\varphi))$$

2. Let $\varphi \in \mathcal{N}^+$, then

$$(3.21) \quad P_k(\varphi) \geq \min \left\{ 4(n_0 - S_k(\varphi)), \frac{2}{5} \left(\|\nabla \varphi\|_{L^2}^2 - \frac{1}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |\varphi(x)|^2 dx \right) \right\}.$$

Proof. For any $\varphi \in H^1(\mathbf{R}^3)$, define $s(\lambda) := S_k(\varphi_\lambda^{3,-2})$, then

$$(3.22) \quad \begin{aligned} s(\lambda) &= \frac{1}{2} e^{4\lambda} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx + \frac{1}{2} \int_{\mathbf{R}^3} V(e^{-2\lambda} x) |\varphi(x)|^2 dx \\ &\quad + \frac{1}{2} \int_{\mathbf{R}^3} |\varphi(x)|^2 dx - \frac{1}{4} e^{6\lambda} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx, \\ s'(\lambda) &= P_k(\varphi_\lambda^{3,-2}) = 2e^{4\lambda} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx - e^{-2\lambda} \int_{\mathbf{R}^3} [x \cdot (\nabla V)(e^{-2\lambda} x)] |\varphi(x)|^2 dx \\ &\quad - \frac{3}{2} e^{6\lambda} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \end{aligned}$$

$$(3.23) \quad \begin{aligned} &\geq 2e^{4\lambda} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx - \frac{3}{2} e^{6\lambda} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx, \\ s''(\lambda) &= 8e^{4\lambda} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx + 2e^{-2\lambda} \int_{\mathbf{R}^3} [x \cdot (\nabla V)(e^{-2\lambda} x)] |\varphi(x)|^2 dx \\ &\quad + 2e^{-4\lambda} \int_{\mathbf{R}^3} [x(\nabla^2 V)(e^{-2\lambda} x)x^T] |\varphi(x)|^2 dx - 9e^{6\lambda} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \\ &\leq 8e^{4\lambda} \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx - 4e^{-2\lambda} \int_{\mathbf{R}^3} [x \cdot (\nabla V)(e^{-2\lambda} x)] |\varphi(x)|^2 dx \end{aligned}$$

$$\begin{aligned}
& -9e^{6\lambda} \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \\
& = 4s'(\lambda) - 3 \int_{\mathbf{R}^3} |\varphi(x)|^4 dx \leq 4s'(\lambda),
\end{aligned}$$

where we have used the inequalities $x \cdot \nabla V$ and $3x \cdot \nabla V + x \nabla^2 V x^T \leq 0$.

1. If $\varphi \in \mathcal{N}^-$, then it follows from (3.22) that $s'(0) = P_k(\varphi) < 0$ and $s'(\lambda) > 0$ for sufficiently small $\lambda < 0$. Thus, by the continuity of $s'(\lambda) = P_k(\varphi_{\lambda}^{3,-2})$ in λ , there exists a negative $\lambda_0 < 0$ such that

$$s'(\lambda_0) = P_k(\varphi_{\lambda_0}^{3,-2}) = 0 \quad \text{and} \quad s'(\lambda) < 0, \quad \text{for } \forall \lambda \in (\lambda_0, 0].$$

By the definition of n_0 (1.14), $s(\lambda_0) = S_k(\varphi_{\lambda_0}^{3,-2}) \geq n_0$. Integrating the inequality (3.23) over $[\lambda_0, 0]$ yields that

$$P_k(\varphi) = s'(0) = s'(0) - s'(\lambda_0) \leq 4(s(0) - s(\lambda_0)) \leq 4(S_k(\varphi) - n_0) = -4(n_0 - S_k(\varphi)).$$

Thus, we complete the proof of (3.20).

2. If $\varphi \in \mathcal{N}^+$, we consider two cases: one is $8P_k(\varphi) \geq 3\|\varphi\|_{L^4}^4$ and the other is $8P_k(\varphi) < 3\|\varphi\|_{L^4}^4$.

For the case $8P_k(\varphi) \geq 3\|\varphi\|_{L^4}^4$, it follows from the definition of P_k (1.12) that

$$2P_k(\varphi) = 4 \int_{\mathbf{R}^3} |\nabla \varphi(x)|^2 dx - 2 \int_{\mathbf{R}^3} (x \cdot \nabla V) |\varphi(x)|^2 dx - 3 \int_{\mathbf{R}^3} |\varphi(x)|^4 dx,$$

and then we have

$$10P_k(\varphi) \geq 4(\|\nabla \varphi\|_{L^2}^2 - \frac{1}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |\varphi(x)|^2 dx),$$

that is,

$$(3.24) \quad P_k(\varphi) \geq \frac{2}{5} \left(\|\nabla \varphi\|_{L^2}^2 - \frac{1}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |\varphi(x)|^2 dx \right).$$

For the other case $8P_k(\varphi) < 3\|\varphi\|_{L^4}^4$, by (3.23), we have

$$(3.25) \quad 0 < 8s'(\lambda) < 3e^{6\lambda} \|\varphi\|_{L^4}^4 \quad \text{and then} \quad s''(\lambda) \leq 4s'(\lambda) - 3e^{6\lambda} \|\varphi\|_{L^4}^4 < -4s'(\lambda)$$

at $\lambda = 0$. Also as $s'(\lambda)$ and $s''(\lambda)$ are continuous, $s'(\lambda)$ decreases as s increases until $s'(\lambda_1) = 0$ for some $0 < \lambda_1 < +\infty$ and (3.25) is true over $[0, \lambda_1]$. Since $P_k(\varphi_{\lambda_1}^{3,-2}) = s'(\lambda_1) = 0$, by the definition of n_0 (1.14), $s(\lambda_1) = S_k(\varphi_{\lambda_1}^{3,-2}) \geq n_0$. Integrating the second inequality in (3.25) over $[0, \lambda_1]$, we have

$$-P_k(\varphi) = s'(\lambda_1) - s'(0) < -4(s(\lambda_1) - s(0)) \leq -4(n_0 - S_k(\varphi)),$$

which is

$$(3.26) \quad P_k(\varphi) \geq 4(n_0 - S_k(\varphi)).$$

Putting (3.24) and (3.26) together yields (3.21).

□

4. CRITERIA FOR GLOBAL WELL-POSEDNESS AND BLOW-UP

In this section, we will give the criteria for global well-posedness and blow-up for the solution u of (NLS_k) , which are partial results of Theorem 1.3. The proof of blow-up part is based on the argument of [4].

Theorem 4.1. *Let u be the solution of (NLS_k) on $(-T_{\min}, T_{\max})$, where $(-T_{\min}, T_{\max})$ is the maximal life-span.*

(i) *If $u_0 \in \mathcal{N}^+$, then u is global well-posedness and $u(t) \in \mathcal{N}^+$ for any $t \in \mathbf{R}$.*

(ii) *If $u_0 \in \mathcal{N}^-$, then $u(t) \in \mathcal{N}^-$ for any $t \in (-T_{\min}, T_{\max})$ and one of the following four statements holds true:*

(a) *$T_{\max} < +\infty$ and $\lim_{t \uparrow T_{\max}} \|\nabla u(t)\|_{L^2} = +\infty$.*

(b) *$T_{\min} < +\infty$ and $\lim_{t \downarrow -T_{\min}} \|\nabla u(t)\|_{L^2} = +\infty$.*

(c) *$T_{\max} = +\infty$ and there exists a sequence $\{t_n\}_{n=1}^{+\infty}$ such that $t_n \rightarrow +\infty$ and $\lim_{t_n \uparrow +\infty} \|\nabla u(t)\|_{L^2} = +\infty$.*

(d) *$T_{\min} = +\infty$ and there exists a sequence $\{t_n\}_{n=1}^{+\infty}$ such that $t_n \rightarrow -\infty$ and $\lim_{t_n \downarrow -\infty} \|\nabla u(t)\|_{L^2} = +\infty$.*

Proof. (i) Define

$$I^+ = \{t \in (-T_{\min}, T_{\max}) : u(t) \in \mathcal{N}^+\}.$$

Obviously, $0 \in I^+ \neq \emptyset$. On one hand, since $S_k(u(t)) = S_k(u_0) < n_0$ and $P_k(u(t))$ is continuous in t , I^+ is a closed subset of $(-T_{\min}, T_{\max})$. On the other hand, by (3.21), we further obtain that, I^+ is an open subset of $(-T_{\min}, T_{\max})$. Therefore, $I^+ = (-T_{\min}, T_{\max})$. Namely, for any $t \in (-T_{\min}, T_{\max})$, $u(t) \in \mathcal{N}^+$. It follows from (3.19) that for any $t \in (-T_{\min}, T_{\max})$,

$$\|u(t)\|_{H^1}^2 \leq \|u(t)\|_{\mathcal{H}_k^1}^2 \leq 4S_k(u(t)) \leq 4n_0.$$

So by local well-posedness result (i) of Theorem 1.1, we have $(-T_{\min}, T_{\max}) = \mathbf{R}$, which implies that u is global well-posedness and $u(t) \in \mathcal{N}^+$ for any $t \in \mathbf{R}$. Thus, we complete the proof of (i).

(ii) Similarly above, we can show that $u(t) \in \mathcal{N}^-$ for any $t \in (-T_{\min}, T_{\max})$ by replacing (3.21) with (3.20).

In the sequel, we only consider the positive time because the negative time can be dealt with similarly. If $T_{\max} < +\infty$, we naturally have $\lim_{t \uparrow T_{\max}} \|\nabla u(t)\|_{L^2} = +\infty$. If $T_{\max} = +\infty$, we shall prove $\lim_{t \uparrow +\infty} \|\nabla u(t)\|_{L^2} = +\infty$ by contradiction. Assume we have

$$C_0 = \sup_{t \in \mathbf{R}^+} \|\nabla u(t)\|_{L^2} < \infty.$$

Consider the localized Virial identity and define

$$(4.1) \quad I(t) := \int_{\mathbf{R}^3} \phi(x) |u(t, x)|^2 dx,$$

then by straight computations, we obtain that for any $\phi \in C^4(\mathbf{R}^3)$ (e.g., see Proposition 7.1 in [8])

$$I'(t) = 2 \operatorname{Im} \int_{\mathbf{R}^3} \nabla \phi \cdot \nabla u \bar{u} dx;$$

$$I''(t) = \int_{\mathbf{R}^3} 4 \operatorname{Re} \nabla \bar{u} \nabla^2 \phi \nabla u dx - \int_{\mathbf{R}^3} 2 \nabla \phi \cdot \nabla V |u|^2 + \Delta \phi |u|^4 dx - \int_{\mathbf{R}^3} \Delta^2 \phi |u|^2 dx.$$

In particular, if ϕ is a radial function ,

$$(4.2) \quad I'(t) = 2 \operatorname{Im} \int_{\mathbf{R}^3} \phi'(r) \frac{x \cdot \nabla u}{r} \bar{u} dx,$$

$$(4.3) \quad \begin{aligned} I''(t) &= 4 \int_{\mathbf{R}^3} \frac{\phi'}{r} |\nabla u|^2 dx + 4 \int_{\mathbf{R}^3} \left(\frac{\phi''}{r^2} - \frac{\phi'}{r^3} \right) |x \cdot \nabla u|^2 dx \\ &\quad - 2 \int_{\mathbf{R}^3} \frac{\phi'}{r} x \cdot \nabla V |u|^2 dx - \int_{\mathbf{R}^3} \left(\phi''(r) + \frac{2}{r} \phi'(r) \right) |u|^4 dx - \int_{\mathbf{R}^3} \Delta^2 \phi |u|^2 dx. \end{aligned}$$

L^2 estimate in the exterior ball

Given $R \gg 1$, which will be determined later. Take ϕ in (4.1) such that

$$\phi = \begin{cases} 0, & 0 \leq r \leq \frac{R}{2}; \\ 1, & r \geq R, \end{cases}$$

and

$$0 \leq \phi \leq 1, \quad 0 \leq \phi' \leq \frac{4}{R}.$$

By (4.2) and Hölder inequality, there holds that

$$\begin{aligned} I(t) &= I(0) + \int_0^t I'(\tau) d\tau \leq I(0) + t \|\phi'\|_{L^\infty} M(u_0) C_0 \\ &\leq \int_{|x| \geq \frac{R}{2}} |u_0|^2 dx + \frac{4M(u_0)C_0 t}{R}. \end{aligned}$$

Note that

$$\int_{|x| \geq \frac{R}{2}} |u_0|^2 dx = o_R(1),$$

and

$$\int_{|x| \geq R} |u(t, x)|^2 dx \leq I(t).$$

So for given $\eta_0 > 0$, if

$$t \leq \frac{\eta_0 R}{4M(u_0)C_0},$$

then we have that

$$(4.4) \quad \int_{|x| \geq \frac{R}{2}} |u(t, x)|^2 dx \leq \eta_0 + o_R(1).$$

Localized Virial identity

$I''(t)$ can be rewritten as

$$(4.5) \quad I''(t) = 4P_k(u) + R_1 + R_2 + R_3 + R_4,$$

where

$$\begin{aligned} R_1 &= 4 \int_{\mathbf{R}^3} \left(\frac{\phi'}{r} - 2 \right) |\nabla u|^2 dx + 4 \int_{\mathbf{R}^3} \left(\frac{\phi''}{r^2} - \frac{\phi'}{r^3} \right) |x \cdot \nabla u|^2 dx, \\ R_2 &= - \int_{\mathbf{R}^3} \left(\phi'' + \frac{2}{r} \phi'(r) - 6 \right) |u|^4 dx, \end{aligned}$$

$$R_3 = -2 \int_{\mathbf{R}^3} \left(\frac{\phi'}{r} - 2 \right) (x \cdot \nabla V) |u|^2 dx,$$

$$R_4 = - \int_{\mathbf{R}^3} \Delta^2 \phi |u|^2 dx.$$

At this stage, we choose another radial function ϕ such that

$$(4.6) \quad 0 \leq \phi \leq r^2, \quad \phi'' \leq 2, \quad \phi^{(4)} \leq \frac{4}{R^2}, \quad \text{and} \quad \phi = \begin{cases} r^2, & 0 \leq r \leq R; \\ 0, & r \geq 2R \end{cases}.$$

First, note that $R_1 \leq 0$. If $\phi'' \leq r^{-1}\phi' \leq 0$, by $\phi' \leq 2r$, it is easy to see that $R_1 \leq 0$. If $\phi'' \leq r^{-1}\phi' \leq 0$, by $\phi'' \leq 2$, it holds that

$$R_1 \leq 4 \int_{\mathbf{R}^3} \left(\frac{\phi'}{r} - 2 \right) |\nabla u|^2 dx + 4 \int_{\mathbf{R}^3} \left(\phi'' - \frac{\phi'}{r} \right) |\nabla u|^2 dx = 4 \int_{\mathbf{R}^3} (\phi'' - 2) |\nabla u|^2 dx \leq 0.$$

Secondly, let $\chi^4(r) = |\phi'' + \frac{2}{r}\phi'(r) - 6|$. it is easy to see that $\text{supp}\chi \subset [R, \infty)$. So by Gagliardo-Nirenberg inequality

$$R_2 \leq \int_{\mathbf{R}^3} \chi^4 |u|^4 dx \lesssim \|\nabla(\chi u)\|_{L^2}^3 \|\chi u\|_{L^2} \leq C(M(u_0), C_0) \|u\|_{L^2(|x|>R)}^{\frac{1}{2}}.$$

By the properties of ϕ ,

$$R_4 \leq CR^{-2} \|u\|_{L^2(|x|>R)}^2.$$

Finally, by $x \cdot \nabla V \leq 0$, and we obtain $R_3 \leq 0$.

Putting all the above estimates together, there holds that for $R \gg 1$,

$$(4.7) \quad I''(t) \leq 4P_k(u) + \tilde{C} \|u\|_{L^2(|x|>R)},$$

where $\tilde{C} > 0$ depending on $M(u_0)$ and C_0 .

Using (4.4) and (4.7) yields that $t \leq T := \eta_0 R / (4M(u_0)C_0)$,

$$I''(t) \leq 4P_k(u) + \tilde{C}(\eta_0^{1/2} + o_R(1)).$$

As $u(t) \in \mathcal{N}^-$, it follows from (3.20) that there exists

$$\beta_0 := -4(n_0 - S_k(u(t))) = -4(n_0 - S_k(u_0)) < 0$$

such that

$$(4.8) \quad I''(t) \leq 4\beta_0 + \tilde{C}(\eta_0^{1/2} + o_R(1)).$$

Choose η_0 sufficiently small and take R sufficiently large such that

$$4\beta_0 + \tilde{C}\eta_0^{1/2} + o_R(1) < \beta_0.$$

Integrating (4.8) over $[0, T]$ twice, we obtain that

$$I(T) \leq I(0) + I'(0)T + \int_0^T \int_0^t \beta_0$$

$$\leq I(0) + I'(0)T + \beta_0 \frac{T^2}{2}.$$

Hence for $T = \eta_0 R / (4M(u_0)C_0)$, we obtain that

$$(4.9) \quad I(T) \leq I(0) + I'(0)\eta_0 R / (4M(u_0)C_0) + \alpha_0 R^2,$$

where the constant

$$\alpha_0 = \beta_0 \eta_0^2 / (4M(u_0)C_0)^2 < 0$$

is independent of R .

At the same time, we note that

$$(4.10) \quad I(0) = o_R(1)R^2, \quad I'(0) = o_R(1)R.$$

In fact,

$$\begin{aligned} I(0) &\leq \int_{|x| < \sqrt{R}} |x|^2 |u_0|^2 dx + \int_{\sqrt{R} < |x| < 2R} |x|^2 |u_0|^2 dx \\ &\leq RM(u_0) + R^2 \int_{|x| > \sqrt{R}} |u_0|^2 dx = o_R(1)R^2. \end{aligned}$$

Similarly, we obtain the second estimate and then prove (4.10).

Putting (4.9) with (4.10) together and choosing R sufficiently enough, we find that

$$I(T) \leq (o_R(1) + \alpha_0)R^2 \leq \frac{1}{4}\alpha_0R^2 < 0,$$

which is impossible since $I \geq 0$. Thus, we conclude the proof of blow-up part. \square

5. SCATTERING RESULT

In this section, we shall show the remaining part of Theorem 1.3. In the previous section, we have obtained that the solution $u(t)$ of (NLS_k) is global and belongs to \mathcal{N}^+ if $u_0 \in \mathcal{N}^+$. To get scattering result, by (iv) of Theorem 1.1, it's enough to get (1.3). To this end, we introduce a definition.

Definition 5.1. *We say that $SC(u_0)$ holds if for $u_0 \in H^1(\mathbf{R}^3)$ satisfying $u_0 \in \mathcal{N}^+$, the corresponding global solution u of (NLS_k) satisfies (1.3).*

We first note that for $u_0 \in \mathcal{N}^+$, there exists $\delta > 0$ such that if $S_k(u_0) < \delta$, then (1.3) holds. In fact, by (3.19), $\|u_0\|_{H^1} \lesssim S_k(u_0)$. Therefore, by (ii) of Theorem 1.1, taking $\delta > 0$ sufficiently small gives (1.3).

Now for each $\delta > 0$, we define the set S_δ as follows:

$$(5.1) \quad S_\delta = \{u_0 \in H^1(\mathbf{R}^3) : S_k(u_0) < \delta \text{ and } u_0 \in \mathcal{N}^+ \Rightarrow (1.3) \text{ holds}\}.$$

We also define

$$(5.2) \quad n_c = \sup\{\delta : u_0 \in S_\delta \Rightarrow SC(u_0) \text{ holds}\}.$$

Hence, $0 < n_c \leq n_0$. Next we shall prove that $n_c < n_0$ is impossible, which implies that $n_c = n_0$. Thus, we assume now

$$n_c < n_0.$$

By the definition of n_c , we can find a sequence of solutions u_n of (NLS_k) with initial data $\phi_n \in \mathcal{N}^+$ such that $S_k(\phi_n) \rightarrow n_c$ and

$$(5.3) \quad \|u_n\|_{L_{\mathbf{R}^+,x}^5} = +\infty \text{ and } \|u_n\|_{L_{\mathbf{R}^-,x}^5} = +\infty.$$

In the subsequent subsection, our goal is to prove the existence of critical element $u_c \in H^1(\mathbf{R}^3)$, which is a global solution of (NLS_k) with initial data $u_{c,0}$ such that $S_k(u_{c,0}) = n_c$, $u_{c,0} \in \mathcal{N}^+$ and

$SC(u_{c,0})$ does not hold. Moreover, we prove that if $\|u_c\|_{L_{t,x}^5} = +\infty$, then $K := \{u_c(t) : t \in \mathbf{R}\}$ is precompact in $H^1(\mathbf{R}^3)$.

Before showing the existence and compactness of critical element u_c , we need a lemma related with the linear profile decomposition Lemma 2.12.

Lemma 5.2. *Let $M \in \mathbf{N}$ and $\psi^j \in H^1(\mathbf{R}^3)$ for any $0 \leq j \leq M$. Suppose that there exist some $\delta > 0$ and $\epsilon > 0$ with $2\epsilon < \delta$ such that*

$$\sum_{j=0}^M S_k(\psi^j) - \epsilon \leq S_k\left(\sum_{j=0}^M \psi^j\right) \leq n_0 - \delta, \quad -\epsilon \leq I_k\left(\sum_{j=0}^M \psi^j\right) \leq \sum_{j=0}^M I_k(\psi^j) + \epsilon.$$

Then $\psi^j \in \mathcal{N}^+$ for any $0 \leq j \leq M$.

Proof. Suppose that for some $0 \leq l \leq M$, $I_k(\psi^l) < 0$. By (3.2) with $(a, b) = (3, 0)$, we have $I_k((\psi^l)_\lambda^{3,0}) > 0$ for sufficiently small $\lambda < 0$. Thus, by continuity of $I_k((\psi^l)_\lambda^{3,0})$ in λ , there exists $\lambda_2 < 0$ such that $I_k((\psi^l)_{\lambda_2}^{3,0}) = 0$. As $n_k^{3,0} = n_0$, by the increasing property of $J_k^{3,0}((\psi^l)_\lambda^{3,0})$ in λ , we have

$$J_k^{3,0}(\psi^l) \geq J_k^{3,0}((\psi^l)_{\lambda_2}^{3,0}) = S_k((\psi^l)_{\lambda_2}^{3,0}) \geq n_0.$$

By the nonnegativity of $J_k^{3,0}(\psi^j)$ for any $0 \leq j \leq M$ and $2\epsilon < \delta$, we have

$$\begin{aligned} n_0 &\leq J_k^{3,0}(\psi^l) \leq \sum_{j=0}^M J_k^{3,0}(\psi^j) = \sum_{j=0}^M \left(S_k(\psi^j) - \frac{1}{2}I_k(\psi^j)\right) \\ &\leq S_k\left(\sum_{j=0}^M \psi^j\right) + \epsilon - \frac{1}{2}I_k\left(\sum_{j=0}^M \psi^j\right) + \frac{1}{2}\epsilon \\ &\leq n_0 - \delta + 2\epsilon < n_0, \end{aligned}$$

which is impossible. Hence, for each $0 \leq j \leq M$, we obtain

$$I_k(\psi^j) \geq 0.$$

So

$$S_k(\psi^j) = J_k^{3,0}(\psi^j) + \frac{1}{2}I_k(\psi^j) \geq 0,$$

which together with

$$\sum_{j=0}^M S_k(\psi^j) \leq S_k\left(\sum_{j=0}^M \psi^j\right) + \epsilon \leq n_0 - \delta + \epsilon < n_0$$

yields that $S_k(\psi^j) < n_0$ for each $0 \leq j \leq M$. By Lemma 3.7, $\psi^j \in \mathcal{N}^+$ for any $0 \leq j \leq M$. □

5.1. Existence and compactness of critical element.

Proposition 5.3. *There exists a $u_{c,0}$ in $H^1(\mathbf{R}^3)$ with $S_k(u_{c,0}) = n_c$, $u_{c,0} \in \mathcal{N}^+$ such that if u_c is the corresponding global solution of (NLS_k) with the initial data $u_{c,0}$, then $\|u_c\|_{L_{t,x}^5(\mathbf{R} \times \mathbf{R}^3)} = +\infty$ and K is precompact in $H^1(\mathbf{R}^3)$.*

Proof. We first note that $\{\phi_n\}_{n=1}^{+\infty}$ be a uniformly bounded sequence in $H^1(\mathbf{R}^3)$. In fact, since $\phi_n \in \mathcal{N}^+$ for any $n \in \mathbf{N}$, by (3.19),

$$\|\phi_n\|_{H^1} \leq \|\phi_n\|_{\mathcal{H}_k^1}^2 \leq 4S_k(\phi_n) < 4n_0.$$

We apply Lemma 2.12 to ϕ_n to get that for each $M \leq M^*$,

$$(5.4) \quad \phi_n = \sum_{j=1}^M \psi_n^j + W_n^M,$$

$$S_k(\phi_n) = \sum_{j=1}^M S_k(\psi_n^j) + S_k(W_n^M) + o_n(1),$$

and

$$I_k(\phi_n) = \sum_{j=1}^M I_k(\psi_n^j) + I_k(W_n^M) + o_n(1),$$

which together with $\phi_n \in \mathcal{N}^+$ yield that there exist some $\delta > 0$ and $\epsilon > 0$ with $2\epsilon < \delta$ such that

$$\sum_{j=1}^M S_k(\psi_n^j) + S_k(W_n^M) - \epsilon \leq S_k(\phi_n) \leq n_0 - \delta, \quad -\epsilon \leq I_k(\phi_n) \leq \sum_{j=1}^M I_k(\psi_n^j) + I_k(W_n^M) + \epsilon.$$

According to Lemma 5.2, we have that for large n and each $1 \leq j \leq M$, $\psi_n^j, W_n^M \in \mathcal{N}^+$, and then

$$(5.5) \quad 0 \leq \overline{\lim}_{n \rightarrow +\infty} S_k(\psi_n^j) \leq \overline{\lim}_{n \rightarrow +\infty} S_k(\phi_n) = n_c,$$

where if equality holds in the last inequality for some j , we must have $M^* = 1$ and $W_n^1 \rightarrow 0$ in H^1 .

We claim that if equality holds in the last inequality of (5.5) for some j (w.l.g. let $j = 1$), $u_{c,0}$ is namely ψ^1 . Indeed, at this time we have

$$(5.6) \quad \phi_n = \psi_n^1 + W_n^1 = e^{it_n H_a} \tau_{x_n^1} \psi^1 + W_n^1,$$

$$\overline{\lim}_{n \rightarrow +\infty} S_k(\psi_n^1) = n_c$$

and

$$(5.7) \quad W_n^1 \rightarrow 0 \text{ in } H^1.$$

Our target is to prove that

$$(5.8) \quad x_n^1 \equiv 0 \quad \text{and} \quad t_n^1 \equiv 0.$$

If (5.8) is true, then we have

$$\phi_n = \psi^1 + W_n^1, \quad S_k(\psi^1) = n_c, \quad \psi^1 \in \mathcal{N}^+$$

and

$$\lim_{n \rightarrow +\infty} \|\phi_n - \psi^1\|_{H^1} = 0.$$

Here ψ^1 is namely our required $u_{c,0}$. Let u_c be the solution of (NLS_k) with the initial data $u_{c,0} = \psi^1$, then u_c is global and $S_k(u_c) = S_k(u_{c,0}) = n_c$. Using Lemma 2.3, it holds that $\|u_c\|_{L_{t,x}^5(\mathbf{R} \times \mathbf{R}^3)} = +\infty$. Otherwise, $\|u_n\|_{L_{t,x}^5} < +\infty$, which contradicts with (5.3).

If (5.8) is false, then either $|x_n^1| \rightarrow +\infty$ or $t_n^1 \rightarrow \pm\infty$. I will see that it leads to $\|u_n\|_{L_{t,x}^5} < +\infty$ or $\|u_n\|_{L_{\mathbb{R}^{\pm,x}}^5} < +\infty$ contradicting with (5.3).

For the case $|x_n^1| \rightarrow +\infty$, by (2.20), we have

$$(5.9) \quad \lim_{n \rightarrow +\infty} \|\psi_n^1\|_{\mathcal{H}_k^1} = \|\psi^1\|_{H^1} > 0,$$

which implies that when $t_n^1 \equiv 0$,

$$S_0(\psi^1) = \overline{\lim}_{n \rightarrow +\infty} S_k(\psi_n^1) = n_c < n_0 \quad \text{and} \quad I_0(\psi^1) = \overline{\lim}_{n \rightarrow +\infty} I_k(\psi_n^1) \geq 0.$$

By (3.6), $P_0(\psi^1) \geq 0$. Hence, when $t_n^1 \equiv 0$, ψ^1 satisfies the condition (2.48). When $t_n^1 \rightarrow \pm\infty$, apply (5.9) and (2.32) to get

$$\frac{1}{2} \|\psi^1\|_{H^1}^2 = \overline{\lim}_{n \rightarrow +\infty} S_k(\psi_n^1) = n_c < n_0,$$

that is, ψ^1 satisfies the condition (2.49). Using Theorem 2.13 yields that the solution $NLS_k(t)\psi_n^1$ of (NLS_k) with initial data ψ_n^1 is global and satisfies

$$\|NLS_k(t)\psi_n^1\|_{S_\alpha^1(t)} \lesssim_{\|\psi^1\|_{H^1}} 1.$$

We know that $W_n^1 \rightarrow 0$ in H^1 , which is

$$\lim_{n \rightarrow +\infty} \|\phi_n - \psi_n^1\|_{H^1} = 0.$$

Using Lemma 2.3 again, we obtain $\|u_n\|_{L_{t,x}^5} < +\infty$.

For the other case $t_n^1 \rightarrow \pm\infty$, we only cope with $t_n^1 \rightarrow -\infty$ since $t_n^1 \rightarrow +\infty$ can be dealt with similarly. Apply (5.6) with $x_n^1 \equiv 0$, (5.7), Strichartz estimates Lemma 2.2 and norm equivalence Lemma 2.1 to get that

$$\begin{aligned} \lim_{n \rightarrow +\infty} \|e^{-itH_\alpha} \phi_n\|_{L_{\mathbb{R}^+,x}^5} &\leq \lim_{n \rightarrow +\infty} \|e^{-i(t-t_n^1)H_\alpha} \psi^1\|_{L_{\mathbb{R}^+,x}^5} + \lim_{n \rightarrow +\infty} \|e^{-itH_\alpha} W_n^1\|_{L_{\mathbb{R}^+,x}^5} \\ &\lesssim \lim_{n \rightarrow +\infty} \|W_n^1\|_{H^1} + \lim_{n \rightarrow +\infty} \|e^{-itH_\alpha} \psi^1\|_{L_{(-t_n^1, +\infty),x}^5} = 0, \end{aligned}$$

which immediately implies that $\lim_{n \rightarrow +\infty} \|u_n\|_{L_{\mathbb{R}^+,x}^5} = 0$ by (ii) of Theorem 1.1. Thus, we obtain that (5.8) holds true.

Next we turn to the other situation that equality doesn't hold in the last inequality of (5.5) for any $1 \leq j \leq M$. So for each $1 \leq j \leq M$ and $\psi_n^j \in \mathcal{N}^+$, there exists $\delta = \delta_j > 0$ such that

$$\overline{\lim}_{n \rightarrow +\infty} S_k(\psi_n^j) \leq n_c - 2\delta, \quad P_k(\psi_n^j) \geq 0 \quad \text{and} \quad I_k(\psi_n^j) \geq 0.$$

We shall use ψ_n^j to constitute approximate solutions of u_n under three cases: $|x_n^j| \rightarrow +\infty$; $x_n^j \equiv 0$ and $t_n^j \equiv 0$; $x_n^j \equiv 0$ and $t_n^j \rightarrow \pm\infty$ and then apply Lemma 2.3 to get a contradiction.

For some j such that $|x_n^j| \rightarrow +\infty$, (5.9) still holds for ψ_n^j . Using the same argument after (5.9), we obtain that ψ^j satisfies (2.48) or (2.49). Therefore, using Theorem 2.13, we can constitute a global solution $v_n^j(t) := NLS_k(t)\psi_n^j$ of (NLS_k) with initial data ψ_n^j such that

$$\|v_n^j\|_{L_{t,x}^5} \leq \|NLS_k(t)\psi_n^j\|_{S_\alpha^1(t)} \lesssim_{\|\psi^j\|_{H^1}} 1.$$

For some j such that $x_n^j \equiv 0$ and $t_n^j \equiv 0$, we apply $\psi^j \in \mathcal{N}^+$ to constitute a global solution $v_n^j(t) := NLS_k(t)\psi^j$ of (NLS_k) with initial data ψ^j .

For some j such that $x_n^j \equiv 0$ and $t_n^j \rightarrow \pm\infty$, by (iii) of Theorem 1.1, there exists $\tilde{\psi}^j \in H^1$ such that

$$(5.10) \quad \|NLS_k(t_n^j)\tilde{\psi}^j - e^{it_n^j H_\alpha} \psi^j\|_{\mathcal{H}_k^1} \sim \|NLS_k(t_n^j)\tilde{\psi}^j - e^{it_n^j H_\alpha} \psi^j\|_{H^1} \rightarrow 0 \text{ as } n \rightarrow +\infty,$$

which implies that for each $1 \leq j \leq M$ and n large enough,

$$S_k(NLS_k(t_n^j)\tilde{\psi}^j) \leq n_c - \delta, \quad P_k(NLS_k(t_n^j)\tilde{\psi}^j) \geq 0 \text{ and } I_k(NLS_k(t_n^j)\tilde{\psi}^j) \geq 0.$$

We set $v_n^j(0) = NLS_k(t_n^j)\tilde{\psi}^j$. Then according to the definition of n_c and $v_n^j(0) \in \mathcal{N}^+$, we obtain that the solution $v_n^j(t) := NLS_k(t + t_n^j)\tilde{\psi}^j$ of (NLS_k) with initial data $v_n^j(0)$ is global and satisfies uniform space-time bounds: $\|v_n^j\|_{L_{t,x}^5} < +\infty$.

As a result, we can construct approximate solutions of (NLS_k):

$$\tilde{u}_n(t) := \sum_{j=1}^M v_n^j + e^{-itH_\alpha} W_n^M$$

and set

$$e := (i\partial_t - H_\alpha)\tilde{u}_n + |\tilde{u}_n|^2\tilde{u}_n.$$

By (5.4) and (5.10), we have

$$(5.11) \quad \|\phi_n - \tilde{u}_n(0)\|_{H^1} = \|u_n(0) - \tilde{u}_n(0)\|_{H^1} \rightarrow 0 \text{ as } n \rightarrow +\infty,$$

which implies that

$$(5.12) \quad \overline{\lim}_{n \rightarrow +\infty} \|\tilde{u}_n(0)\|_{H^1} \text{ has a uniform bound independent of } M.$$

Using the same argument of Lemma 7.3 in [14] and replacing H_α and homogeneous fractional operator (e.g., $|\nabla|^{\frac{1}{2}}$) with $1 + H_\alpha$ and inhomogeneous fractional operator (e.g., $(1 + \Delta)^{\frac{1}{4}}$), respectively, we also obtain the same results as there for $\tilde{u}_n(t)$, that is,

$$(5.13) \quad \overline{\lim}_{n \rightarrow +\infty} \|\tilde{u}_n\|_{L_{t,x}^5} \text{ has a uniform bound independent of } M$$

and

$$(5.14) \quad \lim_{M \rightarrow M^*} \overline{\lim}_{n \rightarrow +\infty} \|(1 + \Delta)^{\frac{1}{4}} e\|_{L_{t,x}^{\frac{10}{7}}} = 0$$

Applying (5.11)-(5.14) to Theorem 2.3 gives $\|u_n\|_{L_{t,x}^5} < +\infty$, which is a contradiction with (5.3). Thus, we have completed the proof of existence of critical element u_c .

Finally, we consider precompactness of K in H^1 . We recall that u_c satisfies the following properties:

$$S_k(u_c(t)) = n_c, \quad u_c(t) \in \mathcal{N}^+ \text{ for } \forall t \in \mathbf{R} \text{ and } \|u_c\|_{L_{t,x}^5} = +\infty.$$

In particular, for any time sequence $\{t_n\}_{n=1}^{+\infty}$, the sequence $\{u_c(t_n)\}_{n=1}^{+\infty}$ also satisfies

$$S_k(u_c(t_n)) = n_c, \quad u_c(t_n) \in \mathcal{N}^+ \text{ for } \forall t \in \mathbf{R} \text{ and } \|u_c\|_{L_{(-\infty, t_n], x}^5} = \|u_c\|_{L_{(t_n, +\infty), x}^5} = +\infty.$$

Hence, regarding $u_c(t_n)$ as the foregoing ϕ_n and noting that the fact ϕ_n converges ψ^1 in H^1 yields that K is precompact in H^1 . Thus, we complete the whole proof. \square

5.2. Precluding the critical element. In this subsection, we shall apply the localized Virial identities (4.2) and (4.5) to preclude the critical element u_c . First by the precompactness of K , we have uniform localization of u_c : For each $\epsilon > 0$, there exists $R = R(\epsilon) > 0$ independent of t such that

$$(5.15) \quad \int_{|x|>R} (|\nabla u_c(t, x)|^2 + |u_c(t, x)|^2 + |u_c(t, x)|^4) dx \leq \epsilon.$$

We next claim that there exists a constant c such that for any $t \in \mathbf{R}$,

$$(5.16) \quad \|\nabla u_c(t)\|_{L^2} \geq c \|u_c(t)\|_{L^2}.$$

Indeed, if it's false, then there exists a time sequence $\{t_n\}_{n=1}^{+\infty}$ such that

$$\|\nabla u_c(t_n)\|_{L^2} \leq \frac{1}{n} \|u_c(t_n)\|_{L^2} = \frac{1}{n} \|u_c(0)\|_{L^2},$$

which means that $u_c(t_n) \rightarrow 0$ in H^1 . However, $\{u_c(t_n)\}_{n=1}^{+\infty}$ is precompact in H^1 . Hence, there exists a subsequence (still denoted by itself) $u_c(t_n) \rightarrow 0$ in H^1 . As $u_c(t_n) \in \mathcal{N}^+$, by (3.19), $n_c = \lim_{n \rightarrow +\infty} S_k(u_c(t_n)) = 0$, which is impossible.

Now we use localized Virial identities (4.2) and (4.5) only with u_c in place of u again, where we still choose the radial function ϕ satisfying (4.6). For R_1, R_2, R_3 and R_4 in (4.5), by (5.15), we have

$$(5.17) \quad \begin{aligned} |R_1 + R_2 + R_3 + R_4| &\lesssim \int_{|x| \geq R} (|\nabla u_c(t, x)|^2 + |u_c(t, x)|^4 + \frac{1}{R^2} |u_c(t, x)|^2 + \frac{1}{R^a} |u_c(t, x)|^2) dx \\ &\rightarrow 0 \text{ as } R \rightarrow +\infty. \end{aligned}$$

And for $P_k(u_c)$ in (4.5), by (3.21), $x \cdot \nabla V \leq 0$, (5.16) and (3.19), we have

$$(5.18) \quad \begin{aligned} P_k(u_c(t)) &\geq \min \left\{ 4(n_0 - S_k(u_c(t))), \frac{2}{5} \left(\|\nabla u_c(t)\|_{L^2}^2 - \frac{1}{2} \int_{\mathbf{R}^3} (x \cdot \nabla V) |u_c(t)|^2 dx \right) \right\} \\ &\gtrsim \|\nabla u_c(t)\|_{L^2}^2 \gtrsim \|u_c(t)\|_{H^1}^2 \gtrsim S_k(u_c(t)) = n_c. \end{aligned}$$

It follows from (5.17) and (5.18) that there exists $\delta_0 > 0$ such that for R large enough,

$$I''(t) \geq \delta_0,$$

which means that $\lim_{t \rightarrow +\infty} I'(t) = +\infty$. However, it is impossible since $I'(t)$ is bounded. In fact, by (4.2),

$$|I'(t)| \lesssim R.$$

Thus, we complete the proof of scattering part of Theorem 1.3.

REFERENCES

- [1] T. Akahori and H. Nawa, Blowup and scattering problems for the nonlinear Schrödinger equations, *Kyoto J. Math.*, 53(2013), pp. 629-672. 3, 4
- [2] N. Burq, F. Planchon, J. Stalker and A. S. Tahvildar-Zadeh, Strichartz estimates for the wave and Schrödinger equations with the inverse-square potential, *J. Funct. Anal.*, 203(2003), pp. 519-549. 7
- [3] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics, 10. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003. 15
- [4] D. Du, Y. Wu and K. Zhang, On blow-up criterion for the nonlinear Schrödinger equation, *Discrete Contin. Dyn. Syst.*, 36(2016), pp. 3639-3650. 1, 4, 5, 20
- [5] T. Duyckaerts, J. Holmer and S. Roudenko, Scattering for the non-radial 3D cubic nonlinear Schrödinger equation, *Math. Res. Lett.*, 15(2008), pp. 1233-1250. 2, 3, 4
- [6] D. Fang, J. Xie and T. Cazenave, Scattering for the focusing energy-subcritical nonlinear Schrödinger equation, *Sci. China Math.*, 54(2011), pp. 2037-2062. 3, 4

- [7] J. Holmer and S. Roudenko, A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation, *Comm. Math. Phys.*, 282(2008), pp. 435-467. 2, 3, 4
- [8] Y. Hong, Scattering for a nonlinear Schrödinger equation with a potential, *Commun. Pure Appl. Anal.*, 15(2016), pp. 1571-1601. 6, 20
- [9] S. Ibrahim, N. Masmoudi and K. Nakanishi, Scattering threshold for the focusing nonlinear Klein-Gordon equation, *Anal. PDE*, 4(2011), pp. 405-460. 1, 3, 4, 18
- [10] M. Ikeda and T. Inui, Global dynamics below the standing waves for the focusing semilinear Schrödinger equation with a repulsive Dirac delta potential. 3, 18
- [11] C. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case, *Invent. Math.*, 166(2006), pp. 645-675. 2
- [12] R. Killip, C. Miao, M. Visan, J. Zhang and J. Zheng, Multipliers and Riesz transforms for the Schrödinger operator with inverse-square potential, arXiv:1503.02716. 6, 7
- [13] R. Killip, C. Miao, M. Visan, J. Zhang and J. Zheng, The energy-critical NLS with inverse-square potential, arXiv: 1509.05822. 8, 9, 11
- [14] R. Killip, J. Murphy, M. Visan and J. Zheng, The focusing cubic NLS with inverse-square potential in three space dimensions, arXiv: 1603.08912. 2, 3, 7, 11, 12, 27
- [15] C. Miao, J. Zhang and J. Zheng, Nonlinear Schrödinger equation with Coulomb potential, arXiv:1809.06685. 3, 6
- [16] H. Mizutani, Strichartz estimates for Schrödinger equations with slowly decaying potential, arXiv:1808.06987. 2, 7
- [17] A. Sikora and J. Wright, Imaginary powers of Laplace operators, *Proc. Amer. Math. Soc.*, 129(2001), pp. 1745-1754. 6
- [18] B. Simon, Schrödinger semigroups, *Bull. Amer. Math. Soc.*, 7(1982), pp.447-526. 2
- [19] J. Zhang and J. Zheng, Scattering theory for nonlinear Schrödinger with inverse-square potential, *J. Funct. Anal.*, 267(2014), pp.2907-2932. 6

QING GUO, COLLEGE OF SCIENCE, MINZU UNIVERSITY OF CHINA, BEIJING, 100081, P.R. CHINA

E-mail address: guoqing0117@163.com

HUA WANG, SCHOOL OF MATHEMATICS AND STATISTICS AND HUBEI PROVINCE KEY LABORATORY OF MATHEMATICAL PHYSICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN, 430079, P.R. CHINA

E-mail address: wanghua_math@126.com

XIAOHUA YAO, SCHOOL OF MATHEMATICS AND STATISTICS AND HUBEI PROVINCE KEY LABORATORY OF MATHEMATICAL PHYSICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN, 430079, P.R. CHINA

E-mail address: yaoxiaohua@mail.ccnu.edu.cn