

ON THE APPROXIMATION PROPERTIES OF CESÀRO MEANS OF NEGATIVE ORDER OF VILENKIN-FOURIER SERIES

TSITSINO TEPNADZE

ABSTRACT. In this paper we establish approximation properties of Cesàro $(C, -\alpha)$ means with $\alpha \in (0, 1)$ of Vilenkin-Fourier series. This result allows one to obtain a condition which is sufficient for the convergence of the means $\sigma_n^{-\alpha}(f, x)$ to $f(x)$ in the L^p -metric.

Let N_+ denote the set of positive integers, $N := N_+ \cup \{0\}$. Let $m := (m_0, m_1, \dots)$ denote a sequence of positive integers not less than 2. Denote by $Z_{m_k} := \{0, 1, \dots, m_k - 1\}$ the additive group of integers modulo m_k . Define the group G_m as the complete direct product of the groups Z_{m_j} with the product of the discrete topologies of Z_{m_j} 's.

The direct product of the measures

$$\mu_k(\{j\}) := \frac{1}{m_k} \quad (j \in Z_{m_k}).$$

Is the Haar measure on G_m with $\mu(G_m) = 1$. If the sequence m is bounded, then G_m is called a bounded Vilenkin group. In this paper we will consider only bounded Vilenkin group. The elements of G_m can be represented by sequences $x := (x_0, x_1, \dots, x_j, \dots)$, $(x_j \in Z_{m_j})$. The group operation $+$ in G_m is given by

$$x + y = ((x_0 + y_0) \bmod m_0, \dots, (x_k + y_k) \bmod m_k, \dots),$$

where $x := (x_0, \dots, x_k, \dots)$ and $y := (y_0, \dots, y_k, \dots) \in G_m$.

The inverse of $+$ will be denoted by $-$. It is easy to give a base for the neighborhoods of G_m :

$$I_0(x) := G_m$$

$$I_n(x) := \{y \in G_m | y_0 = x_0, \dots, y_{n-1} = x_{n-1}\}.$$

for $x \in G_m$, $n \in N$ define $I_n = I_n(0)$. Set $e_n := (0, \dots, 0, 1, 0, \dots) \in G_m$ the n th coordinate of which is 1 and the rest are zeros ($n \in N$).

If we define the so-called generalized number system based on m in the following way: $M_0 := 1$, $M_{k+1} := m_k M_k$ ($k \in N$), then every $n \in N$ can

⁰2010 Mathematics Subject Classification 42C10 .

Key words and phrases: Vilenkin system, Cesàro means, Convergence in norm

be uniquely expressed as $n = \sum_{j=0}^{\infty} n_j M_j$, where $n_j \in Z_{mj}$ ($j \in N_+$) and only a finite number of n_j 's differ from zero. We use the following notation. Let $|n| := \max\{k \in N : n_k \neq 0\}$ (that is, $M_{|n|} \leq n < M_{|n|+1}$). Next, we introduce of G_m an orthonormal system which is called Vilenkin system. At first define the complex valued functions $r_k(x) : G_m \rightarrow C$. the generalized Rademacher functions in this way

$$r_k(x) := \exp\left(\frac{2\pi i x_k}{m_k}\right), \quad (i^2 = -1, x \in G_m, k \in N).$$

Now define the Vilenkin system $\psi := (\psi_n : n \in N)$ on G_m as follows.

$$\psi_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}(x), \quad (n \in N).$$

In particular, we call the system the Walsh-Paley if $m = 2$. The Vilenkin system is orthonormal and complete in $L^1(G_m)$. Now, introduce analogues of the usual definitions of the Fourier analysis. If $f \in L^1(G_m)$ we can establish the following definitions in the usual way:

Fourier coefficients:

$$\hat{f}(k) := \int_{G_m} f \overline{\psi_k} d\mu, \quad (k \in N),$$

partial sums:

$$S_n f := \sum_{k=0}^{n-1} \hat{f}(k) \psi_k, \quad (n \in N_+, S_0 f := 0),$$

Fejér means:

$$\sigma_n f := \sum_{k=0}^n \left(1 - \frac{k}{n+1}\right) \hat{f}(k) \psi_k, \quad (n \in N_+).$$

Dirichlet kernels:

$$D_n := \sum_{k=0}^{n-1} \psi_k, \quad (n \in N_+).$$

Fejér kernels:

$$K_n(x) := \frac{1}{n} \sum_{k=1}^n D_k(x).$$

Recall that (see [3] or [13])

$$(1) \quad D_{M_n}(x) = \begin{cases} M_n, & \text{if } x \in I_n, \\ 0, & \text{if } x \notin I_n. \end{cases}$$

It is well Known that

$$\sigma_n f(x) = \int_{G_m} f(t) K_n(x-t) d\mu(t).$$

The $(C, -\alpha)$ means of the Vilenkin-Fourier series are defined as

$$\sigma_n^{-\alpha}(f, x) = \frac{1}{A_n^{-\alpha}} \sum_{k=0}^n A_{n-k}^{-\alpha} \widehat{f}(k) \psi_k(x),$$

where

$$A_0^\alpha = 1, \quad A_n^\alpha = \frac{(\alpha+1)\dots(\alpha+n)}{n!}.$$

It is well Known that [16]

$$(2) \quad A_n^\alpha = \sum_{k=0}^n A_k^{\alpha-1}.$$

$$(3) \quad A_n^\alpha - A_{n-1}^\alpha = A_n^{\alpha-1}.$$

$$(4) \quad A_n^\alpha \sim n^\alpha.$$

The norm of the space $L^p(G_m)$ is defined by

$$\|f\|_p := \left(\int_{G_m} |f(x)|^p d\mu(x) \right)^{1/p}, \quad (1 \leq p < \infty).$$

Denote by $C(G_m)$ the class of continuous functions on the group G_m , endowed with the supremum norm.

For the sake of brevity in notation, we agree to write $L^\infty(G_m)$ instead of $C(G_m)$.

Let $f \in L^p(G_m)$, $1 \leq p \leq \infty$. The expression

$$\omega\left(\frac{1}{M_n}, f\right)_p = \sup_{h \in I_n} \|f(\cdot + h) - f(\cdot)\|_p$$

is called the modulus of continuity.

The problems of summability of partial sums and Cesàro means for Walsh-Fourier series were studied in [2], [5]-[12],[14]. In his monography [15] Zhizhinashvili investigated the behavior of Cesàro method of negative order for trigonometric Fourier series in detail. Goginava [5] studied analogaical question in case fo the Walsh system. In particular, the following theorem is proved.

Theorem G. [5] Let f belong to $L^p(G_2)$ for some $p \in [1, \infty]$ and $\alpha \in (0, 1)$. Then for any $2^k \leq n < 2^{k+1}$ ($k, n \in N$) the inequality

$$\|\sigma_n^{-\alpha}(f) - f\|_p \leq c(p, \alpha) \left\{ 2^{k\alpha} \omega\left(1/2^{k-1}, f\right)_p + \sum_{r=0}^{k-2} 2^{r-k} \omega\left(1/2^r, f\right)_p \right\}$$

holds true.

In this paper we establish approximation properties of Cesàro $(C, -\alpha)$ means with $\alpha \in (0, 1)$ of Vilenkin-Fourier series.

Theorem 1. Let f belong to $L^p(G_m)$ for some $p \in [1, \infty]$ and $\alpha \in (0, 1)$. Then for any $M_k \leq n < M_{k+1}$ ($k, n \in N$) the inequality

$$\|\sigma_n^{-\alpha}(f) - f\|_p \leq c(p, \alpha) \left\{ M_k^\alpha \omega\left(1/M_{k-1}, f\right)_p + \sum_{r=0}^{k-2} \frac{M_r}{M_k} \omega\left(1/M_r, f\right)_p \right\}$$

holds true.

This result allows one to obtain the condition which is sufficient for the convergence of the means $\sigma_n^{-\alpha}(f, x)$ to $f(x)$ in the L^p -metric.

Corollary 1. Let f belong to $L^p(G_m)$ for some $p \in [1, +\infty]$ and let

$$M_k^\alpha \omega\left(1/M_{k-1}, f\right)_p \rightarrow 0, \quad \text{as } k \rightarrow \infty, \quad (0 < \alpha < 1).$$

Then

$$\|\sigma_n^{-\alpha}(f) - f\|_p \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

In order to prove Theorem 1 we need the following lemmas.

Lemma 1. [1] Let $\alpha_1, \dots, \alpha_n$ be real numbers. Then

$$\frac{1}{n} \int_G \left| \sum_{k=1}^n \alpha_k D_k(x) \right| d\mu(x) \leq \frac{c}{\sqrt{n}} \left(\sum_{k=1}^n \alpha_k^2 \right)^{1/2}.$$

where c is an absolute constant.

Lemma 2. Let $f \in L^p(G_m)$ for some $p \in [1, \infty]$. Then for every $\alpha \in (0, 1)$ the following estimations holds

$$\begin{aligned} \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=0}^{M_{k-1}-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\ \leq c(p, \alpha) \sum_{r=0}^{k-1} \frac{M_r}{M_k} \omega\left(1/M_k, f\right)_p, \end{aligned}$$

where $M_k \leq n \leq M_{k+1}$.

Proof of Lemma 2. Applying Abel's transformation, from (3) we get

$$\begin{aligned}
(5) \quad & \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=0}^{M_{k-1}-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\
& = \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=1}^{M_{k-1}} A_{n-v-1}^{-\alpha} \psi_{v-1}(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\
& \leq \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=1}^{M_{k-1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\
& \quad + \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} A_{n-M_{k-1}-1}^{-\alpha} D_{M_{k-1}}(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\
& \quad = I_1 + I_2.
\end{aligned}$$

From the generalized Minkowski's inequality, and by (1) and (4) we obtain

$$\begin{aligned}
(6) \quad I_2 & \leq c(\alpha) M_{k-1} \int_{I_{k-1}} \| [f(\cdot + u) - f(\cdot)] \|_p d\mu(u) \\
& = O(\omega(1/M_{k-1}, f))_p,
\end{aligned}$$

$$\begin{aligned}
(7) \quad I_1 & \leq \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \left\| \int_{G_m} \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\
& \leq \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \left\| \int_{G_m} \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [f(\cdot + u) - S_{M_r}(\cdot + u, f)] d\mu(u) \right\|_p \\
& \quad + \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \left\| \int_{G_m} \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [S_{M_r}(\cdot + u, f) - S_{M_r}(\cdot, f)] d\mu(u) \right\|_p \\
& \quad + \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \left\| \int_{G_m} \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [S_{M_r}(\cdot, f) - f(\cdot)] d\mu(u) \right\|_p
\end{aligned}$$

$$= I_{11} + I_{12} + I_{13}.$$

Since

$$\|f - S_{M_r}(f)\|_p \leq \omega(1/M_r, f)_p,$$

using Lemma 1 for I_{11} we can write

$$(8) \quad I_{11}$$

$$\begin{aligned} &\leq \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \int_{G_m} \left| \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) \right| \|f(\cdot + u) - S_{M_r}(\cdot + u, f)\|_p d\mu(u) \\ &\leq \frac{1}{A_n^{-\alpha}} \sum_{r=0}^{k-2} \omega(1/M_r, f)_p \int_{G_m} \left| \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) \right| d\mu(u) \\ &\leq c(\alpha) n^\alpha \sum_{r=0}^{k-2} \omega(1/M_r, f)_p \sqrt{M_{r+1}} \left(\sum_{v=M_r}^{M_{r+1}-1} (n-v-1)^{-2\alpha-2} \right)^{1/2} \\ &\leq c(\alpha) n^\alpha \sum_{r=0}^{k-2} \omega(1/M_r, f)_p \sqrt{M_{r+1}} (n-M_{r+1})^{-\alpha-1} \sqrt{M_{r+1}} \\ &\leq c(\alpha) \sum_{r=0}^{k-2} \frac{M_r}{M_k} \omega(1/M_r, f)_p. \end{aligned}$$

Analogously, we can prove that

$$(9) \quad I_{13} \leq c(\alpha) \sum_{r=0}^{k-2} \frac{M_r}{M_k} \omega(1/M_r, f)_p.$$

It is evident that

$$\begin{aligned} (10) \quad & \int_{G_m} \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} D_v(u) [S_{M_r}(x+u, f) - S_{M_r}(x, f)] d\mu(u) \\ &= \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} \int_{G_m} S_{M_r}(x+u, f) D_v(u) d\mu(u) - \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} S_{M_r}(x, f) \\ &= \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} S_v(x, S_{M_r}(f)) - \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} S_{M_r}(x, f) \end{aligned}$$

$$= \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} S_{M_r}(x, f) - \sum_{v=M_r}^{M_{r+1}-1} A_{n-v-1}^{-\alpha-1} S_{M_r}(x, f) = 0.$$

Hence,

$$(11) \quad I_{12} = 0.$$

Combining (5)-(11) we receive the proof of Lemma 2. \square

Lemma 3. *Let $f \in L^p(G_m)$ for some $p \in [1, \infty]$. Then for every $\alpha \in (0, 1)$ the following estimations hold*

$$\begin{aligned} & \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\ & \leq c(p, \alpha) \omega(1/M_{k-1}, f)_p M_k^\alpha \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_k}^n A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\ & \leq c(p, \alpha) \omega(1/M_k, f)_p M_k^\alpha \end{aligned}$$

where $M_k \leq n < M_{k+1}$.

Proof of Lemma 3. We can write

$$\begin{aligned} (12) \quad II &= \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - f(\cdot)] d\mu(u) \right\|_p \\ &= \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) f(\cdot + u) d\mu(u) \right\|_p \\ &\leq \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot + u) - S_{M_{k-1}}(\cdot + u, f)] d\mu(u) \right\|_p \\ &+ \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) S_{M_{k-1}}(\cdot + u, f) d\mu(u) \right\|_p = II_1 + II_2. \end{aligned}$$

Since

$$(13) \quad \begin{aligned} & \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) S_{M_{k-1}}(x+u, f) d\mu(u) \\ &= \sum_{j=0}^{M_{k-1}-1} \widehat{f}(j) \psi_j(x) \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \int_{G_m} \psi_v(u) \psi_j(u) d\mu(u) = 0, \end{aligned}$$

for II_2 we obtain

$$(14) \quad II_2 = 0.$$

Using generalized Minkowski's inequality we have

$$(15) \quad \begin{aligned} II_1 &\leq \frac{1}{A_n^{-\alpha}} \int_{G_m} \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| \\ &\quad \times \| [f(\cdot \oplus u) - S_{M_{k-1}}(\cdot \oplus u, f)] \|_p d\mu(u) \\ &\leq c(\alpha) n^\alpha \omega(1/M_{k-1}, f)_p \int_{G_m} \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| d\mu(u). \end{aligned}$$

Let $t \in I_{A-1} \setminus I_A$, $A = 1, 2, \dots, k-1$ and $M_k = pM_A + q$, where $0 \leq q < M_A$. Then we have

$$(16) \quad \begin{aligned} & \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(t) \\ &= \sum_{v=M_{k-1}}^{pM_A-1} A_{n-v}^{-\alpha} \psi_v(t) + \sum_{v=pM_A}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(t) \\ &= \sum_{r=M_{k-1}/M_A}^{p-1} \sum_{v=rM_A}^{(r+1)M_A-1} A_{n-v}^{-\alpha} \psi_v(t) \\ &\quad + \sum_{v=0}^{q-1} A_{n-v-pM_A}^{-\alpha} \psi_{v+pM_A}(t) \\ &= \sum_{r=M_{k-1}/M_A}^{p-1} \sum_{v=0}^{M_A-1} A_{n-v-rM_A}^{-\alpha} \psi_{v+rM_A}(t) \\ &\quad + \sum_{v=0}^{q-1} A_{n-v-pM_A}^{-\alpha} \psi_{v+pM_A}(t) \end{aligned}$$

$$\begin{aligned}
&= \sum_{r=M_{k-1}/M_A}^{p-2} \psi_{rM_A}(t) \sum_{v=0}^{M_A-1} A_{n-v-rM_A}^{-\alpha} \psi_v(t) \\
&\quad + \psi_{(p-1)M_A}(t) \sum_{v=0}^{M_A-1} A_{n-v-(p-1)M_A}^{-\alpha} \psi_v(t) \\
&\quad + \psi_{rM_A}(t) \sum_{v=0}^{q-1} A_{q-v}^{-\alpha} \psi_v(t) \\
&= A_1 + A_2 + A_3.
\end{aligned}$$

Since $D_{M_A}(t) = 0, t \in I_{A-1} \setminus I_A$ and $|D_k(t)| \leq k$, from Abel's transformation, we have

$$\begin{aligned}
(17) \quad |A_1| &= \left| \sum_{r=M_{k-1}/M_A}^{p-2} \psi_{rM_A}(t) \sum_{v=0}^{M_A-2} A_{n-v-rM_A}^{-\alpha-1} D_v(t) \right| \\
&\leq c(\alpha) M_A \sum_{r=M_{k-1}/M_A}^{p-2} \sum_{v=0}^{M_A} (n - rM_A - v)^{-\alpha-1} \\
&\leq c(\alpha) M_A (n - (p-1)M_A)^{-\alpha} \\
&\leq c(\alpha) M_A^{1-\alpha}
\end{aligned}$$

For A_2 we have

$$\begin{aligned}
(18) \quad |A_2| &\leq c(\alpha) \sum_{v=0}^{M_A-1} (n - (p-1)M_A - v)^{-\alpha} \\
&\leq c(\alpha) \sum_{v=0}^{M_A-1} (M_A + q - v)^{-\alpha} \leq c(\alpha) M_A^{1-\alpha}.
\end{aligned}$$

We can write

$$\begin{aligned}
(19) \quad |A_3| &\leq c(\alpha) \sum_{v=0}^{q-1} (q - v)^{-\alpha} \\
&\leq c(\alpha) q^{1-\alpha} \leq c(\alpha) M_A^{1-\alpha}.
\end{aligned}$$

Combining (16)-(19) we obtain

$$(20) \quad \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| \leq c(\alpha) M_A^{1-\alpha}, \quad t \in I_{A-1} \setminus I_A, A = 1, \dots, k-1.$$

Hence, we can write

$$\begin{aligned}
(21) \quad & \int_{G_m} \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| d\mu(u) \\
&= \sum_{A=1}^{k-1} \int_{I_{A-1} \setminus I_A} \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| d\mu(u) \\
&\quad + \int_{I_{k-1}} \left| \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) \right| d\mu(u) \\
&\leq c(\alpha) \sum_{A=1}^k \frac{1}{M_A} M_A^{1-\alpha} + \frac{c(\alpha)}{M_k} M_k^{1-\alpha} \\
&\leq c(\alpha).
\end{aligned}$$

Combining (20)-(21) we have

$$(22) \quad II_1 \leq c(\alpha) \omega(1/M_{k-1}, f)_p M_k^\alpha.$$

Combining (15), (12) and (22) we conclude that

$$\begin{aligned}
& \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot+u) - f(\cdot)] d\mu(u) \right\|_p \\
&\leq c(\alpha) \omega(1/M_{k-1}, f)_p M_k^\alpha.
\end{aligned}$$

Analogously, we can prove that

$$\begin{aligned}
& \frac{1}{A_n^{-\alpha}} \left\| \int_{G_m} \sum_{v=M_k}^n A_{n-v}^{-\alpha} \psi_v(u) [f(\cdot+u) - f(\cdot)] d\mu(u) \right\|_p \\
&\leq c(\alpha) \omega(1/M_k, f)_p M_k^\alpha.
\end{aligned}$$

Lemma 3 proved. □

Proof of Theorem 1. We can write

$$\begin{aligned}
& \sigma_n^{-\alpha}(f, x) - f(x) \\
= & \frac{1}{A_n^{-\alpha}} \int_{G_m} \sum_{v=0}^{M_{k-1}-1} A_{n-v}^{-\alpha} \psi_v(x) [f(\cdot + u) - f(\cdot)] d\mu(u) \\
& + \frac{1}{A_n^{-\alpha}} \int_{G_m} \sum_{v=M_{k-1}}^{M_k-1} A_{n-v}^{-\alpha} \psi_v(x) [f(\cdot + u) - f(\cdot)] d\mu(u) \\
& + \frac{1}{A_n^{-\alpha}} \int_{G_m} \sum_{v=M_k}^n A_{n-v}^{-\alpha} \psi_v(x) [f(\cdot + u) - f(\cdot)] d\mu(u) \\
= & I + II + III
\end{aligned}$$

Since

$$\|\sigma_n^{-\alpha}(f, \cdot) - f(\cdot)\|_p \leq \|I\|_p + \|II\|_p + \|III\|_p$$

From Lemmas 2 and 3 the proof of theorem is complete. \square

REFERENCES

- [1] G. N. Agaev, N.Ya. Vilenkin, G.M. Dzhafarli, and A.I. Rubinshtejn, Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehlm, 1981 (in Russian).
- [2] N.J.fine,Cesàro summability of Walsh-Fourier series, *Proc. Nat.Acad. Sci. U.S.A* 41 (1995), 558-591.
- [3] B. I. Golubov, A.V.Efimov, and V.A.Skvortsov,"series and transformation of Walsh,"Nauka,Moscow,1987 [In Russian];English translation,Kluwer Academic,Dordrecht,1991.
- [4] U. Goginava, On the uniform convergence of Walsh-Fourier series. *Acta Math. Hungar.* 93 (2001), no. 1-2, 59–70.
- [5] U. Goginava, On the approximation properties of Cesàro means of negative order of Walsh-Fourier series. *J. Approx. Theory* 115 (2002), no. 1, 9–20.
- [6] U. Goginava, Uniform convergence of Cesàro means of negative order of double Walsh-Fourier series. *J. Approx. Theory* 124 (2003), no. 1, 96–108.
- [7] U. Goginava and K. Nagy, On the maximal operator of Walsh-Kaczmarz-Fejér means. *Czechoslovak Math. J.* 61(136) (2011), no. 3, 673–686.
- [8] G. Gát and U. Goginava, A weak type inequality for the maximal operator of (C, α) -means of Fourier series with respect to the Walsh-Kaczmarz system. *Acta Math. Hungar.* 125 (2009), no. 1-2, 65–83.
- [9] G. Gát and K. Nagy, Cesàro summability of the character system of the p-series field in the Kaczmarz rearrangement. *Anal. Math.* 28 (2002), no. 1, 1–23.
- [10] K. Nagy, Approximation by Cesàro means of negative order of Walsh-Kaczmarz-Fourier series. *East J. Approx.* 16 (2010), no. 3, 297–311.
- [11] P. Simon and F. Weisz, Weak inequalities for Cesàro and Riesz summability of Walsh-Fourier series. *J. Approx. Theory* 151 (2008), no. 1, 1–19.
- [12] F. Schipp, Über gewisse Maximaloperatoren, *Ann. Univ. Sci. Budapest. Sect. Math.* 18(1975), 189-195.
- [13] F. Schipp, W.R. Wade, P. Simon and J. Pál, "Walsh Series, Introduction to Dyadic Harmonic Analysis," Hilger, Bristol, 1990.

- [14] V. I. Tevzadze, Uniform $(C, -\alpha)$ summability of Fourier series with respect to the Walsh-Paley system. *Acta Math. Acad. Paedagog. Nyhazi. (N.S.)* 22 (2006), no. 1, 41–61 (electronic).
- [15] L. V. Zhizhiashvili, Trigonometric Fourier series and their conjugates, Tbilisi, 1993 (Russian); English transl.: Kluwer Acad. publ; 1996.
- [16] A.Zygmund , Trigonometric series.Vo1.1,Cambridge University Press, Cambridge,UK, 1959.

T.TEPNADZE, DEPARTMENT OF MATHEMATICS, FACULTY OF EXACT AND NATURAL SCIENCES, IVANE JAVAKHISHVILI TBILISI STATE UNIVERSITY, CHAVCHAVADZE STR. 1, TBILISI 0128, GEORGIA

E-mail address: tsitsinotefnadze@gmail.com