

A COMPREHENSIVE SUBCLASS OF ANALYTIC AND BI-UNIVALENT FUNCTIONS ASSOCIATED WITH SUBORDINATION

A. A. AMOURAH AND MOHAMED ILLAFE

ABSTRACT. In the present paper, we define a new general subclass of bi-univalent functions involving a differential operator in the open unit disk \mathbb{U} . For this purpose, we use the Faber polynomial expansions. Several connections to some of the earlier known results are also pointed out.

1. INTRODUCTION

Let \mathcal{A} denote the class of all analytic functions f defined in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ and normalized by the conditions $f(0) = 0$ and $f'(0) = 1$. Thus each $f \in \mathcal{A}$ has a Taylor-Maclaurin series expansion of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (z \in \mathbb{U}). \quad (1.1)$$

Further, let \mathcal{S} denote the class of all functions $f \in \mathcal{A}$ which are univalent in \mathbb{U} (for details, see [10]; see also some of the recent investigations [3, 6, 7, 8, 23]). And let \mathcal{C} be the class of functions $\Phi(z) = 1 + \sum_{n=1}^{\infty} \Phi_n z^n$ that are analytic in \mathbb{U} and satisfy the condition $\operatorname{Re}(\Phi(z)) > 0$ in \mathbb{U} . By the Caratheodory's lemma (see [10]) we have $|\Phi(z)| \leq 2$.

Let the functions f, g be analytic in \mathbb{U} . If there exists a Schwarz function ϖ , is analytic in \mathbb{U} under the conditions

$$\varpi(0) = 0, \quad |\varpi(z)| \leq 1,$$

such that

$$f(z) = g(\varpi(z)), \quad z \in \mathbb{U},$$

then, the function f is subordinate to g in \mathbb{U} , and we write $f(z) \prec g(z)$.

The Koebe one-quarter theorem (for details, (see [10]), we know that the image of \mathbb{U} under every function $f \in \mathcal{A}$ contains a disk of radius $\frac{1}{4}$. According to this, every function $f \in \mathcal{A}$ has an inverse map f^{-1} that satisfies the following conditions:

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U}),$$

and

2010 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Faber polynomials, bi-univalent functions, subordination, upper bound.

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4} \right).$$

In fact, the inverse function is given by

$$g(w) = f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2 a_3 + a_4)w^4 + \dots \quad (1.2)$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in \mathbb{U} if both $f(z)$ and $f^{-1}(z)$ are univalent in \mathbb{U} . Let Σ denote the class of bi-univalent functions in \mathbb{U} given by (1.1). Examples of functions in the class Σ are

$$\frac{z}{1-z}, \quad -\log(1-z), \quad \frac{1}{2} \log\left(\frac{1+z}{1-z}\right), \dots$$

It is worth noting that the familiar Koebe function is not a member of Σ , since it maps the unit disk \mathbb{U} univalently onto the entire complex plane except the part of the negative real axis from $-1/4$ to $-\infty$. Thus, clearly, the image of the domain does not contain the unit disk \mathbb{U} . For a brief history and some intriguing examples of functions and characterization of the class Σ , see Srivastava et al. [20], Yousef et al. [24, 25, 26], and Frasin and Aouf [12].

In 1967, Lewin [18] investigated the bi-univalent function class Σ and showed that $|a_2| < 1.51$. Subsequently, Brannan and Clunie [9] conjectured that $|a_2| \leq \sqrt{2}$. On the other hand, Netanyahu [19] showed that $\max_{f \in \Sigma} |a_2| = \frac{4}{3}$. The best known estimate for functions in Σ has been obtained in 1984 by Tan [21], that is, $|a_2| < 1.485$. The coefficient estimate problem for each of the following Taylor-Maclaurin coefficients $|a_n|$ ($n \in \mathbb{N} \setminus \{1, 2\}$) for each $f \in \Sigma$ given by (1.1) is presumably still an open problem.

The Faber polynomials introduced by Faber [11] play an important role in various areas of mathematical sciences, especially in geometric function theory. The recent publications [13] and [14] applying the Faber polynomial expansions to meromorphic bi-univalent functions motivated us to apply this technique to classes of analytic bi-univalent functions. In the literature, there are only a few works determining the general coefficient bounds $|a_n|$ for the analytic bi-univalent functions given by (1.1) using Faber polynomial expansions (see for example, [15, 16, 17]). Hamidi and Jahangiri [15] considered the class of analytic bi-close-to-convex functions. Jahangiri and Hamidi [17] considered the class defined by Frasin and Aouf [12], and Jahangiri et al. [16] considered the class of analytic bi-univalent functions with positive real-part derivatives.

2. THE CLASS $\mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$

Yousef et al. [25] have introduced and studied the following subclass of analytic bi-univalent functions:

Definition 2.1. For $\lambda \geq 1, \mu \geq 0, \delta \geq 0$ and $0 \leq \alpha < 1$, a function $f \in \Sigma$ given by (1.1) is said to be in the class $\mathfrak{B}_\Sigma^\mu(\alpha, \lambda, \delta)$ if the following conditions hold for all $z, w \in \mathbb{U}$:

$$\operatorname{Re} \left((1 - \lambda) \left(\frac{f(z)}{z} \right)^\mu + \lambda f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \xi \delta z f''(z) \right) > \alpha \tag{2.1}$$

and

$$\operatorname{Re} \left((1 - \lambda) \left(\frac{g(w)}{w} \right)^\mu + \lambda g'(w) \left(\frac{g(w)}{w} \right)^{\mu-1} + \xi \delta w g''(w) \right) > \alpha, \tag{2.2}$$

where the function $g(w) = f^{-1}(w)$ is defined by (1.2) and $\xi = \frac{2\lambda + \mu}{2\lambda + 1}$.

Using the Faber polynomial expansion of functions $f \in \mathcal{A}$ of the form (1.1), the coefficients of its inverse map $g = f^{-1}$ may be expressed as in [1]:

$$g(w) = f^{-1}(w) = w + \sum_{n=2}^{\infty} \frac{1}{n} K_{n-1}^{-n}(a_2, a_3, \dots) w^n, \tag{2.3}$$

where

$$\begin{aligned} K_{n-1}^{-n} = & \frac{(-n)!}{(-2n+1)!(n-1)!} a_2^{n-1} + \frac{(-n)!}{(2(-n+1))!(n-3)!} a_2^{n-3} a_3 + \frac{(-n)!}{(-2n+3)!(n-4)!} a_2^{n-4} a_4 \\ & + \frac{(-n)!}{(2(-n+2))!(n-5)!} a_2^{n-5} [a_5 + (-n+2)a_3^2] + \frac{(-n)!}{(-2n+5)!(n-6)!} a_2^{n-6} \\ & [a_6 + (-2n+5)a_3 a_4] + \sum_{j \geq 7} a_2^{n-j} V_j, \end{aligned} \tag{2.4}$$

such that V_j with $7 \leq j \leq n$ is a homogeneous polynomial in the variables a_2, a_3, \dots, a_n [2].

In particular, the first three terms of K_{n-1}^{-n} are

$$K_1^{-2} = -2a_2, \quad K_2^{-3} = 3(2a_2^2 - a_3), \quad K_3^{-4} = -4(5a_2^3 - 5a_2 a_3 + a_4). \tag{2.5}$$

In general, for any $p \in \mathbb{N} := \{1, 2, 3, \dots\}$, an expansion of K_n^p is as in [1],

$$K_n^p = p a_n + \frac{p(p-1)}{2} D_n^2 + \frac{p!}{(p-3)!3!} D_n^3 + \dots + \frac{p!}{(p-n)!n!} D_n^n, \tag{2.6}$$

where $D_n^p = D_n^p(a_2, a_3, \dots)$, and by [22], $D_n^m(a_1, a_2, \dots, a_n) = \sum_{i_1, \dots, i_n}^{\infty} \frac{m!}{i_1! \dots i_n!} a_1^{i_1} \dots a_n^{i_n}$ while $a_1 = 1$, and the sum is taken over all non-negative integers i_1, \dots, i_n satisfying $i_1 + i_2 + \dots + i_n = m$, $i_1 + 2i_2 + \dots + ni_n = n$, it is clear that $D_n^m(a_1, a_2, \dots, a_n) = a_1^n$.

Now, we are ready to establish a new subclass of analytic and bi-univalent functions based on subordination.

Definition 2.2. For $\lambda \geq 1$, $\mu \geq 0$, and $\delta \geq 0$, A function $f \in \Sigma$ is said to be in the class $\mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$, if the following subordinations are satisfied:

$$(1 - \lambda) \left(\frac{f(z)}{z} \right)^\mu + \lambda f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \xi \delta z f''(z) \prec \Phi(z) \quad (2.7)$$

and

$$(1 - \lambda) \left(\frac{g(w)}{w} \right)^\mu + \lambda g'(w) \left(\frac{g(w)}{w} \right)^{\mu-1} + \xi \delta w g''(w) \prec \Phi(w) \quad (2.8)$$

where the function $g(w) = f^{-1}(w)$ is defined by (1.2) and $\xi = \frac{2\lambda + \mu}{2\lambda + 1}$.

3. COEFFICIENT BOUNDS FOR THE FUNCTION CLASS $\mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$

Theorem 3.1. For $\lambda \geq 1$, $\mu \geq 0$, and $\delta \geq 0$, let the function $f \in \mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$ be given by (1.1). Then

$$|a_2| \leq \min \left\{ \frac{2}{\mu + \lambda + 2\xi\delta}, \sqrt{\frac{8}{(\mu + 2\lambda + 6\xi\delta)(\mu + 1)}} \right\}$$

and

$$|a_3| \leq \min \left\{ \frac{\frac{1}{(1 + \frac{6\delta}{2\lambda + 1})} \left[\frac{4}{(\mu + \lambda + 2\xi\delta)^2} + \frac{2}{(\mu + 2\lambda + 6\xi\delta)} \right]}{\frac{1}{(1 + \frac{6\delta}{2\lambda + 1})} \left[\frac{8}{(\mu + 2\lambda + 6\xi\delta)(\mu + 1)} + \frac{2}{(\mu + 2\lambda + 6\xi\delta)} \right]} \right\}.$$

Proof. Let $\mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$. The inequalities (2.7) and (2.8) imply the existence of two positive real part functions

$$\varpi(z) = 1 + \sum_{n=1}^{\infty} t_n z^n$$

and

$$\varphi(w) = 1 + \sum_{n=1}^{\infty} s_n z^n$$

where $\operatorname{Re}(\varpi(z)) > 0$ and $\operatorname{Re}(\varphi(w)) > 0$ in \mathcal{C} so that

$$(1 - \lambda) \left(\frac{f(z)}{z} \right)^\mu + \lambda f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \xi \delta z f''(z) = \Phi(\varpi(z)), \quad (3.1)$$

$$(1 - \lambda) \left(\frac{g(w)}{w} \right)^\mu + \lambda g'(w) \left(\frac{g(w)}{w} \right)^{\mu-1} + \xi \delta w g''(w) = \Phi(\varphi(w)). \quad (3.2)$$

It follows from (3.1) and (3.2) that

$$(\mu + \lambda + 2\xi\delta) a_2 = \Phi_1 t_1 \quad (3.3)$$

$$(\mu + 2\lambda + 6\xi\delta) \left[\frac{\mu - 1}{2} a_2^2 + \left(1 + \frac{6\delta}{2\lambda + 1} \right) a_3 \right] = \Phi_1 t_2 + \Phi_2 t_1^2, \quad (3.4)$$

and

$$-(\mu + \lambda) a_2 = \Phi_1 s_1, \quad (3.5)$$

$$(\mu + 2\lambda + 6\xi\delta) \left[\frac{\mu + 3}{2} a_2^2 - \left(1 + \frac{6\delta}{2\lambda + 1} \right) a_3 \right] = \Phi_1 s_2 + \Phi_2 s_1^2. \quad (3.6)$$

From (3.3) and (3.5), we find

$$|a_2| \leq \frac{|\Phi_1 t_1|}{\mu + \lambda + 2\xi\delta} = \frac{|\Phi_1 s_1|}{\mu + \lambda + 2\xi\delta} \leq \frac{2}{\mu + \lambda + 2\xi\delta}. \quad (3.7)$$

From (3.4) and (3.6), we get

$$(\mu + 2\lambda + 6\xi\delta) (\mu + 1) a_2^2 = \Phi_1 (t_2 + s_2) + \Phi_2 (t_1^2 + s_1^2)$$

or, equivalently

$$|a_2| \leq \sqrt{\frac{8}{(\mu + 2\lambda + 6\xi\delta) (\mu + 1)}}. \quad (3.8)$$

Next, in order to find the bound on the coefficient $|a_3|$, we subtract (3.6) from (3.4). We thus get

$$2(\mu + 2\lambda + 6\xi\delta) \left[\left(1 + \frac{6\delta}{2\lambda + 1} \right) a_3 - a_2^2 \right] = \Phi_1 (t_2 - s_2) + \Phi_2 (t_1^2 - s_1^2) \quad (3.9)$$

or

$$\begin{aligned} |a_3| &\leq \frac{1}{\left(1 + \frac{6\delta}{2\lambda + 1} \right)} \left[|a_2|^2 + \frac{|\Phi_1 (t_2 - s_2)|}{2(\mu + 2\lambda + 6\xi\delta)} \right] \\ &= \frac{1}{\left(1 + \frac{6\delta}{2\lambda + 1} \right)} \left[|a_2|^2 + \frac{2}{(\mu + 2\lambda + 6\xi\delta)} \right]. \end{aligned} \quad (3.10)$$

Upon substituting the value of a_2^2 from (3.7) and (3.8) into (3.10), it follows that

$$|a_3| \leq \frac{1}{\left(1 + \frac{6\delta}{2\lambda + 1} \right)} \left[\frac{4}{(\mu + \lambda + 2\xi\delta)^2} + \frac{2}{(\mu + 2\lambda + 6\xi\delta)} \right]$$

and

$$|a_3| \leq \frac{1}{\left(1 + \frac{6\delta}{2\lambda + 1} \right)} \left[\frac{8}{(\mu + 2\lambda + 6\xi\delta) (\mu + 1)} + \frac{2}{(\mu + 2\lambda + 6\xi\delta)} \right]$$

Which completes the proof of 3.1. □

Theorem 3.2. Let $\mathfrak{B}_\Sigma(\mu, \lambda, \Phi, \xi)$. If $a_m = 0$ with $2 \leq m \leq n - 1$, then

$$|a_n| \leq \frac{2}{\mu + (n - 1)\lambda + n(n - 1)\xi\delta} \quad (n \geq 4). \quad (3.11)$$

Proof. By using the Faber polynomial expansion of functions $f \in \mathcal{A}$ of the form (1.1) and its inverse map $g = f^{-1}$, we can write

$$(1 - \lambda) \left(\frac{f(z)}{z} \right)^\mu + \lambda f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \xi \delta z f''(z) = 1 + \sum_{n=2}^{\infty} F_{n-1}(a_2, a_3, \dots, a_n) z^{n-1} \quad (3.12)$$

and

$$(1 - \lambda) \left(\frac{g(w)}{w} \right)^\mu + \lambda g'(w) \left(\frac{g(w)}{w} \right)^{\mu-1} + \xi \delta w g''(w) = 1 + \sum_{n=2}^{\infty} F_{n-1}(A_2, A_3, \dots, A_n) w^{n-1} \quad (3.13)$$

where

$$F_1 = (\mu + \lambda + 2\xi\delta)a_2, \quad F_2 = (\mu + 2\lambda + 6\xi\delta) \left[\frac{\mu - 1}{2}a_2^2 + \left(1 + \frac{6\delta}{2\lambda + 1}\right)a_3 \right] \quad (3.14)$$

and, in general (see [5])

$$F_{n-1}(a_2, a_3, \dots, a_n) = [\mu + (n-1)\lambda + n(n-1)\xi\delta] \times [(\mu-1)!] \quad (3.15)$$

$$\times \left[\sum_{n=2}^{\infty} \frac{a_2^{i_1} a_3^{i_2} \dots a_n^{i_{n-1}}}{i_1! i_2! \dots i_n! [\mu - (i_1 + i_2 + \dots + i_{n-1})!]} \right] \quad (3.16)$$

Next, by using the Faber polynomial expansion of functions $\varpi, \varphi \in \mathcal{C}$, we also obtain

$$\Phi(\varpi(z)) = 1 + \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \Phi_n F_n^k(t_1, t_2, \dots, t_n) z^n, \quad (3.17)$$

and

$$\Phi(\varphi(z)) = 1 + \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \Phi_n F_n^k(s_1, s_2, \dots, s_n) w^n. \quad (3.18)$$

Comparing the corresponding coefficients yields

$$[\mu + (n-1)\lambda + n(n-1)\xi\delta] a_n = \sum_{k=1}^{n-1} \Phi_n F_{n-1}^k(t_1, t_2, \dots, t_{n-1}) \quad (n \geq 2)$$

and

$$[\mu + (n-1)\lambda + n(n-1)\xi\delta] A_n = \sum_{k=1}^{n-1} \Phi_n F_{n-1}^k(s_1, s_2, \dots, s_{n-1}) \quad (n \geq 2). \quad (3.19)$$

Note that for $a_m = 0$, $2 \leq m \leq n-1$, we have $A_n = -a_n$ and so

$$\begin{aligned} [\mu + (n-1)\lambda + n(n-1)\xi\delta] a_n &= \Phi_1 t_{n-1}, \\ -[\mu + (n-1)\lambda + n(n-1)\xi\delta] a_n &= \Phi_1 s_{n-1}, \end{aligned} \quad (3.20)$$

Now taking the absolute values of either of the above two equations and using the facts that $|\Phi_1| \leq 2$, $|t_{n-1}| \leq 1$, and $|s_{n-1}| \leq 1$, we obtain

$$|a_n| \leq \frac{|\Phi_1 t_{n-1}|}{\mu + (n-1)\lambda + n(n-1)\xi\delta} = \frac{|\Phi_1 s_{n-1}|}{\mu + (n-1)\lambda + n(n-1)\xi\delta} \quad (3.21)$$

$$\leq \frac{2}{\mu + (n-1)\lambda + n(n-1)\xi\delta} \quad (3.22)$$

This evidently completes the proof of 3.2. \square

Remark 3.3. As a final remark, for $\delta = 0$ in

(i) Theorem 3.1 we obtain Theorem 1 in [4].

(ii) Theorem 3.2 we obtain Theorem 2 in [4].

REFERENCES

- [1] H. Airault; A. Bouali, *Differential calculus on the Faber polynomials*, Bull. Sci. Math. 130 (3) (2006) 179–222.
- [2] H. Airault; J. Ren, *An algebra of differential operators and generating functions on the set of univalent functions*, Bull. Sci. Math. 126 (5) (2002) 343–367.
- [3] T. Al-Hawary; B.A. Frasin; F. Yousef, *Coefficients estimates for certain classes of analytic functions of complex order*, Afrika Matematika 29(7-8) (2018), 1265-1271.
- [4] S. Altinkaya; S. Y. Tokgöz, *On the bounds of general subclasses of analytic and bi-univalent functions associated with subordination*, In 4th International Conference on Analysis and its Applications, September 11-14, 2018, Kirsehir/Turkey, p. 135.
- [5] A. Amourah, *Faber polynomial coefficient estimates for a class of analytic bi-univalent functions*, arXiv preprint arXiv:1810.07018 (2018).
- [6] A.A. Amourah; F. Yousef, *Some properties of a class of analytic functions involving a new generalized differential operator*, Boletim da Sociedade Paranaense de Matemática, In press.
- [7] A.A. Amourah; F. Yousef; T. Al-Hawary, M. Darus, *A certain fractional derivative operator for p -valent functions and new class of analytic functions with negative coefficients*, Far East Journal of Mathematical Sciences 99(1) (2016) 75-87.
- [8] A.A. Amourah; F. Yousef; T. Al-Hawary; M. Darus, *On $H_3(p)$ Hankel determinant for certain subclass of p -valent functions*, Ital. J. Pure Appl. Math 37 (2017) 611-618.
- [9] D.A. Brannan; J.G. Clunie, *Aspects of contemporary complex analysis* (Proceedings of the NATO Advanced Study Institute held at the University of Durham, Durham; July 120, 1979), Academic Press, New York and London, 1980.
- [10] P.L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, Berlin, Heidelberg and Tokyo, 1983.
- [11] G. Faber, *Über polynomische entwickelungen*, Mathematische Annalen 57(3) (1903) 389-408.
- [12] B.A. Frasin; M.K. Aouf, *New subclasses of bi-univalent functions*, Appl. Math. Lett. 24(9) (2011) 1569–1573.
- [13] S.G. Hamidi; S.A. Halim; J.M. Jahangiri, *Coefficient estimates for a class of meromorphic bi-univalent functions*, C. R. Acad. Sci. Paris, Ser. I 351 (9–10) (2013) 349–352.
- [14] S.G. Hamidi; T. Janani; G. Murugusundaramoorthy, J.M. Jahangiri, *Coefficient estimates for certain classes of meromorphic bi-univalent functions*, C. R. Acad. Sci. Paris, Ser. I 352 (4) (2014) 277–282.
- [15] S.G. Hamidi; J.M. Jahangiri, *Faber polynomial coefficient estimates for analytic bi-close-to-convex functions*, C. R. Acad. Sci. Paris, Ser. I 352 (1) (2014) 17–20.
- [16] J.M. Jahangiri; S.G. Hamidi; S.A. Halim, *Coefficients of bi-univalent functions with positive real part derivatives*, Bull. Malays. Math. Soc., in press, <http://math.usm.my/bulletin/pdf/acceptedpapers/2013-04-050-R1.pdf>.
- [17] J.M. Jahangiri; S.G. Hamidi, *Coefficient estimates for certain classes of bi-univalent functions*, Int. J. Math. Math. Sci. (2013), Article ID 190560, 4 p.
- [18] M. Lewin, *On a coefficient problem for bi-univalent functions*, Proc. Amer. Math. Soc. 18 (1967) 63–68.
- [19] E. Netanyahu, *The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$* , Arch. Rational Mech. Anal. 32 (1969) 100–112.
- [20] H.M. Srivastava; A.K. Mishra; P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett. 23(10) (2010) 1188–1192.
- [21] D.L. Tan, *Coefficient estimates for bi-univalent functions*, Chin. Ann. Math. Ser. A 5 (1984) 559–568.
- [22] P.G. Todorov, *On the Faber polynomials of the univalent functions of class Σ* , J. Math. Anal. Appl. 162 (1) (1991) 268–276.
- [23] F. Yousef; A.A. Amourah; M. Darus, *Differential sandwich theorems for p -valent functions associated with a certain generalized differential operator and integral operator*, Italian Journal of Pure and Applied Mathematics 36 (2016) 543-556.
- [24] F. Yousef; B.A. Frasin; T. Al-Hawary, *Fekete-Szegő Inequality for Analytic and Bi-univalent Functions Subordinate to Chebyshev Polynomials*, arXiv preprint arXiv:1801.09531 (2018).
- [25] F. Yousef; S. Alroud; M. Illafe, *New subclasses of analytic and bi-univalent functions endowed with coefficient estimate problems*, arXiv preprint arXiv:1808.06514 (2018).

- [26] F. Yousef; S. Alroud; M. Illafe, *A Comprehensive Subclass of Bi-Univalent Functions Associated with Chebyshev Polynomials of the Second Kind*, arXiv preprint arXiv:1809.09365 (2018).

A. A. AMOURAH: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, IRBID NATIONAL UNIVERSITY, IRBID, JORDAN.

E-mail address: alaammour@yahoo.com

MOHAMED ILLAFE: SCHOOL OF ENGINEERING, MATH, & TECHNOLOGY, NAVAJO TECHNICAL UNIVERSITY, CROWNPOINT, NM 87313, USA.

E-mail address: millafe@navajotech.edu