PAINLEVÉ-II PROFILE OF THE SHADOW KINK IN THE THEORY OF LIGHT-MATTER INTERACTION IN NEMATIC LIQUID CRYSTALS

CHRISTOS SOURDIS

ABSTRACT. We confirm a prediction that the recently introduced shadow kink defect in the theory of light-matter interaction in nematic liquid crystals is described to main order by a solution of the Painlevé-II equation which changes sign once in the whole real line. Our result implies the existence of such a solution to the latter equation which is energy minimizing with respect to compactly supported perturbations.

1. INTRODUCTION

Motivated by experiments in light-matter interaction in nematic liquid crystals, the authors of [5] considered energy minimizing in $H^1(\mathbb{R})$ solutions of the following singular perturbation problem:

$$\epsilon^2 v''(x) + \mu(x)v(x) - v^3(x) + \epsilon a f(x) = 0, \quad x \in \mathbb{R},$$
(1.1)

where a > 0 is a parameter, and the fixed functions μ , f satisfy

$$\begin{cases} \mu \in C^{1}(\mathbb{R}) \cap L^{\infty}(\mathbb{R}) \text{ is even, } \mu' < 0 \text{ in } (0, \infty), \\ \mu(\xi) = 0 \text{ for some } \xi > 0; \\ f \in L^{1}(\mathbb{R}) \cap L^{\infty}(\mathbb{R}) \cap C(\mathbb{R}) \text{ is odd, } f > 0 \text{ in } (0, \infty). \end{cases}$$
(1.2)

It was shown that, for any $\epsilon, a > 0$, minimizers v_{ϵ} always exist and have a unique zero ρ_{ϵ} . The main result of [5] can be summarized as follows. There exist positive numbers $a_* \leq a^*$ (independent of ϵ) that can be characterized variationally as

$$a^{*} = \sup_{x \in [-\xi,0]} \frac{\sqrt{2} \left((\mu(0))^{\frac{3}{2}} - (\mu(x))^{\frac{3}{2}} \right)}{3 \int_{x}^{0} |f| \mu} < \infty,$$

$$a_{*} = \inf_{x \in (-\xi,0]} \frac{\sqrt{2} (\mu(x))^{\frac{3}{2}}}{3 \int_{-\xi}^{x} |f| \mu} \in (0,\infty),$$
(1.3)

such that:

Date: April 17, 2019.

- If $a \in (a^*, \infty)$, then the minimizers v_{ϵ} converge pointwise to $\operatorname{sgn}(x)\sqrt{\mu^+(x)}$ as $\epsilon \to 0$ (the convergence being uniform away from the origin, and so $\rho_{\epsilon} \to 0$). There is a transition layer around the origin of width $O(\epsilon)$, where the behaviour of v_{ϵ} is governed by a usual squeezed hyperbolic tangent profile (see also (2.22) below).
- If $a \in (0, a_*)$, then v_{ϵ} converges uniformly to $\sqrt{\mu^+(x)}$ and $\rho_{\epsilon} \to -\xi$ as $\epsilon \to 0$ (without loss of generality).

In any case, v_{ϵ} has steep corner layers of width $O(\epsilon^{\frac{2}{3}})$ around $\pm\xi$, where the behaviour of v_{ϵ} is governed by a suitable blow-down of a minimizing (in the sense of Morse) solution of the Painlevé-II equation:

$$y''(s) - sy(s) - 2y^{3}(s) - \alpha = 0, \quad s \in \mathbb{R},$$
(1.4)

with

$$\alpha = \frac{af(-\xi)}{\sqrt{2\mu'(-\xi)}} < 0, \tag{1.5}$$

after appropriate reflections and rescalings of constants. More precisely, assuming that $v_{\epsilon} \to \sqrt{\mu^+}$ (without loss of generality), it holds

$$2^{-\frac{1}{2}} \left(\mu'(-\xi)\epsilon\right)^{-\frac{1}{3}} v_{\epsilon} \left(-\xi - \frac{\epsilon^{\frac{2}{3}}s}{\left(\mu'(-\xi)\right)^{\frac{1}{3}}}\right) \to -Y(s) \text{ in } C^{1}_{loc}(\mathbb{R}) \text{ as } \epsilon \to 0, \quad (1.6)$$

where Y is an energy minimizing solution of (1.4) with α as in (1.5). We clarify that here minimality is understood in the sense of Morse or De Giorgi, that is with respect to compact perturbations. In fact, such solutions automatically satisfy either the plus or the minus set of asymptotic conditions in (1.7) below (see again [5]).

It is known that for any $\alpha \leq 0$ the equation (1.4) admits a unique positive solution Y_+ which is defined in the whole real line (see [10] for $\alpha = 0$, and [5, 8] for $\alpha < 0$). In fact, it holds $Y'_+ < 0$, which implies that Y_+ is linearly nondegenerate, in the sense that the linearization of (1.4) about Y_+ does not have bounded elements in its kernel. On the other hand, for $\alpha < 0$ in some neighborhood of zero, the problem (1.4) has also a solution Y_- which changes sign once [17] (we refer to the discussion following Corollary 1.1 for more details and references). We point out that it holds

$$Y_{\pm}(s) \sim \pm \sqrt{-\frac{s}{2}} + O\left(\frac{1}{s}\right) \text{ as } s \to -\infty; \quad Y_{\pm}(s) \sim \frac{|\alpha|}{s} \text{ as } s \to +\infty.$$
 (1.7)

A natural question that arises is whether the minimizing solution Y of (1.4) that appears in (1.6) is Y_+ or a solution that changes sign exactly once (recall that v_{ϵ} changes sign exactly once but its zero ρ could escape at infinity in the blow-up (1.6)). In this paper we will show that it is indeed the second, and more interesting, scenario that occurs. Thus, we describe the microstructure of v_{ϵ} near $-\xi$ (at least in an inner zone of width $O(\epsilon^{\frac{2}{3}})$). We confirm the prediction that

the microscopic phase transition that takes place near $-\xi$ is a new type of defect that does not involve the standard hyperbolic tangent. The fact that this is an open problem was emphasized at the end of the introduction in [6].

Our main result is the following.

Theorem 1.1. If in addition to (1.2) we assume that μ , $f \in C^2$ near $-\xi$, then the minimizing solution Y of (1.4) that appears in (1.6) changes sign exactly once, provided that $a \in (0, a_*)$ with a_* as in (1.3).

Our result implies the following.

Corollary 1.1. If $\alpha \in (-\frac{1}{2}, 0)$, the problem (1.4) admits a solution Y_{-} with exactly one sign change that is energy minimizing with respect to compactly supported perturbations. Moreover, the solution Y_{-} satisfies the corresponding conditions in (1.7).

Let us compare our above result with some related recent ones in the literature. In [4, 8], by solving the corresponding Riemann-Hilbert problem, it was shown that there exists a unique solution of (1.4)- $(1.7)_{-}$ for $\alpha \in (-\frac{1}{2}, 0)$. Moreover, it was shown in the latter reference that this solution is a meromorphic function with no poles on the real line. More recently, the existence of a solution to (1.4)- $(1.7)_{-}$ with exactly one sign change was shown by a shooting argument in [17] for $\alpha \in (\alpha_*, 0)$, where $\alpha_* < 0$ is some number that is sufficiently close to zero. The approach in the latter reference provides a qualitative description of the obtained solution. In particular, it changes monotonicity only once. It is worth mentioning that the existence and uniqueness of such a nonmonotone solution for $\alpha \in (-\frac{1}{2}, 0)$ were indicated in [9]. Interestingly enough, it was also indicated therein that there are no such solutions for $\alpha = -\frac{1}{2}$.

In relation to the aforementioned papers, where different approaches are employed, our main contribution is that we associated to the solution of (1.4)-(1.7) the strong property of energy minimality. In general, in problems of the calculus of variations, the standard method for showing this property is by constructing a calibration. However, it is not clear to us how to apply this considerably more direct approach to the problem at hand. Nevertheless, we note in passing that this is indeed possible for Y_+ , based on the fact that it is sign definite. Actually, it follows from our proof that the validity of Corollary 1.1 holds for $\alpha \in (\alpha_*, 0)$, where α_* is as in the left hand side of (3.1). In this regard, we point out that it was mentioned at the end of the introduction of [5] that numerical evidence suggests that the range of α in Corollary 1.1 should be larger. Lastly, we should mention that by the Bäcklund transformation (see for instance [8]) our result implies the existence of a solution to (1.4)- $(1.7)_-$ for $\alpha \in (-k - \frac{1}{2}, -k), k \in \mathbb{N}$.

Our proof of Theorem 1.1 goes by contradiction. Assuming that the assertion is false puts us sufficiently far from $-\xi$, in some sense, which allows us to compare

 v_{ε} to a sign definite solution η_{ϵ} of (1.1) by examining the quotient

$$w_{\epsilon} = \frac{v_{\epsilon}}{\eta_{\epsilon}}.$$

This division trick was introduced by [13] in a different context and has been used extensively in the study of vortices in Bose-Einstein condensates (see [1] and the references therein); the simplest case being (1.1) with a = 0 and $-\mu$ being a trapping potential. In this regard, we point out that the connection between (1.1) and the aforementioned studies was already discussed in [5]. However, to the best of our knowledge, this important division trick is applied here for the first time in this context. Thankfully, the required estimates for η and its derivatives are readily derivable from [12] or [15]. These rely on the fact that the positive solution Y_+ of (1.4) is linearly nondegenerate. The quotient w satisfies a weighted Allen-Cahn equation with a weight that degenerates at $-\xi$ (see (2.15)) and the discussion leading to (2.24)). However, as we remarked, we will be working sufficiently away from this degeneracy. It then boils down to showing that w cannot have a sharp transition layer from 1 to -1 at ρ . For this purpose, there are several approaches in the literature for related problems of Allen-Cahn type. For example see [3, Thm. 1.5] for a nonlinear Schrödinger equation with a potential, [2, Thm 4.1] for a phase field model of phase transitions, and [14, Thm. 1] for a spatially inhomogeneous Allen-Cahn equation. On the one hand, the equation for w resembles more the aforementioned phase field model. On the other hand, armed with our estimates for w, we found it more convenient to adapt the corresponding proof of [3] with our own twist.

We close this introduction by expressing our hope that some of our arguments can be extended to describe the core of the 'shadow vortex' defect in the recent paper [7] or the 'shadow domain wall' in [6].

In the rest of the paper we will prove our main results. Following the proof of Theorem 2.1, which implies Theorem 1.1, we will present in Remark 2.1 an applied mathematicians point of view which partly motivated our rigorous analysis.

2. Proof of the main results

2.1. **Proof of Theorem 1.1.** In order to prove Theorem 1.1, as was already observed at the end of Section 3 of [5], it suffices to establish the following (recall also the preamble to the aforementioned theorem).

Theorem 2.1. Let μ , f satisfy (1.2) with μ , $f \in C^2$ near $-\xi$. Then, if a is as in Theorem 1.1, the unique root ρ_{ϵ} of v_{ϵ} satisfies

$$\rho_{\epsilon} + \xi = O(\epsilon^{\frac{2}{3}}) \quad as \ \epsilon \to 0.$$

Proof. Throughout the proof, we will denote by C/c > 0 a large/small generic constant that is independent of small $\epsilon > 0$, and whose value will increase/decrease as we proceed.

Firstly, we will show that

$$\rho_{\epsilon} \ge -\xi - C\epsilon^{\frac{2}{3}}.\tag{2.1}$$

To this end, let us consider the algebraic equation that comes from (1.1) when we ignore the term $\epsilon^2 v''$:

$$\mu(x)\nu - \nu^3 + \epsilon a f(x) = 0.$$
(2.2)

Based on the fact that $\mu'(-\xi) > 0$, it is straightforward to verify that there exists a large R > 0 such that the following properties hold. The above equation admits a unique solution ν_{ϵ} for $x \leq -\xi - R\epsilon^{\frac{2}{3}}$, provided that $\epsilon > 0$ is sufficiently small. In fact, it holds

$$\nu_{\epsilon}(x) = -\epsilon a \frac{f(x)}{\mu(x)} + O\left(\frac{\epsilon^3}{|x+\xi|^4}\right) < 0,$$
(2.3)

uniformly on $\left[-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}\right]$ for some small d > 0 (such that $f \in C^2(-\xi - 2d, -\xi + 2d)$), as $\epsilon \to 0$, and

$$|\nu_{\epsilon}'(x)| \le C \frac{\epsilon}{|x+\xi|^2}, \quad |\nu_{\epsilon}''(x)| \le C \frac{\epsilon}{|x+\xi|^3}, \quad x \in [-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}].$$

Let

$$\varphi_{\epsilon}(x) = v_{\epsilon}(x) - \nu_{\epsilon}(x), \quad x \in [-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}]. \tag{2.4}$$

It follows readily that φ_{ϵ} satisfies the following linear equation:

$$-\epsilon^2 \varphi'' + Q(x)\varphi = O\left(\frac{\epsilon^3}{|x+\xi|^3}\right), \qquad (2.5)$$

with

$$Q(x) := \frac{v_{\epsilon}^{3}(x) - \mu(x)v_{\epsilon}(x) - a\epsilon f(x)}{v_{\epsilon}(x) - \nu_{\epsilon}(x)}$$

$$= \varphi^{2} + 3\nu_{\epsilon}^{2} + 3\nu_{\epsilon}\varphi - \mu$$

$$\geq -\mu$$

$$\geq c|x + \xi|, \qquad (2.6)$$

for $x \in [-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}].$

It follows from Lemma 3.1 in [5] that

$$\left| v_{\epsilon} \left(-\xi - R\epsilon^{\frac{2}{3}} \right) \right| \le C\epsilon^{\frac{1}{3}}.$$
(2.7)

On the other side, it follows as in Proposition 2.1 in [11] that

$$\varphi_{\epsilon}(-\xi - d) = O(\epsilon^2) \text{ as } \epsilon \to 0.$$
 (2.8)

In light of (2.5), (2.6), (2.7), (2.8), we deduce by a standard barrier argument that

$$|\varphi_{\epsilon}(x)| \le C\epsilon^{\frac{1}{3}} e^{c\left\{\frac{x+\xi}{\epsilon^{3}}\right\}} + C\epsilon^{2} + C\frac{\epsilon^{3}}{|x+\xi|^{4}}, \quad x \in [-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}].$$

By combining (2.3), (2.4) and the above relation, we arrive at

$$v_{\epsilon}(x) = -\epsilon a \frac{f(x)}{\mu(x)} \left[1 + O\left(\frac{\epsilon^2}{|x+\xi|^3}\right) + O\left(\frac{|x+\xi|}{\epsilon^{\frac{2}{3}}}e^{c\left\{\frac{x+\xi}{\epsilon^{\frac{2}{3}}}\right\}}\right) + O\left(\epsilon|x+\xi|\right) \right],$$

uniformly on $[-\xi - d, -\xi - R\epsilon^{\frac{2}{3}}]$, as $\epsilon \to 0$. Assuming that (2.1) does not hold, then we can evaluate the above relation at $x = \rho_{\epsilon}$ along a contradicting sequence of ϵ 's that tend to zero such that

$$\frac{\rho_{\epsilon} + \xi}{\epsilon^{\frac{2}{3}}} \to -\infty.$$

This gives us that

$$1 + O\left(\frac{\epsilon^2}{|\rho_{\epsilon} + \xi|^3}\right) + O\left(e^{c\left\{\frac{\rho_{\epsilon} + \xi}{2}\right\}}\right) + o\left(\epsilon\right) = 0,$$

which is clearly a contradiction. We have thus established the desired lower bound in (2.1).

The main effort will be to establish the 'opposite direction' of (2.1). To this end, we will argue by contradiction. Assuming that the assertion of the theorem is false would give us a sequence $\epsilon_n \to 0$ such that

$$\frac{\rho_{\epsilon_n} + \xi}{\epsilon_n^{\frac{2}{3}}} \to +\infty.$$
(2.9)

Abusing notation, we will drop from now on the subscript n and assume that all the following $\epsilon's$ are along this sequence or a subsequence of it.

Let η_{ϵ} be the unique negative solution of

$$\begin{cases} \epsilon^2 \eta''(x) + \mu(x)\eta(x) - \eta^3(x) + \epsilon af = 0, \ x < 0; \\ \eta(-\infty) = 0, \ \eta'(0) = 0. \end{cases}$$

In passing, we note that uniqueness follows as in Remark 9 in [16]. For future purposes, we point out that the method of the aforementioned reference yields that

$$\frac{v_{\epsilon}(x)}{\eta_{\epsilon}(x)} < 1, \quad x < \rho_{\epsilon}.$$
(2.10)

Fine estimates for the convergence $\eta_{\epsilon} \to -\sqrt{\mu^+}$ uniformly as $\epsilon \to 0$ are available from [12] (because the positive solution of (1.4) is nondegenerate, as we have remarked). In particular, η_{ϵ} satisfies (1.6) with $Y = Y_+$. Essentially we will only need that there exist constants $C, \delta > 0$ such that

$$-C\sqrt{x+\xi} \le \eta_{\epsilon}(x) \le -\frac{1}{C}\sqrt{x+\xi}, \quad \epsilon^{\frac{2}{3}} \le x+\xi \le \delta,$$

$$\eta_{\epsilon}(x) \sim -\sqrt{\mu'(-\xi)(x+\xi)}, \quad \epsilon^{\frac{2}{3}} \ll x+\xi \le \delta,$$

(2.11)

and that the above relations can be differentiated with respect to x in the obvious way (see also Appendix A in [1]). For a bit more refined estimates we refer to Remark 2.1 below.

Using a well known trick from [13], suppressing the dependence on ϵ , we write

$$v = \eta w, \tag{2.12}$$

for some w such that

$$w > 0 \text{ for } x < \rho; \quad w < 0 \text{ for } x > \rho.$$
 (2.13)

For future purposes, let us note that the upper bound

$$|v(x)| \le C\left(\sqrt{x+\xi} + \epsilon^{\frac{1}{3}}\right), \quad x \in (-\xi, 0),$$

which follows at once from Lemma 3.1 in [5], together with the identical lower bound which follows from (2.11) imply that

$$|w(x)| \le C, \ x \in (-\xi, 0].$$
 (2.14)

We find that w satisfies

$$\epsilon^2 w'' + 2\epsilon^2 \frac{\eta'}{\eta} w' + \eta^2 \left(w - w^3 \right) + \epsilon a \frac{f(x)}{\eta} (1 - w) = 0, \quad x < 0.$$
 (2.15)

Next, we stretch variables around $x = \rho$, setting

$$W(y) = w(x)$$
 where $y = \frac{x - \rho}{\tilde{\epsilon}}$

with $\tilde{\epsilon} = \tilde{\epsilon}(\epsilon) > 0$ to be determined such that

$$\tilde{\epsilon} \ll \epsilon^{\frac{2}{3}}.$$
 (2.16)

We will denote with the derivative with respect to y. We obtain from (2.15) that

$$\ddot{W} + 2\tilde{\epsilon}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{W} + \frac{\tilde{\epsilon}^2}{\epsilon^2}\eta^2\left(\rho + \tilde{\epsilon}y\right)\left(W - W^3\right) + a\frac{\tilde{\epsilon}^2}{\epsilon}\frac{f}{\eta}\left(\rho + \tilde{\epsilon}y\right)\left(1 - W\right) = 0$$
(2.17)

for $y < -\rho/\tilde{\epsilon}$. In light of (2.9), (2.11) and the working assumption (2.16), it holds

$$\frac{\tilde{\epsilon}^2}{\epsilon^2}\eta^2(\rho+\tilde{\epsilon}y) \sim \mu'(-\xi)\frac{\tilde{\epsilon}^2}{\epsilon^2}(\rho+\xi) \text{ as } \epsilon \to 0$$
(2.18)

(for fixed y). Therefore, we choose

$$\tilde{\epsilon} = \frac{\epsilon}{\sqrt{\rho + \xi}}.$$
(2.19)

Recalling (2.9), this choice clearly satisfies (2.16). The point is that we want the third term in (2.17) to be of the same order as the first one; the other terms will turn out to be of smaller order. Then, equation (2.17) becomes

$$\ddot{W} + 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{W} + \frac{\eta^2\left(\rho + \tilde{\epsilon}y\right)}{\rho + \xi}\left(W - W^3\right) + a\frac{\epsilon}{\rho + \xi}\frac{f}{\eta}\left(\rho + \tilde{\epsilon}y\right)\left(1 - W\right) = 0$$
(2.20)

By virtue of (2.14), we have that W is bounded locally with respect to ϵ . Hence, using standard elliptic estimates and the usual diagonal argument, keeping in mind (2.13), passing to a further subsequence if necessary, we find that

$$W(y) \to W_0(y) \text{ in } C^1_{loc}(\mathbb{R}) \text{ as } \epsilon \to 0,$$
 (2.21)

where W_0 satisfies

$$\frac{1}{\mu'(-\xi)}\ddot{W}_0 + W_0 - W_0^3 = 0, \quad y \in \mathbb{R}; \quad yW_0(y) \le 0, \quad y \in \mathbb{R}.$$

We claim that W_0 is nontrivial, which would imply that

$$W_0(y) \equiv -\tanh\left(\sqrt{\frac{\mu'(-\xi)}{2}}y\right). \tag{2.22}$$

In passing, we note that the explicit formula for W_0 is not important for our purposes, we will essentially use that

$$W_0(y) \to \mp 1 \text{ as } y \to \pm \infty.$$
 (2.23)

To this end, let Z_{ϵ} be the unique positive solution of the following boundary value problem:

$$\ddot{Z} + 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{Z} + \frac{\eta^2\left(\rho + \tilde{\epsilon}y\right)}{\rho + \xi}\left(Z - Z^3\right) = 0, \quad y \in \left(\frac{-\xi + \epsilon^{\frac{2}{3}} - \rho}{\tilde{\epsilon}}, 0\right);$$
$$Z\left(\frac{-\xi + \epsilon^{\frac{2}{3}} - \rho}{\tilde{\epsilon}}\right) = \frac{1}{4}, \quad Z(0) = 0.$$

As will become apparent shortly, the specific value 1/4 is not of importance. The existence of such a Z_{ϵ} follows by directly minimizing the associated energy functional

$$E_{\epsilon}(Z) = \int_{(-\xi+\epsilon^{\frac{2}{3}}-\rho)\tilde{\epsilon}^{-1}}^{0} \left\{ \frac{\eta^2 \left(\rho + \tilde{\epsilon}y\right)}{2} (\dot{Z})^2 + \frac{\eta^4 \left(\rho + \tilde{\epsilon}y\right)}{\rho + \xi} \frac{\left(1 - Z^2\right)^2}{4} \right\} dy \quad (2.24)$$

subject to the above boundary conditions. Moreover, simple energy considerations give that $0 < Z_{\epsilon} < 1$. The uniqueness comes from the observation that $\{tZ_{\epsilon} : t \in (0,1)\}$ is a family of lower solutions to the above boundary value problem and Serrin's sweeping principle (see [16] and the references therein). In fact, by virtue of (1.2), we note that these are also lower solutions to the equation (2.20) in the same interval. As a consequence of the assumption (2.9), the blow-up analysis in [5] yields that η_{ϵ} and v_{ϵ} share the same first order term in their corresponding inner expansions around $-\xi$. Hence, both η_{ϵ} and v_{ϵ} satisfy (1.6) with $Y = -Y_{+}$. In particular, given any L > 0, it holds

$$W_{\epsilon}\left(\frac{-\xi - \rho + L\epsilon^{\frac{2}{3}}}{\tilde{\epsilon}}\right) \to 1 \text{ as } \epsilon \to 0.$$
 (2.25)

Thus, since $W_{\epsilon}\left((-\xi + \epsilon^{\frac{2}{3}} - \rho)\tilde{\epsilon}^{-1}\right) > 1/2$, $W_{\epsilon}(0) = 0$ and $W_{\epsilon} > 0$ in between, we deduce by Serrin's sweeping principle that

$$W_{\epsilon}(y) > Z_{\epsilon}(y), \quad y \in \left((-\xi + \epsilon^{\frac{2}{3}} - \rho)\tilde{\epsilon}^{-1}, 0 \right).$$

On the other hand, as in Proposition 2.3 in [14], we have

$$Z_{\epsilon} \to -\tanh\left(\sqrt{\frac{\mu'(-\xi)}{2}}y\right)$$
 in $C^{1}_{loc}\left((-\infty,0]\right)$ as $\epsilon \to 0$.

The above two relations imply that W_0 is nontrivial, and thus is given by (2.22) as claimed.

Let us consider now (2.20) without derivatives, which recalling (2.19) and after dividing by 1 - W becomes

$$\Sigma^2 + \Sigma + a\epsilon \frac{f}{\eta^3}(\rho + \tilde{\epsilon}y) = 0.$$

The above algebraic equation can be solved explicitly and has the following two solutions:

$$\Sigma_{\pm}(y) = \frac{-1 \pm \sqrt{1 - 4a\epsilon \frac{f}{\eta^3}(\rho + \tilde{\epsilon}y)}}{2}$$

It follows from (2.9) and (2.11) that there is some fixed large D > 0 such that these are real valued for

$$-\xi + D\epsilon^{\frac{2}{3}} \le \rho + \tilde{\epsilon}y \le \delta,$$

provided that $\epsilon > 0$ is sufficiently small. From now on let us fix a D such that

$$0 < 1 + \Sigma_{-}(y) < \frac{1}{100}$$
 and $-\frac{1}{100} < \Sigma_{+}(y) < 0$ (2.26)

in the aforementioned interval, for sufficiently small $\varepsilon > 0$ (the number 100 has no significance here).

Let

$$\Phi(y) = 1 - W(y), \quad y \in \left[D\sqrt{\rho + \xi} \epsilon^{-\frac{1}{3}} - \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}, 0 \right].$$
(2.27)

We know that

$$0 < \Phi < 1 \tag{2.28}$$

(recall (2.10)). We can write (2.20) as

$$-\ddot{\Phi} - 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{\Phi} + \frac{\eta^2(\rho + \tilde{\epsilon}y)}{\rho + \xi}\left(W - \Sigma_{-}(y)\right)\left(W - \Sigma_{+}(y)\right)\Phi = 0.$$
(2.29)

By virtue of (2.21), (2.23), (2.25) with L = D, and (2.26), we deduce from (2.20) via the maximum principle that there exists an M > 0 such that

$$0 < 1 - W(y) < \frac{1}{100}, \quad y \in \left[D\sqrt{\rho + \xi} \epsilon^{-\frac{1}{3}} - \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}, -M \right], \quad (2.30)$$

provided that $\epsilon > 0$ is sufficiently small. More precisely, we observe that $\underline{W}_t = 1 - t$, $\frac{1}{100} < t \leq 1$, is a family of lower solutions to (2.20) which allows to use Serrin's sweeping principle since W > 0, see [16] and the references therein. Concerning the second term in (2.29), it follows from (2.11) that

$$0 < \frac{\epsilon}{\sqrt{\rho + \xi}} \frac{\eta'}{\eta} \left(\rho + \tilde{\epsilon}y\right) \le C \frac{\epsilon^{\frac{1}{3}}}{\sqrt{\rho + \xi}}, \quad y \in \left[D\sqrt{\rho + \xi}\epsilon^{-\frac{1}{3}} - \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}, 0\right]; \quad (2.31)$$

concerning the last term, we obtain from (2.11) and (2.30) that

$$C \ge \frac{\eta^2(\rho + \tilde{\epsilon}y)}{\rho + \xi} \left(W - \Sigma_- \right) \left(W - \Sigma_+ \right) \ge \frac{1}{C}, \quad y \in \left[-\frac{2(\rho + \xi)^{\frac{3}{2}}}{3\epsilon}, -M \right].$$
(2.32)

So, by a standard barrier argument, we deduce from (2.28), (2.29), (2.31) and the above relation that

$$0 < \Phi(y) \le e^{cy} + e^{-c\left(y + \frac{2(\rho+\xi)^{\frac{3}{2}}}{3\epsilon}\right)}, \quad y \in \left[-\frac{2(\rho+\xi)^{\frac{3}{2}}}{3\epsilon}, -M\right].$$

Consequently, by recalling the definition of Φ from (2.27), we infer that

$$0 < 1 - W(y) \le e^{cy} + e^{-c\left\{\frac{\rho+\xi}{2}\right\}}, \quad y \in \left[-\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon} - 1, 0\right].$$
(2.33)

In turn, by standard elliptic estimates, via (2.29), (2.31) and the upper bound in (2.32), we get that

$$\left|\dot{W}(y)\right| \le Ce^{cy} + Ce^{-c\left\{\frac{\rho+\xi}{2}\right\}}, \quad y \in \left[-\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon}, 0\right].$$
 (2.34)

Let us write (2.20) as

$$\ddot{W} + 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{W} + \frac{\partial}{\partial W}G(W, y) = 0, \qquad (2.35)$$

with

$$G(W,y) = -\frac{\eta^2(\rho + \tilde{\epsilon}y)}{\rho + \xi} \frac{(1 - W^2)^2}{4} - a \frac{\epsilon}{\rho + \xi} \frac{f}{\eta} \left(\rho + \tilde{\epsilon}y\right) \frac{(1 - W)^2}{2}.$$
 (2.36)

When multiplied by \dot{W} , equation (2.35) reads as

$$\frac{d}{dy}\left(\frac{(\dot{W})^{2}}{2}\right) + 2\frac{\epsilon}{\sqrt{\rho+\xi}}\frac{\eta'}{\eta}\left(\rho+\tilde{\epsilon}y\right)(\dot{W})^{2} + \frac{d}{dy}\left(G(W,y)\right) \\ + \frac{\epsilon}{(\rho+\xi)^{\frac{3}{2}}}\eta\eta'\left(\rho+\tilde{\epsilon}y\right)\frac{\left(1-W^{2}\right)^{2}}{2} \\ -a\frac{\epsilon^{2}}{(\rho+\xi)^{\frac{3}{2}}}\frac{f\eta'}{\eta^{2}}\left(\rho+\tilde{\epsilon}y\right)\frac{(1-W)^{2}}{2} + a\frac{\epsilon^{2}}{(\rho+\xi)^{\frac{3}{2}}}\frac{f'}{\eta}\left(\rho+\tilde{\epsilon}y\right)\frac{(1-W)^{2}}{2} = 0.$$
(2.37)

We shall next integrate the above relation over

$$y \in I = \left(-\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon}, (\delta-\rho)\frac{\sqrt{\rho+\xi}}{\epsilon}\right) = (\alpha, \beta).$$

The following fact will be useful in the sequel. By the definition (2.12) of w, and standard estimates for v and η that hold uniformly away from their corner layer at $-\xi$ (see Proposition 2.1 in [11]), we infer that

$$\|w(x) - \Sigma_{-}\left(\frac{x-\rho}{\tilde{\epsilon}}\right)\|_{C^{1}\left[-\xi+\frac{\delta}{2},-\xi+2\delta\right]} \le C\epsilon^{2}.$$
(2.38)

The integral of the first term in (2.37) is plainly the half of

$$\begin{split} (\dot{W})^2 \left(\beta\right) - (\dot{W})^2 \left(\alpha\right) &= \frac{\epsilon^2}{\rho + \xi} (w')^2 \left(-\xi + \delta\right) - (\dot{W})^2 \left(-\frac{(\rho + \xi)^3}{2\epsilon}\right) \\ &= O\left(\frac{\epsilon^4}{\rho + \xi}\right) + O\left(\left(\frac{\epsilon^3}{\rho + \xi}\right)^{10}\right) \\ &= O\left(\left(\frac{\epsilon^3}{\rho + \xi}\right)^6\right), \end{split}$$

where we used (2.34), and (2.38) which implies that $w'(-\xi + \delta) = O(\epsilon)$ (recall also the formula for Σ_{-}). We point out in passing that the rough estimate $w'(-\xi + \delta) = O(1)$ would have actually sufficed for our purposes. Regarding the second term in (2.37), the fact that it is nonnegative will suffice for the time being. Using (2.33) and (2.38), we find from the definition of G in (2.36) that the integral of the third term in (2.37) can be estimated as follows:

$$\int_{I} \frac{d}{dy} \left(G(W, y) \right) dy = G \left(W(\beta), \beta \right) - G \left(W(\alpha), \alpha \right) = O \left(\frac{\epsilon}{\rho + \xi} \right).$$

Concerning the fourth term in (2.37), thanks to (2.11), we find that

$$\int_{I} \eta \eta' \left(\rho + \tilde{\epsilon}y\right) \frac{\left(1 - W^2\right)^2}{2} dy \ge c \int_{I} \left(1 - W^2\right)^2 dy.$$

Concerning the fifth term in (2.37), recalling (2.14) and using once more (2.11), we have

$$\left| \int_{I} \frac{f\eta'}{\eta^{2}} \left(\rho + \tilde{\epsilon}y \right) \frac{(1-W)^{2}}{2} dy \right| \leq C \int_{I} \left(\rho + \xi + \tilde{\epsilon}y \right)^{-\frac{3}{2}} dy$$
$$\leq C \tilde{\epsilon}^{-1} (\rho + \xi)^{-\frac{1}{2}}$$
(2.39)

 $\leq \frac{C}{\epsilon}.$

Finally, concerning the last term in (2.37), we obtain similarly that

$$\left| \int_{I} \frac{f'}{\eta} \left(\rho + \tilde{\epsilon} y \right) \frac{(1-W)^2}{2} dy \right| \le \frac{C}{\epsilon}$$

Collecting all of the above, we infer from the integration of (2.37) over I that

$$\int_{I} \left(1 - W^2\right)^2 dy \le C.$$
 (2.40)

We claim that (2.40) implies that there exist c, N > 0 such that

$$W(y) \le -c, \quad y \in \left[N, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right],$$

$$(2.41)$$

if $\epsilon > 0$ is sufficiently small. Indeed, if not, then there would exist points

$$p_{\epsilon} \in \left(0, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right] \quad \text{with} \quad p_{\epsilon} \to +\infty$$
 (2.42)

such that

$$W(p_{\epsilon}) \to 0.$$

Let

$$\tilde{W}(y) = W(y+p).$$

Then, we have

$$\ddot{\tilde{W}} + 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}p + \tilde{\epsilon}y\right)\dot{\tilde{W}} + \frac{\eta^2\left(\rho + \tilde{\epsilon}p + \tilde{\epsilon}y\right)}{\rho + \xi}\left(\tilde{W} - \tilde{W}^3\right) + a\frac{\epsilon}{\rho + \xi}\frac{f}{\eta}\left(\rho + \tilde{\epsilon}p + \tilde{\epsilon}y\right)\left(1 - \tilde{W}\right) = 0$$
(2.43)

(at least) for $y \in \left[-1, 100 \frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right]$, $\tilde{W} < 0$ therein, and $\tilde{W}(0) \to 0$. Then, since $\tilde{\epsilon}p \to 0$, we can argue as we did for (2.21) to find that $\tilde{W} \to \tilde{W}_0$ in $C^1_{loc}\left([-1,\infty)\right)$,

where

$$\frac{1}{\mu'(-\xi)}\ddot{\tilde{W}}_0 + \tilde{W}_0 - \tilde{W}_0^3 = 0, \quad y > -1; \quad \tilde{W}_0(y) \le 0, \quad y > -1, \quad \tilde{W}_0(0) = 0.$$

It is clear that $\tilde{W}_0 \equiv 0$. So, given any K > 1, it holds

$$\int_{I} (1 - W^{2})^{2} dy \ge \int_{p}^{p+K} (1 - W^{2})^{2} dy = \int_{0}^{K} (1 - \tilde{W}^{2})^{2} dy \ge \frac{K}{2},$$

provided that $\epsilon > 0$ is sufficiently small, which contradicts (2.40) and establishes the validity of (2.41).

The estimate (2.40) yields that

$$\int_{5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}}^{10\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}} \left(1-W^2\right)^2 dy \le C,$$

3

which implies that

$$|-1 - W(q)| < \frac{1}{100}$$
 for some $q \in \left(5\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}, 10\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right).$ (2.44)

We may further assume that the N in (2.41) is such that

$$-1 < W_0(N) < -1 + \frac{1}{1000},$$

(recall (2.23)). Then, thanks to (2.21), (2.26) and (2.44), as we did for (2.30) we have that

$$W(y) > -1 - \frac{1}{100}, \ y \in \left[N, 5\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right]$$

On the other side, the relations (2.41) and (2.44) allow us to apply the same argument in the opposite direction, and thus arrive at

$$|W(y) + 1| < \frac{1}{100}, \quad y \in \left[N, 5\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right].$$
 (2.45)

Let now

$$\Psi(y) = W(y) - \Sigma_{-}(y), \quad y \in \left[0, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right].$$

We can write (2.20) as

$$\ddot{\Psi} + 2\frac{\epsilon}{\sqrt{\rho+\xi}}\frac{\eta'}{\eta}\left(\rho+\tilde{\epsilon}y\right)\dot{\Psi} + \frac{\eta^2(\rho+\tilde{\epsilon}y)}{\rho+\xi}\left(1-W\right)\left(W-\Sigma_+(y)\right)\Psi = -\ddot{\Sigma}_- - 2\frac{\epsilon}{\sqrt{\rho+\xi}}\frac{\eta'}{\eta}\left(\rho+\tilde{\epsilon}y\right)\dot{\Sigma}_-.$$
(2.46)

We note that

$$-1 - \Sigma_{-}(y) \leq C\epsilon(\rho + \xi + \tilde{\epsilon}y)^{-\frac{3}{2}}, \qquad (2.47)$$

$$\left|\dot{\Sigma}_{-}(y)\right| \le C \frac{\epsilon^2}{\sqrt{\rho+\xi}} (\rho+\xi+\tilde{\epsilon}y)^{-\frac{5}{2}}, \quad \left|\ddot{\Sigma}_{-}(y)\right| \le C \frac{\epsilon^3}{\rho+\xi} (\rho+\xi+\tilde{\epsilon}y)^{-\frac{7}{2}}, \quad (2.48)$$

for $y \in \left[0, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right]$. Thus, since $0 < \frac{\eta'}{\eta} \left(\rho + \tilde{\epsilon}y\right) \le \frac{C}{\rho+\xi}, \quad y \in \left[-\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon}, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right], \quad (2.49)$

equation (2.46) becomes

$$\ddot{\Psi} + 2\frac{\epsilon}{\sqrt{\rho + \xi}}\frac{\eta'}{\eta}\left(\rho + \tilde{\epsilon}y\right)\dot{\Psi} + \frac{\eta^2(\rho + \tilde{\epsilon}y)}{\rho + \xi}\left(1 - W\right)\left(W - \Sigma_+(y)\right)\Psi = O\left(\frac{\epsilon^3}{\rho + \xi}\left(\rho + \xi + \tilde{\epsilon}y\right)^{-\frac{\tau}{2}}\right),$$
(2.50)

for $y \in \left[0, 5\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right]$. By virtue of (2.45), as we did previously for Φ (also keep in mind (2.14)), we get that

$$|W - \Sigma_{-}| \le C \frac{\epsilon^{3}}{(\rho + \xi)^{\frac{9}{2}}} + Ce^{-cy} + Ce^{c\left(y - 5\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right)}, \quad y \in \left[0, 5\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right].$$

Thus, it holds

$$|W - \Sigma_{-}| \le C \frac{\epsilon^{3}}{(\rho + \xi)^{\frac{9}{2}}} + Ce^{-cy}, \quad y \in \left[0, 4\frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right].$$
(2.51)

In turn, by using the upper bound

$$\frac{\eta^2(\rho + \tilde{\epsilon}y)}{\rho + \xi} \le C, \quad y \le 5 \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon},$$

we deduce from (2.49), (2.50), (2.51), and standard elliptic estimates that

$$\left|\dot{W} - \dot{\Sigma}_{-}\right| \le C \frac{\epsilon^{3}}{(\rho + \xi)^{\frac{9}{2}}} + Ce^{-cy}, \quad y \in \left[0, \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon}\right].$$
 (2.52)

By combining (2.47) and (2.51), we find that

$$|W+1| \le C \frac{\epsilon}{(\rho+\xi)^{\frac{3}{2}}} + Ce^{-cy}, \ y \in \left[0, \frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right].$$
 (2.53)

By combining (2.48) and (2.52), we get that

$$\left|\dot{W}\right| \le C \frac{\epsilon^2}{(\rho+\xi)^3} + Ce^{-cy}, \quad y \in \left[0, \frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon}\right]. \tag{2.54}$$

We shall integrate the above relation over

$$y \in J = \left(-\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}, \frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}\right) = (\tilde{\alpha}, \tilde{\beta}),$$

with N to be chosen sufficiently large but independent of ϵ . Below we will carefully estimate each term in the resulting relation. Thanks to (2.34) and (2.54), in regards to the first term in (2.37), we find that

$$(\dot{W})^2 \left(\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}\right) - (\dot{W})^2 \left(-\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}\right) = O\left(\frac{\epsilon^4}{(\rho+\xi)^6}\right),$$

as $\epsilon \to 0$. Regarding the second term in (2.37), thanks to (2.11) and (2.21), we have

$$0 < \frac{(\rho + \xi)^{\frac{3}{2}}}{\epsilon} \frac{\epsilon}{\sqrt{\rho + \xi}} \frac{\eta'}{\eta} \left(\rho + \tilde{\epsilon}y\right) (\dot{W})^2 \to \frac{1}{2} (\dot{W}_0)^2 \text{ in } C_{loc}(\mathbb{R}) \text{ as } \epsilon \to 0.$$

Therefore, we obtain from (2.34), (2.49), (2.54), and Lebesgue's dominated convergence theorem that

$$\frac{(\rho+\xi)^{\frac{3}{2}}}{\epsilon} \frac{\epsilon}{\sqrt{\rho+\xi}} \int_{J} \frac{\eta'}{\eta} \left(\rho+\tilde{\epsilon}y\right) (\dot{W})^{2} dy \to \frac{1}{2} \int_{-\infty}^{\infty} (\dot{W}_{0})^{2} dy.$$
(2.55)

Recalling (2.33), (2.36) and (2.53), the integral of the third term in (2.37) can be estimated as follows:

$$\int_{J} \frac{d}{dy} \left(G(W, y) \right) dy = G\left(W(\tilde{\beta}), \tilde{\beta} \right) - G\left(W(\tilde{\alpha}), \tilde{\alpha} \right)$$
$$= \frac{2a}{\sqrt{1 + 1/N}} \frac{f(-\xi)}{\sqrt{\mu'(-\xi)}} \frac{\epsilon}{(\rho + \xi)^{\frac{3}{2}}} \left(1 + o(1) \right),$$

as $\epsilon \to 0$. Concerning the fourth term in (2.37), by working as we did for (2.55), using (2.33) and (2.53), we find that

$$\int_{J} \eta \eta' \left(\rho + \tilde{\epsilon}y\right) \frac{\left(1 - W^{2}\right)^{2}}{2} dy \to \frac{\mu'(-\xi)}{4} \int_{-\infty}^{\infty} \left(1 - W_{0}^{2}\right)^{2} dy = \frac{1}{2} \int_{-\infty}^{\infty} (\dot{W}_{0})^{2} dy.$$

Concerning the fifth term in (2.37), keeping in mind (2.11) and (2.14), we have

$$\int_{J} \frac{f\eta'}{\eta^{2}} \left(\rho + \tilde{\epsilon}y\right) \frac{(1-W)^{2}}{2} dy = \int_{-\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}}^{\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}} \frac{f\eta'}{\eta^{2}} \left(\rho + \tilde{\epsilon}y\right) \frac{(1-W)^{2}}{2} dy$$

$$\leq \frac{C}{(\rho+\xi)^{\frac{3}{2}}} \int_{-\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}}^{\frac{(\rho+\xi)^{\frac{3}{2}}}{N\epsilon}} 1 dy$$

$$\leq \frac{C}{N\epsilon},$$
(2.56)

3

where the above constant C is independent of both large N and small ϵ . Concerning the last term in (2.37), by using (2.14) and that $-\eta \ge c\sqrt{\rho + \xi}$ in J, we get that

$$\left| \int_{J} \frac{f'}{\eta} \left(\rho + \tilde{\epsilon} y \right) \frac{(1-W)^2}{2} dy \right| \le C \frac{\rho + \xi}{\epsilon}.$$

Putting all the above in the integral of (2.37) over J gives us that

$$\frac{\epsilon}{(\rho+\xi)^{\frac{3}{2}}} \left| \frac{3}{2} \int_{-\infty}^{\infty} (\dot{W}_0)^2 dy + \frac{2a}{\sqrt{1+1/N}} \frac{f(-\xi)}{\sqrt{\mu'(-\xi)}} + o(1) \right| \le C \frac{\epsilon}{\sqrt{\rho+\xi}} + \frac{C}{N} \frac{\epsilon}{(\rho+\xi)^{\frac{3}{2}}}$$
(2.57)

as $\epsilon \to 0$, where the above constant C is independent of both large N and small ϵ . By letting first $\epsilon \to 0$ and then $N \to \infty$ in the above relation, we infer that

$$\frac{3}{2} \int_{-\infty}^{\infty} (\dot{W}_0)^2 dy + 2a \frac{f(-\xi)}{\sqrt{\mu'(-\xi)}} = 0.$$

We note that

$$\int_{-\infty}^{\infty} (\dot{W}_0)^2 dy = -\frac{\sqrt{\mu'(-\xi)}}{\sqrt{2}} \int_{-\infty}^{\infty} (1 - W_0^2) \dot{W}_0 dy = \sqrt{2}\sqrt{\mu'(-\xi)}.$$

So, we have been led to a contradiction unless

$$a = -\frac{3}{2\sqrt{2}} \frac{\mu'(-\xi)}{f(-\xi)}.$$
(2.58)

Therefore, to complete the proof, it just remains to exclude the above possibility. To this end, we will estimate the integral in (2.39) over $J \subset I$ from below with, say, N = 2 in the definition of the interval J. We find as above that

$$\int_{J} \frac{f\eta'}{\eta^{2}} \left(\rho + \tilde{\epsilon}y\right) \frac{(1-W)^{2}}{2} dy \geq \int_{0}^{\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon}} \frac{f\eta'}{\eta^{2}} \left(\rho + \tilde{\epsilon}y\right) \frac{(1-W)^{2}}{2} dy$$
$$\geq \frac{c}{(\rho+\xi)^{\frac{3}{2}}} \int_{0}^{\frac{(\rho+\xi)^{\frac{3}{2}}}{2\epsilon}} 1 dy$$
$$\geq \frac{c}{\epsilon}.$$

Hence, keeping in mind the upper bound in (2.39) and that $J \subset I$, we get that

$$\epsilon \int_J \frac{f\eta'}{\eta^2} \left(\rho + \tilde{\epsilon}y\right) \frac{(1-W)^2}{2} dy \to c$$

for some $c_* \in (0, \infty)$. Now, assuming to the contrary that *a* was as in (2.58), instead of (2.57) we end up with

$$\frac{\epsilon}{(\rho+\xi)^{\frac{3}{2}}}c_* = O\left(\frac{\epsilon}{\sqrt{\rho+\xi}}\right) + \frac{o(\epsilon)}{(\rho+\xi)^{\frac{3}{2}}},$$

as $\epsilon \to 0$. Clearly we have been led again to a contradiction, which completes the proof of the theorem.

Remark 2.1. Let us give the formal argument that led us to the rigorous analysis following (2.21). According to folklore, we write

 $W(y) = W_0(y) + \hat{\epsilon}W_1(y) + higher \text{ order terms},$

and try to find $\hat{\epsilon}$ and W_1 by plugging this into (2.20).

Firstly, we will need the following estimates for $-\eta$ near $-\xi$ that follow in analogy to [12, Thm. 1.1]. It holds

$$\eta_{\epsilon}(x) = \nu_{\epsilon}(x) + O\left(\epsilon^{2}\right) x^{-\frac{5}{2}}, \ x \in \left(-\xi + \epsilon^{\frac{2}{3}}, -\xi + \delta\right),$$

uniformly as $\epsilon \to 0$, for some fixed small $\delta > 0$, where $\nu_{\epsilon} < 0$ solves the algebraic equation (2.2) near $-\xi$. It follows readily that

$$\nu_{\epsilon}(x) = -\sqrt{\mu'(-\xi)(x+\xi)} + \frac{af(-\xi)}{2\mu'(-\xi)}\frac{\epsilon}{x+\xi} + O(\epsilon)$$

The above together with the relations (2.9) and (2.11) yield the following:

$$\begin{aligned} \frac{1}{\rho+\xi}\eta^2\left(\rho+\tilde{\epsilon}y\right) &= \mu'(-\xi)+\mu'(-\xi)\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}}y - \frac{af(-\xi)}{\sqrt{\mu'(-\xi)}}\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}} \\ &+ o\left(\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}}\right)\left(|y|+1\right), \\ \frac{\epsilon}{\sqrt{\rho+\xi}}\frac{\eta'}{\eta}\left(\rho+\tilde{\epsilon}y\right) &= \frac{\epsilon}{2(\rho+\xi)^{\frac{3}{2}}} + o\left(\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}}\right)\left(|y|+1\right), \\ \frac{\epsilon}{\rho+\xi}\frac{f}{\eta}\left(\rho+\tilde{\epsilon}y\right) &= -\frac{f(-\xi)}{\sqrt{\mu'(-\xi)}}\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}} + o\left(\frac{\epsilon}{\left(\rho+\xi\right)^{\frac{3}{2}}}\right)\left(|y|+1\right), \end{aligned}$$

uniformly for $|\tilde{\epsilon}y| \leq \delta$, as $\epsilon \to 0$.

Taking the above into account, we find that

$$\hat{\epsilon} = \frac{\epsilon}{\rho + \xi}$$

and

$$\ddot{W}_1 + (1 - 3W_0^2)W_1 = -\frac{\mu'(-\xi)}{2}y(W_0 - W_0^3) - (\dot{W}_0)^2 + \frac{af(-\xi)}{\sqrt{\mu'(-\xi)}}(W_0 - W_0^3) + \frac{af(-\xi)}{\sqrt{\mu'(-\xi)}}(1 - W_0)$$

for $y \in \mathbb{R}$. However, the last term does not decay to zero as $y \to +\infty$, which is necessary in order to get a solution W_1 that decays to zero as $|y| \to \infty$.

3. Proof of Corollary 1.1

Proof. In view of (1.3) and (1.5), for $\alpha \in (\alpha_*, 0)$ with

$$\alpha_* = \alpha_*(\xi, \mu, f) = \frac{a_*f(-\xi)}{\sqrt{2}\mu'(-\xi)} = \frac{f(-\xi)}{3\mu'(-\xi)} \inf_{x \in (-\xi, 0]} \frac{(\mu(x))^{\frac{1}{2}}}{\int_{-\xi}^x |f|\sqrt{\mu}},$$

our Theorem 1.1 implies the existence of a solution to (1.4), given by appropriate reflections and modulations of Y from (1.6), that changes sign exactly once and is energy minimizing with respect to compactly supported perturbations. The latter property then implies that this solution also satisfies the asymptotic behaviour (1.7) (see [5, Sec. 4]).

It thus remains to show that

$$\inf_{\{(\xi,\mu,f) \text{ as in Thm. } 1.1\}} \alpha_*(\xi,\mu,f) \le -\frac{1}{2}.$$
(3.1)

To this end, as was observed in [5], the special choice $f = -\frac{\mu'}{2}$ yields that $a_* = \sqrt{2}$. So, this choice gives $\alpha_* = -\frac{1}{2}$ which suffices for the proof of the corollary.

References

- Amandine Aftalion, Benedetta Noris, and Christos Sourdis. Thomas-Fermi approximation for coexisting two component Bose–Einstein condensates and nonexistence of vortices for small rotation. *Communications in Mathematical Physics*, 336(2):509–579, 2015.
- [2] Nicholas D Alikakos and Peter W Bates. On the singular limit in a phase field model of phase transitions. In Annales de l'Institut Henri Poincare (C) Non Linear Analysis, volume 5, pages 141–178. Elsevier, 1988.
- [3] Antonio Ambrosetti, Andrea Malchiodi, and Wei-Ming Ni. Singularly perturbed elliptic equations with symmetry: existence of solutions concentrating on spheres, part I. Communications in mathematical physics, 235(3):427–466, 2003.
- [4] Tom Claeys, Arno BJ Kuijlaars, and Maarten Vanlessen. Multi-critical unitary random matrix ensembles and the general Painlevé II equation. Annals of Mathematics, 168:601– 641, 2008.
- [5] Marcel G Clerc, Juan Diego Dávila, Michał Kowalczyk, Panayotis Smyrnelis, and Estefania Vidal-Henriquez. Theory of light-matter interaction in nematic liquid crystals and the second Painlevé equation. *Calculus of Variations and Partial Differential Equations*, 56(4):93, 2017.
- [6] Marcel G Clerc, Michal Kowalczyk, and Panayotis Smyrnelis. Gradient theory of domain walls in thin, nematic liquid crystals films. arXiv preprint arXiv:1809.01034, 2018.

- [7] Marcel G Clerc, Michał Kowalczyk, and Panayotis Smyrnelis. Symmetry breaking and restoration in the Ginzburg–Landau model of nematic liquid crystals. *Journal of Nonlinear Science*, 28(3):1079–1107, 2018.
- [8] Athanassios S Fokas, Alexander R Its, Andrei A Kapaev, Andrei I Kapaev, Victor Yu Novokshenov, and V IU Novokshenov. *Painlevé transcendents: the Riemann-Hilbert approach*. Number 128. American Mathematical Soc., 2006.
- Bengt Fornberg and JAC Weideman. A computational exploration of the second Painlevé equation. Foundations of Computational Mathematics, 14(5):985–1016, 2014.
- [10] Stuart P Hastings and John Bryce Mcleod. A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation. Archive for Rational Mechanics and Analysis, 73(1):31–51, 1980.
- [11] Radu Ignat and Vincent Millot. The critical velocity for vortex existence in a two-dimensional rotating Bose–Einstein condensate. *Journal of Functional Analysis*, 233(1):260–306, 2006.
- [12] Georgia Karali and Christos Sourdis. The ground state of a Gross-Pitaevskii energy with general potential in the Thomas-Fermi limit. Archive for Rational Mechanics and Analysis, 217(2):439–523, 2015.
- [13] Lotfi Lassoued and Petru Mironescu. Ginzburg-landau type energy with discontinuous constraint. Journal d'Analyse Mathématique, 77(1):1–26, 1999.
- [14] Kimie Nakashima. Multi-layered stationary solutions for a spatially inhomogeneous Allen-Cahn equation. Journal of Differential Equations, 191(1):234–276, 2003.
- [15] Stephen Schecter and Christos Sourdis. Heteroclinic orbits in slow-fast hamiltonian systems with slow manifold bifurcations. *Journal of Dynamics and Differential Equations*, 22(4):629–655, 2010.
- [16] C Sourdis. On the uniqueness of solutions to a class of semilinear elliptic equations by Serrin's sweeping principle. *Rend. Sem. Mat. Univ. Politec. Torino Bruxelles-Torino Talks* in PDEs Turin, May 25, 2016, 74(2).
- [17] William C Troy. The role of Painleve II in predicting new liquid crystal self-assembly mechanisms. Archive for Rational Mechanics and Analysis, 227(1):367–385, 2018.

INSTITUTE OF APPLIED AND COMPUTATIONAL MATHEMATICS, FORTH, GR711 10 HER-AKLION, GREECE.

E-mail address: sourdis@uoc.gr