Vacuum Constraints for Realistic Strongly Coupled Heterotic M-Theories

Burt A. Ovrut *

Department of Physics, University of Pennsylvania Philadelphia, PA 19104–6396

Abstract

The compactification from the 11-dimensional Horava-Witten orbifold to 5dimensional heterotic M-theory on a Schoen Calabi-Yau threefold is reviewed, as is the specific SU(4) vector bundle leading to the "heterotic standard model" in the observable sector. A generic formalism for a consistent hidden sector gauge bundle, within the context of strongly coupled heterotic M-theory, is presented. Anomaly cancellation and the associated bulk space 5-branes are discussed in this context. The further compactification to a 4-dimensional effective field theory on a linearized BPS double domain wall is then presented to order $\kappa_{11}^{4/3}$. Specifically, the generic constraints required for anomaly cancellation and by the linearized domain wall solution, as well as restrictions imposed by the necessity to have positive squared gauge couplings to order $\kappa_{11}^{4/3}$ are presented in detail. Finally, the expression for the Fayet-Iliopoulos term associated with any anomalous U(1)gauge connection is presented and its role in N = 1 spontaneous supersymmetry breaking in the low energy effective theory is discussed.

^{*} Submitted to the special issue "Supersymmetric Field Theory 2018" in *Symmetry* ovrut@elcapitan.hep.upenn.edu

Contents

1	Intr	oduction	1
2	The	Compactification Vacuum	3
	2.1	The $d=11 \rightarrow d=5$ Compactification	3
		2.1.1 The Calabi-Yau Threefold	3
		2.1.2 The Observable Sector Gauge Bundle	5
		2.1.3 The Hidden Sector Gauge Bundle	6
		2.1.4 Bulk Space Five-Branes	8
		2.1.5 Anomaly Cancellation	8
	2.2		10
		2.2.1 The Linearized Double Domain Wall	10
3	The	$d=4 E_8 \times E_8$ Effective Theory	13
	3.1	The $\kappa_{11}^{2/3}$ Lagrangian	13
	3.2	The $\kappa_{11}^{4/3}$ Lagrangian	17
	0	3.2.1 Gauge Threshold Corrections	18
		3.2.2 Corrections to a Fayet-Iliopoulos Term	20
	3.3	A Specific Class of Examples	$\frac{-\circ}{22}$
Bibliography			26

1 Introduction

One of the major prerogatives of the Large Hadron Collider (LHC) at CERN is to search for low-energy N = 1 supersymmetry. It has long been known that specific vacua of both the weakly coupled [1,2] and strongly coupled [3–5] $E_8 \times E_8$ heterotic superstring can produce effective theories with at least a quasi-realistic particle spectrum exhibiting N = 1 supersymmetry [6–10]. It has also been shown that the various moduli associated with these theories can, in principle, be stabilized [11]. It is of considerable interest, therefore, to examine these low-energy theories in detail and to confront them, and their predictions, with present and up-coming CERN data.

There are several important criterion that such theories should possess. First, they must exhibit a low-energy spectrum in the observable sector that is not immediately in conflict with known phenomenology. In this paper, we will choose the so-called "heterotic standard model" introduced in [12–16]. The observable sector of this theory, associated with the first E_8 gauge factor, contains exactly the matter spectrum of the minimal supersymmetric standard model (MSSM) augmented by three right-handed neutrino chiral superfields, one per family, and a single pair of Higgs-Higgs conjugate superfields. There are no vector-like pairs of superfields or exotic matter of any kind. The low-energy gauge group of this theory is the $SU(3)_C \times SU(2)_L \times U(1)_Y$ of the standard model enhanced by a single additional gauged U(1) symmetry, which can be associated with B - L. Interestingly, the observable sector of this model-derived from the "top down" in the $E_8 \times E_8$ heterotic superstring—is identical to the minimal, anomaly free B - L extended MSSM, derived from the "bottom up" in [17, 18]. The requisite radiative breaking of both the B - L and electroweak symmetry, the associated mass hierarchy [19–23], and various phenomenological [24, 25] and cosmological [26–29] aspects of this low-energy theory have been discussed. Their detailed predictions for LHC observations are currently being explored [30–34]. The reader is also referred to works on heterotic standard models in [35–38]

The second important criterion is that there be a hidden sector, associated with the second E_8 gauge factor, that, prior to its spontaneous breaking, exhibits N = 1supersymmetry. The original papers on the heterotic standard model, in analogy with the KKLT mechanism [39] in Type II string theory, allowed the hidden sector to contain an anti-5-brane-thus explicitly breaking N = 1 supersymmetry [40, 41]. This was done so that the potential energy could admit a meta-stable de Sitter space vacuum. Such a vacuum was shown to exist, and its physical properties explored, in [42]. Be this as it may, it is important to know whether or not completely N = 1supersymmetric hidden sectors can be constructed in this context. It was demonstrated in [15] that the hidden sector of the heterotic standard model satisfies the Bogomolov inequality, a non-trivial necessary condition for supersymmetry [43] (see also [44–46]). More recently, within weakly coupled $E_8 \times E_8$ heterotic superstring theory, an N = 1 supersymmetric hidden sector was explicitly constructed using holomorphic line bundles [47]. This vacuum satisfies all conditions required to have a consistent compactification in the weakly coupled context-and is an explicit proof that completely N = 1 supersymmetric heterotic standard models exist. Be that as it may, it remains important to construct a completely N = 1 supersymmetric vacuum state of *strongly* coupled heterotic string theory that is totally consistent with all physical requirements. In this paper, we present the precise constraints required by such a vacuum within the context of strongly coupled heterotic M-theory.

Specifically, we do the following. In Section 2, the focus is on the compactification from the 11-dimensional Horava-Witten orbifold to 5-dimensional heterotic M-theorypresenting the relevant details of the explicit Calabi-Yau threefold and the observable sector SU(4) vector bundle of the heterotic standard model. Following a formalism first presented in [48-53], we discuss the generic structure of a large class of the hidden sector gauge bundles-a Whitney sum of a non-Abelian SU(N) bundle and line bundles. Bulk space 5-branes [54–56] and the anomaly cancellation constraint are presented in this context. The compactification of 5-dimensional heterotic Mtheory to the 4-dimensional low-energy effective field theory on a BPS linearized double domain wall is then described. In particular, we derive the constraints that must be satisfied in order for the linearized approximation to be valid. Section 3 is devoted to discussing the 4-dimensional $E_8 \times E_8$ effective theory; first the lowest order $\kappa_{11}^{2/3}$ Lagrangian [57]-along with the slope-stability criteria for the observable sector SU(4) vector bundle-and then the exact form of the order $\kappa_{11}^{4/3}$ corrections [58]. These corrections for both the observable and hidden sector gauge couplings and the Fayet-Iliopoulos terms are presented in a unified formalism. They are shown to be of the

same form as in the weakly coupled string with the weak coupling constants replaced by a moduli-dependent expansion parameter. The constraints required for there to be positive squared gauge coupling parameters, as well as the constraints so that the low energy theory be either N = 1 supersymmetric or to exhibit spontaneously broken N = 1 supersymmetry, are then specified. Finally, we consider a specific set of vacua for which the hidden sector vector bundle is restricted to be the Whitney sum of one non-Abelian SU(N) bundle with a single line bundle, while allowing only one fivebrane in the bulk space. Under these circumstances, the constraint equations greatly simplify and are explicitly presented. We demonstrate how the parameters associated with these bundles are computed for a specific choice of these objects.

2 The Compactification Vacuum

N = 1 supersymmetric heterotic M-theory on four-dimensional Minkowski space M_4 is obtained from eleven-dimensional Horava-Witten theory via two sequential dimensional reductions: First with respect to a Calabi-Yau threefold X whose radius is assumed to be smaller than that of the S^1/\mathbb{Z}_2 orbifold and second on a "linearized" BPS double domain wall solution of the effective five-dimensional theory. Let us present the relevant information for each of these within the context of the heterotic standard model.

2.1 The $d=11 \rightarrow d=5$ Compactification

2.1.1 The Calabi-Yau Threefold

The Calabi-Yau manifold X is chosen to be a torus-fibered threefold with fundamental group $\pi_1(X) = \mathbb{Z}_3 \times \mathbb{Z}_3$. Specifically, it is a fiber product of two rational elliptic $d\mathbb{P}_9$ surfaces, that is, a self-mirror Schoen threefold [13, 59], quotiented with respect to a freely acting $\mathbb{Z}_3 \times \mathbb{Z}_3$ isometry. Its Hodge data is $h^{1,1} = h^{1,2} = 3$ and, hence, there are three Kähler and three complex structure moduli. The complex structure moduli will play no role in the present paper. Relevant here is the degree-two Dolbeault cohomology group

$$H^{1,1}(X,\mathbb{C}) = \operatorname{span}_{\mathbb{C}}\{\omega_1,\omega_2,\omega_3\}$$
(2.1)

where $\omega_i = \omega_{ia\bar{b}}$ are harmonic (1, 1)-forms on X with the property

$$\omega_3 \wedge \omega_3 = 0, \quad \omega_1 \wedge \omega_3 = 3 \ \omega_1 \wedge \omega_1, \quad \omega_2 \wedge \omega_3 = 3 \ \omega_2 \wedge \omega_2 \ . \tag{2.2}$$

Defining the intersection numbers as

$$d_{ijk} = \frac{1}{v} \int_X \omega_i \wedge \omega_j \wedge \omega_k, \quad i, j, k = 1, 2, 3$$
(2.3)

where v is a reference volume of dimension $(length)^6$, it follows from (2.2) that

$$(d_{ijk}) = \begin{pmatrix} (0, \frac{1}{3}, 0) & (\frac{1}{3}, \frac{1}{3}, 1) & (0, 1, 0) \\ (\frac{1}{3}, \frac{1}{3}, 1) & (\frac{1}{3}, 0, 0) & (1, 0, 0) \\ (0, 1, 0) & (1, 0, 0) & (0, 0, 0) \end{pmatrix}.$$
(2.4)

The (i, j)-th entry in the matrix corresponds to the triplet $(d_{ijk})_{k=1,2,3}$.

Our analysis will require the Chern classes of the tangent bundle TX. Noting that the associated structure group is $SU(3) \subset SO(6)$, it follows that $\operatorname{rank}(TX) = 3$ and $c_1(TX) = 0$. Furthermore, the self-mirror property of this specific threefold implies $c_3(TX) = 0$. Finally, we find that

$$c_2(TX) = \frac{1}{v^{2/3}} (12\omega_1 \wedge \omega_1 + 12\omega_2 \wedge \omega_2).$$
 (2.5)

We will use the fact that if one chooses the generators of SU(3) to be hermitian, then the second Chern class of the tangent bundle can be written as

$$c_2(TX) = -\frac{1}{16\pi^2} \operatorname{tr}_{SO(6)} R \wedge R, \qquad (2.6)$$

where R is the Lie algebra valued curvature two-form.

Note that $H^{2,0} = H^{0,2} = 0$ on a Calabi-Yau threefold. It follows that $H^{1,1}(X, \mathbb{C}) = H^2(X, \mathbb{R})$ and, hence, ω_i , i = 1, 2, 3 span the real vector space $H^2(X, \mathbb{R})$. Furthermore, it was shown in [14] that the curve Poincare dual to each two-form ω_i is effective. It follows that the Kähler cone is the positive octant

$$\mathcal{K} = H^2_+(X, \mathbb{R}) \subset H^2(X, \mathbb{R}).$$
(2.7)

The Kähler form, defined to be $\omega_{a\bar{b}} = ig_{a\bar{b}}$ where $g_{a\bar{b}}$ is the Calabi-Yau metric, can be any element of \mathcal{K} . That is, suppressing the Calabi-Yau indices, the Kähler form can be expanded as

$$\omega = a^{i}\omega_{i}, \quad a^{i} > 0, \quad i = 1, 2, 3.$$
(2.8)

The real, positive coefficients a^i are the three (1, 1) Kähler moduli of the Calabi-Yau threefold. Here, and throughout this paper, upper and lower $H^{1,1}$ indices are summed unless otherwise stated. The dimensionless volume modulus is defined by

$$V = \frac{1}{v} \int_X \sqrt[6]{g} \tag{2.9}$$

and, hence, the dimensionful Calabi-Yau volume is $\mathbf{V} = vV$. Using the definition of the Kähler form and (2.3), V can be written as

$$V = \frac{1}{6v} \int_X \omega \wedge \omega \wedge \omega = \frac{1}{6} d_{ijk} a^i a^j a^k.$$
(2.10)

It is useful to express the three (1,1) moduli in terms of V and two additional independent moduli. This can be accomplished by defining the scaled shape moduli

$$b^i = V^{-1/3} a^i, \quad i = 1, 2, 3.$$
 (2.11)

It follows from (2.10) that they satisfy the constraint

$$d_{ijk}b^i b^j b^k = 6 \tag{2.12}$$

and, hence, represent only two degrees of freedom. Finally, note that all moduli defined thus far, that is, the a^i , V and b^i are functions of the five coordinates x^{α} , $\alpha = 0, \ldots, 3, 11$ of $M_4 \times S^1/\mathbb{Z}_2$, where $x^{11} \in [0, \pi\rho]$.

2.1.2 The Observable Sector Gauge Bundle

On the observable orbifold plane, the vector bundle $V^{(1)}$ on X is chosen to be holomorphic with structure group $SU(4) \subset E_8$, thus breaking

$$E_8 \longrightarrow Spin(10).$$
 (2.13)

To preserve N = 1 supersymmetry in four-dimensions, $V^{(1)}$ must be both slopestable and have vanishing slope [14,15]. In the context of this paper, these constraints are most easily examined in the d = 4 effective theory and, hence, will be discussed in Section 3 below. Finally, when two *flat* Wilson lines are turned on, each generating a different \mathbb{Z}_3 factor of the $\mathbb{Z}_3 \times \mathbb{Z}_3$ holonomy of X, the observable gauge group can be further broken to *

$$Spin(10) \longrightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}.$$
 (2.14)

Our analysis will require the Chern classes of $V^{(1)}$. Since the structure group is SU(4), it follows immediately that $\operatorname{rank}(V^{(1)}) = 4$ and $c_1(V^{(1)}) = 0$. The heterotic standard model is constructed so as to have the observed three chiral families of quarks/leptons and, hence, $V^{(1)}$ is constructed so that $c_3(V^{(1)}) = 3$. Finally, we found in [12, 14] that

$$c_2(V^{(1)}) = \frac{1}{v^{2/3}} (\omega_1 \wedge \omega_1 + 4 \,\omega_2 \wedge \omega_2 + 4 \,\omega_1 \wedge \omega_2).$$
(2.15)

Here, and below, it will be useful to note the following. Let \mathcal{V} be an arbitrary vector bundle on X with structure group \mathcal{G} , and $\mathcal{F}^{\mathcal{V}}$ the associated Lie algebra valued twoform gauge field strength. If the generators of \mathcal{G} are chosen to be hermitian, then

$$\frac{1}{8\pi^2} \operatorname{tr}_{\mathcal{G}} \mathcal{F}^{\mathcal{V}} \wedge \mathcal{F}^{\mathcal{V}} = ch_2(\mathcal{V}) = \frac{1}{2}c_1(\mathcal{V}) \wedge c_1(\mathcal{V}) - c_2(\mathcal{V}), \qquad (2.16)$$

where $ch_2(\mathcal{V})$ is the second Chern character of \mathcal{V} . Furthermore, we denote by $tr_{\mathcal{G}}$ the trace in the fundamental representation of the structure group \mathcal{G} of the bundle. When applied to the vector bundle $V^{(1)}$ in the observable sector, it follows from $c_1(V^{(1)}) = 0$ that

$$c_2(V^{(1)}) = -\frac{1}{8\pi^2} \operatorname{tr}_{SU(4)} F^{(1)} \wedge F^{(1)} = -\frac{1}{16\pi^2} \operatorname{tr}_{E_8} F^{(1)} \wedge F^{(1)}, \qquad (2.17)$$

where $F^{(1)}$ is the gauge field strength for the visible sector bundle $V^{(1)}$ and tr_{E_8} indicates the trace is over the fundamental **<u>248</u>** representation of E_8 . Note that the conventional normalization of the trace tr_{E_8} includes a factor of $\frac{1}{30}$, the inverse of the dual Coxeter number of E_8 . We have expressed $c_2(V^{(1)})$ in terms of tr_{E_8} since the fundamental SU(4) representation must be embedded into the adjoint representation of E_8 in the observable sector.

For the visible sector bundle $V^{(1)}$ with structure group SU(4), the group-theoretic embedding is simply the standard $SU(4) \subset SU(9) \hookrightarrow E_8$.

^{*}As discussed in [17,18], the two U(1) factor groups depend on the explicit choice of Wilson lines. For the renormalization group analysis of the low-energy d=4 theory, it is more convenient to choose $U(1)_{T_{3R}} \times U(1)_{B-L}$. However, since this is not our concern in this paper, we present the more canonical choice $U(1)_Y \times U(1)_{B-L}$.

2.1.3 The Hidden Sector Gauge Bundle

On the hidden orbifold plane, we will consider more general vector bundles and group embeddings. Specifically, in this paper, we will restrict any choice of hidden sector bundle to have the generic form of a Whitney sum

$$V^{(2)} = \mathcal{V}_N \oplus \mathcal{L} , \qquad \mathcal{L} = \bigoplus_{r=1}^R L_r$$
 (2.18)

where \mathcal{V}_N is a slope-stable, non-Abelian bundle and each L_r , $r = 1, \ldots, R$ is a holomorphic line bundle with structure group U(1). Note that a subset of hidden sector vector bundles might have no non-Abelian factor at all, being composed entirely of the sum of one or more line bundles. On the other hand, one could choose the hidden sector bundle to be composed entirely of a non-Abelian vector bundle, that is, with no line bundle factors. Should the hidden sector bundle contain a non-Abelian factor, one could generically choose it to possess an arbitrary structure group. However, in this paper, for specificity, we will assume that the structure group of the non-Abelian factor is SU(N) for some N. The explicit embeddings of the SU(N) and individual U(1) structure groups into the hidden sector E_8 gauge group will be discussed below. Finally, to preserve N = 1 supersymmetry in four-dimensions, $V^{(2)}$, being a Whitney sum of vector bundles, must be poly-stable–generically with vanishing slope (but, importantly, see Section 3.2.1 below). As with the observable sector vector bundle, these constraints are most easily examined in the d = 4 effective theory and, hence, will be discussed in Section 3 below. Let us first examine the non-Abelian factor.

• Hidden Sector SU(N) Factor

Since the structure group of \mathcal{V}_N is SU(N), it follows immediately that rank $(\mathcal{V}_N) = N$ and $c_1(\mathcal{V}_N) = 0$. The precise form of the second Chern class depends on the type of \mathcal{V}_N bundle one chooses. Since this bundle is no longer constrained to give any particular spectrum it, and its associated second Chern class, can be quite general. The generic form for the second Chern class is given by

$$c_2(\mathcal{V}_N) = \frac{1}{v^{2/3}} (c_N^{ij} \omega_i \wedge \omega_j)$$
(2.19)

where c_N^{ij} are, a priori, arbitrary real coefficients. Finally, note from (2.16) that since $c_1(\mathcal{V}_N) = 0$,

$$ch_2(\mathcal{V}_N) = -c_2(\mathcal{V}_N) . \tag{2.20}$$

Let us now consider the line bundle factors.

• Hidden Sector Line Bundles

Let us briefly review the properties of holomorphic line bundles on our specific geometry. Line bundles are classified by the divisors of X and, hence, equivalently by

the elements of the integral cohomology

$$H^{2}(X,\mathbb{Z}) = \{a\omega_{1} + b\omega_{2} + c\omega_{3} | a, b, c \in \mathbb{Z}\}.$$
(2.21)

It is conventional to denote the line bundle associated with the element $a\omega_1 + b\omega_2 + c\omega_3$ of $H^2(X,\mathbb{Z})$ as

$$\mathcal{O}_X(a,b,c). \tag{2.22}$$

Furthermore, in order for these bundles to arise from $\mathbb{Z}_3 \times \mathbb{Z}_3$ equivariant line bundles on the covering space of X, they must satisfy the additional constraint that

$$a + b = 0 \mod 3.$$
 (2.23)

Finally, as discussed in [47], for the purposes of constructing a heterotic gauge bundle from $\mathcal{O}_X(a, b, c)$, (2.23) is the only constraint required on the integers a, b, c. Specifically, it is not necessary to impose that these integers be even for there to exist a spin structure on $V^{(2)}$.

We will choose the Abelian factor of the hidden bundle to be

$$\mathcal{L} = \bigoplus_{r=1}^{R} L_r, \quad L_r = \mathcal{O}_X(\ell_r^1, \ell_r^2, \ell_r^3)$$
(2.24)

where

$$(\ell_r^1 + \ell_r^2) \mod 3 = 0, \quad r = 1, \dots, R$$
 (2.25)

for any positive integer R. The structure group is $U(1)^R$, where each U(1) factor has a specific embedding into the hidden sector E_8 gauge group. It follows from the definition that rank(\mathcal{L}) = R and that the first Chern class is

$$c_1(\mathcal{L}) = \sum_{r=1}^R c_1(L_r), \quad c_1(L_r) = \frac{1}{v^{1/3}} (\ell_r^1 \omega_1 + \ell_r^2 \omega_2 + \ell_r^3 \omega_3).$$
(2.26)

Note that since \mathcal{L} is a sum of holomorphic line bundles, $c_2(\mathcal{L}) = c_3(\mathcal{L}) = 0$. However, the relevant quantity for the hidden sector vacuum is the second Chern character defined in (2.16). For \mathcal{L} this becomes

$$ch_2(\mathcal{L}) = \sum_{r=1}^R ch_2(L_r)$$
 (2.27)

Since $c_2(L_r) = 0$, it follows that

$$ch_2(L_r) = 2a_r \frac{1}{2}c_1(L_r) \wedge c_1(L_r)$$
 (2.28)

where

$$a_r = \frac{1}{4 \cdot 30} \operatorname{tr}_{E_8} Q_r^2 \tag{2.29}$$

with Q_r the generator of the *r*-th U(1) factor embedded into the <u>**248**</u> representation of the hidden sector E_8 . Computation of this *a* coefficient depends on the choice of the hidden sector and will be discussed in more detail in Section 3.3.

The relevant topological object in the analysis of this paper will be the second Chern character of the complete hidden sector bundle

$$ch_2(V^{(2)}) = ch_2(\mathcal{V}_N \oplus \mathcal{L}) = ch_2(\mathcal{V}_N) + ch_2(\mathcal{L}) . \qquad (2.30)$$

Using (2.20) and (2.27), (2.28) this becomes

$$ch_2(V^{(2)}) = -c_2(\mathcal{V}_N) + \sum_{r=1}^R a_r c_1(L_r) \wedge c_1(L_r)$$
 (2.31)

with a_r given in (2.29). Note from (2.16), the explicit embedding of the structure group of $V^{(2)}$ into E_8 and (2.31) that

$$\frac{1}{16\pi^2} \operatorname{tr}_{E_8} F^{(2)} \wedge F^{(2)} = -c_2(\mathcal{V}_N) + \sum_{r=1}^R a_r c_1(L_r) \wedge c_1(L_r) \ . \tag{2.32}$$

2.1.4 Bulk Space Five-Branes

In addition to the holomorphic vector bundles on the observable and hidden orbifold planes, the bulk space between these planes can contain five-branes wrapped on twocycles $C_2^{(n)}$, $n = 1, \ldots, N$ in X. Cohomologically, each such five-brane is described by the (2, 2)-form Poincare dual to $C_2^{(n)}$, which we denote by $W^{(n)}$. Note that to preserves N = 1 supersymmetry in the four-dimensional theory, these curves must be holomorphic and, hence, each $W^{(n)}$ is an effective class.

2.1.5 Anomaly Cancellation

As discussed in [60, 61], anomaly cancellation in heterotic M-theory requires that

$$\sum_{n=0}^{N+1} J^{(n)} = 0, \qquad (2.33)$$

where

$$J^{(0)} = -\frac{1}{16\pi^2} \Big(\operatorname{tr}_{E_8} F^{(1)} \wedge F^{(1)} - \frac{1}{2} \operatorname{tr}_{SO(6)} R \wedge R \Big)$$

$$J^{(n)} = W^{(n)}, \quad n = 1, \dots, N,$$

$$J^{(N+1)} = -\frac{1}{16\pi^2} \Big(\operatorname{tr}_{E_8} F^{(2)} \wedge F^{(2)} - \frac{1}{2} \operatorname{tr}_{SO(6)} R \wedge R \Big)$$
(2.34)

Note that the indices n = 0 and n = N + 1 denote the observable and hidden sector domain walls respectively, and *not* the location of a five-brane. Using (2.6), (2.17) and (2.32), the anomaly cancellation condition can be expressed as

$$c_2(TX) - c_2(V^{(1)}) - c_2(\mathcal{V}_N) + \sum_{r=1}^R a_r c_1(L_r) \wedge c_1(L_r) - W = 0, \qquad (2.35)$$

where $W = \sum_{n=1}^{N} W^{(n)}$ is the total five-brane class.

Condition (2.35) is expressed in terms of four-forms in $H^4(X, \mathbb{R})$. We find it easier to analyze its consequences by writing it in the dual homology space $H_2(X, \mathbb{R})$. In this case, the coefficient of the *i*-th vector in the basis dual to $(\omega_1, \omega_2, \omega_3)$ is given by wedging each term in (2.35) with ω_i and integrating over X. Using (2.5), (2.15) and the intersection numbers (2.3), (2.4) gives

$$\frac{1}{v^{1/3}} \int_X \left(c_2(TX) - c_2(V^{(1)}) \right) \wedge \omega_i = \left(\frac{4}{3}, \frac{7}{3}, -4\right)_i, \quad i = 1, 2, 3.$$
 (2.36)

For $c_2(\mathcal{V}_N)$, it follows from (2.3) and (2.19) that

$$\frac{1}{v^{1/3}} \int_X \left(-c_2(\mathcal{V}_N) \right) \wedge \omega_i = -d_{ijk} c_N^{jk} .$$
(2.37)

Similarly, (2.3), (2.4) and (2.26) imply

$$\frac{1}{v^{1/3}} \int_X c_1(L_r) \wedge c_1(L_r) \wedge \omega_i = d_{ijk} \ell_r^j \ell_r^k , \quad i = 1, 2, 3.$$
 (2.38)

Defining

$$W_i = \frac{1}{v^{1/3}} \int_X W \wedge \omega_i , \qquad (2.39)$$

it follows that the anomaly condition (2.35) can be expressed as

$$W_{i} = \left(\frac{4}{3}, \frac{7}{3}, -4\right)\Big|_{i} - d_{ijk}c_{N}^{jk} + \sum_{r=1}^{R} a_{r}d_{ijk}\ell_{r}^{j}\ell_{r}^{k} \ge 0 , \quad i = 1, 2, 3.$$
 (2.40)

The positivity constraint on W follows from the requirement that it be an effective class to preserve N = 1 supersymmetry.

Finally, it is useful to define the charges

$$\beta_i^{(n)} = \frac{1}{v^{1/3}} \int_X J^{(n)} \wedge \omega_i , \quad i = 1, 2, 3.$$
(2.41)

For example, it follows from (2.34), using (2.5), (2.6), (2.15), (2.17) and the intersection numbers (2.3), (2.4), that

$$\beta_i^{(0)} = \left(\frac{2}{3}, -\frac{1}{3}, 4\right)\big|_i \,. \tag{2.42}$$

Note that the anomaly condition (2.33) can now be expressed as

$$\sum_{n=0}^{N+1} \beta_i^{(n)} = 0 . (2.43)$$

2.2 The $d=5 \rightarrow d=4$ Compactification

2.2.1 The Linearized Double Domain Wall

The five-dimensional effective theory of heterotic M-theory, obtained by dimensionally reducing Horava-Witten theory on the above Calabi-Yau threefold, admits a BPS double domain wall solution with five-branes in the bulk space [54, 56, 57, 60, 62, 63]. This solution depends on the previously defined moduli V and b^i as well as the a, b functions of the five-dimensional metric

$$ds_5^2 = a^2 dx^{\mu} dx^{\nu} \eta_{\mu\nu} + b^2 dy^2 , \qquad (2.44)$$

all of which are dependent on the five coordinates x^{α} , $\alpha = 0, \ldots, 3, 11$ of $M_4 \times S^1/\mathbb{Z}_2$. Denoting the reference radius of S^1 by ρ , then $x^{11} \in [0, \pi\rho]$. These moduli can all be expressed in terms of functions f^i , i = 1, 2, 3 satisfying the equations

$$d_{ijk}f^j f^k = H_i av{2.45}$$

where each H_i is a linear function of $z = \frac{x^{11}}{\pi \rho}$ with $z \in [0, 1]$ and whose exact form depends on the number and position of five-branes in the bulk space. As a simple, and relevant, example, let us consider the case when there are no five-branes in the vacuum. Then

$$H_{i} = 4k\epsilon'_{S}\beta_{i}^{(0)}\left(z - \frac{1}{2}\right) + k_{i},$$
(2.46)

where

$$\epsilon'_S = \pi \epsilon_S \quad , \ \epsilon_S = \left(\frac{\kappa_{11}}{4\pi}\right)^{2/3} \frac{2\pi\rho}{v^{2/3}} \tag{2.47}$$

and the charge $\beta_i^{(0)}$ is given in (2.42). The k, k_i are independent of z, but otherwise arbitrary functions of the four-dimensional moduli.

We are unable to give an exact analytic solution of (2.45) and (2.46). However, one can obtain an approximate solution by expanding to linear order in $\epsilon'_S \beta_i^{(0)} (z - \frac{1}{2})$. It is clear from (2.46) that this approximation will be valid under the conditions that

$$2\epsilon_S'|k\beta_i^{(0)}| \ll |k_i| \tag{2.48}$$

for each i = 1, 2, 3. This *linearized* solution was discussed in detail in [54,56,57]. Here we present only the results required in this paper. For an arbitrary dimensionless function f of the five $M_4 \times S^1/\mathbb{Z}_2$ coordinates, define its average over the S^1/\mathbb{Z}_2 orbifold interval as

$$\langle f \rangle_{11} = \frac{1}{\pi \rho} \int_0^{\pi \rho} dx^{11} f ,$$
 (2.49)

where ρ is a reference length. Then $\langle f \rangle_{11}$ is a function of the four coordinates x^{μ} , $\mu = 0, \ldots, 3$ of M_4 only. The linearized solution is expressed in terms of orbifold average functions

$$V_0 = \langle V \rangle_{11}, \quad b_0^i = \langle b^i \rangle_{11}, \quad \frac{\hat{R}_0}{2} = \langle b \rangle_{11}.$$
 (2.50)

We have defined $\frac{\hat{R}_0}{2} = \langle b \rangle_{11}$ to conform to specific normalization later in the paper. One then finds that the linearized solution specifies that

$$k = \frac{\hat{R}_0}{2V_0^{2/3}} , \ k_i = d_{ijk} b_0^j b_0^k V_0^{1/3} .$$
(2.51)

In terms of these averaged moduli, the conditions (2.48) for the validity of the linearized approximation can be written as

$$\epsilon_S' \frac{\widehat{R}_0}{V_0} \left| \beta_i^{(0)} \right| \ll d_{ijk} b_0^j b_0^k , \qquad (2.52)$$

where we have removed the absolute value of the right-hand side since all elements of d_{ijk} given in (2.4) and each field b^i are non-negative. Equivalently, one can write

$$\epsilon'_S \frac{\widehat{R}_0}{V_0} \ll d_{ijk} b_0^j b_0^k \left| \beta_i^{(0)} \right|^{-1}$$
 (2.53)

for each i = 1, 2, 3. These conditions are actually much simpler than they first appear. This can be seen by writing out each equation for i = 1, 2, 3 respectively. Using (2.42), we find that

$$i = 1:$$
 $q \quad \frac{2}{3}\epsilon'_S \frac{\hat{R}_0}{V_0} \ll \frac{2}{3}b_0^1 b_0^2 + \frac{1}{3}(b_0^2)^2 + 2b_0^2 b_0^3$ (2.54)

$$i = 2: \qquad \frac{1}{3}\epsilon'_{S}\frac{\dot{R}_{0}}{V_{0}} \ll \frac{1}{3}(b_{0}^{1})^{2} + \frac{2}{3}b_{0}^{1}b_{0}^{2} + 2b_{1}^{2}b_{3}^{3}$$
(2.55)

$$i = 3:$$
 $4\epsilon'_S \frac{R_0}{V_0} \ll 2b_0^1 b_0^2$ (2.56)

Clearly, if the equation for i = 3 is satisfied then equations i = 1, 2 are automatically satisfied as well. Hence, one can replace the constraint (2.53) by the simpler requirement that

$$2\epsilon'_S \frac{\hat{R}_0}{V_0} \ll b_0^1 b_0^2 . (2.57)$$

Assuming that these conditions are fulfilled, the linearized solution for V, b^i , a and b can be determined in terms of the orbifold average functions. For example, assuming there are no five-branes in the bulk space, the linearized solution for V is given by

$$V = V_0 \left(1 + \epsilon'_S \frac{\hat{R}_0}{V_0} b_0^i \beta_i^{(0)} \left(z - \frac{1}{2} \right) \right) .$$
 (2.58)

The linearized expressions for b^i and a, b are similar expansions in the moduli dependent quantity $(\epsilon'_S \frac{\hat{R}_0}{V_0}) b_0^i \beta_i^{(0)}$. It follows that another check on the validity of these expansions is that

$$\frac{1}{2} \epsilon'_{S} \frac{\widehat{R}_{0}}{V_{0}} \left| b^{i}_{0} \beta^{(0)}_{i} \right| \ll 1 .$$
(2.59)

However, using (2.57) it is straight forward to show that if constraint (2.52) is satisfied then (2.59) will be fulfilled automatically. To see this, note, using (2.42) and constraint (2.57), that

$$\frac{1}{2}\epsilon'_{S}\frac{\widehat{R}_{0}}{V_{0}}\left|b_{0}^{i}\beta_{i}^{(0)}\right| = \frac{1}{2}\epsilon'_{S}\frac{\widehat{R}_{0}}{V_{0}}\left|\frac{2}{3}b_{0}^{1} - \frac{1}{3}b_{0}^{2} + 4b_{0}^{3}\right| \ll \frac{b_{0}^{1}b_{0}^{2}}{4}\left|\frac{2}{3}b_{0}^{1} - \frac{1}{3}b_{0}^{2} + 4b_{0}^{3}\right| .$$
(2.60)

That is,

$$\frac{1}{2}\epsilon'_{S}\frac{\widehat{R}_{0}}{V_{0}}\left|b_{0}^{i}\beta_{i}^{(0)}\right| \ll \left|\frac{1}{6}(b_{0}^{1})^{2}b_{0}^{2} - \frac{1}{12}b_{0}^{1}(b_{0}^{2})^{2} + b_{0}^{1}b_{0}^{2}b_{0}^{3}\right|$$
(2.61)

However, it follows from (2.4) for d_{ijk} that

$$\frac{1}{6}d_{ijk}b_0^i b_0^j b_0^k = \frac{1}{6}(b_0^1)^2 b_0^2 + \frac{1}{6}b_0^1 (b_0^2)^2 + b_0^1 b_0^2 b_0^3 = \frac{1}{6}(b_0^1)^2 b_0^2 - \frac{1}{12}b_0^1 (b_0^2)^2 + b_0^1 b_0^2 b_0^3 + \frac{1}{4}b_0^1 (b_0^2)^2 .$$
(2.62)

Therefore

$$\left|\frac{1}{6}(b_0^1)^2 b_0^2 - \frac{1}{12}b_0^1(b_0^2)^2 + b_0^1 b_0^2 b_0^3\right| = \left|\frac{1}{6}d_{ijk}b_0^i b_0^j b_0^k - \frac{1}{4}b_0^1(b_0^2)^2\right| = \left|1 - \frac{1}{4}b_0^1(b_0^2)^2\right| , \quad (2.63)$$

where we have used expression (2.12). However, it is clear from (2.61) that

$$0 < \frac{1}{4}b_0^1(b_0^2)^2 < \frac{3}{2} \tag{2.64}$$

and therefore

$$\frac{1}{2} \epsilon'_S \frac{\widehat{R}_0}{V_0} \left| b_0^i \beta_i^{(0)} \right| \ll \left| 1 - \frac{1}{4} b_0^1 (b_0^2)^2 \right| < 1 .$$
(2.65)

Hence, condition (2.59) is automatically satisfied if constraint (2.52) is.

Thus far, we have considered the case when there are no five-branes in the bulk space. Including an arbitrary number of five-branes in the linearized BPS solution is straightforward and was presented in [60,62,63]. Here, it will suffice to generalize the above discussion to the case of one five-brane located at $z_1 \in [0, 1]$. The conditions for the validity of the linear approximation then break into two parts. Written in terms of the averaged moduli, these are

$$2\epsilon'_{S}\frac{R_{0}}{V_{0}}\left|\beta_{i}^{(0)}\left(z-\frac{1}{2}\right)-\frac{1}{2}\beta_{i}^{(1)}(1-z_{1})^{2}\right| \ll \left|d_{ijk}b_{0}^{j}b_{0}^{k}\right|, \quad z \in [0, z_{1}]$$
(2.66)

and

$$2\epsilon'_{S}\frac{\widehat{R}_{0}}{V_{0}}\left| (\beta_{i}^{(0)} + \beta_{i}^{(1)})(z - \frac{1}{2}) - \frac{1}{2}\beta_{i}^{(1)}z_{1}^{2} \right| \ll \left| d_{ijk}b_{0}^{j}b_{0}^{k} \right|, \quad z \in [z_{1}, 1].$$
(2.67)

Assuming these conditions are satisfied, the linearized solution for V, b_i and a, b can be determined in each region. For example, the linearized solution for V is given by

$$V = V_0 \left(1 + \epsilon'_S \frac{\hat{R}_0}{V_0} b_0^i \left[\beta_i^{(0)} \left(z - \frac{1}{2} \right) - \frac{1}{2} \beta_i^{(1)} (z_1 - 1)^2 \right] \right), \quad z \in [0, z_1]$$
(2.68)

and

$$V = V_0 \left(1 + \epsilon'_S \frac{\widehat{R}_0}{V_0} b_0^i \left[(\beta_i^{(0)} + \beta_i^{(1)}) \left(z - \frac{1}{2} \right) - \frac{1}{2} \beta_i^{(1)} z_1^2 \right] \right), \quad z \in [z_1, 1] .$$
 (2.69)

It follows that the conditions for the validity of this linearized solution for V are given by

$$\epsilon'_{S} \frac{R_{0}}{V_{0}} b_{0}^{i} \Big| \beta_{i}^{(0)} \left(z - \frac{1}{2} \right) - \frac{1}{2} \beta_{i}^{(1)} (z_{1} - 1)^{2} \Big| \ll 1, \quad z \in [0, z_{1}]$$

$$(2.70)$$

and

$$\epsilon'_{S} \frac{\hat{R}_{0}}{V_{0}} b_{0}^{i} \Big| (\beta_{i}^{(0)} + \beta_{i}^{(1)}) (z - \frac{1}{2}) - \frac{1}{2} \beta_{i}^{(1)} z_{1}^{2} \Big| \ll 1, \quad z \in [z_{1}, 1] .$$

$$(2.71)$$

Note that if the five-brane is located near the hidden wall, that is, $z_1 \rightarrow 1$, conditions (2.66) and (2.67) for the validity of the linear approximation both revert to (2.52), and, hence, inequality (2.57), as they must for consistency. Similarly, the conditions (2.70) and (2.71) for the validity of the solutions for V-as well as for the expansions of b_i and a, b- simply reduce to (2.59). Again, we find that conditions (2.70) and (2.71) will be automatically satisfied if the strong coupling constraints (2.66) and (2.67) are. For this reason, we henceforth consider the strong coupling constraints only.

When dimensionally reduced on this linearized BPS solution, the four-dimensional functions a_0^i , V_0 , b_0^i and \hat{R}_0 will become moduli of the d = 4 effective heterotic M-theory. The geometric role of a_0^i and V_0 , b_0^i will remain the same as above—now, however, for the averaged Calabi-Yau threefold. For example, the dimensionful volume of the averaged Calabi-Yau manifold will be given by vV_0 . The new dimensionless quantity \hat{R}_0 will be the length modulus of the orbifold. The dimensionful length of S^1/\mathbb{Z}_2 is given by $\pi\rho\hat{R}_0$. Finally, since the remainder of this paper will be within the context of the d = 4 effective theory, we will, for simplicity, drop the subscript "0" on all moduli henceforth.

3 The d=4 $E_8 \times E_8$ Effective Theory

When d = 5 heterotic M-theory is dimensionally reduced to four dimensions on the linearized BPS double domain wall with five-branes, the result is an N = 1 super-symmetric effective four-dimensional theory with (potentially spontaneously broken) $E_8 \times E_8$ gauge group. The Lagrangian will break into two distinct parts. The first contains terms of order $\kappa_{11}^{2/3}$ in the eleven-dimensional Planck constant κ_{11} , while the second consists of terms of order $\kappa_{11}^{4/3}$.

3.1 The $\kappa_{11}^{2/3}$ Lagrangian

This Lagrangian is well-known and was presented in [57]. Here we discuss only those properties required in this paper. In four dimensions, the moduli must be organized into the lowest components of chiral supermultiplets. Here, we need only consider the real part of these components. Additionally, one specifies that these chiral multiplets have *canonical* Kähler potentials in the effective Lagrangian. The dilaton is simply given by

$$\operatorname{Re} S = V . \tag{3.1}$$

However, neither a^i nor b^i have canonical kinetic energy. To obtain this, one must define the rescaled moduli

$$t^i = \widehat{R}b^i = \widehat{R}V^{-1/3}a^i , \qquad (3.2)$$

where we have used (2.11), and choose the complex Kähler moduli T^i so that

$$\operatorname{Re} T^i = t^i \ . \tag{3.3}$$

Denote the real modulus specifying the location of the *n*-th five-brane in the bulk space by $z_n = \frac{x_n^{11}}{\pi\rho}$ where n = 1, ..., N. As with the Kähler moduli, it is necessary to define the fields

$$\operatorname{Re} Z^n = \beta_i^{(n)} t^i z_n \ . \tag{3.4}$$

These rescaled Z^n five-brane moduli have canonical kinetic energy.

The gauge group of the d = 4 theory has two E_8 factors, the first associated with the observable sector and the second with the hidden sector. As discussed previously, both vector bundles must be chosen so as to preserve N = 1 supersymmetry in fourdimensions. We now explicitly discuss the conditions under which this will be true. We begin with the observable sector.

• Stability of the Observable Sector Vector Bundle

To preserve N = 1 supersymmetry in four-dimensions the holomorphic SU(4) vector bundle $V^{(1)}$ associated with the observable E_8 gauge group must be both slopestable and have vanishing slope [64–66]. The slope of any bundle or sub-bundle \mathcal{F} is defined as

$$\mu(\mathcal{F}) = \frac{1}{\operatorname{rank}(\mathcal{F})v^{2/3}} \int_X c_1(\mathcal{F}) \wedge \omega \wedge \omega , \qquad (3.5)$$

where ω is the Kähler form in (2.8)—now, however, written in terms of the a^i moduli averaged over S^1/\mathbb{Z}_2 . Since $c_1(V^{(1)}) = 0$, $V^{(1)}$ has vanishing slope. But, is it slopestable? As proven in detail in [15], this will be the case in a subspace of the Kähler cone defined by seven inequalities required for all sub-bundles of $V^{(1)}$ to have negative slope. These can be slightly simplified into the statement that the moduli a^i , i = 1, 2, 3must satisfy at least one of the two inequalities

$$\begin{pmatrix} a^1 < a^2 \le \sqrt{\frac{5}{2}}a^1 & \text{and} & a^3 < \frac{-(a^1)^2 - 3a^1a^2 + (a^2)^2}{6a^1 - 6a^2} \end{pmatrix} \text{ or } \\ \left(\sqrt{\frac{5}{2}}a^1 < a^2 < 2a^1 & \text{and} & \frac{2(a^2)^2 - 5(a^1)^2}{30a^1 - 12a^2} < a^3 < \frac{-(a^1)^2 - 3a^1a^2 + (a^2)^2}{6a^1 - 6a^2} \end{pmatrix}$$
(3.6)

The subspace \mathcal{K}^s satisfying (3.6) is a full-dimensional subcone of the Kähler cone \mathcal{K} defined in (2.7). It is a cone because the inequalities are homogeneous. In other




Figure 1: The visible sector stability region in the Kähler cone.

words, only the angular part of the Kähler moduli are constrained, but not the overall volume. Hence, it is best displayed as a two-dimensional "star map" as seen by an observer at the origin. This is shown in Figure 1. For Kähler moduli restricted to this subcone, the four-dimensional low energy theory in the observable sector is N = 1 supersymmetric.

• Poly-Stability of the Hidden Sector Vector Bundle

To preserve N = 1 supersymmetry in four-dimensions, the hidden sector vector bundle must satisfy two conditions, First, since it is generically a Whitney sum, the vector bundle must be poly-stable. That is, each factor of the Whitney sum must be slope-stable and, in addition, all factors in the sum must have the same slope. Second, generically, this slope must vanish identically–but with one important caveat discussed in Section 3.3.2. In order to make this more concrete, we now present three non-trivial examples to illustrate the these two conditions. As a first example, let us choose

1. $V^{(2)} = \mathcal{V}_N$:

First, in this case, since for a single vector bundle slope-stability implies polystability, one need only check that \mathcal{V}_N is slope-stable. For example, one could choose \mathcal{V}_N to be identical to the SU(4) bundle in the observable sector, $V^{(1)}$, presented above. Note that, since we are restricting all hidden sector non-Abelian bundles to have structure group SU(N), it follows that $\mu(\mathcal{V}_N)$ must vanish, thus satisfying the second condition for N = 1 supersymmetry. As with the observable sector bundle SU(4) bundle, stability of a generic non-Abelian vector bundle will only occur within a specific region of Kähler moduli space.

2. $V^{(2)} = L$:

As in the previous case, one need only check that the line bundle L is slopestable, which will imply poly-stability. Fortunately, every line bundle is trivially slope-stable, so any line bundle can be used. It is important to note that the slope of a line bundle which appears as a lone factor in the Whitney sum has, a priori, no further constraints-that is, $\mu(L)$ need not vanish. Using (3.5), (2.26) and (2.4), it follows that the slope of an arbitrary line bundle specified by $L = \mathcal{O}_X(\ell^1, \ell^2, \ell^3)$ is given by

$$\mu(L) = d_{ijk}\ell^i a^j a^k$$

= $\frac{1}{3} \Big(a^2(a^2 + 6a^3)\ell^1 + (a^1)^2\ell^2 + 6a^1a^3\ell^2 + 2a^1a^2(\ell^1 + \ell^2 + 3\ell^3) \Big).$ (3.7)

That is, its value is a highly specific function of the Kähler moduli. We will discuss the requirements that such a line bundle lead to four-dimensional N = 1 supersymmetry in Section 3.2.2 below.

3.
$$V^{(2)} = \mathcal{V}_N \oplus L$$
:

As specified above, the non-Abelian vector bundle \mathcal{V}_N must be slope-stable in a region of Kähler moduli space. Furthermore, since we are restricting the structure group in our discussion to be SU(N), it follows that $\mu(\mathcal{V}_N) = 0$. As we just indicated, any line bundle L will be slope-stable everywhere in Kähler moduli space. However, the full Whitney sum $V^{(2)} = \mathcal{V}_N \oplus L$ will be poly-stableand, hence, preserve N = 1 supersymmetry-if and only if $\mu(L) = \mu(\mathcal{V}_N) = 0$. That is, because of the existence of a non-Abelian SU(N) factor, the line bundle L now has the additional constraint that its slope vanish identically. It is clear from (3.7) that this will be the case only in a restricted region of Kähler moduli space. It follows that the full Whitney sum $V^{(2)} = \mathcal{V}_N \oplus L$ will only be a viable hidden sector bundle if the region of stability of \mathcal{V}_N has a non-vanishing intersection with the region where the slope of L vanishes. This is a very nontrivial requirement. To give a concrete example, let us choose $\mathcal{V}_N = V^{(1)}$, where $V^{(1)}$ is the SU(4) observable sector bundle specified above. Recall that the region of slope-stability of this bundle in Kähler moduli space is delineated by the inequalities in (3.6) and shown in Figure 1. Plotted in 3-dimensions, this region of slope-stability over a limited region of Kähler moduli space is shown in Figure 2(a). Furthermore, let us specify, for example, that $L = \mathcal{O}_X(1,2,-3)$. Note that L satisfies condition (2.23), as it must. It follows from (3.7) that the region of moduli space in which $\mu(L) = 0$ is given by the equation

$$\frac{2}{3}(a^1)^2 - 4a^1a^2 + \frac{1}{3}(a^2)^2 + 4a^1a^3 + 2a^2a^3 = 0.$$
 (3.8)

Plotted over a limited region of Kähler moduli space in 3-dimensions, the region where $\mu(L) = 0$ is shown in Figure 2(b). Figure 2(c) then shows that these two regions have a substantial overlap in Kähler moduli space. Furthermore, since \mathcal{V}_N was chosen to be $V^{(1)}$, it follows that Figure 2(c) also represents the overlap with the stability region of the observable sector vector bundle. We conclude that the specific choice of $V^{(2)} = V^{(1)} \oplus \mathcal{O}_X(1,2,-3)$ is, potentially, a suitable choice for a poly-stable hidden sector vector bundle.




Figure 2: The region of poly-stability for the hidden sector vector bundle $V^{(2)} = V^{(1)} \oplus \mathcal{O}_X(1,2,-3)$. The red volume in Figure 2(a) is a sub-region of Kähler moduli space where the bundle $V^{(1)}$ is slope stable, whereas the green volume of Figure 2(b) is the sub-region where $\mu(\mathcal{O}_X(1,2,-3)) = 0$. They have a substantial region of overlap in Kähler moduli space, indicated by the yellow volume in Figure 3(c).

These three examples give the rules for constructing specific poly-stable vector bundles. They can easily be generalized to construct generic poly-stable Whitney sum hidden sector vector bundles.

3.2 The $\kappa_{11}^{4/3}$ Lagrangian

The terms in the BPS double domain wall solution proportional to ϵ'_S lead to order $\kappa_{11}^{4/3}$ additions to the d = 4 Lagrangian. These have several effects. The simplest is that the five-brane location moduli now contribute to the definition of the dilaton,

which becomes

$$\operatorname{Re} S = V + \epsilon'_{S} \sum_{n=1}^{N} \beta_{i}^{(n)} t^{i} z_{n}^{2} , \qquad (3.9)$$

where the fields t^i are defined in (3.2). More profoundly, these $\kappa_{11}^{4/3}$ terms lead, first, to threshold corrections to the gauge coupling parameters and, second, to additions to the Fayet-Iliopoulos (FI) term associated with any anomalous U(1) factor in the low energy gauge group. Let us analyze these in turn.

3.2.1 Gauge Threshold Corrections

The gauge couplings of the non-anomalous components of the d = 4 gauge group, in both the observable and hidden sectors, have been computed to order $\kappa_{11}^{4/3}$ in [54]. Written in terms of the fields b^i defined in (2.11) and including five-branes in the bulk space, these are given by

$$\frac{4\pi}{(g^{(1)})^2} \propto V(1 + \epsilon'_S \frac{\widehat{R}}{2V} \sum_{n=0}^N (1 - z_n)^2 b^i \beta_i^{(n)})$$
(3.10)

and

$$\frac{4\pi}{(g^{(2)})^2} \propto V(1 + \epsilon'_S \frac{\widehat{R}}{2V} \sum_{n=1}^{N+1} z_n^2 b^i \beta_i^{(n)})$$
(3.11)

respectively. The positive definite constant of proportionality is identical for both gauge couplings and is not relevant to the present discussion. It is important to note that the effective parameter of the $\kappa_{11}^{2/3}$ expansion in (3.10) and (3.11), namely $\epsilon'_S \hat{R}$, is identical to 1) the parameter appearing in (2.52) (and its five-brane extension (2.66) and (2.67)) for the validity of the linearized approximation, as well as 2) the $\kappa_{11}^{2/3}$ expansion parameter for V presented in (2.58) (and its five-brane extension (2.68) and (2.70)). That is, the effective strong coupling parameter of the $\kappa_{11}^{2/3}$ expansion is given by

$$\epsilon_S^{\text{eff}} = \epsilon'_S \frac{\widehat{R}}{V} \ . \tag{3.12}$$

We point out that this is, up to a constant factor of order one, precisely the strong coupling parameter presented in equation (1.3) of [67].

Recall that n = 0 and n = N + 1 correspond to the observable and hidden sector domain walls—not to five-branes. Therefore, $z_0 = 0$ and $z_{N+1} = 1$. Using (2.34) and (2.41), one can evaluate the $\beta_i^{(n)}$ coefficients in terms of the the $a^i, i = 1, 2, 3$ Kähler moduli defined in (2.8). Rewritting the above expressions in terms of these moduli using (2.6), (2.8), (2.10), (2.11), (2.27), (2.28), as well as redefining the five-brane moduli to be

$$\lambda_n = z_n - \frac{1}{2}, \quad \lambda_n \in \left[-\frac{1}{2}, \frac{1}{2}\right], \tag{3.13}$$

we find that

$$\frac{4\pi}{(g^{(1)})^2} \propto \frac{1}{6v} \int_X \omega \wedge \omega \wedge \omega - \epsilon'_S \frac{\widehat{R}}{2V^{1/3}} \frac{1}{v^{1/3}} \\ \times \int_X \omega \wedge \left(-c_2(V^{(1)}) + \frac{1}{2}c_2(TX) - \sum_{n=1}^N (\frac{1}{2} - \lambda_n)^2 W^{(n)} \right)$$
(3.14)

and

$$\frac{4\pi}{(g^{(2)})^2} \propto \frac{1}{6v} \int_X \omega \wedge \omega \wedge \omega - \epsilon'_S \frac{\widehat{R}}{2V^{1/3}} \frac{1}{v^{1/3}} \\ \times \int_X \omega \wedge \left(-c_2(\mathcal{V}_N) + \sum_{r=1}^R a_r c_1(L_r) \wedge c_1(L_r) + \frac{1}{2} c_2(TX) - \sum_{n=1}^N (\frac{1}{2} + \lambda_n)^2 W^{(n)} \right)$$
(3.15)

where a_r is given in (2.29). The first term on the right-hand side, that is, the volume V defined in (2.10), is the order $\kappa_{11}^{2/3}$ result. The remaining terms are the $\kappa_{11}^{4/3}$ M-theory corrections first presented in [54].

Clearly, consistency of the d = 4 effective theory requires both $(g^{(1)})^2$ and $(g^{(2)})^2$ to be positive. It follows that the moduli of the four-dimensional theory are constrained to satisfy

$$\frac{1}{v} \int_{X} \omega \wedge \omega \wedge \omega - 3\epsilon'_{S} \frac{\widehat{R}}{V^{1/3}} \frac{1}{v^{1/3}} \int_{X} \omega \wedge (-c_{2}(V^{(1)}) + \frac{1}{2}c_{2}(TX) - \sum_{n=1}^{N} (\frac{1}{2} - \lambda_{n})^{2} W^{(n)}) > 0 \quad (3.16)$$

and

$$\frac{1}{v} \int_{X} \omega \wedge \omega \wedge \omega - 3\epsilon'_{S} \frac{\widehat{R}}{V^{1/3}} \frac{1}{v^{1/3}} \int_{X} \omega \wedge \left(-c_{2}(\mathcal{V}_{N}) + \sum_{r=1}^{R} a_{r}c_{1}(L_{r}) \wedge c_{1}(L_{r}) + \frac{1}{2}c_{2}(TX) - \sum_{n=1}^{N} (\frac{1}{2} + \lambda_{n})^{2}W^{(n)} \right) > 0. \quad (3.17)$$

One can use (2.3), (2.4), (2.5), (2.8), (2.15), (2.19), (2.26) and (2.39) to rewrite these expressions as

$$d_{ijk}a^{i}a^{j}a^{k} - 3\epsilon'_{S}\frac{\widehat{R}}{V^{1/3}}\Big(-(\frac{8}{3}a^{1} + \frac{5}{3}a^{2} + 4a^{3}) + 2(a^{1} + a^{2}) - \sum_{n=1}^{N}(\frac{1}{2} - \lambda_{n})^{2}a^{i}W_{i}^{(n)}\Big) > 0$$

$$(3.18)$$

and

$$d_{ijk}a^{i}a^{j}a^{k} - 3\epsilon'_{S}\frac{\widehat{R}}{V^{1/3}} \Big(-d_{ijk}a^{i}c_{N}^{jk} + d_{ijk}a^{i}\sum_{r=1}^{R}a_{r}\ell_{r}^{j}\ell_{r}^{k} + 2(a^{1} + a^{2}) - \sum_{n=1}^{N}(\frac{1}{2} + \lambda_{n})^{2}a^{i}W_{i}^{(n)} \Big) > 0$$

$$(3.19)$$

respectively. It is of interest to compare the $(g^{(1)})^2$, $(g^{(2)})^2 > 0$ conditions calculated to order $\kappa_{11}^{4/3}$ in strongly coupled heterotic M-theory, that is, (3.16) and (3.17), to the one-loop corrected conditions computed in the weakly coupled heterotic string [51]. Assuming the same observable and hidden sector vector bundles used in this paper, we find that the weakly coupled conditions for $(g^{(1)})^2$, $(g^{(2)})^2 > 0$, derived using equation (3.103) in [51], are identical to (3.16) and (3.17) if one replaces

$$g_s^2 l_s^4 \longrightarrow \epsilon_S' \frac{\hat{R}}{V^{1/3}} v^{2/3}$$
 (3.20)

in the weak coupling formulas, where g_s and $l_s = 2\pi\sqrt{\alpha'}$ are the weak coupling parameter and the string length respectively and ϵ'_S is defined in (2.47).

3.2.2 Corrections to a Fayet-Iliopoulos Term

In the heterotic standard model vacuum, the observable sector vector bundle $V^{(1)}$ has structure group SU(4). Hence, it does not lead to an anomalous U(1) gauge factor in the observable sector of the low energy theory. However, the hidden sector bundle $V^{(2)}$ introduced above, in addition to a possible non-Abelian bundle \mathcal{V}_N , consists of a sum of line bundles with the additional structure group $U(1)^R$. Each U(1) factor leads to an anomalous U(1) gauge group in the four-dimensional effective field theory and, hence, an associated *D*-term. Let L_r be any one of the irreducible line bundles of $V^{(2)}$. The string one-loop corrected Fayet-Iliopoulos (FI) term for L_r was computed in [48] within the context of the weakly coupled heterotic string. Comparing various results in the literature, it is straightforward to show that strong coupling results to order $\kappa_{11}^{4/3}$ can be obtained from string one-loop weak coupling expressions using the same replacement $g_s^2 l_s^4 \longrightarrow \epsilon'_S \frac{\hat{R}}{V^{1/3}} v^{2/3}$ presented in (3.20). Making this substitution, we find that the expression for the FI term associated with L_r in strongly coupled heterotic M-theory to order $\kappa_{11}^{4/3}$ is given by

$$FI_r = \frac{3}{16} \frac{\epsilon_S \epsilon_r^2}{\kappa_4^2} \frac{1}{\widehat{R} V^{2/3}} \Big(\mu(L_r) + \epsilon'_S \frac{\widehat{R}}{V^{1/3}} \int_X c_1(L_r) \wedge \Big(J^{(N+1)} + \sum_{n=1}^N z_n^2 J^{(n)} \Big) \Big), \quad (3.21)$$

where $\mu(L_r)$ is given in (3.5). We note that the $\kappa_{11}^{2/3}$ part of this expression is identical to that derived in [58]. Inserting (2.34), (2.6), (2.32) and, following the conventions

of [48, 51], redefining the five-brane moduli as in (3.13), we find that

$$FI_{r} = \frac{3}{16} \frac{\epsilon_{S} \epsilon_{r}^{2}}{\kappa_{4}^{2}} \frac{1}{\hat{R} V^{2/3}} \left(\mu(L_{r}) - \epsilon_{S}' \frac{\hat{R}}{V^{1/3}} \right)$$
$$\int_{X} c_{1}(L_{r}) \wedge \left(-c_{2}(\mathcal{V}_{N}) + \sum_{s=1}^{R} a_{s} c_{1}(L_{s}) \wedge c_{1}(L_{s}) + \frac{1}{2} c_{2}(TX) - \sum_{n=1}^{N} (\frac{1}{2} + \lambda_{n})^{2} W^{(n)} \right),$$
(3.22)

where a_s is given in (2.29). The first term on the right-hand side, that is, the slope of L_r , is the order $\kappa_{11}^{2/3}$ result. The remaining terms are the $\kappa_{11}^{4/3}$ M-theory corrections first presented in [54]. Note that the dimensionless parameter $\epsilon'_S \frac{\hat{R}}{V^{1/3}}$ of the $\kappa_{11}^{4/3}$ term is identical to the expansion coefficient of the linearized solution—when expressed in term of the a^i moduli—discussed in Subsection 2.2. See, for example, (2.52). Finally, recalling definition (3.5) of the slope, using (2.3), (2.4), (2.8), (2.26), (2.39) and noting from (2.5) that

$$\frac{1}{v^{1/3}} \int_X \frac{1}{2} c_2(TX) \wedge \omega_i = (2, 2, 0)_i, \quad i = 1, 2, 3,$$
(3.23)

it follows that for each L_r the associated Fayet-Iliopoulos factor FI_r in (3.22) can be written as

$$FI_{r} = \frac{3}{16} \frac{\epsilon_{S} \epsilon_{r}^{2}}{\kappa_{4}^{2}} \frac{1}{\widehat{R} V^{2/3}} \left(d_{ijk} \ell_{r}^{i} a^{j} a^{k} - \epsilon_{S}^{\prime} \frac{\widehat{R}}{V^{1/3}} \right) \\ \left(- d_{ijk} \ell_{r}^{i} c_{N}^{jk} + d_{ijk} \ell_{r}^{i} \sum_{s=1}^{R} a_{s} \ell_{s}^{j} \ell_{s}^{k} + \ell_{r}^{i}(2,2,0) |_{i} - \sum_{n=1}^{N} (\frac{1}{2} + \lambda_{n})^{2} \ell_{r}^{i} W_{i}^{(n)} \right)$$
(3.24)

where

$$V = \frac{1}{6} d_{ijk} a^i a^j a^k.$$
 (3.25)

As discussed in [54], the general form of each *D*-term in the low energy fourdimensional theory is the sum of 1) the moduli dependent FI parameter (3.24) and 2) terms quadratic in the four-dimensional scalar fields charged under the associated U(1) gauge symmetry weighted by their specific charge. For each line bundle L_r , $r = 1, \ldots, R$ on the Calabi-Yau threefold, there is an anomalous $U(1)_r$ symmetry in the four-dimensional low energy theory on the hidden sector. Written in terms of the simplified notation introduced in [56, 58], the associated D-term is given by

$$D_r = FI_r - \sum_{L,\bar{M}} Q_r^L G_{L\bar{M}} C_r^L \bar{C}_r^{\bar{M}}.$$
 (3.26)

Here $G_{L\bar{M}}$ is an hermitian metric on the $U(1)_r$ reducible space of all charged, mass dimension one, scalar matter fields C^L , which block diagonalizes into the allowed irreducible representations for the r-th line bundle. The indices L and M each run over the full reducible representation, breaking into a sum of the indices over each irreducible sector–each such sector with a unique charge Q. The metric $G_{L\bar{M}}$ is, in general, a complicated function of the Kähler moduli with positive definite eigenvalues. Note from (3.24) that FI_r has mass dimension two–consistent with expression (3.26). As is well-known, a necessary condition for a static vacuum state of the effective four-dimensional theory to be N = 1 supersymmetric is that the D term associated with each line bundle L_r must identically vanish. Generically, this will be the case if

$$\sum_{L,\bar{M}} Q_r^L G_{L\bar{M}} C_r^L \bar{C}_r^{\bar{M}} = F I_r .$$
(3.27)

These C_r^L scalars break into two distinct types– 1) those that transform only under the Abelian group $U(1)_r$ and 2) those which, in addition, transform non-trivially under the non-Abelian gauge factor of the hidden sector low energy theory. This second type of scalar field will also appear in the *D*-term associated with the non-Abelian group–which cannot contain a FI term. Hence, the demand that the vacuum be supersymmetric generically sets their vacuum expectation values to zero. It follow that one can, henceforth, ignore such fields and restrict the scalars in (3.27) to those that transform under the Abelian $U(1)_r$ symmetry only.

In the weakly coupled heterotic case discussed in [47], it was assumed, for simplicity, that the vacuum expectation values $\langle C_r^L \rangle$ all vanish, even for the scalars not transforming under the low energy non-Abelian gauge factor. In that case, each D_r will vanish if and only if $FI_r = 0$. This restriction puts very strong constraints on the choice of the hidden sector vector bundle. Be that as it may, the assumption that all $\langle C_r^L \rangle$ vanish and that $FI_r = 0$ remains a valid constraint for strongly coupled vacua. However, in the strongly coupled case we are now considering, an *alternative* set of constraints can be also be adopted. That is, one can assume that the scalar fields that only transform under the low energy $U(1)_r$ groups are, in general, non-vanishing and that each D_r is set to zero by the associated vacuum expectation values $\langle C_r^L \rangle$ becoming non-zero. For this to be the case, it is essential to specify the hidden sector vector bundle and to compute the pure $U(1)_r$ low energy scalar fields C_r^L and their associated charges Q_r^L . This is essential because, should the $U(1)_r$ charge be positive, then the associated D_r term can vanish if and only if $FI_r > 0$. On the other hand, if the associated charge is negative, then D_r can vanish if and only if $FI_r < 0$. That is, the condition one needs to impose on the Fayet-Iliopoulos terms will depend on the sign of the charges in the scalar spectrum.

3.3 A Specific Class of Examples

The constraint equations listed above are technically rather complicated. Therefore, as we did when discussing poly-stability in subsection 3.1, we now analyze the constraint equations within the context of a specific class of N = 1 supersymmetric hidden sector vector bundles. To do this, one must specify the non-Abelian bundle \mathcal{V}_N with structure group SU(N), the number of line bundles L_r and their exact embeddings into the hidden E_8 vector bundle. We will, henceforth, consider hidden sector bundles

that may, or may not, contain a non-Abelian factor and, for simplicity, are restricted to contain at most a single line bundle

$$L = \mathcal{O}_X(\ell^1, \ell^2, \ell^3) \tag{3.28}$$

where

$$\ell^1, \ell^2, \ell^3 \in \mathbb{Z}, \quad (\ell^1 + \ell^2) \mod 3 = 0.$$
 (3.29)

In this case, there is only a single a_r coefficient—which we denote simply by a. In addition, one must specify the number of five-branes in the bulk space. Again, for simplicity, we assume that there is only one five-brane in this example. It then follows from (2.40), (3.18), (3.19) and (3.24)that the constraints for this restricted class of examples are given by

$$W_i = \left(\frac{4}{3}, \frac{7}{3}, -4\right)\Big|_i - d_{ijk}c_N^{jk} + ad_{ijk}\ell^j\ell^k \ge 0, \quad i = 1, 2, 3, \tag{3.30}$$

$$d_{ijk}a^{i}a^{j}a^{k} - 3\epsilon'_{S}\frac{R}{V^{1/3}}\left(-\left(\frac{8}{3}a^{1} + \frac{5}{3}a^{2} + 4a^{3}\right) + 2(a^{1} + a^{2}) - \left(\frac{1}{2} - \lambda\right)^{2}a^{i}W_{i}\right) > 0$$

$$(3.31)$$

$$d_{ijk}a^{i}a^{j}a^{k} - 3\epsilon'_{S}\frac{\widehat{R}}{V^{1/3}} \Big(-d_{ijk}a^{i}c_{N}^{jk} + ad_{ijk}a^{i}\ell^{j}\ell^{k} + 2(a^{1} + a^{2}) - (\frac{1}{2} + \lambda)^{2}a^{i}W_{i} \Big) > 0$$
(3.32)

$$d_{ijk}\ell^{i}a^{j}a^{k} - \epsilon'_{S}\frac{\widehat{R}}{V^{1/3}} \Big(-d_{ijk}\ell^{i}c_{N}^{jk} + ad_{ijk}\ell^{i}\ell^{j}\ell^{k} + \ell^{i}(2,2,0)|_{i} - (\frac{1}{2} + \lambda)^{2}\ell^{i}W_{i} \Big) \leq 0, = 0$$
(3.33)

where \widehat{R} is an independent modulus and V satisfies relation (3.25). Note that the expression on the left-hand side of (3.33) is a) > 0 or < 0 if one assumes that some $\langle \phi_{\alpha} \rangle \neq 0$ and that the associated scalar charge q_{α} is positive or negative respectively or b) = 0 if, alternatively, one assumes that all $\langle \phi_{\alpha} \rangle = 0$.

To proceed, one must specify the the coefficient a, as well as the coefficients c_N^{jk} of the second Chern class of \mathcal{V}_N . We begin with the coefficient a. Recall from (2.29) that

$$a = \frac{1}{4 \cdot 30} \operatorname{tr}_{E_8} Q^2 , \qquad (3.34)$$

where Q is the generator of the U(1) structure group of the line bundle. Hence, the value of coefficient a will depend entirely on the explicit embedding of this U(1) into the **<u>248</u>** representation of the hidden sector E_8 . Here, we will present one explicit example of such an embedding, although, as will discussed elsewhere, this specific type of embedding is not unique. First, assume that the hidden sector gauge bundle is the Whitney sum of a non-Abelian bundle \mathcal{V}_N and a line bundle L. Choosing the structure group of \mathcal{V}_N to be SU(N), embed

$$SU(N) \times U(1) \subset SU(N+1) . \tag{3.35}$$

The structure group of the line bundle L is identified with the specific U(1) generator in SU(N + 1) which commutes with the generators of the chosen SU(N). In the fundamental representation of SU(N + 1), the generator of U(1) can then be written as

$$\operatorname{diag}\left(\underbrace{1,\ldots,1}_{N},-N\right). \tag{3.36}$$

This specifies the exact embedding of U(1) into SU(N + 1). Now choose SU(N + 1) to be a factor of a maximal subgroup of E_8 . The decomposition of the **<u>248</u>** of E_8 with respect to this maximal subgroup, together with (3.36), then determines the generator Q.

This is most easily explained by giving a simple explicit example. Let us assume there is *no* non-Abelian bundle–only a single line bundle. That is,

$$V^{(2)} = L (3.37)$$

The explicit embedding of L into E_8 is chosen as follows. First, recall that

$$SU(2) \times E_7 \subset E_8 \tag{3.38}$$

is a maximal subgroup. With respect to $SU(2) \times E_7$, the **<u>248</u>** representation of E_8 decomposes as

$$\underline{\mathbf{248}} \longrightarrow (\underline{\mathbf{1}}, \underline{\mathbf{133}}) \oplus (\underline{\mathbf{2}}, \underline{\mathbf{56}}) \oplus (\underline{\mathbf{3}}, \underline{\mathbf{1}}). \tag{3.39}$$

Now choose the generator of the U(1) structure group in the fundamental representation of SU(2) to be (1, -1). It follows that under $SU(2) \rightarrow U(1)$

$$\underline{2} \longrightarrow 1 \oplus -1 \tag{3.40}$$

and, hence, under $U(1) \times E_7$

$$\underline{\mathbf{248}} \longrightarrow (0, \underline{\mathbf{133}}) \oplus \left((1, \underline{\mathbf{56}}) \oplus (-1, \underline{\mathbf{56}}) \right) \oplus \left((2, \underline{\mathbf{1}}) \oplus (0, \underline{\mathbf{1}}) \oplus (-2, \underline{\mathbf{1}}) \right).$$
(3.41)

The generator Q of this embedding of the line bundle can be read off from expression (3.41). Inserting this into (3.34), we find that

$$a = 1. \tag{3.42}$$

For the choice of a non-Abelian bundle \mathcal{V}_N with structure group SU(N), similar calculations give

$$N = 2 \Rightarrow a = 3, \quad N = 3 \Rightarrow a = 6, \quad N = 4 \Rightarrow a = 10, \quad N = 5 \Rightarrow a = 15$$
 (3.43)

As discussed previously, one may or may not include a non-Abelian factor in the hidden sector vector bundle. If a non-Abelian factor \mathcal{V}_N is to be included, one must specify it exactly. Generically, there are many possibilities for such a bundle. As an explicit example, let us choose this to be precisely the same SU(4) bundle as in the observable sector described in Subsection 2.1.2. Doing this greatly simplifies

the analysis since this \mathcal{V}_4 bundle is slope-stable with vanishing slope in the same region of Kähler moduli space as the observable sector bundle $V^{(1)}$ -that is, when the inequalities (3.6) are satisfied. Since N = 4, it follows from (3.43) that coefficient

$$a = 10 \tag{3.44}$$

and, since \mathcal{V}_4 is identical to $V^{(1)}$, it follows from (2.15) that

$$c_2(\mathcal{V}_4) = \frac{1}{v^{2/3}} \big(\omega_1 \wedge \omega_1 + 4 \, \omega_2 \wedge \omega_2 + 4 \, \omega_1 \wedge \omega_2 \big). \tag{3.45}$$

Hence, from (2.19) the only non-vanishing c_N^{jk} coefficients are

$$c_4^{11} = 1, \quad c_4^{22} = 4, \quad c_4^{12} = c_4^{21} = 2.$$
 (3.46)

Inserting these coefficients, along with a=10, into (3.30),(3.31),(3.32),(3.33) give the appropriate constraint equations for this class of vacua. As discussed above, if one assumes the VEVs of all scalar fields vanish, then the left-hand side of (3.33) must be zero. However, if not all $\langle \phi_{\alpha} \rangle$ vanish, then to determine whether the left-hand side of the FI inequality (3.31) should be > 0 or < 0 depends on the sign of the charge of the associated low energy U(1) charged scalars. Since the charge can be different for different choices of the hidden sector bundle, this can only be determined within the context of an explicit example. This will be presented elsewhere.

Of course, these constraints have to be solved simultaneously with the condition (3.6) for the slope-stability of both the observable and hidden sector non-Abelian vector bundles; that is

$$\left(a^{1} < a^{2} \le \sqrt{\frac{5}{2}}a^{1} \quad \text{and} \quad a^{3} < \frac{-(a^{1})^{2} - 3a^{1}a^{2} + (a^{2})^{2}}{6a^{1} - 6a^{2}} \right) \quad \text{or}$$

$$\left(\sqrt{\frac{5}{2}}a^{1} < a^{2} < 2a^{1} \quad \text{and} \quad \frac{2(a^{2})^{2} - 5(a^{1})^{2}}{30a^{1} - 12a^{2}} < a^{3} < \frac{-(a^{1})^{2} - 3a^{1}a^{2} + (a^{2})^{2}}{6a^{1} - 6a^{2}} \right)$$

$$(3.47)$$

Finally, it is essential to implement equations (2.66) and (2.67) for the validity of the linear approximation. These equations depend sensitively on the sign of each component of $\beta_i^{(0)}$, the value of $\beta_i^{(1)} = W_i$ and the five-brane location z_1 . For the specific class of models presented in this section, (2.66) and (2.67) become

$$2\epsilon'_{S}\frac{\widehat{R}}{V^{1/3}}\left|\beta_{i}^{(0)}\left(z-\frac{1}{2}\right)-\frac{1}{2}W_{i}\left(\frac{1}{2}-\lambda\right)^{2}\right|\ll\left|d_{ijk}a^{j}a^{k}\right|,\quad z\in[0,\lambda+\frac{1}{2}]$$
(3.48)

and

$$2\epsilon'_{S}\frac{\widehat{R}}{V^{1/3}}\left|\left(\beta_{i}^{(0)}+W_{i}\right)\left(z-\frac{1}{2}\right)-\frac{1}{2}W_{i}\left(\frac{1}{2}+\lambda\right)^{2}\right|\ll\left|d_{ijk}a^{j}a^{k}\right|,\quad z\in\left[\lambda+\frac{1}{2},1\right].$$
 (3.49)

where, as defined in (3.13), $\lambda = z_1 - \frac{1}{2}$.

Acknowledgments

The author would like to thank Yang-Hui He and Rehan Deen for many informative conversations. Burt Ovrut is supported in part by DOE No. DE-SC0007901 and SAS Account 020-0188-2-010202-6603-0338.

Bibliography

- David J. Gross, Jeffrey A. Harvey, Emil J. Martinec, and Ryan Rohm. Heterotic string theory. 1. the free heterotic string. *Nucl. Phys.*, B256:253, 1985.
- [2] David J. Gross, Jeffrey A. Harvey, Emil J. Martinec, and Ryan Rohm. Heterotic string theory. 2. the interacting heterotic string. Nucl. Phys., B267:75, 1986. 1
- [3] Edward Witten. Strong coupling expansion of Calabi-Yau compactification. Nucl. Phys., B471:135–158, 1996. 1
- [4] Petr Horava and Edward Witten. Heterotic and type I string dynamics from eleven dimensions. Nucl. Phys., B460:506–524, 1996. 1
- [5] Petr Horava and Edward Witten. Eleven-dimensional supergravity on a manifold with boundary. Nucl. Phys., B475:94–114, 1996. 1
- [6] Brian R. Greene, Kelley H. Kirklin, Paul J. Miron, and Graham G. Ross. A superstring inspired standard model. *Phys. Lett.*, B180:69, 1986. 1
- [7] Brian R. Greene, Kelley H. Kirklin, Paul J. Miron, and Graham G. Ross. A three generation superstring model. 1. compactification and discrete symmetries. *Nucl. Phys.*, B278:667, 1986. 1
- [8] Brian R. Greene, Kelley H. Kirklin, Paul J. Miron, and Graham G. Ross. A three generation superstring model. 2. symmetry breaking and the low-energy theory. *Nucl. Phys.*, B292:606, 1987. 1
- [9] Takeo Matsuoka and Daijiro Suematsu. Realistic models from the $E_8 \times E'_8$ superstring theory. *Prog. Theor. Phys.*, 76:886, 1986. 1
- [10] Brian R. Greene, K. H. Kirklin, P. J. Miron, and Graham G. Ross. 27³ Yukawa couplings for a three generation superstring model. *Phys. Lett.*, B192:111, 1987. 1
- [11] Lara B. Anderson, James Gray, Andre Lukas, and Burt Ovrut. Stabilizing All Geometric Moduli in Heterotic Calabi-Yau Vacua. *Phys. Rev.*, D83:106011, 2011.
- [12] Volker Braun, Yang-Hui He, Burt A. Ovrut, and Tony Pantev. The Exact MSSM spectrum from string theory. JHEP, 0605:043, 2006. 1, 2.1.2

- [13] Volker Braun, Burt A. Ovrut, Tony Pantev, and Rene Reinbacher. Elliptic Calabi-Yau threefolds with Z(3) x Z(3) Wilson lines. JHEP, 0412:062, 2004. 1, 2.1.1
- [14] Volker Braun, Yang-Hui He, Burt A. Ovrut, and Tony Pantev. Vector Bundle Extensions, Sheaf Cohomology, and the Heterotic Standard Model. Adv. Theor. Math. Phys., 10:4, 2006. 1, 2.1.1, 2.1.2, 2.1.2
- [15] Volker Braun, Yang-Hui He, and Burt A. Ovrut. Stability of the minimal heterotic standard model bundle. JHEP, 0606:032, 2006. 1, 2.1.2, 3.1
- [16] Volker Braun, Yang-Hui He, and Burt A. Ovrut. Yukawa couplings in heterotic standard models. JHEP, 0604:019, 2006. 1
- [17] Vernon Barger, Pavel Fileviez Perez, and Sogee Spinner. Minimal gauged U(1)(B-L) model with spontaneous R-parity violation. *Phys.Rev.Lett.*, 102:181802, 2009.
 1, *
- [18] Pavel Fileviez Perez and Sogee Spinner. Spontaneous R-Parity Breaking in SUSY Models. *Phys. Rev.*, D80:015004, 2009. 1, *
- [19] Michael Ambroso and Burt Ovrut. The B-L/Electroweak Hierarchy in Heterotic String and M-Theory. JHEP, 0910:011, 2009. 1
- [20] Michael Ambroso and Burt A. Ovrut. The B-L/Electroweak Hierarchy in Smooth Heterotic Compactifications. Int.J.Mod.Phys., A25:2631–2677, 2010. 1
- [21] Michael Ambroso and Burt A. Ovrut. The Mass Spectra, Hierarchy and Cosmology of B-L MSSM Heterotic Compactifications. *Int.J. Mod. Phys.*, A26:1569–1627, 2011. 1
- [22] Burt A. Ovrut, Austin Purves, and Sogee Spinner. Wilson Lines and a Canonical Basis of SU(4) Heterotic Standard Models. JHEP, 1211:026, 2012. 1
- [23] Burt A. Ovrut, Austin Purves, and Sogee Spinner. The minimal SUSY BL model: from the unification scale to the LHC. JHEP, 06:182, 2015. 1
- [24] Pavel Fileviez Perez and Sogee Spinner. TeV Scale Spontaneous R-Parity Violation. AIP Conf. Proc., 1200:529–532, 2010. 1
- [25] Pavel Fileviez Perez and Sogee Spinner. The Minimal Theory for R-parity Violation at the LHC. JHEP, 1204:118, 2012. 1
- [26] Tamaz Brelidze and Burt A. Ovrut. B-L Cosmic Strings in Heterotic Standard Models. JHEP, 1007:077, 2010. 1
- [27] Lorenzo Battarra, Michael Koehn, Jean-Luc Lehners, and Burt A. Ovrut. Cosmological Perturbations Through a Non-Singular Ghost-Condensate/Galileon Bounce. JCAP, 1407:007, 2014. 1

- [28] Michael Koehn, Jean-Luc Lehners, and Burt A. Ovrut. Cosmological superbounce. Phys. Rev., D90(2):025005, 2014. 1
- [29] Michael Koehn, Jean-Luc Lehners, and Burt A. Ovrut. Higher-Derivative Chiral Superfield Actions Coupled to N=1 Supergravity. *Phys. Rev.*, D86:085019, 2012.
- [30] Zachary Marshall, Burt A. Ovrut, Austin Purves, and Sogee Spinner. Spontaneous *R*-Parity Breaking, Stop LSP Decays and the Neutrino Mass Hierarchy. *Phys. Lett.*, B732:325–329, 2014. 1
- [31] Zachary Marshall, Burt A. Ovrut, Austin Purves, and Sogee Spinner. LSP Squark Decays at the LHC and the Neutrino Mass Hierarchy. *Phys. Rev.*, D90(1):015034, 2014. 1
- [32] Sebastian Dumitru, Burt A. Ovrut, and Austin Purves. The *R*-parity Violating Decays of Charginos and Neutralinos in the B-L MSSM. *JHEP*, 02:124, 2019. 1
- [33] Sebastian Dumitru, Burt A. Ovrut, and Austin Purves. *R*-parity Violating Decays of Wino Chargino and Wino Neutralino LSPs and NLSPs at the LHC. *JHEP*, 06:100, 2019. 1
- [34] Sebastian Dumitru, Christian Herwig, and Burt A. Ovrut. *R*-parity Violating Decays of Bino Neutralino LSPs at the LHC. *JHEP*, 12:042, 2019. 1
- [35] Philip Candelas, Xenia de la Ossa, Yang-Hui He, and Balazs Szendroi. Triadophilia: A Special Corner in the Landscape. Adv. Theor. Math. Phys., 12:429– 473, 2008. 1
- [36] Lara B. Anderson, James Gray, Andre Lukas, and Eran Palti. Two Hundred Heterotic Standard Models on Smooth Calabi-Yau Threefolds. *Phys.Rev.*, D84:106005, 2011. 1
- [37] Lara B. Anderson, James Gray, Andre Lukas, and Eran Palti. Heterotic Line Bundle Standard Models. JHEP, 1206:113, 2012. 1
- [38] Lara B. Anderson, James Gray, Andre Lukas, and Eran Palti. Heterotic standard models from smooth Calabi-Yau three-folds. PoS, CORFU2011:096, 2011. 1
- [39] Shamit Kachru, Renata Kallosh, Andrei D. Linde, and Sandip P. Trivedi. De Sitter vacua in string theory. *Phys.Rev.*, D68:046005, 2003. 1
- [40] James Gray, Andre Lukas, and Burt Ovrut. Perturbative anti-brane potentials in heterotic M-theory. *Phys. Rev.*, D76:066007, 2007. 1
- [41] James Gray, Andre Lukas, and Burt Ovrut. Flux, gaugino condensation and anti-branes in heterotic M-theory. *Phys. Rev.*, D76:126012, 2007. 1

- [42] Volker Braun and Burt A. Ovrut. Stabilizing moduli with a positive cosmological constant in heterotic M-theory. JHEP, 0607:035, 2006. 1
- [43] F. A. Bogomolov. Holomorphic tensors and vector bundles on projective manifolds. Izv. Akad. Nauk SSSR Ser. Mat., 42(6):1227–1287, 1439, 1978. 1
- [44] Michael R. Douglas, Rene Reinbacher, and Shing-Tung Yau. Branes, bundles and attractors: Bogomolov and beyond. 2006. 1
- [45] Bjorn Andreas and Gottfried Curio. Spectral Bundles and the DRY-Conjecture. J.Geom.Phys., 62:800–803, 2012. 1
- [46] Bjorn Andreas and Gottfried Curio. On the Existence of Stable bundles with prescribed Chern classes on Calabi-Yau threefolds. 2011. 1
- [47] Volker Braun, Yang-Hui He, and Burt A. Ovrut. Supersymmetric Hidden Sectors for Heterotic Standard Models. JHEP, 1309:008, 2013. 1, 2.1.3, 3.2.2
- [48] Ralph Blumenhagen, Gabriele Honecker, and Timo Weigand. Loop-corrected compactifications of the heterotic string with line bundles. *JHEP*, 0506:020, 2005. 1, 3.2.2, 3.2.2
- [49] Ralph Blumenhagen, Gabriele Honecker, and Timo Weigand. Non-Abelian brane worlds: The Heterotic string story. JHEP, 0510:086, 2005. 1
- [50] Ralph Blumenhagen, Sebastian Moster, and Timo Weigand. Heterotic GUT and standard model vacua from simply connected Calabi-Yau manifolds. *Nucl. Phys.*, B751:186–221, 2006. 1
- [51] T. Weigand. Compactifications of the heterotic string with unitary bundles. Fortsch.Phys., 54:963–1077, 2006. 1, 3.2.1, 3.2.2
- [52] Ralph Blumenhagen, Sebastian Moster, Rene Reinbacher, and Timo Weigand. Massless Spectra of Three Generation U(N) Heterotic String Vacua. JHEP, 0705:041, 2007. 1
- [53] Ralph Blumenhagen, Gabriele Honecker, and Timo Weigand. Supersymmetric (non-)Abelian bundles in the Type I and SO(32) heterotic string. *JHEP*, 0508:009, 2005. 1
- [54] Andre Lukas, Burt A. Ovrut, and Daniel Waldram. Nonstandard embedding and five-branes in heterotic M theory. *Phys.Rev.*, D59:106005, 1999. 1, 2.2.1, 2.2.1, 3.2.1, 3.2.1, 3.2.2, 3.2.2
- [55] Ron Donagi, Burt A. Ovrut, and Daniel Waldram. Moduli spaces of five-branes on elliptic Calabi-Yau threefolds. JHEP, 11:030, 1999. 1
- [56] Andre Lukas, Burt A. Ovrut, and Daniel Waldram. Five-branes and supersymmetry breaking in M theory. JHEP, 9904:009, 1999. 1, 2.2.1, 2.2.1, 3.2.2

- [57] Andre Lukas, Burt A. Ovrut, and Daniel Waldram. On the four-dimensional effective action of strongly coupled heterotic string theory. *Nucl. Phys.*, B532:43– 82, 1998. 1, 2.2.1, 2.2.1, 3.1
- [58] Lara B. Anderson, James Gray, Andre Lukas, and Burt Ovrut. Stability Walls in Heterotic Theories. JHEP, 0909:026, 2009. 1, 3.2.2, 3.2.2
- [59] Chad Schoen. On fiber products of rational elliptic surfaces with section. Math. Z., 197(2):177–199, 1988. 2.1.1
- [60] Andre Lukas, Burt A. Ovrut, K.S. Stelle, and Daniel Waldram. Heterotic M theory in five-dimensions. Nucl. Phys., B552:246–290, 1999. 2.1.5, 2.2.1, 2.2.1
- [61] Ron Donagi, Andre Lukas, Burt A. Ovrut, and Daniel Waldram. Nonperturbative vacua and particle physics in M theory. JHEP, 9905:018, 1999. 2.1.5
- [62] Andre Lukas, Burt A. Ovrut, K.S. Stelle, and Daniel Waldram. The Universe as a domain wall. *Phys. Rev.*, D59:086001, 1999. 2.2.1, 2.2.1
- [63] Ron Donagi, Andre Lukas, Burt A. Ovrut, and Daniel Waldram. Holomorphic vector bundles and nonperturbative vacua in M theory. *JHEP*, 9906:034, 1999. 2.2.1, 2.2.1
- [64] K. Uhlenbeck and S.-T. Yau. On the existence of hermitian-Yang-Mills connections in stable vector bundles. *Comm. Pure Appl. Math.*, 39(S, suppl.):S257–S293, 1986. Frontiers of the mathematical sciences: 1985 (New York, 1985). 3.1
- [65] S. K. Donaldson. Anti self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. Proc. London Math. Soc. (3), 50(1):1–26, 1985. 3.1
- [66] Michael B. Green, J. H. Schwarz, and Edward Witten. Superstring theory. vol.
 2: Loop amplitudes, anomalies and phenomenology. Cambridge, Uk: Univ. Pr. (1987) 596 P. (Cambridge Monographs On Mathematical Physics). 3.1
- [67] Tom Banks and Michael Dine. Couplings and scales in strongly coupled heterotic string theory. Nucl. Phys., B479:173–196, 1996. 3.2.1